1 **Age-linked heterogeneity among oligodendrocyte precursor cells in the cerebral**
2 **cortex of mice and human**

3 **Authors:** Bribián A[1,2], Medina-Rodríguez EM [1,7], Josa-Prado F[2], García-Álvarez I[1,3],
4 Machín-Díaz I[1], Murcia-Belmonte V[1,4], Vega-Zelaya L[5],Pastor J[5], Garrido L[6]and de
5 Castro, F[1,2]

6
[1]Hospital Nacional de Parapléjicos, Finca "La Peraleda" s/n, 45071 Toledo, Spain.
[2]Instituto Cajal-CSIC, Avda. Dr. Arce 37,28002 Madrid, Spain.
[3]Institutode Química Orgánica General (IQOG-CSIC), CSIC, Juan de la Cierva 3, 28006 Madrid, Spain.
[4]Instituto de Neurociencias, Universidad Miguel Hernández-CSIC, Campus San Juan de Alicante, 03550
Alicante, Spain.
[5]Neurofisiología Clínica; Instituto de Investigación Sanitaria, Hospital Universitario de la Princesa,
Madrid, Spain.
[6]Instituto de Ciencia y Tecnología de Polímeros (ICTP-CSIC), CSIC, Juan de la Cierva 3, 28006 Madrid,
Spain.
[7]University of Miami, Coral Gables, FL 33124, USA.

17
18 **Corresponding author:**
19
F. de Castro PhD, MD
Grupo de Neurobiología del Desarrollo-GNDe
Instituto Cajal-CSIC
Avda. Dr. Arce 37. 28002, Madrid, Spain.
Phone: 91-5854650
fdecastro@cajal.csic.es

26
27

ABSTRACT

Oligodendrocyte precursor cells (OPCs) are responsible for spontaneous remyelination after a demyelinating lesion. They are present in large parts of the mouse and human central nervous system, both during development and in adulthood, yet how their physiological behaviour is modified throughout life remains largely unknown. Moreover, the activity of adult human OPCs is still not fully understood. Significantly, most of the molecules involved in OPC-mediated remyelination are also involved in their development, a phenomenon that may be clinically relevant. In this article, we have systematically analyzed the intrinsic properties of OPCs isolated from the cerebral cortex of neonatal, postnatal and adult mice, as well as those recovered from neurosurgical adult human cerebral cortex tissue. We also analyze the response of these cells to two molecules that have known effects on OPC biology during development and that are overexpressed in individuals with Multiple Sclerosis (MS): FGF2 and anosmin-1. By analyzing intact OPCs for the first time with H-1 HR-MAS NMR spectroscopy, we show that these cells behave distinctly and that they have different metabolic patterns in function of their stage of maturity. Moreover, their response to FGF-2 and anosmin-1 differs in relation to their developmental stage and in function of the species. Our data reveal that the behaviour of adult human and mouse OPCs differs in a very dynamic way that should be considered when testing drugs and for the proper design of effective pharmacological and/or cell therapies for MS.

KEYWORDS

oligodendrocyte, myelin, Multiple Sclerosis, human, remyelination.

ABBREVIATIONS

CNS: Central nervous system

DMEM: Dulbecco's Modified Eagle's Medium

ECM: extracellular matrix

FBS: Fetal Bovine Serum

HRMAS: high-resolution magic angle spinning

MS: Multiple Sclerosis

OPC: Oligodendrocyte precursor cell

INTRODUCTION

Oligodendrocyte precursor cells (OPCs) are the sole source of oligodendrocytes, the myelin-forming cells in the central nervous system (CNS), and they are widely distributed throughout both the gray and white matter [1-5]. The vast majority of mature oligodendrocytes are born during postnatal life but in adults, oligodendrocytes can also be generated from OPCs that continue to divide and generate new-myelinating cells during adulthood [5-11]. In the adult brain, OPCs constitute 2–3% of the cells inthe gray matter and 5–8% in the white matter [1]. These cells are maintained as a resident population by self-renewal, representing the main proliferative cell type outside the neurogenic regions of the CNS [1,7,12,13]. However, the adult OPC population is not homogeneous as OPCs cycle less intensely with age, their proliferation rate is higher in the whiter matter than in the grey matter, and their differentiation rate declines progressively in postnatal and adult life [1,5,8-11,14,15]. Although the functions of adult-born oligodendrocytes remain largely unclear, they apparently maintain physiological oligodendroglial homeostasis and under pathological circumstances, they mediate long-term repair in the white matter after injury and disease, even in MS where they can partially restore the lost myelin sheaths [16,17]. In demyelinating diseases, there is an up-regulation of different molecules in the microenvironment in and around the lesions, including several factors that are important for developmental oligodendrogliogenesis, such as secreted semaphorins, growth factors, mitogens and adhesion molecules [18-24].

In the present study, we focused on cerebral cortex OPCs to avoid any variability introduced by anatomical origin or location. We examined the degree of intrinsic heterogeneity (both heterochronic and interspecies) of these cortical OPCs and we described the dynamic changes in their intrinsic properties when assessed by NMR (i.e, metabolic patterns), and how these changes affect their response to two molecules known to influence OPC biology during development and remyelination: FGF2 and anosmin-1 [19,25-33]. We systematically studied mouse cortical OPCs at different ages and OPCs isolated from the human adult cerebral cortex obtained from neurosurgery biopsies (resection margins of tumors and/or epilepsy surgery [34]). In terms of the latter, the scarcity of articles dealing with human OPCs is especially remarkable [21,34-38].

The data we obtained highlight the intrinsic physiological differences between mouse
OPCs isolated at different ages. Indeed, while OPC proliferation and migration were
enhanced similarly by FGF2, this factor produced different effects on OPC
differentiation. By contrast, anosmin-1 elicited diverse effects on differentiating, and
especially on migrating OPCs, yet not on the proliferation of these cells. While FGF2
was a chemoattractant for human and murine OPCs, anosmin-1 only attracted tumor-
related cells and it repelled non-tumor OPCs. Neither FGF2 nor anosmin-1 affected
human adult OPC differentiation. These observations confirm the chronological and
inter-species heterogeneity of OPCs, adversity that will probably be relevant when
designing effective pharmacological and/or cell therapies to repair myelin loss in MS
and other demyelinating diseases.

MATERIALS AND METHODS

Animals

CD-1 mice were obtained from (Charles River Laboratories, Inc., Wilmington, MA) and maintained in the animal facilities of the Hospital Nacional de Parapléjicos (Toledo, Spain). OPCs were obtained from different age animals: embryonic (E16); postnatal (P0, P7, P15); and adult (P60). All experimental procedures were carried out in accordance with Spanish (RD223/88) and European (2010/63/EU) guidelines, and were approved by the Animal Review Board at the Hospital Nacional de Parapléjicos (SAPA001).

Human biopsies

Biopsies of human tumor (resection of the safety margins but not the tumor itself) and non-tumor tissue (epilepsy surgery) were obtained from the Neurosurgery Service at the Hospital La Princesa (Madrid, Spain). All human samples were obtained from the adult brain cortex and the biopsies were predominantly from the temporal lobe (Table 1). They were transported at 4°C as rapidly as possible in AQIX® RS-I (AQIX Ltd., Imperial College BioIncubator) to reduce cell damage. All experiments involving human samples were carried out in accordance with the guidelines of the Research Ethics Committee of our institution, which approved our research, and all the subjects provided their written informed consent.

HOG cells

An immortalized oligodendroglioma cell line, HOG, was grown at 37ºC in an atmosphere of 5% CO_2, and in Dulbecco's Modified Eagle's Medium (DMEM, Gibco) supplemented with 10% Fetal Bovine Serum (FBS, Gibco), 100 U/mL penicillin and 100 µg/mL streptomycin (Sigma).

Oligodendrocyte culture

OPCs were isolated as described previously [19,34,38]. Briefly, the cerebral cortex of mice aged E16, P0, P7, P15 and P60 was dissected out and the meninges were carefully removed. The cortical tissue was mechanically disaggregated and it was then enzymatically dissociated in a papain solution, filtered using a 100 µm nylon mesh strainer (BD Biosciences), and seeded in polyornithine-treated flasks in DMEM

containing 10% FBS (BioWhittaker) and an antibiotic anti-mycotic solution (100 U/mL penicillin/0.1 mg/mL streptomycin and 0.25 µg/ml Amphotericin B; Sigma). The cultures were maintained at 37°C and in 5%CO_2, and the medium was changed every 4 days, adding10 ng/ml of human PDGF-AA (Millipore) to the cultures from adult mice. When the cultures reached confluence, they were shaken overnight at 250 rpm to detach the oligodendrocyte progenitors growing on the top of the confluent astrocyte monolayer. The medium was then filtered through a 40µm nylon mesh strainer (BD Biosciences) and centrifuged at 900 rpm. The cells recovered were seeded twice (45 minutes each) in bacterial grade Petri dishes (Sterilin) to remove the adherent microglia and after another round of centrifugation, the resulting pellet was resuspended and the OPCs were counted and seeded.

For cells obtained from human biopsies, the same protocol was used with certain modifications published previously [34]. This protocol allowed us to obtain sufficient human OPCs to perform the necessary assays. Briefly, rather than 75 cm^2flasks, 25 cm^2 flasks were used with a final volume of 5 ml rather than 10 ml OPC medium per flask. In addition, the medium was supplemented with 10 ng/ml of human PDGF-AA (Millipore), as in the cultures obtained from adult mice. Cultures reached confluence after 30-40 days and the flasks were shaken at 230 rpm instead of 250 rpm.

Preparation of extracts for anosmin-1

Proteins from the extracellular matrix (ECM) of CHO cells (CT) and CHO cells stably transfected with an expression plasmid carrying a C-terminal hemaglutinin-tagged (HA) version of the human KAL1 cDNA (A1 [39]) were extracted as described previously [32]. Briefly, the cells were washed once with calcium/magnesium-free Hank's Balanced Salt Solution (Gibco) and they were then incubated at 4ºC for 30 minutes in a gently rocking 10 cm diameter culture dish in 1 ml of 20 mM phosphate buffer (PB, pH 7.4), containing 350 mM NaCl and complete EDTA free protease inhibitor (Roche). The ECM proteins released into the buffer were concentrated ten-fold with Amicon Ultra-4 Ultracel-30k (Millipore Corporation) and used in the chemotaxis experiments (ECM extracts). The presence of anosmin-1 was confirmed in western blots probed with a peroxidase-conjugated rat monoclonal antibody against HA (High Affinity 3F10; Roche). Equal amounts of protein from untransfected and transfected cells were used in all the *in vitro* experiments described below.

Chemotaxis

To test chemoattraction, cells (murine and human OPCs, and HOG cells) were seeded in the upper chamber of a chemotaxis chamber (40,000 cells/transwell), and FGF2 (20ng/ml; R&D Systems) or 0.1 mg/ml of CT and A1 cell extracts (see below) were added to the lower chamber in Bottenstein-Sato (BS) medium, as reported previously [19,27,32,40]. After 20 hours at 37ºC, the cells were fixed in 4% paraformaldehyde (PFA) for 10 minutes at room temperature and the OPCs were stained with antibodies against Ganglioside GT3 (A2B5, 1:20; Developmental Studies Hybridoma Bank) and the Olig-2 oligodendroglial transcription factor (1:200, Millipore), and HOG cells were stained with Hoechst. After immunostaining, the filters were examined using an In Cell Analyzer 1000 (GE-HealthCare) and 15-20 microphotographs from each membrane were taken randomly. The In Cell Analyzer 1000 Work station software was used (GE-HealthCare) to quantify the number of transmigrated OPCs per field, expressing the data as the percentage of migrating OPCs relative to the controls±s.e.m. (Considered as 100% [32]). The data were analyzed with the Sigma Stat software package (SPSS).

Time-Lapse Imaging

For a detailed study of OPC migration, time-lapse video-microscopy (Leica, DMI 6000B) was performed to observe the behaviour of the cortical OPCs. Purified OPCs were placed on poly-L-lysine and laminin coated coverslips (10,000 cells/coverslip; Sigma) and FGF2 (20ng/ml; R&D Systems) or CT CHO or A1-expressing CHO cell extracts (0.1 mg/ml, see above) were added in BS medium. Images were taken every 20 minutes for 18 hours and the speed of cell motility was assessed with the manual tracking plugin of the ImageJ software. The values represent the percentage of the mean velocity relative to the respective controls (± s.e.m.).

Differentiation assay

Purified OPCs were placed on coverslips coated with poly-L-lysine and laminin (20,000 cells/coverslip), and FGF2 (20ng/ml; R&D Systems) or total protein from CT CHO cells or A1-expressing CHO cells (0.1 mg/ml,see above) were added in differentiation medium (BME:F12 media [1:1] supplemented with 100 μg/ml transferrin, 20 μg/ml putrescine, 12.8 ng/ml progesterone, 10.4 ng/ml selenium, 25 μg/ml insulin, 0.8 μg/ml thyroxine, 0.6% glucose, and 6.6 mM glutamine, as reported previously [34,38,41].

AraC (5µM; Sigma-Aldrich) was added to the cultures to inhibit proliferation and after

5-7days *in vitro* (DIV), the cells were fixed for immunocytochemical detection of

CNPase (1:200; Covance) and Olig-2 (1:200; Millipore). Finally, 10 random fields per

coverslip were photographed under a Leica microscope using a20X objective and the

proportion of CNPase[+] cells was assessed relative to the respective controls (± s.e.m.).

Pharmacological treatments

In order to study the mechanism underlying the activity of the FGF2/FGFR1/anosmin-1

system in specific aspects of cortical OPC biology (chemoattraction, motogenicity,

proliferation and differentiation), these processes were analyzed in the presence of the

specific FGFR inhibitor SU5402 [19,27,32,40]. The SU5402 inhibitor was reconstituted

in DMSO and the controls were administered the vehicle alone at the same

concentration to account for any negative effects of DMSO on OPCs.

Differential FGFR expression

FGFR1 expression in murine and human OPCs was assessed by dual

immunofluorescence with the OPC marker NG2 (1:200; Millipore) and an antibody

against FGFR1 (1:200; Santa Cruz Biotechnology). In order to understand the response

to FGF2 and anosmin-1 observed in OPCs from different ages, we studied the

expression of different FGFRs: FGFR1, FGFR2 and FGFR3. Cells were plated in 96-

well tissue culture plates (10,000 cells/well) and maintained at 37°C in a 5% CO_2

atmosphere and with 95% relative humidity in BS medium. After 1 DIV, the cells were

fixed for 10 min at room temperature with 4% PFA. The OPCs were immunostained

with antibodies against FGFR1 (1:200; Santa Cruz Biotechnology), FGFR2 (1:50; Santa

Cruz Biotechnology) and FGFR3 (1:20; Santa Cruz Biotechnology), and against tubulin

(1:40,000; Sigma-Aldrich). Antibody binding was detected using 680 or 800 IRDye

conjugated secondary antibodies and the plates were scanned using the Odyssey

Infrared Imaging System (LICOR). Fluorescence intensity was measured according to

the manufacturer's instructions and the amount of FGFRs relative to the amount of

tubulin was normalized to the FGFR expression in P0 samples. The values represent the

mean (± s.e.m.).

Proliferation assays

Purified OPCs were seeded on poly-L-lysine and laminin coated coverslips (20,000

cells/coverslip), and FGF2 (20ng/ml; R&D Systems) or CT CHO or A1-expressing

CHO cell extracts (0.1 mg/ml, see above) were added in BS medium. After 42 hours in

culture, a BrdU pulse (6 hours) was administered and 24 hours later, the cells were fixed

for immunocytochemical detection of BrdU and Olig-2, as described previously [27].

Finally, 10 random fields per coverslip were photographed under a Leica microscope

using a 20X objective and the data were expressed as the percentage of proliferating

OPCs relative to the controls (±s.e.m.) considered as 100%.

234 H-1 HR-MAS NMR spectroscopy of OPCs

H-1 HR-MAS NMR measurements were performed on a Bruker spectrometer operating

at 9.4 T (proton Larmor frequency of 400.14 MHz). The spectra were acquired at 4 ºC

to minimize the effect of temperature on cell stability during the acquisition time and to

reduce the variations in some amino acids [42]. Sample spinning at the magic angle was

applied at a speed of 2.4 kHz. The acquisition sequence and parameters were selected to

assess the presence of metabolites of interest and their mobility, essentially using T2-

filtered (CPMG) 1D experiments recorded with relaxation times of 2 and 60 ms. Low

power presaturation during the interscan delay of 3 s was applied for water suppression.

Typically, 256 scans were accumulated using a spectral width of 20 ppm, and the total

acquisition time per spectrum was 13 min.

The spectra were processed using MestReNova version 8.1 software (Mestrelab

Research, Santiago de Compostela, Spain) and all free induction decays were processed

with exponential multiplication (0.5 Hz line-broadening) prior to Fourier

transformation, followed by baseline correction. The chemical shifts were referenced as

follows: a small amount (5 mL) of 10 mM DSS in D_2O was added to one sample of

intact cells and referenced to DSS = 0 ppm, after which all the spectra were processed

identically and aligned (using the creatine methyl resonance at 3.026 ppm) with respect

to the sample with added DSS. Of the various methods to estimate or measure the

intensity of NMR signals [43], we compared peak intensity ratios using the methyl

signal of creatine as an internal reference. Creatine is commonly used as an internal

concentration reference for in vivo H-1 NMR because its concentration correlates with

the number of metabolically active cells and it can be used as a measure of viable cell

number [44]. Cells were washed (three times) with a deuterated PB solution,

centrifuged, and the cell pellet was frozen in liquid nitrogen and stored at -80 ºC.
Typically, approximately 60 µL of the cell pellet was placed into a 4 mm Ø zirconia
rotor for each sample.

Imaging analysis

Fluorescence digital images were obtained with a DFC480 FX digital camera (Leica)
coupled to a Leica DM5000 B microscope or using a SP5 resonant scanner (Leica
Microsystems).

Statistical Analysis

For all experimental conditions, the results were analyzed using at-test (SigmaPlot 11
software (SPSS Science, Inc.) and critical values of: *$P<0.05$, **$P<0.01$ and
***$P<0.001$.

RESULTS

Changes in the intrinsic properties of OPCs with aging

The intrinsic properties and metabolic pathways of murine OPCs isolated from the cerebral cortex were assessed at different ages (P0, P15 and P60). For the first time, we assessed these changes using H-1 high-resolution (HR)-magic angle spinning (MAS) NMR spectroscopy, which allows a large number of metabolites or small molecules in the cytoplasm to be systematically evaluated and their interaction with the local environment probed [45-47]. This approach may therefore provide valuable information on cellular physico-chemical characteristics that may be correlated with cellular processes of interest. We obtained 3 spectra for each group of OPCs, with two different echo-times TE ("T2 filters T2": 2, 10 and 60 ms), analyzing differences in signal intensities of peaks associated to the metabolites present in intact OPCs (see Table 2, Fig.1a). If we compared the proton spectra corresponding to P0, P15 and P60, for a given T_2-filter (TE), we could see that there were not large differences between ages (Fig.1b). However, at TE of 60 ms, a relative increase in most peak intensities at P15 is observed (Fig 1c-d). At long echo times (TE: 60 ms), the intensity of the peaks associated to macromolecules or molecules with low mobility decreases (i.e., molecules bound to membranes or intracellular structures). It is remarkable that one the most significant ($p < 0.05$) signal change corresponds to myo-inositol, a metabolite known to be present in cell cytoplasm. Signal intensities depend on their T_2 relaxation time, which is directly related to the molecular mobility in the cytoplasm. The higher metabolite mobility at P15 suggests a lower viscosity in the cytoplasm and low interaction between metabolites and the cytoskeleton, which as shown later, are associated with a decreased in the migratory capability and a decrease proliferation rate. These results indicate that OPCs isolated at P15 display intrinsic differences to those isolated at P0 or P60.

Since the differences in the metabolic patterns may give rise to differences in other aspects of OPC behaviour, such as proliferation and migration, it was not surprising that the rate of proliferation was significantly higher at P0 and lowest at P15 (Fig. 2a). Notably, OPCs isolated from adults (P60) proliferated at an intermediate rate (Fig. 2a) and in general, there was a significant decrease in the migratory capacities of OPCs with age (Fig. 2b). This was corroborated by video time-lapse analysis, where P15 (the slowest) and P60 OPCs migrated slower than perinatal OPCs (Fig. 2c). Moreover, P60

OPCs stopped least often during their migration (Table 2) and in total; P0 OPCs

migrated further while P15 OPCs migrated over the shortest distance (P0, 332μm; P15,

128μm; P60, 210 μm). These data highlight the heterogeneity in OPC behaviour (i.e.,

proliferation rate and migratory properties) under basal conditions, which could be

related to differences in their metabolic pattern and variation in relaxation times of

metabolites (i.e., cytoplasm viscosity). The strongest differences were detected in P15

OPCs, which may reflect the large scale myelination that is ongoing at that time [48,49].

Migrating OPCs respond dynamically to FGF2 and anosmin-1 in an age-
dependent manner

FGF-2 and anosmin-1 exert distinct chemotropic effects on rat SVZ neuroblasts, and on

rodent embryonic and postnatal OPCs, mainly acting via FGFR1 [19,27,32,33,40,50].

Hence, these molecules represent a good tool to systematically study the physiological

heterogeneity in OPC populations, avoiding the variability in structure and origin by

restricting our study to cerebral cortex OPCs. Exposure of OPCs isolated at different

stages (E16, P0, P7, P15 and P60) to FGF2 homogeneously increased the number of

transmigrating cells, confirming the previously described chemoattractive effect of this

factor (and its potential motogenic effect as well: Fig. 3a). The effect of FGF2 on

migration was significantly stronger on OPCs isolated from E16 and early postnatal

stages (P0 and P7) than on those from later stages, reflecting a gradual loss of

sensitivity with age and CNS maturation. Interestingly, anosmin-1 induced different

OPC responses depending on the age of the cells: neutral for E16 OPCs; significant

chemoattraction of OPCs isolated at P0 and P7; chemorepulsion in young adults (P15

and P60: Fig.3a). At all ages except P0, FGF2 triggered a stronger OPC response than

anosmin-1 (Fig. 3a).

To more clearly distinguish between the chemotropic and motogenic effects of FGF2

and anosmin-1 (direction or rate of migration), we followed OPCs from different aged

mice by video time-lapse microscopy. FGF2 increased the motility of OPCs at all the

stages studied and it increased the velocity of OPCs by around 20% relative to their

counterparts in control conditions (Fig. 3b). By contrast, anosmin-1 augmented the

motility of P0 OPCs, yet it clearly reduced the speed of P15 and P60 OPCs (Fig.3b).

Interestingly, P0 OPCs migrated further in the presence of either of the molecules tested

than in control conditions, and they stopped less frequently (Table 2). On the other

hand, P15 and P60 OPCs behaved similarly, migrating further and with fewer stops in

the presence of FGF2, and anosmin-1 exerted exactly the opposite effects (Fig. 3c-h;

Table 2). Together, our results suggest that both FGF2 and anosmin-1 mainly influence

the motility of OPCs (motogenic effects), consistent with previous data from our group

[19,27].

Migration of human OPCs in the presence of FGF2 and anosmin-1

As indicated above, molecules involved in the migration of OPCs during development

are up-regulated in the adult in conjunction with demyelination (including that

occurring in MS lesions in humans), suggesting the need to test the effects of these

molecules in adult OPCs. To complement this analysis, we sought to obtain information

from human adult OPCs, evaluating the effects of both anosmin-1 and FGF2on three

kinds of human OPCs in chemotaxis chambers: an oligodendroglioma cell line (HOG

cells), and on OPCs isolated from tumor tissueand non-tumor tissue obtained through

neurosurgery (the exact origin of these samples is indicated in the Methods). We first

confirmed that these three cell types expressed FGFR1, the receptor thought to mediate

the effects of FGF2 and anosmin-1 on OPC migration (Fig. 4a-f; [27,32,51-53]). It was

not surprising to find that the tumor-derived OPCs were more motile in control

conditions than non-tumoral OPCs (Table 3). However, exposure to the two factors

tested also produced differences, whereby the number of transmigrated non-tumor

OPCs significantly increased in the presence of FGF2 and was significantly lower in the

presence of anosmin-1(Fig. 4g, j-l), response similar to those of P60 mouse OPCs (Fig.

3a). In addition, both proteins were motogenic and acted as chemoattractants for tumor-

derived OPCs and HOG cells (Fig. 4h-l), as also reported for P0 murine OPCs (Fig. 3a).

It should be noted that the effect of anosmin-1 was stronger on OPCs isolated from

tumor samples than on the immortalized cell line. Video time-lapse analysis of non-

tumor isolated OPCs revealed that anosmin-1 reduced their migration velocity and the

distance they migrated, while increasing the number of stops, the opposite effect to that

observed with FGF2 for the same three parameters (Fig.5a-j, Table 3), reflecting the

contribution of pathological environment to the heterogeneity of OPC behaviour.

Together, these results from adult OPCs demonstrated that they are a heterogeneous

population of cells, in which the physiological heterogeneity depends on the species and

age. Indeed, this highlights the risk of directly extrapolating from drug testing on OPCs

isolated from neonatal/early postnatal rodent brains to the human adult scenario.

The biological effects of FGF2 and anosmin-1 on OPC proliferation and differentiation

Rather than limiting our study to cell migration, we also studied how FGF2 and anosmin-1 affected OPC proliferation and differentiation. Both these factors had a constant effect on cell proliferation and while FGF2 significantly increased the number of newly-generated OPCs at each developmental stage analyzed and in young mature OPCs, anosmin-1 had no evident effect whatsoever on cell proliferation (Supplementary Fig.1a). Regarding OPC differentiation towards myelin-forming phenotypes, more heterogeneous results were obtained. In OPCs from embryonic (E16) and young adult (P60) stages, neither of these factors altered the rate of differentiation relative to the control conditions (Fig.6a-c, j-l, m). By contrast, FGF2 and anosmin-1 exerted opposite effects on P0 cells, while both these factors drove P15 OPCs towards a myelinating phenotype (Fig.6d-f, m).

The effects of FGF2 and anosmin-1 on OPCs are FGFR-dependent

FGF2 and its receptors have been implicated in numerous processes, including cell proliferation, migration, differentiation and survival (for reviews see: [54-56]). Although we previously demonstrated that FGF2 and anosmin-1 exert their effects on OPC migration via FGFR1 [19,27,33,50,53], we analyzed the expression of the 3 FGFRs present in the oligodendroglial lineage [57,58] in our cultures of OPCs obtained from mice of different ages (E16, P0, P15 and P60). The FGFR1 was expressed by murine OPCs at every stage analyzed (Fig.7 a-l). Thus, to confirm that FGFR1 influences OPC migration, we repeated the chemotaxis assays on P0 (both anosmin-1 and FGF2 acting as chemoattractants) and P15 OPCs (anosmin-1 no effect; FGF2 chemoattractant), in the presence or absence of an FGFR signaling inhibitor (SU5402). This inhibitor abrogated both the chemoattraction and chemorepulsion provoked by FGF2 and/or anosmin-1, confirming FGFR1 as the FGF receptor involved in P0 and P15 OPC migration (Fig. 7m).

In terms of OPC differentiation, SU5402 abolished the effects of both molecules at P0, reducing the percentage of CNPase$^+$ cells to control levels, although it failed to achieve the same effect on P15 OPCs exposed to either FGF2 or anosmin-1 (Fig.7n). As FGF2 binds to every FGFR but anosmin-1 binds exclusively to FGFR1 [52,53], it suggests that other FGFRs expressed by OPCs at this stage are likely to be involved in their

differentiation [57-59]. Finally, when we analyzed the effect of SU5402 on OPC

proliferation, we noted that the increase in proliferation observed in the presence of

FGF2 was abolished with the inhibitor (Supplementary Fig. 1b).

Finally, we found a decrease in FGFR1 expression with age, which was most strongly

expressed at the embryonic stage analyzed (Fig. 7o). This gradual decline mirrors the

decrease in migration speed and chemotaxis in the presence of FGF2 and anosmin-

1with age (Fig. 1). In parallel, FGFR2 expression was weakest at P0, with no

remarkable changes at the other three stages studied (Fig. 7o). Finally, FGFR3

expression gradually increasing with age, from very weak expression at E16 to

approximately 1.5-fold stronger expression at P15 and P60 than at P0 (Fig. 7o),

consistent with previous reports [58].

DISCUSSION

Given their relative abundance in the adult CNS, their ability to move towards lesion sites and their effectiveness in remyelination, there is ever increasing interest in adult OPCs. However, this is a highly heterogeneous cell population and in practical terms, the capacity of these cells to remyelinate lesions is limited [12,19,21,60-63]. Different developmental signaling cues, including growth factors, are over expressed locally in damaged areas, and they determine the survival, proliferation, migration and correct differentiation of endogenous adult OPCs [18,19,23,24,64] (for a review see [65,66]). Here we systematically studied heterochronic OPCs isolated from the mouse cerebral cortex at E16, P0, P15 and P60, as well as OPCs isolated from samples of adult human cerebral neocortex, avoiding the interference of the potential differences derived from the different developmental origins of the cells [2].

The heterogeneity in the responses of the OPCs also varied in function of the factor to which the cells were exposed. In response to FGF-2, OPCs responded in a more consistent manner when proliferation and migration were explored than when differentiation was studied. At each developmental stage studied, as well as in OPCs isolated from the cerebral cortex of adult mice and humans, FGF2 significantly enhanced cell proliferation (consistent with previous reports [55,67,68]) and certain migratory parameters, reflecting a motogenic and chemoattractive effect as reported previously [19,27,32,57,69]. It is interesting that the effect on human OPCs isolated from the safety-margins of tumors was, in absolute numbers, stronger than that on human adult OPCs isolated from non-tumor samples or even HOG cells (of human origin). It is also true that P15 OPCs had both a lower rate of proliferation and poorer mobility than the OPCs isolated from P0 and P60 mice. Our data on OPC differentiation towards myelin-forming phenotypes substantially differed with age. As such, while FGF2 interfered with OPC differentiation of P0 OPCs (as indicated elsewhere [59,70]), it favored the differentiation of P15 OPCs and it had a neutral effect on OPCs isolated from the adult cerebral cortex, independent of the species (similar results were obtained with rat P0 OPCs –data not shown- and adult human OPCs). NMR on murine OPCs also identified different metabolic patterns, and the less viscous cytoplasm of P15 OPCs compared to that of those isolated at P0 or P60 could explain the decreased in the migratory activity and proliferation OPCs at these stages [71-73]. This novel technique

was previously used to analyze extracts of different cells types, including stem cells
[74,75], although it was not applied to intact cells as occurred here.

In contrast to FGF2, anosmin-1 had no effect on OPC proliferation and it only
potentiated OPC differentiation at early stages (P0, P15), having a neutral effect on
differentiation at adult stages. However, the effects of anosmin-1 on OPC migration
differed dramatically in function of both species and age, increasing the motility of P0
and tumor-related human OPCs (cells isolated from the safety-margins of tumors and
also the HOG cell line), while it slowed down the migration of P15 and P60/adult
OPCs, and that of OPCs from human non-tumor samples. These latter experiments
confirmed earlier data and to some extent they explain the factors that condition OPC
responses to anosmin-1 [27,32,33].

With respect to the different effects observed with human OPCs and other tumor cells,
anosmin-1 seems to promote/attract the migration of tumor cells and immortalized cell
lines ([52,76-79]; data herein). This might contribute to cancer cell dispersion and
metastasis, while anosmin-1 can contribute to the immobilization of cells in both
physiological conditions (data herein) and in non-tumoral diseases like MS [19].
Nevertheless, it was suggested elsewhere that the *Kal-1* gene would be a suppressor for
other cancer types [78,80], which merits more systematic study. Likewise, the opposing
or cooperative effects of FGF2 and anosmin-1 on tumor cells are still to be evaluated.

Undoubtedly, our study provides particularly interesting results in terms of the
similarities and differences between OPCs isolated from the murine and human adult
brain. The migratory properties of adult OPCs isolated from both species displayed
parallel responses to FGF2 and anosmin-1, which is far from similar to the responses
observed in early postnatal OPCs. Thus, our data demonstrate that OPC heterogeneity
depends not only on the anatomical region from where they are isolated and on the way
they respond to injury [21,81] but also, on the species, the stage in the life cycle and the
functional system to which they pertain. This phenomenon should be taken into account
in future studies that search for pharmacological agents to promote the motility of
human OPCs isolated from different origins (the fetal or adult brain) in order to drive
remyelination [66,82,83]. As such, and irrespective of their origin (murine or human),
adult OPCs behave distinctly to OPCs isolated at earlier stages at least in part because
their metabolic profiles are very different. In addition, their response to different cues,

such as FGF2 and anosmin-1, is modified with age. Although not as systematic as our current work, earlier data also suggest that the effects triggered by growth factors and motogenic cues (PDGF, FGF2) are more constant than those driven by chemotropic molecules and/or cytokines (semaphorins, netrin, anosmin-1, CCLs [35,65,84,85]. Hence, these data together probably reflect a hierarchy among the different molecules that influence the biology of OPCs during development and in different species.

In conclusion, the data presented here help understand the response of endogenous OPCs to lesion in the adult CNS. As such, therapeutic approaches for neurodegenerative diseases should focus on the potentiation of the endogenous OPCs present in the adult brain, enhancing their spontaneous remyelinating capacity. Accordingly, it should also be borne in mind that the modulation of FGF family members might also improve myelin repair in diseases like MS.

REFERENCES

[1] Dawson MR, Polito A, Levine JM, Reynolds R (2003) NG2-expressing glial
progenitor cells: an abundant and widespread population of cycling cells in the
adult rat CNS. Mol Cell Neurosci 24: 476-488. doi: 10.1016/S1044-
7431(03)00210-0

[2] de Castro F, Zalc B (2013) Migration of Myelin-Forming Cells in the CNS. In:
Rubenstein JLR, Rakic P, editors. Comprehensive Developmental Neuroscience:
Cellular Migration and Formation of Neuronal Connections. Elsevier,
Amsterdam, p 417-429.

[3] McKenzie IA, Ohayon D, Li H, de Faria JP, Emery B, Tohyama K, Richardson
WD (2014) Motor skill learning requires active central myelination. Science
346: 318-322. doi: 10.1126/science.1254960

[4] Richardson WD, Kessaris N, Pringle N (2006) Oligodendrocyte wars. Nat Rev
Neurosci 7: 11-18. doi: 10.1038/nrn1826

[5] Young KM, Psachoulia K, Tripathi RB, Dunn SJ, Cossell L, Attwell D,
Tohyama K, Richardson WD (2013) Oligodendrocyte dynamics in the healthy
adult CNS: evidence for myelin remodeling. Neuron 77: 873-885. doi:
10.1016/j.neuron.2013.01.006

[6] Dimou L, Simon C, Kirchhoff F, Takebayashi H, Gotz M (2008) Progeny of
Olig2-expressing progenitors in the gray and white matter of the adult mouse
cerebral cortex. J Neurosci 28: 10434-10442. doi: 10.1523/JNEUROSCI.2831-
08.2008

[7] Dimou L, Gallo V (2015) NG2-glia and their functions in the central nervous
system. Glia 63: 1429-1451. doi: 10.1002/glia.22859

[8] Kang SH, Fukaya M, Yang JK, Rothstein JD, Bergles DE (2010) NG2+ CNS
glial progenitors remain committed to the oligodendrocyte lineage in postnatal
life and following neurodegeneration. Neuron 68: 668-681. doi:
10.1016/j.neuron.2010.09.009

[9] Psachoulia K, Jamen F, Young KM, Richardson WD (2009) Cell cycle
dynamics of NG2 cells in the postnatal and ageing brain. Neuron Glia Biol 5:
57-67. doi: 10.1017/S1740925X09990354

[10] Rivers LE, Young KM, Rizzi M, Jamen F, Psachoulia K, Wade A, Kessaris N,
Richardson WD (2008) PDGFRA/NG2 glia generate myelinating
oligodendrocytes and piriform projection neurons in adult mice. Nat Neurosci
11: 1392-1401. doi: 10.1038/nn.2220

[11] Zhu Q, Whittemore SR, Devries WH, Zhao X, Kuypers NJ, Qiu M (2011)
Dorsally-derived oligodendrocytes in the spinal cord contribute to axonal
myelination during development and remyelination following focal
demyelination. Glia 59: 1612-1621. doi: 10.1002/glia.21203

[12] Hill RA, Nishiyama A (2014) NG2 cells (polydendrocytes): listeners to the
neural network with diverse properties. Glia 62: 1195-1210. doi:
10.1002/glia.22664

[13] Richardson WD, Young KM, Tripathi RB, McKenzie I (2011) NG2-glia as
multipotent neural stem cells: fact or fantasy? Neuron 70: 661-673. doi:
10.1016/j.neuron.2011.05.013

[14] Calver AR, Hall AC, Yu WP, Walsh FS, Heath JK, Betsholtz C, Richardson
WD (1998) Oligodendrocyte population dynamics and the role of PDGF in vivo.
Neuron 20: 869-882. doi: 10.1016/S0896-6273(00)80469-9

[15] Simon C, Gotz M, Dimou L (2011) Progenitors in the adult cerebral cortex: cell
cycle properties and regulation by physiological stimuli and injury. Glia 59:
869-881. doi: 10.1002/glia.21156

[16] Nishiyama A, Komitova M, Suzuki R, Zhu X (2009) Polydendrocytes (NG2
cells): multifunctional cells with lineage plasticity. Nat Rev Neurosci 10: 9-22.
doi: 10.1038/nrn2495

[17] Seo JH, Miyamoto N, Hayakawa K, Pham LD, Maki T, Ayata C, Kim KW, Lo
EH, Arai K (2013) Oligodendrocyte precursors induce early blood-brain barrier
opening after white matter injury. J Clin Invest 123: 782-786. doi:
10.1172/JCI65863

[18] Boyd A, Zhang H, Williams A (2013) Insufficient OPC migration into
demyelinated lesions is a cause of poor remyelination in MS and mouse models.
Acta Neuropathol 125: 841-859. doi: 10.1007/s00401-013-1112-y

[19] Clemente D, Ortega MC, Arenzana FJ, de Castro F (2011) FGF-2 and Anosmin-
1 are selectively expressed in different types of multiple sclerosis lesions. J
Neurosci 31: 14899-14909. doi: 10.1523/JNEUROSCI.1158-11.2011

[20] Ferent J, Zimmer C, Durbec P, Ruat M, Traiffort E (2013) Sonic Hedgehog
signaling is a positive oligodendrocyte regulator during demyelination. J
Neurosci 33: 1759-1772. doi: 10.1523/JNEUROSCI.3334-12.2013

[21] Moyon S, Dubessy AL, Aigrot MS, Trotter M, Huang JK, Dauphinot L, Potier
MC, Kerninon C, Melik PS, Franklin RJ, Lubetzki C (2015) Demyelination
causes adult CNS progenitors to revert to an immature state and express immune
cues that support their migration. J Neurosci 35: 4-20. doi:
10.1523/JNEUROSCI.0849-14.2015

[22] Ortega JA, Radonjic NV, Zecevic N (2013) Sonic hedgehog promotes
generation and maintenance of human forebrain Olig2 progenitors. Front Cell
Neurosci 7: 254. doi: 10.3389/fncel.2013.00254

[23] Piaton G, Williams A, Seilhean D, Lubetzki C (2009) Remyelination in multiple
sclerosis. Prog Brain Res 175: 453-464. doi: 10.1016/S0079-6123(09)17530-1

[24] Williams A, Piaton G, Aigrot MS, Belhadi A, Theaudin M, Petermann F,
Thomas JL, Zalc B, Lubetzki C (2007) Semaphorin 3A and 3F: key players in

myelin repair in multiple sclerosis? Brain 130: 2554-2565. doi:
10.1093/brain/awm202

[25] Díaz-Balzac CA, Lázaro-Pena MI, Ramos-Ortiz GA, Bulow HE (2015) The
Adhesion Molecule KAL-1/anosmin-1 Regulates Neurite Branching through a
SAX-7/L1CAM-EGL-15/FGFR Receptor Complex. Cell Rep 11: 1377-1384.
doi: 10.1016/j.celrep.2015.04.057

[26] Azin M, Mirnajafi-Zadeh J, Javan M (2015) Fibroblast Growth Factor-2
Enhanced The Recruitment of Progenitor Cells and Myelin Repair in
Experimental Demyelination of Rat Hippocampal Formations. Cell J 17: 540-
456.

[27] Bribián A, Barallobre MJ, Soussi-Yanicostas N, de Castro F (2006) Anosmin-1
modulates the FGF-2-dependent migration of oligodendrocyte precursors in the
developing optic nerve. Mol Cell Neurosci 33: 2-14. doi:
10.1016/j.mcn.2006.05.009

[28] Bribián A, Esteban PF, Clemente D, Soussi-Yanicostas N, Thomas JL, Zalc B,
de Castro F (2008) A novel role for anosmin-1 in the adhesion and migration of
oligodendrocyte precursors. Dev Neurobiol 68: 1503-1516. doi:
10.1002/dneu.20678

[29] Endo Y, Ishiwata-Endo H, Yamada KM (2012) Extracellular matrix protein
anosmin promotes neural crest formation and regulates FGF, BMP, and WNT
activities. Dev Cell 23: 305-316. doi: 10.1016/j.devcel.2012.07.006

[30] Hu Y, Poopalasundaram S, Graham A, Bouloux PM (2013) GnRH neuronal
migration and olfactory bulb neurite outgrowth are dependent on FGF receptor 1
signaling, specifically via the PI3K p110alpha isoform in chick embryo.
Endocrinology 154: 388-399. doi: 10.1210/en.2012-1555

[31] Mierzwa AJ, Zhou YX, Hibbits N, Vana AC, Armstrong RC (2013) FGF2 and
FGFR1 signaling regulate functional recovery following cuprizone
demyelination. Neurosci Lett 548: 280-285. doi: 10.1016/j.neulet.2013.05.010

[32] Murcia-Belmonte V, Medina-Rodríguez EM, Bribián A, de Castro F, Esteban
PF (2014) ERK1/2 signaling is essential for the chemoattraction exerted by
human FGF2 and human anosmin-1 on newborn rat and mouse OPCs via
FGFR1. Glia 62: 374-386. doi: 10.1002/glia.22609

[33] Murcia-Belmonte V, Esteban PF, Martínez-Hernández J, Gruart A, Lujan R,
Delgado-Garcia JM, de Castro F (2016) Anosmin-1 over-expression regulates
oligodendrocyte precursor cell proliferation, migration and myelin sheath
thickness. Brain Struct Funct 221: 1365-1385. doi: 10.1007/s00429-014-0977-4

[34] Medina-Rodríguez EM, Arenzana FJ, Bribián A, de Castro F (2013) Protocol to
isolate a large amount of functional oligodendrocyte precursor cells from the
cerebral cortex of adult mice and humans. PLoS One 8: e81620. doi:
10.1371/journal.pone.0081620

[35] Bin JM, Rajasekharan S, Kuhlmann T, Hanes I, Marcal N, Han D, Rodrigues
SP, Leong SY, Newcombe J, Antel JP, Kennedy TE (2013) Full-length and
fragmented netrin-1 in multiple sclerosis plaques are inhibitors of
oligodendrocyte precursor cell migration. Am J Pathol 183: 673-680. doi:
10.1016/j.ajpath.2013.06.004

[36] Cui QL, Kuhlmann T, Miron VE, Leong SY, Fang J, Gris P, Kennedy TE,
Almazan G, Antel J (2013) Oligodendrocyte progenitor cell susceptibility to
injury in multiple sclerosis. Am J Pathol 183: 516-525. doi:
10.1016/j.ajpath.2013.04.016

[37] John GR (2012) Investigation of astrocyte - oligodendrocyte interactions in
human cultures. Methods Mol Biol 814: 401-414. doi: 10.1007/978-1-61779-
452-0_27

[38] Medina-Rodríguez EM, Arenzana FJ, Pastor J, Redondo M, Palomo V, García
de Sola R, Gil C, Martínez A, Bribián A, de Castro F (2013) Inhibition of
endogenous phosphodiesterase 7 promotes oligodendrocyte precursor
differentiation and survival. Cell Mol Life Sci 70: 3449-3462. doi:
10.1007/s00018-013-1340-2

[39] Soussi-Yanicostas N, Hardelin JP, Arroyo-Jiménez MM, Ardouin O, Legouis R,
Levilliers J, Traincard F, Betton JM, Cabanie L, Petit C (1996) Initial
characterization of anosmin-1, a putative extracellular matrix protein
synthesized by definite neuronal cell populations in the central nervous system. J
Cell Sci 109 (Pt 7): 1749-1757.

[40] García-González D, Clemente D, Coelho M, Esteban PF, Soussi-Yanicostas N,
de Castro F (2010) Dynamic roles of FGF-2 and Anosmin-1 in the migration of
neuronal precursors from the subventricular zone during pre- and postnatal
development. Exp Neurol 222: 285-295. doi: 10.1016/j.expneurol.2010.01.006

[41] Casaccia-Bonnefil P, Tikoo R, Kiyokawa H, Friedrich V, Jr., Chao MV, Koff A
(1997) Oligodendrocyte precursor differentiation is perturbed in the absence of
the cyclin-dependent kinase inhibitor p27Kip1. Genes Dev 11: 2335-2346. doi:
10.1101/gad.11.18.2335

[42] Rocha CM, Barros AS, Gil AM, Goodfellow BJ, Humpfer E, Spraul M, Carreira
IM, Melo JB, Bernardo J, Gomes A, Sousa V, Carvalho L, Duarte IF (2010)
Metabolic profiling of human lung cancer tissue by 1H high resolution magic
angle spinning (HRMAS) NMR spectroscopy. J Proteome Res 9: 319-332. doi:
10.1021/pr9006574

[43] Moestue S, Sitter B, Bathen TF, Tessem MB, Gribbestad IS (2011) HR MAS
MR spectroscopy in metabolic characterization of cancer. Curr Top Med Chem
11: 2-26. doi: 10.2174/156802611793611869

[44] Govindaraju V, Young K, Maudsley AA (2000) Proton NMR chemical shifts
and coupling constants for brain metabolites. NMR Biomed 13: 129-153.

[45] Cheng LL, Lean CL, Bogdanova A, Wright SC, Jr., Ackerman JL, Brady TJ,
Garrido L (1996) Enhanced resolution of proton NMR spectra of malignant

lymph nodes using magic-angle spinning. Magn Reson Med 36: 653-658. doi:
10.1002/mrm.1910360502

[46] Griffin JL, Bollard M, Nicholson JK, Bhakoo K (2002) Spectral profiles of
cultured neuronal and glial cells derived from HRMAS (1)H NMR
spectroscopy. NMR Biomed 15: 375-384. doi: 10.1002/nbm.792

[47] Lindon JC, Beckonert OP, Holmes E, Nicholson JC (2009) High resolution
magic angle spinning NMR spectroscopy: Application to biomedical studies.
Prog. NMR Spectrosc 55: 79-100. doi: 10.1016/j.pnmrs.2008.11.004

[48] Flores AI, Narayanan SP, Morse EN, Shick HE, Yin X, Kidd G, Avila RL,
Kirschner DA, Macklin WB (2008) Constitutively active Akt induces enhanced
myelination in the CNS. J Neurosci 28: 7174-7183. doi:
10.1523/JNEUROSCI.0150-08.2008

[49] Lu Z, Ku L, Chen Y, Feng Y (2005) Developmental abnormalities of myelin
basic protein expression in fyn knock-out brain reveal a role of Fyn in
posttranscriptional regulation. J Biol Chem 280: 389-395. doi:
10.1074/jbc.M405973200

[50] Esteban PF, Murcia-Belmonte V, García-González D, de Castro F (2013) The
cysteine-rich region and the whey acidic protein domain are essential for
anosmin-1 biological functions. J Neurochem 124: 708-720. doi:
10.1111/jnc.12104

[51] González-Martínez D, Kim SH, Hu Y, Guimond S, Schofield J, Winyard P,
Vannelli GB, Turnbull J, Bouloux PM (2004) Anosmin-1 modulates fibroblast
growth factor receptor 1 signaling in human gonadotropin-releasing hormone
olfactory neuroblasts through a heparan sulfate-dependent mechanism. J
Neurosci 24: 10384-10392. doi: 10.1523/JNEUROSCI.3400-04.2004

[52] Hu Y, Guimond SE, Travers P, Cadman S, Hohenester E, Turnbull JE, Kim SH,
Bouloux PM (2009) Novel mechanisms of fibroblast growth factor receptor 1
regulation by extracellular matrix protein anosmin-1. J Biol Chem 2009; 284:
29905-29920. doi: 10.1074/jbc.M109.049155

[53] Murcia-Belmonte V, Esteban PF, García-González D, de Castro F (2010)
Biochemical dissection of Anosmin-1 interaction with FGFR1 and components
of the extracellular matrix. J Neurochem 115: 1256-1265. doi: 10.1111/j.1471-
4159.2010.07024.x

[54] Bansal R, Pfeiffer SE (1997) FGF-2 converts mature oligodendrocytes to a
novel phenotype. J Neurosci Res 50: 215-228. doi: 10.1002/(SICI)1097-
4547(19971015)50:2%3C215::AID-JNR10%3E3.3.CO;2-8

[55] Bansal R (2002) Fibroblast growth factors and their receptors in oligodendrocyte
development: implications for demyelination and remyelination. Dev Neurosci
24: 35-46. doi: 10.1159/000064944

[56] Oh LY, Yong VW (1996) Astrocytes promote process outgrowth by adult
human oligodendrocytes in vitro through interaction between bFGF and

astrocyte extracellular matrix. Glia 17: 237-253. doi: 10.1002/(SICI)1098-
1136(199607)17:3%3C237::AID-GLIA6%3E3.0.CO;2-Y

[57] Bansal R, Kumar M, Murray K, Morrison RS, Pfeiffer SE (1996) Regulation of
FGF receptors in the oligodendrocyte lineage. Mol Cell Neurosci 7: 263-275.
doi: 10.1006/mcne.1996.0020

[58] Fortin D, Rom E, Sun H, Yayon A, Bansal R (2005) Distinct fibroblast growth
factor (FGF)/FGF receptor signaling pairs initiate diverse cellular responses in
the oligodendrocyte lineage. J Neurosci 25: 7470-7479. doi:
10.1523/JNEUROSCI.2120-05.2005

[59] Furusho M, Dupree JL, Nave KA, Bansal R (2012) Fibroblast growth factor
receptor signaling in oligodendrocytes regulates myelin sheath thickness. J
Neurosci 2012; 32: 6631-6641. doi: 10.1523/JNEUROSCI.6005-11.2012

[60] Franklin RJ, Gallo V (2014) The translational biology of remyelination: past,
present, and future. Glia 2014; 62: 1905-1915. doi: 10.1002/glia.22622

[61] Maki T, Liang AC, Miyamoto N, Lo EH, Arai K (2013) Mechanisms of
oligodendrocyte regeneration from ventricular-subventricular zone-derived
progenitor cells in white matter diseases. Front Cell Neurosci 7: 275. doi:
10.3389/fncel.2013.00275

[62] Nishiyama A (2007) Polydendrocytes: NG2 cells with many roles in
development and repair of the CNS. Neuroscientist 13: 62-76. doi:
10.1177/1073858406295586

[63] Tepavcevic DK, Pekmezovic T, Stojsavljevic N, Kostic J, Basuroski ID,
Mesaros S, Drulovic J (2014) Change in quality of life and predictors of change
among patients with multiple sclerosis: a prospective cohort study. Qual Life
Res 23: 1027-1037. doi: 10.1007/s11136-013-0535-1

[64] Syed YA, Baer A, Hofer MP, González GA, Rundle J, Myrta S, Huang JK, Zhao
C, Rossner MJ, Trotter MW, Lubec G, Franklin RJ, Kotter MR (2013) Inhibition
of phosphodiesterase-4 promotes oligodendrocyte precursor cell differentiation
and enhances CNS remyelination. EMBO Mol Med 5:1918-1934. doi:
10.1002/emmm.201303123

[65] de Castro F, Bribián A, Ortega MC (2013) Regulation of oligodendrocyte
precursor migration during development, in adulthood and in pathology. Cell
Mol Life Sci 70: 4355-4368. doi: 10.1007/s00018-013-1365-6

[66] Huang JK, Franklin RJ (2011) Regenerative medicine in multiple sclerosis:
identifying pharmacological targets of adult neural stem cell differentiation.
Neurochem Int 59: 329-332. doi: 10.1016/j.neuint.2011.01.017

[67] Azim K, Raineteau O, Butt AM (2012) Intraventricular injection of FGF-2
promotes generation of oligodendrocyte-lineage cells in the postnatal and adult
forebrain. Glia 60: 1977-1990. doi: 10.1002/glia.22413

[68] Murtie JC, Zhou YX, Le TQ, Vana AC, Armstrong RC (2005) PDGF and FGF2
pathways regulate distinct oligodendrocyte lineage responses in experimental
demyelination with spontaneous remyelination. Neurobiol Dis 19: 171-182. doi:
10.1016/j.nbd.2004.12.006

[69] McKinnon RD, Smith C, Behar T, Smith T, Dubois-Dalcq M (1993) Distinct
effects of bFGF and PDGF on oligodendrocyte progenitor cells. Glia 7: 245-254.
doi: 10.1002/glia.440070308

[70] Furusho M, Roulois AJ, Franklin RJ, Bansal R (2015) Fibroblast growth factor
signaling in oligodendrocyte-lineage cells facilitates recovery of chronically
demyelinated lesions but is redundant in acute lesions. Glia 63: 1714-1728. doi:
10.1002/glia.22838

[71] Lee K, Fodor WL, Machaty Z. 2007. Dynamics of lamin A/C in porcine
embryos produced by nuclear transfer. MolReprod Dev. 74:1221-1227. doi:
10.1002/mrd.20681

[72] Maniotis AJ, Chen CS, Ingber DE (1997) Demonstration of mechanical
connections between integrins, cytoskeletal filaments, and nucleoplasm that
stabilize nuclear structure.Proc Natl AcadSci U S A. 94:849-854. doi:
10.1073/pnas.94.3.849

[73] Pollard TD, Borisy GG (2003) Cellular motility driven by assembly and
disassembly of actin filaments. Cell 112: 453-465. doi: 10.1016/S0092-
8674(03)00120-X

[74] Jansen JF, Shamblott MJ, van Zijl PC, Lehtimäki KK, Bulte JW, Gearhart JD,
Hakumäki JM (2006) Stem cell profiling by nuclear magnetic resonance
spectroscopy.MagnReson Med. 56:666-670. doi: 10.1002/mrm.20968

[75] Urenjak J, Williams SR, Gadian DG, Noble M (1993) Proton nuclear magnetic
resonance spectroscopy unambiguously identifies different neural cell types. J
Neurosci 13: 981-989.

[76] Cariboni A, Pimpinelli F, Colamarino S, Zaninetti R, Piccolella M, Rumio C,
Piva F, Rugarli EI, Maggi R (2004) The product of X-linked Kallmann's
syndrome gene (KAL1) affects the migratory activity of gonadotropin-releasing
hormone (GnRH)-producing neurons. Hum Mol Genet 13: 2781-2791. doi:
10.1093/hmg/ddh309

[77] Choy CT, Kim H, Lee JY, Williams DM, Palethorpe D, Fellows G, Wright AJ,
Laing K, Bridges LR, Howe FA, Kim SH (2014) Anosmin-1 contributes to brain
tumor malignancy through integrin signal pathways. Endocr Relat Cancer 21:
85-99. doi: 10.1530/ERC-13-0181

[78] Jian B, Nagineni CN, Meleth S, Grizzle W, Bland K, Chaudry I, Raju R (2009)
Anosmin-1 involved in neuronal cell migration is hypoxia inducible and cancer
regulated. Cell Cycle 8: 3770-3776. doi: 10.4161/cc.8.22.10066

[79] Kanda M, Shimizu D, Fujii T, Sueoka S, Tanaka Y, Ezaka K, Takami H, Tanaka
H, Hashimoto R, Iwata N, Kobayashi D, Tanaka C, Yamada S, Nakayama G,

Sugimoto H, Koike M, Fujiwara M, Kodera Y (2016) Function and diagnostic
value of Anosmin-1 in gastric cancer progression. Int J Cancer 138: 721-730.
doi: 10.1002/ijc.29803

[80] Liu J, Cao W, Chen W, Xu L, Zhang C (2015) Decreased expression of
Kallmann syndrome 1 sequence gene (KAL1) contributes to oral squamous cell
carcinoma progression and significantly correlates with poorly differentiated
grade. J Oral Pathol Med 44: 109-114. doi: 10.1111/jop.12206

[81] Nishiyama A, Suzuki R, Zhu X (2014) NG2 cells (polydendrocytes) in brain
physiology and repair. Front Neurosci 8: 133. doi: 10.3389/fnins.2014.00133

[82] Dubois-Dalcq M, Ffrench-Constant C, Franklin RJ (2005) Enhancing central
nervous system remyelination in multiple sclerosis. Neuron 48: 9-12. doi:
10.1016/j.neuron.2005.09.004

[83] Leong SY, Rao VT, Bin JM, Gris P, Sangaralingam M, Kennedy TE, Antel JP
(2014) Heterogeneity of oligodendrocyte progenitor cells in adult human brain.
Ann Clin Transl Neurol 1: 272-283. doi: 10.1002/acn3.55

[84] Mitew S, Hay CM, Peckham H, Xiao J, Koenning M, Emery B (2014)
Mechanisms regulating the development of oligodendrocytes and central
nervous system myelin. Neuroscience 276: 29-47. doi:
10.1016/j.neuroscience.2013.11.029

[85] Tepavcevic V, Kerninon C, Aigrot MS, Meppiel E, Mozafari S, rnould-Laurent
R, Ravassard P, Kennedy TE, Nait-Oumesmar B, Lubetzki C (2014) Early
netrin-1 expression impairs central nervous system remyelination. Ann Neurol
76: 252-268. doi: 10.1002/ana.24201
793

AKNOWLEDGEMENTS

We thank Miss Iris Sánchez and Mr Rafael Lebrón for their technical support, as well as

Dr José Ángel Rodríguez-Alfaro and Dr Javier Mazarío for their help with the imaging.

This work was supported by grants from the Spanish Ministerio de Economía y

Competitividad (MINECO: grant numbers SAF2009–07842, SAF2012-40023, RD07-

0060-2007; and the Red Española de Esclerosis Múltiple: RD12-0032-12) and the

''Fundación Eugenio Rodríguez Pascual'' to FdC. ABA was hired under MINECO

grants ADE10-0010 andSAF2012-40023. EMMR is the recipient of a predoctoral

fellowship from the MINECO FPI program (BES-2010-042593 associated to

SAF2009–07842). IM is contracted by SESCAM. FdC is a CSIC Staff Scientist.

    FIGURE LEGENDS

Figure 1: NMR of intact OPCs. Proton HR-MAS NMR spectra with water suppression

corresponding to a sample of OPCs (P15) illustrating the chemical shift assignment of

selected metabolites with T_2-filters of 2 ms (a) and 60 ms (c), and the intensities of the

metabolites indicated normalized with respect to creatine for T_2-filters of 2 ms (b) and

60 ms (d). An increase in the relative intensities of P15 metabolites with respect to those

of P0 and P60, for the T2-filter of 60 ms, is observed suggesting a decrease in

cytoplasm viscosity.

Figure 2: OPC proliferation and migration rates differ with age. a) Proliferation

rate of neonatal, postnatal and adult OPCs. The number of proliferating OPCs also

decreases with age. b) Transmigrated cells of different ages after 20 hours in control

conditions: E16, P0, P15 and P60. The number of transmigrated cells decreases with

age. c) Histogram represents the migration speed at P0, P15 and P60. The migration

speed is higher at P0, and it diminishes in OPCs from P15 and P60 mice. For all

experimental groups, the results were analyzed using ANOVA and a Student´s*t*-test:

*P<0.05, **P<0.01, and ***P<0.001.

Figure 3: FGF2 and anosmin-1 control OPC migration. a) Histograms showing the

proportion of OPCs relative to the controls quantified in the presence or absence of

FGF2 or anosmin-1 at different stages: E16, P0, P7, P15 and P60. ANOVA showed

anosmin-1 had no effect on migration at E16 and P7, in contrast to other stages, while

and effect of FGF2 was evident at all stages. b) Histogram from the video time lapse

analysis representing the speed of migration at P0, P15 and P60 in the presence of either

factor. c-e) Images show the manual tracking at the end of the video time lapse

recording for P0 OPCs, in control conditions (c) and in the presence of anosmin-1 (d) or

FGF-2 (e). f-h) Analysis of the total distance migrated after 18hoursin all conditions: P0

(f), P15 (g) and P60 (h).All experimental groups were analyzed with a Student´s *t*-test:

*P<0.05, **P<0.01, and ***P<0.001.

Figure 4: Effect of FGF2 and anosmin-1 on human OPC migration. Adult human

OPCs (a-c) and HOG cells (d-f) expressed FGFR1. In both types of tumor cell analyzed,

tumor cells (g) and HOG cells (h), the response to FGF2 and anosmin-1 was the same.

Non-tumor cells responded to FGF2 like tumor cells, while the response to anosmn-1 is

the opposite in these cell types (i). j-l) Images of tumor OPCs in chemotaxis chambers

stained with A2B5 (red). Scale bar represents 7.5μm for a-f and 25 μm for j-l. The
results were analyzed with a Student´s *t*-test: *P<0.05, **P<0.01, and ***P<0.001.

Figure 5: Video time lapse imaging of human OPC migration.(a-f) The manual
tracking of non-tumor human OPCs (a-c) and tumor OPCs (d-f). Non-tumor cells
migrate faster and further in the presence of FGF2 (g, i), while anosmin-1 has the
opposite effect. Anosmin-1 and FGF2 both induce faster (j) and further migration (k) of
tumor OPCs. The results were analyzed with a Student´s *t*-test: *P<0.05, **P<0.01, and
***P<0.001.

Figure 6: The effects of FGF2 and anosmin-1 on OPC differentiation. Images show
mature oligodendrocytes from E16 (a-c), P0 (d-f), P15 (g-i) and P60 (j-l) mice labeled
with CNPase (green) in control conditions and in the presence of FGF2 or anosmin-1.
m) Histogram represents the proportion of CNPase$^+$ cells relative to the controls at all
the stages analyzed. Scale bar represents 25 μm for a-l and the results were analyzed
with a Student´s *t*-test: *P<0.05, **P<0.01, and ***P<0.001.

Figure 7: FGFR expression and its influence on OPCs. a-l) FGFR1 is expressed by
OPCs at all the stages analyzed: E16 (a-c), P0 (d-f), P15 (g-i) and P60 (j-l). m)
Histogram showing the results of chemotaxis at P0 and P15 in the presence of SU5402,
which reverts the effects of anosmin-1 and FGF2. n) CNPase$^+$ cells at P0 and P15 in
cultures treated with SU5402 or the vehicle alone. No changes were observed in the
presence of SU5402. o) FGFR1, R2 and R3 expression by OPCs isolated at E16, P0,
P15 and P60. Scale bar represents 7.5μm for a-c, g-i, and 10 μm for d-f, j-l. The results
were analyzed with a Student´s *t*-test: *P<0.05, **P<0.01, and ***P<0.001.

Table 1: Summary of human samples used for in vitro experiments

Patient	Age (years)/sex	Diagnostic	Lobe	Sample weight (gr)
540	51/Female	Epilepsy	Temporal	4,9
573	24/Male	Epilepsy	Temporal	2,9
T-590	47/Female	Tumoral	Temporal	1,9
554	50/Female	Epilepsy	Temporal	2,6
574	45/Female	Epilepsy	Temporal	1,4
559	45/Female	Epilepsy	Temporal	1,1
579	50/Male	Epilepsy	Temporal	2,6
603	53/Female	Epilepsy	Temporal	1,1
549	32/Female	Epilepsy	Temporal	2
641	27/Female	Epilepsy	Temporal	1,3
T-604	37/Female	Tumoral	Temporal	0,7
363	51/Female	Epilepsy	Temporal	4
583	40/Male	Epilepsy	Temporal	0,8
T-1-SC	62/Female	Tumoral	Temporal	0,2
T-2-SC	68/Male	Tumoral	Temporal	0,6
T-3-SC	61/Female	Tumoral	Temporal	1,5
T-5-SC	51/Female	Tumoral	Temporal	3,3

Table 2. Chemical shifts of main metabolites observed in OPCs

Metabolite	δ (ppm), multiplicity, number H
myo-inositol	3.93, m
myo-inositol	3.77, m
glycine	3.54, s 2H
taurine	3.42, t, 2H
hipotaurine	3.30, t, 2H
total-choline	3.23, s, 9H
glutamine	2.44, m, 2H
glutamate	2.34, m, 2H
glutamine/glutamate/NAA	2.12
acetate	1.91, s, 3H
lipids	1.70
alanine	1.47, d, 3H
lactate and lipids (methylene)	1.31
lipids and aminoacids (methyl)	0.94, t, 3H

Table 3: Analysis of OPCs migration. The table shows the number of stops of migrating OPCs expressed as: number of stops/number of analyzed cells, in control conditions and in the presence of FGF2 and anosmin-1.

Specie and age	CONDITION		
	CT	FGF2	Anosmin-1
Mouse P0	6.2	3.5	0.85
Mouse P15	11.4	1.52	20.7
Mouse P60	3.76	2.63	8.42
Non tumoral human, adult	3.61	8.8	17.6
Tumoral human, adult	23.7	19.7	23.1

Adult human

a NG2

b FGFR1

c Merge

HOG

d CNPase

e FGFR1

f Merge

g

% A2B5$^+$/Olig2$^+$ cells relative to CT

Non tumoral cells

*

**

h

% A2B5$^+$/Olig2$^+$ cells relative to CT

Tumoral cells

**

***

i

% A2B5$^+$/Olig2$^+$ cells relative to CT

HOG cells

*

**

CT
A1
FGF2

Non tumoral cells

j A2B5 Hoechst

k A2B5 Hoechst

l A2B5 Hoechst

CT

A1

FGF2