1 The secretome of a parasite alters its host's behaviour but does not
2 recapitulate the behavioural response to infection
3
4 Running head: Secretome behavioural effects
5

Chloé Suzanne Berger[1] and Nadia Aubin-Horth[1*]

[1]Département de Biologie and Institut de Biologie Intégrative et des Systèmes (IBIS),
Université Laval, Québec, QC, Canada G1V 0A6

*corresponding author: Nadia.Aubin-Horth@bio.ulaval.ca

Key words: parasite manipulation, behaviour, *Schistocephalus solidus*, threespine
stickleback, secretome, co-evolution

ABSTRACT

Parasites with complex life cycles have been proposed to manipulate the behaviour of their intermediate hosts to increase the probability of reaching their final host. The cause of these drastic behavioural changes could be manipulation factors released by the parasite in its environment (the secretome), but this has rarely been assessed. We studied a non-cerebral parasite, the cestode *Schistocephalus solidus*, and its intermediate host, the threespine stickleback (*Gasterosteus aculeatus*), whose response to danger becomes significantly diminished when infected. These altered behaviours appear only during late infection, when the worm is ready to reproduce in its final avian host. Sympatric host-parasite pairs show higher infection success for parasites, suggesting that the secretome effects could differ for allopatric host-parasite pairs with independent evolutionary histories. We tested the effects of secretome exposure on behaviour by using secretions from the early and late infection of *S. solidus* and by injecting them in healthy sticklebacks from a sympatric and allopatric population. Contrary to our prediction, secretome from late infection worms did not result in more risky behaviours, but secretome from early infection resulted in more cautious hosts, only in fish from the allopatric population. Our results suggest that the secretome of *Schistocephalus solidus* contains molecules that can affect host behaviour, that the causes underlying the behavioural changes in infected sticklebacks are multifactorial, and that local adaptation between host-parasite pairs may extend to the response to the parasite's secretome content.

INTRODUCTION

Parasites have strong impacts on the phenotype of their hosts, including modifications of the host morphology (1), physiology (2), immunity (3) and behaviour (4). Parasites with complex life cycles have been proposed to manipulate the behaviour of their hosts, in order to increase their transmission rates to their final host (5). If parasites are manipulative, they should disturb the functioning of their host central nervous system and physiological systems (hormonal and immune systems) by the production, storage, and release of molecules called manipulation factors (6-7). Parasites release a wide variety of molecules into their external environment, including nucleic acids (8) and proteins (9), which are sometimes included in extracellular vesicles (10). These molecules are collectively referred to as the secretome. While the effects of exposing hosts to the secretome of their parasite on their immune response has been demonstrated in some host-parasite pairs (11, 12), there is currently no experimental evidence that the secretome content from a parasite directly affects behaviour in the host in the same way that the whole parasite infection does (13).

We studied the effect of the parasite secretome on host behaviour using the interaction between the threespine stickleback and the parasite *Schistocephalus solidus*. *Schistocephalus solidus* is a freshwater cestode that is trophically transmitted to three hosts (14). It enters the body cavity of its first host, a copepod, after being ingested, and then negatively affects its growth and reproduction (15). In the copepod, the expression of risky behaviours shows a pattern of "protection then facilitation" (16-17): infected copepods exhibit reduced activity in the early phase of infection, then increased activity and a changed swimming pattern, leading to higher detection and ingestion by the second host, the threespine stickleback (18). Infection in this host results in changes in morphology, physiology, immune response and behaviour (19). The worm life cycle is completed when it reaches its final avian host. (19).

The parasite's life stage within the body cavity of the threespine stickleback is

characterised by two phases with distinct phenotypes in both the parasite and its host.

During the early days of infection (the non-infective phase), the worm shows high growth

and is too small to mature if switched to its final bird host (20). The host does not show

the expected adaptive immune response (21) and no behavioural changes are known

(22). At the infective phase, the worm is ready to reproduce in its final avian host (20).

The transition from the non-infective to the infective phase is characterized by major

genome-wide reprogramming events in the worm, including the activation of genes

predicted to be involved in neural pathways and sensory perception (23). In this infective

phase, drastic behavioural changes that result in a loss of the anti-predator response

appear in sticklebacks (22). Compared to non-infected individuals, they are more

exploratory (24), less anxious (25) bolder in the presence of a predator (26), they recover

more quickly from a frightening overhead stimulus, and feed following the stimulus (26).

These behavioural perturbations could result in part from the combined effects of

modifying the serotonin pathway and the immune response in infected fish (21, 27, 28).

At the infective phase, secretome collected from *S. solidus* has been shown to modulate

the stickleback immune response *in vitro* (29). Whether this modulation of the immune

system is also involved in behavioural changes is unclear (27, 28).

Altogether, the timing and the nature of these changes in both host and parasite support

the hypothesis of a manipulation of the threespine stickleback by *Schistocephalus solidus*

to reach its final avian host. The tegument of *S. solidus* is known to include vesicles (30).

However, whether the worm secretes any molecule that can be classified as manipulation

factors that are able to alter fish behaviour is unknown (7). This system offers a unique

opportunity to functionally test if the secretome is responsible for the behavioural

changes. We predict that only exposure to secretome from worms at the late, infective

phase should induce behavioural perturbations similar to what is seen in infected

sticklebacks, and not exposure to secretome from worms at the early, non-infective

phase.

The interaction between *S. solidus* and the threespine stickleback is highly specific, since it is the only fish host that this parasite can infect (31). *S. solidus* infects freshwater sticklebacks (19) and a combination of studies of wild populations and of experimental reciprocal cross infection data suggests local adaptation between sympatric ("native") host-parasite pairs (32-34). On the other hand, marine sticklebacks are rarely exposed to *S. solidus*, as the worm does not tolerate high salinity levels when its hatches (35). Potentially reflecting the lower probability of co-evolution between these allopatric ("naïve") marine fish and *S. solidus*, infection success of experimental infections is highly variable, ranging from 1.4% to 99% depending on the population (34, 36). Marine fish infected by *S. solidus* show morphological changes (37) and the immune response of allopatric fish exposed *in vitro* to *S. solidus* antigens at the infective phase differs from the one of sympatric sticklebacks (38). However, the effects on behaviour of infecting an allopatric stickleback with *S. solidus* have never been studied.

The occurrence of threespine stickleback populations from freshwater and marine environments and their variation in infection susceptibility allow us to functionally test if *S. solidus* and its secretome co-evolved with its sympatric freshwater host to specifically modify its behaviour. Strong co-evolution between a manipulative parasite and its sympatric host was previously reported in the *Ophiocordyceps*-ant system, where *ex vivo* essays demonstrated that the fungus *Ophiocordyceps* secretes a specific array of metabolites only in presence of the host brain it has specifically evolved to manipulate (39). We had two opposite but plausible predictions according to the manipulation hypothesis: the first one was based on the observation that marine fish can be highly susceptible to *S. solidus* infection (34). If allopatric sticklebacks do not have adaptive mechanisms to fight against the secretome effects because they did not evolve with the parasite, we predicted that they would be more sensitive to the secretome than the sympatric population and would show stronger behavioural changes. The second prediction was that marine fish would activate their immune response earlier than

sympatric fish following infection, which would reduce the effects of secretome on fish
physiology and behaviour (40). In this case, we predicted that the allopatric population
would be less sensitive to the secretome from the infective phase than sympatric fish and
that they would show reduced or no behavioural changes. In both cases, we predicted
that the injection of secretome from the non-infective phase would not modify behaviour
of allopatric fish.

The objective of this study was to test if the secretome of a parasite located outside the
brain is sufficient to induce behavioural changes in its vertebrate host. To reach this
objective, we used a functional approach (Figure 1).

MATERIALS AND METHODS

Collecting secretome from the non-infective and infective phases

The collection of the secretome from *S. solidus* was performed according to a protocol
adapted from (41). We caught threespine stickleback fish from Lake Témiscouata
(Québec, 47°80' N 68°87' O), in June and July 2016 using minnow traps. One day after
fishing, fish (n=41) were euthanized with an overdose of MS-222 (75 mg/L mg/kg) and
dissected to determine if they were infected by *S. solidus*. Fish sex, size and mass, and
S. solidus mass and number in each fish were noted. Twenty-eight fish were infected by
worms, 8 at the non-infective phase (worm mass ≤ 50 mg) and 20 at the infective phase
(worm mass > 50 mg;). Following dissection, each *S. solidus* worm was washed to
remove fish blood and placed in a saline solution protected from light to collect its
secretome for 2 hours (see Supplementary material). The secretome was stored at -20°C.
For subsequent injections, we only used secretomes sampled from fish harbouring a
single worm (n=21, including 7 non-infective secretomes and 14 infective secretomes) in
order to remove potential effects of multiple infection on secretome content and function.

Sampling of sympatric and allopatric fish hosts

We performed the injections in two populations of threespine sticklebacks. The first

population, from Lake Témiscouata, Québec, was composed of fish that lived in sympatry

with the worms used to obtain secretome samples. The second population was composed

of fish from a salt march at Isle-Verte, Québec (48°00' N 69°40' O; salinity 22-26 ppt),

which thus lived in allopatry with the worms used to obtain the secretome. Fish in this

population are anadromous, meaning that they live in saltwater and migrate to the lower

salinities of the salt marsh to reproduce. Sticklebacks in this population are not expected

to be infected by *S. solidus,* because the worm does not tolerate high salinity (42).

Detailed sampling in Supplementary material.

Experimental treatments

We used four treatments for each population: an injection control treatment, an "infective

secretome" treatment, a "non-infective secretome" treatment and a "denatured infective

secretome" treatment. For these four treatments, we used healthy non-infected fish from

the two populations. In addition, in the Témiscouata population, we added a group of fish

infected by *S. solidus* that were not injected as a positive control, to validate that the

behavioural test could discriminate between healthy and *S. solidus* infected fish (n=7).

The control treatment involved fish injected with phosphate-buffered saline (PBS), to test

handling effects on fish behaviour. The three other treatments involved injections of

secretome collected in Lake Témiscouata. For each of these three treatments, we

prepared a mixture of secretome that insured sufficient volume for injections, for a total

of 3 mixtures ("infective secretome", "non-infective secretome" and "denatured infective

secretome"). For each treatment, we used one secretome mixture from 7 worms to

remove potential individual effects from a specific worm: if a worm secreted proteins with

aberrant effects on fish behaviour, then their effects would be more likely to be buffered

by the proteins secreted by the other six worms. For each treatment, the same mixture

was injected in fish from the Témiscouata and the Isle-verte populations, which allowed

to compare behavioural effects between them. Each mixture was made of 1600 µL of

secretome from each worm. The infective secretome treatment included fish injected with
secretome from worms whose weight was above 50 mg (the mass at which the worm
becomes infective (19)). Mass of the worms ranged from 80 mg to 700 mg with a mean
of 368.6 mg. The non-infective secretome treatment used fish injected with secretome
from worms whose weight was equal to or under 50 mg. Worms had a mass around 10
mg (minimal value detected by the balance) to 50 mg with a mean of 27.1 mg. This
treatment was used as a control for which behavioural changes were not expected. The
denatured secretome treatment used fish injected with secretome from worms whose
weight was above 50 mg, but for which the proteins included in the secretome were
denatured by heat at 95°C for 20 minutes before each injection. This treatment was also
added as a negative control, as proteins lose their secondary and tertiary structures
following heat treatment, and consequently their biological activities. If we found
differences in behaviours between fish injected with the intact and the denatured
secretome from large worms, this would suggest that the activity of proteins included in
the secretome is involved in the behavioural alteration. Worm mass for this last treatment
was comprised between 170 mg and 730 mg with a mean of 464.3 mg. Each treatment
was composed of n=11 fish in each population, except for the infective secretome
treatment in the Témiscouata population (n=10). We verified protein concentration and
content of the secretome mixtures using a Bradford assay and SDS-Page
(Supplementary material, supp. Fig 1).

Injection protocol

At the beginning of the protocol, we isolated 51 fish from the Témiscouata population and
44 fish from the Isle-Verte population in 2-L tanks and assigned an identification number
(ID). The first day following isolation, we verified infection by *S. solidus* using the
environmental DNA method (43). Forty-four fish from the Témiscouata population were
detected as non-infected and could be used for injections, while 7 fish were infected by
S. solidus and were used as control infected fish. As expected, none of the fish from the
allopatric population was infected by *S. solidus* and all could be used for injections.

We randomly assigned each fish to an injection treatment (PBS, "infective secretome",

"non-infective secretome" or "denatured infective secretome"). During the first week

(control week), we inserted a needle into the abdominal cavity of each fish (no liquid

injected), once a day for four days. On the 4^{th} day, we tested behaviour of each fish

between 30 and 45 minutes after manipulation. During the second week (injection week),

we injected each fish in the abdominal cavity with 100 µL of the appropriate treatment,

once a day for four days (total of 400 µL injected per fish). On the 4^{th} day, we tested

behaviour of each fish between 30 and 45 minutes after the last injection. Denatured

secretome was heated at 95°C for 20 minutes and was cooled before each injection.

Infected fish used as a positive control were exposed to the same protocol, except that

during the injection week the needle was only inserted. Fish were not anesthetized, as

manipulation took less than 1 minute and induced minor stress to fish. We fed fish daily

with brine shrimps following needle insertion or injection, except on day 4. Each fish was

given 3 days of rest between the control and the injection weeks. One fish for the infective

secretome treatment of the Témiscouata population died between the two weeks and was

removed from analysis.

Behavioural tests and analysis

The behaviours of each fish were measured in a 45 L test aquarium using standard tests

to measure exploration, thigmotaxis and boldness in threespine sticklebacks (22, 25, 27)

(see Supplementary material). Statistical analyses were performed using the R software

version 3.5.0 (44). See supplementary material for additional information on statistical

tests.

233   **RESULTS**

234   **Behaviours that differentiated infected and non-infected fish**

We measured seven behaviours in order to quantify exploration, thigmotaxis (wall-hugging tendency (25)) and boldness in infected and non-infected fish. We found that in the sympatric population, infected and non-infected fish differed significantly in two behaviours associated with boldness: latency to feed before a predator attack and time spent frozen after a predator attack. Infected fish ate two times more rapidly before a predator attack (average measured in control week: 113 sec) than non-infected fish (average measured in control week: 224 sec) (p = 0.049) (Figure 2 panel 1.A). Infected fish almost did not freeze after the predator attack (average measured in control week: 6 sec) while non-infected fish froze for a longer time (average measured in control week: 107 sec) (p = 0.009) (Figure 2 panel 1.B). For the other five behaviours, infected fish did not differ significantly from non-infected fish, although trends were apparent (Figure 2, panel 2).

Behavioural effects of secretome injections in non-infected fish from two populations of threespine sticklebacks

Sympatric population: Lake Témiscouata

We tested behaviour twice in each individual: during the control week and during the injection week. We first tested control non-infected individuals in which we injected only a saline solution, in order to quantify any changes related to the disturbance of the experiment in these two weeks. For all behaviours assessed, we did not find any significant differences between the two weeks in these control fish, suggesting that handling and injections did not significantly affect them. Similarly, infected fish did not change their behaviour between the first and second tests (Supp Tables 1 to 7).

We were not able to reproduce the typical behaviours of infected fish using secretome injections from the infective phase when injecting healthy individuals from the Témiscouata freshwater population (Supp Tables 1 to 7, Supp Figures 2 to 7). For example, in this sympatric population, none of the secretome injections had significant

effects on feeding latency before a predator attack, while it did affect fish from the

allopatric population (see below). Sympatric fish injected with the infective secretome did

not significantly change the time they took before eating between the two trials (p = 0.975,

Supp Table 4, figure 3 panel 1.C). The same was observed in fish injected with the non-

infective secretome (p = 0.852, Supp Table 4, figure 3 panel 1.D) and following injection

of denatured secretome (p = 0.526, Supp Table 4, figure 3 panel 1.E).

Allopatric population: The Isle-Verte salt marsh

In the allopatric population, we first tested control individuals in which we injected only a

saline solution, in order to quantify any changes related to the disturbance of the

experiment in these two trials. For all behaviours assessed, we did not find any significant

differences between the two weeks in these control fish, suggesting that handling and

injections did not significantly affect them (Supp Tables 8 to 14). For example, the control

non-infected fish had similar latency to eat before the predator attack in the first and

second weeks (p = 0.762, Supp Table 11) (Figure 3 panel 2.A).

We found that behaviour was changed after secretome injections in the allopatric

population in some treatments. The latency to eat before a predator attack was not

changed after injection of the infective secretome in healthy individuals (p = 0.642, Supp

Table 11, figure 3 panel 2.B). However, fish injected with the non-infective secretome

significantly increased their latency to approach food, taking more than twice longer

before resuming feeding (p < 0.001, Supp Table 11, figure 3 panel 2.C). After injection of

denatured secretome, fish significantly delayed or refrained entirely from eating before

the predator attack compared to the control week (p=0.025, Supp Table 11, figure 3 panel

2.D). We also found that allopatric fish took significantly longer to resume feeding after a

predator attack following injection of the non-infective secretome (p=0.037, Supp Table

14). After a predator attack, they stopped feeding entirely during the injection week (Supp

Figure 7 panel 2.C). This behaviour was not significantly affected in any other treatments

(Supp Table 14, supp figure 7 panel 2). Additionally, we found that allopatric fish

significantly spent more time swimming after injection of denatured secretome (p=0.016,

Supp Table 9), while other secretome treatments did not affect this behaviour (Supp

Figure 3 panel 2.D). None of the other behaviours showed changes after injections of

secretome in this population (Supp Tables 8, 10, 12 and 13 and Supp Figures 2 to 7).

297   **DISCUSSION**

It has been proposed that parasites manipulate the behaviour of their host by secreting

molecules (*i.e.* manipulation factors) that ultimately impact the physiology and the

behaviour of their host. Here, we tested if the secreted molecules of *S. solidus* could be

responsible (entirely or in part) for the behavioural changes in infected sticklebacks, by

injecting secretome in fish from a native, sympatric population and from a naïve, allopatric

one. We were not able to recreate the typical risky behaviours of infected fish in healthy

individuals using solely the secretome of *S. solidus* from the infective phase. However,

the secretome of the parasite was sufficient to significantly alter two risk-taking

behaviours, the latency to feed before and after a predator attack, in fish injected with the

secretome from early infection worms, but only in the allopatric fish population that did

not evolve in presence of the parasite.

The lack of behavioural change in fish injected with infective secretome can be interpreted

in two ways: either the secretome is not sufficient to alter behaviour, or our protocol did

not recreate the biological alterations that happen in the host during the weeks of

infection. In both populations, fish injected with the infective secretome did not differ in

behaviour from the control non-infected fish, despite the fact that proteins were detected

in the secretome. It is possible that *S. solidus* does not act on the behaviour of its host

through its secretome. If such is the case, the probability that this host behaviour alteration

is the result of active manipulation by the parasite would be low. If this is the correct

interpretation, behavioural changes would be the result of a side effect of infection, such

as the parasite mass burden, or the result of a host response (25). However, it would be
improbable to observe the significant changes in behaviour we quantified when injecting
the non-infective secretome. Moreover, as our behavioural assay could detect significant
differences in latency to feed before a predator attack between infected and non-infected
fish, we expect that we also had the statistical power to detect differences in the infective
secretome treatment.

An alternative explanation would be that our exposure protocol was not sufficient to test
our prediction. First, it is possible that we did not reach a threshold level of manipulation
factors, as it has been proposed that they may be accumulated slowly and stored by the
parasite (7). Because secretomes were sampled from the wild, the amount of secretome
collected was limited by the number of fish caught that were parasitized by infective *S.*
solidus. We thus injected a small dose every 24 hours for 4 days. In contrast, if
manipulative, *S. solidus* probably secretes molecules continuously within its host.
Increasing the dose and exposure duration should definitely be tested in order to better
understand the role, if any, of the parasitic secretome in host manipulation. Second, a
potential explanation is that *S. solidus* may secrete molecules during the non-infective
phase that would act to "prime" the host (13). One could test the importance of the
interaction between the effects of molecules secreted in the non-infective and infective
phases in inducing behavioural changes by sequentially injecting a fish with the non-
infective and infective secretomes. Third, our protocol did not include the potential
combined effects on behaviour of the secretome and of the physical deformations induced
by the worm in the fish abdominal cavity. This could be tested by simultaneously injecting
the infective secretome with silicone, which was previously used to re-create the parasitic
mass burden in non-infected fish (25). Finally, it is also possible that the infective
secretome acts in combination with molecules of the stress response, which was
previously proposed to be involved in the host's behavioural changes (27, 28). A next
step will require to adopt an integrative functional approach to simultaneously test the
effects of the secretome, of the activation of the stress response, and of the parasitic
mass burden on host behaviour.

The significant increase in latency to eat before and after a predator attack induced by the secretome from worms at the non-infective phase is similar to the altered behaviour seen in infected copepods, the first intermediate host of *S. solidus.* The copepod host exhibits a behaviour pattern of "protection then facilitation", first showing reduced activity in the early phase of infection, presumably reducing detection by predators, then an increased activity, which leads to higher detection by threespine sticklebacks (16-18). The induced behavioural perturbations in sticklebacks are surprising, as behavioural changes have not been reported in infected sticklebacks when *S. solidus* is in the early non-infective phase (22). It is possible that the same shift in behaviour exists in the fish host but has gone undetected. Using single experimental infection in 5 female sticklebacks, Barber et al. tracked changes in their antipredator behaviour for 16 weeks post-parasite exposure and reported significant behaviour changes only when the worm reached 50 mg (22). This could be explained in part by their low sample size, and testing more fish infected by worms at the non-infective stage could provide a better understanding of the potential behavioural sequence of "protection then facilitation". We can hypothesize that the secretome contributes to the worm protection at the non-infective stage, while other mechanisms (including the host stress response and the parasitic mass burden) could act independently or in combination with the infective secretome to alter behaviours that facilitate worm transfer to the final host. Also, it is important to keep in mind that secretomes were sampled from naturally infected fish, which prevented to precisely estimate the developmental stage of each worm. For example, small worms (≤ 50 mg), which we considered non-infective, might have grown more slowly because they invested more resources into molecules responsible for behavioural changes. Finally, the divergence in behavioural effects between the non-infective and the infective secretomes suggests that the secretome content between these two phases is different. In accordance with our results, the analysis of the transcriptome of *S. solidus* previously demonstrated that the genes expressed by the worm in the non-infective and infective phases were associated with contrasting biological functions, with thousands of genes differing in activity (23). However, it is also possible that behavioural differences result solely from concentration effects, if non-infective worms produced less concentrated

secretomes compared to infective worms when their secretions were collected in a saline

solution of constant volume. Analysing the nature of the protein content of the *S. solidus*

secretome at each stage would provide functional information that could shed light on the

nature of the molecular interaction between the threespine stickleback and *S. solidus*.

Studies in several host-parasite pairs from different branches of the tree of life have

shown that behavioural alteration in the host affects specific behaviours rather than

disrupting the host response to all cues (5, 7, 19, 45). Not all behaviours were altered by

the non-infective secretome in our study. The specificity of effects on cautiousness in

presence of a predator, but not on exploration of a novel and risky environment, or of

anxiety-like behaviour such as wall hugging, is in accordance with previous studies that

have shown that not all behaviours are modified in infected sticklebacks (45) and that not

all behaviours are changed when healthy sticklebacks are experimentally manipulated to

behave like infected ones (27). Indeed, manipulating serotonin levels in healthy

sticklebacks to mimic the changes observed in the brain of infected ones lead to changes

in schooling and the tendency to swim close to the surface, but not the response to a

predator attack, which was significantly altered in the opposite direction to an infected fish

in the present study (27). This result points to a probable multifactorial nature of host

altered behaviours, as no experimental manipulation leads to changes in all the

behaviours observed in *Schistocephalus* infected fish. Here, the non-infective secretome

exposure resulted in more cautious fish that took longer to feed under risk. Identifying the

content of the secretome will potentially uncover molecules that affect feeding directly or

indirectly, as found in another host-parasite system where feeding is affected (13).

Our results show that the secretome of *S. solidus* had significant effects on fish risky

behaviour only in an allopatric (naïve) population. This result is in accordance with the

hypothesis that since allopatric sticklebacks did not evolve with *S. solidus*, they do not

possess the adaptive "generalised response" against *S. solidus* and its effects on fish

physiology and behaviour, as proposed for this parasite and others (34, 46). Local

adaptation and potential co-evolution between the threespine stickleback and *S. solidus*
was previously suggested with cross-infection studies (33, 43, 63). Our results point to
the need for further studies on local adaptation in this host-parasite pair, by including
secretome exposure of fish that have evolved in the presence of another worm genotype,
i.e. sticklebacks of freshwater origin that can recognize the parasite but did not co-evolve
specifically with that genotype (34, 47). For instance, non-infected freshwater sticklebacks
from the west Coast of Canada (known to be infected by *S. solidus* (34)) could be exposed
to the secretome from Lake Temiscouata worms.

The significant increase in the latency to eat before an attack, which was reported after
the injection of the non-infective secretome in the allopatric population, was also found in
fish injected with heat-denatured secretome, suggesting that these impacts on behaviour
were not related solely to the activity of proteins. Indeed, since these secretome samples
were denatured by heat in order to lose the secondary and tertiary structures of proteins,
and consequently their biological activities, it is possible that this behavioural change
might be induced by other molecules included in the secretome that were not impacted
by the heat treatment. It is likely that *Schistocephalus* secretes and/or excretes other
molecules, such as extracellular vesicles including micro-RNAs (11). These molecules
could have their biological activities reduced in the presence of the active secretome
proteins, and heat treatment possibly revealed their actions. To confirm this hypothesis
in the future, we could replace the heat treatment of secretome by a protease exposure,
in order to specifically break down and inactivate proteins in the secretome. Heat
treatment and protease exposure could also be applied to the non-infective secretome to
better understand the role of proteins and other molecules in the reported behavioural
change.

In this study, secretome alone did not recapitulate the behavioural changes of infected
sticklebacks in non-infected sticklebacks. Yet, the effects of secretome from the non-
infective phase on boldness in the allopatric population suggest that there may be a

"protection then facilitation" effect at play in the threespine stickleback host, as seen in

the first copepod host. Our findings demonstrate that the secretome of *S. solidus* is an

important component of the molecular cross-talk between the parasite and its threespine

stickleback host.

Acknowledgements

The authors thank the personnel of the Laboratoire Aquatique de Recherche en Sciences Environnementales et Médicales at Université Laval for their help with fish rearing. We thank Christian Landry for his comments and advice during the development of this project. We thank Marie-Pier Brochu, Verônica Alves, Florent Sylvestre and Sann Delaive for comments on previous versions of this manuscript. This project was funded by an NSERC Discovery grant to NAH. The authors declare no competing or financial interests.

Author contributions

C.S.B. designed the study with input from N.A.H.. C.S.B. performed the experiments. C.S.B. analysed the data with input from N.A.H.. C.S.B. and N.AH. wrote the manuscript.

Ethics

This study was approved by the Comité de Protection des Animaux de l'Université Laval (CPAUL 2014069-3).

Data availability

The complete dataset is available as a supplementary tab-delimited .txt file. Data on seven behaviours measured during the control and treatment weeks, mass (g), size (mm) and sex of each fish are reported, as well as the infection status (no=0; yes=1), the number and the total mass of *S. solidus* (g) for individual fish in each treatment.

Electronic supplementary material

A) Supplementary tables. Results of all the statistical comparisons performed with the lmer function to determine if the secretome injections change the behaviour of non-infected fish of the Témiscouata population and of the allopatric population are presented. Results are reported in separate tables for each of the seven behaviours and for each population (supplementary tables 1 to 14). B) Supplementary figures. Seven supplementary figures are presented. C) Supplementary information on methods.

REFERENCES

1.   Sandland GJ, Goater CP. Parasite-induced variation in host morphology: Brain-
encysting trematodes in fathead minnows. *The Journal of Parasitology*. 2001**;87**(2):267-
72.

2.   Garvin MC, Szell CC, Moore FR. Blood parasites of nearctic-neotropical migrant
passerine birds during spring trans-gulf migration: Impact on host body condition. *The*
Journal of Parasitology. 2006;**92**(5):990-6.

3.   Rigano R, Profumo E, Bruschi F, Carulli G, Azzara A, Ioppolo S, et al. Modulation
of human immune response by *Echinococcus granulosus* antigen B and its possible role
in evading host defenses. *Infection and Immunity*. 2001;**69**(1):288.

4.   Watson DW, Mullens BA, Petersen JJ. Behavioral fever response of *Musca*
domestica (Diptera: Muscidae) to infection by *Entomophthora muscae* (Zygomycetes:
Entomophthorales). *Journal of Invertebrate Pathology*. 1993;**61**(1):10-6.

5.   Lafferty K, Shaw J. Comparing mechanisms of host manipulation across host and
parasite taxa. *The Journal of Experimental Biology*. 2013;**216**(1):56-66.

6.   Biron DG, Loxdale HD. Host-parasite molecular cross-talk during the manipulative
process of a host by its parasite. *The Journal of Experimental Biology*. 2013;**216**(1):148-
60.

7.   Herbison R, Lagrue C, Poulin R. The missing link in parasite manipulation of host
behaviour. *Parasites & Vectors*. 2018;**11**(1): 222.

8.   Twu O, Johnson PJ. Parasite extracellular vesicles: Mediators of intercellular
communication. *PLoS Pathogens*. 2014;**10**(8):e1004289.

9.   Grimwood J, Smith JE. *Toxoplasma gondii*: the role of parasite surface and
secreted proteins in host cell invasion. *International journal for parasitology*.
1996;**26**(2):169-73.

10.   Coakley G, Maizels RM, Buck AH. Exosomes and other extracellular vesicles: the
new communicators in parasite infections. *Trends in parasitology*. 2015;**31**(10):477-89.

11.   Harnett W. Secretory products of helminth parasites as immunomodulators.
Molecular & Biochemical Parasitology. 2014;**195**(2):130-6.

12.   Hewitson JP, Grainger JR, Maizels RM. Helminth immunoregulation: The role of
parasite secreted proteins in modulating host immunity. *Molecular & Biochemical*
Parasitology. 2009;**167**(1):1-11.

13.   Adamo SA. Turning your victim into a collaborator: Exploitation of insect
behavioural control systems by parasitic manipulators. *Current opinion in insect science*.
2019;**33**:25-29.

14.   Hammerschmidt K, Kurtz J. Chapter 5 Ecological immunology of a tapeworms'
interaction with its two consecutive hosts. *Advances in Parasitology*. 2009;**68**:111-37.

15.   Wedekind C. The infectivity, growth, and virulence of the cestode *Schistocephalus*
solidus in its first intermediate host, the copepod *Macrocyclops albidus*. *Parasitology*.
1997;**115**(3):317-24.

16.   Dianne L, Perrot-Minnot MJ, Bauer A, Gaillard M, Léger E, Rigaud T. Protection
first then facilitation: a manipulative parasite modulates the vulnerability to predation of
its intermediate host according to its own developmental stage. *Evolution: International*
Journal of Organic Evolution. 2011;**65**(9):2692-8.

17.   Wedekind C, Milinski M. Do three-spined sticklebacks avoid consuming copepods,
the first intermediate host of *Schistocephalus solidus*? An experimental analysis of
behavioural resistance. *Parasitology*. 1996;**112**(4):371-83.

18.   Weinreich F, Benesh D, Milinski M. Suppression of predation on the intermediate
host by two trophically-transmitted parasites when uninfective. *Parasitology*.
2013;**140**(1):129-35.

19.   Barber I. Sticklebacks as model hosts in ecological and evolutionary parasitology.
Trends in Parasitology. 2013;**29**(11):556-66.

20.   Tierney JF, Crompton DWT. Infectivity of plerocercoids of *Schistocephalus solidus*
(Cestoda: Ligulidae) and fecundity of the adults in an experimental definitive host, *Gallus*
gallus. *The Journal of Parasitology*. 1992;**78**(6):1049-54.

21.   Scharsack JP, Koch K, Hammerschmidt K. Who is in control of the stickleback
immune system: interactions between *Schistocephalus solidus* and its specific vertebrate
host. *Proceedings of the Royal Society B: Biological Sciences.* 2007**;274**(1629):3151-8.

22.   Barber J, Walker P, Svensson PA. Behavioural responses to simulated avian
predation in female three spined sticklebacks: The effect of experimental *Schistocephalus*
solidus infections. *Behaviour*. 2004;**141**:1425-40.

23.   Hebert FO, Grambauer S, Barber I, Landry CR, Aubin-Horth N. Major host
transitions are modulated through transcriptome-wide reprogramming events in
Schistocephalus solidus, a threespine stickleback parasite. *Molecular ecology*.
2017;**26**(4):1118-30.

24.   Arber B, Huntingford FA, Crompton DWT. The effect of hunger and cestode
parasitism on the shoaling decisions of small freshwater fish. *Journal of Fish Biology*.
1995;**47**(3):524-36.

25.   Grécias L, Valentin J, Aubin-Horth N. Testing the parasite mass burden effect on
alteration of host behaviour in the -stickleback system. *The Journal of Experimental*
Biology. 2018;**221**: jeb174748.

26.   Giles N. Behavioural effects of the parasite *Schistocephalus solidus* (Cestoda) on
an intermediate host, the three-spined stickleback, *Gasterosteus aculeatus* L. Animal
Behaviour. 1983;**31**(4):1192-4.

27.   Grécias L, Hébert F-O, Berger CS, Barber I, Aubin-Horth N. Can the behaviour of
threespine stickleback parasitized with *Schistocephalus solidus* be replicated by
manipulating host physiology? *The Journal of Experimental Biology*. 2016;**220**(2):237-46.

28.   Overli O, Pll M, Borg B, Jobling M, Winberg S. Effects of *Schistocephalus solidus*
infection on brain monoaminergic activity in female three-spined sticklebacks
Gasterosteus aculeatus. *Proceedings of the Royal Society B: Biological Sciences*.
2001;**268**(1474):1411-5.

29.   Scharsack JP, Gossens A, Franke F, Kurtz J. Excretory products of the cestode,
Schistocephalus solidus, modulate in vitro responses of leukocytes from its specific host,
the three-spined stickleback (*Gasterosteus aculeatus*). *Fish & shellfish immunology*.
2013;**35**(6):1779-87.

30.   Yoneva A, Scholz T, Kuchta R. Comparative morphology of surface ultrastructure
of diphyllobothriidean tapeworms (Cestoda: Diphyllobothriidea). *Invertebrate biology*.
2018;**137**(1):38-48.

31. Orr TSC, Hopkins CA, Charles GH. Host specificity and rejection of
Schistocephalus solidus. *Parasitology*. 1969;**59**(3):683-90.

32. Hamley M, Franke F, Kurtz J, Scharsack JP. An experimental approach to the
immuno-modulatory basis of host-parasite local adaptation in tapeworm-infected
sticklebacks. *Experimental parasitology*. 2017;**180**:119-32.

33. Kalbe M, Eizaguirre C, Scharsack JP, Jakobsen PJ. Reciprocal cross infection of
sticklebacks with the diphyllobothriidean cestode *Schistocephalus solidus* reveals
consistent population differences in parasite growth and host resistance. *Parasites &*
vectors. 2016;**9**(1):130.

34. Weber JN, Kalbe M, Shim KC, Erin NI, Steinel NC, Ma L, et al. Resist Globally,
Infect Locally: A transcontinental test of adaptation by stickleback and their tapeworm
parasite. *The American naturalist*. 2017;**189**(1):43.

35. Simmonds NE, Barber I. The effect of salinity on egg development and viability of
Schistocephalus solidus (Cestoda: Diphyllobothriidea). *Journal of Parasitology*.
2016;**102**(1):42-7.

36. Confer A, Vu V, Drevecky C, Aguirre W. Occurrence of *Schistocephalus solidus* in
anadromous threespine stickleback. *Journal of Parasitology*. 2012;**98**(3):676-8.

37. Macat Z, Bednarik A, Rulik M. The three-spined stickleback (*Gasterosteus*
aculeatus) infection with *Schistocephalus solidus* in Hel marina (Puck Bay, Baltic Sea,
Poland). *Oceanological and hydrobiological studies*. 2015;**44**(1):11-7.

38. Franke F, Rahn AK, Dittmar J, Erin N, Rieger JK, Haase D, et al. In vitro leukocyte
response of three-spined sticklebacks (*Gasterosteus aculeatus*) to helminth parasite
antigens. *Fish & shellfish immunology*. 2014;**36**(1):130-40.

39. De Bekker C, Quevillon LE, Smith PB, Fleming KR, Ghosh D, Patterson AD, et al.
Species-specific ant brain manipulation by a specialized fungal parasite. *BMC*
evolutionary biology. 2014;**14**(1):166.

40. Franceschi N, Cornet S, Bollache L, Dechaume-Moncharmont FX, Bauer A,
Motreuil S, et al. Variation between populations and local adaptation in acanthocephalan-
induced parasite manipulation. *Evolution*. 2010;**64**(8):2417-30.

41.   Cao X, Fu Z, Zhang M, Han Y, Han Q, Lu K, et al. Excretory/secretory proteome

of 14-day schistosomula, *Schistosoma japonicum*. *Journal of Proteomics*.

2016;**130**(C):221-30.

42.   Poulin R, FitzGerald GJ. Risk of parasitism and microhabitat selection in juvenile

sticklebacks. *Canadian Journal of Zoology*. 1989;**67**(1):14-8.

43.   Berger CS, Aubin-Horth N. An eDNA-qPCR assay to detect the presence of the

parasite *Schistocephalus solidus* inside its threespine stickleback host. *The Journal of*

Experimental Biology. 2018; **221**: jeb178137.

44.   RCoreTeam. R: A language and environment for statistical computing. *R*

Foundation for Statistical Computing, Vienna, Austria. 2018.

45.   Talarico M, Seifert F, Lange J, Sachser N, Kurtz J, Scharsack JP. Specific

manipulation or systemic impairment? Behavioural changes of three-spined sticklebacks

(*Gasterosteus aculeatus*) infected with the tapeworm *Schistocephalus solidus*.

Behavioral ecology and sociobiology. 2017;**71**(2):36.

46.   Rosenkranz M, Poulin R, Selbach C. Behavioural impacts of trematodes on their

snail host: Species-specific effects or generalised response? *Ethology*.

2018;**124**(11):790-5.

47.   Piecyk A, Roth O, Kalbe M. Specificity of resistance and geographic patterns of

virulence in a vertebrate host-parasite system. *BMC evolutionary biology*. 2019;**19**(1):80.

 FIGURES

Figure 1. Experimental design used to determine if the secretome of a parasite is
sufficient to induce behavioural changes in its vertebrate host. Secretomes from the
non-infective and infective phases of *S. solidus* were injected in non-infected sticklebacks
from a sympatric and an allopatric population. Only injection of the infective secretome
was predicted to induce behavioural changes. Saline (PBS), non-infective secretome, and
denatured secretome treatments were used as controls for which no behavioural changes
were expected. In the sympatric population, infected fish were also tested to validate that
the behavioural tests could discriminate between healthy and *S. solidus* infected fish.

Figure 2. Two boldness-related behaviours differ between *S. solidus*-infected and
non-infected sticklebacks. 1. Behaviours that were significantly different between
infected and non-infected sticklebacks ($p < 0.05$). A. latency to feed before a predator
attack (sec). B. time spent frozen after a predator attack (sec). **2.** Behaviours that were
non-significantly different between infected and non-infected sticklebacks. A. latency to
leave the acclimation zone (sec). B. time spent swimming (sec). C. time in the centre
(sec). D. time spent fleeing after a predator attack (sec). E. latency to feed after a predator
attack (sec). Each comparison was performed between infected fish and non-infected fish
(from the saline (PBS) group) of the control week. Black lines represent the median, the
green triangles the average, and each dot one individual.

Figure 3. Injections of non-infective secretome significantly lower the expression
of boldness-related behaviours, only in fish from the allopatric population. 1.
Latency to feed before a predator attack (sec) in the Témiscouata population. A. Infected
fish. B. Fish injected with control saline (PBS). C. Fish injected with infective secretome.
D. Fish injected with non-infective secretome. E. Fish injected with denatured secretome.
2. Latency to feed before a predator attack (sec) in the allopatric population. A. Fish

injected with control saline (PBS). B. Fish injected with infective secretome. C. Fish

injected with non-infective secretome. D. Fish injected with denatured secretome. Black

lines represent the median, the green triangles the average, each dot one individual, and

grey lines the reaction norm of one individual between the control and injection weeks.

Secretome collection:

Fish infected by *Schistocephalus solidus*
Lake Témiscouata

In saline for 2 hours

Experimental treatments:

Saline

Effects of handling?

Infective secretome
from 7 worms > 50 mg

Effects of secretome?

Non-infective secretome
from 7 worms ≤ 50 mg

Effects of size of worm?

Denatured secretome
from 7 worms > 50 mg
95°C for 20 min before injection

Effects of reducing
activity of proteins?

Behavioural tests:

Sympatric population:
Lake Témiscouata

n=11 fish per treatment
(except infective secretome n=10)
+ n=7 infected fish

Allopatric population:
The Isle-Verte salt marsh

n=11 fish per treatment

First week: Control

Needle insertion
during 4 days

Day 4: Behavioural tests

Second week: Injection

Injection (100 µL)
of the appropriate treatment
during 4 days

Day 4: Behavioural tests