The secretome of a parasite is sufficient to alter its host's behaviour

Running title: Behavioural effects of a parasite secretome

Chloé Suzanne Berger[1] and Nadia Aubin-Horth[1*]

[1]Département de Biologie and Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval, Québec, QC, Canada G1V 0A6

*corresponding author: Nadia.Aubin-Horth@bio.ulaval.ca

Key words: parasite manipulation, behaviour, *Schistocephalus solidus*, threespine stickleback, secretome, co-evolution

ABSTRACT

Parasites with complex life cycles have been proposed to manipulate the behaviour of
their intermediate hosts to increase the probability of reaching their final host. The cause
of these drastic behavioural changes could be manipulation factors released by the
parasite in its external environment (*i.e.* the secretome), but this has rarely been
assessed. Here, we studied a non-cerebral parasite, the cestode *Schistocephalus*
solidus, and its intermediate host, the threespine stickleback (*Gasterosteus aculeatus*),
whose response to danger becomes significantly diminished when infected. These
altered behaviours appear only when the worm is ready to reproduce in its final avian
host. We tested the effects of secretome exposure on behaviour by using secretions from
the early and late infection of *S. solidus* and by injecting it in healthy sticklebacks from a
sympatric and allopatric population. Contrary to our prediction, secretome from late
infection worms did not result in more risky behaviours in sticklebacks, but secretome
from early infection parasite resulted in more cautious host, only in fish from the allopatric
population. Our results suggest that the secretome of *Schistocephalus solidus* contains
molecules that can affect host behaviour, that the causes underlying the behavioural
changes in infected sticklebacks are multifactorial, and that local adaptation between
host-parasite pairs may extend to the response to the parasite's secretome content.

INTRODUCTION

Parasites have strong impacts on the phenotype of their hosts, including modifications of

the host morphology (1), physiology (2), immunity (3) and behaviour (4). Parasites with

complex life cycles have been proposed to manipulate the behaviour of their hosts, in

order to increase their transmission rates to their final host (5). For example, ants infected

by the fungus *Ophiocordyceps* leave their normal activities and climb into the vegetation,

which facilitates the propagation of the parasite's spores (6). If parasites are manipulative,

they should disturb the functioning of their host central nervous system and physiological

systems (hormonal and immune systems) by the production, storage, and release of

molecules called manipulation factors (7-9). To demonstrate that behavioural changes in

the host are the result of direct parasitic manipulation, it is crucial to find the manipulation

factors produced by the parasite and their source, and to test how they impact the

physiology and behaviour of the host. Such a link has almost never been established in

previous studies of parasitic manipulation (9). Parasites release a wide variety of

molecules into their external environment, including lipids (10), nucleic acids (11) and

proteins (12), which are sometimes included in extracellular vesicles (13). These

molecules are collectively referred to as the secretome. While the effects of exposing

hosts to the secretome of their parasite on their immune response has been

demonstrated in some host-parasite pairs (14, 15), there is currently no experimental

evidence that the secretome content from a parasite directly affects behaviour in the host

in the same way that the whole parasite infection does (16).

We studied the effect of the parasite secretome on host behaviour using the interaction

between the threespine stickleback and the parasite *Schistocephalus solidus*.

Schistocephalus solidus is a freshwater cestode that is trophically transmitted to three

hosts (17). It enters the body cavity of its first host, a copepod, after being ingested, and

then negatively affects its growth and reproduction (18). In the copepod, the expression

of risky behaviours shows a pattern of "protection then facilitation" (19): infected copepods

exhibit reduced activity in the early phase of infection, then increased activity and a

changed swimming pattern, leading to higher detection and ingestion by the second host,

the threespine stickleback (20-22). Infection in the second host results in changes in

morphology, physiology, immune response and behaviour (23, 24). The worm life cycle

is completed when it reaches its final host, usually a bird (23).

The parasite's life stage within the threespine stickleback lasts 7 to 16 weeks, is spent

within the body cavity of its host, and is characterised by two phases with distinct

phenotypes in both the parasite and its host. During the early days of infection (the non-

infective phase), the worm shows high growth and is too small to mature if switched to its

final bird host (25). In this first phase of infection, the host does not show the expected

adaptive immune response (26) and no behavioural changes are known (24, 27). At the

infective phase, the worm is ready to reproduce in its final avian host (24, 25). The

transition from the non-infective to the infective phase is characterized by major genome-

wide reprogramming events in the worm, including a shift from growth related genes to

the activation of genes predicted to be involved in neural pathways and sensory

perception (28). In this infective phase, drastic behavioural changes that result in a loss

of the anti-predator response appear in sticklebacks (24, 27). They are more exploratory

(29), less anxious (30) bolder in the presence of a predator (31), they recover more

quickly from a frightening overhead stimulus than non-infected individuals, and they feed

following the stimulus, while non-parasitized fish do not (31). These behavioural

perturbations could result from the combined effects of modifying the serotonin pathway

and the immune responses in infected fish (32, 33). The infective phase also corresponds

to the activation of the immune response in the host, when up-regulation of the respiratory

burst activity occurs (26). At this phase, secretome collected from *S. solidus* has been

shown to modulate the stickleback immune response *in vitro* (34). Whether this

modulation of the immune system is also involved in behavioural changes is unclear (32,

33).

Altogether, the timing and the nature of these changes in both host and parasite support

the hypothesis of a manipulation of the threespine stickleback by *Schistocephalus solidus*

to reach its final avian host. The tegument of *S. solidus* is known to include vesicles (35).

However, whether the worm secretes any molecule that can be classified as manipulation

factors that are able to alter fish behaviour is unknown (9, 36). This system offers a unique

opportunity to functionally test if the secretome is responsible for the behavioural

changes. We predict that only exposure to secretome from worms at the late, infective

phase should induce behavioural perturbations similar to what is seen in infected

sticklebacks, and not exposure to secretome from worms at the early, non-infective

phase.

The interaction between *S. solidus* and the threespine stickleback is highly specific, since

it is the only fish host that this parasite can infect and it dies rapidly when ingested by

other fish species (37). *S. solidus* infects freshwater sticklebacks (24, 38) and a

combination of studies of wild populations and of experimental reciprocal cross infection

data suggests local adaptation between sympatric ("native") host-parasite pairs (39-43).

For example, in distinct host-parasite population pairs, the ratio of worm to host weight,

an indicator of parasite virulence, is highly similar, suggesting an optimal virulence and

local adaptation (40). On the other hand, marine sticklebacks are rarely exposed to *S.*

solidus, as the worm does not tolerate high salinity levels when its hatches (44).

Potentially reflecting the lower probability of co-evolution between these allopatric

("naïve") marine fish and *S. solidus*, infection success of experimental infections are

highly variable, ranging from 1.4% to 99% depending on the population (43, 45). Marine

fish infected by *S. solidus* show morphological changes (46) and the immune response

of allopatric fish *in vitro* exposed to *S. solidus* antigens at the infective phase differs from

the one of sympatric sticklebacks (47). However, the effects on behaviour of infecting an

allopatric stickleback with *S. solidus* have never been studied.

The occurrence of threespine stickleback populations from freshwater and marine
environments, and their variation in infection susceptibility, allow us to functionally test if
S. solidus and its secretome co-evolved with its sympatric freshwater host to specifically
modify its behaviour. Strong co-evolution between a manipulative parasite and its
sympatric host was previously reported in the *Ophiocordyceps*-ant system, where *ex vivo*
essays demonstrated that the fungus *Ophiocordyceps* secretes a specific array of
metabolites only in presence of the host brain it has specifically evolved to manipulate
(48). We had two opposite but plausible predictions according to the manipulation
hypothesis: the first one was based on the observation that marine fish can be highly
susceptible to *S. solidus* infection (43). If allopatric sticklebacks do not have adaptive
mechanisms to fight against the effects of secretome on fish physiology and behaviour
because they did not co-evolve with the parasite (49), we predicted that they would be
more sensitive to the secretome from the infective phase than the sympatric population
and that they would show stronger behavioural changes. Conversely, if allopatric fish
have an early activation of their immune response that would reduce the effects of
secretome on fish physiology and behaviour (50), which sympatric hosts did not show
because of the coevolution arms race ongoing with its parasite, we predicted that the
allopatric population would be less sensitive to the secretome from the infective phase
than sympatric fish and that they would show reduced or no behavioural changes. For
both results, we predicted that the injection of secretome from the non-infective phase
would not modify behaviour of allopatric fish.

The objective of this study was to determine if a parasite located outside the brain can
manipulate the behaviour of its vertebrate host by releasing manipulation factors through
its secretome. We tested if the secretome could be sufficient to induce behavioural
changes in the vertebrate host, by injecting secretions from the non-infective and infective
phases of *S. solidus* in non-infected sticklebacks from a sympatric (native) population and
an allopatric (naïve) population.

MATERIALS AND METHODS

Collecting secretome from the non-infective and infective phases

The collection of the secretome from *S. solidus* was performed according to a protocol
adapted from (51). We caught threespine stickleback fish from Lake Témiscouata
(Québec, 47°80' N 68°87' O), in June and July 2016 using minnow traps. One day after
fishing, fish (n=41) were euthanized with an overdose of MS-222 (75 mg/L mg/kg) and
dissected to determine if they were infected by *S. solidus*. Fish sex, size and mass, and
S. solidus mass and number in each fish were noted. Twenty-eight fish were infected by
worms at the non-infective (worm mass ≤ 50 mg; 8 fish) and infective (worm mass > 50
mg; 20 fish) phases (see "experimental treatments" for details). Following dissection, we
put each *S. solidus* worm in a 2 mL tube of PBS (Phosphate Buffered Saline, pH 7.4 Life
Technologies) after removing fish blood by washing the worm with 1 mL of PBS when
necessary. The tube was covered with aluminium foil to protect the worm from light and
placed in a water recipient from the lake for 2 hours. We then removed the worm from the
tube. The secretome tube obtained was stored at -20°C.

Sampling of sympatric and allopatric fish hosts

We performed the injections in two populations of threespine sticklebacks. The first
population, from Lake Témiscouata, Québec, was composed of fish that lived in sympatry
with the worms used to obtain secretome samples. The second population was composed
of fish from a salt march at Isle-Verte, Québec (48°00' N 69°40' O; salinity 22-26 ppt) and
that thus lived in allopatry with the worms used to obtain the secretome. Fish in this
population are anadromous, meaning that they live in saltwater and migrate to the lower
salinities of the salt marsh to reproduce. Sticklebacks in this population are not expected
to be infected by *S. solidus* because the worm does not tolerate high levels of salinity
(52).

We caught sticklebacks from Lake Témiscouata using a seine in August 2016. Fish from

the anadromous Isle-Verte population were sampled in tide pools in July 2016 using hand

nets. Fish from both populations were juveniles at the time of capture. We brought them

to the "Laboratoire Aquatique de Recherche en Sciences Environnementales et

Médicales" at Université Laval. Sticklebacks from both populations were maintained in

separate 80 L tanks under a Light : Dark photoperiod of 12 h:12 h and a temperature of

15°C reflecting their natural environment conditions (Québec, Canada). For anadromous

fish, salinity was at 28 ppt. We fed fish daily with brine shrimps. All sticklebacks were

mature at the start of the injections.

Experimental treatments

We used four treatments for each population: an injection control treatment, an "infective

secretome" treatment, a "non-infective secretome" treatment and a "denatured infective

secretome" treatment. For these four treatments, we used healthy non-infected fish from

the two populations. In addition, in the Témiscouata population, we added a group of fish

infected by *S. solidus* that were not injected as a positive control, to validate that the

behavioural test could discriminate between healthy and *S. solidus* infected fish (n=7).

The control treatment involved fish injected with phosphate-buffered saline (PBS), to test

handling effects on fish behaviour. The three other treatments involved injections of

secretome collected in Lake Témiscouata. For each of these three treatments, we

prepared a mixture of secretome coming from 7 worms in order to remove individual

effects from a specific worm and to insure sufficient volume for injections. The mixture

was made of 1600 μL of secretome from each worm. The infective secretome treatment

included fish injected with secretome from worms whose weight was above 50 mg (the

mass at which the worm becomes infective, i.e. it can reproduce in its final host and it

induces behavioural perturbation in fish (24)). Mass of the worms ranged from 80 mg to

700 mg with a mean of 368.6 mg. The non-infective secretome treatment used fish

injected with secretome from worms whose weight was equal to or under 50 mg (the mass

at which the worm is non-infective, i.e. it does not induce behavioural changes in fish).

Worms had a mass around 10 mg (minimal value detected by the balance) to 50 mg with

a mean of 27.1 mg. This treatment was used as a control for which behavioural changes

were not expected. The denatured secretome treatment used fish injected with secretome

from worms whose weight was above 50 mg, but for which the proteins included in the

secretome were denatured by heat before each injection. This treatment was also added

as a negative control, as proteins lose their secondary and tertiary structures following

heat treatment, and consequently their biological activities. If we found differences in

behaviours between fish injected with the intact and the denatured secretome from large

worms, this would suggest that the activity of proteins included in the secretome is

involved in the behavioural alteration. Worm mass for this last treatment was comprised

between 170 mg and 730 mg with a mean of 464.3 mg. Each treatment was composed

of n=11 fish in each population, except for the infective secretome treatment in the

Témiscouata population (n=10).

Protein content of secretome

Prior to injections, we determined the protein concentration of each secretome mixture

using a Pierce Coomassie (Bradford) Protein Assay Kit (Fisher Scientific). Bovine Serum

Albumin (BSA) (2 mg/mL) was used as standard. The concentrations obtained for the

infective secretome, the non-infective secretome and the denatured secretome were

respectively 24.08 µg/mL, 6.47 µg/mL and 40.48 µg/mL. We then diluted the denatured

secretome to the concentration of the infective secretome in order to remove potential

concentration effects. We visualized proteins for each secretome mixture on a SDS-

PAGE (resolving gel 12%) using a Pierce Silver Stain Kit (Fisher Scientific). Before

running a gel, each mixture (100 µL) was precipitated with Trichloroacetic Acid (TCA) (53)

to facilitate protein visualization at low concentration. A yeast protein extract was used as

a positive control and a Blue pre-stained protein standard (BioLabs) was added on the

gel. A negative control including the SDS-PAGE loading buffer and water was used.

Migration was performed during 60 min at 175 V followed by silver staining. For the three

mixtures, proteins of various sizes were effectively separated on gel (Supp Figure 1).

Injection protocol

At the beginning of the protocol, we isolated 51 fish from the Témiscouata population and 44 fish from the Isle-Verte population in 2 L tanks and assigned an identification number (ID). We applied the environmental DNA method developed to non-invasively detect the infection by *S. solidus* (54). On 2 consecutive days following isolation, we injected 100 µL of PBS in the abdominal cavity of an individual and we immediately aspirated it back into the syringe, to sample the eDNA present in the abdominal cavity fluids of each fish. Forty-four fish from the Témiscouata population were detected as non-infected and could be used for injections, while 7 fish were infected by *S. solidus* and were used as controlled infected, non-injected fish. As expected, none of the fish from the anadromous population was infected by *S. solidus* and all could be used for injections.

Following the environmental DNA injection protocol, we randomly assigned each fish to an injection treatment (PBS, "infective secretome", "non-infective secretome" or "denatured infective secretome"). During the first week (control week), we inserted a needle into the abdominal cavity of each fish (no injection), once a day for four days. On 4th day, we tested behaviour of each fish between 30 and 45 minutes after the last needle insertion. During the second week (injection week), we injected each fish in the abdominal cavity with 100 µL of the appropriate treatment (PBS, "infective secretome", "non-infective secretome" or "denatured secretome"), once a day for four days (total of 400 µL injected per fish). On the 4th day, we tested behaviour of each fish between 30 and 45 minutes after the last injection. Denatured secretome was heated at 95°C for 20 minutes and was allowed to be cooled before each injection. Infected fish used as a positive control were exposed to the same protocol, except that during the injection week the needle was only inserted as during the control week. We fed fish daily with brine shrimps following needle insertion or injection, except on day 4. Each fish was given 3 days of rest between the control week and the injection week. One fish for the infective secretome treatment of the

Témiscouata population died between the control week and the injection week and was
not taken into account in the analysis.

Behavioural tests and analysis

We performed behavioural tests in an aquarium (30 Width x 30 Depth x 60 Length cm)
inside which a white box (25 W x 36 D x 57 L cm) was inserted in order to prevent fish
reflections on the glass of the aquarium and to optimize tracking during analyses. We
filled the box with 25 cm of water depth and water was changed between each fish. The
box was opened on the top to allow recording by a camera. The day before the
behavioural test of a fish, we enclosed the 2L individual tank with a small white box (14
W x 17 D x 30 L cm) opened on top, to acclimatize the fish to a white environment. From
this point the fish was not fed. The day of the test, at the start of the experiment, we put
the fish in the acclimation zone inside the white box of the aquarium for 600 seconds. The
acclimation zone, which was delimited by a white wall, represented one third of the white
box. After 600 seconds, the wall was removed, and 3 behaviours were measured.

<u>Exploration</u>

Exploration of a novel environment was measured using 2 variables: 1) Latency to leave
the acclimation zone (sec). As soon as the wall was removed, the fish was given 600
seconds to leave the acclimation zone. If the fish never left the acclimation zone, it was
given the worst score for exploration (*i.e.* 600 sec). 2) Time spent swimming (sec). As
soon as the fish left the acclimation zone (or after 600 sec), time spent swimming by the
fish was recorded for 300 seconds.

<u>Thigmotaxis</u>

Wall hugging was measured with 1 variable according to a previously developed protocol:
time spent in the centre of the aquarium (sec) (30). We quantified thigmotaxis as soon as
the fish left the acclimation zone (or after 600 sec) and measured for 300 seconds. The
centre of the aquarium was defined as a zone at more than 5 cm of the edge of the white
box.

<u>Boldness</u>

The inclination to take risks was measured using 4 variables: 1) Latency to feed before a
predator attack (sec). We deposited brine shrimps with a pipette in front of the fish so that
it could see it. The fish was given 300 seconds to feed. 2) Time spent fleeing after a
predator attack (sec). 3) Time spent frozen after a predator attack (sec). 4) Latency to
feed after a predator attack (sec). For these last three variables, as soon as the fish fed
after food deposition, or 300 seconds after food deposition, we simulated a predator
attack using a heron head model (beak of the model was going back and forth into the
water during 5 seconds near the fish head). Time spent fleeing corresponds to the time
during which the fish escapes just after the heron head model was removed. Time spent
frozen indicates the time during which the fish stays motionless (*i.e.* does not swim a
distance greater than its body length) just after the heron head model was removed.
Latency to feed after an attack was also measured just after the heron head model was
removed. These three variables were determined for 300 seconds.

Following the behavioural tests, we euthanized fish on day 4 of the injection week with an
overdose of MS-222 (75 mg/L) and dissected them to confirm that they were infected by
S. solidus or not. Fish sex, size and mass, and *S. solidus* mass and number in each fish
were noted.

Two behaviours were automatically analysed using a tracking module with the Ethovision
software (Ethovision XT 11.5, Noldus Information Technology) (55): time spent swimming
and time spent in the centre of the aquarium. All the other behaviours were analysed
manually. All the videos were analysed blinded by C.S.B.

Statistical analyses

Statistical analyses were performed using the R software version 3.5.0 (56). We used the
package lme4 (57) to create a linear mixed model for each population separately including
the week (control or injection), the treatment (PBS, "infective secretome", "non-infective
secretome", "denatured secretome" or infected) and their interaction (week*treatment),
the mass, the size and the sex of the fish as fixed effects and the ID of the fish as random
effect. For each analysis, we checked homoscedasticity by plotting residuals and fitted
data. We checked for homogeneity of variances using a Levene Test. We tested normality
of residuals using graphical assessment (QQ-plot and normal distribution histogram) and
a Shapiro-Wilk test. When the data were not normally distributed, we used the Boxcox
test from the package MASS (58) to find the best transformation to fit normality (log,
square root, square or no transformation). To obtain p-values, we performed pairwise
comparisons between weeks and treatments using the package lmerTest (59). In the
sympatric Témiscouata population, the first analysis was performed with two reference
treatments (infected and PBS) to confirm that there was no difference in behaviour before
(PBS-control week) and after (PBS-injection week) control injections, and to determine if
infected fish (infected-control week) and non-infected fish (PBS-control week) showed
significantly different behaviours as expected. The second analysis aimed to test our main
predictions and was performed with the secretome treatments only ("infective secretome",
'non-infective secretome" and "denatured secretome") to determine if the secretome
injections were able to reproduce the behaviours of infected fish, using each individual as
their own control. In the allopatric anadromous population, fish were not infected by *S.*
solidus. The first analysis was thereby performed with all reference (PBS) and secretome
treatments, to test if PBS injections changed behaviours. To test our predictions about

the effects of secretome on behaviour, we performed a second analysis using fish from
the three secretome treatments ("infective secretome", "non-infective secretome" and
"denatured secretome"), using each individual as their own control.

RESULTS

Behaviours that differentiated infected and non-infected fish

We measured seven behaviours in order to quantify exploration, thigmotaxis and
boldness in infected and non-infected fish. We found that in the sympatric population,
infected and non-infected fish differed significantly in two behaviours associated with
boldness: latency to feed before a predator attack and time spent frozen after a predator
attack. Infected fish ate two times more rapidly before a predator attack (average for
infected fish, measured in control week: 113 sec) than non-infected fish (average for non-
infected fish, measured in control week: 224 sec) ($p = 0.049$) (Figure 1 panel 1.A).
Infected fish almost did not freeze after the predator attack (average measured in control
week: 6 sec) while non-infected fish froze for a longer time (average measured in control
week: 107 sec) ($p = 0.009$) (Figure 1 panel 1.B). For the other five behaviours, infected
fish did not differ significantly from non-infected fish, although trends were apparent.
Infected fish took 12 seconds on average to leave the acclimation zone, while non-
infected fish took 98 seconds ($p = 0.958$) (Figure 1 panel 2.A) and spent 30% more time
swimming (average measured in control week: 152 sec) than non-infected fish (average
measured in control week: 116 sec) ($p = 0.596$) (Figure 1 panel 2.B). On average, infected
fish spent 57% more time in the centre of the aquarium compared to non-infected fish
(average of 179 sec versus 114 sec) ($p = 0.998$) (Figure 1 panel 2.C). Infected fish spent
6.1 seconds fleeing on average after the predator attack, while non-infected fish spent
3.6 seconds ($p = 0.543$) (Figure 1 panel 2.D). Infected fish took 176 seconds before
resuming feeding on average after the predator attack, while non-infected fish mostly
forewent eating, with an average feeding latency close to the maximum given time of 300
seconds (average: 274 sec) ($p = 0.214$) (Figure 1 panel 2.E). To confirm these results,
we performed additional statistical analyses to validate that our model was not overfitted

and that the obtained p-values were accurate. We found that mass of fish was correlated
with behaviour, but p-values were not changed when removing this effect from the model.

Behavioural effects of secretome injections in non-infected fish from two
populations of threespine sticklebacks

Sympatric population: Lake Témiscouata

We tested behaviour twice in each individual: during the control week and during the
injection week. We first tested control non-infected individuals in which we injected only
a saline solution, in order to quantify any changes related to the disturbance of the
experiment in these two trials. For all behaviours assessed, we did not find any significant
differences between the two weeks in these control fish, suggesting that handling did not
significantly affect them (Supp Tables 1 to 7).For the latency to feed before a predator
attack, infected and healthy individuals differed in their behaviour on both the control (p =
0.049) and injection week (p<0.001), as stated above (Supp Table 4). During the first
week, infected fish took 113 seconds on average before eating, and started eating almost
twice as fast on the second week (average: 60 sec), although there were no statistically
significant differences between these two tests (average between weeks, p = 0.619, Supp
Table 4) (Figure 2 panel 1.A). Non-infected fish showed much greater latency to eat than
infected fish and did not change their behaviour between the first and second tests
(average: 224 sec vs 256 sec) (average between weeks, p = 0.781, Supp Table 4) (Figure
2 panel 1.B).

We were not able to reproduce the typical behaviours of infected fish using secretome
injections from the infective phase when injecting healthy individuals from the
Témiscouata freshwater population (Supp Tables 1 to 7). For example, in this sympatric
population, none of the secretome injections had significant effects on feeding latency
before a predator attack. Fish injected with the infective secretome did not significantly
change the time they took before eating between the two trials. Their average latency in

the control week was 191 seconds, then 222 seconds in the injection week (p = 0.975,

Supp Table 4) (Figure 2 panel 1.C). The same was observed in fish injected with the non-

infective secretome (average: 154 sec vs 201 sec) (p = 0.852, Supp Table 4) (Figure 2

panel 1.D) and following injection of denatured secretome (average: 206 sec vs 276 sec)

(p = 0.526, Supp Table 4) (Figure 2 panel 1.E). None of the other behaviours showed

changes after injections in the Témiscouata population (Supp Figures 2 to 7).

<u>Allopatric population: The Isle-Verte salt marsh</u>

In the allopatric population, we first tested control individuals in which we injected only a

saline solution, in order to quantify any changes related to the disturbance of the

experiment in these two trials. For all behaviours assessed, we did not find any significant

differences between the two weeks in these control fish, suggesting that handling did not

significantly affect them (Supp Tables 8 to 14). For example, the control non-infected fish

had similar latency to eat before the predator attack in the first and second weeks

(average: 194 sec vs 243 sec) (p = 0.762, Supp Table 11) (Figure 2 panel 2.A).

We found that behaviour was changed after secretome injections in the allopatric

population in some treatments. The latency to eat before a predator attack was not

changed after injection of the infective secretome in healthy individuals. Their latency to

feed on the control week was 145 seconds and 189 seconds after the secretome injection

(p = 0.642, Supp Table 11) (Figure 2 panel 2.B). However, fish injected with the non-

infective secretome significantly increased their latency to approach food, taking more

than twice longer before resuming feeding (average: 101 sec vs 269 sec) (p < 0.001,

Supp Table 11) (Figure 2 panel 2.C). After injection of denatured secretome, fish

significantly delayed or refrained entirely from eating before the predator attack compared

to the control week (average: 165 sec vs 281 sec) (p=0.025, Supp Table 11) (Figure 2

panel 2.D). We also found that anadromous fish took significantly longer to resume

feeding after a predator attack following injection of the non-infective secretome (p=0.037,

Supp Table 14). After a predator attack, their latency to feed was 227 seconds during the
control week, while they stopped feeding entirely during the injection week, with an
average feeding latency of 300 seconds, the maximum possible value (Supp Figure 7
panel 2.C). This behaviour was not significantly affected in the two other secretome
treatments nor in the control fish injected with PBS (Supp Table 14). Additionally, we
found that anadromous fish significantly spent more time swimming after injection of
denatured secretome (average: 128 sec vs 176 sec) (p=0.016, Supp Table 9) while other
secretome treatments did not affect this behaviour (Supp Figure 3 panel 2.D). None of
the other behaviours showed changes after injections of secretome in this population
(Supp Tables 8, 10, 12 and 13 and Supp Figures 2 to 7).

DISCUSSION

It has been proposed that parasites manipulate the behaviour of their host by secreting
molecules (*i.e.* manipulation factors) that ultimately impact the physiology and the
behaviour of their host. Here, we tested if the secreted molecules of *S. solidus* could be
responsible (entirely or in part) for the behavioural changes in infected sticklebacks, by
injecting secretome in fish from a native, sympatric population and from a naïve, allopatric
one. We were not able to recreate the typical risky behaviours of infected fish in healthy
individuals using solely the secretome of *S. solidus* from the infective phase. However,
the secretome of the parasite was sufficient to significantly alter two risk-taking
behaviours, the latency to feed before and after a predator attack, in fish injected with the
secretome from early infection worms, but only in the allopatric fish population that did
not evolve in presence of the parasite.

The lack of behavioural change in fish injected with infective secretome can be interpreted
in two ways: either the secretome is not sufficient to alter behaviour, or our protocol did
not recreate the biological alterations that happen in the host during the weeks of
infection. In both population, fish injected with the infective secretome did not differ in

behaviour from the control non-infected fish that were injected with PBS, despite the fact
that proteins were detected in the secretome. Clearly, it is possible that *S. solidus* does
not act on the behaviour of its host through its secretome, and if such is the case, the
probability that this host behaviour alteration is the result of active manipulation by the
parasite would be low. If this is the correct interpretation, behavioural changes would be
only the result of a side effect of infection, such as the effect of the parasite mass burden,
or the result of a host response (30). However, if this was the case, it would be improbable
to observe the significant changes in behaviour we quantified when injecting the non-
infective secretome. An alternative explanation would be that our exposure protocol was
not sufficient to test our prediction. First, it is possible that we did not reach a threshold
level of manipulation factors, as it has been proposed that they may be accumulated
slowly and stored by the parasite (9, 60). Because secretomes were sampled from the
wild, the amount of secretome collected, and thus available for injection, was limited by
the number of caught fish that were effectively infected by *S. solidus* at the infective
phase. We consequently performed injections for 4 days, with a small dose injected every
24h. In contrast, if manipulative, *S. solidus* probably secretes molecules continuously in
the abdominal cavity of the threespine stickleback. Increasing the dose and the duration
of exposure in non-infected fish should definitely be tested in order to better understand
the role, if any, of the parasitic secretome in host manipulation. Second, a potential
explanation is that *S. solidus* may secrete molecules during the non-infective phase,
which lasts several weeks, that would act to "prime" the host (16). The importance of the
interaction between the effects of molecules of the non-infective phase and of the infective
phase to induce behavioural changes should be tested by sequentially injecting the non-
infective secretome and the infective secretome in non-infected fish. Finally, it is also
possible that the molecules of the secretome at the infective phase act in combination
with molecules of the stress response, which was previously proposed to be involved in
the behavioural changes of the threespine stickleback (32, 33), to induce the complex
behavioural changes. A next step will require to adopt an integrative functional approach
to simultaneously test the effects of the secretome and of the activation of the stress
response on the behaviour of the host.

The significant increase in the fish latency to eat before and after a predator attack induced by the secretome from worms at the non-infective phase is similar to the altered behaviour seen in infected copepods, the first intermediate host of *S. solidus.* The copepod host exhibits a behaviour pattern of "protection then facilitation", where the copepod first shows reduced activity in the early phase of infection by *S. solidus*, presumably reducing detection by predators, then an increased activity, which leads to higher detection by the threespine stickleback (19-22). The induced behavioural perturbations in sticklebacks are surprising, as behavioural changes have not been reported in infected sticklebacks when *S. solidus* is in the early non-infective phase (24, 27). It is possible that the same shift in behaviour exists in the fish host but has gone undetected. Studying the possible behavioural sequence of "protection then facilitation" in threespine sticklebacks infected by *S. solidus* thus represents a promising opportunity to better understand behavioural manipulation. Furthermore, the divergence in behavioural effects between the non-infective and the infective secretome suggests that the secretome content between these two phases is different. In accordance with our results, the analysis of the transcriptome of *S. solidus* previously demonstrated that the genes expressed by the worm are significantly different between non-infective and infective phases, with thousands of genes differing in activity (28). Prior to injections, proteins were detected in secretomes from both phases. Analysing the nature of the protein content of the *S. solidus* secretome would provide functional information that could shed light on the nature of the molecular interaction between the threespine stickleback and *S. solidus*.

Studies in several host-parasite pairs from different branches of the tree of life have shown that behavioural alteration in the host affects specific behaviours rather than disrupting the host response to all cues. (5, 7, 23, 36, 61). Not all behaviours were altered by the non-infective secretome in our study. The specificity of effects on cautiousness in presence of a predator, but not on exploration of a novel and risky environment, or of anxiety-like behaviour such as wall hugging, is in accordance with previous studies that have shown that not all behaviours are modified in infected sticklebacks (61) and that not

all behaviours are changed when healthy sticklebacks are experimentally manipulated to
behave like infected ones (32). Indeed, manipulating serotonin levels in healthy
sticklebacks to mimic the changes observed in the brain of infected ones lead to changes
in schooling and the tendency to swim close to the surface, but not the response to a
predator attack, which was significantly altered (in the opposite direction to an infected
fish) in the present study (32). This result points to a probable multifactorial nature of host
altered behaviours, as no experimental manipulation leads to changes in all the
behaviours observed in *Schistocephalus* infected fish.Here, the non-infective secretome
exposure resulted in more cautious fish that took longer to feed under risk (or refrained
for feeding). Identifying the content of the secretome will potentially uncover molecules
that affect directly or indirectly feeding, as found in another host-parasite system where
feeding is affected (16).

Our results show that the secretome of *S. solidus* had significant effects on fish risky
behaviour only in an allopatric (naïve) population and that allopatric fish are therefore
more sensitive to the secretome. These results are in accordance with the hypothesis
that since allopatric sticklebacks did not co-evolve with *S. solidus*, they did not develop
the adaptive "generalised response" against *S. solidus* and its effects on fish physiology
and behaviour, as proposed for this parasite and others (43, 49, 62, 63). Local adaptation
and potential co-evolution between the threespine stickleback and *S. solidus* was
previously suggested with cross-infection studies (40, 43, 64). We provide here some new
insights that point to the need for further studies that include exposure of fish that have
evolved in the presence of another worm genotype, *i.e.* sticklebacks of freshwater origin
that can recognize the parasite but did not co-evolve specifically with that genotype, to
reveal further information on local adaptation in this host-parasite pair (43, 64).

The significant increase in the latency to eat before an attack, which was reported after
the injection of the non-infective secretome in the allopatric population, was also found in
fish injected with heat-denatured secretome, suggesting that these impacts on behaviour

were not related solely to the activity of proteins. Indeed, since these secretome samples

were denatured by heat in order to lose the secondary and tertiary structures of proteins,

and consequently their biological activities, it is possible that this behavioural change

might be induced by other molecules included in the secretome that were not as impacted

by the heat treatment. It is likely that *Schistocephalus* secretes and/or excretes other

molecules, such as extracellular vesicles including micro-RNA (65, 66). To confirm this

hypothesis in the future, we could replace the heat treatment of secretome by a protease

exposure, in order to specifically break down and inactivate proteins in the secretome.

Our findings demonstrate that the secretome of *S. solidus* is an important component of

the molecular cross-talk between the parasite and its threespine stickleback host.

Secretome alone is not able to reproduce the behavioural changes of infected

sticklebacks in non-infected sticklebacks. Yet, its effects on boldness in the allopatric

population suggest that the secretome may act in combination with other processes, like

the activation of the stress response in fish, to induce the complex behaviour of infected

sticklebacks. Future studies could explore how the secretome of *S. solidus*, in

combination with other factors linked to the activation of the stress response in

sticklebacks, could induce this complex set of behavioural changes.

Acknowledgements

The authors thank the personnel of the Laboratoire Aquatique de Recherche en Sciences Environnementales et Médicales at Université Laval for their help with fish rearing. We thank Christian Landry for his comments and advice during the development of this project. We thank Marie-Pier Brochu, Verônica Alves, Florent Sylvestre and Sann Delaive for comments on previous versions of this manuscript. This project was funded by an NSERC Discovery grant to NAH.

The authors declare no competing or financial interests.

Author contributions

C.S.B. designed the study with input from N.A.-H.. C.S.B. performed the experiments. C.S.B. analyzed the data with input from N.A.-H.. C.S.B. and N.A-H. wrote the manuscript.

Data availability

The complete dataset is available as a supplementary tab-delimited supp_dataset_final.txt file.

Ethics

This study was approved by the Comité de Protection des Animaux de l'Université Laval (CPAUL 2014069-3).

REFERENCES

1. Sandland GJ, Goater CP. Parasite-induced variation in host morphology: Brain-encysting trematodes in fathead minnows. *The Journal of Parasitology*. 2001**;87**(2):267-72.

2. Garvin MC, Szell CC, Moore FR. Blood parasites of nearctic-neotropical migrant passerine birds during spring trans-gulf migration: Impact on host body condition. *The Journal of Parasitology*. 2006;**92**(5):990-6.

3. Rigano R, Profumo E, Bruschi F, Carulli G, Azzara A, Ioppolo S, et al. Modulation of human immune response by *Echinococcus granulosus* antigen B and its possible role in evading host defenses. *Infection and Immunity*. 2001;**69**(1):288.

4. Watson DW, Mullens BA, Petersen JJ. Behavioral fever response of *Musca domestica* (Diptera: Muscidae) to infection by *Entomophthora muscae* (Zygomycetes: Entomophthorales). *Journal of Invertebrate Pathology*. 1993;**61**(1):10-6.

5. Lafferty K, Shaw J. Comparing mechanisms of host manipulation across host and parasite taxa. *The Journal of Experimental Biology*. 2013;**216**(1):56-66.

6. De Bekker C, Merrow M, Hughes DP. From behavior to mechanisms: An integrative approach to the manipulation by a parasitic fungus (*Ophiocordyceps unilateralis* s.l.) of its host ants (*Camponotus* spp .). *Integrative and Comparative Biology*. 2014;**54**(2):166-76.

7. Adamo SA. Parasites: evolution's neurobiologists. *The Journal of Experimental Biology*. 2013;**216**(1):3-10.

8. Biron DG, Loxdale HD. Host-parasite molecular cross-talk during the manipulative process of a host by its parasite. *The Journal of Experimental Biology*. 2013;**216**(1):148-60.

9. Herbison R, Lagrue C, Poulin R. The missing link in parasite manipulation of host behaviour. *Parasites & Vectors*. 2018;**11**(1): 222.

10. Coppens I, Vielemeyer O. Insights into unique physiological features of neutral lipids in *Apicomplexa*: from storage to potential mediation in parasite metabolic activities. *International journal for parasitology*. 2005;**35**(6):597-615.

11. Fromm B, Ovchinnikov V, Høye E, Bernal D, Hackenberg M, Marcilla A. On the
presence and immunoregulatory functions of extracellular micro RNA s in the trematode
Fasciola hepatica. *Parasite immunology*. 2017;**39**(2):e12399.

12. Grimwood J, Smith JE. *Toxoplasma gondii*: the role of parasite surface and
secreted proteins in host cell invasion. *International journal for parasitology*.
1996;**26**(2):169-73.

13. Coakley G, Maizels RM, Buck AH. Exosomes and other extracellular vesicles: the
new communicators in parasite infections. *Trends in parasitology*. 2015;**31**(10):477-89.

14. Harnett W. Secretory products of helminth parasites as immunomodulators.
Molecular & Biochemical Parasitology. 2014;**195**(2):130-6.

15. Hewitson JP, Grainger JR, Maizels RM. Helminth immunoregulation: The role of
parasite secreted proteins in modulating host immunity. *Molecular & Biochemical
Parasitology*. 2009;**167**(1):1-11.

16. Adamo SA. Turning your victim into a collaborator: Exploitation of insect
behavioural control systems by parasitic manipulators. *Current opinion in insect science*.
2019;**33**:25-29.

17. Hammerschmidt K, Kurtz J. Chapter 5 Ecological immunology of a tapeworms'
interaction with its two consecutive hosts. *Advances in Parasitology*. 2009;**68**:111-37.

18. Wedekind C. The infectivity, growth, and virulence of the cestode *Schistocephalus
solidus* in its first intermediate host, the copepod *Macrocyclops albidus*. *Parasitology*.
1997;**115**(3):317-24.

19. Dianne L, Perrot-Minnot MJ, Bauer A, Gaillard M, Léger E, Rigaud T. Protection
first then facilitation: a manipulative parasite modulates the vulnerability to predation of
its intermediate host according to its own developmental stage. *Evolution: International
Journal of Organic Evolution.* 2011;**65**(9):2692-8.

20. Hafer-Hahmann N. Experimental evolution of parasitic host manipulation.
Proceedings of the Royal Society B: Biological Sciences. 2019;**286**(1895):20182413.

21. Wedekind C, Milinski M. Do three-spined sticklebacks avoid consuming copepods,
the first intermediate host of *Schistocephalus solidus*? An experimental analysis of
behavioural resistance. *Parasitology*. 1996;**112**(4):371-83.

22. Weinreich F, Benesh D, Milinski M. Suppression of predation on the intermediate
host by two trophically-transmitted parasites when uninfective. *Parasitology*.
2013;**140**(1):129-35.

23. Barber I. Sticklebacks as model hosts in ecological and evolutionary parasitology.
Trends in Parasitology. 2013;**29**(11):556-66.

24. Barber I, Scharsack JP. The three-spined stickleback- *Schistocephalus solidus*
system: an experimental model for investigating host-parasite interactions in fish.
Parasitology. 2010;**137**(3):411-24.

25. Tierney JF, Crompton DWT. Infectivity of plerocercoids of *Schistocephalus solidus*
(Cestoda: Ligulidae) and fecundity of the adults in an experimental definitive host, *Gallus*
gallus. The Journal of Parasitology. 1992;**78**(6):1049-54.

26. Scharsack JP, Koch K, Hammerschmidt K. Who is in control of the stickleback
immune system: interactions between *Schistocephalus solidus* and its specific vertebrate
host. *Proceedings of the Royal Society B: Biological Sciences.* 2007**;274**(1629):3151-8.

27. Barber J, Walker P, Svensson PA. Behavioural responses to simulated avian
predation in female three spined sticklebacks: The effect of experimental *Schistocephalus*
solidus infections. *Behaviour*. 2004;**141**:1425-40.

28. Hebert FO, Grambauer S, Barber I, Landry CR, Aubin-Horth N. Major host
transitions are modulated through transcriptome-wide reprogramming events in
Schistocephalus solidus, a threespine stickleback parasite. *Molecular ecology*.
2017;**26**(4):1118-30.

29. Arber B, Huntingford FA, Crompton DWT. The effect of hunger and cestode
parasitism on the shoaling decisions of small freshwater fish. *Journal of Fish Biology*.
1995;**47**(3):524-36.

30. Grécias L, Valentin J, Aubin-Horth N. Testing the parasite mass burden effect on
alteration of host behaviour in the -stickleback system. *The Journal of Experimental*
Biology. 2018;**221**: jeb174748.

31. Giles N. Behavioural effects of the parasite *Schistocephalus solidus* (Cestoda) on
an intermediate host, the three-spined stickleback, *Gasterosteus aculeatus* L. Animal
Behaviour. 1983;**31**(4):1192-4.

32.   Grécias L, Hébert F-O, Berger CS, Barber I, Aubin-Horth N. Can the behaviour of
threespine stickleback parasitized with *Schistocephalus solidus* be replicated by
manipulating host physiology? *The Journal of Experimental Biology*. 2016;**220**(2):237-46.
33.   Overli O, Pll M, Borg B, Jobling M, Winberg S. Effects of *Schistocephalus solidus*
infection on brain monoaminergic activity in female three-spined sticklebacks
Gasterosteus aculeatus. *Proceedings of the Royal Society B: Biological Sciences*.
2001;**268**(1474):1411-5.
34.   Scharsack JP, Gossens A, Franke F, Kurtz J. Excretory products of the cestode,
Schistocephalus solidus, modulate in vitro responses of leukocytes from its specific host,
the three-spined stickleback (*Gasterosteus aculeatus*). *Fish & shellfish immunology*.
2013;**35**(6):1779-87.
35.   Yoneva A, Scholz T, Kuchta R. Comparative morphology of surface ultrastructure
of diphyllobothriidean tapeworms (Cestoda: Diphyllobothriidea). *Invertebrate biology*.
2018;**137**(1):38-48.
36.   Poulin R. Parasite manipulation of host behavior-Chapter 5: An update and
frequently asked questions. *Advances in the Study of Behavior*. 2010;**41**(C):151-86.
37.   Orr TSC, Hopkins CA, Charles GH. Host specificity and rejection of
Schistocephalus solidus. *Parasitology*. 1969;**59**(3):683-90.
38.   Kennedy C. A checklist of British and Irish freshwater fish parasites with notes on
their distribution. *Journal of Fish Biology*. 1974;**6**(5):613-44.
39.   Hamley M, Franke F, Kurtz J, Scharsack JP. An experimental approach to the
immuno-modulatory basis of host-parasite local adaptation in tapeworm-infected
sticklebacks. *Experimental parasitology*. 2017;**180**:119-32.
40.   Kalbe M, Eizaguirre C, Scharsack JP, Jakobsen PJ. Reciprocal cross infection of
sticklebacks with the diphyllobothriidean cestode *Schistocephalus solidus* reveals
consistent population differences in parasite growth and host resistance. *Parasites &*
vectors. 2016;**9**(1):130.
41.   Nishimura N, Heins DC, Andersen RO, Barber I, Cresko WA. Distinct lineages of
Schistocephalus parasites in threespine and ninespine stickleback hosts revealed by
DNA sequence analysis. *PLoS One*. 2011;**6**(7):e22505.

42. Piecyk A, Ritter M, Kalbe M. The right response at the right time: Exploring
helminth immune modulation in sticklebacks by experimental coinfection. *Molecular*
ecology. 2019;**28**(10): 2668-2680.

43. Weber JN, Kalbe M, Shim KC, Erin NI, Steinel NC, Ma L, et al. Resist Globally,
Infect Locally: A transcontinental test of adaptation by stickleback and their tapeworm
parasite. *The American naturalist*. 2017;**189**(1):43.

44. Simmonds NE, Barber I. The effect of salinity on egg development and viability of
Schistocephalus solidus (Cestoda: Diphyllobothriidea). *Journal of Parasitology*.
2016;**102**(1):42-7.

45. Confer A, Vu V, Drevecky C, Aguirre W. Occurrence of *Schistocephalus solidus* in
anadromous threespine stickleback. *Journal of Parasitology*. 2012;**98**(3):676-8.

46. Macat Z, Bednarik A, Rulik M. The three-spined stickleback (*Gasterosteus*
aculeatus) infection with *Schistocephalus solidus* in Hel marina (Puck Bay, Baltic Sea,
Poland). *Oceanological and hydrobiological studies*. 2015;**44**(1):11-7.

47. Franke F, Rahn AK, Dittmar J, Erin N, Rieger JK, Haase D, et al. In vitro leukocyte
response of three-spined sticklebacks (*Gasterosteus aculeatus*) to helminth parasite
antigens. *Fish & shellfish immunology*. 2014;**36**(1):130-40.

48. De Bekker C, Quevillon LE, Smith PB, Fleming KR, Ghosh D, Patterson AD, et al.
Species-specific ant brain manipulation by a specialized fungal parasite. *BMC*
evolutionary biology. 2014;**14**(1):166.

49. Thompson JN. The role of coevolution. *Science*. 2012;**335**(6067):410-1.

50. Franceschi N, Cornet S, Bollache L, Dechaume-Moncharmont FX, Bauer A,
Motreuil S, et al. Variation between populations and local adaptation in acanthocephalan-
induced parasite manipulation. *Evolution*. 2010;**64**(8):2417-30.

51. Cao X, Fu Z, Zhang M, Han Y, Han Q, Lu K, et al. Excretory/secretory proteome
of 14-day schistosomula, *Schistosoma japonicum*. *Journal of Proteomics*.
2016;**130**(C):221-30.

52. Poulin R, FitzGerald GJ. Risk of parasitism and microhabitat selection in juvenile
sticklebacks. *Canadian Journal of Zoology*. 1989;**67**(1):14-8.

53. Link AJ, Labaer J. Trichloroacetic acid (TCA) precipitation of proteins. *Cold Spring*
Harbor protocols. 2011;**2011**(8):993.

54.   Berger CS, Aubin-Horth N. An eDNA-qPCR assay to detect the presence of the
parasite *Schistocephalus solidus* inside its threespine stickleback host. *The Journal of*
Experimental Biology. 2018; **221**: jeb178137.

55.   Noldus L, Spink A, Tegelenbosch R. EthoVision: A versatile video tracking system
for automation of behavioral experiments. *Behavior Research Methods, Instruments, &*
Computers. 2001;**33**(3):398-414.

56.   RCoreTeam. R: A language and environment for statistical computing. *R*
Foundation for Statistical Computing, Vienna, Austria. 2018.

57.   Bates D, Maechler M, Bolker B, Walker S. Fitting linear mixed-effects models using
lme4. *Journal of Statistical Software*. 2015;**67**(1):1-48.

58.   Venables WN, Ripley BD. Modern Applied Statistics with S. Fourth Edition.
Springer, New York. 2002.

59.   Kuznetsova A, Brockhoff P, Christensen R. lmerTest package: Tests in linear
mixed effects models. *Journal of Statistical Software*. 2017;**82**(13):1–26.

60.   Ditgen D, Anandarajah EM, Meissner KA, Brattig N, Wrenger C, Liebau E.
Harnessing the helminth secretome for therapeutic immunomodulators. *BioMed research*
international. 2014;**2014**: 964350.

61.   Talarico M, Seifert F, Lange J, Sachser N, Kurtz J, Scharsack JP. Specific
manipulation or systemic impairment? Behavioural changes of three-spined sticklebacks
(*Gasterosteus aculeatus*) infected with the tapeworm *Schistocephalus solidus*.
Behavioral ecology and sociobiology. 2017;**71**(2):36.

62.   Fayard M, Cezilly F, Perrot-Minnot M-J. Inter-population variation in the intensity
of host manipulation by the fish acanthocephalan *Pomphorhynchus tereticollis*: are
differences driven by predation risk? *Parasitology*. 2019;**146**(10):1296-304.

63.   Rosenkranz M, Poulin R, Selbach C. Behavioural impacts of trematodes on their
snail host: Species-specific effects or generalised response? *Ethology*.
2018;**124**(11):790-5.

64.   Piecyk A, Roth O, Kalbe M. Specificity of resistance and geographic patterns of
virulence in a vertebrate host-parasite system. *BMC evolutionary biology*. 2019;**19**(1):80.

65.   Twu O, Johnson PJ. Parasite extracellular vesicles: Mediators of intercellular
communication. *PLoS Pathogens*. 2014;**10**(8):e1004289.

66. Zaborowski MP, Balaj L, Breakefield XO, Lai CP. Extracellular vesicles:

Composition, biological relevance, and methods of study. *BioScience*. 2015;**65**(8):783-

97.

 FIGURES

Figure 1. Two boldness-related behaviours differ between *S. solidus*-infected and
non-infected sticklebacks. 1. Behaviours that were significantly different between
infected and non-infected sticklebacks ($p < 0.05$). A. latency to feed before a predator
attack (sec). B. time spent frozen after a predator attack (sec). **2.** Behaviours that were
non-significantly different between infected and non-infected sticklebacks. A. latency to
leave the acclimation zone (sec). B. time spent swimming (sec). C. time in the center
(sec). D. time spent fleeing after a predator attack (sec). E. latency to feed after a predator
attack (sec). Each comparison was performed between infected fish and non-infected fish
(from the PBS group) of the control week. For each boxplot, the black line represents the
median and the green triangle represents the average. Each dot stands for one individual.

Figure 2. Injections of non-infective secretome significantly lowers the expression
of boldness-related behaviours, only in fish from the allopatric population. 1.
Latency to feed before a predator attack (sec) in the Témiscouata population. A. Infected
fish. B. Fish injected with control PBS. C. Fish injected with "infective secretome". D. Fish
injected with "non-infective secretome". E. Fish injected with "denatured secretome". **2.**
Latency to feed before a predator attack (sec) in the anadromous population. A. Fish
injected with control PBS. B. Fish injected with "infective secretome". C. Fish injected with
"non-infective secretome". D. Fish injected with "denatured secretome". For each boxplot,
the black line represents the median and the green triangle represents the average. Each
dot stands for one individual. Each grey line is the reaction norm of one individual between
the control week and the injection week.

Figure 1.

Figure 2.

796 **Supplementary Figure 1. Visualization of protein content of secretome on SDS-**
797 **PAGE gel. 1.** "Infective secretome". **2.** "Non-infective secretome". **3.** "Denatured
798 secretome". A. Ladder. B. Yeast extract. C. and D. Secretome.

Supplementary Figure 2. Latency to leave the acclimation zone (sec) 1. in the Témiscouata population. A. Infected fish. B. Fish injected with control PBS. C. Fish injected with "infective secretome". D. Fish injected with "non-infective secretome". E. Fish injected with "denatured secretome". **2.** in the anadromous population. A. Fish injected with control PBS. B. Fish injected with "infective secretome". C. Fish injected with "non-infective secretome". D. Fish injected with "denatured secretome". For each boxplot, the black line represents the median and the green triangle represents the average. Each dot stands for one individual. Each grey line is the reaction norm of one individual between the control week and the injection week.

Supplementary Figure 3. Time spent swimming (sec) 1. in the Témiscouata population. A. Infected fish. B. Fish injected with control PBS. C. Fish injected with "infective secretome". D. Fish injected with "non-infective secretome". E. Fish injected with "denatured secretome". **2.** in the anadromous population. A. Fish injected with control PBS. B. Fish injected with "infective secretome". C. Fish injected with "non-infective secretome". D. Fish injected with "denatured secretome". For each boxplot, the black line represents the median and the green triangle represents the average. Each dot stands for one individual. Each grey line is the reaction norm of one individual between the control week and the injection week.

Supplementary Figure 4. Time spent in the center of the aquarium (sec) 1. in the
Témiscouata population. A. Infected fish. B. Fish injected with control PBS. C. Fish
injected with "infective secretome". D. Fish injected with "non-infective secretome". E.
Fish injected with "denatured secretome". **2.** in the anadromous population. A. Fish
injected with control PBS. B. Fish injected with "infective secretome". C. Fish injected with
"non-infective secretome". D. Fish injected with "denatured secretome". For each boxplot,
the black line represents the median and the green triangle represents the average. Each
dot stands for one individual. Each grey line is the reaction norm of one individual between
the control week and the injection week.

Supplementary Figure 5. Time spent fleeing after a predator attack (sec) 1. in the Témiscouata population. A. Infected fish. B. Fish injected with control PBS. C. Fish injected with "infective secretome". D. Fish injected with "non-infective secretome". E. Fish injected with "denatured secretome". **2.** in the anadromous population. A. Fish injected with control PBS. B. Fish injected with "infective secretome". C. Fish injected with "non-infective secretome". D. Fish injected with "denatured secretome". For each boxplot, the black line represents the median and the green triangle represents the average. Each dot stands for one individual. Each grey line is the reaction norm of one individual between the control week and the injection week.

Supplementary Figure 6. Time spent frozen after a predator attack (sec) 1. in the Témiscouata population. A. Infected fish. B. Fish injected with control PBS. C. Fish injected with "infective secretome". D. Fish injected with "non-infective secretome". E. Fish injected with "denatured secretome". **2.** in the anadromous population. A. Fish injected with control PBS. B. Fish injected with "infective secretome". C. Fish injected with "non-infective secretome". D. Fish injected with "denatured secretome". For each boxplot, the black line represents the median and the green triangle represents the average. Each dot stands for one individual. Each grey line is the reaction norm of one individual between the control week and the injection week.

Supplementary Figure 7. Latency to feed after a predator attack (sec) 1. in the Témiscouata population. A. Infected fish. B. Fish injected with control PBS. C. Fish injected with "infective secretome". D. Fish injected with "non-infective secretome". E. Fish injected with "denatured secretome". **2.** in the anadromous population. A. Fish injected with control PBS. B. Fish injected with "infective secretome". C. Fish injected with "non-infective secretome". D. Fish injected with "denatured secretome". For each boxplot, the black line represents the median and the green triangle represents the average. Each dot stands for one individual. Each grey line is the reaction norm of one individual between the control week and the injection week.

 SUPPLEMENTARY MATERIAL

Supplementary tables in PDF format. Results of all the statistical comparisons
performed with the lmer function to determine if the secretome injections change
the behaviour of non-infected fish of the Témiscouata population and of the
anadromous population. Results are reported in separate tables for each of the seven
behaviours and for each population (supplementary tables 1 to 14).

Supplementary dataset in tab-delimited format. Complete dataset of behavioural
and phenotypic results for the Témiscouata and the anadromous populations of
threespine stickleback. Data on seven behaviours measured during the control and
treatment weeks, mass (g), size (mm) and sex of each fish are reported, as well as the
infection status (no=0; yes=1), the number and the total mass of *S. solidus* (g) for
individual fish in each treatment.