Microstructures amplify seemingly honest signals in carotenoid-colored tanagers

Authors: Dakota E. McCoy[1*], Allison J. Shultz[1,2,3,4], Charles Vidoudez[5], Emma van der Heide[1], Sunia A.Trauger[5], David Haig[1]

1. Department of Organismic and Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, MA 02138
2. Informatics Group, Harvard University, 38 Oxford Street, Cambridge, MA 02138
3. Museum of Comparative Zoology, Harvard University, 26 Oxford Street, Cambridge, MA 02138
4. Ornithology Department, Natural History Museum of Los Angeles County, 900 Exposition Blvd, Los Angeles, CA 90007
5. Small Molecule Mass Spectrometry Facility, Harvard University, 52 Oxford Street (B2), Cambridge, MA 02138
      * corresponding author: dakotamccoy@g.harvard.edu

Abstract:
Red, orange, and yellow carotenoid-colored plumages have been considered honest signals of condition. To investigate this interpretation, we comprehensively quantify carotenoid signals in the social, sexually-dimorphic tanager genus *Ramphocelus* using scanning electron microscopy (SEM), finite-difference time-domain (FDTD) optical modeling, liquid chromatography–mass spectrometry (LC-MS), and spectrophotometry. Males and females within a species have equivalent amounts and types of carotenoids, which is surprising if carotenoids are an honest-because-costly signal. Male, but not female, feathers have elaborate microstructures which amplify color appearance. Expanded barbs enhance color saturation (for the same amount of pigment) by increasing the transmission of optical power through the feather. Vertically-angled, strap-shaped barbules generate "super black" plumage, an optical illusion to enhance nearby color. Together, our results suggest that a widely cited index of honesty—carotenoid pigments—cannot fully explain male appearance. We propose that males are selected to evolve amplifiers of honest signals.

Keywords: sexual selection, honest signaling, Goodhart's law, carotenoid pigments, microstructure, LCMS, FDTD

Introduction

      Why are so many male birds beautiful? To investigate this evolutionary "why," we study both physical mechanisms of color (pigments and structures) and the evolutionary mechanisms which favor colorful signals over evolutionary time (selective forces such as honest signaling, aesthetic preference, and sensory bias). We must fully understand the physical basis of beautiful traits in order to infer their evolutionary history.

      A complete understanding of the physical cause of colorful ornaments may help us determine the relative importance of three overlapping selective pressures within mate choice: species identity, aesthetic beauty, and individual quality (Hill 2015). Coloration may facilitate species identification, essential to avoid sterile hybrids and wasted mating efforts. Beautiful ornaments may reflect arbitrary aesthetic preferences

in the choosing sex (Prum 2012), may be maintained through a Fisherian runaway process (Fisher 1999),
or may occur as a side effect of selection on another domain such as foraging—termed "sensory bias"
(Dawkins and Guilford 1996). Finally, color may be an indicator of individual quality ("honest signaling
theory") due to physiological linkage, resource-trade-offs, or direct/indirect costs. In birds, bright feathers
may indicate nutritional condition (Hill and Montgomerie 1994), parasite resistance (Folstad and Karter
1992), immunocompetence (Simons et al. 2012), or ability to survive despite an extreme "handicap" of
beautiful ornaments (Zahavi 1975). Of the three selective pressures, research to date has largely focused on
individual quality, as captured through a large body of literature on honest signaling theory. Frequently,
researchers use the physical, pigmentary basis of colorful signals to draw conclusions about honest
signaling.
Carotenoid-pigmented plumages, ranging in hue from yellow to red, have been considered an
honest signal because carotenoids must be eaten by vertebrates rather than synthesized, may be scarce in
nature, and serve important immunological functions (Olson and Owens 1998). However, the relationship
between carotenoids, signal appearance, and individual quality is not yet fully understood. Carotenoids are
correlated with only some, but not all, individual quality measures (Simons et al. 2012; Koch et al. 2018;
Weaver et al. 2018). Importantly, carotenoids are consumed in a yellow (dietary) form and then metabolized
within vertebrate bodies to become immunologically useful molecules that are redder in color. A large
meta-analysis found that metabolized, not dietary, carotenoids drive the (weak) overall correlation between
appearance and health (Weaver et al. 2018), suggesting that carotenoids are an index of proper metabolic
function rather than a costly signal (Weaver et al. 2017). Further, it is often thought that carotenoids are a
limiting resource in the immune system, but counter to this view, canaries with a mutation knocking out
tissue carotenoids show no difference in immune system function (Koch et al. 2018). The "honesty" of
carotenoid-based signals is not yet fully understood, nor is the complete physical basis of colorful
carotenoid plumages (although we know that nano- and microstructures can alter the appearance of
carotenoid-colored organisms (Shawkey and Hill 2005; Iskandar et al. 2016)). Is it possible that
microstructural elements to carotenoid coloration in birds complicate the link between signal and individual
quality, contributing to mixed research results?
The physical basis of color, including microstructural contributions, is not only intrinsically
interesting but is also critical to our understanding of sexual selection and evolutionary dynamics.
Nanostructures and pigments produce colors through well-understood pathways (Hill and McGraw 2006),
but only recently have researchers begun to describe how microstructures can substantially enhance or alter
pigmentary colors. Carotenoid pigments produce reds, oranges, and yellows, but microstructures make the
colors glossy or matte (Iskandar et al. 2016). The African Emerald Cuckoo *Chrysococcyx cupreus* produces
stunning color with milli-scale emerald mirrors (Harvey et al. 2013). Birds-of-paradise (McCoy et al. 2018)
and birds from 14 other families (McCoy and Prum 2019) convergently evolved super black plumage with
vertically-oriented barbules that enhance melanin-based absorption by multiply scattering light. Outside of
birds, many organisms use microstructures to enhance the impact of pigments, from super black peacock
spiders (McCoy et al. 2019), stick insects (Maurer et al. 2017), and butterflies (Vukusic et al. 2004) to
flowers with conical epidermal cells that generate richer petal colors (Kay et al. 1981; Gorton and
Vogelmann 1996; Gkikas et al. 2015; Wilts et al. 2018). Intriguingly, some researchers have described
substantial microstructural variation between male and female birds (Enbody et al. 2017). To what extent
may microstructural, rather than pigmentary, differences explain why so many males are brilliantly colored
while many females are drab? The answer may help us determine which selective forces cause so many
male birds to be beautiful.
Herein, we aim to better understand the physical basis of color and thus draw inferences about the
evolutionary dynamics of colorful signals, honest or otherwise. We focus on the sexually dimorphic
Ramphocelus tanagers, a useful clade for questions of visual signaling (Burns and Shultz 2012; Shultz and
Burns 2017). These tanagers have carotenoid-based coloration ranging from bright yellow to deep velvet
red in males, while females are relatively duller (Figure 1). *Ramphocelus* tanager behavior is not well-
known, but behavioral and morphological observations suggest that they are subject to sexual selection and
mate choice. The genus is composed of social, brightly colored birds, traits that tend to align with mate

choice (Hill 2006). *Ramphocelus costaricensis* has high rates of extra-pair matings (Krueger et al. 2008),
and sexual displays have been described in *R. bresilius* (Harris 1987), *R. dimidiatus*, and *R. carbo*
(Moynihan 1962).

Using scanning electron microscopy (SEM), finite-difference time-domain (FDTD) optical
modeling, pigment extraction, liquid chromatography–mass spectrometry (LC-MS), and
spectrophotometry, we comprehensively document the physical basis of color in both male and female
Ramphocelus tanagers. From this, we make inferences about the dynamics of mate choice over evolutionary
time. We find evidence that carotenoid-based signals in these birds are altered by substantial microstructural
influence, and thus are not classically honest. Specifically, we suggest that males, under intense selection
to satisfy female testing criteria, evolve "amplifiers" to honest signals (microstructures that enhance light-
pigment interactions, generating a more saturated color). Females may prefer bright reds as a signal of male
quality, but males are incentivized to find non-costly means to amplify their appearance. As they say in
economics, "when a measure becomes a target, it ceases to be a good measure (Strathern 1997).

100
Figure 1. Male *Ramphocelus* tanagers (top row) have more striking carotenoid-based coloration
than females (bottom row). A. *R. flammigerus icteronotus* male with vivid yellow and velvety black
plumage. B. *R. flammigerus icteronotus* female. C. *R. flammigerus* male with vivid orange and velvety
black plumage. D. *R. flammigerus* female. E. *R. bresilius* male with bright red plumage. F. *R. bresilius*
female. G. *R. carbo* male with deep, velvety red and black plumage. H. *R. carbo* female. All photos are
credit Nick Athanas - www.antpitta.com.

Material and Methods
110
Specimens
Ramphocelus tanager specimens were selected from the Ornithology Collections at the Harvard
Museum of Comparative Zoology (MCZ; specimen numbers are listed in Table S1). We selected intact
male and female specimens from each species in the genus *Ramphocelus,* plus one subspecies with visually
divergent plumage that has had full species status in the past (Griscom 1932): Crimson-collared tanager,
Ramphocelus sanguinolentus; Masked crimson tanager, *Ramphocelus nigrogularis;* Crimson-backed
tanager, *Ramphocelus dimidiatus;* Huallaga tanager, *Ramphocelus melanogaster;* Silver-beaked tanager,
Ramphocelus carbo; Brazilian tanager, *Ramphocelus bresilius;* Passerini's tanager, *Ramphocelus*
passerinii; Cherrie's tanager, *Ramphocelus costaricensis;* Flame-rumped tanager, *Ramphocelus*
flammigerus; Lemon-rumped tanager, *Ramphocelus flammigerus icteronotus.* Taxonomy is according to
the Clements Checklist v.2017 (Clements et al. 2017).
We designate *R. flammigerus, R. f. icteronotus*, *R. passerinii*, and *R. costaricensis* to be the
"rumped" tanagers because they form a clade with vivid color restricted to the rump. We designate the
clade of tanagers with color on their whole body to be the "whole body" clade (*Ramphocelus nigrogularis*,
Ramphocelus dimidiatus, *Ramphocelus melanogaster*, *Ramphocelus carbo*, *Ramphocelus bresilius*). *R.*
sanguinolentis is the sister to all others.
127
Spectrophotometry
We performed spectrophotometry using an OceanOptics PX2 with a pulsed xenon light source, and
OceanOptics USB4000, and an OceanOptics Spectralon White Standard. We used an integration time of
100 ms, averaged 5 scans, and set a boxcar width of 5. We measured three replicates per plumage patch.
For each specimen, we measured the bird at 7 locations: crown, back, rump, dorsal tail, throat, breast, and
belly. Using an OceanOptics reflectance probe holder, we measured reflectance for 90° incident light and
for 45° incident light. These two angles of incidence allow us to determine the directionality and structural
absorption potential of the plumage patches.
136
Mass Spectrometry
We prepared feathers for pigment extraction using a simple mechanical procedure. First, we washed
all feathers in hexane and allowed them to air dry. Next, we trimmed off the entirety of colored portions of
barbules from three feathers. We carefully weighed them, then placed these barbules with 0.3 mL methanol
in a screw cap micro tube in the presence of 10 ceramic beads (2mm). We subjected these tubes to bead
beating in a FastPrep FP120 for three cycles, each for 45 seconds at level 6.0. We centrifuged the resulting
mixture in an Eppendorf centrifuge 5417R for 5 minutes at 3000 RCF, then transferred the supernatant and
dried the barbules under a stream of nitrogen. The samples were resuspended in 100 µL of
acetonitrile:methanol (1:1) and transferred to micro inserts in amber glass vials. All samples were kept at -
80 °C until analysis.
Initially, we took 9 feathers per species (three feathers from each of three specimens for two
species) to ensure that we had enough material; for all additional species we took 3 feathers from a single
individual, which proved sufficient. In order to ensure that results were repeatable between individuals of
a single species, we extracted pigments from two additional individuals for each of two species—*R.*
flammigerus and *R. flammigerus icteronotus*. The pigment profiles were significantly correlated (Figure
S3, Linear regression output for *R. flammigerus*: slope = 0.88, SE = 0.066, R^2 = 0.87, p < 0.0005. Linear
regression output for *R. f. icteronotus*: slope = 0.96, SE = 0.054, R^2 = 0.92, p < 0.0005), indicating that our
extraction and characterization procedure was repeatable across individuals of a species. See Statistics
section for details.
We acquired pigment standards for 8-apo carotenal, Canthaxanthin, Astaxanthin (Sigma Aldrich),
Lycopene (Santa Cruz Biotechnology), and Alloxanthin (ChromaDex). Carotenoid contents were analyzed
using an Ultimate 3000 LC coupled with a Q-Exactive Plus hybrid quadrupole-orbitrap mass spectrometer
(Thermo Fisher Scientific). The LC was fitted with a Kinetex C18 (2.6 µm, 100 Å, 150 mm x 2.1 mm,

Phenomenex) column and the mobile phases used were MA: Acetontrile:Methanol 85:15 and MB: 2-
propanol. The gradient elution started with 4 min of 5% MB, then to 100% MB in 11 min, followed by 4
min at 100% MB, 1 min to get back to 5% MA and 4 min re-equilibration at 5% MB. The flow was kept
constant at 0.185 mL min^{-1}, and 10 µL of samples were injected. Electrospray ionization was used in
positive ion mode. The mass spectrometer was operated at 70,000 resolving power, scanning the m/z 150-
1500 range, with data-dependent (top 5) MS/MS scan at 17500 resolution, 1 m/z isolation and fragmentation
at stepped normalized collision energy 15, 35 and 55. The data were first manually analyzed to identify
peaks of ions with accurate masses corresponding to potential carotenoids within a 5ppm mass accuracy
threshold. The potential carotenoids included both known molecules, and those with a similar molecular
formula based on their accurate mass and mass defect. MS/MS fragmentation spectra were then compared
with the standards available or to MS/MS database entries within METLIN (-metlin.scripps.edu) to assign
identity when standards were available or putative identity based on fragmentation pattern similarity.
Lastly, peaks corresponding to carotenoids or carotenoids-candidates were integrated using Quan Browser
(Xcalibur, Thermo Fisher Scientific). Pigments were grouped into "families", based on their accurate mass
and relation to standards and fragmentation similarities.
175
SEM & Feather Microstructure
In preparation for SEM, we mounted a single feather from each specimen on a pin using a carbon
adhesive tab. We sputter-coated the feathers with ~12 nm of Pt/Pd at a current of 40 mA. We performed
scanning electron microscopy on an FESEM Ultra55.
Using ImageJ, we made multiple measurements on each SEM image, including maximum barb
width, top-down barb width, barbule width, inter-barbule distance, barb-barbule angle, and barbule length.
We also coded feathers according to whether the barbules were strap shaped or cylindrical, and whether the
barbules emerged from the plane of the feather (i.e., had "3-D" structure).
184
Optical Modelling
To approximate the optical effect of the observed microstructures, we performed finite-difference
time-domain (FDTD) simulations using the commercially available software Lumerical FDTD. We
constructed idealized two-dimensional feather cross sections for two apparently important microstructural
features, as determined by the PCA described below): (i) increasing barbule angle (a purported cause of
low-reflectance, super black plumage), (ii) oblong and expanded barb, as is typical in colorful male feathers
but not colorful female feathers (a purported enhancer of color). We simulated the interaction of light with
these structures to understand the impact of microstructure on plumage appearance. For angled barbules (i),
we assessed total reflectance based on structure alone without considering the contribution of melanin. In
this manner we can calculate the quantitative decrease in feather reflectance from structure alone. For
oblong expanded barbs (ii), we calculated the total optical power transmission through the pigmented
portion of the feather. This is a proxy for light-pigment interactions, which strongly influences color: the
greater the total optical power transmission through pigment, the greater the pigmentary action. In this
manner we can approximate the structural enhancement of pigment due to barb size and shape in males.
FDTD uses the Yee cell method (Taflove and Hagness 2005) to calculate the spatio-temporal
electromagnetic field distribution resulting from an initial pulse launched into the simulation domain. For
each simulation, a plane wave of light (ranging from 400-700nm) was normally incident (y-direction) on
an idealized feather cross-section in the x-y plane. We performed 2D simulations without incorporating
pigment because (i) we wished to isolate the effect of structure in a simple, interpretable manner and (ii)
the real and imaginary components of carotenoid absorption are not well-understood, as would be required
for simulations incorporating pigment. The simulation domain was bounded at the horizontal edges of the
y plane by perfectly matched layers (PML) and at the vertical edges by periodic boundary conditions, to
simulate a feather plumage. Frequency domain field monitors were placed above and below the structure
to collect the reflected and transmitted light, respectively. For our simulations of the oblong, expanded barb,

we included a 2D square monitor to capture optical power transmission in a 7.5 x 7.5 µm^2 square (the
purported pigmented region). We used a mesh size of 0.025 x 0.025 µm^2.
To focus only on the impact of surface reflections and light's path through the feather after surface
diffraction, we extended the bottom feather surface to beyond the bottom vertical PML boundary in the y-
plane. This was a measure to eliminate reflections from the underside of the feather for two reasons. First,
in life, feathers are arranged in complex stacks in a plumage, which we did not simulate; feathers are flush
with other feathers (i.e., the bottom of a feather would be a feather-feather interface, not a feather-air
interface). Second, had we included reflections from the underside of a feather, this may over-exaggerate
the extent to which microstructure enhances light-pigment interactions.
For our simulations of the male-typical barb vs. female-typical barb, we simulated an average
female feather barb based on our measurements: 22.2 µm tall by 18.9 µm wide—and an average male
feather barb: 39.8 µm by 27.7 µm. For our simulations of angled barbules, we simulated a feather with
hemispherical barb (radius 12.5 µm) and barbules 80 µm long varying in angle from 0 to 80°.
222
Phylogeny
Several recent studies have examined the phylogeny of *Ramphocelus* tanagers (Hackett 1996;
Burns and Racicot 2009; Burns et al. 2014). Unfortunately Burns and Racicot (2009) and Burns et al.
(2014) did not include *R. f. icteronotus*, and Hackett (1996) did not include *R. melanogaster* or *R.
dimidiatus*, but the two studies combined had *cytochrome b* (cyt b) sequences available for all species. To
construct a phylogeny, we downloaded cyt b sequences from NCBI for all *Ramphocelus* species, and six
outgroups from the Tachyphoninae clade (*Tachyphonus coronatus, T. rufus, T. phoenicius, Eucometis
penicillate, Lanio fulvus, and T. luctuosus*). NCBI accession numbers are available in Table S2. We
aligned the sequences with MAFFT v. 7.313 (Katoh and Standley 2013), and constructed the best-scoring
maximum likelihood phylogeny with Raxml v. 8.2.10 using the GTRCAT model with 100 bootstrap
replicates (Stamatakis 2006, 2014). The resulting phylogeny confirmed the relationships found by
previous studies, and so we extracted the monophyletic *Ramphocelus* clade for use in downstream
analyses (Figure 3).
236
Statistics
We performed principal component analyses (PCAs) on microstructure and pigment levels in males
and females, using both standard and phylogenetic PCA methods in the software program R, version 3.4.3.
We log transformed and centered all data. For standard PCAs we used function prcomp in R, both centering
and scaling the data, on measurements of feather microstructure (Table S3) and pigment signal strengths
for all feathers from all species, both males and females. PC1 and PC2 were extracted and used for
visualizations; PC1 was used to assess correlations between males and females.
For phylogenetic PCAs, we used function phyl.pca in the R package phytools (Revell 2012), with
a lambda method of correlation, and PCA mode "cov" (covariance). For phylogenetic PCAs, we had to
reduce the character matrix, so we summed the pigment signals within each pigment family to generate 8,
rather than 29, individual characters. Phylogenetic PCAs cannot incorporate more than one measurement
per species, so we had to run separate phylogenetic PCAs for males and females.
We tested for correlations between the proportions of carotenoids detected (carotenoid signal, as
peak area) in two replicates each of two species, *R. flammigerus* and *R. f. icteronotus*. First, for each
individual, we normalized the signal for each carotenoid to account for variation in overall amounts of
carotenoid detected (versus proportion) by dividing the signal of each carotenoid by the sample weight and
then by the maximum signal detected for that individual. We then tested for a correlation between the two
individuals of each species separately with a linear model. We performed the same procedure to test for
correlations between the upper and lower throat of male *R. melanogaster* and the throat and rump of male
R. dimidiatus.
We also tested for correlations among the proportions of male and female carotenoids and male
and female feather structure. Because it is not possible to test more than one character per species at a time
in a phylogenetic framework, we decided to test for correlations between male and female measurements

with the PCA scores, which captured a large proportion of the variance. For both carotenoids and structure,
we tested PC1 and PC2, and used phylogenetic generalized least squares (PGLS; (Grafen 1989; Martins
and Hansen 1997)) with a Brownian motion model to account for phylogeny. We could not use PGLS for
multiple patches when measured, so we randomly selected one patch for each species. Analyses using the
alternative patch were qualitatively similar, but not presented here. Results are also robust to PGLS model
choice for phylogeny transformation. Note that these PC scores were taken from the non-phylogenetic PCA,
and thus are directly comparable between males and females (both of which were included in the PCA).
Finally, we reconstructed the evolutionary history of carotenoid evolution for each pigment family
for males and females. Because there might have been slight variations in isoform for different species, we
decided to focus on carotenoid families as the most biologically meaningful variables. First, for each
pigment family, we summed all isoforms for males and females, and log-transformed them (base 10). We
removed any species that were missing data for both males and female for that family. We did the
reconstruction with the contMap function in phytools v. 0.6-44 (Revell 2012, 2013), which uses maximum
likelihood to estimate states at internal nodes and interpolate these states along internal branches
(Felsenstein 1985).
275
276
277
278 **Results**
279

*Male **Ramphocelus** tanagers generally have darker blacks and more brilliant colors than females.*
Spectrophotometry revealed a wide range of yellows, oranges, and reds in males and females
(complete results in Figure S6). The males from multiple species had extremely low-reflectance black
plumage, with broadband flat reflectance below 0.5%, in areas adjacent to bright color patches (Figure 2A,
Figure S6). Females had typical black plumage reflectances, with a slight increase in reflectance at higher
wavelengths (Figure 2A, Figure S6). Vivid, highly saturated color patches in males typically reflected
almost no light for short wavelengths before sloping sharply upwards to reflect long wavelengths in the
yellow-orange-red space (Figure 2B, Figure S6). In contrast, females of most species had colorful patches
with a relatively more gradual upward slope, a greater-than-0 reflectance over a broad range of wavelengths,
and a relatively lower peak reflectance (Figure 2B, Figure S6).
One species, *R. carbo*, has dark red velvet color on its whole body. Spectrophotometry showed that
this plumage reflected very little light (Figure S6) in a manner similar to the velvet black feathers in other
species (Figure 2A).
In general, many of the colored feathers were strong directional reflectors, such that reflected light
differed in quantity (percent total reflectance) from 90° incident light to 45° incident light (Figure S6). For
example, the red velvet patches on *R. carbo* increased their reflectance from ~0-1% to roughly 5-7% when
the angle of light incidence changed from 90° to 45°. This characteristic, reflecting more light at a lower
angle of incidence, is a hallmark of feathers with microstructures that impact absorbance and reflection
(McCoy et al. 2018).
299

300
Figure 2. Male *Ramphocelus* tanagers generally have darker blacks and more vivid, saturated colors
than females. Spectrophotometry results. **A:** Males (solid lines) have a darker black color, with a flatter
reflectance curves, on their backs compared to females (dashed lines) for all species except *R.*
sanguinolentis. We measured the black back of each species except *R. dimidiatus*, for which we measured
the black throat. We excluded *R. bresilius*, which is all red except for its tail and wings. **B.** Males (solid
lines) have more vividly-colored rumps for all species except *R. flammigerus* and *R. f. icteronotus*. We
excluded *R. carbo* from this plot, because it is dark red rather than brilliant. Complete spectrophotometry
results for all body regions are in Figure S6.
309

Many dietary and metabolized carotenoids generate color in both males and females

We identified 29 distinct molecules that matched the absorption spectra of carotenoids. These fell into 8 groups by monoisotopic molecular mass, with 1-6 different molecules in each pigment group, which we refer to as pigment families (Table S4). We mapped the relative abundances of all described molecules onto a tree for both males and females (Figure 3), performed ancestral state reconstructions for each pigment family (Figure S1), and mapped all identified pigments onto a metabolic network that shows how different molecules can be modified within the body (Figure S2).

Five of the molecules matched our purchased pigment standards and could be conclusively identified. For the remaining pigments, the molecule's identity was inferred based on accurate mass, retention time, MS/MS spectra, and pigments commonly found in bird feathers as described in the literature. For example, it is difficult to distinguish between the two dietary carotenoids lutein and zeaxanthin. Using the metabolic networks presented in Morrison and Badyaev (2016) and LaFountain et al. (2015), we identify the molecules with the accurate mass 568.43 as zeaxanthin because none of the common avian derivates from lutein were found in any of our samples, while many derivates of zeaxanthin (and related metabolites) were found. If some or all of the detected molecules with this mass were lutein, it would not change the conclusions herein, because both are dietary pigments.

We found that dietary pigments zeaxanthin, β-cryptoxanthin, and apo-8-carotenal were found primarily in the rumped tanager clade (Figure 3, blue stars in Figure S2) while the whole-body tanagers and *R. sanguinolentis* had primarily metabolized carotenoids. The "rumped" tanager clade had less metabolized pigments than the "whole-body" clade and *R. sanguinolentis* (Figure 3). This was affirmed through ancestral state reconstructions (Figure S1). These mappings also showed qualitative concordance between male and female pigment profiles within a species, and qualitative concordance between pigment profiles of different regions within a bird (Figure 3).

Males and females (within a species) have significantly correlated carotenoid profiles

We found that pigment profiles were significantly correlated between males and females within species (Figure 4A; PGLS with pigmentary PC1 scores: coefficient = 0.46, SE = 0.15, p= 0.013). In other words, we performed a PCA of all pigment signal strengths in both males and females to generate a single PC1 score that represented pigment content (Figure 6A). We then assessed the correlation between male pigmentary PC1 scores and female pigmentary PC1 scores, controlling for phylogeny with PGLS, to determine whether males and females have similar carotenoid pigment profiles. To visualize this result, we plotted the relative signal strengths (a proxy for amount) of each pigment in feathers from a male and female of the same species (Figure 4A), along with example pigment profiles from male and female *R. bresilius* (Figure 4B). These plots of the raw data (Figures 3, 4) qualitatively support our quantitative conclusions based on PC1: male and female tanagers within a species have concordant pigment profiles. Further detailed chemical analyses of the pigments which are differentially expressed between males and females (Figure 4) would be valuable.

Figure 3. Many dietary and metabolized carotenoids generate color in both males and females.
Relative log-transformed presence of pigments across birds for A. males and B. females. Color saturation
indicates normalized signal strength of a given pigment molecule within a bird, where pale is least and
rich is most. Because standards were not available for all molecules, absolute quantification and response
factors determination was not possible. Relative intensity comparison can therefore only be made within
each molecule (column). Some comparisons are possible within each family, because response factors of
isomers are expected to be similar. White indicates that the pigment was not detected. + indicates the
exact molecule that matched a pigment standard; * indicates that the molecules in this family were
identified based on accurate mass, retention time, MS/MS spectra, and the literature (no pigment standard
was available); for complete identification information see Table S4. Abbreviations: t: throat; ut: upper
throat; r: rump; b: breast; c: chest. Artwork in male silhouettes credit Gabriel Ugueto; female silhouettes
are modified by Allison Shultz from original art by Gabriel Ugueto.

362
Figure 4: Males and females (within a species) have significantly correlated carotenoid profiles. A.
Male versus female carotenoid pigments; each point represents the signal strength (a proxy for presence
and amount) of a given pigment in both a male and female of one species. All values are log-transformed
and normalized; since 7 was the minimum detectable threshold for our LC-MS setup, we set all values of
0 equal to 7 to better display the data. **B.** Example carotenoid pigment profile for male (top) and female
(bottom) *R. bresilius*, demonstrating the similarity. Artwork in bird silhouettes credit Gabriel Ugueto;
female silhouettes are modified by Allison Shultz from original art by Gabriel Ugueto.
370

Males, but not females, have diverse and elaborate feather microstructure

Female *Ramphocelus* tanager feathers were mostly of standard microstructural appearance. Feather microstructure usually looks the same as feather macrostructure, with simple cylindrical barbules extending from the central cylindrical barb, all in a single plane (Figure 5A-B, Table 1, Table S3). In this manner feathers are similar to fractals; as you zoom in, the smaller features look the same as the overall large feather. The only female to exhibit broad variation on this theme was *R. sanguinolentis* (Table 1, Figure S5, Table S3).

In contrast, male *Ramphocelus* tanagers exhibited wide variation in feather microstructure (Figure 5, Table 1, Table S3, Figure S5), included widely expanded barbs, oblong barbs, strap-shaped barbules (rather than cylindrical), and 3D barbules that projected from the plane of the feather (Table 1). Among the 3D barbules, we observed a dihedral morphology in velvety black and velvety red feathers as described in *R. flammigerus* (McCoy and Prum 2019). Two of these unusual male morphological traits, which we assess through optical simulations, deserve special mention (Figure 5):

Dihedral. In the crown of *R. carbo*, feathers have densely packed, strap-shaped barbules pointing upwards out of the plane of the feather to form a dihedral structure, while the central barb was shaped like a razor or narrow triangular prism (Figure 5D-F). *Ramphocelus dimidiatus* shared the "red velvet" morphology present in *R. carbo* in its upper throat feathers, which are dark velvety red (Figure S5). Both are reminiscent of the structurally absorbing "super black" feather morphology present in *Ramphocelus flammigerus* (McCoy and Prum 2019). Similar morphology, to a lesser degree, was observed in *R. melanogaster* dark velvet throat feathers and both male and female *R. sanguinolentis*. To the human eye, dihedral feather structure generates a velvety appearance.

Expanded barb. Multiple species had expanded barbs, including the broad and roughly cylindrical barbs of the rump feathers of *R. flammigerus* (e.g., Figure5 G-H, *R. nigrogularis*, and *R. passerinii*. These feathers also had flatter, strap-shaped barbules shorter in length than that of females but wider in width. Additionally, the rump and body feathers of *R. dimidiatus* are vivid, glittery red feathers with a more extremely expanded central barb with a featureless surface (Figure 5G-I). They also have upward-slanting, densely packed, strap-shaped barbules (Figure 5G-I)**.** Together, this generates (to the human eye) strong specular reflection at an angle, so that the bird glitters when rotated in the hand.

401
402
Figure 5: Males, but not females, have diverse and elaborate feather microstructure. A-B. Female *R.*
carbo; SEMs show red chest feather. C-D. Male *R. carbo;* SEMs show velvet red-black back feather with
dihedral morphology. E-F: Male *R. dimidiatus*; SEMs show shiny red rump feather with "expanded barb"
morphology. G-H: Male *R. passerinii* bright orange rump feather with "expanded barb" morphology.
Scale bars in left column (A, C, E, G) are 200 µm; scale bars in right column (B, D, F, H) are 50 µm.
Artwork in inset silhouettes credit Gabriel Ugueto.
409

410
411
412

species	sex	region	3D barbules	strap-shaped barbules	oblong barb	wide barb
R. bresilius	f	rump	--	--	--	yes
R. carbo	f	chest	--	--	yes	--
R. costaricensis	f	throat	--	--	--	--
R. dimidatus	f	rump	--	--	--	--
R. dimidatus	f	throat	--	--	yes	--
R. flammigerus	f	breast	--	--	--	--
R. flammigerus	f	rump	--	yes	--	--
R. f. icteronotus	f	rump	--	yes	--	--
R. melanogaster	f	chest	--	--	--	--
R. melanogaster	f	throat	--	yes	--	--
R. nigrogularis	f	rump	--	--	yes	--
R. passerinii	f	rump	--	--	yes	--
R. sanguinolentis	f	rump	yes	yes	yes	yes
R. bresilius	m	rump	yes	yes	yes	--
R. carbo	m	back	yes	yes	yes	yes
R. carbo	m	breast	yes	yes	yes	yes
R. costaricensis	m	rump	yes	yes	--	yes
R. dimidatus	m	throat	yes	yes	--	yes
R. dimidiatus	m	rump	yes	yes	yes	yes
R. flammigerus	m	rump	yes	yes	--	yes
R. f. icteronotus	m	rump	yes	yes	yes	--
R. melanogaster	m	upper throat	yes	yes	yes	yes
R. melanogaster	m	lower throat	yes	yes	yes	yes
R. nigrogularis	m	rump	yes	yes	--	yes
R. passerinii	m	rump	yes	--	--	yes
R. sanguinolentis	m	rump	yes	yes	yes	yes

413
Table 1: Males, but not females, have diverse and elaborate feather microstructure. Male
Ramphocelus tanagers have many atypical microstructural features compared to females. "3D barbules"
refers to whether or not barbules extended upwards from the plane of the feather; "strap-shaped barbules"
refers to whether barbules were flattened rather than classically cylindrical; "oblong barb" refers to
whether barbs were taller than they were wide (numerical threshold: > median (max barb width)/(top-
down barb width)); wide barb refers to whether the barb was expanded (numerical threshold: > median
barb width value). Complete numerical measurements can be found in Table S3.

Microstructure, not carotenoids, differs between sexes

Males and females have differentiated microstructures but not pigments. A PCA of carotenoid pigments in all feathers from both males and females demonstrated that birds do not separate by sex but instead by phylogeny (rumped clade separate from whole-body clade; Figure 6A; PC1 31.5%, PC2 17.3%) which was supported by separate phylogenetic PCAs of males and females (Figure S4, Table S6). A PCA of feather microstructure measurements demonstrated that males and female cluster separately and are not correlated (Figure 6B; PC1 55.6%, PC2 20.8%; PGLS with microstructure PC1 scores: coefficient = -0.023, SE = 0.17, p= 0.90); a phylogenetic PCA was not possible to compare males and females, because such analyses can consider only one value per species, but we performed sex-specific phylogenetic PCAs of one patch per species and observed some clustering by clade (whole-body versus rumped, Figure S4) and that females cluster tightly with the exception of outliers *R. sanguinolentis* and *R. bresilius,* both of which are females with wider-than-median barbs (Table 1). *R. sanguinolentis* is also the female with the most male-like scores in the regular PCA (Figure 6B). Barb width, barbule width, barb oblongness, and interbarbule distance were primary axes along which males and females separated (Table S5). No species-specific or region-specific (e.g., rump feather versus chest feather) clustering was observed in these microstructural measurements (Figure S4).

Microstructure, not carotenoids, differs between bright and dark red patches within a bird

To better assess the relative contribution of microstructure and pigment, we compared within-bird pigment profiles of shiny saturated red patches versus dark velvety patches for both *R. melanogaster* and *R. dimidiatus* (Figure 7). For each bird, these two patches had significantly correlated pigment profiles (Figure 7A,D), but the SEM photos revealed large differences in feather microstructure between velvety and bright patches (Figure 7B,C,E,F). Expanded barbs were associated with bright saturated color while vertically-angled or dihedral barbules were associated with velvety color. We suggest that microstructure – possibly with associated changes in nanostructure -- may explain this optical difference, a hypothesis we explored directly through optical modelling.

Figure 6: Microstructure, not carotenoids, differs between sexes. A. PCA of log-transformed centered pigments in all patches from all males (triangles) and females (circles), showing clustering by phylogenetic position (rumped clade separates from whole-body clade). Male and female pigment profiles are significantly correlated (PGLS with pigmentary PC1 scores: coefficient = 0.46, SE = 0.15, p= 0.013). **B.** PCA of log-transformed centered microstructural measurements for all patches from all males and all females, showing clustering by sex (males separate from females). Male and female microstructures are not correlated (PGLS with microstructure PC1 scores: coefficient = -0.023, SE = 0.17, p= 0.90).

473
Figure 7: Microstructure, not carotenoids, differs between bright and dark red patches within a bird. **A:** Pigment profiles of *R. dimidiatus* dark velvet throat versus bright red rump, where each point represents the signal strength for one pigment molecule. **B:** *R. dimidiatus* with SEM images of dark red throat feather (top) compared to bright red rump feather (bottom). **C:** *R. melanogaster* dark velvet red throat versus brighter red lower throat. **D:** *R. melanogaster* with SEM images of dark red throat feather (top) compared to brighter red lower throat feather (bottom). All values are normalized such that the largest value (strongest signal) was set equal to one. Scale bars are 50 µm. Artwork in bird silhouettes credit Gabriel Ugueto.

Optical simulations show that male-typical microstructures can enhance blackness and colorfulness
Using 2D finite-difference time-domain (FDTD) simulations of idealized feather cross-sections,
we isolated the effect of two male-typical structure on reflectance: dihedral barbules and expanded barb.
We selected these two features because they were associated with (i) within-bird color changes from bright,
saturated red to velvety red (Figure 7) and (ii) male-female differences in color appearance within a species.
We found that "dihedral" barbules (McCoy and Prum 2019), projecting out of the plane of the
feather and associated with velvety blacks and reds in our study species, contribute to a lower reflectance
(Figure 8A). Total reflectance based on light-keratin interactions alone, without considering the
contribution of melanin or carotenoids, drops from ~4% to 0.5% as barbules increase in angle from 0° to
80°. That is, the dihedral feather morphology in super black regions of these tanagers are made blacker in
part due to structure alone. Likewise, the velvet red feathers in *R. carbo* and *R. dimidiatus* are made darker
and velvety by structure alone.
Further, we found that male-typical expanded feather barbs, which are wider than female barbs in
80% of species and more oblong than female barbs in 60% of our species (Table 1), substantially enhance
optical power flow through the pigmented portion of the feather (Figure 8B-C). In other words, this male-
typical barbule shape causes greater light-pigment interactions, even though females and males have the
same amount of pigment (Figure 4). Therefore, the structure alone may contribute to a richer, more saturated
color in the same manner as flower petals' conical epidermal cells (Gkikas et al. 2015).
506

508
**Figure 8. Optical simulations show that male-typical microstructures can enhance blackness and
colorfulness**. All simulations are finite-difference time-domain (FDTD) simulations on 2D idealized
feather cross-sections, simulating structural effects only without accounting for pigmentary absorption. **A:**
A male-typical black feather, with angled barbules, has structurally enhanced blackness compared to a
female-typical feather with flat barbules. As the barbule angle increases out of the plane of the feather,
percent reflectance arising from the keratin structure alone decreases from 4% to 0.5%. We did not simulate
absorption effects of melanin, but instead we isolated the effect of structure alone. **B:** For a female-typical
colored feather, optical power flow through the purported pigmented area (white square) ranges from 0.004-
0.005 W/m^2.   **C**: For a male-typical colored feather, an oblong, expanded central barb increases optical
power transmission through the purported pigmented area (white square; 0.005-0.016 W/m^2). This
enhanced power transmission through the pigmented region enhances pigmentary activity for the same
amount of incident light. For B-C, optical power transmission is shown for 700nm light, but the results are
similar for other wavelengths of light.

522
523 **Discussion**
524
Summary
Carotenoid-based coloration has often been invoked as an example of honest signaling. By the
simplest version of honest signaling theory, male appearance is determined by carotenoid pigments which
honestly indicate quality because they are rare, are costly, require a trade-off, and/or are an index of proper
metabolic function (Hill and Montgomerie 1994; Olson and Owens 1998; Faivre et al. 2003; Saks et al.
2003; Simons et al. 2012; Weaver et al. 2017, 2018; Koch et al. 2018). In contrast to expectations of honest
signaling, we show that differences between the sexes in coloration of *Ramphocelus* cannot be attributed
primarily to carotenoids. Males and females have the same amounts and types of carotenoid pigments.
Instead, unusual microstructures in male (but not female) feathers augment male appearance in two ways.
First, wide, oblong barbs in bright red and orange male feathers enhance light-pigment interactions to create
a more vivid, saturated color from structure alone without requiring more pigment. Second, dihedral
barbules produce (i) low-reflectance super black near colorful patches, a proposed optical illusion for color
emphasis (McCoy et al. 2018, 2019; McCoy and Prum 2019)), and (ii) velvety red feathers adjacent to
brilliantly reflective beaks (Figure 1). We observe that structures contribute significantly to color signal
appearance, rather than pigment, which is inconsistent with traditional carotenoids-as-honest-signals
theory.
Why did male feathers evolve elaborate microstructures? If percepts of color influence female
choices of mates, whether the percept is an arbitrary preference (Prum 2012) or an index of male quality,
then males will be selected to produce this percept in females whether this is by changes to the chemical
composition or microstructure of their feathers. From this perspective, the evolutionary dynamics of female
preferences and male traits has aspects of an arms race between the conflicting interests of the sexes
(Dawkins and Krebs 1978; Krebs and Dawkins 1984; Burk 1988; Hill 1994; Funk and Tallamy 2000).
Females establish tests of quality and males are selected to pass these tests whether by 'honest' displays or
'gaming' the test. Rather than eating and metabolizing more carotenoids to appear more intensely red, males
could use microstructural amplifiers to make their red plumage appear brighter and more saturated. Perhaps
red carotenoid-based colors were favored by female selection because they were correlated with male
quality (Olson and Owens 1998), but selection on males to appear attractive by whatever means gradually
degrades the information about male quality conveyed by the male signal (Hill 1994). As males exaggerate
their appearance, females are selected to develop additional quality tests to separate the wheat from the
chaff (e.g., females may prefer redder feathers or additional ornaments). Because females cannot
unilaterally abandon a prior quality test without dooming their sons to being unattractive, male ornaments
may pile atop one another, potentially causing extreme and elaborate signals. Signals cannot be pigeonholed
as purely honest nor purely deceptive, because we are always viewing a snapshot in evolutionary time
wherein species are subject to many competing selective pressures.
559
Pigments: males and females within a species have the same carotenoid profiles
We found a surprising 29 distinct carotenoid molecules in the plumages of *Ramphocelus* tanagers
(Figure 3), some of which are dietary carotenoids- consumed and directly deposited into feathers- while
others are metabolized. If carotenoids were present in feathers as an honest (because costly) signal, it
would be surprising to see the same levels in females (the choosing sex) as in males (the displaying sex).
Unexpectedly, males and females of the same species had similar carotenoid profiles in feathers, with
similar levels and types in both sexes (Figure 2). A correlation between male and female signals despite
dimorphism can result from general constraint and opportunities of a shared genome (Lassance 2009). For
example, female club-winged manakins (*Machaeropterus deliciosus*, Pipridae) select males who have
altered wing feathers and bones in order to produce musical sound; oddly, the females have modified
bones as well, which may hurt their adaptive survival (Prum 2017). Male and female tanagers have been
able to evolve different feather structures so it is difficult to argue that the presence of carotenoids in

feathers is costly, is favored in males because it is costly, but females who gain no benefit therefrom have
been unable to exclude carotenoids from their feathers.
However, it is still possible that carotenoids are an index of metabolic function, rather than an
example of a costly investment or trade-off (Weaver et al. 2017). It is worth dedicating a few sentences to
the significance of dietary versus metabolized carotenoids. Dietary carotenoids are typically yellow, while
metabolized carotenoids are redder. A recent meta-analysis showed that metabolized, rather than dietary,
carotenoids drive correlations between male appearance and health (Weaver et al. 2018). Perhaps red
metabolized carotenoids are thus a signal of a functioning metabolism. However, in *Ramphocelus* it
appears that rumped tanagers evolved to use dietary carotenoids (yellow & orange) from a likely ancestral
state of metabolized (red) carotenoids (Figure 3, Figure S1; this is the most parsimonious explanation,
particularly given that the *Ramphocelus* clade has red-colored sister species (Burns and Racicot 2009)). In
fringillid finches, a model system for plumage evolution, there are no documented reversions from
metabolized to dietary pigments (Ligon et al. 2016). New world blackbirds (Icteridae) convergently
evolved red plumage from yellow six times but never in the reverse direction (Friedman et al. 2011).
Reversions to yellow seem rare (i.e., less likely than gains of red), and our tree may indicate two
independent origins of red rather than one loss. However, our data demonstrates that yellow and orange
birds do have some red metabolized pigments (Figure 3), just in lower quantities than in the red birds, and
are thus capable of the metabolic processes that convert dietary to metabolized pigments. The possible
reversion from red to yellow and orange documented herein could be evidence that *Ramphocelus* males
were not selected to display metabolically indicative or costly traits through carotenoid pigmentation.
Further research should be done on the behavioural ecology of *Ramphocelus* to clarify female
preferences, particularly regarding dietary (yellow/orange) versus metabolized (orange/red) pigment
signals. In a hybrid zone, Morales-Rozo and colleagues demonstrate that the pigment-poor, yellow *R. f.*
icteronotus outcompetes the comparatively pigment-rich, orange *R. flammigerus* in areas where both male
phenotypes occur (i.e., where females have a choice). Specific mate-choice experiments would be needed
to demonstrate this conclusively, but if females do indeed prefer this less-elaborate, less-costly trait, this
aligns well with Hill's criteria for an arbitrary, rather than honest, model of trait evolution: Hill predicted
that under honest advertisement models, "in lineages in which a less elaborate form of a trait is derived
from a more elaborate form, females should show a preference for the primitive, extreme expression of the
trait" (Hill 1994).
In addition to the pigments we document, the relative amounts of other pigment types, such as
melanin, or internal nanostructural changes likely also play a role in the different appearances of males and
females. This is a worthwhile focus of future studies, as well as the physiological cost or importance of
other pigments and nanostructures.
606
Structures: microstructures amplify plumage appearance in males
Feathers in *Ramphocelus* males, but not females, vary remarkably in microscopic structure (Table
1, Figure 5, Figure S5). Relative to the typically-proportioned female feathers, many males have extremely
expanded barbs and strap-shaped, upward-angled barbules. It is microstructure, not carotenoid pigment,
that differs between males and females (Figure 6). With optical modelling, we show that expanded barbs
enhance color and upward-angled barbules generate velvety red and super black (Figure 8)—a deep black
which frames and amplifies nearby color. That is, structures alone enhance appearance independently of
pigment content.
Oblong, expanded central barbs enhance color because they increase the optical power transmitted
through the feather (Figure 8B-C). Male feathers have more oblong and more expanded barbs (Table 1).
More light interacts with the same amount of pigment to generate a more saturated, vivid color in males
(Figures 1-2). This is analogous to a pigmented solution in a glass test tube, which appears more-richly-
colored from certain angles due to the differences in path length of light through the solution.
Upward-angled barbs in a dihedral arrangement generate a velvety, super black appearance
(Figure 8A)—an optical illusion that enhances nearby colors. These "super black" plumage patches
reflect almost no 90° incident light and have a broadband, featureless reflectance spectrum in contrast to a

rise in reflectance at higher wavelengths characteristic of melanin pigmentary reflection alone (Figure
2A). Normally, vertebrates use white gleams (mirror-like specular reflections) to control for the quantity
of ambient light, thus accurately perceiving color across many lighting conditions. Super black enhances
the perceived brilliance of adjacent colors by removing nearly all specular reflection, thus influencing the
innate control capacities of vertebrate eyes and brains (Kreezer 1930; Brainard et al. 1993; Speigle and
Brainard 1996; McCoy et al. 2018). In the intriguing case of *R. carbo*, velvety red plumage shares its
microstructure (and reflectance curve) with super black plumages (Figure S6); this velvet red, absorptive
plumage is adjacent to an extremely bright white bill (Figure 1D). This dark, antireflective frame would
thus enhance the perceived brilliance of the "silver" beak, which may play an important role in female
choice.
Further evidence for the importance of microstructure in color appearance comes from within-bird
plumage variation. Within single birds that have both bright red and dark velvety red plumage (e.g., upper
throat versus lower throat in both *R. dimidiatus* and *R. melanogaster*), the relative amount of pigment
molecules in bright and dark red patches are highly correlated (Figure 7A,D), suggesting that pigment alone
is not responsible for these differences in appearance. By contrast, the feather microstructure is strikingly
different between these types of patches (Figure 7B-C,E-F), suggesting that microstructure, not pigment, is
responsible for the differences in appearance of bright and dark velvety plumage. These feathers may also
have internal nanostructural changes, which have been shown to make carotenoid-based colors brighter
(Shawkey and Hill 2005). Research on microstructural variation in colorful displays, including sex
differences, is expanding rapidly (Lee et al. 2009; Iskandar et al. 2016; Enbody et al. 2017; McCoy et al.
2018; McCoy and Prum 2019). To gain more insight into evolutionary dynamics, we require a complete
understanding of the physical basis of color. This often means accounting for the optical effects of
microstructure.
646
An arms race: honest signals can be gamed, causing trait elaboration
Microstructural modifications allow male *Ramphocelus* tanagers to amplify (Hasson 1989) their
carotenoid display independent of the amount of carotenoids consumed and metabolized (Olson 1970; Hill
1994). Mutant melanic scarlet tanagers (*Piranga olivacea*, Cardinalidae) retain the feather structure typical
of carotenoid-containing feathers, suggesting that at least some birds have independent genetic control
systems for pigment deposition and feather structure (Brush 1970). Thus, it is possible for selection to act
on feather structure without impacting pigment. Male *Ramphocelus* tanagers may have been selected for
structural modifications which alter their visual appearance and distort the link between appearance and
carotenoid content.
Mate choice is akin to an examination. Under honest signaling theory, female preference is
selected to gather information about the quality of a potential mate: she establishes an examination. Males
are selected to pass her test whether or not their performance correlates with intrinsic quality (McCoy and
Haig, unpublished manuscript). If the male examinee is indeed of high quality, both parties benefit when
this information is conveyed accurately, but, if the examinee is of poor quality, his best interest may be to
cheat (Haig 2010). The moment a quality test is established, selection favors individuals who can pass the
test. Direct evidence for deceptive elements to honest signaling has been reported in many invertebrates
(e.g., Wilson et al. 2007; Christy and Rittschof 2010; Ghislandi et al. 2017) and a few vertebrates
(Candolin 1999; Bro-Jørgensen and Pangle 2010; Summers 2014). Goodhart's Law in economics and the
analogous Campbell's Law describe this phenomenon: pithily phrased, "when a measure becomes a
target, it ceases to be a good measure" (Strathern 1997). The classic case is high-stakes testing. When
standardized test scores are used as a proxy for how well middle school teachers educate their pupils,
determining funding, teachers 'teach to the test' rather than improving students' reading skills (Goldstein
2004). This does not mean tests are useless: for example, the SAT does convey some true information—
but this signal is distorted by socioeconomic background, teacher strategy, culture, and test prep
programs.
What about the testing bed of mate selection in birds? Many years of research have relied upon
hypotheses that sexual signals are honest; that is, honest correlations between signal traits and individual

quality primarily drive the evolution of beautiful signals. But many have also questioned how such a model
could explain the maintenance and, particularly, *elaboration* of honest traits (Hill 1994; Weaver et al. 2017).
A model of sexual selection that allows for deception can better explain trait elaboration (Krebs and
Dawkins 1984; Hill 1994; Haig 2010). Hill (1994) proposed that the elaboration of sexually selected traits
follows a model of sexual conflict. He writes, "…female demand for display characters that reliably signal
male quality leads to selection for male strategies to express these display characters independent of quality.
The result can be an evolutionary race leading to the elaboration of male characteristics while maintaining
a correlation between character expression and male quality." Additionally, females cannot abandon test
criteria because then their preferences will be in the minority (put simply, their sons will not be sexy (Fisher
1999)). To preserve honest tests of quality, females may either raise the bar (feathers must be 25% redder)
or establish new criteria while retaining old criteria; thus proceeds an arms race. Herein, we find support
for this hypothesis; microstructure and an associated optical illusion (super black) enhance color
independent of carotenoid content. Whatever criteria females establish, males will be selected to satisfy
female preferences— by hook or by crook.

688
689

Author Contributions: All authors conceived the research design. DEM and EH performed all pigment
extractions, SEM work, and spectrophotometry. CV and ST designed and implemented the LC-MS
procedure for carotenoid characterization. DEM performed the optical simulations. AJS constructed the
phylogeny and performed all regressions. DH provided conceptual groundwork. All authors jointly wrote
the paper.

Acknowledgements:
We are very grateful to Jeremiah Trimble and Kate Eldridge for their help with the ornithological
collections. Special thanks to Dan Utter and Professor Colleen Cavanaugh for their methodological
assistance and use of their lab equipment. Thank you to Sofia Prado-Irwin and Dan Utter for helpful
comments on the manuscript. Thank you to Gabriel Ugueto for the outstanding artwork, and to Nick
Athanas – antpitta.com – for the beautiful photos. This work was performed in part at the Harvard
University Center for Nanoscale Systems (CNS), a member of the National Nanotechnology Coordinated
Infrastructure Network (NNCI), which is supported by the National Science Foundation under NSF
ECCS award no. 1541959. DEM's research is conducted with Government support under and awarded by
DoD, Air Force Office of Scientific Research, National Defense Science and Engineering Graduate
(NDSEG) Fellowship, 32 CFR 168a, and DEM and AJS were supported by research funds from the
Harvard University Department of Organismic and Evolutionary Biology (OEB).

708
709 **Conflict of Interest Statement:** The authors declare no conflicts of interest.

710
711 **Data Accessibility Statement:** All code and data will be made available on Dryad.

712
713
714
715
<u>References</u>

Brainard, D. H., B. A. Wandell, and E.-J. Chichilnisky. 1993. Color constancy: from physics to
appearance. Curr. Dir. Psychol. Sci. 165–170. JSTOR.
Bro-Jørgensen, J., and W. M. Pangle. 2010. Male Topi Antelopes Alarm Snort Deceptively to Retain
Females for Mating. Am. Nat. 176:E33–E39.
Brush, A. H. 1970. Pigments in hybrid, variant and melanic tanagers (birds). Comp. Biochem. Physiol.
36:785–793. Elsevier.

Burk, T. 1988. Acoustic signals, arms races and the costs of honest signalling. Florida Entomol. 71:400.

Burns, K. J., and R. A. Racicot. 2009. Molecular Phylogenetics of a Clade of Lowland Tanagers:
Implications for Avian Participation in the Great American Interchange. Auk 126:635–648. BioOne.

Burns, K. J., and A. J. Shultz. 2012. Widespread cryptic dichromatism and ultraviolet reflectance in the
largest radiation of Neotropical songbirds: implications of accounting for avian vision in the study
of plumage evolution. Auk 129:211–221. BioOne.

Burns, K. J., A. J. Shultz, P. O. Title, N. A. Mason, F. K. Barker, J. Klicka, S. M. Lanyon, and I. J.
Lovette. 2014. Phylogenetics and diversification of tanagers (Passeriformes: Thraupidae), the largest
radiation of Neotropical songbirds. Mol. Phylogenet. Evol. 75:41–77. Elsevier Inc.

Candolin, U. 1999. The relationship between signal quality and physical condition : is sexual signalling
honest in the three-spined stickleback ? 1261–1267.

Christy, J. H., and D. Rittschof. 2010. Deception in visual and chemical communication in crustaceans.
Pp. 313–333 *in* Chemical communication in crustaceans. Springer.

Clements, J. F., T. S. Schulenberg, M. J. Iliff, D. Roberson, T. A. Fredericks, B. L. Sullivan, and C. L.
Wood. 2017. The eBird/Clements checklist of birds of the world: v2017.

Dawkins, M. S., and T. Guilford. 1996. Sensory bias and the adaptiveness of female choice. Am. Nat.
937–942. JSTOR.

Dawkins, R., and J. R. Krebs. 1978. Animal signals: information or manipulation. Behav. Ecol. An Evol.
approach 2:282–309.

Enbody, E. D., S. M. Lantz, and J. Karubian. 2017. Production of plumage ornaments among males and
females of two closely related tropical passerine bird species. Ecol. Evol. 7:4024–4034. Wiley
Online Library.

Faivre, B., A. Grégoire, M. Préault, F. Cézilly, and G. Sorci. 2003. Immune activation rapidly mirrored in
a secondary sexual trait. Science (80-.). 300:103. American Association for the Advancement of
Science.

Felsenstein, J. 1985. Phylogenies and the Comparative Method. Am. Nat. 125:1–15. JSTOR.

Fisher, R. A. 1999. The genetical theory of natural selection: a complete variorum edition. Oxford
University Press.

Folstad, I., and A. J. Karter. 1992. Parasites, Bright Males, and the Immunocompetence Handicap. Am.
Nat. 139:603–622.

Friedman, N. R., L. M. Kiere, and K. E. Omland. 2011. Convergent gains of red carotenoid-based
coloration in the new world blackbirds. Auk 128:1–10.

Funk, D. H., and D. W. Tallamy. 2000. Courtship role reversal and deceptive signals in the long-tailed
dance fly, Rhamphomyia longicauda. Anim. Behav. 59:411–421. Citeseer.

Ghislandi, P. G., M. Beyer, P. Velado, and C. Tuni. 2017. Silk wrapping of nuptial gifts aids cheating
behaviour in male spiders. Behav. Ecol. 28:744–749.

Gkikas, D., A. Argiropoulos, and S. Rhizopoulou. 2015. Epidermal focusing of light and modelling of
reflectance in floral-petals with conically shaped epidermal cells. Flora Morphol. Distrib. Funct.
Ecol. Plants 212:38–45.

Goldstein, H. 2004. Education for all: The globalization of learning targets. Comp. Educ. 40:7–14.

Gorton, H. L., and T. C. Vogelmann. 1996. Effects of epidermal cell shape and pigmentation on optical
properties of Antirrhinum petals at visible and ultraviolet wavelengths. Plant Physiol. 112:879–888.
Am Soc Plant Biol.

Grafen, A. 1989. The phylogenetic regression. Phil. Trans. R. Soc. B 326:119–157.

Griscom, L. 1932. Notes on imaginary species of Ramphocelus. Auk 49:199–203. JSTOR.

Hackett, S. J. 1996. Molecular phylogenetics and biogeography of tanagers in the genus Ramphocelus
(Aves). Mol. Phylogenet. Evol. 5:368–382.

Haig, D. 2010. Fertile soil or no man's land: cooperation and conflict in the placental bed. Pp. 165–173 *in*
R. Pijnenborg, I. Brosens, and R. Romero, eds. Placental Bed Disorders: Basic Science and its
Translation to Obstetrics. Cambridge University Press, Cambridge.

Harris, E. 1987. Breeding, Husbandry and Management of Tanager Species. AFA Watchb. 14:20–27.

Harvey, T. A., K. S. Bostwick, and S. Marschner. 2013. Directional reflectance and milli-scale feather morphology of the African Emerald Cuckoo, *Chrysococcyx cupreus*. J. R. Soc. Interface 10:20130391. The Royal Society.

Hasson, O. 1989. Amplifiers and the handicap principle in sexual selection: a different emphasis. Proc. R. Soc. London B Biol. Sci. 235:383–406. The Royal Society.

Hill, G. E. 2006. Female mate choice for ornamental coloration. Bird Color. 2:137–200.

Hill, G. E. 2015. Sexiness, Individual Condition, and Species Identity: The Information Signaled by Ornaments and Assessed by Choosing Females. Evol. Biol. 42:251–259.

Hill, G. E. 1994. Trait elaboration via adaptive mate choice: sexual conflict in the evolution of signals of male quality. Ethol. Ecol. Evol. 6:351–370. Taylor & Francis.

Hill, G. E., and K. J. McGraw. 2006. Bird coloration: mechanisms and measurements. Harvard University Press, Cambridge.

Hill, G. E., and R. Montgomerie. 1994. Plumage colour signals nutritional condition in the house finch. Proc. R. Soc. B Biol. Sci. 258:47–52.

Iskandar, J.-P., C. M. Eliason, T. Astrop, B. Igic, R. Maia, and M. D. Shawkey. 2016. Morphological basis of glossy red plumage colours. Biol. J. Linn. Soc. 119:477–487. Oxford University Press.

Katoh, K., and D. M. Standley. 2013. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Mol. Biol. Evol. 30:772–780.

Kay, Q. O. N., H. S. Daoud, and C. H. Stirton. 1981. Pigment distribution, light reflection and cell structure in petals. Bot. J. Linn. Soc. 83:57–83. Oxford University Press.

Koch, R. E., A. N. Kavazis, D. Hasselquist, W. R. Hood, Y. Zhang, M. B. Toomey, and G. E. Hill. 2018. No evidence that carotenoid pigments boost either immune or antioxidant defenses in a songbird. Nat. Commun. 9:491. Nature Publishing Group.

Krebs, J. R., and R. Dawkins. 1984. Animal signals: mindreading and manipulation. Behav. Ecol. an Evol. approach 380–402. Blackwell Scientific Publication.

Kreezer, G. 1930. Luminous appearances. J. Gen. Psychol. 4:247–281. Taylor & Francis.

Krueger, T. R., D. A. Williams, and W. A. Searcy. 2008. The genetic mating system of a tropical tanager. Condor 110:559–562. BioOne.

LaFountain, A. M., R. O. Prum, and H. A. Frank. 2015. Diversity, physiology, and evolution of avian plumage carotenoids and the role of carotenoid–protein interactions in plumage color appearance. Arch. Biochem. Biophys. 572:201–212. Elsevier.

Lassance, J. M. 2009. Concerted evolution of male and female display traits in the European corn borer, Ostrinia nubilalis. BMC Biol. 7:10.

Lee, E., M. Aoyama, and S. Sugita. 2009. Microstructure of the feather in Japanese Jungle Crows (Corvus macrorhynchos) with distinguishing gender differences. Anat. Sci. Int. 84:141–147. Springer.

Ligon, R. A., R. K. Simpson, N. A. Mason, G. E. Hill, and K. J. McGraw. 2016. Evolutionary innovation and diversification of carotenoid-based pigmentation in finches. Evolution (N. Y). 70:2839–2852.

Martins, E. P., and T. F. Hansen. 1997. Phylogenies and the comparative method: a general approach to incorporating phylogenetic information into the analysis of interspecific data. Am. Nat. 149:646–667.

Maurer, D. L., T. Kohl, and M. J. Gebhardt. 2017. Cuticular microstructures turn specular black into matt black in a stick insect. Arthropod Struct. Dev. 46:147–155. Elsevier.

McCoy, D. E., T. Feo, T. A. Harvey, and R. O. Prum. 2018. Structural absorption by barbule microstructures of super black bird of paradise feathers. Nat. Commun. 9:1–8. Nature Publishing Group.

McCoy, D. E., V. E. McCoy, N. K. Mandsberg, A. V Shneidman, J. Aizenberg, R. O. Prum, and D. Haig. 2019. Structurally assisted super black in colourful peacock spiders. Proc. R. Soc. B 286:20190589. The Royal Society.

McCoy, D. E., and R. O. Prum. 2019. Convergent evolution of super black plumage near bright color in 15 bird families. J. Exp. Biol. 222:jeb208140. The Company of Biologists Ltd.

Morrison, E. S., and A. V Badyaev. 2016. Structuring evolution: biochemical networks and metabolic

diversification in birds. BMC Evol. Biol. 16:168. BioMed Central.

Moynihan, M. 1962. Display Patterns of Tropical American" Nine-Primaried" Songbirds II. Some
Species of Ramphocelus. Auk 79:655–686. JSTOR.

Olson, S. L. 1970. Specializations of some carotenoid-bearing feathers. Condor.

Olson, V. A., and I. P. F. Owens. 1998. Costly sexual signals: Are carotenoids rare, risky or required?

Prum, R. O. 2012. Aesthetic evolution by mate choice: Darwin's really dangerous idea. Philos. Trans. R.
Soc. B Biol. Sci. 367:2253–2265. The Royal Society.

Prum, R. O. 2017. The Evolution of Beauty: How Darwin's Forgotten Theory of Mate Choice Shapes the
Animal World-and Us. Doubleday.

Revell, L. J. 2012. phytools: An R package for phylogenetic comparative biology (and other things).
Methods Ecol. Evol. 3:217–223.

Revell, L. J. 2013. Two new graphical methods for mapping trait evolution on phylogenies. Methods
Ecol. Evol. 4:754–759.

Saks, L., K. McGRAW, and P. Hõrak. 2003. How feather colour reflects its carotenoid content. Funct.
Ecol. 17:555–561. Wiley Online Library.

Shawkey, M. D., and G. E. Hill. 2005. Carotenoids need structural colours to shine. Biol. Lett. 1:121–
124.

Shultz, A. J., and K. J. Burns. 2017. The role of sexual and natural selection in shaping patterns of sexual
dichromatism in the largest family of songbirds (Aves: Thraupidae). Evolution (N. Y). Wiley Online
Library.

Simons, M. J. P., A. A. Cohen, and S. Verhulst. 2012. What does carotenoid-dependent coloration tell?
Plasma carotenoid level signals immunocompetence and oxidative stress state in birds-a meta-
analysis. PLoS One 7.

Speigle, J. M., and D. H. Brainard. 1996. Luminosity thresholds: Effects of test chromaticity and ambient
illumination. JOSA A 13:436–451. Optical Society of America.

Stamatakis, A. 2006. RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with
thousands of taxa and mixed models. Bioinformatics 22:2688–2690.

Stamatakis, A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics 30:1312–1313.

Strathern, M. 1997. 'Improving ratings': audit in the British University system. Eur. Rev. 5:305–321.
Cambridge University Press.

Summers, K. 2014. Sexual conflict and deception in poison frogs. Curr. Zool. 60:37–42. Oxford
University Press Oxford, UK.

Taflove, A., and S. C. Hagness. 2005. Computational electromagnetics: the finite-difference time-domain
method. Artech House, Norwood.

Vukusic, P., J. R. Sambles, and C. R. Lawrence. 2004. Structurally assisted blackness in butterfly scales.
Proc. R. Soc. London. Ser. B Biol. Sci. 271:S237–S239. The Royal Society.

Weaver, R. J., R. E. Koch, and G. E. Hill. 2017. What maintains signal honesty in animal colour displays
used in mate choice? Phil. Trans. R. Soc. B 372:20160343. The Royal Society.

Weaver, R. J., E. S. A. Santos, A. M. Tucker, A. E. Wilson, and G. E. Hill. 2018. Carotenoid metabolism
strengthens the link between feather coloration and individual quality. Nat. Commun. 9:73. Nature
Publishing Group.

Wilson, R. S., M. J. Angilletta Jr, R. S. James, C. Navas, and F. Seebacher. 2007. Dishonest signals of
strength in male slender crayfish (Cherax dispar) during agonistic encounters. Am. Nat. 170:284–
291. The University of Chicago Press.

Wilts, B. D., P. J. Rudall, E. Moyroud, T. Gregory, Y. Ogawa, S. Vignolini, U. Steiner, and B. J. Glover.
2018. Ultrastructure and optics of the prism-like petal epidermal cells of Eschscholzia californica
(California poppy). New Phytol. 219:1124–1133. Wiley Online Library.

Zahavi, A. 1975. Mate selection—a selection for a handicap. J. Theor. Biol. 53:205–214. Elsevier.

875