Genetic relatedness cannot explain social bonds in black-and-white ruffed lemurs (*Varecia variegata*).

Andrea L. Baden[1,2,3*], Timothy H. Webster[4*], Brenda J. Bradley[5]

[1]Department of Anthropology, Hunter College, New York, NY 10065

[2]The Graduate Center of the City University of New York, New York, NY 10065

[3]The New York Consortium in Evolutionary Primatology (NYCEP)

[4]Department of Anthropology, University of Utah, Salt Lake City, UT 84112

[5] Department of Anthropology, The George Washington University, Washington DC

*Corresponding authors: Andrea L. Baden, andrea.baden@hunter.cuny.edu, Hunter College, Department of Anthropology, 695 Park Avenue, New York, NY 10065, +001-212-396-6818; Timothy H. Webster, timothy.h.webster@utah.edu, Department of Anthropology, University of Utah, 260 S Central Drive, Carolyn and Kem Gardner Commons, Suite 4625, Salt Lake City, UT 84112

Word count: 8,258

ABSTRACT

Several primates exhibit fission-fusion social dynamics, yet the factors shaping variable associations among subgroup members have not been widely addressed. Associations may occur simply because of shared habitat preferences; however, social ties may also be influenced by genetic relatedness (kinship) or social attraction. Here, we investigate the association patterns of wild black-and-white ruffed lemurs, *Varecia variegata*, in Ranomafana National Park, Madagascar using behavioural, spatial (home range), and genetic data from twenty-four individually identified animals. We collected 40,840 records of group composition over a 17-month period and from this calculated pairwise association indices. We also used ranging coordinates and genetic samples to estimate patterns of spatial overlap and kinship, and then related these measures to patterns of affiliation using Mantel tests and MRQAP. From these analyses, we found that dyadic ruffed lemur social associations were generally sparse and weak; that home range overlap was minimal; and that average relatedness within the community was low. We found no evidence that kinship was related to patterns of either spatial overlap or social association; instead, associations were primarily driven by space use. While home range overlap explained most of the variation seen in social association, some variation remains unaccounted for, suggesting that other social, ecological, and biological factors such as shared resource defense or communal breeding might also play a role in social attraction among members of this species.

Keywords: social network analysis, range use, spatial overlap, Association Index, kinship, fission-fusion, microsatellites, primates

INTRODUCTION

Animal social systems reflect non-random relationships among neighbouring conspecifics, the content, quality, and patterning of which define their social structure (Hinde, 1976). In fact, there is growing evidence that taxa as diverse as fishes, lizards, birds, cetaceans, and equids exhibit social associations that are not only non-random, but highly structured (Augusto, Frasier, & Whitehead, 2017; Croft et al., 2012, 2005; Firth et al., 2017; Spiegel, Sih, Leu, & Bull, 2018; Stanley, Mettke-Hofmann, Hager, & Shultz, 2018). Individuals vary in the numbers, strengths, and stabilities of their social ties (Croft et al., 2005; McDonald, 2007; Silk, Alberts, & Altmann, 2006; Silk, Altmann, & Alberts, 2006; Silk et al., 2010a; Stanley et al., 2018), as well as in their preferences for social associates (e.g., Cords, 2002; Gero, Gordon, & Whitehead, 2015; Kohn, Meredith, Magdaleno, King, & West, 2015; Mourier, Vercelloni, & Planes, 2012; Perry, 2012; Schülke, Bhagavatula, Vigilant, & Ostner, 2010)—outcomes of which can be evolutionarily significant (Seyfarth & Cheney, 2012; Silk, 2007). For example, the extent and/or strength of females' social connections have been positively linked to higher fertility (*Macaca mulatta*: Brent et al., 2013; humans: Balbo & Barban, 2014), longer lifespan (*Papio cynocephalus*: Alberts, 2019; Archie, Tung, Clark, Altmann, & Alberts, 2014; *Papio ursinus:* Silk et al., 2010b), and greater offspring survival (*Equus caballus*: Cameron, Setsaas, & Linklater, 2009; *Papio*

cynocephalus: Silk, Alberts, & Altmann, 2003; *Papio ursinus:* Cheney, Silk, & Seyfarth, 2016; Silk et al., 2009; *Physeter microcephalus*: Whitehead, 1996). Similarly, males that form strong social bonds have been shown to be more successful at forming coalitions, achieving high rank, and siring more offspring than males with weaker social bonds (e.g., *Chiroxiphia linearis*: McDonald, 2007; *Macaca assamensis*: Schülke et al., 2010; *Pan troglodytes*: Gilby et al., 2013). Nevertheless, while the long-term health and fitness outcomes of social bonds have been the focus of much investigation, the proximate factors shaping social preferences have received comparatively little attention.

Several hypotheses have been proposed to explain social preference, the most popular being kinship, whereby group members show preferential associations with and, in the case of kin selection, are more likely to direct costly altruistic behaviours toward close genetic relatives (Hamilton, 1964). For instance, many female mammals exhibit social preferences toward maternal kin (Smith, 2014), a pattern observed in primates in particular (reviewed in Langergraber, 2012), but also in spotted hyaenas (Wahaj et al., 2004), African elephants (Archie, Moss, & Alberts, 2006), dolphins (Frère et al., 2010), and mountain goats (Godde, Côté, & Réale, 2015), among others. Social preferences toward kin can be driven by the benefits of associating with relatives, such as allomaternal care (reviewed in Briga, Pen, & Wright, 2012; Pope, 2000; and in some cases even brood parasitism, Andersson, 2017), reduced aggression and infanticide risk (reviewed in Brown & Brown, 1996; Agrell, Wolff, & Ylonen, 1998), foraging advantages (Griffiths & Armstrong, 2002; Nystrand, 2007), and shared social and ecological

knowledge (McComb, Moss, Durant, Baker, & Sayialel, 2001; Salpeteur et al.,

2015). However, social integration has also been shown to have strong direct

fitness consequences between nonrelatives (Baden, Wright, Louis, & Bradley,

2013; Cameron et al., 2009; G. G. Carter & Wilkinson, 2015; Riehl, 2011),

suggesting that social bonds can evolve based on these direct benefits alone.

Moreover, evidence for kin-biased relationships in other species, particularly those

with higher fission-fusion dynamics, is weak (e.g., Best, Dwyer, Seddon, &

Goldizen, 2014; K. D. Carter, Seddon, Frère, Carter, & Goldizen, 2013; Hirsch,

Prange, Hauver, & Gehrt, 2013; Langergraber, Mitani, & Vigilant, 2007, 2009;

Moscovice et al., 2017; Wilkinson, Carter, Bohn, & Adams, 2016), raising

questions as to the ubiquity of kinship in the formation and maintenance of social

ties.

One complicating factor in determining whether and how kinship shapes

social bonds is that it can be difficult to decouple social associations due to kinship

from associations due to other social and spatial contexts (reviewed in Doreian &

Conti, 2012; Wey, Blumstein, Shen, & Jordán, 2008). For example, in some

species it is difficult to differentiate whether strong associations are due to strong

genetic ties or to other factors, such as sex or age (Lusseau & Newman, 2004;

Mourier et al., 2012), habitat utilization (Wiszniewski, Allen, & Möller, 2009), or

foraging specializations (Daura-Jorge, Cantor, Ingram, Lusseau, & Simões-Lopes,

2012; Gilby & Wrangham, 2008; Griffiths & Armstrong, 2002; Mitani, Merriwether,

& Zhang, 2000). Indeed, patterns of social association have been found to

correlate with spatial overlap (i.e., overlapping home ranges) more strongly than,

or to the exclusion of, kinship in a number of species (Best et al., 2014; K. D. Carter

et al., 2013; Frère et al., 2010; Strickland, Gardiner, Schultz, & Frère, 2014).

Moreover, although individuals living in closer physical proximity are more prone

to interact (e.g., Clutton-Brock, 1989; Kossinets & Watts, 2006), conspecifics with

high levels of spatial overlap may also mediate their interactions by temporally

modifying range use, such that animals with highly overlapping ranges may

interact minimally, if at all (e.g., temporal avoidance) (Leu, Bashford, Kappeler, &

Bull, 2010; Ramos-Fernández, Boyer, Aureli, & Vick, 2009; Strickland et al., 2017).

While still rare, studies that consider the simultaneous effects of spatial overlap

and kinship on association patterns are thus important to understanding the role

of social preference in the evolution of social systems and sociality (e.g., Best et

al., 2014; K. D. Carter et al., 2013; Frère et al., 2010; Lusseau et al., 2006; Maher,

2009; Podgórski, Lusseau, Scandura, Sönnichsen, & Jędrzejewska, 2014;

Strickland et al., 2014).

Ruffed lemurs (Genus *Varecia*) are moderately-sized frugivores (Baden,

Brenneman, & Louis Jr., 2008; Balko & Underwood, 2005) that live in groups

(hereafter "communities") comprising as many as 30 adult and subadult individuals

and their offspring (Baden, Webster, & Kamilar, 2016 and references therein).

From earlier work, we know that communities exhibit high fission-fusion dynamics,

with members varying in their degrees of home range overlap and number and

strength of social ties (Baden, 2011; Baden & Gerber, n.d.; Baden et al., 2016;

Vasey, 2006). Moreover, studies of microsatellite markers and mitochondrial DNA

have shown that while average pairwise relatedness within communities is low

(i.e., both sexes disperse: Baden et al., 2014), communities nevertheless contain

networks of closely related kin (e.g, Baden, 2011; Baden et al., 2013). Taken

together, these lines of evidence suggest that community members may be

actively choosing social associates, and that these associations may be shaped

by a number of ecological, social, and biological factors. Here, we investigate the

relative influences of space use and kinship on the strength of associations

amongst members of a black-and-white ruffed lemur (*Varecia variegata*)

community. Specifically, we quantify dyadic measures of association, home range

overlap, and genetic relatedness to address the overarching question: What

influences social association in ruffed lemurs?

METHODS

Study site and subjects

We collected data from one *V. variegata* community at Mangevo [21°220

6000 S, 47°280 000 E], a mid-elevation (660-1,200 m) primary rainforest site

located in the southeastern parcel of Ranomafana National Park, Madagascar

(Wright et al., 2012) during 16 months of study (August–December 2007;

February– December 2008). At the time of the study, the community included 24

adults and subadults (8 adult females, 11 adult males, 5 subadult males). All

individuals in this study were habituated and individually identified via radio-collars

or unique collar-tag combinations prior to behavioural observations (see Baden et

al., 2016 for specific details). Animals were collared under veterinary supervision

following a strict protocol (Glander, 1993). Nineteen infants were born in the 2008

birth season and were present from October to December 2008, when the study

ended. Sampling efforts resulted in a total of 4,044 focal observation hours.

Ethical note

Permission to conduct research was granted by Madagascar's National

Parks (ANGAP/MNP, #084/07-041/08), and research protocols were in

compliance with Stony Brook University IACUC #2005-20081449. All animal

protocols followed the Guidelines for the treatment of animals for teaching and

research recommended by ASAB/ABS (2019).

Data collection

Behavioural monitoring

We collected data during dawn to dusk follows of focal individuals. We

located focal subjects at the beginning of each observation period via radio-

telemetry and selected new subjects daily. We never sampled focal subjects on

consecutive days and every effort was made to follow all subjects at least once per

month. If an individual with a collar-tag was located in association with a radio-

collared focal individual prior to 10:00 h, this individual became the new focal

subject for that observation period. Observational periods ranged in duration

between 8 to 11 hours depending on seasonal differences in day length and time

needed to locate animals at dawn.

Upon initial contact with the focal individual, we recorded the number and

identities of all other individuals present within the subgroup. To do so, one

observer remained with the focal individual while the remaining three team

members spread out to locate and identify all other members of the subgroup. *A*

priori, individuals were considered members of a subgroup only if they were within

50 m of the group center and were seen associating, traveling with and maintaining

proximity to members of the subgroup being followed (see Baden et al., 2016 for

details). After initial contact, we monitored subsequent changes in subgroup size,

composition (age/sex class, individual identity), and cohesion (i.e., the greatest

distance between any two subgroup members), as well as activity state of the focal

subject using instantaneous scan sampling techniques collected at 5 min intervals

(Altmann, 1974). We recorded a total of 40,840 group scans.

We collected simultaneous GPS coordinates at 10-min intervals from as

close to the focal individual as possible to document daily individual range use. We

recorded spatial coordinates only if estimated positional error was less than 10 m.

Genetic sampling

In addition, we collected genetic samples from 38 black-and-white ruffed

lemurs from the Mangevo population, including all subjects in this study, during

five capture seasons spanning four consecutive years (2005-2008; see Baden,

2011 for details). Sample collection occurred under veterinary supervision and

followed a strict protocol outlined by Glander (1993). All capture procedures

occurred during non-reproductive seasons in the absence of infants and dependent offspring.

For each individual captured, one of us (ALB) collected approximately 1 ml/kg of whole blood (~4 cc) from the femoral vein and four 2mm tissue biopsies from ear pinnae. All samples were stored in 5ml of lysis buffer solution (0.1 M Tris-HCl pH, 8.0, 0.1 M EDTA, 0.01 M NaCl, and 0.5% w/v SDS) at ambient temperature in the field (7 to 21 days) (Longmire, Gee, Hardekopf, Graham, & Mark, 1992). We then banked samples in a −80°C freezer at the Madagascar Biodiversity Partnership headquarters in Antananarivo, Madagascar and subsequently at the Yale Molecular Anthropology Lab in New Haven, CT until analysis.

Data analysis

Association strength

Sampling was biased toward radio-collared females; we therefore subsampled our data prior to association analyses. Details are presented in Baden et al. (2016), but briefly, we divided our dataset into monthly periods and according to the sex of the focal individual. Using only scans for which all individuals were known, we randomized point scans and selected X scans to include in the dataset for each sex, where X is 90% of the point scans for the sex with the fewest scans in a given month. This procedure resulted in the inclusion of a total of 11,784 point scans, with equal numbers of scans targeting male and female focal subjects in each month (Baden et al., 2016).

We used SOCPROG 2.9 (Whitehead, 2009) to calculate association indices

between all pairs of individuals using the "simple ratio" index, which quantifies the

proportion of time that two individuals are observed together relative to their total

observation time overall. This index is appropriate when individuals are equally

likely to be correctly identified (Cairns & Schwager, 1987; Ginsberg & Young, 1992;

Whitehead, 2008), which is the case for this population. Given that subgroup

composition in this community changes approximately every 90 minutes (Baden

et al., 2016), we used a 6-hour sampling interval to reduce autocorrelation among

periods. We also removed any subject observed in fewer than 10 sampling

periods. Our final dataset included 8 adult females and 9 adult males observed in

over 11,171 point scans.

To examine seasonal variation in associations, we organized the subsetted

data from 2008 into three seasons according to female reproductive state as

defined by Baden et al. (2013). Using the protocols described above, our final

datasets included: 8 females and 9 males over 5,051 scans during the

nonbreeding season (January through June); 7 females and 8 males over 3,267

scans during the mating/gestation season (July through September); and 8

females and 6 males over 2,664 scans during the birth/lactation season (October

through December).

Spatial overlap

We calculated home range overlap between all pairs of individuals using a

utilization distribution overlap index (UDOI; Fieberg & Kochanny, 2005)

implemented in the R package adehabitat (Calenge, 2011), as described in Baden

& Gerber (n.d.). The UDOI is an index of space-use sharing between two utilization

distributions (UDs). UDOI values can range from 0 to 1, with a UDOI of 0 indicating

no home range overlap and a UDOI of 1 indicating that home ranges are uniformly

distributed and have 100% overlap. Values can also be >1 if both UDs are

nonuniformly distributed and also have a high degree of overlap. Values <1

indicate less overlap relative to uniform space, whereas values >1 indicate higher

than normal overlap relative to uniform space. We calculated four UDOIs for all

pairs of individuals: one annual UDOI, and three seasonal UDOIs according to

female reproductive state, as for association analyses described above.

Relatedness

We genotyped individuals at a suite of 15 polymorphic microsatellite loci (see

Baden, 2011 and Baden et al., 2014 for details). We extracted total genomic DNA

from blood and/or tissue samples using standard nucleic acid extraction kits

(QIAamp DNA Mini Kit; Qiagen) automated on a QiaCube (Qiagen). Extraction

procedures followed the manufacturer's protocols, with the following modification

to the tissue extraction procedures: samples were allowed to lyse initially in ASL

buffer for 24-48 hours rather than 10 minutes.

We carried out PCR amplifications in a total reaction volume of 25 µl

consisting of 2 µl template, 12.5 µl Qiagen HotStar Taq Master Mix, and 10 µM of

each primer. Amplification conditions were as follows: initial denaturation at 95 °C

for 15 min; 35 cycles of 30 s at 94 °C, 40 s at 54 to 60 °C (see Louis et al., 2005),

1 min at 72 °C, and a final extension of 7 min at 72 °C. The 5′ end of the forward

primer was fluorescently labeled, and amplification products were separated and

visualized using capillary electrophoresis (ABI 3730xl Genetic Analyzer). We

assessed allele sizes relative to an internal size standard (ROX-500) using Gene

Mapper software (Applied Biosystems), and scored final genotypes based on

multiple independent reactions (Taberlet, 1996). Panels yielded PIsib (Queller &

Goodnight, 1989) values of 2.7×10^{-5}, demonstrating the very low probability that

two individuals would share the same multilocus genotype by chance. We further

tested the robusticity of this suite of loci for estimating relatedness with a

rarefaction analysis as in Altmann & Alberts (1996) and de Ruiter & Geffen (1998)

using the program RE-RAT (http://people.musc.edu/~schwaclh/). We estimated

pairwise relatedness among individuals (*r*) following Queller & Goodnight (1989) using

the program GenAlEx 6.5 (Peakall & Smouse, 2012). Relatedness was based on allele

frequencies derived from a larger population of 38 adult multilocus genotypes (Baden,

2011). Fine scale dyadic relatedness assessments (e.g., distinguishing between

full and half-sibs) are not possible in most microsatellite studies, and in fact, the

ability to differentiate relatedness disjunctions on such a scale would probably

require 30 to 60 microsatellite loci (Stone & Björklund, 2001). We therefore

consider "related dyads" to be those with r-values ≥ 0.25; we made no attempt to

further distinguish categories or degrees of relatedness.

Correlations among association strengths, spatial overlap, and relatedness

We used Mantel tests to examine whether association indices were related

to either spatial overlap or relatedness, as well as test for a correlation between

spatial overlap and relatedness (Mantel, 1967). We further used a multiple

regression quadratic assignment procedure (MRQAP; Dekker, Krackhardt, &

Snijders, 2007) to determine relationships between the response variable—

association index—and predictor variables—sex, spatial overlap, and relatedness.

For sex, we used a matrix of sex similarity, which used values of 1 (same sex) and

0 (different sex). MRQAP tests each pairwise combination of response and

predictor matrices while holding the remaining predictor matrices constant. We

performed both Mantel tests and MRQAP for the overall dataset as well as in each

of the three seasons using SOCPROG 2.9 (Whitehead, 2009).

Quantifying relatedness among dyads with "preferred" association strengths or

spatial overlap

In addition to the correlations described above, we further investigated

patterns of kinship among individuals identified as "preferred" associates (AIs) and

those with "preferred" spatial overlap (UDOIs) following methods described in

Frère et al. (2010) and Carter et al. (2013). We first divided relationships into

"preferred" (>95th percentile of association indices or UDOIs) and "non-preferred"

(<=95th percentile). Note that the descriptor "preferred" is clear for associates

(dyads observed together more often than others), but dyads with high spatial

overlap were not necessarily utilizing shared ranges in coordinated ways. Unlike

Frère et al. (2010) and Carter et al. (2013), we did not include an 'avoided' category

because the number of dyads that did not associate and/or overlap spatially

exceed the 5th percentile, which was used as a threshold in previous studies. We

then calculated mean observed relatedness for each of the two categories

("preferred" and "non-preferred") for both AI and UDOI datasets. Next, we

randomly resampled relatedness values with replacement for each of the two

datasets, using the sample sizes for each category in the empirical data (i.e.,

sample sizes for "preferred" and "nonpreferred" were held constant). We repeated

this procedure 10,000 times to generate a distribution, and then calculated

significance as the fraction of replicates (out of 10,000) in which relatedness of

"preferred" AI or UDOI was greater than the observed, empirical value. The script

in which we implemented this procedure is available at

https://github.com/thw17/Varecia_social_preferences.

RESULTS

Association strength and spatial overlap

The average overall association index (AI) was 0.05 ± 0.03 (Baden et al.,

2016). We observed similar mean AIs in each of the three reproductive seasons

(nonbreeding = 0.04 ± 0.02; mating/gestation = 0.06 ± 0.03; birth/lactation = 0.07

$\pm$ 0.03). Within seasons, however, individuals varied substantially in their AIs.

Average individual AIs varied from 0.01 to 0.11 (nonbreeding = 0.01-0.08;

mating/gestation = 0.01-0.11; birth/lactation = 0.01-0.10), while maximum

individual AIs ranged from 0.07 to 0.82 (nonbreeding = 0.08-0.82; mating/gestation

= 0.07-0.79; birth/lactation = 0.08-0.82).

Average home range overlap (UDOI) was 0.211 ± 0.357. Some dyads did

not overlap at all (UDOI = 0), while the maximum UDOI observed was 1.895

(Figure 1). As with AIs, we observed similar mean UDOIs in each of the three

reproductive seasons (nonbreeding = 0.17 ± 0.27; mating/gestation = 0.15 ± 0.29;

birth/lactation = 0.20 ± 0.43; Table 1)

a) No overlap   **b) Some overlap**   **c) Extreme overlap**

Figure 1. Examples of home range overlap among pairs of individuals within the Mangevo ruffed lemur
community: a) Females Pink-Yellow and Radio-Yellow have no spatial overlap among home ranges (i.e.,
UDOI = 0.00), whereas b) females Radio-Yellow and Radio-Red and c) females Radio-Blue and Radio-Green
share some (UDOI = 0.106) to nearly complete home range overlap (UDOI = 0.951), respectively.

Table 1. Summary of association indices, utilization distribution overlap indices

(UDOIs), and relatedness overall and by reproductive season.

Association Indices	n	Mean	SD	Min	Max
Overall	136	0.05	0.03	0.00	0.80
Nonreproductive	136	0.04	0.12	0.00	0.82
Mating/Gestation	105	0.06	0.13	0.00	0.79
Birth/Lactation	91	0.07	0.15	0.00	0.82
UDOIs	n	Mean	SD	Min	Max
Overall	136	0.21	0.36	0.00	1.90
Nonreproductive	136	0.17	0.27	0.00	1.97
Mating/Gestation	120	0.15	0.29	0.00	1.74
Birth/Lactation	78	0.20	0.43	0.00	2.13
Relatedness	n	Mean	SD	Min	Max
Overall	171	-0.06	0.02	-0.72	0.65
Female-female	45	-0.13	0.04	-0.60	0.52
Female-male	90	-0.15	0.02	-0.72	0.28
Male-male	36	-0.13	0.04	-0.54	0.60

354
355
Relatedness
Genotypes were 93% complete; all subjects (n = 38) were scored for at least

12 loci (average = 14, range = 12 to 15). Allelic richness was 4.33 and average

observed heterozygosity was 0.400. There were no significant deviations from

Hardy-Weinberg Equilibrium for any of the loci examined, nor was there evidence

of null alleles.

Results of the rarefaction curve (y = 0.7991, r^2 =0.9992) showed average

relatedness values stabilizing after 5 loci, with the difference between mean

relatedness using 5 and 6 loci changing by only 0.95% (0.023), and the difference

between using 6 and 7 loci changing by only 0.56% (0.016). Thus, subsequent

dyadic r-value calculations included all possible dyads (n = 703 dyads), as all

individuals could be compared at 5 or more loci. Average pairwise relatedness

among adults within the community was -0.06 ± 0.02 and ranged from -0.72 to

0.65 (Table 1; Figure 2).

Figure 2. Pairwise genetic relatedness (r >= 0.25) among a) female-female dyads, b) male-male dyads, and
c) female-male dyads within the Mangevo social community. "?" indicates individuals for which genotype data
are unavailable. Nodes are organized according to individual home range centroids.

Relationships between association strength, spatial overlap, and relatedness

Using the full dataset, Mantel tests revealed that kinship was unrelated to

either home range overlap (UDOI; n = 25, r = 0.047 p = 0.724) or association

indices (AI; n = 14, r = 0.120, p = 0.294), whereas UDOI and AI were significantly

correlated (n = 17, r = 0.789, p << 0.001; Figure 3). This pattern held across all

three reproductive seasons (Table 2). It is worth noting, however, that the

correlation coefficient between UDOI and AI in the nonbreeding season was much

lower, albeit still significantly positive, than the mating/gestation and birth/lactation

seasons, which were very similar (Table 2). Similarly, when we used MRQAPs to

jointly analyze the effects of UDOI, kinship, and sex on AI, in every time period we

analyzed, the partial correlations between UDOI and AI were always significantly

positive, while kinship and sex were never significantly correlated with AI (Table

3).

Figure 3. Scatterplot illustrating the relationship between social association (AI) and home range overlap
(UDOI). AIs range from 0 (no association) to 1 (100% association). Similarly, UDOIs range from 0 (no overlap)
to 1 (complete overlap); values of >1 are possible if both UDs are nonuniformly distributed and also have a
high degree of overlap. Values <1 indicate less overlap relative to uniform space, whereas values >1 indicate
higher than normal overlap relative to uniform space. White circles indicate kin (r>=0.25); black triangles
indicate non-kin (r<0.25).

397   Table 2. Results from Mantel tests correlating Association Indices with utilization

398   distribution overlap indices (UDOIs), and relatedness overall and by reproductive

399   season.

Season	Variable	Records	N	Correlation	p-value
Overall	UDOI	11,171	17	0.789	<<0.001
Overall	Relatedness	11,171	14	0.120	0.294
Mating	UDOI	3,267	13	0.891	<<0.001
Mating	Relatedness	3,267	13	0.099	0.392
Nonbreeding	UDOI	5,051	14	0.566	<<0.001
Nonbreeding	Relatedness	5,051	14	0.149	0.184
Birth/gestation	UDOI	2,664	11	0.892	<<0.001
Birth/gestation	Relatedness	2,664	11	0.159	0.254

401   Note: Overall includes 2007-2008, whereas seasonal measures are from 2008
402   only. N indicates the number of individuals included in each analysis; individuals
403   had to be represented in both datasets in order to be included in the analyses for
404   a given season.
405
406

408   Table 3. Results from a multiple regression quadratic assignment procedure (MRQAP) investigating the partial relationships

409   of association (AI) with utilization distribution overlap indices (UDOI), relatedness, and sex.

Season	N	UDOI		Relatedness		Sex	
		Partial corr	p-value	Partial corr	p-value	Partial corr	p-value
Overall	14	0.920	<<0.001	0.035	0.754	0.056	0.617
Nonbreeding	11	0.655	<<0.001	0.123	0.356	-0.113	0.460
Mating/gestation	10	0.873	<<0.001	-0.049	0.742	0.281	0.720
Birth/lactation	8	0.941	<<0.001	0.296	0.144	0.102	0.530

Relatedness among dyads with "preferred" association strengths or spatial overlap

Average relatedness among "preferred" dyads, defined as falling in the >95[th] percentile of AIs (5 of 91 dyads), was significantly higher than among dyads classified as "not preferred" (mean $R_{xAI\_preferred}$ = 0.109 versus mean $R_{xAI\_non-preferred}$ = -0.095; p = 0.049). In contrast, average relatedness of dyads with the greatest home range overlap, defined as falling in the >95[th] percentile of UDOI (10 of 190 dyads), was not statistically different from other dyads (mean $R_{xUDOI\_preferred}$ = -0.015326 versus mean $R_{xUDOI\_non-preferred}$ = -0.03464, p = 0.3978).

Discussion

Results from this study illustrate complex relationships among social association, space use, and kinship in wild black-and-white ruffed lemurs, patterns that—while unusual for primates—are well aligned with much of the broader mammalian literature. We found that ruffed lemur social associations varied immensely, ranging from no association between some individuals to dyads observed together more than 75% of the time. From a community perspective, the social network was sparse, with weak associations (AIs) being common. Similarly, home range overlap was minimal and average relatedness among community members was low. These patterns were consistent throughout the year and across reproductive seasons. Together, these and earlier lines of evidence (e.g., Baden, 2011; Baden et al., 2016) can be used to characterize ruffed lemurs as having a

spatiotemporally dispersed fission-fusion social organization with weak social and

kinship ties.

Kin selection theory (Hamilton, 1964) has long been invoked to explain the

social preferences observed among mammals (e.g., Archie et al., 2006; Frère et

al., 2010; Godde et al., 2015; Smith, 2014; Wahaj et al., 2004), particularly in

primates (reviewed in Langergraber, 2012; Silk, 2002). In this study, however, we

found no evidence that kinship structured either spatial overlap or social

association overall in the community. There were exceptions, however, as some,

but not all, of the very closest social relationships were between closely related

individuals. The closest social ties in the community appear to occur primarily

between females and their preferred (often unrelated male) social partners,

followed by mothers and their pre-dispersal-aged subadult and adult offspring.

Nevertheless, although fewer than ten percent (8.8%) of adults within the

community were genetic relatives, nearly three-quarters of adult females (71%)

and all of adult males (100%) were closely related to at least one, and up to as

many as three other same-sex relative(s) within the community. Thus, while overall

patterns of dyadic relatedness suggest that ruffed lemur communities are

composed of both kin and non-kin, and that preferred associates are sometimes

close relatives, related dyads are not forming spatial networks of kin within the

larger communal range. These patterns contrast with the spatially structured

matrilines described in *Microcebus murinus* (Eberle & Kappeler, 2006; Radespiel,

Lutermann, Schmelting, Bruford, & Zimmermann, 2003), another communally

breeding strepsirrhine. Some authors have used this model to hypothesize that, at

least in part, kin networks have facilitated the evolution of cooperative infant care

(Eberle & Kappeler, 2006). That kin networks don't exist in ruffed lemurs, who also

engage in cooperative infant care (Baden et al., 2013), further highlights the

uniqueness of their communal breeding strategies.

These results build on growing evidence that spatial overlap is more

strongly associated with social association than kinship in many taxa characterized

by high fission-fusion dynamics (Best et al., 2014; K. D. Carter et al., 2013; Frère

et al., 2010; Strickland et al., 2014). Importantly, not all dyads with a high degree

of home range overlap were close social associates. Indeed, many dyads with

nonzero spatial overlap were never observed together. Similar patterns have been

observed in dolphins, giraffes, and water dragons, wherein subjects did not

associate, despite sharing complete or near complete home range overlap (i.e.,

'social avoidance' in dolphins: Frère et al., 2010; giraffes: K. D. Carter et al., 2013;

water dragons: Strickland et al., 2014). Thus, spatial overlap does not necessitate

social association. Instead, these results raise important questions related to

causation. What motivates social association? Are individuals that bias their time

toward overlapping areas simply more likely to associate? Or, in cases where

patterns of spatial overlap and social association do not align, is there some

additional force shaping these spatial and social decisions? Evaluating these

alternatives requires further investigation.

Moreover, while home range overlap explained most of the variation seen

in social association in ruffed lemurs, some variation remains unaccounted for,

suggesting that other social, ecological, and biological factors must also be at play.

One possibility is that even weak or infrequent social associations may facilitate

cooperative resource defense against other frugivorous competitors. For instance,

ruffed lemurs actively defend fruit-bearing trees against larger brown lemur (Genus

Eulemur) groups for days or even weeks during the resource scarce austral winter

(Baden, personal observation). Perhaps social association during these times

better equips otherwise solitary individuals to defend valuable fruit resources

against interspecific competitors. If this was the case, one might expect higher AIs

during resource scarce seasons, periods that correspond primarily with

mating/gestation, but also birth/lactation seasons. AIs were lowest in the

nonbreeding, resource abundant periods, though not significantly so, lending

minimal support for this hypothesis. In addition, communal breeding plays an

important role in female reproductive success in this species (Baden et al., 2013)

and might also be important in driving social preferences. For instance, recent work

suggests social networks in guppies may be structured by the propensity for non-

kin to cooperate (Croft et al., 2009), which could lead to the maintenance of

cooperation in the absence of kin assortment (Fletcher & Doebeli, 2009). These

lines of research offer exciting opportunities to better understand the myriad factors

shaping social preferences in fission-fusion species.

Acknowledgements

Research would not have been possible without the logistical support of The

Institute for the Conservation of Tropical Environments (ICTE), MICET and Centre

ValBio, or permission from ANGAP/Madagascar's National Parks. Drs. Randy

Junge, Felicia Knightly, Angie Simai, Edward E. Louis, Jr., and staff of the

Madagascar Biodiversity Project provided invaluable assistance during capture

procedures. Solo Justin, Telo Albert, Lahitsara Pierre, Razafindrakoto Georges,

Leroa, Velomaro, Reychell Chadwick, Lindsay Dytham, and A.J. Lowin assisted

with data collection and provided invaluable support and advice on the ground,

while Gary Aronsen provided support in the Yale Molecular Anthropology Lab. We

are further grateful for resources and support from the Center for High

Performance Computing at the University of Utah. Funding was generously

provided by National Science Foundation DDIG (BSC-0725975), the US

Department of State Fulbright Fellowship, The Leakey Foundation, Primate

Conservation, Inc., Conservation International's Primate Action Fund, Stony Brook

University, Henry Doorly Zoo, and Yale University.

References

Agrell, J., Wolff, J. O., & Ylonen, H. (1998). Counter-strategies to infanticide in
mammals: costs and consequences. *Oikos*, *83*, 507–517.
Alberts, S. C. (2019). Social influences on survival and reproduction: Insights
from a long-term study of wild baboons. *Journal of Animal Ecology*, *88*(1),
47–66. https://doi.org/10.1111/1365-2656.12887
Altmann, J. (1974). Observational Study of Behavior: Sampling Methods.
Behaviour, *49*(3), 227–267. https://doi.org/10.1163/156853974X00534
Altmann, J., & Alberts, S. (1996). Behavior predicts genes structure in a wild
primate group. *Proceedings of the National Academy of Sciences of the*
United States of America, *93*(June), 5797–5801. Retrieved from
http://www.pnas.org/content/93/12/5797.short
Andersson, M. (2017). Helping relatives survive and reproduce: inclusive fitness
and reproductive value in brood parasitism. *American Naturalist*, *189*(2),
138–152. https://doi.org/10.1086/689991
Archie, E. A., Moss, C. J., & Alberts, S. C. (2006). The ties that bind: genetic
relatedness predicts the fission and fusion of social groups in wild African
elephants. *Proceeding of the Royal Society B.*
https://doi.org/10.1098/rspb.2005.3361

Archie, E. A., Tung, J., Clark, M., Altmann, J., & Alberts, S. C. (2014). Social affiliation matters: Both same-sex and opposite-sex relationships predict survival in wild female baboons. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1793). https://doi.org/10.1098/rspb.2014.1261

Augusto, J. F., Frasier, T. R., & Whitehead, H. (2017). Social structure of long-finned pilot whales (Globicephala melas) off northern Cape Breton Island, Nova Scotia. *Behaviour*, *154*(5), 509–540. https://doi.org/10.1163/1568539X-00003432

Baden, A. L. (2011). *Communal infant care in black-and-white ruffed lemurs (Varecia variegata). Ph.D Thesis*. Stony Brook University.

Baden, A. L., Brenneman, R. A., & Louis Jr., E. E. (2008). Morphometries of wild black-and-white ruffed lemurs [Varecia variegata; Kerr, 1792]. *American Journal of Primatology*, *70*(10). https://doi.org/10.1002/ajp.20583

Baden, A. L., & Gerber, B. D. (n.d.). Spatial ecology of black-and-white ruffed lemurs (Varecia variegata) in southeastern Madagascar. *American Journal of Primatology*.

Baden, A. L., Holmes, S. M., Johnson, S. E., Engberg, S. E., Louis, E. E., & Bradley, B. J. (2014). Species-level view of population structure and gene flow for a critically endangered primate (Varecia variegata). *Ecology and Evolution*, *4*(13). https://doi.org/10.1002/ece3.1119

Baden, A. L., Webster, T. H., & Kamilar, J. M. (2016). Resource seasonality and reproduction predict fission-fusion dynamics in black-and-white ruffed lemurs (Varecia variegata). *American Journal of Primatology*, *78*(2), 256–279. https://doi.org/10.1002/ajp.22507

Baden, A. L., Wright, P. C., Louis, E. E., & Bradley, B. J. (2013). Communal nesting, kinship, and maternal success in a social primate. *Behavioral Ecology and Sociobiology*, *67*(12), 1939–1950. https://doi.org/10.1007/s00265-013-1601-y

Balbo, N., & Barban, N. (2014). Does fertility behavior spread among friends? *American Sociological Review*, *79*(3), 412–431. https://doi.org/10.1177/0003122414531596

Balko, E. A., & Underwood, H. B. (2005). Effects of forest structure and composition on food availability for Varecia variegata at Ranomafana National Park, Madagascar. *American Journal of Primatology*, *66*(1), 45–70. https://doi.org/10.1002/ajp.20127

Best, E. C., Dwyer, R. G., Seddon, J. M., & Goldizen, A. W. (2014). Associations are more strongly correlated with space use than kinship in female eastern grey kangaroos. *Animal Behaviour*, *89*, 1–10. https://doi.org/10.1016/j.anbehav.2013.12.011

Brent, L. J. N., Heilbronner, S. R., Horvath, J. E., Gonzalez-Martinez, J., Ruiz-Lambides, A., Robinson, A. G., … Platt, M. L. (2013). Genetic origins of social networks in rhesus macaques. *Scientific Reports*, *3*, 1–8. https://doi.org/10.1038/srep01042

Briga, M., Pen, I., & Wright, J. (2012). Care for kin: Within-group relatedness and allomaternal care are positively correlated and conserved throughout the mammalian phylogeny. *Biology Letters*, *8*(4), 533–536.

https://doi.org/10.1098/rsbl.2012.0159

Brown, G. E., & Brown, J. A. (1996). Does kin-biased territorial behavior increase
kin-biased foraging in juvenile salmonids? *Behavioral Ecology*, *7*(1), 24–29.
https://doi.org/10.1093/beheco/7.1.24

Cairns, S. J., & Schwager, S. J. (1987). A comparison of association indices.
Animal Behaviour, *35*(5), 1454–1469. https://doi.org/10.1016/S0003-
3472(87)80018-0

Calenge, C. (2011). Home range estimation in R : the adehabitatHR package. *R*
Package, 1–60. Retrieved from http://cran.r-
project.org/web/packages/adehabitatHR/vignettes/adehabitatHR.pdf

Cameron, E. Z., Setsaas, T. H., & Linklater, W. L. (2009). Social bonds between
unrelated females increase reproductive success in feral horses.
Proceedings of the National Academy of Sciences of the United States of
America, *106*(33), 13850–13853. https://doi.org/10.1073/pnas.0900639106

Carter, G. G., & Wilkinson, G. S. (2015). Social benefits of non-kin food sharing
by female vampire bats. *Proceedings of the Royal Society B: Biological*
Sciences, *282*(1819). https://doi.org/10.1098/rspb.2015.2524

Carter, K. D., Seddon, J. M., Frère, C. H., Carter, J. K., & Goldizen, A. W. (2013).
Fission-fusion dynamics in wild giraffes may be driven by kinship, spatial
overlap and individual social preferences. *Animal Behaviour*, *85*(2), 385–
394. https://doi.org/10.1016/j.anbehav.2012.11.011

Cheney, D. L., Silk, J. B., & Seyfarth, R. M. (2016). Network connections, dyadic
bonds and fitness in wild female baboons. *Royal Society Open Science*,
3(7). https://doi.org/10.1098/rsos.160255

Clutton-Brock, T. H. (1989). Mammalian mating systems. *Proceedings of the*
Royal Society of London. Series B. Biological Sciences, *236*(1285), 339–
372. https://doi.org/10.1098/rspb.1989.0027

Cords, M. (2002). Friendship among adult female blue monkeys (Cercopithecus
mitis). *Behaviour*, *139*(2), 291–314.

Croft, D. P., Hamilton, P. B., Darden, S. K., Jacoby, D. M. P., James, R.,
Bettaney, E. M., & Tyler, C. R. (2012). The role of relatedness in structuring
the social network of a wild guppy population. *Oecologia*, *170*(4), 955–963.
https://doi.org/10.1007/s00442-012-2379-8

Croft, D. P., James, R., Ward, A. J. W., Botham, M. S., Mawdsley, D., & Krause,
J. (2005). Assortative interactions and social networks in fish. *Oecologia*,
143(2), 211–219. https://doi.org/10.1007/s00442-004-1796-8

Croft, D. P., Krause, J., Darden, S. K., Ramnarine, I. W., Faria, J. J., & James, R.
(2009). Behavioural trait assortment in a social network: Patterns and
implications. *Behavioral Ecology and Sociobiology*, *63*(10), 1495–1503.
https://doi.org/10.1007/s00265-009-0802-x

Daura-Jorge, F. G., Cantor, M., Ingram, S. N., Lusseau, D., & Simões-Lopes, P.
C. (2012). The structure of a bottlenose dolphin society is coupled to a
unique foraging cooperation with artisanal fishermen. *Biology Letters*, *8*(5),
702–705. https://doi.org/10.1098/rsbl.2012.0174

de Ruiter, J. R., & Geffen, E. (1998). Relatedness of matrilines, dispersing males
and social groups in long-tailed macaques (Macaca fascicularis).

Proceedings of the Royal Society B: Biological Sciences, *265*, 79–87. https://doi.org/10.1017/CBO9781107415324.004

Dekker, D., Krackhardt, D., & Snijders, T. A. B. (2007). Sensitivity of MRQAP tests to collinearity and autocorrelation conditions. *Psychometrika*, *72*(4), 563–581. https://doi.org/10.1007/s11336-007-9016-1

Doreian, P., & Conti, N. (2012). Social context, spatial structure and social network structure. *Social Networks*, *34*(1), 32–46. https://doi.org/10.1016/j.socnet.2010.09.002

Eberle, M., & Kappeler, P. M. (2006). Family insurance: Kin selection and cooperative breeding in a solitary primate (Microcebus murinus). *Behavioral Ecology and Sociobiology*, *60*(4), 582–588. https://doi.org/10.1007/s00265-006-0203-3

Fieberg, J., & Kochanny, C. O. (2005). Quantifying home-range overlap: the importance of the utilization distribution. *The Journal of Wildlife Management*, *69*(4), 1346–1359.

Firth, J. A., Voelkl, B., Crates, R. A., Aplin, L. M., Biro, D., Croft, D. P., & Sheldon, B. C. (2017). Wild birds respond to flockmate loss by increasing their social network associations to others. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1854). https://doi.org/10.1098/rspb.2017.0299

Fletcher, J. A., & Doebeli, M. (2009). A simple and general explanation for the evolution of altruism. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1654), 13–19. https://doi.org/10.1098/rspb.2008.0829

Frère, C. H., Krützen, M., Mann, J., Watson-Capps, J. J., Tsai, Y. J., Patterson, E. M., … Sherwin, W. B. (2010). Home range overlap, matrilineal and biparental kinship drive female associations in bottlenose dolphins. *Animal Behaviour*, *80*(3), 481–486. https://doi.org/10.1016/j.anbehav.2010.06.007

Gero, S., Gordon, J., & Whitehead, H. (2015). Individualized social preferences and long-term social fidelity between social units of sperm whales. *Animal Behaviour*, *102*, 15–23. https://doi.org/10.1016/j.anbehav.2015.01.008

Gilby, I. C., Brent, L. J. N., Wroblewski, E. E., Rudicell, R. S., Hahn, B. H., Goodall, J., & Pusey, A. E. (2013). Fitness benefits of coalitionary aggression in male chimpanzees. *Behavioral Ecology and Sociobiology*, *67*(3), 373–381. https://doi.org/10.1007/s00265-012-1457-6

Gilby, I. C., & Wrangham, R. W. (2008). Association patterns among wild chimpanzees (Pan troglodytes schweinfurthii) reflect sex differences in cooperation. *Behavioral Ecology and Sociobiology*, *62*(11), 1831–1842. https://doi.org/10.1007/s00265-008-0612-6

Ginsberg, J. R., & Young, T. P. (1992). Measuring association between individuals or groups in behavioural studies. *Animal Behaviour*, *44*(PART 2), 377–379. https://doi.org/10.1016/0003-3472(92)90042-8

Glander, K. E. (1993). Capture and marking techniques for arboreal primates. In A. Estrada, E. Rodriguez-Luna, R. Lopez-Wilchis, & R. Coates-Estrada (Eds.), *Estudios Primatologicos en Mexico, Vol. 1* (pp. 299–304).

Godde, S., Côté, S. D., & Réale, D. (2015). Female mountain goats, Oreamnos americanus, associate according to kinship and reproductive status. *Animal Behaviour*, *108*, 101–107. https://doi.org/10.1016/j.anbehav.2015.07.005

Griffiths, S. W., & Armstrong, J. D. (2002). Kin-biased territory overlap and food
sharing among Atlantic salmon juveniles. *Journal of Animal Ecology*, *71*(3),
480–486. https://doi.org/10.1046/j.1365-2656.2002.00614.x
Hamilton, W. D. (1964). The genetical evolution of social behaviour. I. *Journal of
Theoretical Biology*, *7*(1), 1–16. https://doi.org/10.1016/0022-
5193(64)90038-4
Hinde, R. A. (1976). Interactions, relationships and social structure. *Man*, *11*(1),
1–17.
Hirsch, B. T., Prange, S., Hauver, S. A., & Gehrt, S. D. (2013). Genetic
relatedness does not predict racoon social network structure. *Animal
Behaviour*, *85*(2), 463–470. https://doi.org/10.1016/j.anbehav.2012.12.011
Kohn, G. M., Meredith, G. R., Magdaleno, F. R., King, A. P., & West, M. J.
(2015). Sex differences in familiarity preferences within fission-fusion brown-
headed cowbird, Molothrus ater, flocks. *Animal Behaviour*, *106*, 137–143.
https://doi.org/10.1016/j.anbehav.2015.05.023
Kossinets, G., & Watts, D. J. (2006). Empirical analysis of an evolving social
network. *Science*, *311*, 88–90. https://doi.org/10.1126/science.1116869
Langergraber, K. E. (2012). Kinship and cooperation in primates. In *The
Evolution of Primate Societies* (pp. 491–513).
Langergraber, K. E., Mitani, J. C., & Vigilant, L. (2007). The limited impact of
kinship on cooperation in wild chimpanzees. *Proceedings of the National
Academy of Sciences*, *104*(19), 7786–7790.
https://doi.org/10.1073/pnas.0611449104
Langergraber, K. E., Mitani, J., & Vigilant, L. (2009). Kinship and social bonds in
female chimpanzees (Pan troglodytes). *American Journal of Primatology*,
71(10), 840–851. https://doi.org/10.1002/ajp.20711
Leu, S. T., Bashford, J., Kappeler, P. M., & Bull, C. M. (2010). Association
networks reveal social organization in the sleepy lizard. *Animal Behaviour*,
79(1), 217–225. https://doi.org/10.1016/j.anbehav.2009.11.002
Longmire, J. L., Gee, G. F., Hardekopf, C. L., Graham, A., & Mark, G. A. (1992).
Establishing paternity in whooping cranes (Grus americana) by DNA
analysis. *The Auk*, *109*(3), 522–529.
Louis, E. E., Ratsimbazafy, J. H., Razakamaharauo, V. R., Pierson, D. J., Barber,
R. C., & Brenneman, R. A. (2005). Conservation genetics of black and white
ruffed lemurs, Varecia variegata, from Southeastern Madagascar. *Animal
Conservation*, *8*(1), 105–111. https://doi.org/10.1017/S1367943004001891
Lusseau, D., & Newman, M. E. J. (2004). Identifying the role that animals play in
their social networks. *Proceedings of the Royal Society B: Biological
Sciences*, *271*(SUPPL. 6), 477–481. https://doi.org/10.1098/rsbl.2004.0225
Lusseau, D., Wilson, B., Hammond, P. S., Grellier, K., Durban, J. W., Parsons, K.
M., … Thompson, P. M. (2006). Quantifying the influence of sociality on
population structure in bottlenose dolphins. *Journal of Animal Ecology*,
75(1), 14–24. https://doi.org/10.1111/j.1365-2656.2005.01013.x
Maher, C. R. (2009). Genetic relatedness and space use in a behaviorally flexible
species of marmot, the woodchuck (Marmota monax). *Behavioral Ecology
and Sociobiology*, *63*(6), 857–868. https://doi.org/10.1007/s00265-009-0726-

5

Mantel, N. (1967). The Detection of Disease Clustering and a Generalized
Regression Approach. *Cancer Research*, *27*(2 Part 1), 209 LP – 220.
Retrieved from
http://cancerres.aacrjournals.org/content/27/2_Part_1/209.abstract

McComb, K., Moss, C., Durant, S. M., Baker, L., & Sayialel, S. (2001). Matriarchs
as repositories of social knowledge in African elephants. *Science*, *292*, 491–
494. https://doi.org/10.1126/science.1057895

McDonald, D. B. (2007). Predicting fate from early connectivity in a social
network. *Proceedings of the National Academy of Sciences of the United*
States of America, *104*(26), 10910–10914.
https://doi.org/10.1073/pnas.0701159104

Mitani, J. C., Merriwether, D. A., & Zhang, C. (2000). Male affiliation, cooperation
and kinship in wild chimpanzees. *Animal Behaviour*, *59*(4), 885–893.
https://doi.org/10.1006/anbe.1999.1389

Moscovice, L. R., Douglas, P. H., Martinez-Iñigo, L., Surbeck, M., Vigilant, L., &
Hohmann, G. (2017). Stable and fluctuating social preferences and
implications for cooperation among female bonobos at LuiKotale, Salonga
National Park, DRC. *American Journal of Physical Anthropology*, *163*(1),
158–172. https://doi.org/10.1002/ajpa.23197

Mourier, J., Vercelloni, J., & Planes, S. (2012). Evidence of social communities in
a spatially structured network of a free-ranging shark species. *Animal*
Behaviour, *83*(2), 389–401. https://doi.org/10.1016/j.anbehav.2011.11.008

Nystrand, M. (2007). Associating with kin affects the trade-off between energy
intake and exposure to predators in a social bird species. *Animal Behaviour*,
74(3), 497–506. https://doi.org/10.1016/j.anbehav.2006.11.031

Peakall, R., & Smouse, P. E. (2012). GenALEx 6.5: Genetic analysis in Excel.
Population genetic software for teaching and research-an update.
Bioinformatics, *28*(19), 2537–2539.
https://doi.org/10.1093/bioinformatics/bts460

Perry, S. (2012). *The Behavior of Wild White-Faced Capuchins. Advances in the*
Study of Behavior (1st ed., Vol. 44). Elsevier Inc.
https://doi.org/10.1016/b978-0-12-394288-3.00004-6

Podgórski, T., Lusseau, D., Scandura, M., Sönnichsen, L., & Jędrzejewska, B.
(2014). Long-lasting, kin-directed female interactions in a spatially structured
wild boar social network. *PLoS ONE*, *9*(6), 1–11.
https://doi.org/10.1371/journal.pone.0099875

Pope, T. R. (2000). Reproductive success increases with degree of kinship in
cooperative coalitions of female red howler monkeys (Alouatta seniculus).
Behavioral Ecology and Sociobiology, *48*(4), 253–267.
https://doi.org/10.1007/s002650000236

Queller, D. C., & Goodnight, K. F. (1989). Estimating relatedness using genetic
markers. *Evolution*, *43*(2), 258–275.

Radespiel, U., Lutermann, H., Schmelting, B., Bruford, M. W., & Zimmermann, E.
(2003). Patterns and dynamics of sex-biased dispersal in a nocturnal
primate, the grey mouse lemur, Microcebus murinus. *Animal Behaviour*,

65(4), 709–719. https://doi.org/10.1006/anbe.2003.2121

Ramos-Fernández, G., Boyer, D., Aureli, F., & Vick, L. G. (2009). Association networks in spider monkeys (Ateles geoffroyi). *Behavioral Ecology and Sociobiology*, *63*(7), 999–1013. https://doi.org/10.1007/s00265-009-0719-4

Riehl, C. (2011). Living with strangers: direct benefits favour non-kin cooperation in a communally nesting bird. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1712), 1728–1735. https://doi.org/10.1098/rspb.2010.1752

Salpeteur, M., Patel, H., Balbo, A. L., Rubio-Campillo, X., Madella, M., Ajithprasad, P., & Reyes-García, V. (2015). When knowledge follows blood: Kin groups and the distribution of traditional ecological knowledge in a community of seminomadic pastoralists, Gujarat (India). *Current Anthropology*, *56*(3), 471–483. https://doi.org/10.1086/681006

Schülke, O., Bhagavatula, J., Vigilant, L., & Ostner, J. (2010). Social bonds enhance reproductive success in male macaques. *Current Biology*, *20*(24), 2207–2210. https://doi.org/10.1016/j.cub.2010.10.058

Seyfarth, R. M., & Cheney, D. L. (2012). The Evolutionary Origins of Friendship. *Annual Review of Psychology*, *63*(1), 153–177. https://doi.org/10.1146/annurev-psych-120710-100337

Silk, J. B. (2002). Kin selection in primate groups. *International Journal of Primatology*, *23*(4), 849–875. https://doi.org/10.1023/A:1015581016205

Silk, J. B. (2007). The adaptive value of sociality in mammalian groups. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *362*(1480), 539–559. https://doi.org/10.1098/rstb.2006.1994

Silk, J. B., Alberts, S. C., & Altmann, J. (2003). Social Bonds of Female Baboons Enhance Infant Survival. *Science*, *302*, 1231–1234. https://doi.org/10.1126/science.1088580

Silk, J. B., Alberts, S. C., & Altmann, J. (2006). Social relationships among adult female baboons (Papio cynocephalus) II. Variation in the quality and stability of social bonds. *Behavioral Ecology and Sociobiology*, *61*(2), 197–204. https://doi.org/10.1007/s00265-006-0250-9

Silk, J. B., Altmann, J., & Alberts, S. C. (2006). Social relationships among adult female baboons (papio cynocephalus) I. Variation in the strength of social bonds. *Behavioral Ecology and Sociobiology*, *61*(2), 183–195. https://doi.org/10.1007/s00265-006-0249-2

Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., … Cheney, D. L. (2009). The benefits of social capital: Close social bonds among female baboons enhance offspring survival. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1670), 3099–3104. https://doi.org/10.1098/rspb.2009.0681

Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., … Cheney, D. L. (2010a). Female chacma baboons form strong, equitable, and enduring social bonds. *Behavioral Ecology and Sociobiology*, *64*(11), 1733–1747. https://doi.org/10.1007/s00265-010-0986-0

Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., … Cheney, D. L. (2010b). Strong and consistent social bonds enhance the longevity of female baboons. *Current Biology*, *20*(15), 1359–1361.

https://doi.org/10.1016/j.cub.2010.05.067
Smith, J. E. (2014). Hamilton's legacy: Kinship, cooperation and social tolerance
inmammalian groups. *Animal Behaviour*, *92*, 291–304.
https://doi.org/10.1016/j.anbehav.2014.02.029
Spiegel, O., Sih, A., Leu, S. T., & Bull, C. M. (2018). Where should we meet?
Mapping social network interactions of sleepy lizards shows sex-dependent
social network structure. *Animal Behaviour*, *136*, 207–215.
https://doi.org/10.1016/j.anbehav.2017.11.001
Stanley, C. R., Mettke-Hofmann, C., Hager, R., & Shultz, S. (2018). Social
stability in semiferal ponies: networks show interannual stability alongside
seasonal flexibility. *Animal Behaviour*, *136*, 175–184.
https://doi.org/10.1016/j.anbehav.2017.04.013
Stone, J., & Björklund, M. (2001). DELRIOUS: A computer program designed to
analyse molecular marker data and calculate delta and relatedness
estimates with confidence. *Molecular Ecology Notes*, *1*(3), 209–212.
https://doi.org/10.1046/j.1471-8278.2001.00034.x
Strickland, K., Gardiner, R., Schultz, A. J., & Frère, C. H. (2014). The social life of
eastern water dragons: Sex differences, spatial overlap and genetic
relatedness. *Animal Behaviour*, *97*, 53–61.
https://doi.org/10.1016/j.anbehav.2014.08.009
Strickland, K., Levengood, A., Foroughirad, V., Mann, J., Krzyszczyk, E., & Frère,
C. H. (2017). A framework for the identification of long-term social avoidance
in longitudinal datasets. *Royal Society Open Science*, *4*(8).
https://doi.org/10.1098/rsos.170641
Taberlet, P. (1996). Reliable genotyping of samples with very low DNA quantities
using PCR. *Nucleic Acids Research*, *24*(16), 3189–3194.
https://doi.org/10.1093/nar/24.16.3189
Vasey, N. (2006). Impact of seasonality and reproduction on social structure,
ranging patterns and fission–fusion social organization in red ruffed lemurs.
In *Lemurs: ecology and adaptation* (pp. 275–305). New York:
Springer/Kluwer.
Wahaj, S. A., Van Horn, R. C., Van Horn, T. L., Dreyer, R., Hilgris, R., Schwarz,
J., & Holekamp, K. E. (2004). Kin discrimination in the spotted hyena
(Crocuta crocuta): Nepotism among siblings. *Behavioral Ecology and
Sociobiology*, *56*(3), 237–247. https://doi.org/10.1007/s00265-004-0783-8
Wey, T., Blumstein, D. T., Shen, W., & Jordán, F. (2008). Social network analysis
of animal behaviour: a promising tool for the study of sociality. *Animal
Behaviour*, *75*(2), 333–344. https://doi.org/10.1016/j.anbehav.2007.06.020
Whitehead, H. (1996). Babysitting, dive synchrony, and indications of alloparental
care in sperm whales. *Behavioral Ecology & Sociobiology*, *38*(4), 237–244.
Whitehead, H. (2008). *Analyzing Animal Societies: Quantitative methods for
vertebrate social analysis.* Chicago: University of Chicago Press.
Whitehead, H. (2009). SOCPROG programs: Analysing animal social structures.
Behavioral Ecology and Sociobiology, *63*(5), 765–778.
https://doi.org/10.1007/s00265-008-0697-y
Wilkinson, G. S., Carter, G. G., Bohn, K. M., & Adams, D. M. (2016). Non-kin

cooperation in bats. *Philosophical Transactions of the Royal Society B:*
Biological Sciences, *371*(1687). https://doi.org/10.1098/rstb.2015.0095
Wiszniewski, J., Allen, S. J., & Möller, L. M. (2009). Social cohesion in a
hierarchically structured embayment population of Indo-Pacific bottlenose
dolphins. *Animal Behaviour*, *77*(6), 1449–1457.
https://doi.org/10.1016/j.anbehav.2009.02.025
Wright, P. C., Erhart, E. M., Tecot, S., Baden, A. L., Arrigo-nelson, S. J., Herrera,
J., … Zohdy, S. (2012). *Long-Term Field Studies of Primates*.
https://doi.org/10.1007/978-3-642-22514-7