Genetic relatedness cannot explain social preferences in black-and-white

ruffed lemurs (*Varecia variegata*).

ABSTRACT

Fission-fusion social dynamics are common among a number of vertebrate taxa,

and yet the factors shaping these variable associations among subgroup

members have not been widely addressed. Associations may occur simply

because of shared habitat preferences; however, social ties may also be

influenced by genetic relatedness (kinship) or social attraction. Here, we

investigate the association patterns of wild black-and-white ruffed lemurs,

Varecia variegata, in Ranomafana National Park, Madagascar using behavioural,

spatial (home range), and genetic data from twenty-four individually identified

animals. We collected 40,840 records of group composition over a 17-month

period and from this calculated pairwise association indices. We also used

ranging coordinates and genetic samples to estimate patterns of spatial overlap

and kinship, and then related these measures to patterns of affiliation. From

these analyses, we found that dyadic ruffed lemur social associations were

generally sparse and weak; that home range overlap was minimal; and that

average relatedness within the community was low. We found no evidence that

kinship was related to patterns of either spatial overlap or social association;

instead, associations were primarily driven by space use. Moreover, social

preferences were unrelated to kinship. While home range overlap explained most

of the variation seen in social association, some variation remains unaccounted

for, suggesting that other social, ecological, and biological factors such as shared

resource defense or communal breeding might also play a role in social attraction.

Our results further highlight the need to consider individual space use and

nuances of species behavior when investigating social preference and social

association more generally.

Keywords: social network analysis, range use, spatial overlap, Association Index,

kinship, fission-fusion, microsatellites, primates

INTRODUCTION

Animal social systems reflect non-random relationships among

neighbouring conspecifics, the content, quality, and patterning of which define

their social structure (Hinde, 1976). In fact, there is growing evidence that taxa as

diverse as fishes, lizards, birds, cetaceans, and equids exhibit social associations

and interactions that are not only non-random, but highly structured (Augusto,

Frasier, & Whitehead, 2017; Croft et al., 2012, 2005; Firth et al., 2017; Spiegel,

Sih, Leu, & Bull, 2018; Stanley, Mettke-Hofmann, Hager, & Shultz, 2018).

Individuals vary in the numbers, strengths, and stabilities of their social ties (Croft

et al., 2005; McDonald, 2007; Silk, Alberts, & Altmann, 2006; Silk, Altmann, &

Alberts, 2006; Silk et al., 2010a; Stanley et al., 2018), as well as in their

preferences for particular social associates (e.g., Cords, 2002; Gero, Gordon, &

Whitehead, 2015; Kohn, Meredith, Magdaleno, King, & West, 2015; Mourier,

Vercelloni, & Planes, 2012; Perry, 2012; Schülke, Bhagavatula, Vigilant, &

Ostner, 2010)—outcomes of which can be evolutionarily significant (Seyfarth &

Cheney, 2012; Silk, 2007). For example, the extent, strength, and nature of

females' social relationships have been positively linked to higher fertility

(*Macaca mulatta*: Brent et al., 2013; humans: Balbo & Barban, 2014), longer

lifespan (*Papio cynocephalus*: Alberts, 2019; Archie, Tung, Clark, Altmann, &

Alberts, 2014; *Papio ursinus:* Silk et al., 2010b), and greater offspring survival

(*Equus caballus*: Cameron, Setsaas, & Linklater, 2009; *Papio cynocephalus*:

Silk, Alberts, & Altmann, 2003; *Papio ursinus:* Cheney, Silk, & Seyfarth, 2016;

Silk et al., 2009; *Physeter microcephalus*: Whitehead, 1996). Similarly, males

that form strong social bonds have been shown to be more successful at forming

coalitions, achieving high rank, and siring more offspring than males with weaker

social bonds (e.g., *Chiroxiphia linearis*: McDonald, 2007; *Macaca assamensis*:

Schülke et al., 2010; *Pan troglodytes*: Gilby et al., 2013). Nevertheless, while the

long-term health and fitness outcomes of sociality have been the focus of much

investigation, until recently the factors shaping social preferences have received

comparatively less attention.

Several hypotheses have been proposed to explain social association, the

most popular being kinship, whereby group members preferentially associate

with and, in the case of kin selection, are more likely to direct costly altruistic

behaviours toward close genetic relatives (Hamilton, 1964). For instance, many

female mammals associate with and exhibit social preferences toward maternal

kin (Smith, 2014), a pattern observed in primates in particular (reviewed in

Langergraber, 2012), but also in spotted hyaenas (Wahaj et al., 2004), African

elephants (Archie, Moss, & Alberts, 2006), dolphins (Frère et al., 2010), and

mountain goats (Godde, Côté, & Réale, 2015), among others. Social preferences

toward kin can be driven by the benefits of associating with relatives, such as

allomaternal care (reviewed in Briga, Pen, & Wright, 2012; Pope, 2000; and in

some cases even brood parasitism, Andersson, 2017), reduced aggression and

infanticide risk (reviewed in Brown & Brown, 1996; Agrell, Wolff, & Ylonen, 1998),

foraging advantages (Griffiths & Armstrong, 2002; Nystrand, 2007), and shared

social and ecological knowledge (McComb, Moss, Durant, Baker, & Sayialel,

2001; Salpeteur et al., 2015). However, social associations have also been linked

to direct fitness consequences among nonrelatives (Baden, Wright, Louis, &

Bradley, 2013; Cameron et al., 2009; G. G. Carter & Wilkinson, 2015; McFarland

et al. 2017; Riehl, 2011), suggesting that social preferences can evolve based on

direct benefits alone. Moreover, evidence for kin-biased relationships in other

species, particularly those with higher fission-fusion dynamics, is weak (e.g.,

Best, Dwyer, Seddon, & Goldizen, 2014; K. D. Carter, Seddon, Frère, Carter, &

Goldizen, 2013; Hirsch, Prange, Hauver, & Gehrt, 2013; Langergraber, Mitani, &

Vigilant, 2007, 2009; Moscovice et al., 2017; Wilkinson, Carter, Bohn, & Adams,

2016), raising questions as to the ubiquity of kinship in the formation and

maintenance of social bonds.

One complicating factor in determining whether and how kinship shapes

social preference is that it can be difficult to decouple social associations due to

kinship from associations due to other social and spatial contexts (reviewed in

Doreian & Conti, 2012; Wey, Blumstein, Shen, & Jordán, 2008). For example, in

some species it is difficult to differentiate whether strong associations are due to

strong genetic ties or to other factors, such as sex or age (Lusseau & Newman,

2004; Mourier et al., 2012), habitat utilization (Wiszniewski, Allen, & Möller,

2009), or foraging specializations (Daura-Jorge, Cantor, Ingram, Lusseau, &

Simões-Lopes, 2012; Gilby & Wrangham, 2008; Griffiths & Armstrong, 2002;

Mitani, Merriwether, & Zhang, 2000). Indeed, patterns of social association have

been found to correlate with spatial overlap (i.e., overlapping home ranges) more

strongly than, or to the exclusion of, kinship in a number of species (Best et al.,

2014; K. D. Carter et al., 2013; Frère et al., 2010; Strickland, Gardiner, Schultz, &

Frère, 2014). Moreover, although individuals living in closer physical proximity

are more prone to interact (e.g., Clutton-Brock, 1989; Kossinets & Watts, 2006),

conspecifics with high levels of spatial overlap may also mediate their

interactions by temporally modifying range use, such that animals with highly

overlapping ranges may interact minimally, if at all (e.g., temporal avoidance)

(Leu, Bashford, Kappeler, & Bull, 2010; Ramos-Fernández, Boyer, Aureli, & Vick,

2009; Strickland et al., 2017). While still uncommon, studies that consider the

simultaneous effects of spatial overlap and kinship on association patterns are

thus important to understanding the role of social preference in the evolution of

social systems and sociality (e.g., Best et al., 2014; K. D. Carter et al., 2013;

Frère et al., 2010; Lusseau et al., 2006; Maher, 2009; Podgórski, Lusseau,

Scandura, Sönnichsen, & Jędrzejewska, 2014; Strickland et al., 2014; Piza-Roca

et al. 2019).

Ruffed lemurs (Genus *Varecia*) are moderately-sized frugivores (Baden,

Brenneman, & Louis Jr., 2008; Balko & Underwood, 2005) that live in groups

(hereafter "communities") comprising as many as 30 adult and subadult

individuals and their offspring (Baden, Webster, & Kamilar, 2016 and references

therein). From earlier work, we know that communities are characterized by high

fission-fusion dynamics (*sensu* Kummer 1971), with members associating in fluid

subgroups (or '"parties") that vary in size, cohesion, membership, and duration

(Baden, Webster, & Kamilar, 2016). Within communities, there is striking

variation in individual ranging behavior, degree of home range overlap with other

community members, and the number and strength of social associations

(Baden, 2011; Baden & Gerber, n.d.; Baden et al., 2016; Vasey, 2006).

Individuals never use the entire communal territory, instead concentrating their

ranging to proportionately smaller areas that overlap with as few as four (25%) to

as many as twelve (75%) other community members (Baden & Gerber, n.d.).

Association patterns are equally variable, with individuals associating in

subgroups with anywhere from one to ten individuals throughout the year (Baden

et al. 2016). Overall, associations are generally weak, and animals spend nearly

half of their time alone (Baden et al. 2016). Nevertheless, certain individuals

appear to preferentially associate and are regularly found together in "core

groups" whose members are characterized by both high association strength and

home range overlap (Vasey 2006; Morland 1991a,b; Baden 2011). Moreover,

studies of microsatellite STR markers and mitochondrial DNA have shown that,

while average pairwise relatedness within communities is low (i.e., both sexes

disperse: Baden et al., 2014), communities nevertheless contain relatives (e.g,

Baden, 2011; Baden et al., 2013). Taken together, these lines of evidence

suggest that during subgroup formation community members may be actively

choosing social associates, and that these associations may be shaped by a

number of ecological, social, and biological factors. Here, we hypothesize (H1)

that social associations among members of a black-and-white ruffed lemur

(*Varecia variegata*) community are driven by space use and kinship. Specifically,

we quantify dyadic measures of association strength, spatial overlap, and

relatedness to test the predictions that members with higher home range overlap

(P1.1) and relatedness (P1.2) will associate more often than dyads which exhibit

disparate range use or are unrelated. Moreover, because previous studies have

found a relationship between social preference and kinship, we further

hypothesize (H2) that relatedness will drive association preferences (P2.1), with

preferred associates being more related than non-preferred dyads.

METHODS

Study site and subjects

We collected data from one *V. variegata* community at Mangevo [21°220

6000 S, 47°280 000 E], a mid-elevation (660-1,200 m) primary rainforest site

located in the southeastern parcel of Ranomafana National Park, Madagascar

(Wright et al., 2012) during 16 months of study (August–December 2007;

February– December 2008). At the time of the study, the community included 24

adults and subadults (8 adult females, 11 adult males, 5 subadult males). All

individuals in this study were habituated and individually identified via radio-

collars or unique collar-tag combinations prior to behavioural observations (see

Baden et al., 2016 for specific details; see also Ethical note below). Animals were

collared under veterinary supervision following a strict protocol (Glander, 1993).

Nineteen infants were born in the 2008 birth season and were present from

October to December 2008, when the study ended. Sampling efforts resulted in a

total of 4,044 focal observation hours.

Ethical note

All animal protocols adhere to the guidelines for the treatment of animals

for teaching and research recommended by ASAB/ABS (2019). Subjects were

habituated to researcher presence as part of a long-term study established in

2005. Since 2005, subjects have been periodically immobilized in the field

(approximately once every 2 to 3 years) for application or replacement of radio-

transmitter collars, which allow researchers to accurately identify individuals and

locate and track them in the field during behavioral observations. Immobilization

and sampling procedures adhered to the protocols established by Glander (1993)

and have been implemented by Dr. Randall Junge (MS, DVM, Dipl American

College of Zoological Medicine, Dipl American College of Animal Welfare),

Director of the Prosimian Biomedical Survey Project (PBSP), since 2000. During

captures, each lemur is immobilized by remote injection with

tiletamine/zolazepam. Animals are examined immediately upon capture, and

once determined to be stable they are transported back to the basecamp for

examination (not more than 15 minutes). Animals are evaluated again for

stability, and monitored by a licensed veterinarian (Dr. Junge) throughout the

anesthesia. Biological samples, including blood (no more than 1 ml per kilogram)

for genetic analysis and biomedical health assessments, are collected for later

analysis. Lemurs are given a balanced electrolyte solution (30 ml)

subcutaneously and are monitored until recovery. Once fully recovered, lemurs

are released at the site of capture. Anesthetic episodes typically last 1 hour or

less. Lemurs that are adequately awake by midafternoon (3 PM) are released the

same day. Those that are not adequately recovered are held overnight in cloth

bags and released before 10 AM the next morning. As ruffed lemurs are diurnal,

releasing animals with residual sedation in the late afternoon could predispose

them to predation. This capture, monitoring, sampling and release protocol has

been used successfully by PBSP veterinarians for 15 years, on 44 trips to 18

sites throughout Madagascar. Biomedical health assessments by Dr. Junge and

population genetic assessments by Baden et al. (2014, 2019) are integral parts

of population management, as they are used to identify heath concerns such as

introduction of human or livestock parasites to the site (as documented in other

parts of Ranomafana National Park) and inbreeding depression. They are also

useful for identifying health parameters in a population inhabiting a pristine forest,

allowing that information to be used as a benchmark to evaluate this and other

populations should they experience epidemic disease or population decline.

Observers never interact with subjects outside of the capture period. Utmost care

was taken to minimize the impact of our presence on our subjects during

captures and subsequent behavioral observations. Observations were conducted

noninvasively at a minimum observer distance of 10 m from the focal subject.

When in the presence of subjects, observers speak quietly and make efforts to

not disturb the individual. No more than four observers are allowed to participate

in observations at any one time. Subjects were target for focal observations, at

most, twice per month. Permission to conduct research was granted by

Madagascar's National Parks (ANGAP/MNP, #084/07-041/08). Research

protocols were approved by and in compliance with Stony Brook University

IACUC #2005-20081449.

Data collection

Behavioural monitoring

We collected data during dawn to dusk follows of focal individuals. We

located focal subjects at the beginning of each observation period via radio-

telemetry and selected new subjects daily. We never sampled focal subjects on

consecutive days and every effort was made to follow all subjects at least once

per month. If an individual with a collar-tag was located in association with a

radio-collared focal individual prior to 10:00 h, this individual became the new

focal subject for that observation period. Observational periods ranged in

duration between 8 to 11 hours depending on seasonal differences in day length

and time needed to locate animals at dawn.

Upon initial contact with the focal individual, we recorded the number and

identities of all other individuals present within the subgroup. To do so, one

observer remained with the focal individual while the remaining three team

members spread out to locate and identify all other members of the subgroup. *A*

priori, individuals were considered members of a subgroup only if they were

within 50 m of the group center and were seen associating, traveling with and

maintaining proximity to members of the subgroup being followed. However, an

earlier study found that average group spread in this population ranged between

6 to 15 m, and rarely exceeded 30 m (Baden et al. 2016). Thus, while subgroup

membership was operationalized to include all members within 50 m, effective

subgroup spread was typically much smaller (see Baden et al., 2016 for details).

After initial contact, we monitored subsequent changes in subgroup size,

composition (age/sex class, individual identity), and cohesion (i.e., the greatest

distance between any two subgroup members), as well as activity state of the

focal subject using instantaneous scan sampling techniques collected at 5 min

intervals (Altmann, 1974). Sampling efforts resulted in a total of 4,044 focal

observation hours, during which time we recorded 40,840 group scans.

We collected simultaneous GPS coordinates at 10 min intervals from as

close to the focal individual as possible to document daily individual range use.

We recorded spatial coordinates only if estimated positional error was less than

10 m.

Genetic sampling

In addition, we collected genetic samples from 38 black-and-white ruffed

lemurs from the Mangevo population, including all subjects in this study, during

five capture seasons spanning four consecutive years (2005-2008; see Baden,

2011 for details). Sample collection occurred under veterinary supervision and

followed a strict protocol outlined by Glander (1993). All capture procedures

occurred during non-reproductive seasons in the absence of infants and

dependent offspring.

For each individual captured, one of us (ALB) collected approximately 1

ml/kg of whole blood (~4 cc) from the femoral vein and four 2mm tissue biopsies

from ear pinnae. All samples were stored in 5 ml of lysis buffer solution (0.1 M

Tris-HCl pH, 8.0, 0.1 M EDTA, 0.01 M NaCl, and 0.5% w/v SDS) at ambient

temperature in the field (7 to 21 days) (Longmire, Gee, Hardekopf, Graham, &

Mark, 1992). We then banked samples in a −80°C freezer at the Madagascar

Biodiversity Partnership headquarters in Antananarivo, Madagascar and

subsequently at the Yale Molecular Anthropology Lab in New Haven, CT until

later genetic analysis (see below).

Data analysis

Social association

In this study, we define social association as individuals being present in

the same subgroup at the same time. We considered social preferences to occur

(that is, the dyad was composed of "preferred associates") when dyads had a

significantly stronger strength of association than other dyads in the community

(see "Relatedness among preferred associates" below). This species is typically

described as being "female-bonded" (Morland 1991a), in line with the definition

that to be 'socially bonded' requires that multiple, independent types of

associations or interactions are significantly higher than expected (Whitehead,

2008). We do not examine social bonds in this study because we only focus on a

single measure of association. Finally, we also use the term "social tie" in the

network sense (e.g., Wasserman and Faust, 1994; Croft et al., 2008): individuals

that have a social tie have a nonzero association index and a "stronger tie" has a

higher association index.

Sampling was biased toward radio-collared females; we therefore

subsampled our data prior to association analyses. Details are presented in

Baden et al. (2016), but briefly, we divided our dataset into monthly periods and

according to the sex of the focal individual. Using only scans for which all

individuals were known, we randomized point scans and selected X scans to

include in the dataset for each sex, where X is 90% of the point scans for the sex

with the fewest scans in a given month. This procedure resulted in the inclusion

of a total of 11,784 point scans, with equal numbers of scans targeting male and

female focal subjects in each month (Baden et al., 2016).

We used SOCPROG 2.9 (Whitehead, 2009) to calculate association

indices (AIs) between all pairs (i.e., dyads) of individuals using the "simple ratio"

index, which quantifies the proportion of time that two individuals are observed

together relative to their total observation time overall. This index is appropriate

when individuals are equally likely to be correctly identified (Cairns & Schwager,

1987; Ginsberg & Young, 1992; Whitehead, 2008), which is the case for this

population. Given that subgroup composition in this community changes

approximately every 90 minutes (Baden et al., 2016), we used a 6-hour sampling

interval to reduce autocorrelation among periods. We also removed any subject

observed in fewer than 10 sampling periods. Our final dataset (overall dataset)

included 8 adult females and 9 adult males totaling 136 dyads observed in over

11,171 point scans.

To examine seasonal variation in associations, we organized the

subsetted data from 2008 into three seasons according to female reproductive

state as defined by Baden et al. (2013). Using the protocols described above, our

final datasets included: 8 females and 9 males over 5,051 scans during the

nonbreeding season (January through June); 7 females and 8 males over 3,267

scans during the mating/gestation season (July through September); and 8

females and 6 males over 2,664 scans during the birth/lactation season (October

through December). We compared distributions of AIs across seasons using a

Kruskal-Wallis test implemented in R (Kruskall and Wallis, 1952).

Spatial overlap

Estimates of home range overlap used in this study come from Baden &

Gerber (n.d.). Briefly, we calculated home range overlap between all pairs of

individuals using a utilization distribution overlap index (UDOI; Fieberg &

Kochanny, 2005) implemented in the R package adehabitat (Calenge, 2011).

The UDOI is an index of space-use sharing between two utilization distributions

(UDs). UDOI values can range from 0 to 1, with a UDOI of 0 indicating no home

range overlap and a UDOI of 1 indicating that home ranges are uniformly

distributed and have 100% overlap. Values can also be >1 if both UDs are

nonuniformly distributed and also have a high degree of overlap. Values <1

indicate less overlap relative to uniform space, whereas values >1 indicate higher

than normal overlap relative to uniform space. We calculated four UDOIs for all

pairs of individuals: one annual UDOI, and three seasonal UDOIs according to

female reproductive state, as for association analyses described above.

Relatedness

We genotyped individuals at a suite of 15 polymorphic microsatellite loci

(see Baden, 2011 and Baden et al., 2014 for details). We extracted total genomic

DNA from blood and/or tissue samples using standard nucleic acid extraction kits

(QIAamp DNA Mini Kit; Qiagen) automated on a QiaCube (Qiagen). Extraction

procedures followed the manufacturer's protocols, with the following modification

to the tissue extraction procedures: samples were allowed to lyse initially in ASL

buffer for 24-48 hours rather than 10 minutes.

We carried out PCR amplifications in a total reaction volume of 25 μl

consisting of 2 μl template, 12.5 μl Qiagen HotStar Taq Master Mix, and 10 μM of

each primer. Amplification conditions were as follows: initial denaturation at 95 °C

for 15 min; 35 cycles of 30 s at 94 °C, 40 s at 54 to 60 °C (see Louis et al., 2005),

1 min at 72 °C, and a final extension of 7 min at 72 °C. The 5′ end of the forward

primer was fluorescently labeled, and amplification products were separated and

visualized using capillary electrophoresis (ABI 3730xl Genetic Analyzer). We

assessed allele sizes relative to an internal size standard (ROX-500) using Gene

Mapper software (Applied Biosystems), and scored final genotypes based on

multiple independent reactions (Taberlet, 1996). Panels yielded PI$_{sib}$ (Queller &

Goodnight, 1989) values of 2.7×10^{-5}, demonstrating the very low probability that

two individuals would share the same multilocus genotype by chance. We further

tested the robusticity of this suite of loci for estimating relatedness with a

rarefaction analysis as in Altmann & Alberts (1996) and de Ruiter & Geffen

(1998) using the program RE-RAT (http://people.musc.edu/~schwaclh/). We

estimated pairwise relatedness among individuals (*r*) following Queller & Goodnight

(1989) using the program GenAlEx 6.5 (Peakall & Smouse, 2012). Relatedness was

based on allele frequencies derived from a larger population of 38 adult multilocus

genotypes (Baden, 2011). Fine scale dyadic relatedness assessments (e.g.,

distinguishing between full and half-sibs) are not possible in most microsatellite

studies, and in fact, the ability to differentiate relatedness disjunctions on such a

scale would probably require 30 to 60 microsatellite loci (Stone & Björklund,

2001). We therefore consider "related dyads" to be those with r-values ≥ 0.25; we

made no attempt to further distinguish categories or degrees of relatedness.

Relationships among social associations, spatial overlap, and relatedness

We used a series of Mantel tests followed by a multiple regression

quadratic assignment procedure (MRQAP) to test our first hypothesis (H1), that

association strength was driven by spatial overlap and relatedness. We first used

Mantel tests to examine whether association indices were independently related

to either spatial overlap (P1.1) or relatedness (P1.2), as well as to test for a

correlation between spatial overlap and relatedness (Mantel, 1967). We further

used a multiple regression quadratic assignment procedure (MRQAP; Dekker,

Krackhardt, & Snijders, 2007) to determine relationships between the response

variable—association index—and predictor variables—sex, spatial overlap

(P1.1), and relatedness (P1.2). For sex, we used a matrix of sex similarity, which

used values of 1 (same sex) and 0 (different sex). MRQAP tests each pairwise

combination of response and predictor matrices while holding the remaining

predictor matrices constant. We performed both Mantel tests and MRQAP for the

overall dataset as well as in each of the three seasons using SOCPROG 2.9

(Whitehead, 2009).

Relatedness among preferred associates

Studies of other taxa have found that relatedness can shape social

preferences, even if it does not structure association among all dyads (Carter et

al., 2013; Best et al., 2014). Therefore, in addition to the analyses described

above, we used generalized linear models to test our second hypothesis (H2)

that relatedness between two individuals in a dyad predicted whether they were

"preferred" associates. Here, we define "preferred associates" (we also use

"social preference") as dyads exhibiting significantly higher association indices

than other dyads in the community.

We identified preferred associates in our overall dataset (described in

"Social association" above) using SOCPROG 2.9's (Whitehead, 2009)

permutation tests based on Bejder et al. (1998) with modifications by Whitehead

(2008). We used the "permute associations within samples" option, along with

1000 flips per permutation and 10,000 permutations. We classified dyads as

preferred associates if the dyad's AI was significantly high (one-tailed α = 0.05).

We next used logistic regression analysis, implemented using Scikit-learn

in Python (Pedregosa et al., 2011) to test if relatedness of a dyad (independent

variable; measured using continuous relatedness values calculated as described

in "Relatedness" above) predicted whether members in that dyad were preferred

associates (P2.1; dependent variable; binary response of "preferred" or "not"

based on SOCPROG permutation analyses). To assess significance, we

compared the coefficient of the regression model to a "null" distribution of

coefficients generated from 10,000 randomizations of the model. During each

replicate, we shuffled relatedness values among dyads before building the

randomized regression model. Our p-value was calculated as: 1 – (*percentile of*

observed coefficient in null distribution). The script in which we implemented this

procedure is available at https://github.com/thw17/Varecia_social_preferences.

RESULTS

Social association

The average overall association index (AI) was 0.05 ± 0.03 (Baden et al.,

2016). While we observed similar mean AIs in each of the three reproductive

seasons (nonbreeding = 0.04 ± 0.02; mating/gestation = 0.06 ± 0.03;

birth/lactation = 0.07 ± 0.03), the distributions of AIs significantly differed across

seasons (Kruskall-Wallis chi-squared = 8.423, df = 2, p = 0.015). Within seasons,

individuals also varied substantially in their AIs. Distributions were right skewed,

with many values at or near 0 and maximum values as high as 0.82. Average

individual AIs (i.e., the average of all of an individual's AIs) varied from 0.01 to

0.11 (nonbreeding = 0.01-0.08; mating/gestation = 0.01-0.11; birth/lactation =

0.01-0.10), while maximum individual AIs ranged from 0.07 to 0.82 (nonbreeding

= 0.08-0.82; mating/gestation = 0.07-0.79; birth/lactation = 0.08-0.82).

Spatial overlap

Average home range overlap (UDOI) was 0.211 ± 0.357 (Baden & Gerber,

n.d.). Some dyads did not overlap at all (UDOI = 0), while the maximum UDOI

observed was 1.895 (Figure 1). As with AIs, we observed similar mean UDOIs in

each of the three reproductive seasons (nonbreeding = 0.17 ± 0.27;

mating/gestation = 0.15 ± 0.29; birth/lactation = 0.20 ± 0.43; Table 1; see also

Baden & Gerber n.d.)

a) No overlap   b) Some overlap   c) Extreme overlap

Figure 1. Examples of home range overlap among pairs of individuals within the
Mangevo ruffed lemur community: a) Females Pink-Yellow and Radio-Yellow
have no spatial overlap among home ranges (i.e., UDOI = 0.00), whereas b)

females Radio-Yellow and Radio-Red and c) females Radio-Blue and Radio-
Green share some (UDOI = 0.106) to nearly complete home range overlap
(UDOI = 0.951), respectively.

435
436 *Relatedness*
437
Genotypes were 93% complete; all subjects (n = 38) were scored for at

least 12 loci (average = 14, range = 12 to 15). Allelic richness was 4.33 and

average observed heterozygosity was 0.400. There were no significant

deviations from Hardy-Weinberg Equilibrium for any of the loci examined, nor

was there evidence of null alleles.

Results of the rarefaction curve (y = 0.7991, r^2 =0.9992) showed average

relatedness values stabilizing after 5 loci, with the difference between mean

relatedness using 5 and 6 loci changing by only 0.95% (0.023), and the

difference between using 6 and 7 loci changing by only 0.56% (0.016). Thus,

subsequent dyadic r-value calculations included all possible dyads (n = 703

dyads), as all individuals could be compared at 5 or more loci. Average pairwise

relatedness among adults within the community was -0.06 ± 0.02 and ranged

from -0.72 to 0.65 (Table 1; Figure 2).

Table 1. Summary of association indices, utilization distribution overlap indices
(UDOIs), and relatedness (*r*) overall and by reproductive season.
454

Association Indices	n	Mean	SD	Min	Max
Overall	136	0.05	0.03	0.00	0.80
Nonreproductive	136	0.04	0.12	0.00	0.82
Mating/Gestation	105	0.06	0.13	0.00	0.79
Birth/Lactation	91	0.07	0.15	0.00	0.82
UDOIs	n	Mean	SD	Min	Max

Overall	136	0.21	0.36	0.00	1.90
Nonreproductive	136	0.17	0.27	0.00	1.97
Mating/Gestation	120	0.15	0.29	0.00	1.74
Birth/Lactation	78	0.20	0.43	0.00	2.13
Relatedness	n	Mean	SD	Min	Max
Overall	171	-0.06	0.02	-0.72	0.65
Female-female	45	-0.13	0.04	-0.60	0.52
Female-male	90	-0.15	0.02	-0.72	0.28
Male-male	36	-0.13	0.04	-0.54	0.60

455
456

Figure 2. Pairwise genetic relatedness (r ≥ 0.25) among adult a) female-female dyads, b) male-male dyads, and c) female-male dyads within the Mangevo social community. "?" indicates individuals for which genotype data are unavailable. Nodes are organized according to individual home range centroids.

Fewer than ten percent (8.8%) of dyads within the community were genetic relatives. Despite this, more than three-quarters of adult females (n=6 of 7 for which genetic data were available; 85.7%) and all of adult males (n=11 of 11; 100%) were related (r ≥ 0.25) to at least one, and up to as many as three other same-sex relative(s) within the community (Figure 2).

Relationships among social associations, spatial overlap, and relatedness

Using the full dataset, Mantel tests revealed that kinship was unrelated to

either home range overlap (UDOI; n = 25, r = 0.047 p = 0.724) or association

indices (AI; n = 14, r = 0.120, p = 0.294), whereas UDOI and AI were significantly

correlated (n = 17, r = 0.789, p << 0.001; Figure 3). This pattern held across all

three reproductive seasons (Table 2). It is worth noting, however, that the

correlation coefficient between UDOI and AI in the nonbreeding season was

much lower, albeit still significantly positive, than the mating/gestation and

birth/lactation seasons, which were very similar (Table 2). Similarly, when we

used MRQAPs to jointly analyze the effects of UDOI, kinship, and sex on AI, in

every time period we analyzed, the partial correlations between UDOI and AI

were always significantly positive, while kinship and sex were never significantly

correlated with AI (Table 3). These results support prediction P1.1, but do not

support prediction P1.2.

*Figure 3. Scatterplot illustrating the relationship between social association (AI)
and home range overlap (UDOI). AIs range from 0 (no association) to 1 (100%
association). Similarly, UDOIs range from 0 (no overlap) to 1 (complete overlap);
values of >1 are possible if both UDs are nonuniformly distributed and also have
a high degree of overlap. Values <1 indicate less overlap relative to uniform
space, whereas values >1 indicate higher than normal overlap relative to uniform
space. White circles indicate kin (r ≥ 0.25); black triangles indicate non-kin
(r<0.25).*

495   Table 2. Results from Mantel tests correlating Association Indices with utilization

496   distribution overlap indices (UDOIs), and relatedness overall and by reproductive

497   season.

Season	Variable	Records	N	Correlation	*p*-value
Overall	UDOI	11,171	17	0.789	<<0.001

Overall	Relatedness	11,171	14	0.120	0.294
Mating	UDOI	3,267	13	0.891	<<0.001
Mating	Relatedness	3,267	13	0.099	0.392
Nonbreeding	UDOI	5,051	14	0.566	<<0.001
Nonbreeding	Relatedness	5,051	14	0.149	0.184
Birth/gestation	UDOI	2,664	11	0.892	<<0.001
Birth/gestation	Relatedness	2,664	11	0.159	0.254

499 Note: Overall includes 2007-2008, whereas seasonal measures are from 2008
500 only. N indicates the number of individuals included in each analysis; individuals
501 had to be represented in both datasets in order to be included in the analyses for
502 a given season.
503
504

506 Table 3. Results from a multiple regression quadratic assignment procedure (MRQAP) investigating the partial

507 relationships of association (AI) with utilization distribution overlap indices (UDOI), relatedness, and sex.

Season	N	UDOI		Relatedness		Sex	
		Partial corr	*p*-value	Partial corr	*p*-value	Partial corr	*p*-value
Overall	14	0.920	<<0.001	0.035	0.754	0.056	0.617
Nonbreeding	11	0.655	<<0.001	0.123	0.356	-0.113	0.460
Mating/gestation	10	0.873	<<0.001	-0.049	0.742	0.281	0.720
Birth/lactation	8	0.941	<<0.001	0.296	0.144	0.102	0.530

Relatedness among preferred associates

Relatedness among preferred associates

We identified 18 (of 136 total) pairs of preferred associates, i.e., dyads

with AIs that were significantly higher than other dyads in the community. Of

these preferred associates, we were able to determine relatedness for 11 (91

total dyads with data for association and relatedness). Our logistic regression

model revealed that relatedness did not predict whether a pair of individuals

became preferred associates (coefficient = 0.087, p = 0.41) and thus did not

support prediction P2.1.

Discussion

Results from this study illustrate complex relationships among social

association, space use, and kinship in wild black-and-white ruffed lemurs,

patterns that—while unusual for primates—are well aligned with much of the

broader mammalian literature. We found that ruffed lemur social associations

varied immensely, ranging from no association between some individuals to

dyads observed together more than 75% of the time. From a community

perspective, the social network was sparse, with weak associations (AIs) being

common. Similarly, home range overlap was minimal and average relatedness

among community members was low. These patterns were consistent throughout

the year and across reproductive seasons. Together, these and earlier lines of

evidence (e.g., Baden, 2011; Baden et al., 2016) can be used to characterize

ruffed lemurs as having a spatiotemporally dispersed fission-fusion social

organization with weak social and kinship ties.

Kin selection theory (Hamilton, 1964) has long been invoked to explain the

social preferences observed among mammals (e.g., Archie et al., 2006; Frère et

al., 2010; Godde et al., 2015; Smith, 2014; Wahaj et al., 2004), particularly in

primates (reviewed in Langergraber, 2012; Silk, 2002). In this study, however, we

found no evidence that kinship structured either spatial overlap or social

association overall in the community. In contrast to other fission-fusion species

exhibiting similar patterns (Carter et al. 2013; Best et al. 2014), we further found

that relatedness did not predict social preference. Instead, the closest social ties

in the community appear to occur primarily between females and their

preferred—and often unrelated—male social partners, followed by mothers and

their pre-dispersal-aged subadult and adult offspring. Nevertheless, although

fewer than ten percent (8.8%) of adult dyads within the community were genetic

relatives, nearly three-quarters of adult females (85.7%) and all of adult males

(100%) were closely related to at least one, and up to as many as three other

same-sex relative(s). Thus, while our relatedness estimates suggest that ruffed

lemur communities include both kin and non-kin, and that preferred associates

are sometimes close relatives, kin are not forming spatial or social networks

across the larger communal range. These patterns contrast with the spatially

structured matrilines described in *Microcebus murinus* (Eberle & Kappeler, 2006;

Radespiel, Lutermann, Schmelting, Bruford, & Zimmermann, 2003), the only

other communally breeding strepsirrhine. Some authors have used this model to

hypothesize that spatially structured kin networks have, at least in part, facilitated

the evolution of cooperative infant care (Eberle & Kappeler, 2006). Cooperative

or communal breeding is exceedingly rare among primates (Tecot et al. 2013).

Only three nonhuman primates engage in this behavior: callitrichines (marmosets

and tamarins), cheirogaleids (mouse lemurs), and *Varecia* (Baden, Raboin,

Tecot, n.d.)*.* Like cheirogaleids, callitrichines also rely on close relatives to share

the burden of infant care (reviewed in Erb & Porter 2017). *Varecia* are the only

group that does not rely on close relatives for care (Baden et al., 2013), thereby

distinguishing *Varecia*'s infant care strategies from other cooperatively breeding

nonhuman primates.

These results build on growing evidence that space use is an important

predictor of social association—perhaps moreso than kinship—particularly in

taxa characterized by high fission-fusion dynamics (Best et al., 2014; K. D. Carter

et al., 2013; Frère et al., 2010; Strickland et al., 2014). Indeed, for two individuals

to interact they must be in relatively close physical proximity (Farine 2015). For

most animals, this means that social associations are therefore shaped by their

movement decisions (Emlen and Oring 1977; Farine et al. 2016; Bonnell et al.

2017), which are, in turn, influenced by aspects of their physical environment

(e.g., habitat structure, natural and/or anthropogenic barriers, resource

availability and distribution; reviewed in He et al. 2019). Among these

environmental variables, resource availability and distribution play a fundamental

role in shaping the space use of animals (e.g., Brown and Orians 1970; Clutton-

Brock and Harvey 1977) and by extension, their patterns of inter-individual

spatial proximity (e.g., Minta 1992; Croft et al. 2011) and social relationships

(e.g., Sterck et al. 1997; Bejder, Fletcher & Brager 1998; Wolf et al. 2007; Carter

et al. 2009).

To this point, we found that while home range overlap explained most of

the variation seen in social association in ruffed lemurs, some variation remained

unaccounted for, suggesting that other social, ecological, and biological factors

must also be at play. However, it is possible that even weak or infrequent social

associations may facilitate cooperative resource defense against other

frugivorous competitors. For instance, ruffed lemurs actively defend fruit-bearing

trees against larger brown lemur (Genus *Eulemur*) groups for days or even

weeks during the resource scarce austral winter (Baden, personal observation).

Perhaps social association during these times better equips otherwise solitary

individuals to defend valuable fruit resources against interspecific competitors.

Under this scenario, we would predict higher AIs during resource scarce

seasons, periods that correspond primarily with mating/gestation, but also

birth/lactation seasons. However, AIs were lowest in the nonbreeding, resource

abundant periods (though not significantly so), lending minimal support for this

hypothesis. It is possible, however, that home range overlap as measured by

UDOIs was insufficient to characterize the spatiotemporal component of

associations at highly contested resources. Future analyses may therefore

benefit from a more nuanced investigation of the spatially explicit role valuable

resources play in the social associations of this species.

In addition, communal breeding plays an important role in female

reproductive success in this species (Baden et al., 2013) and might also be

important in driving social preferences. For instance, recent work suggests social

networks in guppies may be structured by the propensity for non-kin to cooperate

(Croft et al., 2009), which could lead to the maintenance of cooperation in the

absence of kin assortment (Fletcher & Doebeli, 2009). These lines of research

offer exciting opportunities to better understand the myriad factors shaping social

preferences in fission-fusion species.

Nevertheless, shared space use does not always necessitate social

association. In this study, we found that not all dyads with a high degree of home

range overlap were close social associates. Indeed, many dyads with nonzero

spatial overlap were never observed together. Similar patterns have been

observed in dolphins, giraffes, and water dragons, wherein subjects did not

associate, despite sharing complete or near complete home range overlap (i.e.,

'social avoidance' in dolphins: Frère et al., 2010; giraffes: K. D. Carter et al.,

2013; water dragons: Strickland et al., 2014, 2017). In such cases, individuals

may actively and consistently avoid one another, despite the spatio-temporal

potential for repeated interactions (reviewed in Strickland et al. 2017). Like

predator avoidance, social avoidance can be costly. Thus, in future studies, it will

be critical to simultaneously consider the consequences of both social attraction

and avoidance when studying the evolution of sociality.

Together, these and earlier results raise important questions related to

causation. What motivates social association? Are individuals that bias their time

toward overlapping areas simply more likely to associate? Or, in cases where

patterns of spatial overlap and social association do not align, is there some

additional force shaping these spatial and social decisions? Evaluating these and

other alternatives require further investigation.

Acknowledgements

Research would not have been possible without the logistical support of

The Institute for the Conservation of Tropical Environments (ICTE), MICET and

Centre ValBio, or permission from ANGAP/Madagascar's National Parks. Drs.

Randy Junge, Felicia Knightly, Angie Simai, Edward E. Louis, Jr., and staff of the

Madagascar Biodiversity Project and Prosimian Biomedical Survey Project

provided invaluable assistance during capture procedures. Solo Justin, Telo

Albert, Lahitsara Pierre, Razafindrakoto Georges, Leroa, Velomaro, Reychell

Chadwick, Lindsay Dytham, and A.J. Lowin assisted with data collection and

provided invaluable support and advice on the ground, while Gary Aronsen

provided support in the Yale Molecular Anthropology Lab. We thank two

anonymous reviewers, as well as James Higham, for their insightful comments

on earlier versions of this manuscript. We are further grateful for resources and

support from the Center for High Performance Computing at the University of

Utah. Funding was generously provided by National Science Foundation DDIG

(BSC-0725975), the US Department of State Fulbright Fellowship, The Leakey

Foundation, Primate Conservation, Inc., Conservation International's Primate

Action Fund, Stony Brook University, Henry Doorly Zoo, and Yale University.

References

Agrell, J., Wolff, J. O., & Ylonen, H. (1998). Counter-strategies to infanticide in mammals: costs and consequences. *Oikos*, *83*, 507–517.

Alberts, S. C. (2019). Social influences on survival and reproduction: Insights from a long-term study of wild baboons. *Journal of Animal Ecology*, *88*(1), 47–66. https://doi.org/10.1111/1365-2656.12887

Altmann, J. (1974). Observational Study of Behavior: Sampling Methods. *Behaviour*, *49*(3), 227–267. https://doi.org/10.1163/156853974X00534

Altmann, J., & Alberts, S. (1996). Behavior predicts genes structure in a wild primate group. *Proceedings of the National Academy of Sciences of the United States of America*, *93*(June), 5797–5801. Retrieved from http://www.pnas.org/content/93/12/5797.short

Andersson, M. (2017). Helping relatives survive and reproduce: inclusive fitness and reproductive value in brood parasitism. *American Naturalist*, *189*(2), 138–152. https://doi.org/10.1086/689991

Archie, E. A., Moss, C. J., & Alberts, S. C. (2006). The ties that bind: genetic relatedness predicts the fission and fusion of social groups in wild African elephants. *Proceeding of the Royal Society B.* https://doi.org/10.1098/rspb.2005.3361

Archie, E. A., Tung, J., Clark, M., Altmann, J., & Alberts, S. C. (2014). Social affiliation matters: Both same-sex and opposite-sex relationships predict survival in wild female baboons. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1793). https://doi.org/10.1098/rspb.2014.1261

Augusto, J. F., Frasier, T. R., & Whitehead, H. (2017). Social structure of long-finned pilot whales (Globicephala melas) off northern Cape Breton Island, Nova Scotia. *Behaviour*, *154*(5), 509–540. https://doi.org/10.1163/1568539X-00003432

Baden, A. L. (2011). *Communal infant care in black-and-white ruffed lemurs (Varecia variegata)*. *Ph.D Thesis*. Stony Brook University.

Baden, A. L., Brenneman, R. A., & Louis Jr., E. E. (2008). Morphometries of wild black-and-white ruffed lemurs [Varecia variegata; Kerr, 1792]. *American Journal of Primatology*, *70*(10). https://doi.org/10.1002/ajp.20583

Baden, A. L., & Gerber, B. D. (n.d.). Spatial ecology of black-and-white ruffed lemurs (V*arecia variegata*) in southeastern Madagascar. *American Journal of Primatology*.

Baden, A. L., Holmes, S. M., Johnson, S. E., Engberg, S. E., Louis, E. E., & Bradley, B. J. (2014). Species-level view of population structure and gene flow for a critically endangered primate (Varecia variegata). *Ecology and Evolution*, *4*(13). https://doi.org/10.1002/ece3.1119

Baden, A. L., Webster, T. H., & Kamilar, J. M. (2016). Resource seasonality and reproduction predict fission-fusion dynamics in black-and-white ruffed lemurs (Varecia variegata). *American Journal of Primatology*, *78*(2), 256–279. https://doi.org/10.1002/ajp.22507

Baden, A. L., Wright, P. C., Louis, E. E., & Bradley, B. J. (2013). Communal nesting, kinship, and maternal success in a social primate. *Behavioral*

Ecology and Sociobiology, *67*(12), 1939–1950. https://doi.org/10.1007/s00265-013-1601-y

Balbo, N., & Barban, N. (2014). Does fertility behavior spread among friends? *American Sociological Review*, *79*(3), 412–431. https://doi.org/10.1177/0003122414531596

Balko, E. A., & Underwood, H. B. (2005). Effects of forest structure and composition on food availability for Varecia variegata at Ranomafana National Park, Madagascar. *American Journal of Primatology*, *66*(1), 45–70. https://doi.org/10.1002/ajp.20127

Bejder, L., Fletcher, D., Brager, S. (1998). A method for testing association patterns of social animals. *Animal Behaviour*, 56(3), 719-725.

Best, E. C., Dwyer, R. G., Seddon, J. M., & Goldizen, A. W. (2014). Associations are more strongly correlated with space use than kinship in female eastern grey kangaroos. *Animal Behaviour*, *89*, 1–10. https://doi.org/10.1016/j.anbehav.2013.12.011

Bonnell, T. R., Clarke, P. M., Henzi, S. P., Barrett, L. (2017). Individual-level movement bias leads to the formation of higher-order social structure in a mobile group of baboons. *Royal Society Open Science,* 4, 170148.

Brent, L. J. N., Heilbronner, S. R., Horvath, J. E., Gonzalez-Martinez, J., Ruiz-Lambides, A., Robinson, A. G., … Platt, M. L. (2013). Genetic origins of social networks in rhesus macaques. *Scientific Reports*, *3*, 1–8. https://doi.org/10.1038/srep01042

Briga, M., Pen, I., & Wright, J. (2012). Care for kin: Within-group relatedness and allomaternal care are positively correlated and conserved throughout the mammalian phylogeny. *Biology Letters*, *8*(4), 533–536. https://doi.org/10.1098/rsbl.2012.0159

Brown, G. E., & Brown, J. A. (1996). Does kin-biased territorial behavior increase kin-biased foraging in juvenile salmonids? *Behavioral Ecology*, *7*(1), 24–29. https://doi.org/10.1093/beheco/7.1.24

Brown, J. L., Orians, G. H. (1970). Spacing patterns in mobile animals. *Annual Review of Ecolocial and Evoluitionary Systems*, 1, 239–262.

Cairns, S. J., & Schwager, S. J. (1987). A comparison of association indices. *Animal Behaviour*, *35*(5), 1454–1469. https://doi.org/10.1016/S0003-3472(87)80018-0

Calenge, C. (2011). Home range estimation in R : the adehabitatHR package. *R Package*, 1–60. Retrieved from http://cran.r-project.org/web/packages/adehabitatHR/vignettes/adehabitatHR.pdf

Cameron, E. Z., Setsaas, T. H., & Linklater, W. L. (2009). Social bonds between unrelated females increase reproductive success in feral horses. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(33), 13850–13853. https://doi.org/10.1073/pnas.0900639106

Carter, A.J., Macdonald, S.L., Thomson, V.A. & Goldizen, A.W. (2009) .Structured association patterns and their energetic benefits in female eastern grey kangaroos, *Macropus giganteus*. *Animal Behaviour*, 77, 839–846.

Carter, G. G., & Wilkinson, G. S. (2015). Social benefits of non-kin food sharing

by female vampire bats. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1819). https://doi.org/10.1098/rspb.2015.2524

Carter, K. D., Seddon, J. M., Frère, C. H., Carter, J. K., & Goldizen, A. W. (2013). Fission-fusion dynamics in wild giraffes may be driven by kinship, spatial overlap and individual social preferences. *Animal Behaviour*, *85*(2), 385–394. https://doi.org/10.1016/j.anbehav.2012.11.011

Cheney, D. L., Silk, J. B., & Seyfarth, R. M. (2016). Network connections, dyadic bonds and fitness in wild female baboons. *Royal Society Open Science*, *3*(7). https://doi.org/10.1098/rsos.160255

Clutton-Brock, T. H. (1989). Mammalian mating systems. *Proceedings of the Royal Society of London. Series B. Biological Sciences*, *236*(1285), 339–372. https://doi.org/10.1098/rspb.1989.0027

Clutton-Brock, T. H., Harvey, P. H. (1977). Primate ecology and social organization. *Journal of Zoology*, 183, 1–39.

Cords, M. (2002). Friendship among adult female blue monkeys (Cercopithecus mitis). *Behaviour*, *139*(2), 291–314.

Croft, D. P., Hamilton, P. B., Darden, S. K., Jacoby, D. M. P., James, R., Bettaney, E. M., & Tyler, C. R. (2012). The role of relatedness in structuring the social network of a wild guppy population. *Oecologia*, *170*(4), 955–963. https://doi.org/10.1007/s00442-012-2379-8

Croft, D. P., James, R., Kraus, J. (2008) Exploring animal social networks. Princeton, NJ: Princeton University Press, 192p.

Croft, D. P., James, R., Ward, A. J. W., Botham, M. S., Mawdsley, D., & Krause, J. (2005). Assortative interactions and social networks in fish. *Oecologia*, *143*(2), 211–219. https://doi.org/10.1007/s00442-004-1796-8

Croft, D. P., Krause, J., Darden, S. K., Ramnarine, I. W., Faria, J. J., & James, R. (2009). Behavioural trait assortment in a social network: Patterns and implications. *Behavioral Ecology and Sociobiology*, *63*(10), 1495–1503. https://doi.org/10.1007/s00265-009-0802-x

Croft, D.P., Madden, J.R., Franks, D.W. & James, R. (2011). Hypothesis testing in animal social networks. *Trends in Ecology & Evolution*, 26, 502–507.

Daura-Jorge, F. G., Cantor, M., Ingram, S. N., Lusseau, D., & Simões-Lopes, P. C. (2012). The structure of a bottlenose dolphin society is coupled to a unique foraging cooperation with artisanal fishermen. *Biology Letters*, *8*(5), 702–705. https://doi.org/10.1098/rsbl.2012.0174

de Ruiter, J. R., & Geffen, E. (1998). Relatedness of matrilines, dispersing males and social groups in long-tailed macaques (Macaca fascicularis). *Proceedings of the Royal Society B: Biological Sciences*, *265*, 79–87. https://doi.org/10.1017/CBO9781107415324.004

Dekker, D., Krackhardt, D., & Snijders, T. A. B. (2007). Sensitivity of MRQAP tests to collinearity and autocorrelation conditions. *Psychometrika*, *72*(4), 563–581. https://doi.org/10.1007/s11336-007-9016-1

Doreian, P., & Conti, N. (2012). Social context, spatial structure and social network structure. *Social Networks*, *34*(1), 32–46. https://doi.org/10.1016/j.socnet.2010.09.002

Eberle, M., & Kappeler, P. M. (2006). Family insurance: Kin selection and

cooperative breeding in a solitary primate (Microcebus murinus). *Behavioral Ecology and Sociobiology*, *60*(4), 582–588. https://doi.org/10.1007/s00265-006-0203-3

Emlen, S. T., Oring, L. W. (1977). Ecology, sexual selection, and the evolution of mating systems. *Science,* 197, 215–223.

Erb, W. M., & Porter, L. M. (2017). Mother's little helpers: What we know (and don't know) about cooperative infant care in callitrichines. *Evolutionary Anthropology,* 26, 5-37. https://doi.org/10.1002/evan.21516

Farine, D. R. (2015) Proximity as a proxy for interactions: issues of scale in social network analysis. *Animal Behaviour*, 104, 1–5.

Farine, D. R., Strandburg-Peshkin, A., Berger-Wolf, T., Ziebart, B., Brugere, I., Li, J., Crofoot, M. C. (2016). Both nearest neighbours and long-term affiliates predict individual locations during collective movement in wild baboons. *Scientific Reports,* 6, 27704.

Fieberg, J., & Kochanny, C. O. (2005). Quantifying home-range overlap: the importance of the utilization distribution. *The Journal of Wildlife Management*, *69*(4), 1346–1359.

Firth, J. A., Voelkl, B., Crates, R. A., Aplin, L. M., Biro, D., Croft, D. P., & Sheldon, B. C. (2017). Wild birds respond to flockmate loss by increasing their social network associations to others. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1854). https://doi.org/10.1098/rspb.2017.0299

Fletcher, J. A., & Doebeli, M. (2009). A simple and general explanation for the evolution of altruism. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1654), 13–19. https://doi.org/10.1098/rspb.2008.0829

Frère, C. H., Krützen, M., Mann, J., Watson-Capps, J. J., Tsai, Y. J., Patterson, E. M., … Sherwin, W. B. (2010). Home range overlap, matrilineal and biparental kinship drive female associations in bottlenose dolphins. *Animal Behaviour*, *80*(3), 481–486. https://doi.org/10.1016/j.anbehav.2010.06.007

Gero, S., Gordon, J., & Whitehead, H. (2015). Individualized social preferences and long-term social fidelity between social units of sperm whales. *Animal Behaviour*, *102*, 15–23. https://doi.org/10.1016/j.anbehav.2015.01.008

Gilby, I. C., Brent, L. J. N., Wroblewski, E. E., Rudicell, R. S., Hahn, B. H., Goodall, J., & Pusey, A. E. (2013). Fitness benefits of coalitionary aggression in male chimpanzees. *Behavioral Ecology and Sociobiology*, *67*(3), 373–381. https://doi.org/10.1007/s00265-012-1457-6

Gilby, I. C., & Wrangham, R. W. (2008). Association patterns among wild chimpanzees (Pan troglodytes schweinfurthii) reflect sex differences in cooperation. *Behavioral Ecology and Sociobiology*, *62*(11), 1831–1842. https://doi.org/10.1007/s00265-008-0612-6

Ginsberg, J. R., & Young, T. P. (1992). Measuring association between individuals or groups in behavioural studies. *Animal Behaviour*, *44*(PART 2), 377–379. https://doi.org/10.1016/0003-3472(92)90042-8

Glander, K. E. (1993). Capture and marking techniques for arboreal primates. In A. Estrada, E. Rodriguez-Luna, R. Lopez-Wilchis, & R. Coates-Estrada (Eds.), *Estudios Primatologicos en Mexico, Vol. 1* (pp. 299–304).

Godde, S., Côté, S. D., & Réale, D. (2015). Female mountain goats, Oreamnos americanus, associate according to kinship and reproductive status. *Animal Behaviour*, *108*, 101–107. https://doi.org/10.1016/j.anbehav.2015.07.005

Griffiths, S. W., & Armstrong, J. D. (2002). Kin-biased territory overlap and food sharing among Atlantic salmon juveniles. *Journal of Animal Ecology*, *71*(3), 480–486. https://doi.org/10.1046/j.1365-2656.2002.00614.x

Hamilton, W. D. (1964). The genetical evolution of social behaviour. I. *Journal of Theoretical Biology*, *7*(1), 1–16. https://doi.org/10.1016/0022-5193(64)90038-4

He, P., Maldonado-Chaparro, A.A. & Farine, D.R. The role of habitat configuration in shaping social structure: a gap in studies of animal social complexity. *Behavioral Ecology Sociobiology*, 73, 9. https://doi:10.1007/s00265-018-2602-7.

Hinde, R. A. (1976). Interactions, relationships and social structure. *Man*, *11*(1), 1–17.

Hirsch, B. T., Prange, S., Hauver, S. A., & Gehrt, S. D. (2013). Genetic relatedness does not predict racoon social network structure. *Animal Behaviour*, *85*(2), 463–470. https://doi.org/10.1016/j.anbehav.2012.12.011

Kohn, G. M., Meredith, G. R., Magdaleno, F. R., King, A. P., & West, M. J. (2015). Sex differences in familiarity preferences within fission-fusion brown-headed cowbird, Molothrus ater, flocks. *Animal Behaviour*, *106*, 137–143. https://doi.org/10.1016/j.anbehav.2015.05.023

Kossinets, G., & Watts, D. J. (2006). Empirical analysis of an evolving social network. *Science*, *311*, 88–90. https://doi.org/10.1126/science.1116869

Kruskal, W.H., Wallis, W.A. (1952) Use of ranks in one-criterion variance analysis. J. Am. Stat. Assoc. *47*, 583–621.

Kummer, H. (1971). Primate societies: group techniques of ecological adaptation. Chicago: Aldine. p 160.

Langergraber, K. E. (2012). Kinship and cooperation in primates. In *The Evolution of Primate Societies* (pp. 491–513).

Langergraber, K. E., Mitani, J. C., & Vigilant, L. (2007). The limited impact of kinship on cooperation in wild chimpanzees. *Proceedings of the National Academy of Sciences*, *104*(19), 7786–7790. https://doi.org/10.1073/pnas.0611449104

Langergraber, K. E., Mitani, J., & Vigilant, L. (2009). Kinship and social bonds in female chimpanzees (Pan troglodytes). *American Journal of Primatology*, *71*(10), 840–851. https://doi.org/10.1002/ajp.20711

Leu, S. T., Bashford, J., Kappeler, P. M., & Bull, C. M. (2010). Association networks reveal social organization in the sleepy lizard. *Animal Behaviour*, *79*(1), 217–225. https://doi.org/10.1016/j.anbehav.2009.11.002

Longmire, J. L., Gee, G. F., Hardekopf, C. L., Graham, A., & Mark, G. A. (1992). Establishing paternity in whooping cranes (Grus americana) by DNA analysis. *The Auk*, *109*(3), 522–529.

Louis, E. E., Ratsimbazafy, J. H., Razakamaharauo, V. R., Pierson, D. J., Barber, R. C., & Brenneman, R. A. (2005). Conservation genetics of black and white ruffed lemurs, Varecia variegata, from Southeastern Madagascar. *Animal*

Conservation, 8(1), 105–111. https://doi.org/10.1017/S1367943004001891

Lusseau, D., & Newman, M. E. J. (2004). Identifying the role that animals play in their social networks. *Proceedings of the Royal Society B: Biological Sciences, 271*(SUPPL. 6), 477–481. https://doi.org/10.1098/rsbl.2004.0225

Lusseau, D., Wilson, B., Hammond, P. S., Grellier, K., Durban, J. W., Parsons, K. M., … Thompson, P. M. (2006). Quantifying the influence of sociality on population structure in bottlenose dolphins. *Journal of Animal Ecology, 75*(1), 14–24. https://doi.org/10.1111/j.1365-2656.2005.01013.x

Maher, C. R. (2009). Genetic relatedness and space use in a behaviorally flexible species of marmot, the woodchuck (Marmota monax). *Behavioral Ecology and Sociobiology, 63*(6), 857–868. https://doi.org/10.1007/s00265-009-0726-5

Mantel, N. (1967). The Detection of Disease Clustering and a Generalized Regression Approach. *Cancer Research, 27*(2 Part 1), 209 LP – 220. Retrieved from http://cancerres.aacrjournals.org/content/27/2_Part_1/209.abstract

McComb, K., Moss, C., Durant, S. M., Baker, L., & Sayialel, S. (2001). Matriarchs as repositories of social knowledge in African elephants. *Science, 292*, 491–494. https://doi.org/10.1126/science.1057895

McDonald, D. B. (2007). Predicting fate from early connectivity in a social network. *Proceedings of the National Academy of Sciences of the United States of America, 104*(26), 10910–10914. https://doi.org/10.1073/pnas.0701159104

McFarland, R., Fuller, A., Hetem, R. S., Mitchell, D., Maloney, S. K., Henzi, S. P., Barrett, L. (2015) Social integration confers thermal benefits in a gregarious primate. *Journal of Animal Ecology, 84*, 871–878.

Minta, S.C. (1992). Tests of spatial and temporal interactions among animals. *Ecological Applications*, 2, 178–188.

Mitani, J. C., Merriwether, D. A., & Zhang, C. (2000). Male affiliation, cooperation and kinship in wild chimpanzees. *Animal Behaviour, 59*(4), 885–893. https://doi.org/10.1006/anbe.1999.1389

Morland, H. (1991a). Social organization and ecology of black andwhite ruffed lemurs (Varecia variegata variegata) inlowland rain forest, Nosy Mangabe, Madagascar [disserta-tion]. New Haven: Yale University.

Morland, H. S. (1991b). Preliminary report on the social organi-zation of ruffed lemurs (Varecia variegata variegata) in anortheast Madagascar rain forest. Folia Primatologica56:157–161.

Moscovice, L. R., Douglas, P. H., Martinez-Iñigo, L., Surbeck, M., Vigilant, L., & Hohmann, G. (2017). Stable and fluctuating social preferences and implications for cooperation among female bonobos at LuiKotale, Salonga National Park, DRC. *American Journal of Physical Anthropology, 163*(1), 158–172. https://doi.org/10.1002/ajpa.23197

Mourier, J., Vercelloni, J., & Planes, S. (2012). Evidence of social communities in a spatially structured network of a free-ranging shark species. *Animal Behaviour, 83*(2), 389–401. https://doi.org/10.1016/j.anbehav.2011.11.008

Nystrand, M. (2007). Associating with kin affects the trade-off between energy intake and exposure to predators in a social bird species. *Animal Behaviour*, *74*(3), 497–506. https://doi.org/10.1016/j.anbehav.2006.11.031

Peakall, R., & Smouse, P. E. (2012). GenALEx 6.5: Genetic analysis in Excel. Population genetic software for teaching and research-an update. *Bioinformatics*, *28*(19), 2537–2539. https://doi.org/10.1093/bioinformatics/bts460

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V. and Vanderplas, J. (2011). Scikit-learn: Machine learning in Python. *Journal of machine learning research*, *12*, 2825-2830.

Perry, S. (2012). *The Behavior of Wild White-Faced Capuchins. Advances in the Study of Behavior* (1st ed., Vol. 44). Elsevier Inc. https://doi.org/10.1016/b978-0-12-394288-3.00004-6

Podgórski, T., Lusseau, D., Scandura, M., Sönnichsen, L., & Jędrzejewska, B. (2014). Long-lasting, kin-directed female interactions in a spatially structured wild boar social network. *PLoS ONE*, *9*(6), 1–11. https://doi.org/10.1371/journal.pone.0099875

Pope, T. R. (2000). Reproductive success increases with degree of kinship in cooperative coalitions of female red howler monkeys (Alouatta seniculus). *Behavioral Ecology and Sociobiology*, *48*(4), 253–267. https://doi.org/10.1007/s002650000236

Queller, D. C., & Goodnight, K. F. (1989). Estimating relatedness using genetic markers. *Evolution*, *43*(2), 258–275.

Radespiel, U., Lutermann, H., Schmelting, B., Bruford, M. W., & Zimmermann, E. (2003). Patterns and dynamics of sex-biased dispersal in a nocturnal primate, the grey mouse lemur, Microcebus murinus. *Animal Behaviour*, *65*(4), 709–719. https://doi.org/10.1006/anbe.2003.2121

Ramos-Fernández, G., Boyer, D., Aureli, F., & Vick, L. G. (2009). Association networks in spider monkeys (Ateles geoffroyi). *Behavioral Ecology and Sociobiology*, *63*(7), 999–1013. https://doi.org/10.1007/s00265-009-0719-4

Riehl, C. (2011). Living with strangers: direct benefits favour non-kin cooperation in a communally nesting bird. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1712), 1728–1735. https://doi.org/10.1098/rspb.2010.1752

Salpeteur, M., Patel, H., Balbo, A. L., Rubio-Campillo, X., Madella, M., Ajithprasad, P., & Reyes-García, V. (2015). When knowledge follows blood: Kin groups and the distribution of traditional ecological knowledge in a community of seminomadic pastoralists, Gujarat (India). *Current Anthropology*, *56*(3), 471–483. https://doi.org/10.1086/681006

Schülke, O., Bhagavatula, J., Vigilant, L., & Ostner, J. (2010). Social bonds enhance reproductive success in male macaques. *Current Biology*, *20*(24), 2207–2210. https://doi.org/10.1016/j.cub.2010.10.058

Seyfarth, R. M., & Cheney, D. L. (2012). The Evolutionary Origins of Friendship. *Annual Review of Psychology*, *63*(1), 153–177. https://doi.org/10.1146/annurev-psych-120710-100337

Silk, J. B. (2002). Kin selection in primate groups. *International Journal of*

Primatology, *23*(4), 849–875. https://doi.org/10.1023/A:1015581016205

Silk, J. B. (2007). The adaptive value of sociality in mammalian groups. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *362*(1480), 539–559. https://doi.org/10.1098/rstb.2006.1994

Silk, J. B., Alberts, S. C., & Altmann, J. (2003). Social Bonds of Female Baboons Enhance Infant Survival. *Science*, *302*, 1231–1234. https://doi.org/10.1126/science.1088580

Silk, J. B., Alberts, S. C., & Altmann, J. (2006). Social relationships among adult female baboons (Papio cynocephalus) II. Variation in the quality and stability of social bonds. *Behavioral Ecology and Sociobiology*, *61*(2), 197–204. https://doi.org/10.1007/s00265-006-0250-9

Silk, J. B., Altmann, J., & Alberts, S. C. (2006). Social relationships among adult female baboons (papio cynocephalus) I. Variation in the strength of social bonds. *Behavioral Ecology and Sociobiology*, *61*(2), 183–195. https://doi.org/10.1007/s00265-006-0249-2

Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., … Cheney, D. L. (2009). The benefits of social capital: Close social bonds among female baboons enhance offspring survival. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1670), 3099–3104. https://doi.org/10.1098/rspb.2009.0681

Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., … Cheney, D. L. (2010a). Female chacma baboons form strong, equitable, and enduring social bonds. *Behavioral Ecology and Sociobiology*, *64*(11), 1733–1747. https://doi.org/10.1007/s00265-010-0986-0

Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., … Cheney, D. L. (2010b). Strong and consistent social bonds enhance the longevity of female baboons. *Current Biology*, *20*(15), 1359–1361. https://doi.org/10.1016/j.cub.2010.05.067

Smith, J. E. (2014). Hamilton's legacy: Kinship, cooperation and social tolerance inmammalian groups. *Animal Behaviour*, *92*, 291–304. https://doi.org/10.1016/j.anbehav.2014.02.029

Spiegel, O., Leu, S. T., Sih, A., Bull, C. M. (2016). Socially interacting or indifferent neighbours? Randomization of movement paths to tease apart social preference and spatial constraints. *Methods in Ecolology and Evolution* 7, 971–979.

Spiegel, O., Sih, A., Leu, S. T., & Bull, C. M. (2018). Where should we meet? Mapping social network interactions of sleepy lizards shows sex-dependent social network structure. *Animal Behaviour*, *136*, 207–215. https://doi.org/10.1016/j.anbehav.2017.11.001

Stanley, C. R., Mettke-Hofmann, C., Hager, R., & Shultz, S. (2018). Social stability in semiferal ponies: networks show interannual stability alongside seasonal flexibility. *Animal Behaviour*, *136*, 175–184. https://doi.org/10.1016/j.anbehav.2017.04.013

Sterck, E. H. M., Watts, D. P., van Schaik, C. P. (1997) The evolution of female social relationships in nonhuman primates. *Behavior Ecology and Sociobiology*, 41, 291-309.

Stone, J., & Björklund, M. (2001). DELRIOUS: A computer program designed to analyse molecular marker data and calculate delta and relatedness estimates with confidence. *Molecular Ecology Notes*, *1*(3), 209–212. https://doi.org/10.1046/j.1471-8278.2001.00034.x

Strickland, K., Gardiner, R., Schultz, A. J., & Frère, C. H. (2014). The social life of eastern water dragons: Sex differences, spatial overlap and genetic relatedness. *Animal Behaviour*, *97*, 53–61. https://doi.org/10.1016/j.anbehav.2014.08.009

Strickland, K., Levengood, A., Foroughirad, V., Mann, J., Krzyszczyk, E., & Frère, C. H. (2017). A framework for the identification of long-term social avoidance in longitudinal datasets. *Royal Society Open Science*, *4*(8). https://doi.org/10.1098/rsos.170641

Taberlet, P. (1996). Reliable genotyping of samples with very low DNA quantities using PCR. *Nucleic Acids Research*, *24*(16), 3189–3194. https://doi.org/10.1093/nar/24.16.3189

Tecot, S. R., Baden, A. L., Romine, N., Kamilar, J. (2013). Reproductivestrategies and infant care in the Malagasy primates. In: Clancy K, Hinde K, Rutherford J, editors. Primate develop-mental trajectories in proximate and ultimate perspectives.Springer developments in primatology: progress and pros-pects book series. New York: Springer. p 321–359.

Vasey, N. (2006). Impact of seasonality and reproduction on social structure, ranging patterns and fission–fusion social organization in red ruffed lemurs. In *Lemurs: ecology and adaptation* (pp. 275–305). New York: Springer/Kluwer.

Wahaj, S. A., Van Horn, R. C., Van Horn, T. L., Dreyer, R., Hilgris, R., Schwarz, J., & Holekamp, K. E. (2004). Kin discrimination in the spotted hyena (Crocuta crocuta): Nepotism among siblings. *Behavioral Ecology and Sociobiology*, *56*(3), 237–247. https://doi.org/10.1007/s00265-004-0783-8

Wasserman, S., & Faust, K. (1994). *Structural analysis in the social sciences. Social network analysis: Methods and applications.* Cambridge University Press. https://doi.org/10.1017/CBO9780511815478

Wey, T., Blumstein, D. T., Shen, W., & Jordán, F. (2008). Social network analysis of animal behaviour: a promising tool for the study of sociality. *Animal Behaviour*, *75*(2), 333–344. https://doi.org/10.1016/j.anbehav.2007.06.020

Whitehead, H. (1996). Babysitting, dive synchrony, and indications of alloparental care in sperm whales. *Behavioral Ecology & Sociobiology*, *38*(4), 237–244.

Whitehead, H. (2008). *Analyzing Animal Societies: Quantitative methods for vertebrate social analysis.* Chicago: University of Chicago Press.

Whitehead, H. (2009). SOCPROG programs: Analysing animal social structures. *Behavioral Ecology and Sociobiology*, *63*(5), 765–778. https://doi.org/10.1007/s00265-008-0697-y

Wilkinson, G. S., Carter, G. G., Bohn, K. M., & Adams, D. M. (2016). Non-kin cooperation in bats. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *371*(1687). https://doi.org/10.1098/rstb.2015.0095

Wiszniewski, J., Allen, S. J., & Möller, L. M. (2009). Social cohesion in a hierarchically structured embayment population of Indo-Pacific bottlenose

1058  dolphins. *Animal Behaviour*, 77(6), 1449–1457.
1059  https://doi.org/10.1016/j.anbehav.2009.02.025
1060 Wolf, J.B.W. & Trillmich, F. (2007). Beyond habitat requirements: individual fine-
1061  scale site fidelity in a colony of the Galapagos sea lion (*Zalophus wollebaeki*)
1062  creates conditions for social structuring. *Oecologia*, 152, 553–567.
1063 Wright, P. C., Erhart, E. M., Tecot, S., Baden, A. L., Arrigo-nelson, S. J., Herrera,
1064  J., … Zohdy, S. (2012). *Long-Term Field Studies of Primates*.
1065  https://doi.org/10.1007/978-3-642-22514-7