1

[3]*Drosophila suzukii* wing spot size is robust to developmental temperature

[6]Ceferino Varón-González[1], Antoine Fraimout[1, 2], Vincent Debat[1]

[7][1]Institut de Systématique, Evolution, Biodiversité (ISYEB), Muséum National d'Histoire Naturelle, [8]CNRS, Sorbonne Université, EPHE, Université des Antilles, 57 rue Cuvier, CP 50, 75005 Paris, France

[9][2]Current affiliation: Ecological Genetics Research Unit, Organismal and Evolutionary Biology Research [10]Programme, Faculty of Biology and Environmental Sciences, Biocenter 3, Viikinkaari 1, FIN-0014, [11]University of Helsinki, Finland.

[14]Contact information:

[15]Ceferino Varón González, Institut de Systématique, Evolution, Biodiversité (ISYEB), Muséum National [16]d'Histoire Naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, 57 rue Cuvier, CP 50, [17]75005 Paris, France. Email: ceferino.vg@gmail.com

24ABSTRACT:

25Phenotypic plasticity is an important mechanism allowing adaptation to new environments and as such
26it has been suggested to facilitate biological invasions. Under this assumption, invasive populations
27are predicted to exhibit stronger plastic responses than native populations. *Drosophila suzukii* is an
28invasive species whose males harbor a spot on the wing tip. In this study, by manipulating
29developmental temperature, we compare the phenotypic plasticity of wing spot size of two invasive
30populations with that of a native population. We then compare the results with data obtained from wild-
31caught flies from different natural populations. While both wing size and spot size are plastic to
32temperature, no difference in plasticity was detected between native and invasive populations,
33rejecting the hypothesis of a role of the wing-spot plasticity in the invasion success. In contrast we
34observed a remarkable stability in the spot-to-wing ratio across temperatures, as well as among
35geographic populations. This stability suggests either that the spot relative size is under stabilizing
36selection, or that its variation might be constrained by a tight developmental correlation between spot
37size and wing size. Our data show that this correlation was lost at high temperature, leading to an
38increased variation in the relative spot size, particularly marked in the two invasive populations. This
39suggests (i) that *D. suzukii*'s development is impaired by hot temperatures, in agreement with the cold-
40adapted status of this species; (ii) that the spot size can be decoupled from wing size, rejecting the
41hypothesis of an absolute constraint and suggesting that the wing color pattern might be under
42stabilizing (sexual) selection; (iii) that such sexual selection might be relaxed in the invasive
43populations. Finally, a subtle but consistent directional asymmetry in spot size was detected in favor of
44the right side in all populations and temperatures, possibly indicative of a lateralized sexual behavior.

47*Keywords:* melanization, wing, spot, plasticity, directional asymmetry, temperature, *Drosophila suzukii*

 Introduction

53Phenotypic plasticity often plays an important role in the adaptation to new environments (Lande,

542015; West-Eberhard, 1989). In particular, it has been repeatedly suggested to facilitate invasions

55(Chown et al., 2007; Gibert et al., 2016; Richards et al., 2006), as genetic variation tends to be limited

56during the first stages of colonization (Geng et al., 2007; Geng et al., 2016). However, this possibility

57has received little empirical support so far (Arnold et al., 2019; Fox Rebecca J. et al., 2019). Invasions

58provide a particularly favorable context for the study of the evolutionary role of plasticity as we can

59compare derived populations repeatedly confronted to new environments with populations from the

60native range, arguably closer from the ancestral state. Differences in reaction norms are expected

61across populations (Pigliucci, 2005): native populations being predicted to be overall less plastic than

62invasive populations (Davidson et al., 2011; Lande, 2015; Lee and Gelembiuk, 2008; Richards et al.,

632006; Yeh and Price, 2004).

64 Biological invasions are processes in which an increasing number of individuals from a

65population colonize one environment where no representative of that population was present. The

66population effective size fluctuates in the new habitat and the population phenotype may change

67during invasions (Travis et al., 2009), as natural selection (in a different environment) would gain

68importance relative to drift. The limited population size during colonization may increase the

69importance of genetic drift and decrease selection on sexual traits (Candolin and Heuschele, 2008;

70Lahti et al., 2009). Such relaxed sexual selection might in turn be associated with an increased

71variability of such traits (Marshall et al., 2008), and in particular an increased plasticity (Price, 2006).

72 *Drosophila suzukii* has recently colonized all continents in a wide and fast dissemination all

73over the world (Fraimout et al., 2017). This species is a particularly suitable model to investigate the

74role of plasticity in the success of an invasion. Remarkably, *D. suzukii* males present a dark spot on

75the wing, as in several other Drosophila species (Kopp and True 2002). A precise understanding of the

76effect of this spot on mating success is lacking (Fuyama, 1979; Roy and Gleason, 2019), but the

77particular courtship behavior (the male confronts the female showing her his dorsal wing area)

78consistently associated with the presence of a spot in males across species, suggests an important

79role in sexual selection (Kopp and True, 2002).

80   The effect of sexual selection on phenotypic plasticity is controversial (Greenfield and

81Rodriguez, 2004; Møller and Alatalo, 1999; Rowe and Houle, 1996). Although sexually selected traits

82are often considered to be plastic (Price, 2006), an increased robustness might evolve through

83stabilizing or directional selection (see Fierst (2013) for a theoretical treatment, and Nieberding et al.

84(2018) for a case study in *Bicyclus anynana*). Stabilizing selection would limit variation around the

85values preferred by females (Mead and Arnold, 2004), while directional selection would limit variation

86on one side of the distribution due to female preferences and on the other side due to developmental

87or physiological constraints (Wiernasz, 1989). We reasoned that *D. suzukii*'s spot might be more

88plastic, or at least less canalized and thus more variable in the invasive populations compared to

89native populations, owing to a hypothetical reduced choosiness in females – and thus reduced sexual

90selection – possibly advantageous in small, peripheral populations (Bleu et al., 2012).

91   Here we quantify the phenotypic variation of the wing spot size in several natural populations

92of *D. suzukii* and assess its plasticity to temperature using samples from three geographic populations

93reared in the lab. Our experimental samples allow us to propose a causal link between temperature

94and phenotypic variation, while our natural populations would confirm the existence of such an

95association in nature. Temperature is one of the main environmental drivers of life history and

96morphological evolution of drosophilids in particular and ectotherms in general (Atkinson, 1994; Crill et

97al., 1996; David et al., 1997; Gibert et al., 1996; Gibert et al., 2007) and the thermal plastic response in

98*D. suzukii* is receiving much attention (Clemente et al., 2018; Fraimout et al., 2018; Shearer et al.,

992016). We test (1) whether natural populations present different spot sizes, and whether this

100differentiation correlates with local temperature at the time of capture, as would be expected for a

101plastic effect; (2) whether invasive and native populations display different plasticities to temperature;

102and (3) whether spot size is less canalized (more variable) in invasive populations. A higher plasticity

103of the spot in invasive populations is expected if plasticity plays a role in the invasion success.

104Combined with a reduced canalization, it might also indicate that sexual selection is relaxed in the

105invasive populations.

107 **Materials and Methods**

108 *Samples*

109Natural populations were sampled in 2014 and 2016 in 13 localities worldwide (see table 1) with baited

110traps and net sweeping, and conserved in ethanol in the dark (Fraimout et al., 2017). The number of

111individuals per locality ranged from 13 to 50 (Table 1). Sample sizes correspond to samples and

112pictures originally collected to run morphometric analyses.

113 The laboratory lines used here were adopted from an earlier study on *D. suzukii*'s wing

114plasticity to developmental temperature (see Fraimout *et al.* 2018 for details on rearing and

115maintenance). Briefly, three natural populations of *D. suzukii were* sampled in 2014 with banana bait

116traps and net swiping: one in the native area (Sapporo, Japan), and two in the invasion range, in Paris

117(France) and Dayton (Oregon, USA). Ten isofemale lines per population were produced from single

118matings of random pairs of individuals performed in separate vials (David et al., 1997). For each line

119from each population, we transferred F1 full-siblings individuals in multiple replicated vials and

120repeated the process for five generations in order to amplify stock populations prior to the temperature

121treatment. Stock populations were kept at 22°C throughout the experiment.

122 At the onset of the temperature treatment, we sampled random individuals from each line and

123transferred them in two sets of new vials. After 24h – and once oviposition was ensured – adult flies

124were removed and one set of vials was placed in an incubator set at 16°C, while the other set was

125placed in an incubator at 28°C. All vials were randomly placed within incubators and kept at each

126experimental temperature for the full development of the flies (i.e. egg to adult). The remaining

127temperature of our treatment (i.e. 22°C) corresponded to the one used for stock populations.

128 Between eight and 17 individuals per line were obtained (Table 2). Due to limitation on

129sample sizes, we did not take into account the lines in the analyses. We mounted wings on slides in a

130mixture of ethanol and glycerin and then we sealed the coverslips with nail polish and small weights to

131keep the wings as flat as possible. Images were taken with a Leica DFC 420 digital camera mounted
132on a Leica Z6 APO microscope controlling for the scale.

134   *Phenotypic data*

135Two phenotypic traits were directly inferred from the pictures: wing size and spot size (Figure 1).
136Because the potential visual effect of the wing spot during courtship may be associated to the
137proportion of the wing occupied by the spot and not its absolute size, we also estimated the relative
138spot size as the ratio between spot size and wing size for each individual. Using the ratio between spot
139and wing size for each individual can lead to erroneous conclusions due to allometry between traits (e.
140g. Packard and Boardman, 1988) but the estimation of residual values from a linear regression
141between wing and spot size gave virtually identical results (see online data).

142   In order to remove the proximal part of the wing, which is usually deteriorated during the
143removal of the wing from the thorax and may bias wing-size estimates, two landmarks were placed in
144ImageJ 1.51j8 (Rasband, 2012), that correspond to landmarks 1 and 3 in Fraimout et al. (2018) and a
145line between them was traced to discard all the proximal part of the wing. Then, a semi-automatized
146script in Python 3.6.1 was written to extract the wing and spot contours. Their respective areas were
147estimated by the Green's theorem in the function contourArea of the OpenCV library (Bradski and
148Kaehler, 2000) as the number of pixels within the wing and spot contours.

149   Our script was based on the manual adjustment of the picture color contrast. Because the
150color difference between the wing and the background is sharp, the wing outline was easily and
151robustly extracted. To the contrary, because the limits of the spot are fuzzy, a subtle difference in the
152image contrast can induce a relatively large difference in the spot outline. The manual adjustment of
153the picture contrast might thus be an important source of measurement error, particularly so for the
154spot size. We thus repeated the estimation of the wing and spot sizes in all the pictures and
155measurement error was quantified using an ANOVA for each population, assessing individual variation
156and measurement residuals.

159Natural populations - We applied a linear model to assess the relationship between absolute spot size

160and wing size, differences among natural populations on the spot size and whether there was natural

161variation in the spot-wing relationship (i. e. spot size ~ wing size + population + wing size x

162population). To have an estimate of natural variation for the other two phenotypic traits, we also

163assessed with one-way ANOVAs the differences among natural populations for wing size and the

164ratio. To explore the potential effect of temperature on those wild samples, we collected the average

165temperature during the month of collection in these places from the National Oceanic and Atmospheric

166Administration database (www.ncdc.noaa.gov/IPS/mcdw/mcdw.html) (Table 1) and ran a linear

167regression for each trait with temperature as a continuous variable.

168Experimental samples - We inferred the effect of temperature and geographic factors on each

169phenotypic trait with two-way ANOVAs (phenotype ~ temperature + population + temperature x

170population). The interaction temperature x population was used to assess the difference in plasticity

171among populations. We also ran a regression to test the effect of wing size on the absolute spot size,

172so that we can experimentally develop a null hypothesis about the relationship between wing and spot

173sizes that can be compared later with the results from natural populations. Because temperature, wing

174size and spot size are correlated (see results), we did not include two of these factors as explanatory

175variables in the same statistical model to avoid collinearity problems. This is the simplest approach

176that allows us to explore the unique contributions of temperature and wing variation to spot variation

177(Graham, 2003). The effect of temperature on the relationship between wing size and spot size was

178assessed by estimating only the effect of the interaction wing size x temperature on spot size in one

179ANCOVA. Differences in the slope of the relationship spot size-wing size among temperature levels

180with three pairwise t-test.

181   Differences in variance might reflect differences in the developmental channeling of

182environmental variation (Debat and David, 2001; Debat et al., 2009). In general, stressful

183environmental conditions have been suggested to alter developmental stability, leading to an increase

184in variance among individuals and asymmetry (Graham et al. 1998).

185   Among individual variation was measured as the coefficient of variation (CV) to account for a

186potential scaling effect on wing and spot size variation, and compared among temperatures within

187each geographic population. The significance of the differences was tested using a modified signed-

188likelihood ratio (MSLR) test for equality of coefficients of variation (Krishnamoorthy and Lee, 2014).For

189all the analyses described we averaged the phenotypic measurements obtained over the two wings

190within individual to avoid pseudo-replication.

191   To explore the hypothesis that temperature affects asymmetry, we measured the difference

192between right and left values for each of the three phenotypic traits (Table 1). That way, positive

193values in asymmetry would reflect larger phenotypes in the right wing and negative values larger

194phenotypes in the left wing. Then, a two-way mixed model ANOVA was applied to each trait (Palmer

195and Strobeck, 1986; Palmer 1994). "Individual" was entered into the model as a random effect, and

196tested for the variation among individuals; "side" was treated as a fixed effect, and tested for a

197systematic deviation from perfect symmetry (directional asymmetry, DA); the interaction "side x

198individual" tested for the significance of non-DA relative to measurement error (Fluctuating asymmetry,

199FA). The replicated measurements were included in the error term. Once the presence or absence of

200significant DA and FA were assessed in each particular group, we assessed the effect of temperature

201and geography on these two traits with an ANOVA (effect on DA). Differences in the slope of the

202relationship temperature-ratio asymmetry among populations with three pairwise t-test. Finally,

203Levene's tests were applied on the asymmetry measures to assess FA.

204   As effect sizes, Cohen's d for differences among geographic populations and r^2 for

205temperature effects were reported due to their simplicity (Rosenthal et al., 1994). All analyses were

206run in R version 3.5.2 (R Core Team 2013). All data, scripts and figures can be found in Dryad and

207Github.

209   **Results**

210*Natural populations*

211In natural populations, we found differences among populations for the spot size (ANOVA: $F_{12, 223}$ =

2122.480, p = 0.005) as well as for the wing (ANOVA: $F_{12, 236}$ = 41.330, p < 0.001) and the ratio (ANOVA:

213$F_{12, 236}$ = 3.092, p < 0.001). The latter showed a remarkable stability in comparison to the wing and the

214absolute spot size (r^2_{RATIO} = 0.14 r^2_{SPOT} = 0.49, $r^2_{WING\_POP}$ = 0.68).

Our linear model (spot size ~ wing size + population + wing size x population) also showed an
association between spot size and wing size in pooled samples (r^2 = 0.66, $F_{1, 223}$ = 525.217, p < 0.001)
(Figure 2). Whether the relationship between spot size and wing size depends on the population is not
conclusive (r^2 = 0.70, ANOVA: $F_{12, 223}$ = 1.748, p = 0.058). We detected a significant effect of
temperature on the three phenotypic traits (Figure 2), but much weaker for the ratio (wing size: r^2 =
0.39, $F_{1, 247}$ = 158.1, p < 0.001; spot size: r^2 = 0.21, $F_{1, 247}$ = 68.88, p < 0.001; ratio: r^2 = 0.03, $F_{1, 247}$ =
8.183, p = 0.005).

Experimental samples

Both wing size and spot size strongly decreased with temperature ($r^2_{WING\_TEMP}$ = 0.93, $F_{2, 116}$ = 886.278,
p < 0.001; $r^2_{SPOT\_TEMP}$ = 0.67, $F_{2, 121}$ = 4.986, p = 0.008) (Figure 3) and both traits are associated (r^2 =
0.73, t = 18.017, p < 0.001) (Figure 3). The ratio exhibited much less variation with temperature (r^2 =
0.08, $F_{2, 116}$ = 4.907, p = 0.009), the values at 22°C being slightly larger and driving the significance (no
difference between 16 and 28°C was detected: $F_{1, 75}$ = 1.243, p = 0.268).

Overall, the three traits exhibited very similar reaction norms across populations (Figure 3a
and 3c). A slight difference was nevertheless detected for wing size plasticity (interaction temperature
x population significant: $F_{4, 116}$ = 4.366, p = 0.003), the Sapporo population being slightly less
responsive to temperature ($r^2_{SAPPORO}$ = 0.92, r^2_{PARIS} = 0.94, r^2_{DAYTON} = 0.95). No difference in plasticity
among populations was found for the either the wing spot ($F_{4, 116}$ = 1.928, p = 0.110) or the ratio ($F_{, 116}$ =
0.508, p = 0.730).

Interestingly, the relationship between absolute spot size and wing size was dependent on
temperature (r^2 = 0.74, $F_{3, 121}$ = 117.4, p < 0.001): the lower the temperature, the higher the slope of the
regression and the tighter the association ($r^2_{16°}$ = 0.50, $r^2_{22°}$ = 0.30, $r^2_{28°}$ = 0.03) (Figure 3b). Although no
significant differences were found among slopes (16°C vs 22°C: $t_{1, 73}$ = -1.472, p = 0.145, 28°C vs
22°C: $t_{1, 88}$ = -0.417, p = 0.678, 28°C vs 16°C: $t_{1, 77}$ = -1.206, p = 0.232), this may be due to the sample
size limitations within temperature.

Within temperature variation tended to be higher at 28°C for both the spot size and the ratio,
although Sapporo did not show significant differences among temperatures for any of the phenotypic

243traits ($MSLRT_{SPOT}$ = 4.25, p = 0.119, $MSLRT_{WING}$ = 0.83, p = 0.661, $MSLRT_{RATIO}$ = 4.60, p = 0.100). For

244absolute spot size, variation was higher at 28°C in comparison to 22°C in all three geographic

245populations: Sapporo (Table 3), Dayton (Table 3, MSLRT = 12.01, p < 0.001) and Paris (Table 3,

246MSLRT = 3.42, p = 0.064). At 16°C, the other extreme temperature, Sapporo population also showed

247increased variation in the spot size (Table 3) but Paris (Table 3, MSLRT = 1.01, p = 0.315) and Dayton

248did not (Table 3, MSLRT = 0.06, p = 0.807). For the ratio, although all three geographic populations

249showed increased variation at 28° compared to 22° (Table 3), Sapporo showed reduced variability

250(Table 3) $F_{2, 40}$ = 1.02, p = 0.369). This might be explained by the fact that Sapporo showed very small

251wing variation among temperatures (Table 3; MSLRT = 0.83, p = 0.661).

252   Measurement error was much lower than the FA detected for all the populations and traits,

253ranging from twice lower for wing size in Dayton at 28°C to near 200 times lower for spot size in Paris

254at 28°. No difference in FA was detected for either the wing or the absolute spot sizes for laboratory

255samples ($F_{WING; 2, 103}$ = 2.449, p = 0.091; $F_{SPOT; 2, 103}$ = 1.166, p = 0.316) and for natural populations no

256difference in FA was found for any trait ($F_{WING; 12, 165}$ = 0.942, p = 0.507; $F_{SPOT; 12, 165}$ = 0.597, p = 0.843;

257$F_{RATIO; 12, 165}$ = 0.712, p = 0.417). For the ratio, the populations at 28° showed more FA than the

258populations at 16° ($SD_{RATIO 28°}$: 0.010, $SD_{RATIO 16°}$: 0.003, $F_{1, 69}$ = 6.070, p = 0.016) but the differences

259between 22° and 16° ($SD_{RATIO 22°}$: 0.005, $F_{1, 62}$ = 2.069, p = 0.155) and 28° and 22° ($F_{1, 75}$ = 3.543, p =

2600.064) were not significant.

261   No DA was detected for wing size. In contrast, significant DA was detected for both absolute

262and ratio, always in favor of the right side. For Paris and Dayton, DA was detected at 22 and 28°C

263while Sapporo only showed significant DA at 16°C. DA subtly increased with temperature for the ratio

264(r^2 = 0.06, $F_{2, 97}$ = 3.323, p = 0.040) and this effect was different depending on the population ($r^2_{TEMP x POP}$

265= 0.15, $F_{8, 97}$ = 2.095, p = 0.043). In particular, native population showed a lower response to

266temperature than invasive populations (Figure 3d, $d_{SAPPORO-PARIS}$ = 0.60, $d_{SAPPORO-DAYTON}$ = 0.49, $d_{DAYTON-}$

267$_{PARIS}$ = 0.21), although pairwise tests did not show significant differences in the slope of the relationship

268temperature-ratio asymmetry between Sapporo with either Paris ($t_{1,65}$ = 0.765, p = 0.447) or Dayton

269($t_{1,68}$ = 1.268, p = 0.209).

270   Most of the natural populations showed similar asymmetry patterns: while no DA was detected

271for wing size, both the absolute spot size and the ratio exhibited right biased DAs (Figure 2). The

272exceptions are the Dayton population, which also showed DA for the wing size, Langfang, which did

273not show DA for either the wing or the spot size, and Paris, which only showed wing size DA. DA was

274not associated with temperature in any trait (Wing: $F_{1,\,165} = 0.090$, $p = 0.765$; absolute spot size ($F_{1,\,165}$

275= 0.488, $p = 0.486$); ratio ($F_{1,\,165} = 0.143$, $p = 0.706$); Figure 2).

277   **Discussion**

278Our results revealed the existence of a relatively large natural variation in wing and spot sizes and

279temperature may explain a substantial part of it. In contrast, the spot-to-wing ratio was quite stable in

280nature and very robust to developmental temperature in the lab. This stability might either reflect a

281tight structural connection between the development of the spot and that of the wing, imposing a

282strong correlation (i.e. constraint hypothesis), or a history of stabilizing selection on the relative spot

283size (i.e. adaptive hypothesis). All three populations exhibited an increase in relative spot size variation

284at 28ºC, indicative of a lesser developmental robustness at high temperature, in agreement with the

285status of cold-adapted species of *D. suzukii* (Enriquez et al., 2018; Jakobs et al., 2015; Shearer et al.,

2862016; Stephens et al., 2015). This increased variation originating from a de-correlation of spot size and

287wing size at 28°C suggests that the stability of the relative spot size is not an absolute structural

288constraint, and therefore points at the adaptive hypothesis. *D. suzukii* males wing color pattern might

289thus be under stabilizing (sexual?) selection. Native and invasive populations showed very similar

290reaction norms, suggesting that spot plasticity is not affected by – and played no role in the success of

291– the invasion history. Relative spot size variability was higher in invasive populations compared to the

292native population, possibly pointing at a relaxed sexual selection during the invasion. Finally, both

293natural and laboratory samples exhibited a slight but significant DA of spot size (both absolute and

294relative), consistently in favor of the right side.

296*Relative spot size robustness: selection or constraint?*

297Although our experimental populations show a slight but significant decrease in the ratio at both high

298and low temperatures, the overall pattern reflects a global stability to developmental temperature,

299contrasting with both wing size and the absolute spot size that are both strongly plastic to temperature.

300   Such robustness might suggest either that spot size is fully determined by wing size, owing to 301the tight connection between wing development and the spot formation – an example of 302developmental constraint – or alternatively, that stabilizing selection is acting on *D. suzukii* males color 303pattern, removing individuals with extreme relative spot sizes.

304   The male-limited presence of the wing spot in all species has been interpreted as a mark of 305sexual selection (Kopp and True 2002). The presence of a spot is indeed phylogenetically tightly 306associated with the occurrence of elaborate courtship behaviors, the males dancing in a stereotypic 307way in front of females and displaying their wings (Kopp and True, 2002; Revadi et al., 2015). 308Although the empirical evidence of an effect of the spot on mating success is scarce (Fuyama, 1979; 309Roy and Gleason, 2019), it is nevertheless conceivable that females choice might be influenced by the 310male wing phenotype. Sexual selection might have favored the stability of the relative spot size and 311reduced its plasticity, a common process (Fierst, 2013) although not universal (Price, 2006). This 312would suggest that the focal trait might not be the spot size itself, but rather its extent on the wing, i.e. 313the wing color pattern.

314   Alternatively, it is possible that spot size plasticity is simply a structural consequence of wing 315size plasticity to temperature, which has been suggested to be adaptive (Crill et al., 1996; David et al., 3161997), and the fact that the boundaries of the expression of the genes involved in the spot formation 317vary according to the positional information of the veins (Arnoult et al., 2013; Gompel et al., 2005). 318Such developmental dependency of spot formation on wing development would thus impose a tight 319correlation between spot size and wing size, leading to a structurally stable spot relative size. This 320would thus be an example of an absolute constraint on the variation of the wing color pattern.

321   The analysis of the ratio variation within temperature provides an element in the discussion. It 322was indeed found that this variation increases at high temperature in the three populations (Figure 3c). 323This effect may indicate that hot temperatures might destabilize the processes involved in the spot 324formation, which is in agreement with the idea that *D. suzukii* is adapted to cold temperatures (Jakobs 325et al. 2015; Stephens et al. 2015; Shearer et al. 2016; Enriquez et al. 2018). The analysis of the 326covariation between wing size and spot size shows that they are tightly correlated. However, this 327correlation decreases at 28°C overall (Figure 3b), leading to the observed increase in variation of the

328ratio. This shows that the developmental link between wing and spot development can be disrupted

329(e.g. under hot temperature) suggesting that the stability of the spot relative size under colder

330(possibly optimal) temperatures might reflect more than a developmental constraint: i.e. might be

331maintained by stabilizing selection.

333*No role for plasticity to temperature in the invasion*

334Because plasticity may give a fitness advantage in early stages of invasion, populations repeatedly

335confronted to changing environments as invasive populations might present larger plastic responses.

336The hypothesis of adaptive plasticity in invasive populations (as opposed to native ones) has

337frequently appeared in the literature (Davidson et al., 2011; Lande, 2015; Lee and Gelembiuk, 2008;

338Richards et al., 2006; Yeh and Price, 2004). However, our results only show very small differences in

339wing size plasticity between invasive and native populations.

340   These results are consistent with some of the previous studies on the invasion of *D. suzukii.*

341No difference in thermal plasticity of wing shape, activity rhythms, gene expression and ovipositor

342shape was detected (Fraimout et al., 2018; Plantamp et al., 2019; Varón-González et al., 2019). There

343is is still scarce evidence that thermal plasticity might have contributed to the success of the invasion

344in this species (but see e. g. Everman et al. 2018). It is possible that *D. suzukii* might have been

345preadapted in its native range to the thermal conditions encountered throughout the invasion (Suarez

346and Tsuitsui, 2008), reducing the importance of temperature as a selective agent, but leaving open the

347hypothesis that plasticity relative to other parameters might have played a role in the invasion success

348(Hamby et al., 2016).

350*Directional asymmetry: affected by heat stress and indicative of a lateralized sexual behavior?*

351Both DA and FA have been suggested to increase under stress and have in turn been used as

352bioindicators of stressful environmental conditions (Graham et al., 1998; Klingenberg and Nijhout,

3531999), but see Houle (1998) for a critical discussion). Our results, showing an increase of the spot DA

354at high temperature may be circumstantial but their congruence with independent suggestions about

355the adaptation of *D. suzukii* populations to cold temperature (Enriquez et al., 2018; Jakobs et al., 2015;

356Shearer et al., 2016; Stephens et al., 2015) might advise further exploration.

357   The systematic DA in favor of the right side for the spot size is surprising. The small

358magnitude of DA might be a consequence of a regular but non adaptive effect (Pélabon and Hansen,

3592008). However, this consistent right-biased DA possibly points at a bias during courtship and mating

360that might be worth investigating, as suggested by QTL analyses linking behavior and wing

361pigmentation in *Drosophila elegans* and *Drosophila gunungcola* (Yeh and True, 2014). By altering a

362trait involved in a sexual behavior, stressful developmental conditions (temperature) might thus

363interfere with sexual selection.

Acknowledgements

366We would like to thank K. Tamura, M. Toda, P. Shearer, S. Fellous and T. Schlenke for their help

367during the fly sampling. We also thank M. Guillaume for her help with the fly stock maintenance and F.

368Peronnet for the rearing medium used. We thank Jennifer Gleason for her words on the reproduction

369of *D. suzukii* and the potential role of directional asymmetry on it. We also thank Thomas Flatt and two

370anonymous reviewers for the comments in a previous version of this article. CVG would like to thank

371Hacklab Almeria for their comments on the Python algorithm to estimate the wing and spot areas.

372CVG and AF were funded by the Agence Nationale de la Recherche (ANR) under the project SWING

373(ANR-16-CE02-0015).

Data accessibility

376All data, R scripts and reports of R scripts can be found in Dryad.

Competing interests

379The authors declare no competing or financial interests.

Author contributions

382Conceptualization: VD and CVG. Flies collection: AF and VD. Experimentation: AF. Statistical

383analysis: CVG. Discussion and writing: CVG and VD.

Literature cited

386**Arnold, P. A., Nicotra, A. B. and Kruuk, L. E. B.** (2019). Sparse evidence for selection on phenotypic
387   plasticity in response to temperature. *Philosophical Transactions of the Royal Society B:*
388   *Biological Sciences* **374**, 20180185.

389**Arnoult, L., Su, K. F. Y., Manoel, D., Minervino, C., Magriña, J., Gompel, N. and Prud'homme, B.**
390   (2013). Emergence and Diversification of Fly Pigmentation Through Evolution of a Gene
391   Regulatory Module. *Science* **339**, 1423.

392**Atkinson, D.** (1994). Temperature and organism size - a biological law for ectotherms. *Advances in*
393   *Ecological Research* **25**, 1–58.

394**Bleu, J., Bessa-Gomes, C. and Laloi, D.** (2012). Evolution of female choosiness and mating frequency:
395   effects of mating cost, density and sex ratio. *Animal Behaviour* **83**, 131–136.

396**Bradski, G. and Kaehler, A.** (2000). OpenCV. *Dr. Dobb's journal of software tools* **25**, 120–125.

397**Candolin, U. and Heuschele, J.** (2008). Is sexual selection beneficial during adaptation to
398   environmental change? *Trends in Ecology & Evolution* **23**, 446–452.

399**Chown Steven L, Slabber Sarette, McGeoch Melodie A, Janion Charlene and Leinaas Hans Petter**
400   (2007). Phenotypic plasticity mediates climate change responses among invasive and
401   indigenous arthropods. *Proceedings of the Royal Society B: Biological Sciences* **274**, 2531–
402   2537.

403**Clemente, M., Fusco, G., Tonina, L. and Giomi, F.** (2018). Temperature-induced phenotypic plasticity
404   in the ovipositor of the invasive species Drosophila suzukii. *Journal of Thermal Biology* **75**,
405   62–68.

406**Crill, W. D., Huey, R. B. and Gilchrist, G. W.** (1996). Within-and between-generation effects of
407   temperature on the morphology and physiology of Drosophila melanogaster. *Evolution* **50**,
408   1205–1218.

409**David, J. R., Gibert, P., Gravot, E., Petavy, G., Morin, J.-P., Karan, D. and Moreteau, B.** (1997).
Phenotypic plasticity and developmental temperature in Drosophila: Analysis and
significance of reaction norms of morphometrical traits. *Journal of Thermal Biology* **22**, 441–
451.

Davidson, A. M., Jennions, M. and Nicotra, A. B. (2011). Do invasive species show higher phenotypic
plasticity than native species and, if so, is it adaptive? A meta-analysis. *Ecology Letters* **14**,
419–431.

Debat, V. and David, P. (2001). Mapping phenotypes: canalization, plasticity and developmental
stability. *TRENDS in Ecology & Evolution* **16**, 555–561.

Debat, V., Debelle, A. and Dworkin, I. (2009). Plasticity, canalization, and developmental stability of
the Drosophila wing: joint effects of mutations and developmental temperature. *Evolution*
63, 2864–2876.

Enriquez, T., Renault, D., Charrier, M. and Colinet, H. (2018). Cold Acclimation Favors Metabolic
Stability in Drosophila suzukii. *Frontiers in Physiology* **9**, 1506.

Everman, E. R., Freda, P. J., Brown, M., Schieferecke, A. J., Ragland, G. J. and Morgan T. J. (2018).
Ovary development and cold tolerance of the invasive pest Drosophila suzukii (Matsumura) in the
centra plains of Kansas. Environmental entomology **47**, 1013-1023.

Fierst, J. L. (2013). Female mating preferences determine system-level evolution in a gene network
model. *Genetica* **141**, 157–170.

Fox Rebecca J., Donelson Jennifer M., Schunter Celia, Ravasi Timothy and Gaitán-Espitia Juan D.
(2019). Beyond buying time: the role of plasticity in phenotypic adaptation to rapid
environmental change. *Philosophical Transactions of the Royal Society B: Biological Sciences*
374, 20180174.

Fraimout, A., Debat, V., Fellous, S., Hufbauer, R. A., Foucaud, J., Pudlo, P., Marin, J.-M., Price, D. K.,
Cattel, J., Chen, X., et al. (2017). Deciphering the Routes of invasion of Drosophila suzukii by
Means of ABC Random Forest. *Molecular Biology and Evolution* **34**, 980–996.

Fraimout, A., Jacquemart, P., Villarroel, B., Aponte, D. J., Decamps, T., Herrel, A., Cornette, R. and
Debat, V. (2018). Phenotypic plasticity of Drosophila suzukii wing to developmental
temperature: implications for flight. *Journal of Experimental Biology* **221**, jeb166868.

Fuyama, Y. (1979). A visual stimulus in the courtship of Drosophila suzukii. *Experientia* **35**, 1327–
1328.

Geng, Y.-P., Pan, X.-Y., Xu, C.-Y., Zhang, W.-J., Li, B., Chen, J.-K., Lu, B.-R. and Song, Z.-P. (2007).
Phenotypic plasticity rather than locally adapted ecotypes allows the invasive alligator weed
to colonize a wide range of habitats. *Biological Invasions* **9**, 245–256.

Geng, Y., van Klinken, R. D., Sosa, A., Li, B., Chen, J. and Xu, C.-Y. (2016). The relative importance of
genetic diversity and phenotypic plasticity in determining invasion success of a clonal weed in
the USA and China. *Frontiers in Plant Sciences* **7**, 2113.

Gibert, P., Moreteau, B., Moreteau, J.-C. and David, J. R. (1996). Growth temperature and adult
pigmentation in two Drosophila sibling species: an adaptive convergence of reaction norms
in sympatric populations? *Evolution* **50**, 2346–2353.

Gibert, J.-M., Peronnet, F. and Schlötterer, C. (2007). Phenotypic plasticity in Drosophila
pigmentation caused by temperature sensitivity of a chromatin regulator network. *PLoS*
genetics **3**, e30.

Gibert, P., Hill, M., Pascual, M., Plantamp, C., Terblanche, J. S., Yassin, A. and Sgrò, C. M. (2016).
Drosophila as models to understand the adaptive process during invasion. *Biological*
Invasions **18**, 1089–1103.

Gompel, N., Prud'homme, B., Wittkopp, P. J., Kassner, V. A. and Carroll, S. B. (2005). Chance caught
on the wing: cis-regulatory evolution and the origin of pigment patterns in Drosophila.
Nature **433**, 481.

Graham, M. H. (2003). Confronting multicollinearity in ecological multiple regression. *Ecology* **84**,
2809–2815.

Graham, J. H., Emlen, J. M., Freeman, D. C., Leamy, L. J. and Kieser, J. A. (1998). Directional
asymmetry and the measurement of developmental instability. *Biological Journal of the*
Linnean Society **64**, 1–16.

Greenfield, M. D. and Rodriguez, R. L. (2004). Genotype–environment interaction and the reliability
of mating signals. *Animal Behaviour* **68**, 1461–1468.

Hamby, K. A., E. Bellamy, D., Chiu, J. C., Lee, J. C., Walton, V. M., Wiman, N. G., York, R. M. and
Biondi, A. (2016). Biotic and abiotic factors impacting development, behavior, phenology,
and reproductive biology of Drosophila suzukii. *Journal of Pest Science* **89**, 605–619.

Houle, D. (1998). High enthusiasm and low R-squared. *Evolution* **52**, 1872–1876.

Jakobs, R., Gariepy, T. D. and Sinclair, B. J. (2015). Adult plasticity of cold tolerance in a continental-
temperate population of Drosophila suzukii. *Journal of Insect Physiology* **79**, 1–9.

Klingenberg, C. P. and Nijhout, H. F. (1999). Genetics of fluctuating asymmetry: A developmental
model of developmental instability. *Evolution* **53**, 358–375.

Kopp, A. and True, J. R. (2002). Evolution of male sexual characters in the oriental Drosophila
melanogaster species group. *Evolution & development* **4**, 278–291.

Krishnamoorthy, K. and Lee, M. (2014). Improved tests for the equality of normal coefficients of
variation. *Computational Statistics* **29**, 215–232.

Lahti, D. C., Johnson, N. A., Ajie, B. C., Otto, S. P., Hendry, A. P., Blumstein, D. T., Coss, R. G.,
Donohue, K. and Foster, S. A. (2009). Relaxed selection in the wild. *Trends in Ecology &*
Evolution **24**, 487–496.

Lande, R. (2015). Evolution of phenotypic plasticity in colonizing species. *Molecular Ecology* **24**,
2038–2045.

Lee, C. E. and Gelembiuk, G. W. (2008). Evolutionary origins of invasive populations. *Evolutionary*
Applications **1**, 427–448.

Marshall, D. J., Bonduriansky, R. and Bussière, L. F. (2008). OFFSPRING SIZE VARIATION WITHIN
BROODS AS A BET-HEDGING STRATEGY IN UNPREDICTABLE ENVIRONMENTS. *Ecology* **89**,
2506–2517.

Mead, L. S. and Arnold, S. J. (2004). Quantitative genetic models of sexual selection. *Trends in*
Ecology & Evolution **19**, 264–271.

Møller, A. P. and Alatalo, R. V. (1999). Good-genes effects in sexual selection. *Proceedings of the*
Royal Society of London. Series B: Biological Sciences **266**, 85–91.

Nieberding, C. M., San Martin, G., Saenko, S., Allen, C. E., Brakefield, P. M. and Visser, B. (2018).
Sexual selection contributes to partial restoration of phenotypic robustness in a butterfly.
Scientific Reports **8**, 14315.

Packard, G. C. and Boardman, T. J. (1988). The misuse of ratios, indices, and percentages in
ecophysiological research. *Physiological Zoology* **61**, 1-9.

Pélabon, C. and Hansen, T. F. (2008). On the adaptive accuracy of directional asymmetry in insect
wing size. *Evolution* **62**, 2855–2867.

Pigliucci, M. (2005). Evolution of phenotypic plasticity: where are we going now? *TRENDS in Ecology*
and Evolution **20**, 481–486.

Plantamp, C., Henri, H., Andrieux, T., Régis, C., Mialdea, G., Dray, S., Gibert, P. and Desouhant, E.
(2019). Phenotypic plasticity in the invasive pest Drosophila suzukii: activity
rhythms and gene expression in response to temperature. *J. Exp. Biol.* **222**, jeb199398.

Price, T. D. (2006). Phenotypic plasticity, sexual selection and the evolution of colour patterns. *J. Exp.*
Biol. **209**, 2368.

Rasband, W. S. (2012). ImageJ: Image processing and analysis in Java. *Astrophysics Source Code*
Library.

Revadi, S., Lebreton, S., Witzgall, P., Anfora, G., Dekker, T. and Becher, G. P. (2015). Sexual Behavior
of Drosophila suzukii. *Insects* **6**,.

Richards, C. L., Bossdorf, O., Muth, N. Z., Gurevitch, J. and Pigliucci, M. (2006). Jack of all trades,
master of some? On the role of phenotypic plasticity in plant invasions. *Ecology Letters* **9**,
981–993.

Rosenthal, R., Cooper, H. and Hedges, L. (1994). Parametric measures of effect size. *The handbook*
of research synthesis **621**, 231–244.

Rowe, L. and Houle, D. (1996). The lek paradox and the capture of genetic variance by condition
dependent traits. *Proceedings of the Royal Society of London. Series B: Biological Sciences*
263, 1415–1421.

Roy, P. R. and Gleason, J. M. (2019). Assessing the use of wing ornamentation and visual display in
female choice sexual selection. *Behavioural Processes* **158**, 89–96.

Shearer, P. W., West, J. D., Walton, V. M., Brown, P. H., Svetec, N. and Chiu, J. C. (2016). Seasonal
cues induce phenotypic plasticity of Drosophila suzukii to enhance winter survival. *BMC*
ecology **16**, 11.

Stephens, A. R., Hutchison, W. D., Asplen, M. K. and Venette, R. C. (2015). Cold Hardiness of Winter-
Acclimated Drosophila suzukii (Diptera: Drosophilidae) Adults. *Environmental Entomology* **44**,
1619–1626.

Suarez, A. P. and Tsuitsui, N. D. (2008). The evolutionary consequences of biological invasions.
Molecular Ecology **17**, 351–360.

Travis, J. M. J., Mustin, K., Benton, T. G. and Dytham, C. (2009). Accelerating invasion rates result
from the evolution of density-dependent dispersal. *Journal of Theoretical Biology* **259**, 151–
158.

Varón-González, C., Fraimout, A., Delapré, A., Debat, V. and Cornette, R. (2019). Limited thermal
plasticity and geographic divergence in the ovipositor of Drosophila suzukii.
bioRxiv 670356.

West-Eberhard, M. J. (1989). Developmental plasticity and the origins of diversity. *Annual Review in*
Ecology and Systematics **20**, 249–278.

Wiernasz, D. C. (1989). Female choice and sexual selection of male wing melanin pattern in Pieris
occidentalis (Lepidoptera). *Evolution* **43**, 1672–1682.

Yeh, P. J. and Price, T. D. (2004). Adaptive Phenotypic Plasticity and the Successful Colonization of a
Novel Environment. *The American Naturalist* **164**, 531–542.

Yeh, S.-D. and True, J. R. (2014). The Genetic Architecture of Coordinately Evolving Male Wing
Pigmentation and Courtship Behavior in Drosophila elegans and Drosophila
gunungcola. *G3: Genes|Genomes|Genetics* **4**, 2079.

Tables

Table 1

Natural populations sample sizes

Collection place	Country	Date (M/Y)	Data place	Temp (°C)	Number of wings	Complete individuals
Barcelona	Spain	07/2014	Barcelona	24.1	49	18
Porto Alegre	Brazil	05/2014	Porto Alegre	16.8	21	7
Geneva	USA	05/2014	Geneva	16.7	30	15
Trento	Italy	09/2014	Paganella	7.3	25	11
Langfang	China	08/2014	Beijing	26.3	30	15
Liaoyuan	China	04/2014	Shenyang	24.2	30	12
Sapporo	Japon	07/2014	Sapporo	22.5	41	16
Shiping	China	05/2014	Mengzi	23.0	28	13
Dayton	USA	10/2014	Portland	15.6	48	20
Tokyo	Japon	07/2014	Tokyo	26.8	13	3
Watsonville	USA	10/2014	Fresno	22.2	38	15
Madison	USA	10/2016	Madison	12.1	25	11
Paris	France	10/2016	Paris	10.9	50	22

549Note: The first three columns represent the nearest city, country and date of the flies collection. Then,
550the closest city to the collection place and its temperature during the month of the collection are
551represented. Finally, the table recalls the total sample size (wings) and the number of complete
552individuals (pairs of wings from the same individual).

554 **Table 2**

555 Lab populations sample sizes

Geographic origin	Temperature (°C)	Number of wings	Complete individuals	Lines
Paris	16	15	7	8
	22	27	12	15
	28	28	13	15
Sapporo	16	25	12	13
	22	26	12	14
	28	31	15	16
Dayton	16	22	10	12
	22	26	11	15
	28	31	14	17

557Note: The first two columns show the geographic origin of each population (first) and the temperature

558set during their development *in-vitro* (second). Then, the total number of wings used to analyze overall

559variation and the number of complete individuals used to analyze asymmetric variation.

561 **Table 3**

562 Coefficient of variation for each trait per population and temperature

		Sapporo	Dayton	Paris
Wing	16°	0.06	0.05	0.04
	22°	0.05	0.07	0.06
	28°	0.06	0.04*	0.04
Spot	16°	0.19	0.10*	0.10*
	22°	0.11	0.10*	0.14**
	28°	0.20	0.29	0.25
Ratio	16°	0.14*	0.09*	0.08*
	22°	0.11*	0.08*	0.12*
	28°	0.20	0.27	0.25

563Note: Asterisks represents a significant difference for a phenotype (*:α = 0.05, **: α = 0.1), within a

564population, between the temperature presenting the largest coefficient of variation and the other two

565temperatures.

Figure legends

569**Figure 1:** *Drosophila suzukii* wing from a population of Sapporo (Japan, native population) raised at

57028°. In blue there is the wing and spot areas identified by our Python algorithm.

572**Figure 2:** Relationship between wing and spot variation in natural populations and temperature.

573Temperature was measured in °C and phenotypic sizes correspond to the number of pixels within

574wing and spot areas in the pictures. The ratio and ratio asymmetry are based on a proportion between

575the former two. For relationships between the variables stated in the diagram axes, *p* and r^2 were

576reported. Association between a) wing size and temperature, b) spot size and temperature , c)

577between the ratio and temperature and d) ratio asymmetry and temperature .

579**Figure 3:** Effect of temperature on the different phenotypic traits in experimental samples.

580Temperature was measured in °C and phenotypic sizes correspond to the number of pixels within

581wing and spot areas in the pictures. The ratio and ratio asymmetry are based on a proportion between

582the former two. For relationships between the variables stated in the diagram axes, *p* and r^2 were

583reported. a) linear regression showing the influence of developmental temperature on wing variation,

584b) linear regression showing the influence of developmental temperature on spot size variation, c)

585effect of temperature on the relationship spot size – wing size, d) differences in the ratio for each

586population among temperatures, e) differences in ratio asymmetry for each population among

587temperatures.