High and stable ATP levels prevent aberrant intracellular protein aggregation

Masak Takaine[1,2]*, Hiromi Imamura[3], Satoshi Yoshida[1,2,4,5]*

[1]Gunma University Initiative for Advanced Research (GIAR), Gunma University, Maebashi

371-8512, Japan

[2]Institute for Molecular and Cellular Regulation (IMCR), Gunma University, Maebashi

371-8512, Japan

[3]Graduate School of Biostudies, Kyoto University, Kyoto 606-8501, Japan

[4]School of International Liberal Studies, Waseda University, Tokyo, 169-8050, Japan

[5]Japan Science and Technology Agency, PREST

*Correspondence should be addressed to: masaktakaine@gunma-u.ac.jp (M.T.) (Lead

contact) or satosh@waseda.jp (S.Y.)

Running title

ATP homeostasis prevents cytotoxic protein aggregation

Keywords

ATP, homeostasis, proteostasis, yeast, neurodegenerative disease, AMPK, adenylate kinase,

protein aggregation

ABSTRACT

ATP at millimolar levels has recently been implicated in the solubilization of cellular proteins. However, the significance of this high ATP level under physiological conditions and the mechanisms that maintain ATP remain unclear. We herein demonstrated that AMP-activated protein kinase (AMPK) and adenylate kinase (ADK) cooperated to maintain cellular ATP levels regardless of glucose levels. Single cell imaging of ATP-reduced yeast mutants revealed that ATP levels in these mutants repeatedly underwent stochastic and transient depletion, which induced the cytotoxic aggregation of endogenous proteins and pathogenic proteins, such as huntingtin and α-synuclein. Moreover, pharmacological elevations in ATP levels in an ATP-reduced mutant prevented the accumulation of α-synuclein aggregates and its cytotoxicity. The removal of cytotoxic aggregates depended on proteasomes, and proteasomal activity cooperated with AMPK or ADK to resist proteotoxic stresses. The present study is the first to demonstrate that cellular ATP homeostasis ensures proteostasis and revealed that suppressing the high volatility of cellular ATP levels prevented cytotoxic protein aggregation, implying that AMPK and ADK are important factors that prevent proteinopathies, such as neurodegenerative diseases.

INTRODUCTION

Adenosine triphosphate (ATP) is the main energy currency used by all living organisms. In addition to its role as energy currency, ATP has recently been suggested to influence the balance between the soluble and aggregated states of proteins, indicating that proteostasis is maintained by energy-dependent chaperones and also by the property of ATP as a hydrotrope to solubilize proteins (Hayes, Peuchen et al., 2018, Patel, Malinovska et al., 2017, Pu, Li et al., 2019, Sridharan, Kurzawa et al., 2019). Furthermore, ATP levels have been shown to regulate the physicochemical properties of the cytoplasm, such as viscosity, macromolecular crowding, and liquid-liquid phase separation (Marini, Nuske et al., 2020, Persson, Ambati et al., 2020). However, the role of ATP was assessed in these studies using artificial ATP depletion. Therefore, it currently remains unclear whether ATP-dependent protein solubilization/desolubilization have physiologically significant cellular roles.

We recently established a reliable imaging technique to quantify intracellular ATP levels in single living yeast cells using the genetically encoded fluorescent ATP biosensor QUEEN (Yaginuma, Kawai et al., 2014), which enables long-term evaluations of ATP homeostasis in individual cells (Takaine, Ueno et al., 2019). The findings obtained demonstrated that intracellular ATP levels did not vary within a yeast population grown in the same culture (Takaine et al., 2019), which was in contrast to the large variations observed in intracellular ATP levels within a bacterial cell population (Yaginuma et al., 2014). Moreover, intracellular ATP levels in individual living yeast cells were stably and robustly maintained at approximately 4 mM, irrespective of carbon sources and cell cycle stages, and temporal fluctuations in intracellular ATP levels were small (Takaine et al., 2019). Based on these findings, we hypothesized that an exceptionally robust mechanism exists to precisely regulate ATP levels in eukaryotes. It currently remains unclear why ATP is stably maintained at a markedly higher level than the K_m (Michaelis constant) required for the enzymatic activities of most ATPases (Edelman, Blumenthal et al., 1987), and the consequences associated with failed ATP homeostasis in living organisms have not yet been elucidated.

The most promising candidate regulator of ATP homeostasis is AMP-activated protein kinase (AMPK). AMPK, which is activated by AMP and inhibited by ATP (Xiao, Heath et al., 2007), has long been regarded as an important regulator of the whole-body and cellular energy status in eukaryotes (Hardie, Schaffer et al., 2016). AMPK is activated by increases in the AMP:ATP or ADP:ATP ratio (*i.e.,* low-energy state), and regulates its downstream effectors by phosphorylation to redirect cell metabolism from an anabolic (ATP-consuming) state to

catabolic (ATP-producing) state (Herzig & Shaw, 2017). In the budding yeast *Saccharomyces*

cerevisiae, the sucrose non-fermenting 1 (Snf1) protein kinase complex is the sole AMPK.

Similar to other AMPKs, the yeast Snf1 complex comprises three subunits: the catalytic α

subunit (*SNF1*), scaffolding β subunit (*SIP1*, *SIP2* or *GAL83*), and regulatory γ subunit

(*SNF4*) (Ghillebert, Swinnen et al., 2011). The role of the Snf1 complex in adaptation to

glucose limitations has been characterized in detail (Hedbacker & Carlson, 2008). The Snf1

complex is inactive in the presence of sufficient glucose levels in media (Wilson, Hawley et

al., 1996). Decreases in glucose levels have been shown to activate the Snf1 complex and

phosphorylate the transcriptional repressor Mig1, which then triggers the transcription of

numerous glucose-repressed genes (Carlson, 1999). However, the contribution of AMPK or

the Snf1 complex to cellular ATP levels remains unknown.

Other possible candidate regulators of ATP homeostasis include genes whose mutation leads

to decreases in the cellular content of ATP. However, based on biochemical analyses of cell

populations, few yeast mutants reduced ATP levels (Gauthier, Coulpier et al., 2008,

Ljungdahl & Daignan-Fornier, 2012). Adenylate kinase (ADK) is a key enzyme that

synthesizes ATP and AMP using two ADP molecules as substrates, and the null mutant of

ADK (*adk1Δ*) was shown to have a reduced cellular ATP level (~70% of the wild type)

(Gauthier et al., 2008). Bas1 is a transcription factor that is required for *de novo* purine

synthesis and *bas1Δ* also has a reduced ATP level (~50% of the wild type) (Gauthier et al.,

2008). However, the regulation of ATP levels and the physiological consequences of reduced

ATP levels in these mutants remain unclear, particularly at the single cell level.

In the present study, we investigated the roles of AMPK, ADK, and Bas1 in ATP homeostasis

using the QUEEN-based single cell ATP imaging technique. We demonstrated for the first

time that AMPK is involved in the regulation of intracellular ATP levels, even under

glucose-rich conditions. Furthermore, time-lapse ATP imaging revealed that cells lacking

both AMPK and ADK frequently underwent transient ATP depletion, while ATP levels

oscillated in those lacking Bas1. These ATP dynamics in the mutants were overlooked in

previous biochemical studies. We found that some intrinsic proteins and aggregation-prone

model proteins, including α-synuclein, which is responsible for Parkinson's disease,

aggregated and were cytotoxic in all of the ATP-reduced mutants tested. The present results

suggest that the stable maintenance of ATP is essential for proteostasis and imply that an ATP

crisis promotes proteinopathies, such as neurodegenerative diseases.

RESULTS

ADK1 cooperates with AMPK to regulate ATP homeostasis

We recently developed a reliable monitoring system for cytoplasmic ATP levels in living yeast cells using the ATP biosensor QUEEN (Takaine et al., 2019). We herein conducted a more detailed examination of ATP dynamics in wild-type and mutant yeast cells using this system. We initially investigated whether the deletion of *SNF1*, which encodes a catalytic subunit of AMPK, affected cellular ATP levels. Cellular ATP levels were significantly lower in *SNF1*-null mutant (*snf1Δ*) cells than in wild-type cells at various glucose levels (Fig. 1A and Fig. S1A, B). It is important to note that in addition to low glucose conditions (0.05% glucose), at which the Snf1 complex is active, *snf1Δ* cells showed significantly reduced ATP levels even under high glucose conditions (2% glucose), at which the Snf1 complex is considered to be inactive. The deletion of *MIG1* had a negligible effect on ATP levels (Fig. S1C), suggesting that as yet unknown factors other than Mig1 primarily regulate ATP levels under the control of the Snf1 complex. Collectively, these results demonstrated for the first time that AMPK/SNF1 affect cellular ATP levels even under glucose-rich conditions.

ADK catalyzes the interconversion of adenine nucleotides (ATP+AMP $\longleftrightarrow$ 2ADP), which is important for *de novo* adenine nucleotide synthesis and the balance between ATP, ADP, and AMP. Previous biochemical studies reported that the deletion of the ADK gene reduced ATP levels in mouse skeletal muscle cells and budding yeasts (Gauthier et al., 2008, Janssen, Dzeja et al., 2000). We confirmed these findings using an ATP imaging method: *adk1Δ* cells showed significantly lower QUEEN ratios than wild-type cells on average in the presence of sufficient carbon sources (Fig. S2).

In addition to being a key enzyme in purine metabolism, ADK has also been suggested to cooperate with AMPK in order to monitor the cellular energy state (Hardie, Carling et al., 1998). Therefore, we compared ATP levels in *snf1Δ adk1Δ* double mutant cells with those in *snf1Δ*, *adk1Δ* cells and wild-type cells (Fig. 1A, B). *snf1Δ adk1Δ* cells had significantly lower average ATP levels than single mutant cells. We also found not only a general reduction, but also a marked variation in ATP levels in the *snf1Δ*, *adk1Δ*, *snf1Δ adk1Δ* cell population, as indicated by the large coefficient of variance (CV) (Fig. 1B). Furthermore, some *snf1Δ adk1Δ* cells had very low ATP levels (Fig. 1A, B). These results suggest that Adk1 and the Snf1 complex both synergistically contribute to ATP homeostasis.

We confirmed the decreases observed in ATP levels in *snf1Δ*, *adk1Δ* and *snf1Δadk1Δ* cells

using a biochemical assay of whole cell extract (Fig. 1C). ADP levels were also reduced in

these mutant cells (Fig. S3). In *adk1Δ* cells, the ATP/ADP ratio increased, whereas the sum of

ATP and ADP levels decreased (Fig. S3), which is consistent with previous findings

(Gauthier et al., 2008). These biochemical data and their relevance to QUEEN data are

discussed later.

A large pool of adenine nucleotides is important for maintaining cellular ATP levels

We examined a *bas1Δ* mutant, which is defective in the expression of genes responsible for

adenine biogenesis (Daignan-Fornier & Fink, 1992, Denis, Boucherie et al., 1998). Consistent

with previous biochemical findings (Gauthier et al., 2008), ATP levels quantified by QUEEN

were reduced by ~50% in *bas1Δ* cells (Fig. 1D, E). We found not only a general reduction,

but also a marked variation in ATP levels in the *bas1Δ* cell population, as indicated by the

large CV (Fig. 1E). The decrease observed in ATP levels was due to reduced adenine

biosynthesis because the addition of extra adenine to media partially restored ATP levels (Fig.

1E). These results suggest that the sufficient production of adenine nucleotides is essential for

the stable maintenance of ATP levels. Moreover, the role of Bas1 in maintaining ATP levels

appeared to be epistatic to that of Snf1 because *bas1Δ snf1Δ* double mutant cells showed a

similar distribution of ATP levels to *bas1Δ* cells (Fig. S4). We confirmed the decrease

observed in ATP levels in *bas1Δ* cells using a biochemical assay (Fig. 1F). ADP levels and

the sum of ATP and ADP levels also significantly decreased in *bas1Δ* cells (Fig. S3), which is

consistent with previous findings (Gauthier et al., 2008).

ATP levels temporally fluctuate in ATP mutant cells

To investigate the mechanisms contributing to the marked variations in ATP levels in *snf1Δ*

adk1Δ cells in more detail, we employed time-lapse ATP imaging (Fig. 2). We found that the

QUEEN ratio often underwent a rapid decline followed by recovery in *snf1Δ adk1Δ* cells (see

116 and 132 min in Fig. 2A, C, and Movie S1, and 180 and 356 min in Fig. 2B, D, and Movie

S2). The sudden decrease in ATP levels (hereafter called "the ATP catastrophe") occurred

within a few minutes without any sign and was never observed in wild-type cells (Takaine et

al., 2019). The ATP catastrophe appeared to be a stochastic event and cell intrinsic: these

events occurred independent of the cell cycle stage or cell size (compare Fig. 2C with D).

Under some conditions, the QUEEN ratio did not recover after the ATP catastrophe and the

cell died, as judged by the loss of QUEEN signals in the cell (Fig. S5). These results suggest

that the marked variations observed in ATP levels in *snf1Δ adk1Δ* cells were not simply due

to a mixed population with different basal ATP levels, but were rather caused by the

stochastic ATP catastrophe in individual cells.

Time-lapse imaging of *bas1Δ* revealed oscillatory cycles in ATP levels (Figs. 2E and S6A,

and Movies S3 and S4): ATP cycling in *bas1Δ* cells was slow (~35 min on average, Fig. S6B)

and distinguishable from that in *snf1Δ adk1Δ* cells; however, the common characteristics of

these mutants were that the level of ATP repeatedly reached close to 0 mM. The ATP

oscillation cycle was unsynchronized in the population and independent of cell cycle

progression, suggesting a unique metabolic rhythm intrinsic to each cell. The oscillatory

nature of ATP cycling in the *bas1Δ* mutant may involve a transcription/translation cycle and

will be described elsewhere.

ATP homeostasis is required for preventing protein aggregation *in vivo*

We recently reported that cellular ATP levels were stably maintained at ~4 mM in budding

yeast (Takaine et al., 2019) and herein demonstrated that Adk1 and Bas1 in the Snf1/AMPK

complex were required for the regulation of ATP homeostasis. However, the physiological

importance of ATP homeostasis remains unknown. To clarify the significance of high ATP

levels, we examined the global genetic interactions of *snf1Δ, adk1Δ,* and *bas1Δ* using

CellMap ((Usaj, Tan et al., 2017), thecellmap.org). An *in silico* analysis identified genes

involved in "protein folding/glycosylation" as common negative genetic interactors with

adk1Δ and *bas1Δ* (Fig. 3A). Negative genetic interactors of *ura6*, a gene encoding uridylate

kinase that also exhibits ADK activity, were enriched in the "protein folding/glycosylation"

category (Fig. 3A). We also found that interactors of *snf1* were implicated in "protein

folding/glycosylation". None of these mutants exhibited apparent genetic interactions with

genes in the "metabolism" category (Fig. 3A). The same analysis using genetic and physical

interactors provided similar results and showed that many interactors were enriched in the

"protein turnover" category (Fig. S7). These results imply that although these three mutants

regulate ATP with distinct mechanisms, all three have a common cellular function.

To examine possible defects in protein folding and turnover (*i.e.*, proteostasis), we challenged

these mutants with various proteotoxic stresses. We found negligible growth defects in ATP

mutants under normal growth conditions with 2% glucose at 30°C (control in Fig. 3B),

suggesting that a high level of ATP is not necessary for cellular growth. However, the *adk1*

and *bas1* mutants both exhibited severe growth defects with a high temperature of 40°C, 1

hour of heat shock at 55°C, or in the presence of 0.5 μg/ml of the glycosylation inhibitor

tunicamycin or 2 mM H_2O_2. The *SNF1* deletion increased the stress sensitivity of *adk1Δ* (Fig.

3B). This sensitivity to proteotoxic stress suggests that ATP homeostasis mutants are

defective in some aspects of proteostasis. We found that all four mutants tested contained

significantly increased numbers of Hsp104-GFP foci, a marker of protein aggregation

(Josefson, Andersson et al., 2017) (Fig. 3C, D). In contrast to Hsp104-GFP foci, Pab1-GFP, a

marker of stress granule (SG) assembly (Hoyle, Castelli et al., 2007), did not form foci in

ATP mutants, suggesting that protein aggregation and SG assembly are regulated in a distinct

manner (Fig. 3D). These analyses identified abnormal protein aggregation as a common

defect associated with ATP homeostasis mutants for the first time.

The transient depletion of ATP leads to the formation of protein aggregates

To examine whether ATP depletion triggers protein aggregation in living yeast, we artificially

depleted cellular ATP levels by replacing glucose with 2-deoxyglucose (2DG), a strong

inhibitor of glycolysis, in media and monitored protein aggregation using Hsp104-GFP as a

marker of protein aggregation (Josefson et al., 2017) in wild-type cells. We previously

showed that ATP levels were almost completely depleted 2 minutes after the 2DG treatment

(Takaine et al., 2019), which was also confirmed biochemically (Fig. S3B). Within 15 min of

the 2DG treatment, more than 20% of cells contained Hsp104-GFP foci indicative of protein

aggregation (Fig. 4A, B). These protein aggregations were retained for hours after refeeding

of glucose (Fig. 4A, B), suggesting that the dissolution kinetics of Hsp104-GFP were

significantly slow. This contrasts intracellular ATP, which recovers to normal levels within 1

min of glucose refeeding (Takaine et al., 2019).

To further confirm whether a high level of ATP is required for protein solubility, we also

tested the Ubc9-ts protein, a model protein that is prone to aggregation (Kaganovich, Kopito

et al., 2008), and found that ATP depletion by the 2DG treatment triggered Ubc9-ts protein

aggregation (Fig. 4C). Therefore, not only Hsp104-GFP-positive intrinsic proteins, but also

extrinsic model proteins aggregate after ATP depletion.

SG are assembled in budding yeast cells under stress conditions, such as glucose depletion

(Hoyle et al., 2007). In contrast to the formation of Hsp104-GFP foci, ATP depletion after the

2DG treatment did not instantly trigger the formation of SG (Fig. S8). Consistent with recent

findings (Jain, Wheeler et al., 2016), the present results suggest that SG formation requires

ATP. We also noted that Hsp104-GFP foci and SG did not co-localize, indicating that these

structures are derived from distinct mechanisms (Fig. S8). Therefore, the artificial depletion

of ATP may trigger abnormal protein aggregation in living yeast cells.

ATP homeostasis is required for the protection of cells from cytotoxicity caused by protein aggregation

Protein aggregation is often associated with neurodegenerative diseases, such as Alzheimer's,

Huntington's, and Parkinson's diseases (Eftekharzadeh, Hyman et al., 2016). Mitochondrial

failure has also been associated with many neurodegenerative diseases; however, it currently

remains unclear whether energy failure causes protein aggregation because mitochondria also

produce cytotoxic reactive oxygen species (ROS). (Bhat, Dar et al., 2015, Pathak, Berthet et

al., 2013). The abnormal aggregation of α-synuclein has been implicated in Parkinson's

disease (Lashuel, Overk et al., 2013). To clarify whether ATP prevents the formation of

cytotoxic protein aggregation, we examined the toxicity of α-synuclein-GFP (Syn-GFP) in

budding yeast. As reported previously, the expression of Syn-GFP exhibited negligible

toxicity against wild-type yeast when expressed under the inducible *GAL1* promotor (Fig. 5A)

(Outeiro & Lindquist, 2003, Sharma, Brandis et al., 2006, Wijayanti, Watanabe et al., 2015).

However, *snf1Δ*, *adk1Δ*, and *bas1Δ* were hypersensitive to the expression of Syn-GFP (Fig.

5A). We also found that *rpn4Δ*, which encodes a key transcription factor for proteasomal

subunits (Xie & Varshavsky, 2001), was very sensitive to Syn-GFP (Fig. 5A), which is

consistent with the concept that Syn-GFP is degraded in the ubiquitin-proteasomal pathway in

yeast (Tofaris, Kim et al., 2011, Wijayanti et al., 2015). We then visualized the cellular

localization of Syn-GFP. Consistent with previous findings (Willingham, Outeiro et al., 2003),

Syn-GFP expressed in yeast mainly localized to the plasma membrane (Fig. 5B, C). In

addition to the plasma membrane, we found that Syn-GFP localized to punctate structures,

most likely corresponding to protein aggregation (Fig. 5B, C). Punctate structures were not as

obvious in *rpn4Δ* cells defective in proteasomes, suggesting that the accumulation of

Syn-GFP puncta was not simply due to defective degradation.

To investigate whether a high level of ATP protects cells from toxic protein aggregation, we

added extra adenine to the medium (Fig. 5D). A previous study demonstrated that the addition

of 300 μM adenine to the medium increased ATP levels from 4 to 5.5 mM in wild-type cells

and from 1 to 4 mM in *bas1Δ* cells (Gauthier et al., 2008) (similar results are shown in Fig.

1E), but induced negligible or no changes in *adk1Δ* cells (from 2 to 2 mM) (Gauthier et al.,

2008). Consistent with our hypothesis, the addition of adenine reduced Syn-GFP toxicity and

aggregation in *bas1Δ,* but not *adk1Δ* cells (Fig. 5D). Thus, a high level of ATP prevented

Syn-GFP aggregation and toxicity.

We examined another model protein involved in neurodegenerative diseases. PolyQ

containing the huntingtin protein is susceptible to aggregation and has been implicated in

Huntington's disease (Jiang, Poirier et al., 2005). We investigated the toxicity of Htt103Q, a

mutant form of the huntingtin protein that is also susceptible to aggregation and causes

cytotoxicity in yeast (Meriin, Zhang et al., 2002). Consistent with the concept that a high level

of ATP prevents protein aggregation, the ATP homeostasis mutants *snf1Δ, adk1Δ, snf1Δ*

adk1Δ, and *bas1Δ* were very sensitive to Htt103Q expression (Fig. S9).

Proteasomes are essential for the removal of protein aggregates induced by ATP depletion

Protein aggregation caused by ATP depletion was cytotoxic (Figs. 3B and 5A) and was not

easily dissolved after ATP repletion (Fig. 4). To identify a pathway that is essential for the

removal of aggregates, we examined the involvement of proteasomes and autophagy.

The deletion of *RPN4*, which encodes a transcription factor of proteasomal genes (Xie &

Varshavsky, 2001), down-regulated proteasomal activity (Kruegel, Robison et al., 2011) and

resulted in synthetic growth defects with *adk1Δ, snf1Δ, bas1Δ* at a high temperature of 38°C

and in the presence of H_2O_2 (Fig. 5E). In contrast to proteasomes, autophagy did not appear to

have genetic interactions with the above mutants (Fig. 5E). The deletion of an essential

component of the autophagic pathway, *ATG1* did not affect the sensitivity of *adk1Δ, snf1Δ,*

bas1Δ to a high temperature of 38°C or to H_2O_2 (Fig. 5E). We also did not observe the

accumulation of Hsp104-GFP foci in the autophagy mutants *atg1Δ, atg8Δ*, and *atg13Δ* (not

shown).

To investigate the involvement of proteasomes in the removal of protein aggregates after the

transient depletion of ATP, we pretreated cells with the proteasomal inhibitor MG132 or

DMSO and examined the kinetics for the formation of Hsp104-GFP foci after the 2DG

treatment (Fig. 5F) using the drug-sensitive yeast strain Y13206 (Piotrowski, Li et al., 2017).

Under both conditions, more than 90% of cells exhibited Hsp104-GFP foci within 30 min of

the 2DG treatment. More than two-thirds of Hsp104-GFP foci dissolved in the DMSO control,

while less than one-third dissolved in MG132-treated samples, indicating that proteasomes

are required for the dissolution process (Fig. 5F).

DISCUSSION

DISCUSSION

In the present study, we demonstrated for the first time that the Snf1 complex, budding yeast

AMPK, is required for the stable maintenance of cellular ATP levels (ATP homeostasis) in

collaboration with Adk1 (Fig. 6A). This function of the Snf1 complex in ATP homeostasis is

independent of glucose levels in the medium and Mig1, the major transcriptional repressor

involved in glucose repression (Fig. S1); therefore, this is distinct from its well-characterized

role in adaptation to glucose limitations. The activity of the Snf1 kinase complex may be

sharply tuned depending on the intracellular levels of adenine nucleotides or other metabolites

indicative of cellular energy to prevent a rapid ATP catastrophe (Fig. 2), even in the presence

of sufficient amounts of glucose. It is important to note that the reductions observed in

intracellular ATP levels in *snf1Δ* cells in the presence of glucose were overlooked in previous

biochemical analyses, again demonstrating the usefulness of QUEEN-based ATP imaging.

Since the deletion of *BAS1* induced the greatest reduction in ATP levels and is epistatic to

snf1Δ, a large pool size of adenine nucleotides is a prerequisite for ATP homeostasis. This

assumption is reasonable because the pool size of recyclable ATP restricts ATP levels based

on the rapid turnover rate of ATP. Bas1 maintains the pool size of ATP by balancing ATP

synthesis and irreversible decreases, such as incorporation into RNA and DNA (following

conversion to deoxy-ATP), degradation, and excretion in rapidly proliferating yeasts.

The decreases observed in ATP levels in the ATP mutant cells were confirmed by our

biochemical measurements (Figs. 1 and S3). The biochemical assay also revealed that ADP

levels decreased in ATP mutants, similar to ATP levels, and, as a consequence, ATP/ADP

ratios, indicators of cellular energy charges, remained largely unchanged. On the other hand,

the sum of ATP and ADP levels, indicators of the pool size of adenine nucleotides, decreased

in the mutants, which may explain the instability observed in ATP levels. While cytosolic

ATP levels were reduced and unstable in *snf1Δadk1Δ* cells, high levels of ATP may

accumulate in intracellular membrane compartments (*e.g.*, vacuoles, lysosomes, and

mitochondria) that were not visualized by the QUEEN, which would increase the average

ATP level of whole cells (Fig. 1B, C).

We also showed that key regulators of ATP homeostasis play roles in preventing cytotoxic

protein aggregation in budding yeast (Fig. 6B). The common feature associated with these

mutants is reduced ATP levels, suggesting that high ATP levels are essential for protein

solubilization.

A proteomic study suggested that the main role of ATP changes depending on its level. At levels lower than 0.5 mM, ATP mainly serves as a substrate for enzymes, such as protein kinases and heat shock protein chaperones, whereas at levels higher than 2 mM, ATP may exert solubilizing effects on disordered proteins (Sridharan et al., 2019). ATP homeostasis may also be required to constantly drive proteasomal protein degradation, which requires high levels of ATP (Benaroudj, Zwickl et al., 2003, Tanaka, Waxman et al., 1983).

Previous biochemical measurements indicated that although ATP levels were lower in *adk1* and *bas1* mutants than in the wild type, these mutants still had ATP levels that were higher than 2 mM (Gauthier et al., 2008). This does not directly explain the accumulation of protein aggregates in these mutants (Fig. 3) because most proteins are expected to be soluble at > 2 mM ATP (Sridharan et al., 2019). In this study, we explained this discrepancy by using the biosensor-based ATP imaging technique we developed (Takaine et al., 2019, Yaginuma et al., 2014). The visualization of ATP dynamics in living cells at the single cell level revealed that ATP repeatedly undergoes transient depletion (the ATP catastrophe) in AMPK and ADK mutants, which explains how aggregation-prone proteins aggregate and cause cytotoxicity. A similar, but distinct, instability in ATP was observed in *bas1Δ*, further confirming that the ATP catastrophe causes protein aggregation. Although severe ATP depletion in these mutants was gradually recovered by as yet uncharacterized negative feedback regulation, the duration period of ~15 min with reduced ATP levels may induce some proteins to form aggregates that last for generations.

A recent study revealed that forced energy (ATP) depletion from budding yeast cells induced extensive cytoplasmic reorganization, including increases in macromolecular (ribosome) crowding, the emergence of numerous membrane-less organelles, and the polymerization of eukaryotic translation initiation factor 2B (Marini et al., 2020). Another recent study showed that the depletion of ATP triggered the "viscoadaptation" of yeast cytoplasm by an as yet unknown mechanism, and induced increases in viscosity and decreases in biomolecular solubility, thereby enhancing phase separation (Persson et al., 2020). These findings imply that malfunctional ATP homeostasis may induce undesirable and dysregulated biomolecular assemblies driven by enhanced liquid-liquid phase separation and molecular crowding. The present study using ATP mutant cells is the first to demonstrate the consequence of failed ATP homeostasis under physiologically relevant conditions and highlights the biological significance of ATP homeostasis.

Many neurodegenerative diseases, such as Alzheimer's, Huntington's, and Parkinson's

diseases, are associated with protein aggregation (Eisele, Monteiro et al., 2015, Josefson et al.,

2017). On the other hand, based on a large body of evidence, mitochondrial dysfunction and

accompanying energy failure in nerve cells may result in many types of neurodegenerative

diseases (Haelterman, Yoon et al., 2014, Pathak et al., 2013). Previous studies demonstrated

that ATP levels in the brain were decreased in patients with early Parkinson's disease (Mochel,

N'Guyen et al., 2012) and also that ATP synthesis in the brain was not properly regulated in

patients with early Huntington's disease (Hattingen, Magerkurth et al., 2009) and in the

corresponding mouse model (Mochel, Durant et al., 2012). Therefore, protein aggregation

induced by the ATP catastrophe, as revealed in the present study, may link energy failure and

protein aggregation, providing a comprehensive insight into the onset of neurodegenerative

diseases. Further studies are warranted to clarify whether the ATP catastrophe also occurs in

the neurons of patients at risk of neurodegenerative diseases or in the elderly. However,

neither biochemical measurements nor mass spectrometry is capable of detecting the ATP

catastrophe because of their insufficient time and space resolution. Therefore, an ATP

imaging approach using the yeast system will be the leading model for elucidating the

molecular mechanisms underlying ATP homeostasis and ATP catastrophe-induced protein

aggregation.

A recent study reported that the activation of AMPK by metformin ameliorated the

progression of Huntington's disease in a mouse model (Arnoux, Willam et al., 2018), and the

potential therapeutic use of metformin for neurodegenerative diseases is being discussed

(Rotermund, Machetanz et al., 2018). Furthermore, the involvement of ATP and ADK in

preventing the manifestation of Parkinson's disease in mouse models and patients has been

proposed (Garcia-Esparcia, Hernandez-Ortega et al., 2015, Nakano, Imamura et al., 2017).

Protein aggregation induced by the ATP catastrophe may be a general mechanism for the

development of proteinopathies. The present study using ATP imaging revealed a

physiological consequence of a failure in ATP homeostasis in living cells for the first time

and suggests that ATP homeostasis has potential as a target for preventing/treating

neurodegenerative diseases.

MATERIALS AND METHODS

Yeast strains and plasmids

The budding yeast strains and plasmids used in the present study are listed in Supplementary Tables S1 and S2, respectively. These strains were constructed by a PCR-based method (Janke, Magiera et al., 2004) and genetic crosses. The yeast knockout strain collection was originally purchased from GE Healthcare (cat. # YSC1053).

Media and cell culture

The standard technique for the yeast culture and manipulation was used (Guthrie & Fink, 2002). Synthetic medium (SC) was prepared as described by Hanscho et al. (Hanscho, Ruckerbauer et al., 2012). 2-Deoxy-D-glucose (2DG), tunicamycin, and MG132 were purchased from FUJIFILM Wako (cat. # 046-06483, 202-08241, and 139-18451, respectively). Tunicamycin and MG132 were dissolved in dimethylsulfoxide (DMSO) to make stock solutions (5 mg/ml and 42 mM, respectively). Cells were grown to the mid-log phase at 30°C in SC before imaging unless otherwise noted.

Microscopy

Cells expressing Hsp104-GFP or GFP-Ubc9ts were concentrated by centrifugation and sandwiched between a slide and coverslip (No. 1.5 thickness, Matsunami, Osaka, Japan). Immobilized cells were imaged using an inverted fluorescent microscope (Eclipse Ti-E, Nikon) equipped with an Apo TIRF 100× Oil DIC N2/NA 1.49 objective lens and electron-multiplying charge-coupled device camera (iXon3 DU897E-CS0-#BV80, Andor) at approximately 25°C. The Hsp104-GFP and GFP-Ubc9ts fluorescent signal was collected from stacks of 11 z-sections spaced by 0.5 μm, and the maximum projections of the optical sections were shown in Figs. 4, 5, and S6. Cells expressing QUEEN were immobilized on a concanavalin A-coated 35-mm glass-bottomed dish (#3971-035, No. 1.5 thickness, IWAKI). The dish was filled with an excess amount of medium (4.5–5 ml) against the cell volume to minimize changes in the chemical compositions of the medium during observations. QUEEN fluorescence was acquired as previously described (Takaine et al., 2019). Cells expressing Syn-GFP were immobilized on a slide glass as described above, and the fluorescence signal was collected from a single z-plane using an inverted fluorescent microscope (Eclipse Ti2-E, Nikon, Tokyo, Japan) equipped with a CFI Plan Apoλ100× Oil DIC/NA1.45 objective lens

and CMOS image sensor (DS-Qi2, Nikon). Images of cells were acquired from several fields

of view for each experimental condition.

Biochemical measurements of ATP and ADP

Whole cell extracts were prepared according to Seo et al. (Seo, Lau et al., 2017) with slight

modifications. Mid-log cells were harvested in a 1.6-ml microtube by centrifugation and

resuspended in 1 ml fresh SC or 40 mM 2DG medium (for ATP depletion). After a 10-min

incubation at 30°C, a small fraction of cells (50 µl) was removed for the measurement of cell

numbers and optical density, and the remaining cells were pelleted and resuspended in 0.75

ml of 90% acetone. The suspension was incubated at 90°C for 15 min to evaporate acetone.

The remaining solution (30–35 µl) was centrifuged at 20,000×g at 4°C for 3 min. The

supernatant was mixed with 450 µl of TE (10 mM Tris-HCl, pH 8.0, and 1 mM

ethylenediaminetetraacetic acid). These extracts were stored at -80°C until analyzed. ATP and

ADP levels were measured using the ATP determination kit (Invitrogen) and EnzyLight ADP

assay kit (EADP-100, Funakoshi), respectively, according to the manufacturers' instructions.

Luminescence was measured using an Enspire multimode plate reader (PerkinElmer). All

samples were assayed in duplicate. ATP and ADP levels were normalized for an optical

density at 600 nm of the initial cell suspension assessed by BioSpectrometer (Eppendorf).

Data analysis

Numerical data were plotted using KaleidaGraph software ver. 4.5.1 (Synergy Software) and

R studio software ver. 3.4.1 (R Core Team, 2017). Means, SDs, and p values were calculated

using Excel software (Microsoft, WA, US). Significance between two sets of data was tested

using the unpaired one-tailed Welch's t-test unless otherwise noted, and was indicated by an

asterisk or p value. Data were sometimes represented by a dot plot that shows distribution

characteristics in extensive detail. The horizontal bar in the dot plot indicates the average of

each population. All measurements were repeated at least twice to confirm reproducibility.

ATP levels in yeast cells were estimated using QUEEN-based ratiometric imaging,

as previously described (Takaine, 2019), (Takaine et al., 2019). The QUEEN ratio is

proportional to ATP levels and pseudo-colored to reflect its value throughout the present

study. The mean QUEEN ratio inside of a cell represents the intracellular ATP level of the

cell.

The autocorrelation functions (ACF) of oscillations in the QUEEN ratio were

calculated using R studio software. The apparent period of oscillation was estimated from the

positive second peak of the correlation coefficient, which was outside the 95% confidence

interval and significant ($p < 0.05$), rejecting the assumption that there is no correlation.

ACKNOWLEDGMENTS

We are grateful to the Yeast Genetic Resource Center, Y. Ohya, K. Ohashi, H. Takagi, D.

Watanabe, and J. Frydman for providing the yeast strains and plasmids. We thank the

members of the Yoshida/Takaine laboratories for their support. This work was supported by

JSPS grants 16H04781 (S.Y. and M.T.), 15K18525 (M.T.), and 19K06654 (M.T.) and the

Takeda Science Foundation (S.Y.). This work was also supported by the joint research

program of the Institute for Molecular and Cellular Regulation, Gunma University, Japan.

AUTHOR CONTRIBUTIONS

M.T. and S.Y. conceived and designed the project. M.T. conducted experiments and the data

analysis. H.I. provided a key reagent and expertise. M.T. and S.Y wrote the manuscript with

input from H.I.

CONFLICT OF INTEREST

The authors declare no competing interests.

REFERENCES

Arnoux I, Willam M, Griesche N, Krummeich J, Watari H, Offermann N, Weber S, Narayan Dey P, Chen C,

Monteiro O, Buettner S, Meyer K, Bano D, Radyushkin K, Langston R, Lambert JJ, Wanker E, Methner A,

Krauss S, Schweiger S et al. (2018) Metformin reverses early cortical network dysfunction and behavior changes

in Huntington's disease. *Elife* 7

Baryshnikova A (2016) Systematic Functional Annotation and Visualization of Biological Networks. *Cell*

Syst 2: 412-21

Benaroudj N, Zwickl P, Seemuller E, Baumeister W, Goldberg AL (2003) ATP hydrolysis by the proteasome

regulatory complex PAN serves multiple functions in protein degradation. *Mol Cell* 11: 69-78

Bhat AH, Dar KB, Anees S, Zargar MA, Masood A, Sofi MA, Ganie SA (2015) Oxidative stress,

mitochondrial dysfunction and neurodegenerative diseases; a mechanistic insight. *Biomed Pharmacother* 74:

101-10

Carlson M (1999) Glucose repression in yeast. *Curr Opin Microbiol* 2: 202-7

Cherry JM, Hong EL, Amundsen C, Balakrishnan R, Binkley G, Chan ET, Christie KR, Costanzo MC,

Dwight SS, Engel SR, Fisk DG, Hirschman JE, Hitz BC, Karra K, Krieger CJ, Miyasato SR, Nash RS, Park J,

Skrzypek MS, Simison M et al. (2012) Saccharomyces Genome Database: the genomics resource of budding

yeast. *Nucleic Acids Res* 40: D700-5

Daignan-Fornier B, Fink GR (1992) Coregulation of purine and histidine biosynthesis by the transcriptional

activators BAS1 and BAS2. *Proc Natl Acad Sci U S A* 89: 6746-50

Denis V, Boucherie H, Monribot C, Daignan-Fornier B (1998) Role of the myb-like protein bas1p in

Saccharomyces cerevisiae: a proteome analysis. *Mol Microbiol* 30: 557-66

Edelman AM, Blumenthal DK, Krebs EG (1987) Protein serine/threonine kinases. *Annu Rev Biochem* 56:

567-613

Eftekharzadeh B, Hyman BT, Wegmann S (2016) Structural studies on the mechanism of protein aggregation

in age related neurodegenerative diseases. *Mech Ageing Dev* 156: 1-13

Eisele YS, Monteiro C, Fearns C, Encalada SE, Wiseman RL, Powers ET, Kelly JW (2015) Targeting

protein aggregation for the treatment of degenerative diseases. *Nat Rev Drug Discov* 14: 759-80

Garcia-Esparcia P, Hernandez-Ortega K, Ansoleaga B, Carmona M, Ferrer I (2015) Purine metabolism gene

deregulation in Parkinson's disease. *Neuropathol Appl Neurobiol* 41: 926-40

Gauthier S, Coulpier F, Jourdren L, Merle M, Beck S, Konrad M, Daignan-Fornier B, Pinson B (2008)

Co-regulation of yeast purine and phosphate pathways in response to adenylic nucleotide variations. *Mol*

Microbiol 68: 1583-94

Ghillebert R, Swinnen E, Wen J, Vandesteene L, Ramon M, Norga K, Rolland F, Winderickx J (2011) The

AMPK/SNF1/SnRK1 fuel gauge and energy regulator: structure, function and regulation. *FEBS J* 278: 3978-90

Guthrie C, Fink GR (2002) *Guide to yeast genetics and molecular and cell Biology: Part C*. Gulf Professional

Publishing,

Haelterman NA, Yoon WH, Sandoval H, Jaiswal M, Shulman JM, Bellen HJ (2014) A Mitocentric View of

Parkinson's Disease. *Annual Review of Neuroscience* 37: 137-159

Hanscho M, Ruckerbauer DE, Chauhan N, Hofbauer HF, Krahulec S, Nidetzky B, Kohlwein SD, Zanghellini

J, Natter K (2012) Nutritional requirements of the BY series of Saccharomyces cerevisiae strains for optimum

growth. *FEMS Yeast Res* 12: 796-808

Hardie DG, Carling D, Carlson M (1998) The AMP-activated/SNF1 protein kinase subfamily: metabolic

sensors of the eukaryotic cell? *Annu Rev Biochem* 67: 821-55

Hardie DG, Schaffer BE, Brunet A (2016) AMPK: An Energy-Sensing Pathway with Multiple Inputs and

Outputs. *Trends Cell Biol* 26: 190-201

Hattingen E, Magerkurth J, Pilatus U, Mozer A, Seifried C, Steinmetz H, Zanella F, Hilker R (2009)

Phosphorus and proton magnetic resonance spectroscopy demonstrates mitochondrial dysfunction in early and

advanced Parkinson's disease. *Brain* 132: 3285-97

Hayes MH, Peuchen EH, Dovichi NJ, Weeks DL (2018) Dual roles for ATP in the regulation of phase

separated protein aggregates in Xenopus oocyte nucleoli. *Elife* 7: e35224

Hedbacker K, Carlson M (2008) SNF1/AMPK pathways in yeast. *Front Biosci* 13: 2408-20

Herzig S, Shaw RJ (2017) AMPK: guardian of metabolism and mitochondrial homeostasis. *Nature Reviews*

Molecular Cell Biology 19: 121

Hoyle NP, Castelli LM, Campbell SG, Holmes LE, Ashe MP (2007) Stress-dependent relocalization of

translationally primed mRNPs to cytoplasmic granules that are kinetically and spatially distinct from P-bodies. *J*

Cell Biol 179: 65-74

Jain S, Wheeler JR, Walters RW, Agrawal A, Barsic A, Parker R (2016) ATPase-Modulated Stress Granules

Contain a Diverse Proteome and Substructure. *Cell* 164: 487-98

Janke C, Magiera MM, Rathfelder N, Taxis C, Reber S, Maekawa H, Moreno-Borchart A, Doenges G,

Schwob E, Schiebel E, Knop M (2004) A versatile toolbox for PCR-based tagging of yeast genes: new

fluorescent proteins, more markers and promoter substitution cassettes. *Yeast* 21: 947-62

Janssen E, Dzeja PP, Oerlemans F, Simonetti AW, Heerschap A, de Haan A, Rush PS, Terjung RR, Wieringa

B, Terzic A (2000) Adenylate kinase 1 gene deletion disrupts muscle energetic economy despite metabolic

rearrangement. *EMBO J* 19: 6371-81

Jiang H, Poirier MA, Ross CA (2005) A structure-based analysis of huntingtin mutant polyglutamine

aggregation and toxicity: evidence for a compact beta-sheet structure. *Human Molecular Genetics* 14: 765-774

Josefson R, Andersson R, Nyström T (2017) How and why do toxic conformers of aberrant proteins

accumulate during ageing? *Essays In Biochemistry* 61: 317

Kaganovich D, Kopito R, Frydman J (2008) Misfolded proteins partition between two distinct quality control

compartments. *Nature* 454: 1088-95

Kruegel U, Robison B, Dange T, Kahlert G, Delaney JR, Kotireddy S, Tsuchiya M, Tsuchiyama S,

Murakami CJ, Schleit J, Sutphin G, Carr D, Tar K, Dittmar G, Kaeberlein M, Kennedy BK, Schmidt M (2011)

Elevated proteasome capacity extends replicative lifespan in Saccharomyces cerevisiae. *PLoS Genet* 7:

e1002253

Lashuel HA, Overk CR, Oueslati A, Masliah E (2013) The many faces of alpha-synuclein: from structure and

toxicity to therapeutic target. *Nat Rev Neurosci* 14: 38-48

Ljungdahl PO, Daignan-Fornier B (2012) Regulation of amino acid, nucleotide, and phosphate metabolism in

Saccharomyces cerevisiae. *Genetics* 190: 885-929

Marini G, Nuske E, Leng W, Alberti S, Pigino G (2020) Reorganization of budding yeast cytoplasm upon

energy depletion. *Mol Biol Cell* 31: 1232-1245

Meriin AB, Zhang X, He X, Newnam GP, Chernoff YO, Sherman MY (2002) Huntington toxicity in yeast

model depends on polyglutamine aggregation mediated by a prion-like protein Rnq1. *J Cell Biol* 157: 997-1004

Mochel F, Durant B, Meng X, O'Callaghan J, Yu H, Brouillet E, Wheeler VC, Humbert S, Schiffmann R,

Durr A (2012) Early alterations of brain cellular energy homeostasis in Huntington disease models. *J Biol Chem*

287: 1361-70

Mochel F, N'Guyen TM, Deelchand D, Rinaldi D, Valabregue R, Wary C, Carlier PG, Durr A, Henry PG

(2012) Abnormal response to cortical activation in early stages of Huntington disease. *Mov Disord* 27: 907-10

Nakano M, Imamura H, Sasaoka N, Yamamoto M, Uemura N, Shudo T, Fuchigami T, Takahashi R,

Kakizuka A (2017) ATP Maintenance via Two Types of ATP Regulators Mitigates Pathological Phenotypes in

Mouse Models of Parkinson's Disease. *EBioMedicine* 22: 225-241

Outeiro TF, Lindquist S (2003) Yeast cells provide insight into alpha-synuclein biology and pathobiology.

Science 302: 1772-5

Patel A, Malinovska L, Saha S, Wang J, Alberti S, Krishnan Y, Hyman AA (2017) ATP as a biological

hydrotrope. *Science* 356: 753-756

Pathak D, Berthet A, Nakamura K (2013) Energy failure: does it contribute to neurodegeneration? *Ann*

Neurol 74: 506-16

Persson LB, Ambati VS, Brandman O (2020) Cellular Control of Viscosity Counters Changes in

Temperature and Energy Availability. *Cell*

Piotrowski JS, Li SC, Deshpande R, Simpkins SW, Nelson J, Yashiroda Y, Barber JM, Safizadeh H, Wilson

E, Okada H, Gebre AA, Kubo K, Torres NP, LeBlanc MA, Andrusiak K, Okamoto R, Yoshimura M,

DeRango-Adem E, van Leeuwen J, Shirahige K et al. (2017) Functional annotation of chemical libraries across

diverse biological processes. *Nature Chemical Biology* 13: 982

Pu Y, Li Y, Jin X, Tian T, Ma Q, Zhao Z, Lin SY, Chen Z, Li B, Yao G, Leake MC, Lo CJ, Bai F (2019)

ATP-Dependent Dynamic Protein Aggregation Regulates Bacterial Dormancy Depth Critical for Antibiotic

Tolerance. *Mol Cell* 73: 143-156 e4

R Core Team (2017) R: A Language and Environment for Statistical Computing. In Vienna, Austria: R

Foundation for Statistical Computing

Rotermund C, Machetanz G, Fitzgerald JC (2018) The Therapeutic Potential of Metformin in

Neurodegenerative Diseases. *Front Endocrinol (Lausanne)* 9: 400

Seo AY, Lau P-W, Feliciano D, Sengupta P, Gros MAL, Cinquin B, Larabell CA, Lippincott-Schwartz J

(2017) AMPK and vacuole-associated Atg14p orchestrate μ-lipophagy for energy production and long-term

survival under glucose starvation. *eLife* 6: e21690

Sharma N, Brandis KA, Herrera SK, Johnson BE, Vaidya T, Shrestha R, DebBurman SK (2006) α-synuclein

budding yeast model. *Journal of Molecular Neuroscience* 28: 161-178

Sridharan S, Kurzawa N, Werner T, Gunthner I, Helm D, Huber W, Bantscheff M, Savitski MM (2019)

Proteome-wide solubility and thermal stability profiling reveals distinct regulatory roles for ATP. *Nat Commun*

10: 1155

Takaine M (2019) QUEEN-based Spatiotemporal ATP Imaging in Budding and Fission Yeast. *Bio-Protocol*

9: e3320

Takaine M, Ueno M, Kitamura K, Imamura H, Yoshida S (2019) Reliable imaging of ATP in living budding

and fission yeast. *J Cell Sci* 132

Tanaka K, Waxman L, Goldberg AL (1983) ATP serves two distinct roles in protein degradation in

reticulocytes, one requiring and one independent of ubiquitin. *J Cell Biol* 96: 1580-5

Tofaris GK, Kim HT, Hourez R, Jung JW, Kim KP, Goldberg AL (2011) Ubiquitin ligase Nedd4 promotes

alpha-synuclein degradation by the endosomal-lysosomal pathway. *Proc Natl Acad Sci U S A* 108: 17004-9

Usaj M, Tan Y, Wang W, VanderSluis B, Zou A, Myers CL, Costanzo M, Andrews B, Boone C (2017)

TheCellMap.org: A Web-Accessible Database for Visualizing and Mining the Global Yeast Genetic Interaction

Network. *G3 (Bethesda)* 7: 1539-1549

Wijayanti I, Watanabe D, Oshiro S, Takagi H (2015) Isolation and functional analysis of yeast ubiquitin

ligase Rsp5 variants that alleviate the toxicity of human α-synuclein. *The Journal of Biochemistry* 157: 251-260

Willingham S, Outeiro TF, DeVit MJ, Lindquist SL, Muchowski PJ (2003) Yeast genes that enhance the

toxicity of a mutant huntingtin fragment or alpha-synuclein. *Science* 302: 1769-72

Wilson WA, Hawley SA, Hardie DG (1996) Glucose repression/derepression in budding yeast: SNF1 protein

kinase is activated by phosphorylation under derepressing conditions, and this correlates with a high AMP:ATP

ratio. *Curr Biol* 6: 1426-34

Xiao B, Heath R, Saiu P, Leiper FC, Leone P, Jing C, Walker PA, Haire L, Eccleston JF, Davis CT, Martin

SR, Carling D, Gamblin SJ (2007) Structural basis for AMP binding to mammalian AMP-activated protein

kinase. *Nature* 449: 496-500

Xie Y, Varshavsky A (2001) RPN4 is a ligand, substrate, and transcriptional regulator of the 26S

proteasome: a negative feedback circuit. *Proc Natl Acad Sci U S A* 98: 3056-61

Yaginuma H, Kawai S, Tabata KV, Tomiyama K, Kakizuka A, Komatsuzaki T, Noji H, Imamura H (2014)

Diversity in ATP concentrations in a single bacterial cell population revealed by quantitative single-cell imaging.

Sci Rep 4: 6522

 FIGURE LEGENDS

Fig. 1 Interconversion and active synthesis of adenine nucleotides are important for

ATP homeostasis

(A) Adk1 and Snf1 synergistically control cellular ATP levels. QUEEN ratio images of ATP

homeostasis mutant cells grown in medium containing 2% glucose. The asterisk indicates an

example of cells with significantly reduced ATP levels. (B) The mean QUEEN ratios of cells

were translated to ATP levels and shown in a dot plot. The horizontal bar indicates the mean

of each population. Asterisks indicate p values versus WT: *=1.5×10^{-59}, **=2.3×10^{-34},

***=2.9×10^{-117}. CV: coefficient of variance. N = 134–276 cells were scored.

(C) Biochemical measurements of cellular ATP levels. ATP levels in cells of the indicated

genotypes were measured as described in the Materials and Methods. Data are the mean ±

1SD (error bars) (N = 4). An asterisk indicates a p value of 0.03 versus WT. (D) QUEEN ratio

images of *bas1Δ* cells grown in 2% glucose medium. Growth in media supplemented with

0.11 mg/ml adenine partially restored the low ATP phenotype of *bas1Δ*. (E) ATP levels in

cells shown in *D* were plotted. Asterisks indicate p values: *=2.3×10^{-160}, **=8.8×10^{-12} (versus

WT in glucose), ***=3.6×10^{-20} (versus *bas1Δ* in glucose). CV: coefficient of variance. N =

186–296 cells were scored. (F) ATP levels in WT and *bas1Δ* cells were measured as

described in *C*. Data are the mean ± 1SD (error bars) (N = 4). An asterisk indicates a p value

of 8.5×10^{-5} versus WT.

Fig. 2 Temporal fluctuations in ATP levels in *snf1Δadk1Δ* and *bas1Δ* cells

(A, C) Time-lapse imaging of QUEEN in *snf1Δ adk1Δ* cells in 2% glucose medium. Images

at the representative time points were shown. The QUEEN ratio decreased twice (indicated by

arrowheads) within a short interval. See also Movie S1. Data were converted into ATP levels

and plotted in (C). (B, D) Another example of the time-lapse imaging of QUEEN in

snf1Δadk1Δ cells in 2% glucose medium. The QUEEN ratio decreased twice (indicated by

arrowheads) with a long interval. See also Movie S2. Data were converted into ATP levels

and plotted in (D). (E) Time-lapse imaging of QUEEN in *bas1Δ* in 2% glucose medium. ATP

levels in the mother (cell-1) and daughter (cell-2) were plotted at the bottom. Images at the

representative time points were shown on the top. Note that the QUEEN ratio is synchronized

until cells undergo separation at the time point of 76 min indicated by an arrow. After

separation, each cell has a unique periodic cycle of ATP. The movie is available in Movie S3.

White scale bar = 5 μm.

Fig. 3 ATP homeostasis is required to prevent protein aggregation

(A) Functional landscape of known interactors of ATP mutants. Negative genetic interactors of the indicated gene were derived from the SGD database (https://www.yeastgenome.org/) (Cherry, Hong et al., 2012) and overlaid on a functional map based on the global genetic interaction network of the yeast genome (Baryshnikova, 2016, Usaj et al., 2017). *URA6* encodes an uridylate kinase that is essential for viability, which also exhibits adenylate kinase activity. (B) Each strain of the indicated genotype was serially diluted (five-fold), spotted on SC + 2% glucose medium, and grown under the indicated stress conditions. Photos were taken after 2-3 days. (C) Formation of Hsp104-GFP foci in ATP homeostasis mutants. The GFP signal (inverted grayscale) was imaged in the log phase culture of the indicated mutant cells expressing Hsp104-GFP at 35°C. (D) Quantification of data shown in (C). Data from similar experiments using strains expressing Pab1-GFP, instead of Hsp104-GFP, were also plotted. Values are the mean ± 1SD (error bars). Asterisks indicate a significant difference from WT ($p < 0.05$) (N=3–4). White scale bar = 5 μm.

Fig. 4 ATP depletion triggers protein aggregation in living yeast cells

(A) The formation of Hsp104-GFP foci after ATP depletion. Wild-type cells expressing Hsp104-GFP were grown to the log phase at 35°C in medium containing 2% glucose. At the time point of 0 min, medium was replaced with 20 mM 2DG (red) or 2% glucose (as a control; blue). Cells were released back to media containing 2% glucose at the time point of 50 min. Cells were imaged at the indicated time points, classified, and scored according to the number of Hsp104-GFP foci. Values are the mean (N = 3). Asterisks indicate a significant difference from the 2% glucose treatment ($p < 0.05$). (B) Representative images of cells analyzed in (A). (C) Formation of Ubc9-ts foci after ATP depletion. Cells expressing GFP-Ubc9-ts under an inducible *GAL* promoter were grown in medium containing 2% galactose (SC-gal) at 33°C, and medium was then exchanged with 2DG or SC-gal. After 30 min, cells were imaged and scored for the number of GFP-Ubc9-ts foci. Representative images (inverted grayscale) are shown on the left and summarized on the right. Values are the mean ± 1SD (error bars) (N = 4). White scale bar = 5 μm.

Fig. 5 Aggregation and cytotoxicity of α-synuclein depends on ATP homeostasis

(A) Each strain of the indicated genotype was transformed with an expression vector carrying α-synuclein-GFP or GFP, serially diluted (five-fold), spotted on SC + 2% galactose plates, and then grown at 30°C for 3 days. (B) The localization of α-synuclein-GFP in ATP mutants. Cells were grown on galactose plates at 30°C for more than 42 h and then imaged.

Representative images of α-synuclein-GFP (inverted grayscale) are shown. (C) Quantification

of the data shown in (B). Cells were classified and scored for the localization pattern of

α-synuclein-GFP. The percentage of cells showing α-synuclein-GFP foci are plotted. Data are

the mean $\pm$ 1SD (error bars) from 3–6 independent observations. N = 33–380 cells were

scored in each measurement. *P* values versus WT are shown. N.S., no significance (*p* value >

0.05). (D) *(top)* Each strain of the indicated genotype was transformed with an expression

vector carrying α-synuclein-GFP and grown on galactose plates containing 0 mM (−

Adenine) or 0.3 mM (+ Adenine) adenine at 30°C for 3 days. *(bottom)* Cells were grown on

galactose plates in the absence or presence of adenine at 30°C for 41-45 h and then imaged.

The percentage of cells showing α-synuclein-GFP foci was plotted. Data are the mean $\pm$ 1SD

(error bars) from 5 independent transformants. N = 53–258 cells were scored in each

measurement. *P* values versus "− Adenine" are shown. (E) Each strain of the indicated

genotype was serially diluted (five-fold), spotted on SC + 2% glucose medium, and grown

under the indicated stress conditions. (F) Cells of the drug-sensitive strain Y13206 were

grown to the log phase at 37°C in medium containing 2% glucose and supplemented with

0.1% DMSO or 0.1% DMSO plus 42 μM MG132 at $t = -30$ min. At $t = 0$ min, these cells

were washed and released in medium containing 20 mM 2DG $\pm$ MG132, and cells were then

washed and released again in medium containing 2% glucose $\pm$ MG132. Cells were imaged at

the indicated time points and scored for the number of Hsp104-GFP foci. Data are the mean $\pm$

1SD (error bars). Asterisks indicate a significant difference from DMSO ($p < 0.02$) (N=3).

Fig. 6 Models for ATP homeostasis and its role in proteostasis

(A) Schematic summary of the roles of Snf1, Adk1, and Bas1 in ATP homeostasis. (B) A

schematic model for ATP homeostasis preventing cytotoxic protein aggregation.

 SUPPLEMENTARY INFORMATION

Supplementary figure legends

Fig. S1 AMPK is involved in the maintenance of cellular ATP levels in non-starving

cells

(A) Visualization of intracellular ATP levels in wild-type (WT) and *snf1Δ* cells using the

ATP sensor QUEEN. Cells were grown to the mid-log phase in SC-H medium containing the

indicated levels of glucose and then imaged. White scale bar = 5 μm. (B) *snf1Δ* cells had

lower ATP levels than wild-type cells. The mean QUEEN ratio inside a single cell (mean

QUEEN ratio/cell), which represents the intracellular ATP level of the cell, was quantified for

each cell from the ratio image. Data are the mean of the cell population ± 1SD (error bar)

normalized to wild-type cells. N = 105–193 cells were scored. Asterisks indicate a significant

difference from WT ($p < 0.05$). (C) *mig1Δ* cells had slightly lower ATP levels than wild-type

cells. Data are the mean of the cell population ± 1SD (error bar) normalized to wild-type cells.

N = 190–231 cells were scored. Asterisks indicate a significant difference between the two

strains.

Fig. S2 Adenylate kinase Adk1 is involved in the maintenance of cellular ATP levels

(A) QUEEN ratio images of wild-type and *adk1Δ* cells grown in 2% glucose or 2% galactose.

White scale bar = 5 μm. (B) Quantification of data shown in (A). The mean QUEEN ratio/cell

was quantified for each cell from ratio images. Values are the mean of the cell population ±

1SD (error bar). N = 182–236 cells were scored. Asterisks indicate a significant difference

from WT ($p < 0.05$).

Fig. S3 Biochemical measurements of ATP and ADP

(A) Biochemical measurements of cellular ADP levels. ADP levels in cells of the indicated

genotypes were measured as described in the Materials and Methods. Data are the mean ±

1SD (error bars) (N = 4). Asterisks indicate *p* values versus WT: *=6.0×10^{-4}, **=1.5×10^{-4}.

(B) 2DG depletes cellular ATP. ATP levels in WT cells incubated with medium containing

2% glucose or 40 mM 2DG for 10 min were biochemically measured. Relative ATP levels

were expressed as the ratios of ATP levels in glucose-treated cells. Data are the mean ± 1SD

(error bars) (N = 4). (C) ATP/ADP ratios in WT and ATP mutant cells. ATP/ADP ratios were

calculated from ATP and ADP levels measured using biochemical assays. Data are the mean

± 1SD (error bars) (N = 4). An asterisk indicates a *p* value of 6.0×10^{-4} versus WT. (D) The

sums of ATP and ADP levels in WT and ATP mutant cells were calculated from ATP and

ADP levels measured using biochemical assays. Data are the mean ± 1SD (error bars) (N = 4).

Asterisks indicate *p* values versus WT: *=1.6×10^{-2}, **=1.0×10^{-4}.

Fig. S4 ATP levels in *bas1Δ snf1Δ* cells

(A) QUEEN ratio images of wild-type, *bas1Δ, snf1Δ*, and *bas1Δ snf1Δ* cells. (B)

Quantification of data shown in (A). The mean QUEEN ratio/cell was quantified for each cell

from ratio images. Data are shown as a dot plot. The horizontal bar in the plot indicates the

mean of each population. Significance between two sets of data was tested using the unpaired

two-tailed Welch's *t*-test and indicated by *p* values. N.S., no significance (*p*-value > 0.05).

Fig. S5 Time-lapse imaging of QUEEN in *snf1Δ adk1Δ* cells

Time-lapse imaging of QUEEN in *snf1Δ adk1Δ* cells in 2% glucose medium. An example of

a *snf1Δ adk1Δ* cell showing an irreversible decrease in the QUEEN ratio (indicated by an

arrowhead). In this case, the cell eventually died (indicated by an arrow). Images at the

representative time points were shown. The ATP level was plotted at the bottom. White scale

bar = 5 μm.

Fig. S6. Oscillatory behavior of the ATP level visualized in *bas1Δ* cells

(A) Another example of *bas1Δ* cells showing an oscillating QUEEN ratio. In this case,

cytokinesis had just been completed at t = 0 min. ATP levels in cell-1 and cell-2 were plotted

at the bottom. See also Movie S4. White scale bar = 5 μm. (B) Autocorrelation function of the

QUEEN ratio calculated from the data on cell-1 in Fig. 3E. Blue dotted lines indicate the 95%

confidence interval. An arrow indicates the second peak of the correlation and corresponds to

the apparent period.

Fig. S7 *In silico* analysis of interactors of ATP mutants

Genetic and physical interactors of the indicated genes were derived from the SGD database

(https://www.yeastgenome.org/) (Cherry et al., 2012) and overlaid on a functional map based

on the global genetic interaction network of the yeast genome (Baryshnikova, 2016, Usaj et

al., 2017). *URA6* encodes an uridylate kinase essential for viability, which also exhibits

adenylate kinase activity.

Fig. S8 Simultaneous observation of Hsp104 and Pab1 foci

Wild-type cells expressing Hsp104-GFP and Pab1-RedStar2 were grown to the log phase at

37°C in medium containing 2% glucose. Cells were washed and released either in medium

containing 20 mM 2DG (top) or in medium lacking glucose (bottom), and then imaged after

30 and 60 min. White scale bar = 5 μm.

Fig. S9 Cytotoxicity of polyQ containing the huntingtin protein in wild-type and

ATP-mutant yeast cells

Each strain of the indicated genotype was transformed with an expression vector carrying

Htt103Q and grown on glucose ((-), no induction) or raffinose ((+), leaky expression) plates

at 30°C for 3 days.

Table S1. *Strains used in the present study*

Name	Genotype	Source	Figure
MTY3008	*snf1Δ::kanMX6 leu2Δ0 lys2Δ0 ura3Δ0*	Lab stock	1C, 3B, 5A-C, 5E, S9
MTY3015	*his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	Lab stock	1C, 1F, 3B, 5A-E, S9
MTY3049	*adk1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	Lab stock	1C, 3B, 5A-E, S9
MTY3118	*bas1Δ::kanMX6 leu2Δ0 lys2Δ0 ura3Δ0*	Lab stock	1F, 3B, 5A-E, S9
MTY3143	*his3Δ1:: 2×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0*	This study	S2
MTY3149	*adk1Δ::kanMX6 his3Δ1:: 2×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0*	This study	S2
MTY3264	*his3Δ1:: 3×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0 MYO1-3mCherry-hphMX6*	Takaine et al., 2019	1A-B, S1, S4
MTY3270	*bas1Δ::kanMX6 his3Δ1:: 3×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0 MYO1-3mCherry-hphMX6*	This study	1D, 1E, 2E, S4, S6
MTY3293	*adk1Δ::kanMX6 his3Δ1:: 3×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0 MYO1-3mCherry-hphMX6*	This study	1A-B
MTY3355	*adk1Δ::kanMX6 snf1Δ::natNT2 his3Δ1:: 3×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0 MYO1-3mCherry-hphMX6*	This study	1A-B, 2A-D, S5
MTY3371	*snf1Δ::kanMX6 his3Δ1:: 3×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0 MYO1-3mCherry-hphMX6*	This study	1A-B, S1, S4
MTY3412	*adk1Δ::kanMX6 snf1Δ::natNT2 leu2Δ0 lys2Δ0 ura3Δ0*	This study	1C, 3B, 5A-E, S9
MTY3417	*mig1Δ::natNT2 his3Δ1:: 3×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0 MYO1-3mCherry-hphMX6*	This study	S1C
MTY3420	*Hsp104-yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	3C, D, 4A, B
MTY3421	*snf1Δ::kanMX6 Hsp104-yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	3C, D
MTY3422	*bas1Δ::kanMX6 Hsp104-yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	3C, D

MTY3424	*adk1Δ::kanMX6 Hsp104-yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	3C-D
MTY3425	*adk1Δ::kanMX6 snf1Δ::natNT2 Hsp104-yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	3C, D
MTY3489	*atg1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	Lab stock	5E
MTY3493	*snq2Δ::KlLeu2; pdr3Δ::Klura3; pdr1Δ::natMX4; can1Δ::STE2prSp_his5 lyp1Δ his3Δ1 leu2Δ0 ura3Δ0 met15Δ0 LYS2+*	From Y. Ohya, (Piotrowski et al., 2017)	Parental strain of MTY3501
MTY3501	*Hsp104-yeGFP-hphNT1 snq2Δ::KlLeu2; pdr3Δ::Klura3; pdr1Δ::natMX4; can1Δ::STE2prSp_his5 lyp1Δ his3Δ1 leu2Δ0 ura3Δ0 met15Δ0 LYS2+*	This study	5F
MTY3503	*Pab1- yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	3D
MTY3504	*Hsp104-yeGFP-hphNT1 Pab1-RedStar2-natNT2 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	S8
MTY3505	*snf1Δ::kanMX6 Pab1- yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	3D
MTY3506	*adk1Δ::kanMX6 Pab1- yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	3D
MTY3507	*adk1Δ::kanMX6 snf1Δ::natNT2 Pab1- yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	3D
MTY3508	*bas1Δ::kanMX6 Pab1- yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	3D
MTY3513	*atg1Δ::natNT2 adk1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	5E
MTY3514	*atg1Δ::natNT2 snf1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	5E
MTY3515	*atg1Δ::natNT2 bas1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	5E
MTY3516	*rpn4Δ::natNT2 adk1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	5E
MTY3517	*rpn4Δ::natNT2 snf1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	5E
MTY3518	*rpn4Δ::natNT2 bas1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	5E
MTY3525	*rpn4Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	Lab stock	5A-C, 5E

816   Table S2. *Plasmids used in the present study*

Name	Structure	Source	Purpose
MTP3091	pESC-Leu-GFP-Ubc9-ts	Judith Frydman (pJF1089)	Fig. 4C
MTP3088	p426 103Q GAL	Addgene (cat.# 1188)	Fig. S9
MTP3108	pYES2-α-synuclein-GFP	H. Takagi (Wijayanti et al., 2015)	Fig. 5A-D
MTP3090	pYES2-GFP	K. Ohashi	Fig. 5A

Captions for supplementary movies

Movie S1

Time-lapse imaging of QUEEN in *snf1Δadk1Δ* cells in 2% glucose medium. Corresponding

to the data shown in Fig. 2A. The QUEEN ratio decreased twice (116 and 132 min) within a

short interval. White scale bar = 5 µm.

Movie S2

Another example of *snf1Δadk1Δ* cells showing a sudden decrease in the QUEEN ratio.

Corresponding to the data shown in Fig. 2B. The QUEEN ratio decreased twice (180 and 356

min) with a long interval. White scale bar = 5 µm.

Movie S3

Oscillatory behavior of the QUEEN ratio in *bas1Δ* cells. Corresponding to the data shown in

Fig. 2E. Left, the QUEEN ratio image; middle, Myo1-mCherry (inverted grayscale image);

right, bright field image. Images were taken every 4 min. White scale bar = 5 µm.

Movie S4

Another example of *bas1Δ* cells showing an oscillating QUEEN ratio. Corresponding to the

data shown in Supplementary Fig. S6A. Left, the QUEEN ratio image; right, bright field

image. In this case, cytokinesis had already been completed at t = 0 min. Images were taken

every 3 min. White scale bar = 5 µm.

Fig. 1

Fig. 2

A *snf1Δadk1Δ* cells in 2% glucose

100' 104' 108' 112' 116' 120' 124' 128' 132' 136' 140' 144' 148' 152'

2.3
0.7

B

172' 176' 180' 184' 188' 192' 196' 200' 204' 208' 212'

336' 340' 344' 348' 352' 356' 360' 364' 368' 372' 376'

3.0
0.5

C

mean [ATP] (mM)

D

Time (min)

E *bas1Δ* cells in 2% glucose

cell-1

cell-2 8' 32' 64' 88' 108' 132' 152' 176' 196' 220' 248' 260'

376' 396' 316' 336' 360' 380' 396' 412' 424' 436' 448'

2
1
0

mean [ATP] (mM)

cell separation

cell1
cell2

Time (min)

Fig. 3

A

Protein folding/Glycosylation

Metabolism

adk1 nega   bas1 nega   snf1 nega   ura6 nega

B

control   40°C   55°C, 1 h   tunicamycin   H$_2$O$_2$

WT
snf1Δ
adk1Δ
snf1Δ adk1Δ
bas1Δ

C

Hsp104-GFP

WT   *snf1Δ*   *adk1Δ*   *snf1Δ adk1Δ*   *bas1Δ*

D

% cells with foci

Hsp104
Pab1

WT *snf1Δ* *adk1Δ* *snf1Δ adk1Δ* *bas1Δ*

Fig. 4

A

B

Hsp104-GFP

glucose	2DG	glucose
0 min	15 min 45 min	110 min

C

GFP-Ubc9ts

mock | 2DG

p=7.3×10⁻⁴

% cells with GFP-Ubc9ts foci

Fig. 5

A

α-synuclein-GFP   GFP

WT
snf1Δ
adk1Δ
bas1Δ
rpn4Δ
snf1Δadk1Δ

B

α-synuclein-GFP

| WT | *snf1Δ* | *adk1Δ* |
| *bas1Δ* | *snf1Δadk1Δ* | *rpn4Δ* |

C

% cells with syn-GFP foci

0   10   20   30   40   50

WT
snf1Δ p=5.1×10⁻³
adk1Δ p=2.2×10⁻⁴
snf1Δadk1Δ p=3.2×10⁻⁶
bas1Δ p=2.3×10⁻⁴
rpn4Δ   N.S.

D

α-synuclein-GFP

− Adenine   + Adenine

WT
bas1Δ
adk1Δ

% cells with syn-GFP foci

0   10   20   30   40   50

WT   N.S.
bas1Δ p=1.2×10⁻³
adk1Δ   N.S.

■ − Adenine
■ + Adenine

E

control   38°C   1.5 mM H₂O₂

wt
snf1Δ
adk1Δ
bas1Δ
rpn4Δ
rpn4Δsnf1Δ
rpn4Δadk1Δ
rpn4Δbas1Δ

wt
snf1Δ
adk1Δ
bas1Δ
atg1Δ
atg1Δsnf1Δ
atg1Δadk1Δ
atg1Δbas1Δ

F

-30   0   30   60   90   120 (min)

glucose | 2DG | glucose
DMSO or MG132

% cells with Hsp104 foci

-30   0   30   60   90   120   0   30   60   90   120

DMSO   MG132

Fig. 6

A

B

Fig. S1

A

B

C

Fig. S2

A

B

Fig. S3

Fig. S4

A

2% glucose

B

Fig. S5

snf1Δadk1Δ cells in 2% glucose

Fig. S6

A

bas1Δ cells in 2% glucose

B

Fig. S7

Fig. S8

Fig. S9