AMP-activated protein kinase and adenylate kinase prevent the ATP catastrophe and

cytotoxic protein aggregation

Masak Takaine[1,2]*, Hiromi Imamura[3], Satoshi Yoshida[1,2,4,5]*

[1]Gunma University Initiative for Advanced Research (GIAR), Gunma University, Maebashi

371-8512, Japan

[2]Institute for Molecular and Cellular Regulation (IMCR), Gunma University, Maebashi

371-8512, Japan

[3]Graduate School of Biostudies, Kyoto University, Kyoto 606-8501, Japan

[4]School of International Liberal Studies, Waseda University, Tokyo, 169-8050, Japan

[5]Japan Science and Technology Agency, PREST

*Correspondence should be addressed to: masaktakaine@gunma-u.ac.jp (M.T.) (Lead

contact) or satosh@waseda.jp (S.Y.)

Running title

ATP homeostasis prevents cytotoxic protein aggregation

Keywords

ATP, homeostasis, proteostasis, yeast, neurodegenerative disease, AMPK, protein

aggregation

ABSTRACT

ATP is the main source of chemical energy in all life and is maintained at several millimolar

in eukaryotic cells. However, the mechanisms responsible for and physiological relevance of

this high and stable concentration of ATP remain unclear. We herein demonstrate that

AMP-activated protein kinase (AMPK) and adenylate kinase (ADK) cooperate to maintain

cellular ATP levels regardless of glucose concentrations. Single cell imaging of ATP-reduced

yeast mutants revealed that ATP concentrations in these mutants underwent stochastic and

transient depletion of ATP repeatedly, which induced the cytotoxic aggregation of

endogenous proteins and pathogenic proteins, such as huntingtin and α-synuclein. Moreover,

pharmacological elevations in ATP levels in an ATP-reduced mutant prevented the

accumulation of α-synuclein aggregates and its cytotoxicity. The removal of cytotoxic

aggregates depended on proteasomes, and proteasomal activity cooperated with AMPK or

ADK to resist proteotoxic stresses. The present results provide the first evidence to show that

cellular ATP homeostasis ensures proteostasis and revealed that stochastic fluctuations in

cellular ATP concentrations contribute to cytotoxic protein aggregation, implying that AMPK

and ADK are important factors that prevent proteinopathies, such as neurodegenerative

diseases.

INTRODUCTION

Adenosine triphosphate (ATP) is the main energy currency used by all living organisms. Due to high demand, the turnover rate of ATP is estimated to be less than one minute in yeast and metazoans (Mortensen, Thaning et al., 2011, Takaine, Ueno et al., 2019). In addition to its role as energy currency, recent studies reported that ATP may influence the balance between the soluble and aggregated states of proteins, suggesting that proteostasis is maintained by energy-dependent chaperones and also by the property of ATP as a hydrotrope to solubilize proteins (Hayes, Peuchen et al., 2018, Patel, Malinovska et al., 2017). Recent proteomic analyses demonstrated that many cellular proteins are insoluble at submillimolar ATP concentrations and become soluble at ATP concentrations > 2 mM (Sridharan, Kurzawa et al., 2019); however, it currently remains unclear whether ATP-dependent protein solubilization/desolubilization play any significant roles in cell physiology.

We recently established a reliable imaging technique to quantify intracellular ATP concentrations in single living yeast cells using the genetically encoded fluorescent ATP biosensor QUEEN (Yaginuma, Kawai et al., 2014), which enables long-term evaluations of ATP homeostasis in individual cells (Takaine et al., 2019). Our findings demonstrated that intracellular ATP concentrations did not vary within a yeast population grown in the same culture (Takaine et al., 2019), which was in contrast to the large variations observed in intracellular ATP concentrations within a bacterial cell population (Yaginuma et al., 2014). Moreover, intracellular ATP concentrations in individual living yeast cells were stably and robustly maintained at approximately 4 mM, irrespective of carbon sources and cell cycle stages, and temporal fluctuations in intracellular ATP concentrations were small (Takaine et al., 2019). Based on these findings, we hypothesized that an exceptionally robust mechanism may exist to precisely regulate ATP concentrations in eukaryotes. It also remains unclear why ATP is stably maintained at a markedly higher concentration than the K_m (Michaelis constant) required for the enzymatic activity of almost all ATPases (Edelman, Blumenthal et al., 1987), and the consequences associated with failed ATP homeostasis in living organisms have not yet been elucidated.

The most promising candidate regulator of ATP homeostasis is AMP-activated protein kinase (AMPK). AMPK, which is activated by AMP and inhibited by ATP (Xiao, Heath et al., 2007), has long been regarded as an important regulator of the whole-body and cellular energy status in eukaryotes (Hardie, Schaffer et al., 2016). AMPK is activated by increases in the AMP:ATP or ADP:ATP ratio (*i.e.,* low-energy state), and regulates its downstream effectors

by phosphorylation to redirect cell metabolism from an anabolic (ATP-consuming) state to

catabolic (ATP-producing) state (Herzig & Shaw, 2017). In the budding yeast *Saccharomyces*

cerevisiae, the sucrose non-fermenting 1 (Snf1) protein kinase complex is the sole AMPK.

Similar to other AMPKs, the yeast Snf1 complex comprises three subunits: the catalytic α

subunit (*SNF1*), scaffolding β subunit (*SIP1*, *SIP2* or *GAL83*), and regulatory γ subunit

(*SNF4*) (Ghillebert, Swinnen et al., 2011). The role of the Snf1 complex in adaptation to

glucose limitations has been characterized in detail (Hedbacker & Carlson, 2008). In the

presence of sufficient glucose in the medium, the Snf1 complex is inactive (Wilson, Hawley

et al., 1996). When glucose concentrations decrease, the Snf1 complex is activated and

phosphorylates the transcriptional repressor Mig1, which then triggers the transcription of

numerous glucose-repressed genes (Carlson, 1999). However, the contribution of AMPK or

the Snf1 complex to cellular ATP levels remains unknown.

Other possible candidate regulators of ATP homeostasis include genes whose mutation leads

to decreases in the cellular content of ATP. However, based on biochemical analyses of cell

populations, few yeast mutants reduced ATP levels (Gauthier, Coulpier et al., 2008,

Ljungdahl & Daignan-Fornier, 2012). Adenylate kinase (ADK) is a key enzyme that

synthesizes ATP and AMP using two ADP molecules as substrates, and the null-mutant of

ADK (*adk1Δ*) was shown to have a reduced cellular ATP concentration (~70% of the wild

type) (Gauthier et al., 2008). Bas1 is a transcription factor that is required for *de novo* purine

synthesis and *bas1Δ* has also a reduced ATP concentration (~50% of the wild type) (Gauthier

et al., 2008). However, the regulation of ATP concentrations and the physiological

consequences of reduced ATP levels in these mutants remain unclear, particularly at the

single cell level.

In the present study, we investigated the roles of AMPK, ADK, and Bas1 in ATP homeostasis

using the QUEEN-based single cell ATP imaging technique. We demonstrated for the first

time that AMPK is involved in the regulation of intracellular ATP concentrations, even under

glucose-rich conditions. Furthermore, time-lapse ATP imaging revealed that cells lacking

both AMPK and ADK frequently underwent transient ATP depletion, while ATP

concentrations oscillated in those lacking Bas1. These ATP dynamics in the mutants were

overlooked in previous biochemical studies. We found that some intrinsic proteins and

aggregation-prone model proteins, including α-synuclein, which is responsible for Parkinson's

disease, aggregated and were cytotoxic in all of the ATP-reduced mutants tested. The present

results suggest that the stable maintenance of ATP is essential for proteostasis and imply that

an ATP crisis promotes proteinopathies, such as neurodegenerative diseases.

RESULTS

AMPK is involved in the maintenance of cellular ATP levels

We recently developed a reliable monitoring system for cytoplasmic ATP concentrations in living yeast cells using the ATP biosensor QUEEN (Takaine et al., 2019). We conducted a more detailed examination of ATP dynamics in wild-type and mutant yeast cells using this system in the present study. We investigated whether the deletion of *SNF1*, which encodes a catalytic subunit of AMPK, affects cellular ATP concentrations. In *SNF1*-null mutant (*snf1Δ*) cells, cellular ATP levels were significantly lower than those in wild-type cells at all glucose concentrations tested (2, 0.2, and 0.05%) (Fig. 1A, B). It is important to note that even under high glucose conditions (2% glucose), in which the Snf1 complex is considered to be inactive, *snf1Δ* cells showed significantly reduced ATP levels. Moreover, the deletion of each gene encoding the β subunit of AMPK (*SIP1*, *SIP2*, or *GAL83*) reduced cellular ATP levels to a similar extent as the deletion of *SNF1*, suggesting that three subtypes of the Snf1 complex are involved in maintaining ATP concentrations (Fig. 1C). On the other hand, the deletion of *MIG1*, which encodes a major downstream target of the Snf1 complex in glucose repression, had a negligible effect on ATP levels (Fig. 1D), suggesting that unknown factors other than Mig1 primarily regulate ATP levels under the control of the Snf1 complex. Collectively, these results demonstrated for the first time that AMPK/SNF1 affect cellular ATP concentrations even under glucose-rich conditions.

Adenylate kinase ADK1 cooperates with AMPK to regulate ATP homeostasis

Adenylate kinase catalyzes the interconversion of adenine nucleotides (ATP+AMP ⟷ 2ADP), which is important for *de novo* adenine nucleotide synthesis and the balance of ATP, ADP, and AMP. Previous biochemical studies reported that the deletion of the adenylate kinase gene reduced ATP concentrations in mouse skeletal muscle cells and budding yeasts (Gauthier et al., 2008, Janssen, Dzeja et al., 2000). We confirmed these findings using an ATP imaging method: *adk1Δ* cells showed significantly lower QUEEN ratios than wild-type cells on average in the presence of sufficient carbon sources (Fig. S1).

In addition to being a key enzyme in purine metabolism, adenylate kinase has also been suggested to cooperate with AMPK in order to monitor the cellular energy state (Hardie, Carling et al., 1998). Therefore, we compared ATP levels in *snf1Δ adk1Δ* double mutant cells with those in *snf1Δ*, *adk1Δ*, and wild-type cells (Fig. 2A, B). The data obtained were represented by a dot plot that shows distribution characteristics in extensive detail. *snf1Δ adk1Δ* cells had significantly lower average ATP levels than single mutant cells. We also

found not only a general reduction, but also a huge variation in ATP levels in the *snf1Δ*,

adk1Δ, snf1Δ adk1Δ cell population, as indicated by the large coefficient of variance (CV)

(Fig. 2B). Furthermore, some cells had extremely low ATP levels in *snf1Δ adk1Δ* (Fig. 2A,

B). These results suggest that Adk1 and the Snf1 complex both synergistically contribute to

ATP homeostasis.

A large pool of adenine nucleotide is important for maintaining cellular ATP

concentrations

We also examined a *bas1Δ* mutant, which is defective in the expression of genes responsible

for adenine biogenesis (Daignan-Fornier & Fink, 1992, Denis, Boucherie et al., 1998).

Consistent with previous biochemical findings (Gauthier et al., 2008), ATP levels quantified

by QUEEN were reduced by ~50% in *bas1Δ* cells (Fig. 2C, D). We found not only a general

reduction, but also a large variation in ATP levels in the *bas1Δ* cell population, as indicated

by the large CV (Fig. 2D). The decrease observed in ATP levels was due to reduced adenine

biosynthesis because the addition of extra adenine to media partially restored ATP levels (Fig.

2D). These results suggest that the sufficient production of adenine nucleotides is essential for

the stable maintenance of ATP levels. Moreover, the role of Bas1 in maintaining ATP levels

appeared to be epistatic to that of Snf1 because *bas1Δ snf1Δ* double mutant cells showed a

similar distribution of ATP levels to *bas1Δ* cells (Fig. S2).

ATP levels temporally fluctuate in ATP mutant cells

To investigate the mechanisms contributing to the large variations in ATP concentrations in

snf1Δ adk1Δ cells in more detail, we employed time-lapse ATP imaging (Fig. 3). We found

that the QUEEN ratio often underwent a rapid decline followed by recovery in *snf1Δ adk1Δ*

cells (see 116 and 132 min in Fig. 3A, C, and Movie S1, and 180 and 356 min in Fig. 3B, D,

and Movie S2). The sudden decrease in ATP concentrations (hereafter called "the ATP

catastrophe") occurred within a few minutes without any sign and was rarely observed in

wild-type cells (Takaine et al., 2019). The ATP catastrophe appeared to be a stochastic event

and cell intrinsic: these events occurred independent of cell cycle stages or cell sizes (compare

Fig. 3C with D). Under some conditions, the QUEEN ratio did not recover after the ATP

catastrophe and the cell died, as judged by the loss of QUEEN signals in the cell (Fig. S3).

These results suggest that the large variations in ATP concentrations observed in *snf1Δ adk1Δ*

cells were not simply due to a mixed population with different basal ATP levels, but were

rather due to the stochastic ATP catastrophe in individual cells.

Time-lapse imaging of *bas1Δ* revealed oscillatory cycles in ATP concentrations (Figs. 3E and

S4A, and Movies S3 and S4): ATP cycling in *bas1Δ* cells was slow (~35 min on average, Fig.

S4B) and distinguishable from that in *snf1Δ adk1Δ* cells; however, the common

characteristics of these mutants were that the concentration of ATP repeatedly reached close

to 0 mM. The ATP oscillation cycle was unsynchronized in the population and independent of

cell cycle progression, suggesting a unique metabolic rhythm intrinsic to each cell. The

oscillatory nature of ATP cycling in the *bas1Δ* mutant may involve a transcription/translation

cycle and will be described elsewhere.

ATP homeostasis is required for preventing protein aggregation.

We recently reported that cellular ATP concentrations are stably maintained at ~4 mM in

budding yeast (Takaine et al., 2019) and herein demonstrated that in the Snf1/AMPK complex,

Adk1 and Bas1 are required for the regulation of ATP homeostasis. However, the

physiological importance of ATP homeostasis remains unknown. To clarify the significance

of high ATP concentrations, we examined the global genetic interactions of *snf1Δ, adk1Δ,*

and *bas1Δ* using CellMap ((Usaj, Tan et al., 2017), thecellmap.org). An *in silico* analysis

identified genes involved in "protein folding/glycosylation" as common negative genetic

interactors with *adk1Δ* and *bas1Δ* (Fig. 4A). Negative genetic interactors of *ura6*, a gene

encoding uridylate kinase that also exhibits adenylate kinase activity, were enriched in the

"protein folding/glycosylation" category (Fig. 4A). We also found that interactors of *snf1*

were implicated in "protein folding/glycosylation. None of these mutants exhibited apparent

genetic interactions with genes in the "metabolism" category (Fig. 4A). The same analysis

using genetic and physical interactors provided similar results and showed that many

interactors were enriched in the "protein turnover" category (Fig. S5). These results imply that

although these three mutants regulate ATP with distinct mechanisms, all three have a

common cellular function.

To examine possible defects in protein folding and turnover (*i.e.*, proteostasis), we challenged

these mutants with various proteotoxic stresses. We found negligible growth defects in ATP

mutants under normal growth conditions with 2% glucose at 30°C (control in Fig. 4B),

suggesting that a high concentration of ATP is not necessary for cellular growth. However,

the *adk1* and *bas1* mutants both exhibited severe growth defects with a high temperature of

40°C, 1 hour of heat shock at 55°C, or in the presence of 0.5 μg/ml of the glycosylation

inhibitor tunicamycin or 2 mM H_2O_2. The *SNF1* deletion increased the stress sensitivity of

adk1Δ (Fig. 4B). This sensitivity to proteotoxic stress suggests that ATP homeostasis mutants

are defective in some aspects of proteostasis. We found that all four mutants tested contained

significantly increased numbers of Hsp104-GFP foci, a marker of protein aggregation

(Josefson, Andersson et al., 2017) (Fig. 4C, D). In contrast to Hsp104-GFP foci, Pab1-GFP, a

marker of stress granule (SG) assembly (Hoyle, Castelli et al., 2007), did not form foci in

ATP mutants, suggesting that protein aggregation and SG assembly are regulated in a distinct

manner (Fig. 4D). These analyses identified abnormal protein aggregation as a common

defect associated with ATP homeostasis mutants for the first time.

The transient depletion of ATP leads to the formation of protein aggregates

To examine whether ATP depletion triggers protein aggregation in living yeast, we artificially

depleted cellular ATP levels by replacing glucose with 2-deoxyglucose (2DG), a strong

inhibitor of glycolysis, in the media and monitored protein aggregation using Hsp104-GFP as

a marker of protein aggregation (Josefson et al., 2017) in wild-type cells. We previously

showed that ATP levels were almost completely depleted 2 minutes after the 2DG treatment

(Takaine et al., 2019). Within 15 min of the 2DG treatment, more than 20% cells contained

Hsp104-GFP foci indicative of protein aggregation (Fig. 5A, B). In contrast to ATP, the

concentration of which recovers to normal levels within 1 min of glucose refeeding (Takaine

et al., 2019), the dissolution kinetics of Hsp104-GFP were significantly slower (up to hours,

Fig. 5A, B).

To further confirm whether a high concentration of ATP is required for protein solubility,

we also tested the Ubc9-ts protein, a model protein that is prone to aggregate (Kaganovich,

Kopito et al., 2008), and found that ATP depletion by the 2DG treatment triggered Ubc9-ts

protein aggregation (Fig. 5C). Thus, not only Hsp104-GFP-positive intrinsic proteins, but also

extrinsic model proteins aggregate after ATP depletion.

SG are assembled in budding yeast cells after stressful conditions, such as glucose depletion

(Hoyle et al., 2007). In contrast to the formation of Hsp104-GFP foci, ATP depletion after the

2DG treatment did not instantly trigger the formation of SG (Fig. S6). Consistent with recent

findings (Jain, Wheeler et al., 2016), the present results suggest that SG formation requires

ATP. We also noted that Hsp104-GFP foci and SG did not co-localize, suggesting that these

structures are derived from distinct mechanisms (Fig. S6). Thus, the artificial depletion of

ATP may trigger abnormal protein aggregation in living yeast cells.

ATP homeostasis is required for the protection of cells from cytotoxicity caused by
protein aggregation

Protein aggregation is often associated with neurodegenerative diseases, such as Alzheimer's,
Huntington's, and Parkinson's diseases (Eftekharzadeh, Hyman et al., 2016). Mitochondrial
failure has also been associated with many neurodegenerative diseases; however, it currently
remains unclear whether energy failure causes protein aggregation because mitochondria also
produce cytotoxic reactive oxygen species (ROS). (Bhat, Dar et al., 2015, Pathak, Berthet et
al., 2013). The abnormal aggregation of α-synuclein has been implicated in Parkinson's
disease (Lashuel, Overk et al., 2013). To clarify whether ATP is preventing the formation of
cytotoxic protein aggregation, we examined the toxicity of α-synuclein-GFP (Syn-GFP) in
budding yeast. As reported previously, the expression of Syn-GFP exhibited negligible
toxicity against wild-type yeast when expressed under the inducible *GAL1* promotor (Fig. 6A)
(Outeiro & Lindquist, 2003, Sharma, Brandis et al., 2006, Wijayanti, Watanabe et al., 2015).
However, *snf1Δ*, *adk1Δ*, and *bas1Δ* are hypersensitive to the expression of Syn-GFP (Fig. 6A).
We also found that *rpn4Δ*, which encodes a key transcription factor for proteasomal subunits
(Xie & Varshavsky, 2001), was very sensitive to Syn-GFP (Fig. 6A), which is consistent with
the concept that Syn-GFP is degraded in the ubiquitin-proteasomal pathway in yeast (Tofaris,
Kim et al., 2011, Wijayanti et al., 2015). We then visualized the cellular localization of
Syn-GFP. Consistent with previous findings (Willingham, Outeiro et al., 2003), Syn-GFP
expressed in yeast mainly localized to the plasma membrane (Fig. 6B, C). In addition to the
plasma membrane, we found that Syn-GFP localized to punctated structures, most likely
corresponding to protein aggregation (Fig. 6B, C). Punctated structures were not as obvious in
rpn4Δ cells defective in proteasomes, suggesting that the accumulation of Syn-GFP puncta
was not simply due to defective degradation.

To investigate whether a high concentration of ATP protects cells from toxic protein
aggregation, we added extra adenine to the medium (Fig. 6D). A previous study demonstrated
that the addition of 300 μM adenine to the medium increased ATP concentrations from 4 to
5.5 mM in wild-type cells and from 1 to 4 mM in *bas1Δ* cells (Gauthier et al., 2008) (similar
results are shown in Fig. 2D), but induced little or no changes in *adk1Δ* cells (from 2 to 2
mM) (Gauthier et al., 2008). Consistent with our hypothesis, the addition of adenine reduced
Syn-GFP toxicity and aggregation in *bas1Δ*, but not *adk1Δ* cells (Fig. 6D). Thus, a high
concentration of ATP prevented Syn-GFP aggregation and toxicity.

We examined another model protein involved in neurodegenerative diseases. PolyQ

containing the huntingtin protein tends to aggregate and has been implicated in Huntington's

disease (Jiang, Poirier et al., 2005). We investigated the toxicity of Htt103Q, a mutant form of

the huntingtin protein, which tends to aggregate and cause cytotoxicity in yeast (Meriin,

Zhang et al., 2002). Consistent with the concept that a high concentration of ATP prevents

protein aggregation, the ATP homeostasis mutants *snf1Δ*, *adk1Δ*, *snf1Δ adk1Δ*, and *bas1Δ*

were very sensitive to Htt103Q expression (Fig. S7).

Proteasomes are essential for the removal of protein aggregates caused by ATP

depletion

Protein aggregation caused by ATP depletion was cytotoxic (Figs. 4B and 6A) and was not

easily dissolved after ATP repletion (Fig. 5). To identify a pathway that is essential for the

removal of aggregates, we examined the involvement of proteasomes and autophagy.

The deletion of *RPN4*, which encodes a transcription factor of proteasomal genes (Xie &

Varshavsky, 2001), down-regulated proteasomal activity (Kruegel, Robison et al., 2011) and

resulted in synthetic growth defects with *adk1Δ*, *snf1Δ bas1Δ* at a high temperature of 38°C

and in the presence of H_2O_2 (Fig. 6E). In contrast to proteasomes, autophagy did not appear to

have genetic interactions with the above mutants (Fig. 6E). The deletion of an essential

component of the autophagic pathway, *ATG1* did not affect the sensitivity of *adk1Δ*, *snf1Δ*,

bas1Δ to a high temperature of 38°C or to H_2O_2 (Fig. 6E). We also did not observe the

accumulation of Hsp104-GFP foci in the autophagy mutants *atg1*, *atg8*, and *atg13* (not

shown).

To investigate the involvement of proteasomes in the removal of protein aggregates after the

transient depletion of ATP, we pretreated cells with the proteasomal inhibitor MG132 or

DMSO and examined the kinetics for the formation of Hsp104-GFP foci after the 2DG

treatment (Fig. 6F) using the drug-sensitive yeast strain Y13206 (Piotrowski, Li et al., 2017).

Under both conditions, more than 90% of cells exhibited Hsp104-GFP foci within 30 min of

the 2DG treatment. More than two-thirds of Hsp104-GFP foci dissolved in the DMSO control,

while less than one-third dissolved in MG132-treated samples, indicating that proteasomes

are required for the dissolution process (Fig. 6F).

DISCUSSION

In the present study, we demonstrated for the first time that the Snf1 complex, budding yeast AMPK, is required for the stable maintenance of cellular ATP concentrations (ATP homeostasis) in collaboration with the adenylate kinase Adk1 (Fig. 7A). This function of the Snf1 complex in ATP homeostasis is independent of glucose concentrations in the medium and Mig1, the major transcriptional repressor involved in glucose repression (Fig. 1); therefore, this is distinct from its well-characterized role in adaptation to glucose limitations. The activity of the Snf1 kinase complex may be sharply tuned depending on the intracellular concentrations of adenine nucleotides or other metabolites indicative of cellular energy to prevent a rapid ATP catastrophe (Fig. 3), even in the presence of sufficient amounts of glucose. It is important to note that the reductions observed in intracellular ATP concentrations in *snf1Δ* cells in the presence of glucose were overlooked in previous biochemical analyses, again demonstrating the usefulness of QUEEN-based ATP imaging.

Since the deletion of *BAS1* caused the greatest reduction in ATP levels and is epistatic to *snf1Δ*, a large pool size of adenine nucleotide is a prerequisite for ATP homeostasis. This assumption is reasonable because the pool size of recyclable ATP restricts ATP concentrations based on the rapid turnover rate of ATP. Bas1 maintains the pool size of ATP by balancing ATP synthesis and irreversible decreases, such as incorporation into RNA and DNA (following conversion to deoxy-ATP), degradation, and excretion in rapidly proliferating yeasts.

We also showed that key regulators of ATP homeostasis play roles in preventing cytotoxic protein aggregation in budding yeast (Fig. 7B). The common feature associated with these mutants is reduced ATP levels, suggesting that high ATP levels are essential for protein solubilization.

A recent study suggested that the requirement for ATP changes depending on its concentration. At concentrations lower than 0.5 mM, ATP mainly serves as a substrate for enzymes, such as protein kinases and heat shock protein chaperones, whereas at concentrations higher than 2 mM, ATP may exert solubilizing effects on disordered proteins (Sridharan et al., 2019). ATP homeostasis may also be required to constantly drive proteasomal protein degradation, which requires large amounts of ATP (Benaroudj, Zwickl et al., 2003, Tanaka, Waxman et al., 1983).

Previous biochemical measurements indicated that although ATP levels were lower in *adk1*

and *bas1* mutants than in the wild type, these mutants still had ATP concentrations that were

higher than 2 mM (Gauthier et al., 2008), which does not directly explain the accumulation of

protein aggregates in these mutants (Fig. 4) because most proteins are expected to be soluble

at > 2 mM ATP (Sridharan et al., 2019). The visualization of ATP dynamics in living cells at

the single cell level revealed that ATP undergoes transient depletion repeatedly (the ATP

catastrophe) in AMPK and ADK mutants, which explains how aggregation-prone proteins

aggregate and cause cytotoxicity. A similar, but distinct instability of ATP was observed in

bas1Δ, further confirming that the ATP catastrophe causes protein aggregation. Although

severe ATP depletion in these mutants was gradually recovered by as yet uncharacterized

negative feedback regulation, the duration period of ~15 min with reduced ATP levels may

induce some proteins to form aggregates that last for generations.

Many neurodegenerative diseases, such as Alzheimer's, Huntington's, and Parkinson's

diseases, are associated with protein aggregation (Eisele, Monteiro et al., 2015, Josefson et al.,

2017). On the other hand, mitochondrial dysfunction and accompanying energy failure in

nerve cells may result in many types of neurodegenerative diseases based on a large body of

evidence (Haelterman, Yoon et al., 2014, Pathak et al., 2013). For example, previous studies

demonstrated that ATP levels in the brain were decreased in patients with early Parkinson's

disease (Mochel, N'Guyen et al., 2012) and also that ATP synthesis in the brain was not

properly regulated in patients with early Huntington's disease (Hattingen, Magerkurth et al.,

2009) and in the corresponding mouse model (Mochel, Durant et al., 2012). Therefore,

protein aggregation induced by the ATP catastrophe, as revealed in the present study, may

link energy failure and protein aggregation, providing a comprehensive insight into the onset

of neurodegenerative diseases. Further studies are warranted to clarify whether the ATP

catastrophe also occurs in the neurons of patients at risk of neurodegenerative diseases or in

the elderly. However, neither biochemical measurements nor mass spectrometry is capable of

detecting the ATP catastrophe because of their insufficient time and space resolution. Thus,

an ATP imaging approach using the yeast model system will be the first choice for

elucidating the molecular mechanisms underlying ATP homeostasis and ATP

catastrophe-induced protein aggregation.

A recent study reported that the activation of AMPK by metformin ameliorated the

progression of Huntington's disease in a mouse model (Arnoux, Willam et al., 2018), and the

potential therapeutic use of metformin for neurodegenerative diseases is being discussed

(Rotermund, Machetanz et al., 2018). Furthermore, the involvement of ATP and ADKs in
preventing the manifestation of Parkinson's disease in mouse models and patients has been
proposed (Garcia-Esparcia, Hernandez-Ortega et al., 2015, Nakano, Imamura et al., 2017).
Protein aggregation induced by the ATP catastrophe may be a general mechanism for the
development of proteinopathies. The present study study using ATP imaging revealed a
physiological consequence of a failure in ATP homeostasis in living cells for the first time
and suggests that ATP homeostasis has potential as a target for preventing/treating
neurodegenerative diseases.

MATERIALS AND METHODS

Yeast strains and plasmids

The budding yeast strains and plasmids used in the present study are listed in Supplementary Tables S1 and S2, respectively. These strains were constructed by a PCR-based method (Janke, Magiera et al., 2004) and genetic crosses. The yeast knockout strain collection was originally purchased from GE Healthcare (cat. # YSC1053).

Media and cell culture

The standard technique for the yeast culture and manipulation was used (Guthrie & Fink, 2002). Synthetic medium (SC) was prepared according to the recipe of Hanscho et al. (Hanscho, Ruckerbauer et al., 2012). 2-Deoxy-D-glucose (2DG), tunicamycin, and MG132 were purchased from FUJIFILM Wako (cat. # 046-06483, 202-08241, and 139-18451, respectively). Tunicamycin and MG132 were dissolved in dimethylsulfoxide (DMSO) to make stock solutions (5 mg/ml and 42 mM, respectively). Cells were grown to the mid-log phase at 30°C in SC before imaging unless otherwise noted.

Microscopy

Cells expressing Hsp104-GFP or GFP-Ubc9ts were concentrated by centrifugation and sandwiched between a slide and coverslip (No. 1.5 thickness, Matsunami, Osaka, Japan). Immobilized cells were imaged using an inverted fluorescent microscope (Eclipse Ti-E, Nikon) equipped with an Apo TIRF 100× Oil DIC N2/NA 1.49 objective lens and electron-multiplying charge-coupled device camera (iXon3 DU897E-CS0-#BV80, Andor) at approximately 25°C. The Hsp104-GFP and GFP-Ubc9ts fluorescent signal was collected from stacks of 11 z-sections spaced by 0.5 μm, and the maximum projections of the optical sections were shown in Figs. 4, 5, and S6. Cells expressing QUEEN were immobilized on a concanavalin A-coated 35-mm glass-bottomed dish (#3971-035, No. 1.5 thickness, IWAKI). The dish was filled with an excess amount of medium (4.5–5 ml) against the cell volume to minimize changes in the chemical compositions of the medium during observations. QUEEN fluorescence was acquired as previously described (Takaine et al., 2019). Cells expressing α-synuclein-GFP were immobilized on a slide glass as described above, and the fluorescence signal was collected from a single z-plane using an inverted fluorescent microscope (Eclipse Ti2-E, Nikon, Tokyo, Japan) equipped with a CFI Plan Apoλ100× Oil DIC/NA1.45 objective

lens and CMOS image sensor (DS-Qi2, Nikon). Images of cells were acquired from several

fields of view for each experimental condition.

Data analysis

Numerical data were plotted using KaleidaGraph software ver. 4.5.1 (Synergy Software).

Means, SDs, and p values were calculated using Excel software (Microsoft, WA, US).

Significance between two sets of data was tested using the unpaired one-tailed Welch's t-test

unless otherwise noted, and was indicated by an asterisk or p value. The horizontal bar in the

dot plot indicates the average of each population. All measurements were repeated at least

twice to confirm reproducibility.

436   ATP concentrations in yeast cells were estimated using QUEEN-based ratiometric

437   imaging, as previously described (Takaine, 2019), (Takaine et al., 2019). The QUEEN ratio is

438   proportional to ATP concentrations and pseudo-colored to reflect its value throughout the

439   present study. The mean QUEEN ratio inside of a cell represents the intracellular ATP

440   concentration of the cell.

441   The autocorrelation functions (ACF) of oscillations in the QUEEN ratio were

442   calculated using R studio software ver. 3.4.1 (R Core Team, 2017). The apparent period of

443   oscillation was estimated from the positive second peak of the correlation coefficient, which

444   was outside the 95% confidence interval and significant ($p < 0.05$), rejecting the assumption

445   that there is no correlation.

ACKNOWLEDGMENTS

447
We are grateful to the Yeast Genetic Resource Center, Y. Ohya, K. Ohashi, H. Takagi, D.
Watanabe, and J. Frydman for providing the yeast strains and plasmids. We thank the
members of the Yoshida/Takaine labs for their support. This work was supported by JSPS
grants 16H04781 (S.Y. and M.T.) and 15K18525 (M.T.) and the Takeda Science foundation
(S.Y.). This work was also supported by the joint research program of the Institute for
Molecular and Cellular Regulation, Gunma University, Japan.
454

455 **AUTHOR CONTRIBUTIONS**

456 M.T. and S.Y. conceived and designed the project. M.T. conducted experiments and the data
457 analysis. H.I. provided a key reagent and expertise. M.T. and S.Y wrote the manuscript with
458 input from H.I.
459

460 **CONFLICT OF INTEREST**

461 The authors declare no competing interests.

REFERENCES

Arnoux I, Willam M, Griesche N, Krummeich J, Watari H, Offermann N, Weber S, Narayan Dey P, Chen C, Monteiro O, Buettner S, Meyer K, Bano D, Radyushkin K, Langston R, Lambert JJ, Wanker E, Methner A, Krauss S, Schweiger S et al. (2018) Metformin reverses early cortical network dysfunction and behavior changes in Huntington's disease. *Elife* 7

Baryshnikova A (2016) Systematic Functional Annotation and Visualization of Biological Networks. *Cell Syst* 2: 412-21

Benaroudj N, Zwickl P, Seemuller E, Baumeister W, Goldberg AL (2003) ATP hydrolysis by the proteasome regulatory complex PAN serves multiple functions in protein degradation. *Mol Cell* 11: 69-78

Bhat AH, Dar KB, Anees S, Zargar MA, Masood A, Sofi MA, Ganie SA (2015) Oxidative stress, mitochondrial dysfunction and neurodegenerative diseases; a mechanistic insight. *Biomed Pharmacother* 74: 101-10

Carlson M (1999) Glucose repression in yeast. *Curr Opin Microbiol* 2: 202-7

Cherry JM, Hong EL, Amundsen C, Balakrishnan R, Binkley G, Chan ET, Christie KR, Costanzo MC, Dwight SS, Engel SR, Fisk DG, Hirschman JE, Hitz BC, Karra K, Krieger CJ, Miyasato SR, Nash RS, Park J, Skrzypek MS, Simison M et al. (2012) Saccharomyces Genome Database: the genomics resource of budding yeast. *Nucleic Acids Res* 40: D700-5

Daignan-Fornier B, Fink GR (1992) Coregulation of purine and histidine biosynthesis by the transcriptional activators BAS1 and BAS2. *Proc Natl Acad Sci U S A* 89: 6746-50

Denis V, Boucherie H, Monribot C, Daignan-Fornier B (1998) Role of the myb-like protein bas1p in Saccharomyces cerevisiae: a proteome analysis. *Mol Microbiol* 30: 557-66

Edelman AM, Blumenthal DK, Krebs EG (1987) Protein serine/threonine kinases. *Annu Rev Biochem* 56: 567-613

Eftekharzadeh B, Hyman BT, Wegmann S (2016) Structural studies on the mechanism of protein aggregation in age related neurodegenerative diseases. *Mech Ageing Dev* 156: 1-13

Eisele YS, Monteiro C, Fearns C, Encalada SE, Wiseman RL, Powers ET, Kelly JW (2015) Targeting protein aggregation for the treatment of degenerative diseases. *Nat Rev Drug Discov* 14: 759-80

Garcia-Esparcia P, Hernandez-Ortega K, Ansoleaga B, Carmona M, Ferrer I (2015) Purine metabolism gene deregulation in Parkinson's disease. *Neuropathol Appl Neurobiol* 41: 926-40

Gauthier S, Coulpier F, Jourdren L, Merle M, Beck S, Konrad M, Daignan-Fornier B, Pinson B (2008) Co-regulation of yeast purine and phosphate pathways in response to adenylic nucleotide variations. *Mol Microbiol* 68: 1583-94

Ghillebert R, Swinnen E, Wen J, Vandesteene L, Ramon M, Norga K, Rolland F, Winderickx J (2011) The AMPK/SNF1/SnRK1 fuel gauge and energy regulator: structure, function and regulation. *FEBS J* 278: 3978-90

Guthrie C, Fink GR (2002) *Guide to yeast genetics and molecular and cell Biology: Part C*. Gulf Professional

Publishing,

Haelterman NA, Yoon WH, Sandoval H, Jaiswal M, Shulman JM, Bellen HJ (2014) A Mitocentric View of

Parkinson's Disease. *Annu Rev Neurosci* 37: 137-159

Hanscho M, Ruckerbauer DE, Chauhan N, Hofbauer HF, Krahulec S, Nidetzky B, Kohlwein SD, Zanghellini

J, Natter K (2012) Nutritional requirements of the BY series of Saccharomyces cerevisiae strains for optimum

growth. *FEMS Yeast Res* 12: 796-808

Hardie DG, Carling D, Carlson M (1998) The AMP-activated/SNF1 protein kinase subfamily: metabolic

sensors of the eukaryotic cell? *Annu Rev Biochem* 67: 821-55

Hardie DG, Schaffer BE, Brunet A (2016) AMPK: An Energy-Sensing Pathway with Multiple Inputs and

Outputs. *Trends Cell Biol* 26: 190-201

Hattingen E, Magerkurth J, Pilatus U, Mozer A, Seifried C, Steinmetz H, Zanella F, Hilker R (2009)

Phosphorus and proton magnetic resonance spectroscopy demonstrates mitochondrial dysfunction in early and

advanced Parkinson's disease. *Brain* 132: 3285-97

Hayes MH, Peuchen EH, Dovichi NJ, Weeks DL (2018) Dual roles for ATP in the regulation of phase

separated protein aggregates in Xenopus oocyte nucleoli. *eLife* 7: e35224

Hedbacker K, Carlson M (2008) SNF1/AMPK pathways in yeast. *Front Biosci* 13: 2408-20

Herzig S, Shaw RJ (2017) AMPK: guardian of metabolism and mitochondrial homeostasis. *Nat Rev Mol Cell*

Biol 19: 121

Hoyle NP, Castelli LM, Campbell SG, Holmes LE, Ashe MP (2007) Stress-dependent relocalization of

translationally primed mRNPs to cytoplasmic granules that are kinetically and spatially distinct from P-bodies. *J*

Cell Biol 179: 65-74

Jain S, Wheeler JR, Walters RW, Agrawal A, Barsic A, Parker R (2016) ATPase-Modulated Stress Granules

Contain a Diverse Proteome and Substructure. *Cell* 164: 487-98

Janke C, Magiera MM, Rathfelder N, Taxis C, Reber S, Maekawa H, Moreno-Borchart A, Doenges G,

Schwob E, Schiebel E, Knop M (2004) A versatile toolbox for PCR-based tagging of yeast genes: new

fluorescent proteins, more markers and promoter substitution cassettes. *Yeast* 21: 947-62

Janssen E, Dzeja PP, Oerlemans F, Simonetti AW, Heerschap A, de Haan A, Rush PS, Terjung RR, Wieringa

B, Terzic A (2000) Adenylate kinase 1 gene deletion disrupts muscle energetic economy despite metabolic

rearrangement. *EMBO J* 19: 6371-81

Jiang H, Poirier MA, Ross CA (2005) A structure-based analysis of huntingtin mutant polyglutamine

aggregation and toxicity: evidence for a compact beta-sheet structure. *Hum Mol Genet* 14: 765-774

Josefson R, Andersson R, Nyström T (2017) How and why do toxic conformers of aberrant proteins

accumulate during ageing? *Essays Biochem* 61: 317

Kaganovich D, Kopito R, Frydman J (2008) Misfolded proteins partition between two distinct quality control

compartments. *Nature* 454: 1088-95

Kruegel U, Robison B, Dange T, Kahlert G, Delaney JR, Kotireddy S, Tsuchiya M, Tsuchiyama S,

Murakami CJ, Schleit J, Sutphin G, Carr D, Tar K, Dittmar G, Kaeberlein M, Kennedy BK, Schmidt M (2011)

Elevated proteasome capacity extends replicative lifespan in Saccharomyces cerevisiae. *PLoS Genet* 7:

e1002253

Lashuel HA, Overk CR, Oueslati A, Masliah E (2013) The many faces of alpha-synuclein: from structure and

toxicity to therapeutic target. *Nat Rev Neurosci* 14: 38-48

Ljungdahl PO, Daignan-Fornier B (2012) Regulation of amino acid, nucleotide, and phosphate metabolism in

Saccharomyces cerevisiae. *Genetics* 190: 885-929

Meriin AB, Zhang X, He X, Newnam GP, Chernoff YO, Sherman MY (2002) Huntington toxicity in yeast

model depends on polyglutamine aggregation mediated by a prion-like protein Rnq1. *J Cell Biol* 157: 997-1004

Mochel F, Durant B, Meng X, O'Callaghan J, Yu H, Brouillet E, Wheeler VC, Humbert S, Schiffmann R,

Durr A (2012) Early alterations of brain cellular energy homeostasis in Huntington disease models. *J Biol Chem*

287: 1361-70

Mochel F, N'Guyen TM, Deelchand D, Rinaldi D, Valabregue R, Wary C, Carlier PG, Durr A, Henry PG

(2012) Abnormal response to cortical activation in early stages of Huntington disease. *Mov Disord* 27: 907-10

Mortensen SP, Thaning P, Nyberg M, Saltin B, Hellsten Y (2011) Local release of ATP into the arterial

inflow and venous drainage of human skeletal muscle: insight from ATP determination with the intravascular

microdialysis technique. *The Journal of Physiology* 589: 1847-1857

Nakano M, Imamura H, Sasaoka N, Yamamoto M, Uemura N, Shudo T, Fuchigami T, Takahashi R,

Kakizuka A (2017) ATP Maintenance via Two Types of ATP Regulators Mitigates Pathological Phenotypes in

Mouse Models of Parkinson's Disease. *EBioMedicine* 22: 225-241

Outeiro TF, Lindquist S (2003) Yeast cells provide insight into alpha-synuclein biology and pathobiology.

Science 302: 1772-5

Patel A, Malinovska L, Saha S, Wang J, Alberti S, Krishnan Y, Hyman AA (2017) ATP as a biological

hydrotrope. *Science* 356: 753-756

Pathak D, Berthet A, Nakamura K (2013) Energy failure: does it contribute to neurodegeneration? *Ann*

Neurol 74: 506-16

Piotrowski JS, Li SC, Deshpande R, Simpkins SW, Nelson J, Yashiroda Y, Barber JM, Safizadeh H, Wilson

E, Okada H, Gebre AA, Kubo K, Torres NP, LeBlanc MA, Andrusiak K, Okamoto R, Yoshimura M,

DeRango-Adem E, van Leeuwen J, Shirahige K et al. (2017) Functional annotation of chemical libraries across

diverse biological processes. *Nature Chemical Biology* 13: 982

R Core Team (2017) R: A Language and Environment for Statistical Computing. In Vienna, Austria: R

Foundation for Statistical Computing

Rotermund C, Machetanz G, Fitzgerald JC (2018) The Therapeutic Potential of Metformin in

Neurodegenerative Diseases. *Front Endocrinol (Lausanne)* 9: 400

Sharma N, Brandis KA, Herrera SK, Johnson BE, Vaidya T, Shrestha R, DebBurman SK (2006) α-synuclein

budding yeast model. *J Mol Neurosci* 28: 161-178

Sridharan S, Kurzawa N, Werner T, Günthner I, Helm D, Huber W, Bantscheff M, Savitski MM (2019)

Proteome-wide solubility and thermal stability profiling reveals distinct regulatory roles for ATP. *Nature*

Communications 10: 1155

Takaine M (2019) QUEEN-based Spatiotemporal ATP Imaging in Budding and Fission Yeast. *Bio-protocol*

9: e3320

Takaine M, Ueno M, Kitamura K, Imamura H, Yoshida S (2019) Reliable imaging of ATP in living budding

and fission yeast. *J Cell Sci* 132

Tanaka K, Waxman L, Goldberg AL (1983) ATP serves two distinct roles in protein degradation in

reticulocytes, one requiring and one independent of ubiquitin. *J Cell Biol* 96: 1580-5

Tofaris GK, Kim HT, Hourez R, Jung JW, Kim KP, Goldberg AL (2011) Ubiquitin ligase Nedd4 promotes

alpha-synuclein degradation by the endosomal-lysosomal pathway. *Proc Natl Acad Sci U S A* 108: 17004-9

Usaj M, Tan Y, Wang W, VanderSluis B, Zou A, Myers CL, Costanzo M, Andrews B, Boone C (2017)

TheCellMap.org: A Web-Accessible Database for Visualizing and Mining the Global Yeast Genetic Interaction

Network. *G3 (Bethesda)* 7: 1539-1549

Wijayanti I, Watanabe D, Oshiro S, Takagi H (2015) Isolation and functional analysis of yeast ubiquitin

ligase Rsp5 variants that alleviate the toxicity of human α-synuclein. *The Journal of Biochemistry* 157: 251-260

Willingham S, Outeiro TF, DeVit MJ, Lindquist SL, Muchowski PJ (2003) Yeast genes that enhance the

toxicity of a mutant huntingtin fragment or alpha-synuclein. *Science* 302: 1769-72

Wilson WA, Hawley SA, Hardie DG (1996) Glucose repression/derepression in budding yeast: SNF1 protein

kinase is activated by phosphorylation under derepressing conditions, and this correlates with a high AMP:ATP

ratio. *Curr Biol* 6: 1426-34

Xiao B, Heath R, Saiu P, Leiper FC, Leone P, Jing C, Walker PA, Haire L, Eccleston JF, Davis CT, Martin

SR, Carling D, Gamblin SJ (2007) Structural basis for AMP binding to mammalian AMP-activated protein

kinase. *Nature* 449: 496-500

Xie Y, Varshavsky A (2001) RPN4 is a ligand, substrate, and transcriptional regulator of the 26S

proteasome: a negative feedback circuit. *Proc Natl Acad Sci U S A* 98: 3056-61

Yaginuma H, Kawai S, Tabata KV, Tomiyama K, Kakizuka A, Komatsuzaki T, Noji H, Imamura H (2014)

Diversity in ATP concentrations in a single bacterial cell population revealed by quantitative single-cell imaging.

Sci Rep 4: 6522

 FIGURE LEGENDS

Fig. 1 AMPK is involved in the maintenance of cellular ATP levels in non-starving cells

(A) Visualization of intracellular ATP levels in wild-type (WT) and *snf1Δ* cells using the

ATP sensor QUEEN. Cells were grown to the mid-log phase in SC-H medium containing the

indicated concentrations of glucose and then imaged. White scale bar = 5 μm. (B) *snf1Δ* cells

had lower ATP levels than wild-type cells. The mean QUEEN ratio inside a single cell (mean

QUEEN ratio/cell), which represents the intracellular ATP level of the cell, was quantified for

each cell from the ratio image. Data are the mean of the cell population ± 1SD (error bar)

normalized to wild-type cells. N = 105–193 cells were scored. Asterisks indicate a significant

difference from WT ($p < 0.05$). (C) ATP levels in single-deletion mutants of AMPK *β*

subunits. Data are the mean of the cell population ± 1SD (error bar) normalized to wild-type

cells. N = 177–223 cells were scored. Asterisks indicate a significant difference between the

two strains (*, $p < 0.05$; **, $p < 0.01$). N.S., no significance. (D) *mig1Δ* cells had slightly

lower ATP levels than wild-type cells. Data are the mean of the cell population ± 1SD (error

bar) normalized to wild-type cells. N = 190–231 cells were scored. Asterisks indicate a

significant difference between the two strains.

Fig. 2 Interconversion and active synthesis of adenine nucleotides are important for

ATP homeostasis

(A) Adk1 and Snf1 synergistically control cellular ATP levels. QUEEN ratio images of ATP

homeostasis mutant cells grown in the medium containing 2% glucose. The asterisk indicates

an example of cells with significantly reduced ATP levels. (B) Mean QUEEN ratios of cells

were translated to ATP concentrations and shown in a dot plot. The horizontal bar indicates

the mean of each population. Asterisks indicate p values versus WT: *=1.5×10^{-59},

=2.3×10^{-34}, *=2.9×10^{-117}. CV: coefficient of variance. N = 134–276 cells were scored.

(C) QUEEN ratio images of *bas1Δ* cells grown in the 2% glucose medium. Growth in media

supplemented with 0.11 mg/ml adenine partially restored the low ATP phenotype of *bas1Δ*.

(D) The ATP concentrations of cells shown in (C) were plotted. Asterisks indicate p values:

*=2.3×10^{-160}, **=8.8×10^{-12} (versus WT in glucose), ***=3.6×10^{-20} (versus *bas1Δ* in glucose).

CV: coefficient of variance. N = 186–296 cells were scored.

Fig. 3 Temporal fluctuations in ATP levels in *snf1Δadk1Δ* and *bas1Δ* cells

(A, C) Time-lapse imaging of QUEEN in *snf1Δ adk1Δ* cells in the 2% glucose medium.

Images at the representative time points were shown. The QUEEN ratio decreased twice

(indicated by arrowheads) within a short interval. See also Movie S1. Data were converted

into ATP concentrations and plotted in (C). (B, D) Another example of time-lapse imaging of

QUEEN in *snf1Δadk1Δ* cells in the 2% glucose medium. The QUEEN ratio decreased twice

(indicated by arrowheads) with a long interval. See also Movie S2. Data were converted into

ATP concentrations and plotted in (D). (E) Time-lapse imaging of QUEEN in *bas1Δ* in the

2% glucose medium. The ATP concentrations of the mother (cell-1) and daughter (cell-2)

were plotted at the bottom. Images at the representative time points were shown on the top.

Note that the QUEEN ratio is synchronized until cells undergo separation at the time point of

76 min indicated by an arrow. After separation, each cell has a unique periodic cycle of ATP.

The movie is available in Movie S3. White scale bar = 5 μm.

Fig. 4 ATP homeostasis is required to prevent protein aggregation

(A) Functional landscape of known interactors of ATP mutants. Negative genetic interactors

of the indicated gene were derived from the SGD database (https://www.yeastgenome.org/)

(Cherry, Hong et al., 2012) and overlaid on a functional map based on the global genetic

interaction network of the yeast genome (Baryshnikova, 2016, Usaj et al., 2017). *URA6*

encodes an uridylate kinase that is essential for viability, which also exhibits adenylate kinase

activity. (B) Each strain of the indicated genotype was serially diluted (five-fold), spotted on

the SC + 2% glucose medium, and grown under the indicated stress conditions. Photos were

taken after 2-3 days. (C) Formation of Hsp104-GFP foci in ATP homeostasis mutants. The

GFP signal (inverted grayscale) was imaged in the log phase culture of the indicated mutant

cells expressing Hsp104-GFP at 35°C. (D) Quantification of data shown in (C). Data from

similar experiments using strains expressing Pab1-GFP, instead of Hsp104-GFP, were also

plotted. Values are the mean ± 1SD (error bars). Asterisks indicate a significant difference

from WT ($p < 0.05$) (N=3–4). White scale bar = 5 μm.

Fig. 5 ATP depletion triggers protein aggregation in living yeast cells

(A) The formation of Hsp104-GFP foci after ATP depletion. Wild-type cells expressing

Hsp104-GFP were grown to the log phase at 35°C in the medium containing 2% glucose. At

the time point of 0 min, the medium was replaced with 20 mM 2DG (red) or 2% glucose (as a

control; blue). Cells were released back to media containing 2% glucose at the time point of

50 min. Cells were imaged at the indicated time points, classified, and scored according to the

number of Hsp104-GFP foci. Values are the mean (N = 3). Asterisks indicate a significant

difference from the 2% glucose treatment ($p < 0.05$). (B) Representative images of cells

analyzed in (A). (C) Formation of Ubc9-ts foci after ATP depletion. Cells expressing

GFP-Ubc9-ts under an inducible *GAL* promoter were grown in the medium containing 2%

galactose (SC-gal) at 33°C, and the medium was then exchanged with 2DG or SC-gal. After

30 min, cells were imaged and scored for the number of GFP-Ubc9-ts foci. Representative

images (inverted grayscale) are shown on the left and summarized on the right. Values are the

mean ± 1SD (error bars) (N = 4). White scale bar = 5 μm.

Fig. 6 Aggregation and cytotoxicity of α-synuclein depends on ATP homeostasis

(A) Each strain of the indicated genotype was transformed with an expression vector carrying

α-synuclein-GFP or GFP, serially diluted (five-fold), spotted on SC + 2% galactose plates,

and then grown at 30°C for 3 days. (B) The localization of α-synuclein-GFP in ATP mutants.

Cells were grown on galactose plates at 30°C for more than 42 h and then imaged.

Representative images of α-synuclein-GFP (inverted grayscale) are shown. (C) Quantification

of the data shown in (B). Cells were classified and scored for the localization pattern of

α-synuclein-GFP. The percentage of cells showing α-synuclein-GFP foci are plotted. Data are

the mean ± 1SD (error bars) from 3–6 independent observations. N = 33–380 cells were

scored in each measurement. *P* values versus WT are shown. N.S., no significance (*p* value >

0.05). (D) *(top)* Each strain of the indicated genotype was transformed with an expression

vector carrying α-synuclein-GFP and grown on galactose plates containing 0 mM (−

Adenine) or 0.3 mM (+ Adenine) adenine at 30°C for 3 days. *(bottom)* Cells were grown on

galactose plates in the absence or presence of adenine at 30°C for 41-45 h and then imaged.

The percentage of cells showing α-synuclein-GFP foci were plotted. Data are the mean ± 1SD

(error bars) from 5 independent transformants. N = 53–258 cells were scored in each

measurement. *P* values versus "− Adenine" are shown. (E) Each strain of the indicated

genotype was serially diluted (five-fold), spotted on the SC + 2% glucose medium, and grown

under the indicated stress conditions. (F) Cells of the drug-sensitive strain Y13206 were

grown to the log phase at 37°C in the medium containing 2% glucose and supplemented with

0.1% DMSO or 0.1% DMSO plus 42 μM MG132 at $t = -30$ min. At $t = 0$ min, these cells

were washed and released in the medium containing 20 mM 2DG ± MG132, and the cells

were then washed and released again in the medium containing 2% glucose ± MG132. Cells

were imaged at the indicated time points and scored for the number of Hsp104-GFP foci.

Data are the mean ± 1SD (error bars). Asterisks indicate a significant difference from DMSO

($p < 0.02$) (N=3).

Fig. 7 Models for ATP homeostasis and its role in proteostasis

(A) Schematic summary of the roles of Snf1, Adk1, and Bas1 in ATP homeostasis. (B) A

schematic model for ATP homeostasis preventing cytotoxic protein aggregation.

 SUPPLEMENTARY INFORMATION

Supplementary figure legends

Fig. S1 Adenylate kinase Adk1 is involved in the maintenance of cellular ATP

concentrations

(A) QUEEN ratio images of wild-type and *adk1Δ* cells grown in 2% glucose or 2% galactose.

White scale bar = 5 μm. (B) Quantification of data shown in (A). The mean QUEEN ratio/cell

was quantified for each cell from ratio images. Values are the mean of the cell population ±

1SD (error bar). N = 182–236 cells were scored. Asterisks indicate a significant difference

from WT ($p < 0.05$).

Fig. S2 ATP levels in *bas1Δ snf1Δ* cells

(A) QUEEN ratio images of wild-type, *bas1Δ, snf1Δ,* and *bas1Δ snf1Δ* cells. (B)

Quantification of data shown in (A). The mean QUEEN ratio/cell was quantified for each cell

from ratio images. Data are shown as a dot plot. The horizontal bar in the plot indicates the

mean of each population. Significance between two sets of data was tested using the unpaired

two-tailed Welch's *t*-test and indicated by *p* values. N.S., no significance (*p*-value > 0.05).

Fig. S3 Time-lapse imaging of QUEEN in *snf1Δ adk1Δ* cells

Time-lapse imaging of QUEEN in *snf1Δ adk1Δ* cells in the 2% glucose medium. An example

of *snf1Δ adk1Δ* cell showing an irreversible decrease in the QUEEN ratio (indicated by an

arrowhead). In this case, the cell eventually died (indicated by an arrow). Images at the

representative time points were shown. The ATP concentration was plotted at the bottom.

White scale bar = 5 μm.

Fig. S4. Oscillatory behavior of the ATP concentration visualized in *bas1Δ* cells

(A) Another example of *bas1Δ* cells showing an oscillating QUEEN ratio. In this case,

cytokinesis had just been completed at t = 0 min. ATP concentrations in cell-1 and cell-2 were

plotted at the bottom. See also Movie S4. White scale bar = 5 μm. (B) Autocorrelation

function of the QUEEN ratio calculated from the data on cell-1 in Fig. 3E. Blue dotted lines

indicate the 95% confidence interval. An arrow indicates the second peak of the correlation

and corresponds to the apparent period.

Fig. S5 *In silico* analysis of interactors of ATP mutants

Genetic and physical interactors of the indicated genes were derived from the SGD database

(https://www.yeastgenome.org/) (Cherry et al., 2012) and overlaid on a functional map based

on the global genetic interaction network of the yeast genome (Baryshnikova, 2016, Usaj et

al., 2017). *URA6* encodes an uridylate kinase essential for viability, which also exhibits

adenylate kinase activity.

Fig. S6 Simultaneous observation of Hsp104 and Pab1 foci

Wild-type cells expressing Hsp104-GFP and Pab1-RedStar2 were grown to the log phase at

37°C in the medium containing 2% glucose. Cells were washed and released either in the

medium containing 20 mM 2DG (top) or in the medium lacking glucose (bottom), and then

imaged after 30 and 60 min. White scale bar = 5 μm.

Fig. S7 Cytotoxicity of polyQ containing the huntingtin protein in wild-type and

ATP-mutant yeast cells

Each strain of the indicated genotype was transformed with an expression vector carrying

Htt103Q and grown on glucose ((-), no induction) or raffinose ((+), leaky expression) plates

at 30°C for 3 days.

Table S1. *Strains used in the present study*

Name	Genotype	Source	Figure
MTY3008	*snf1Δ::kanMX6 leu2Δ0 lys2Δ0 ura3Δ0*	Lab stock	4B, 6A-C, 6E, S7
MTY3015	*his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	Lab stock	4B, 6A-E, S7
MTY3049	*adk1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	Lab stock	4B, 6A-E, S7
MTY3118	*bas1Δ::kanMX6 leu2Δ0 lys2Δ0 ura3Δ0*	Lab stock	4B, 6A-E, S7
MTY3143	*his3Δ1:: 2×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0*	This study	S1
MTY3149	*adk1Δ::kanMX6 his3Δ1:: 2×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0*	This study	S1
MTY3264	*his3Δ1:: 3×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0 MYO1-3mCherry-hphMX6*	Takaine et al., 2019	2A, B, S2
MTY3270	*bas1Δ::kanMX6 his3Δ1:: 3×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0 MYO1-3mCherry-hphMX6*	This study	2C, D, 3E, S2, S4
MTY3293	*adk1Δ::kanMX6 his3Δ1:: 3×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0 MYO1-3mCherry-hphMX6*	This study	2A, B
MTY3355	*adk1Δ::kanMX6 snf1Δ::natNT2 his3Δ1:: 3×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0 MYO1-3mCherry-hphMX6*	This study	2A, B, 3A-D, S3
MTY3371	*snf1Δ::kanMX6 his3Δ1:: 3×pRS303-P$_{TEF}$-QUEEN-2m-T$_{CYC1}$ leu2Δ0 lys2Δ0 ura3Δ0 MYO1-3mCherry-hphMX6*	This study	2A, B, S2
MTY3412	*adk1Δ::kanMX6 snf1Δ::natNT2 leu2Δ0 lys2Δ0 ura3Δ0*	This study	4B, 6A-E, S7
MTY3420	*Hsp104-yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	4C, D, 5A, B
MTY3421	*snf1Δ::kanMX6 Hsp104-yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	4C, D
MTY3422	*bas1Δ::kanMX6 Hsp104-yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	4C, D
MTY3424	*adk1Δ::kanMX6 Hsp104-yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	4C-D
MTY3425	*adk1Δ::kanMX6 snf1Δ::natNT2 Hsp104-yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	4C, D
MTY3489	*atg1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	Lab stock	6E
MTY3493	*snq2Δ::KlLeu2; pdr3Δ::Klura3; pdr1Δ::natMX4; can1Δ::STE2prSp_his5 lyp1Δ*	From Y.	Parental

	his3Δ1 leu2Δ0 ura3Δ0 met15Δ0 LYS2+	Ohya, (Piotrowski et al., 2017)	strain of MTY3501
MTY3501	*Hsp104-yeGFP-hphNT1 snq2Δ::KILeu2; pdr3Δ::KIura3; pdr1Δ::natMX4; can1Δ::STE2prSp_his5 lyp1Δ his3Δ1 leu2Δ0 ura3Δ0 met15Δ0 LYS2+*	This study	6F
MTY3503	*Pab1- yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	4D
MTY3504	*Hsp104-yeGFP-hphNT1 Pab1-RedStar2-natNT2 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	S6
MTY3505	*snf1Δ::kanMX6 Pab1- yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	4D
MTY3506	*adk1Δ::kanMX6 Pab1- yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	4D
MTY3507	*adk1Δ::kanMX6 snf1Δ::natNT2 Pab1- yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	4D
MTY3508	*bas1Δ::kanMX6 Pab1- yeGFP-hphNT1 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	4D
MTY3513	*atg1Δ::natNT2 adk1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	6E
MTY3514	*atg1Δ::natNT2 snf1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	6E
MTY3515	*atg1Δ::natNT2 bas1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	6E
MTY3516	*rpn4Δ::natNT2 adk1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	6E
MTY3517	*rpn4Δ::natNT2 snf1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	6E
MTY3518	*rpn4Δ::natNT2 bas1Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	This study	6E
MTY3525	*rpn4Δ::kanMX6 his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	Lab stock	6A-C, 6E

757 Table S2. *Plasmids used in the present study*

Name	Structure	Source	Purpose
MTP3091	pESC-Leu-GFP-Ubc9-ts	Judith Frydman (pJF1089)	Fig. 5C
MTP3088	p426 103Q GAL	Addgene (cat.# 1188)	Fig. S7
MTP3108	pYES2-α-synuclein-GFP	H. Takagi (Wijayanti et al., 2015)	Fig. 6A-D
MTP3090	pYES2-GFP	K. Ohashi	Fig. 6A

Captions for supplementary movies

Movie S1

Time-lapse imaging of QUEEN in *snf1Δadk1Δ* cells in the 2% glucose medium.

Corresponding to the data shown in Fig. 3A. The QUEEN ratio decreased twice (116 and 132

min) within a short interval. White scale bar = 5 μm.

Movie S2

Another example of *snf1Δadk1Δ* cells showing a sudden decrease in the QUEEN ratio.

Corresponding to the data shown in Fig. 3B. The QUEEN ratio decreased twice (180 and 356

min) with a long interval. White scale bar = 5 μm.

Movie S3

Oscillatory behavior of the QUEEN ratio in *bas1Δ* cells. Corresponding to the data shown in

Fig. 3E. Left, the QUEEN ratio image; middle, Myo1-mCherry (inverted grayscale image);

right, bright field image. Images were taken every 4 min. White scale bar = 5 μm.

Movie S4

Another example of *bas1Δ* cells showing an oscillating QUEEN ratio. Corresponding to the

data shown in Supplementary Fig. S4A. Left, the QUEEN ratio image; right, bright field

image. In this case, cytokinesis had already been completed at t = 0 min. Images were taken

every 3 min. White scale bar = 5 μm.

Fig. 1

A

B

C

D

Fig. 2

A

WT   snf1Δ

adk1Δ   snf1Δ adk1Δ *

cells in 2% glucose

B

CV: 0.11   0.223   0.243   0.282

mean [ATP] (mM)

WT   snf1Δ   adk1Δ   snf1Δ adk1Δ

C

glucose   glucose + adenine

bas1Δ cells in 2% glucose

D

CV: 0.14   0.41   0.10   0.45

mean [ATP] (mM)

● WT
◆ bas1Δ

glucose   glucose + adenine

Fig. 3

A *snf1Δadk1Δ* cells in 2% glucose

B

C

D

E *bas1Δ* cells in 2% glucose

cell separation

Fig. 4

A

Protein folding/Glycosylation

Metabolism

adk1 nega   bas1 nega   snf1 nega   ura6 nega

B

	control	40°C	55°C, 1 h	tunicamycin	H₂O₂

WT
snf1Δ
adk1Δ
snf1Δ adk1Δ
bas1Δ

C

Hsp104-GFP

WT   *snf1Δ*   *adk1Δ*   *snf1Δ adk1Δ*   *bas1Δ*

D

Fig. 5

A

B

Hsp104-GFP

glucose 0 min | 2DG 15 min | 2DG 45 min | glucose 110 min

C

GFP-Ubc9ts

mock | 2DG

$p=7.3 \times 10^{-4}$

% cells with GFP-Ubc9ts foci

Fig. 6

A

α-synuclein-GFP   GFP

WT
snf1Δ
adk1Δ
bas1Δ
rpn4Δ
snf1Δadk1Δ

B

α-synuclein-GFP

WT   snf1Δ   adk1Δ
bas1Δ   snf1Δadk1Δ   rpn4Δ

C

% cells with syn-GFP foci

WT
snf1Δ   p=5.1 × 10⁻³
adk1Δ   p=2.2 × 10⁻⁴
snf1Δadk1Δ   p=3.2 × 10⁻⁶
bas1Δ   p=2.3 × 10⁻⁴
rpn4Δ   N.S.

D

α-synuclein-GFP
− Adenine   + Adenine

WT
bas1Δ
adk1Δ

% cells with syn-GFP foci

WT   N.S.
bas1Δ   p=1.2 × 10⁻³
adk1Δ   N.S.

■ − Adenine
■ + Adenine

E

control   38°C   1.5 mM H₂O₂

wt
snf1Δ
adk1Δ
bas1Δ
rpn4Δ
rpn4Δsnf1Δ
rpn4Δadk1Δ
rpn4Δbas1Δ

wt
snf1Δ
adk1Δ
bas1Δ
atg1Δ
atg1Δsnf1Δ
atg1Δadk1Δ
atg1Δbas1Δ

F

-30   0   30   60   90   120 (min)
glucose   2DG   glucose
DMSO or MG132

% cells with Hsp104 foci

-30 0 30 60 90 120 0 30 60 90 120
DMSO   MG132

Fig. 7

A

B

Fig. S1

A

B

Fig. S2

A

2% glucose

B

Fig. S3

A

$bas1\Delta$ cells in 2% glucose

B

Fig. S5

adk1 interactors   bas1 interactors   snf1 interactors

ura6 interactors

Fig. S6

Fig. S7