AMP-activated protein kinase is a key regulator of acute neurovascular

permeability

Silvia Dragoni[1], Bruna Caridi[1], Eleni Karatsai[2], Thomas Burgoyne[1], Mosharraf H. Sarker[1,3], Patric

Turowski[1,4]

1. Institute of Ophthalmology, University College London, 11-43 Bath Street, London EC1V 9EL, UK.

2. NIHR Moorfields Biomedical Research Centre, Moorfields Eye Hospital, London, UK.

3. School of Science, Engineering & Design, Teesside University, Stephenson Street, Middlesbrough

TS1 3BA, UK.

4. Corresponding author: p.turowski@ucl.ac.uk

ABSTRACT

Many neuronal and retinal disorders are associated with pathological

hyperpermeability of the microvasculature. We have used explants of rodent retinae

to measure and manipulate acute neurovascular permeability and signal transduction

to study the role of AMP-activated protein kinase (AMPK). Following stimulation with

either vascular endothelial growth factor (VEGF-A) or bradykinin (BK), AMPK was

rapidly and strongly phosphorylated and acted as a key mediator of permeability

downstream of Ca^{2+} and Ca^{2+}/calmodulin-dependent protein kinase kinase (CAMKK).

Accordingly, AMPK agonists potently induced acute retinal vascular leakage. AMPK

activation led to phosphorylation of endothelial nitric oxide synthase (eNOS), which in

turn increased VE-cadherin phosphorylation on Y685. In parallel, AMPK also mediated

phosphorylation of p38 MAP kinase and HSP27, indicating that it regulated

paracellular junctions and cellular contractility, both previously associated with

endothelial permeability. Endothelial AMPK provided a missing link in neurovascular

permeability, connecting Ca^{2+} transients to the activation of eNOS and p38,

irrespective of the permeability-inducing factor used. Collectively, the ex-vivo retina

model was easily accessible and, due to its compatibility with small molecule

antagonists/agonists and siRNA, constitutes a reliable tool to study regulators and

mechanism of neurovascular permeability.

KEY WORDS

Neurovascular leakage; AMP-activated protein kinase; VEGF-A, bradykinin; retina

INTRODUCTION

Leakage within the vascular system can be the cause or a significant co-morbidity of

a variety of pathologies and is the consequence of endothelial hyperpermeability,

which leads to extravasation of fluids and proteins, resulting in interstitial oedema[1]. In

the nervous system, where the vasculature is uniquely impermeable and is referred to

as the blood-brain barrier (BBB), vascular leakage accompanies stroke, multiple

sclerosis, Parkinsons Disease as well as various forms of dementia[2, 3]. Neurovascular

leakage also affects the blood-retinal barrier (BRB), where it is a hallmark feature of

diabetic retinopathy and neovascular age-related macular degeneration[4]. Leakage

during retinopathies is driven by permeability-inducing factors (PIFs), most

prominently by the angiogenic growth factor VEGF-A; and VEGF-A antagonists are

successfully used to reduce oedema and abnormal vessel growth, and restore

neuronal dysfunction[5]. Meta-analysis of the clinical use of anti-VEGFs in diabetic

macular oedema suggests that PIFs other than VEGF-A play an important role in the

pathogenesis of retinal leakage disease[6], including angiopoeitin-2[7], lyso-

phosphatidylcholine[8] and BK[9].

PIFs induce both acute and chronic vascular leakage. For instance, exposure of the

vascular endothelium to VEGF-A leads to acute permeability that usually lasts less

than 30 minutes. If not resolved persistent leakage ensues that chronically impairs

vascular integrity[1, 10]. Different PIFs bind to distinct endothelial cell (EC) surface

receptors, but ultimately all permeability responses involve paracellular junction

modulation or formation of transport vesicles[1, 2], suggesting that ECs regulate

hyperpermeability through a core molecular machinery and common downstream

signalling. Indeed, e.g. Ca^{2+} transients, phosphorylation of the MAP kinase p38 and

enhanced actin contractility are associated with all vascular permeability, as is

activation of eNOS[11,12,13,14]. Adherens and tight junction modulation is associated with

paracellular permeability and the phosphorylation of VE-cadherin (VE-cad) is

associated with acute permeability in the periphery and the retina[15, 16]. In the retina the

phosphorylation of occludin on S490 and its subsequent internalisation also plays an

important role, at least in a more chronic setting[17]. Identifying core signalling, on which

all leakage responses depend, is desirable for therapeutic development as it would

allow treatment without prior knowledge of specific known (or unknown) extracellular

PIF.

In the brain and the retina, entirely different signalling is induced by VEGF-A in ECs

when it is added luminally (from the blood side) or abluminally (from the tissue side),

with leakage-inducing signalling entirely restricted to abluminal stimulation[12]. For

instance, leakage-associated p38 activation is triggered by abluminal (basal) VEGF-A

stimulation, whilst activation of the PI3K/AKT pathway, which does not mediate

permeability, is only seen following luminal (apical) stimulation. Thus, signalling

specific to leakage can be inferred by comparing cellular stimulation following luminal

and abluminal addition of VEGF-A. Conversely, BK efficiently induces permeability

from the basal as well as apical side of cerebral or retinal ECs.

AMP-activated protein kinase (AMPK) is a phylogenetically conserved energy sensor

that regulates energy homeostasis by coordinating metabolic pathways and thus

balancing energy requirement with nutrient supply[18]. Previous studies suggest that

AMPK acts as a protector of the BBB integrity, for instance by preventing LPS-

enhanced NAD(P)H oxidase expression in ECs and the consequent barrier

dysfunction and enhanced permeability[19]. Moreover, AMPK mediates up-regulation of

BBB functions induced in vitro by metformin, a drug used for the treatment of

diabetes[20]. Nevertheless, in the retina AMPK activation can lead to the breakdown of

the outer, non-vascular BRB, which is constituted by the retinal pigment epithelium[21].

However, if and how AMPK contributes to permeability induction by agonists such as

VEGF-A or BK in neural microvessels is unknown.

To identify and validate core components mediating acute permeability in

neurovascular ECs, we adopted an ex vivo retinal preparation, originally described for

rats[22]. Development of this method allowed measurement of real time changes of

permeability and signalling in intact BRB vessels from rat and mouse. Importantly, this

model system was compatible with precise pharmacokinetic agonist studies, parallel

IHC staining and manipulation using siRNA. Our workflow can be used to identify core

regulators of CNS endothelial hyperpermeability and was validated by identifying

AMPK as a novel, key regulator linking VEGF-A or BK-induced Ca^{2+} transients to

eNOS activation and VE-cad phosphorylation.

MATERIALS AND METHODS

Materials

Recombinant rat VEGF-A (165) was purchased from R&D Systems (Abingdon, United Kingdom). Bradykinin, Sulforhodamine-B, Evans blue, SU-1498, SB-203580, Compound-C, STO-609, L-NAME, BAPTA-AM, AICAR and A769662 were purchased from MERCK (Dorset, United Kingdom). Polyclonal antibodies specific for p38, Hsp27, AMPKα, eNOS and their phosphorylated forms (p38 Thr180/Tyr182, pHSP27 Ser82, AMPK Thr172 and eNOS Ser1177) were from Cell Signaling Technology (Beverly, MA). Polyclonal antibodies against phosphorylated VE-cad (p-Y658-VEC and p-Y685-VEC) were a gift from E. Dejana (Milan, Italy).

Animals

Wistar female rats (7-10 weeks old) and C75BL/6J mice (7-12 weeks old) were purchased from Charles River Laboratories. All animal procedures were performed in accordance with Animal Welfare Ethical Review Body (AWERB) and Association for Research in Vision and Ophthalmology (ARVO) Statement for the Use of Animals in Ophthalmic and Vision Research guidelines and under a UK Home Office licence.

Methods

Brain microvascular EC isolation and culture

Microvessels were isolated from rat cortical grey matter by collagenase dispase digestion and BSA and Percoll density gradient centrifugation[12]. Purified vessels were seeded onto collagen IV/fibronectin-coated Costar Transwells (Fisher Scientific) at high density (vessels from 6 rat brains per 40 cm^2). Cells were grown in EGM2-MV

(Lonza), with 5 mg/ml puromycin from the second day for 3 days, for 2–3 weeks until

their TEER plateaued at values above 200 $\Omega.cm^2$.

The human brain MVEC line hCMEC/D3 was also grown in EGM2-MV as previously

described[23].

Transendothelial Electrical Resistance (TEER)

Changes in the TEER were determined by time-resolved impedance spectroscopy of

primary cerebral rat brain microvascular ECs grown on 12 mm Transwells, using a

CellZscope (Nanoanalytics). Before the addition of VEGF-A and BK, TEER values

were 500-800 $\Omega.cm^2$.

Immunocytochemistry

Primary cerebral rat brain microvascular ECs were fixed with methanol (−20°C).

Staining was performed as previously described using antibodies against occludin

(OC-3F10, Invitrogen)[24] or VE-cad[25].

Immunogold Electron Microscopy

Cryo-immuno electron microscopy (EM) was performed as previously described[12].

Briefly, hCMEC/D3 cells were fixed in 4% PFA and 0.1% glutaraldehyde. Sections

were stained using antibodies against the extracellular domain (TEA 1.31; Serotech)

or the C terminus (sc-6458; Santa Cruz) of VE-cad. Image J (NIH) was used to process

images and measure the distance among the gold particles and the interendothelial

junctions.

Retinal explant preparation

Retinal explants were prepared essentially as described before[22]. A Wistar female rats or C75BL/6J mouse was killed by overdose of CO_2. The common carotid artery was carefully exposed and cannulated with a glass microcannula. The head vasculature was then flushed first with heparinized saline (300U/L heparin in 0.9% NaCl. Mouse: 5 ml; rat: 20 ml), then with stabilizing solution (10 mM Mg^{2+}, 110 mM NaCl, 8 mMKCl, 10 mM HEPES, 1 mM CaCl2, pH 7.0. 10 µM Isoproterenol to be added before use. Mouse: 5 ml; rat: 20 ml), also referred to as cardioplegic solution containing isoproterenol, and finally with the same solution supplemented with 5 g/L Evans blue dye in 10% albumin (mouse: 5 ml; rat: 20 ml) for subsequent visualisation of the vasculature. Next, an eye was surgically removed and and the retina isolated, together with the attached sclera. The retina was flattened onto a transparent silicone medium (SYLGARD® 184 SILICONE ELASTOMER KIT by Dow Corning), kept in position by a metal ring, and the resulting well sealed by grease. Throughout the procedure the retina was continuously superfused with Krebs solution (124 mM sodium chloride, 5 mM potassium chloride, 2 mM MgSO4, 0.125 mM NaH2 PO4, 22 mM NaHCO3, 2 mM CaCl2, pH 7.4. 5 mM glucose and 0.1% BSA to be added before use).

Permeability measurements

Retinal explants were mounted for visualisation and further experimentation on an upright Zeiss Axiophot fluorescent microscope. A radial vessel of a superfused retinal explant was cannulated at ca. 150 µm from the optic nerve using a microinjection needle (tip diameter 1-5 µm, sharpened to a bevel of < 30°) and the entire retinal vasculature injected with sulforhodamine-B (1 mg/ml in Krebs solution). Illumination was switched to fluorescence and the vessels were visualized under a TRITC filter

using a Olympus 40X water immersion objective. For permeability measurements a

microvessel was chosen at least 200 µm away from the cannulated radial vessel.

Fluorescent content of the vessel was recorded continuously by time lapse (1 frame/2

s) on a Hamamatsu CCD camera for at least 2 min. A baseline was recorded for ca.

30 s, before VEGF-A or BK (in Krebs solution) was added on the top of the retina.

Time lapse series were analysed using ImageJ. Time-dependent fluorescence

intensity data of the chosen vessel was derived from a square region of interest (ca.

18 x 18 pixels) (Supplemental Figure 1a and b). Fluorescence in the immediate vicinity

of the microvessel was measured and subtracted from the vessel fluorescence

measurements. Pixel intensity measurements were charted against time, and

permeability values were computed by fitting data to the exponential equation $C_t =$

$C_0 * e^{-kt}$, where $k = 4P/d$ and d is the diameter of the vessel[12].   The difference in

permeability between pre-treatment and post-treatment resulted in the absolute

permeability change associated with the treatment regimen.

Immunohistochemistry

After dissection, retinae from rat or mouse were fixed with 4% PFA at room

temperature for 1 h. After 30 min of blocking (3% Triton X-100, 1% Tween, 0.5%

Bovine Serum Albumin in 2x PBS), retinae were incubated with primary antibodies

against Isolectin B4 (IB-4), P-p38, P-HSP27, P-eNOS, P-AMPKα, claudin-5 and P-VE-

Cadherin at 4 ºC o/n. Retinae were washed and incubated with matching Alexa Fluor

conjugated secondary antibodies at room temperature for 2 h. Finally, retinae were

washed and mounted using Mowiol 4-88 mounting medium (Sigma). More details can

be found in ref 12.

siRNA-mediated Knockdown of claudin-5 and AMPKα

Specific SiRNA sequences targeting claudin-5 or the α subunit of AMPK were purchased from Dharmacon (Chicago, IL). Mice were anesthetised by intraperitoneal injection of 100 ul of 6% Narketan (ketamine) and 10% Dormitor (medetomidine) in sterile water. 2 µl of the siRNA (1ng/ml in sterile PBS) were injected intravitreously in the right eye under a stereomicroscope, using a Hamilton syringe with a 3 degrees Hamilton RN needle (Esslab). Two microliters of scrambled siRNA were injected into the left eye as a control. To inject, an initial puncture was made to the superior nasal sclera, at the level of the pars plana. Then, the tip of the needle was further introduced through the puncture hole with a 45-degree angle into the vitreous body. Retinae were isolated 72 h after the injection.

Phosphoantibody array

Mature monolayers of primary, unpassaged brain microvascular ECs grown on 24 mm Costar Transwells were stimulated with VEGF-A (50 ng/ml) from the apical or basal side for 5 or 30 min. Cells from 2 Transwells were combined by lysis in 200 µL of lysis buffer and subjected to screening using Human Phospho-Kinase Array Proteome Profiler Array (R&D Systems; ARY003B) exactly according to the manufacturer's instructions. Arrays were exposed for varying amounts of time to capture signals in the linear range and quantified by densitometric scanning and ImageJ (NIH). Signals were normalised using array-internal controls. Results were expressed as fold-differences between apical versus basal signals.

Western Blots

Cell lysates were prepared as previously described[12]. Proteins were separated by

SDS-PAGE and transferred to nitrocellulose by semidry electrotransfer. Membranes

were blocked o/n and then incubated with the appropriate antibody diluted at 1:2,000.

Membranes were washed three times with TBS/0.1% Tween-20 before 1h incubation

with an anti-mouse or anti-rabbit HRP-conjugated IgG (GE Healthcare) at a dilution of

1:10,000 and 1:5,000, respectively. Membranes were developed using the ECL

reagents (Roche) and exposed to X-ray film. Protein bands were evaluated by

densitometric quantification, normalized against the amount of total protein, and

GADPH or Tubulin.

Statistics

TEER measurements of three independent cell monolayers were combined and

expressed as mean ± SD. Significant differences were determined by two ways

ANOVA with replication, with significance levels set at 0,05, followed by post-hoc

Bonferroni's multiple comparison test.

Densitometric quantification of four independent immunoblots were determined by

changes in phosphoprotein content normalized to tubulin or GADPH/total protein

loading controls, with values expressed as fold increase. Data were presented as

mean ± SD. Statistics were performed using one-way ANOVA with significance levels

set at 0.05, followed by post-hoc Dunnett's tests.

Permeability measurements from at least 4 different ex vivo retinae were combined

and expressed as mean ± SD. Significant differences were determined via t-test

between the control and each inhibitor.

Significance levels were set to *, $p < 0.05$; **, $0.001 < p < 0.01$, ***, $p \leq 0.001$.

RESULTS

VEGF-A and BK-induced permeability and junctional changes in brain microvascular ECs.

Treatment of primary rat brain microvascular ECs with VEGF-A or BK significantly reduced TEER, indicating that paracellular permeability was induced (Figure 1a and b). TEER dropped immediately and reached a minimum within less than 5 min after addition of either VEGF-A or BK before reverting to control levels within 1 h. Thereafter, another significant, but more modest reduction in TEER was observed, indicative of a more chronic change in cell monolayer permeability. In order to correlate TEER changes with paracellular junction breakdown, the distribution of occludin and VE-Cad was analysed in VEGF-A- and BK-stimulated primary brain microvascular ECs (Figure 1c). As judged by confocal microscopy, occludin expression and distribution remained unchanged for up to 2 h of VEGF-A or BK stimulation. VE-cad levels also remained unchanged, but a broadening of the staining was observed within 5 min of the addition of the PIF, in particular at and around tricellular junctions. In agreement, cryo-immuno–EM of hCMEC/D3 cells revealed a significant relocation of VE-cad from the junctions to the cell interior by an average distance of 55 and 66 nm following a 5-min stimulation with VEGF-A and BK, respectively (Figure 1d-g). These results showed that single addition of either VEGF-A or BK induced acute and chronic permeability, and that the acute response was accompanied by VE-cad redistribution away from the junctions.

Validation of a modified ex-vivo retinal model

In order to study the acute phase of VEGF-A- and BK-induced permeability in intact retinal microvessels in real time, we adopted an ex-vivo retinal model[22]. Permeability measurements used rat retinal explants, in which the vasculature was stabilised with a cardioplegic solution. To assess if the ex vivo preparation and perfusions led to alterations of the retinal vasculature and to determine stability of the preparation, we compared directly perfused fixed retinae with others, perfused with cardioplegic solution and left under superfusion with Krebs solution for 1 h before fixation. Subsequent whole mount staining for the tight junction protein claudin-5 and the adherens junction protein VE-cad revealed characteristic strands of continuous paracellular staining (Figure 2a, b) in both preparations. Importantly, the staining pattern was indistinguishable between the two different preparations, indicating that the perfusion did not cause significant disturbances of endothelial junctions. Permeability measurements were carried out by monitoring sulforhodamine B loss from individual microvessels (Supplemental Figure 1a, b). Baseline permeability to sulforhodamine-B was very low and on average $0.2 \pm 0.16 \times 10^{-6}$ cm/s. Taken together these data showed that morphological and barrier properties of the retinal microvasculature were well preserved in these preparations.

Stimulation of the ex-vivo retina with VEGF-A or BK induced an immediate, marked loss of sulforhodamine B from the microvessel lumen, which was similar for both stimuli and amounted to a ca. 3-fold increase in microvessel permeability (Figures 2c-f). Preincubation of the ex-vivo retina with VEGFR2 inhibitor SU-1498 for 15 min prevented VEGF-A- but not BK-induced permeability, confirming the role of VEGFR2 in VEGF-A-induced permeability and indicating that BK acted through a different receptor. Whole mount retinal staining showed that VEGF-A- and BK-induced

permeability coincided with the phosphorylation of p38 on T180/Y182, its downstream

effector HSP27 (on S82), as well as eNOS (on S1177) (Figure 2g, h). Phosphorylation

of all three downstream effectors in response to VEGF-A but not BK was abolished

following preincubation with SU-1498.

This experimental model was also used for mouse retinae. Baseline permeability in

mouse preparations was $0.15 \pm 0.1 \times 10^{-6}$ cm/s. VEGF-A stimulation increased

permeability to $0.65 \pm 0.2 \times 10^{-6}$ cm/s, and this was again sensitive to SU-1498 (Figure

2i). Furthermore, we observed SU-1498-sensitive phosphorylation of p38, HSP27 and

eNOS in mouse retinal microvessels within 2 min of VEGF-A stimulation (Figure 2j).

To assess the compatibility of the ex vivo retina with knockdown technology, mouse

eyes were injected intravitreously with siRNA against CLDN5. Western blot analysis

of retinal lysates, harvested 72 h after the injection, showed that CLDN5 expression

was significantly reduced by 65% (Supplemental Figure 1c, d) and this was

corroborated by whole mount staining of the retina (Supplemental Figure 1e).

Microvessel permeability following knock-down of CLDN5 increased ca. 3-fold (to

0.41 ± 0.03 cm/s) (Supplemental Figure 1f, g). Taken together, these results

demonstrated that the ex-vivo retinal preparation was a reliable model to measure

retinal paracellular microvessel permeability in rats and mice.

VEGF-A and Bradykinin induce AMPK phosphorylation

In order to find new regulators of permeability, primary rat brain microvascular ECs

were stimulated for 5 or 30 min with VEGF-A (50 ng/ml) from either the apical (non-

permeability inducing) or basal (permeability-inducing) side (Figure 3a) and cell

lysates analysed by a phospho-protein antibody array. In response to VEGF-A

stimulation many signalling components were phosphorylated, as exemplified by p38,

HSP27, AMPK, eNOS, SRC, ERK and AKT (Figure 3b, c).

Differentially phosphorylated proteins were categorised into three groups; group I

phosphorylated only after basal stimulation with VEGF-A (such as p38, HSP27 and

AMPK), group II phosphorylated regardless of the side of the stimulation (such as

eNOS, SRC and ERK) and group III phosphorylated only when VEGF-A was applied

apically (such as AKT). Phosphorylation of proteins exclusively in response to basally

applied VEGF-A suggested they played a role in hyperpermeability (group I). Among

these, AMPK stood out as its role in acute endothelial hyperpermeability has not yet

been studied in detail. Additionally, AMPK was also among the 10 proteins, for which

phosphorylation increased the most in response to basally applied VEGF-A, when

analysed by phospho-peptide mass spectrometry (Dragoni and Turowski, unpublished

results). A previous study also demonstrates that AMPK link Ca^{2+} transients to VE-cad

phosphorylation in response to ICAM-1 engagement in brain microvascular ECs[26].

AMPK phosphorylation in response to VEGF-A was confirmed by Western blot

analysis. Stimulation of primary rat brain ECs from the basal side led to rapid, transient

phosphorylation of AMPKα on T172, which peaked after ca. 2 min (Figure 3d). A very

similar activation pattern was observed in the intact rat retina when VEGF-A was

applied directly to the top of the isolated retina (corresponding to the basal side of the

endothelium) (Figure 3e). BK induced similar phosphorylation of AMPKα T172, both

in the cultured primary brain microvascular ECs and intact retina (Figure 3f, g). It was

notable that AMPK phosphorylation was maximal after around 2 min in response to

both VEGF-A and BK. However, BK clearly induced more sustained phosphorylation.

AMPK mediates VEGF-A/Bradykinin-induced permeability

In order to specify the role of AMPK during VEGF-A- and BK-induced vascular permeability, AMPK activity was neutralised in the ex vivo retina. Preincubation of the ex-vivo retina with compound C, a widely used AMPK antagonist, significantly decreased VEGF-A- and BK- induced permeability by 80% and 93%, respectively (Figure 4a-d). Moreover, whole mount staining showed that pre-treatment with compound C prevented the VEGF-A- and BK-induced activation of p38, HSP27 and eNOS (Figure 4e, f).

Next AMPKα was knocked down specifically by intravitreous injection of siRNA 72 h prior to preparing the retinae for ex vivo permeability measurements. Western blot results showed that AMPKα protein expression was significantly reduced by 70% (Figures 4g, h). Neither VEGF-A nor BK stimulation led to significant permeability in AMPK knocked-down retinae (Figures 4i-l).

Finally, when AMPK was directly activated by treating ex-vivo retinae with A769662 or AICAR, specific AMPK agonists that directly bind to and activate AMPK without any significant change in cellular ATP, ADP or AMP levels, baseline permeability was induced (Figure 5a-d) as well as the phosphorylation of AMPK, p38, HSP27 and eNOS (Figure 5e). Taken together these results confirmed the central role of AMPK in VEGF-A- and BK-induced acute retinal leakage and showed that AMPK acted upstream of p38 and eNOS.

VEGF-A- and BK-induced permeability requires Ca^{2+} and CAMKK, p38 and eNOS

Next, we aimed at placing AMPK within established PIF signalling cascades. Ca^{2+} is critical for the activation of both p38 and eNOS[26, 27,28]. To test its role in VEGF-A-induced vascular leakage the ex-vivo retina was incubated with BAPTA, a

Ca^{2+}chelant, prior to VEGF-A administration. BAPTA treatment significantly reduced

VEGF-A-induced permeability by 94%, and also prevented the phosphorylation of p38,

Hsp27 and eNOS (Figures 6a, e), suggesting that Ca^{2+} acted upstream to these

molecules. CAMKK is able to phosphorylate and activate AMPK in a Ca^{2+}-dependent

manner[26, 29]. Treatment of retinae with the CAMKK inhibitor STO-609 significantly

reduced VEGF-A-induced permeability by 86% and also prevented the activation of

p38, Hsp27 and eNOS (Figure 6b, e). The role of eNOS in permeability was re-

confirmed by preincubation of the ex-vivo retinae with L-NAME, which reduced VEGF-

A-induced permeability by 87% but it did not have any effect on p38 or HSP27

phosphorylation (Figure 6c, e), indicating that eNOS was not upstream of p38. Finally,

the ex-vivo retina was preincubated with the p38 inhibitor SB203580, which

significantly reduced the VEGF-A-induced permeability by 80% as well as activation

of Hsp27 but not of eNOS (Figure 6d, e).

Similar results were obtained for BK-induced stimulation of retinae. BAPTA or STO

nearly completely abolished BK-induced permeability, together with the activation of

p38, Hsp27 and eNOS (Figure 6f, g, j). Pre-treatment of the ex-vivo retina with L-

NAME reduced BK-induced permeability by 67% but did not affect p38 or HSP27

phosphorylation (Figure 6h, j). Finally, SB203580 prevented BK-induced permeability

as well as phosphorylation of HSP27 but not eNOS (Figure 6i, j).

Importantly, VEGF-A- or BK-induced phosphorylation of AMPKα was completely

abolished by BAPTA or STO. VEGF-A- but not BK-induced AMPKα phosphorylation

was also abolished by SU1498 (Figure 7a). These results indicated that in the retina,

VEGF-A and BK induced Ca^{2+} transients and consequent activation of CAMKK and

AMPK. At this point signalling diverged into either activation of p38 or eNOS, which

both contributed to permeability (Figure 7c).

VEGF-A and BK stimulate VE-Cadherin phosphorylation

Lastly, we wanted to analyse VE-cad internalisation in the VEGF-A and BK stimulated ex vivo retina. However, systematic correlation of cryo-immuno EM to leaky microvessels proved impractical. We therefore used phosphorylation of VE-cad on Y685 as a surrogate marker for retinal microvessel permeability[16], in particular since such VE-cad phosphorylation is also a prerequisite for internalisation after BK treatment[15]. VEGF-A or BK stimulation of ex vivo retinae induced tyrosine phosphorylation of VE-cad on Y685 in intact microvessels (Figure 7b). This phosphorylation was completely abolished following pre-incubation with compound C, L-NAME but not SB203580, indicating that VE-cad was downstream of AMPK/eNOS but not AMPK/p38 (Figure 7c).

DISCUSSION

Measuring retinal microvascular permeability has been mainly restricted to Miles-type assays using either fluorescent tracers or Evans Blue/albumin[8, 30]. Chronic leakage can also be visualised in real time by fluorescein angiography[31]. However, collectively these methods, which measure the amount of extravasated tracer, do not only reflect the degree of leakage but are also strongly influenced by dye concentration in the vasculature and dye clearance from the tissue[1, 2]. Further disadvantages of these methods are that compound concentrations and timing cannot be controlled accurately. Thus, they are inadequate to measure acute permeability and associated signalling accurately and in a controlled manner.

The ex-vivo retinal platform described herein addressed most of these issues. It constituted a significant advance to EC cultures, since it used a complete and intact

neurovascular unit. The functionality of the retinal vasculature was preserved, with

both VE-cad and claudin-5 distribution indistinguishable from that in vivo. Similar

preparations of the brain and the retina also display full cellular functionality of e.g.

pericytes[32]. Importantly, permeability to sulforhodamine-B was very low and within the

range of that of other small non-ionic molecules at the intact BBB in vivo[33] and notably

ca. 10 x lower than in preparations of pial microvessels[34], indicating that this model

was highly suitable for permeability measurement at an intact neurovascular unit.

Vascular barrier properties in these retinal explants was, as expected, dependent on

tight junction integrity. The reported speed, at which compounds such as VEGF-A

induce permeability[10], was fully recapitulated. Leakage measurements were then

combined with whole tissue staining and analyses of the phosphorylation status of key

mediators of permeability using phospho-specific antibodies to gain mechanistic

insight. Furthermore, the preparations could be interrogated using small molecule

antagonists and agonists at defined concentrations and times, and allowed for the

identification of key downstream regulators, common to both VEGF-A and BK

stimulation. Even more specific neutralisation of key proteins was achieved through

prior intraocular injection of siRNA. Conceivably this model is compatible for use with

genetically modified mice and disease models, further broadening its applicability. The

ex vivo platform could also be used to measure Ca^{2+} transients or localised production

of reactive oxygen or nitrogen species in response to vasoactive compounds such as

VEGF-A or BK.

Measurements in retinae were done ex vivo, in the absence of blood flow. Blood flow

and associated shear stress may influence EC biology, such as cell-cell adhesion and

inflammatory dysfunction[15, 35, 36] and their absence in our preparation must be taken

into account when evaluating results. However, it should also be noted that

permeability regulation by shear stress appears to be remembered in ECs in vitro for

at least 24 h[37].

For the purpose of developing and validating the ex vivo retinal platform we have

focused on PIFs, which induce permeability when added to the abluminal (tissue) side

of the endothelium, such as VEGF-A and BK[12], as these are readily applied on top of

the retinal explants. Other PIFs that act only from the luminal side, such as

lysophosphatidic acid[12] or lysophosphatidylcholine[8], could also potentially be

investigated in this system. However, this would require the use of a manifold injection

system[38], which allows switching between injection of sulforhodamine-B with or

without permeability factor into radial vessels.

VEGF-A and BK induced acute leakage in retinal microvessels, which was associated

with and dependent on Ca^{2+}, the MAPK p38 and eNOS, in agreement with published

data[11, 12, 14]. We also identified AMPK as a novel key mediator of both VEGF-A- and

BK-induced permeability, indicating it is a core regulator of acute vascular

permeability. AMPK is primarily known to regulate energy requirements of the cell,

but has also been implicated in other seemingly unrelated cellular processes such as

migration, cell growth and apoptosis[39]. This protein kinase has been studied before in

relation to its protective role of the BBB[19, 20], whereas in the retinal pigment epithelium

it has been shown to be responsible for the permeability induced by IL-1β[21]. However,

all these studies address chronic changes and do not focus on the role of AMPK for

acute permeability. Whilst its canonical activation is dependent on cellular AMP: ATP

concentrations and phosphorylation on T172 by LKB1, we found that, in the regulation

of endothelial permeability, AMPK was activated downstream of Ca^{2+} and CAMKK.

This activation pathway has previously been described as non-canonical[40] and is also

operational when CNS ECs facilitate the transmigration of lymphocytes[26]. Notably,

VEGF-A has also been reported before to induce NO production via a pathway

requiring Ca^{2+} and AMPK[41]. Indeed, eNOS phosphorylation on S1177 can be

mediated by AKT[42] or AMPK[26, 43]. However, the PI3K/Akt pathway is not relevant to

VEGF-A-induced permeability induction in neurovascular ECs[12]. We confirmed the

phosphorylation of eNOS on S1177 downstream of AMPK both during VEGF-A and

BK permeability induction. Interestingly, the Ca^{2+}/AMPK/eNOS pathway resulted in the

phosphorylation of VE-cad on Y685, identified previously as key for vascular

permeability in the periphery[15] and the retina[16], thus providing a direct link between

AMPK and paracellular junction regulation. For retinal permeability, occludin

phosphorylation and internalisation also plays an important role. However, judging

from the published time courses, it is likely to be a later event, not captured by our

experiments, and thus either secondary to VE-cad phosphorylation or, with its

dependency on PKCβ activation, outside of the signalling network we have

investigated here[17].

In response to VEGF-A and BK, AMPK also regulated the phosphorylation of the

MAPK p38 and its substrate HSP27, both previously implicated in actin rearrangement

during endothelial barrier disruption[44, 45]. P38 is a bona fide regulator of VEGF-A

responses[46] and its activation downstream of cdc42 and PAK and subsequent

modulation of the actin cytoskeleton occurs during VEGF-A-induced endothelial

migration downstream of VEGFR2 phosphorylation on Y1214[47]. Whilst it is possible

that additional, parallel AMPK regulation of this cascade was operational during

VEGF-A or BK-induced permeability, we favour an alternative model of direct

activation of p38 by AMPK via TAB1, a pathway described in apoptotic lymphocytes

and the ischemic heart[48, 49]. By switching between two different p38 activation modes

(Ca^{2+}/AMPK/TAB1 versus cdc42/PAK) ECs could adapt cytoskeletal regulation to the

specific requirement of EC migration or permeability.

The ex-vivo retina proved to be a reliable model and demonstrated its usefulness in

identifying key regulators of acute permeability. Whilst AMPK clearly emerged as such

a key regulator, it is unlikely to be exploitable as a target for anti-leakage treatments:

its central role in regulating cellular energy demands throughout the body hints at

many potential side effects. Activation of AMPK is currently investigated as a

therapeutic option to treat cancer, metabolic syndrome and diabetes[50, 51]. However, in

light of the strong induction of permeability we observed in response to at least two

AMPK agonists, we propose that these avenues should be explored cautiously, since

at least acute microvascular leakage may accompany such treatment modalities.

Nevertheless, our data collectively indicated that the ex vivo retina platform can play

an important part in elucidating mechanisms and signalling of neurovascular leakage.

ACKNOWLEDGEMENTS

We thank Dr Paul Fraser (King's College London) for introducing us to his early ex-

vivo retinal preparations and Prof. John Greenwood (UCL) for critical comments on

the manuscript. siRNAs targeting CLDN5 were kindly provided by Dr Matthew

Campbell (Trinity College Dublin). Phospho-VE-Cad antibodies were kindly provided

by Dr Fabrizio Orsenigo and Prof. Elisabetta Dejana (FIRC Institute of Molecular

Oncology, Milan and Uppsala University).

This work was supported by a Moorfields Eye Charity grant to S.D. and a Diabetes

UK grant to P.T.

The data that support the findings of this study are available from the corresponding

author upon reasonable request.

Supplemental material for this paper can be found at the journal website:

https://eur01.safelinks.protection.outlook.com/?url=http%3A%2F%2Fjournals.sagepub.com%

2Fhome%2Fjcb&data=02%7C01%7C%7Cc27df537f41f419d715808d795e9c351%7C1faf

88fea9984c5b93c9210a11d9a5c2%7C0%7C0%7C637142703553285613&sdata=J8jD0ho

MtCdGE1jbrGR5QDN1QGWyjbw0hnEBOenTqj0%3D&reserved=0

AUTHOR CONTRIBUTION STATEMENT

S.D.: designed experiments; performed experiments; analysed the data; wrote the

manuscript

B.C.; E.K.; T.B; M.H.S.: performed experiments

P.T.: designed experiments; performed experiments; analysed the data; wrote the

manuscript

DISCLOSURE/CONFLICT OF INTEREST

None.

REFERENCES

1.   Nagy JA, Benjamin L, Zeng H, et al. Vascular permeability, vascular hyperpermeability and
angiogenesis. *Angiogenesis* 2008; 11: 109-119. 2008/02/23. DOI: 10.1007/s10456-008-9099-z.
2.   Turowski P. Leakage at Blood-Neural Barriers. In: Link S (ed) *The Blood Brain Barrier and*
Inflammation. 2017, pp.81-102.
3.   Sweeney MD, Zhao Z, Montagne A, et al. Blood-Brain Barrier: From Physiology to Disease and
Back. *Physiol Rev* 2019; 99: 21-78. 2018/10/04. DOI: 10.1152/physrev.00050.2017.

4.   Klaassen I, Van Noorden CJ and Schlingemann RO. Molecular basis of the inner blood-retinal
barrier and its breakdown in diabetic macular edema and other pathological conditions. *Prog Retin*
Eye Res 2013; 34: 19-48. 2013/02/19. DOI: 10.1016/j.preteyeres.2013.02.001.
5.   Brown DM, Nguyen QD, Marcus DM, et al. Long-term outcomes of ranibizumab therapy for
diabetic macular edema: the 36-month results from two phase III trials: RISE and RIDE. *Ophthalmology*
2013; 120: 2013-2022. 2013/05/28. DOI: 10.1016/j.ophtha.2013.02.034.
6.   Ford JA, Lois N, Royle P, et al. Current treatments in diabetic macular oedema: systematic
review and meta-analysis. *BMJ Open* 2013; 3 2013/03/05. DOI: 10.1136/bmjopen-2012-002269.
7.   Campochiaro PA and Peters KG. Targeting Tie2 for Treatment of Diabetic Retinopathy and
Diabetic Macular Edema. *Curr Diab Rep* 2016; 16: 126. 2016/10/26. DOI: 10.1007/s11892-016-0816-
5.
8.   Canning P, Kenny BA, Prise V, et al. Lipoprotein-associated phospholipase A2 (Lp-PLA2) as a
therapeutic target to prevent retinal vasopermeability during diabetes. *Proc Natl Acad Sci U S A* 2016;
113: 7213-7218. 2016/06/15. DOI: 10.1073/pnas.1514213113.
9.   Kita T, Clermont AC, Murugesan N, et al. Plasma Kallikrein-Kinin System as a VEGF-
Independent Mediator of Diabetic Macular Edema. *Diabetes* 2015; 64: 3588-3599. 2015/05/17. DOI:
10.2337/db15-0317.
10.   Bates DO. Vascular endothelial growth factors and vascular permeability. *Cardiovasc Res*
2010; 87: 262-271. 2010/04/20. DOI: 10.1093/cvr/cvq105.
11.   Wu HM, Yuan Y, Zawieja DC, et al. Role of phospholipase C, protein kinase C, and calcium in
VEGF-induced venular hyperpermeability. *Am J Physiol* 1999; 276: H535-542. 1999/02/10. DOI:
10.1152/ajpheart.1999.276.2.H535.
12.   Hudson N, Powner MB, Sarker MH, et al. Differential apicobasal VEGF signaling at vascular
blood-neural barriers. *Dev Cell* 2014; 30: 541-552. 2014/09/02. DOI: 10.1016/j.devcel.2014.06.027.
13.   Weidert E, Pohler SE, Gomez EW, et al. Actinomyosin contraction, phosphorylation of VE-
cadherin, and actin remodeling enable melanoma-induced endothelial cell-cell junction disassembly.
PLoS One 2014; 9: e108092. 2014/09/17. DOI: 10.1371/journal.pone.0108092.
14.   Di Lorenzo A, Lin MI, Murata T, et al. eNOS-derived nitric oxide regulates endothelial barrier
function through VE-cadherin and Rho GTPases. *J Cell Sci* 2013; 126: 5541-5552. 2013/09/21. DOI:
10.1242/jcs.115972.
15.   Orsenigo F, Giampietro C, Ferrari A, et al. Phosphorylation of VE-cadherin is modulated by
haemodynamic forces and contributes to the regulation of vascular permeability in vivo. *Nat Commun*
2012; 3: 1208. 2012/11/22. DOI: 10.1038/ncomms2199.
16.   Smith RO, Ninchoji T, Gordon E, et al. Vascular permeability in retinopathy is regulated by
VEGFR2 Y949 signaling to VE-cadherin. *Elife* 2020; 9 2020/04/22. DOI: 10.7554/eLife.54056.
17.   Murakami T, Frey T, Lin C, et al. Protein kinase cbeta phosphorylates occludin regulating tight
junction trafficking in vascular endothelial growth factor-induced permeability in vivo. *Diabetes* 2012;
61: 1573-1583. 2012/03/23. DOI: 10.2337/db11-1367.
18.   Hardie DG. Keeping the home fires burning: AMP-activated protein kinase. *J R Soc Interface*
2018; 15 2018/01/19. DOI: 10.1098/rsif.2017.0774.
19.   Zhao Z, Hu J, Gao X, et al. Activation of AMPK attenuates lipopolysaccharide-impaired integrity
and function of blood-brain barrier in human brain microvascular endothelial cells. *Exp Mol Pathol*
2014; 97: 386-392. 2014/09/16. DOI: 10.1016/j.yexmp.2014.09.006.
20.   Takata F, Dohgu S, Matsumoto J, et al. Metformin induces up-regulation of blood-brain barrier
functions by activating AMP-activated protein kinase in rat brain microvascular endothelial cells.
Biochem Biophys Res Commun 2013; 433: 586-590. 2013/03/26. DOI: 10.1016/j.bbrc.2013.03.036.
21.   Villarroel M, Garcia-Ramirez M, Corraliza L, et al. Fenofibric acid prevents retinal pigment
epithelium disruption induced by interleukin-1beta by suppressing AMP-activated protein kinase
(AMPK) activation. *Diabetologia* 2011; 54: 1543-1553. 2011/03/04. DOI: 10.1007/s00125-011-2089-5.

22.   Warboys CM, Toh HB and Fraser PA. Role of NADPH oxidase in retinal microvascular
permeability increase by RAGE activation. *Invest Ophthalmol Vis Sci* 2009; 50: 1319-1328. 2008/11/11.
DOI: 10.1167/iovs.08-2730.
23.   Weksler B. B. SEA, Perrière N., Charneau P., Holloway K., Leveque M., Tricoire-Leignel H.,
Nicotra A., Bourdoulous S, Turowski P., Male D. k., Roux F., Greenwood J., Romero I. A., and Couraud
P. O. Blood-brain barrier-specific properties of a human adult brain endothelial cell line. *FASEB J* 2005;
19: 1872–1874.
24.   Turowski P, Adamson P, Sathia J, et al. Basement membrane-dependent modification of
phenotype and gene expression in human retinal pigment epithelial ARPE-19 cells. *Invest Ophthalmol
Vis Sci* 2004; 45: 2786-2794. 2004/07/28. DOI: 10.1167/iovs.03-0943.
25.   Martins T, Burgoyne T, Kenny BA, et al. Methamphetamine-induced nitric oxide promotes
vesicular transport in blood-brain barrier endothelial cells. *Neuropharmacology* 2013; 65: 74-82.
2012/09/11. DOI: 10.1016/j.neuropharm.2012.08.021.
26.   Martinelli R, Gegg M, Longbottom R, et al. ICAM-1-mediated endothelial nitric oxide synthase
activation via calcium and AMP-activated protein kinase is required for transendothelial lymphocyte
migration. *Mol Biol Cell* 2009; 20: 995-1005. 2008/12/17. DOI: 10.1091/mbc.E08-06-0636.
27.   Takeda K, Matsuzawa A, Nishitoh H, et al. Involvement of ASK1 in Ca2+-induced p38 MAP
kinase activation. *EMBO Rep* 2004; 5: 161-166. 2004/01/30. DOI: 10.1038/sj.embor.7400072.
28.   da Silva CG, Specht A, Wegiel B, et al. Mechanism of purinergic activation of endothelial nitric
oxide synthase in endothelial cells. *Circulation* 2009; 119: 871-879. 2009/02/04. DOI:
10.1161/CIRCULATIONAHA.108.764571.
29.   Hurley RL, Anderson KA, Franzone JM, et al. The Ca2+/calmodulin-dependent protein kinase
kinases are AMP-activated protein kinase kinases. *J Biol Chem* 2005; 280: 29060-29066. 2005/06/28.
DOI: 10.1074/jbc.M503824200.
30.   Radu M and Chernoff J. An in vivo assay to test blood vessel permeability. *J Vis Exp* 2013:
e50062. 2013/03/26. DOI: 10.3791/50062.
31.   Couturier A, Rey PA, Erginay A, et al. Widefield OCT-Angiography and Fluorescein Angiography
Assessments of Nonperfusion in Diabetic Retinopathy and Edema Treated with Anti-Vascular
Endothelial Growth Factor. *Ophthalmology* 2019 2019/08/07. DOI: 10.1016/j.ophtha.2019.06.022.
32.   Mishra A, O'Farrell FM, Reynell C, et al. Imaging pericytes and capillary diameter in brain slices
and isolated retinae. *Nat Protoc* 2014; 9: 323-336. 2014/01/18. DOI: 10.1038/nprot.2014.019.
33.   Shi L, Zeng M, Sun Y, et al. Quantification of blood-brain barrier solute permeability and brain
transport by multiphoton microscopy. *J Biomech Eng* 2014; 136: 031005. 2013/11/07. DOI:
10.1115/1.4025892.
34.   Easton AS, Sarker MH and Fraser PA. Two components of blood-brain barrier disruption in the
rat. *J Physiol* 1997; 503 (Pt 3): 613-623. 1997/10/23. DOI: 10.1111/j.1469-7793.1997.613bg.x.
35.   Chistiakov DA, Orekhov AN and Bobryshev YV. Effects of shear stress on endothelial cells: go
with the flow. *Acta Physiol (Oxf)* 2017; 219: 382-408. 2016/06/02. DOI: 10.1111/apha.12725.
36.   Ho RX, Tahboub R, Amraei R, et al. The cell adhesion molecule IGPR-1 is activated by, and
regulates responses of endothelial cells to shear stress. *J Biol Chem* 2019 2019/07/26. DOI:
10.1074/jbc.RA119.008548.
37.   Walsh TG, Murphy RP, Fitzpatrick P, et al. Stabilization of brain microvascular endothelial
barrier function by shear stress involves VE-cadherin signaling leading to modulation of pTyr-occludin
levels. *J Cell Physiol* 2011; 226: 3053-3063. 2011/02/09. DOI: 10.1002/jcp.22655.
38.   Liu B, Shao Y and Mirkin MV. Dual-pipet techniques for probing ionic reactions. *Anal Chem*
2000; 72: 510-519. 2000/03/01. DOI: 10.1021/ac990771p.
39.   Hardie DG. AMP-activated protein kinase: an energy sensor that regulates all aspects of cell
function. *Genes Dev* 2011; 25: 1895-1908. 2011/09/23. DOI: 10.1101/gad.17420111.
40.   Carling D, Sanders MJ and Woods A. The regulation of AMP-activated protein kinase by
upstream kinases. *Int J Obes (Lond)* 2008; 32 Suppl 4: S55-59. 2008/09/18. DOI: 10.1038/ijo.2008.124.

41.   Reihill JA, Ewart MA, Hardie DG, et al. AMP-activated protein kinase mediates VEGF-
stimulated endothelial NO production. *Biochem Biophys Res Commun* 2007; 354: 1084-1088.
2007/02/06. DOI: 10.1016/j.bbrc.2007.01.110.
42.   Fulton D, Gratton JP, McCabe TJ, et al. Regulation of endothelium-derived nitric oxide
production by the protein kinase Akt. *Nature* 1999; 399: 597-601. 1999/06/22. DOI: 10.1038/21218.
43.   Youn JY, Wang T and Cai H. An ezrin/calpain/PI3K/AMPK/eNOSs1179 signaling cascade
mediating VEGF-dependent endothelial nitric oxide production. *Circ Res* 2009; 104: 50-59.
2008/11/29. DOI: 10.1161/CIRCRESAHA.108.178467.
44.   Sawada J, Li F and Komatsu M. R-Ras Inhibits VEGF-Induced p38MAPK Activation and HSP27
Phosphorylation in Endothelial Cells. *J Vasc Res* 2015; 52: 347-359. 2016/03/31. DOI:
10.1159/000444526.
45.   Kasa A, Csortos C and Verin AD. Cytoskeletal mechanisms regulating vascular endothelial
barrier function in response to acute lung injury. *Tissue Barriers* 2015; 3: e974448. 2015/04/04. DOI:
10.4161/21688370.2014.974448.
46.   Corre I, Paris F and Huot J. The p38 pathway, a major pleiotropic cascade that transduces
stress and metastatic signals in endothelial cells. *Oncotarget* 2017; 8: 55684-55714. 2017/09/15. DOI:
10.18632/oncotarget.18264.
47.   Lamalice L, Houle F, Jourdan G, et al. Phosphorylation of tyrosine 1214 on VEGFR2 is required
for VEGF-induced activation of Cdc42 upstream of SAPK2/p38. *Oncogene* 2004; 23: 434-445.
2004/01/16. DOI: 10.1038/sj.onc.1207034.
48.   Li J, Miller EJ, Ninomiya-Tsuji J, et al. AMP-activated protein kinase activates p38 mitogen-
activated protein kinase by increasing recruitment of p38 MAPK to TAB1 in the ischemic heart. *Circ*
Res 2005; 97: 872-879. 2005/09/24. DOI: 10.1161/01.RES.0000187458.77026.10.
49.   Lanna A, Henson SM, Escors D, et al. The kinase p38 activated by the metabolic regulator
AMPK and scaffold TAB1 drives the senescence of human T cells. *Nat Immunol* 2014; 15: 965-972.
2014/08/26. DOI: 10.1038/ni.2981.
50.   Hardie DG. Targeting an energy sensor to treat diabetes. *Science* 2017; 357: 455-456.
2017/08/05. DOI: 10.1126/science.aao1913.
51.   Kim J, Yang G, Kim Y, et al. AMPK activators: mechanisms of action and physiological activities.
Exp Mol Med 2016; 48: e224. 2016/04/02. DOI: 10.1038/emm.2016.16.

FIGURE LEGENDS

Figure 1. VEGF-A and BK induced permeability in cultured brain microvascular

ECs.

(a, b) Primary rat brain microvascular ECs were grown on permeable Transwell inserts

to confluence and until they reached full electrical barrier (500-800 Ω.cm2). VEGF-A

(a) or BK (b) were added at time 0. Shown are means ± SD of normalized resistance

changes (n=3). Significant changes were detected in the short- and long-term, as well

as overall responses. **(c)** Changes in the distribution of VE-Cad and Occludin in response to basal (corresponding to abluminal) stimulation with VEGF-A or BK were analysed by confocal microscopy in post-confluent primary rat brain microvascular ECs. Whites arrows indicate the broadening of the VE-Cad staining. Scale bar, 10 µM. **(d-f)** Cryo-immuno-EM of VE-Cad distribution in control (d) and VEGF-A (e) or BK (f) stimulated human hCMEC/D3 cells. Shown are interendothelial junction areas with the two adjacent membranes (red arrowheads). White arrowheads point out gold labelled VE-Cad, which in control cells was found predominantly associated with abutting plasma membranes (within 20 nm; i.e. the distance expected by the primary and the secondary bridging Ab). Scale bar, 100 nm. **(g)** Distances of VE-Cad gold particles from cell-cell junctions determined from three independent preparations as shown in (d-f). **p < 0.01, ***p < 0.001.

Figure 2. Validation of the ex vivo retina model in rats.

(a, b) Control retinae were from animals directly perfused fixed with 4% PFA. Ex vivo retinae were isolated as described, flat mounted and left submerged with Krebs solutions for 1 h before PFA fixation. Whole mounts were stained for Isolectin B4 (IB4), claudin-5 (CLDN5) and VE-cad as indicated. **(c-f)** Sulforhodamine-B fluorescent intensities were recorded in single retinal capillaries. 50 ng/ml VEGF-A (c, d) or 10 µM Bradykinin (BK) (e, f) were applied on top of the retina (abluminal side) at times indicated. Optionally retinae were preincubated with the VEGFR2-selective antagonist SU1498 (10 µM) for 15 min prior to recording. Mean (± SD) permeability changes recorded from at least four retinae are shown in (d) and (f). **(g, h)** Ex vivo retinal explants were incubated with VEGF-A (g) or BK (h) for 2 min, fixed using 4% PFA and then stained using IB4 (green) and for phospho-p38 (pT180/Y182), phospho-Hsp27

(pS82) or phospho-eNOS (pS1177) (magenta). **(i)** Microvessel permeability changes

were recorded in mouse retinae as described in (e, f). **(j)** Mouse retinal explants were

stimulated and stained as described in (g) using phospho-specific antibodies to p38,

HSP27 and eNOS. ns non-significant, ***$p < 0.001$. Scale bars, 10 μm.

Figure 3. VEGF-A and Bradykinin induced AMPK phosphorylation.

(a-c) Primary brain microvascular ECs were stimulated with VEGF-A from the apical

or basal side for 5 min triggering apically (group I), basally specific (group III) or mixed

(group II) responses (see text or ref 12 for more details) (a). Cells were lysed and

phosphorylation of indicated molecules assessed by phosphoantibody array analysis

(b-c). **(d-g)** brain microvascular ECs (d, f) or ex-vivo rat retinae (e, g) were stimulated

with 50 ng/ml VEGF-A (d, e) or 10 μM BK (f, g) for the indicated length of time and

AMPKα phosphorylation (pT172) analysed. Representative results and quantification

of AMPKα activation from three independent experiments are shown as normalised

means ± SD. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 4. AMPK mediated VEGF-A/Bradykinin-induced permeability.

(a-f) Rat ex-vivo retinae were preincubated with or without Compound-C (CC, 10 μM)

for 15 minutes. 10 ng/ml VEGF-A (a, b) or 10 μM BK (c, d) were then applied to the

top of the retina and changes in permeability were recorded. Alternatively (e, f), retinae

were immunostained using IB4 (green) and anti-phospho-p38, -HSP27 or -eNOS

(magenta) as detailed in Figure 2. **(g-l)** AMPKα specific siRNA or scrambled control

was injected into the vitreous of mouse eyes. After 48 h retinae were isolated, lysed

and subjected to immunoblotting as indicated (g). Shown in (h) is the densitometric

quantification of 3 independent experiments as shown in (g). Alternatively, after 48 h

retinae were prepared for ex vivo permeability measurements and stimulated using

VEGF-A (50 ng/ml) (i, j) or BK (10 µM) (k, l). Note that neither PIF induced any

permeability in the knocked down ex-vivo retina. Representative results and

quantifications (normalized mean ± SD) from three independent experiments are

shown. **p < 0.01, ***p < 0.001. Scale bars, 10 µm.

Figure 5. Stimulation of AMPK induced permeability in the ex vivo retina.

(a-d) Ex vivo preparations were stimulated with the two different AMPK activators

A769662 (10 µM) and AICAR (10 µM). Both agonists induced strong and immediate

permeability in the ex vivo retinal microvessels. Mean (± SD) permeability changes

recorded from at least four retinae are shown in (b, d). **(e)** Ex vivo retinae were

stimulated as in (a-d) and after 2 min fixed using 4% PFA and then immunostained

using IB4 (green) and or anti-phospho-p38, -HSP27 or -eNOS (magenta) as detailed

in Figure 2 (magenta). *p < 0.05. Scale bars, 10 µm.

Figure 6. VEGF-A- and Bradykinin-induced permeability requires Ca2+, CaMKK,

p38 and eNOS.

(a-e) Ex vivo retinae were preincubated with 20 µM BAPTA (Ca^{2+} chelator) (a, e as

indicated), 10 µM STO-609 (STO, CaMKK inhibitor) (b, e as indicated), 10 µM L-NAME

(NOS inhibitor) (c, e as indicated), or 10 µM SB202190 (SB, p38 inhibitor) (d, e as

indicated) for 15 minutes. Then VEGF-A (50 ng/ml) was applied to the top of the retina

(abluminal side) and changes in microvessel permeability were recorded as described

in Figure 2. Alternatively (e), retinae were fixed after 2 min using 4 % PFA and then

immunostained with IB4 (green) and for phosphorylation of p38, HSP27 and eNOS

(magenta). **(f-j)** As in panels (a-e), except that ex vivo retinae were stimulated with BK

(10 µM). Representative results and quantifications (mean ± SD) from four

independent experiments are shown. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. Scale bars,

10 µm.

Figure 7. VEGF-A and Bradykinin induced VE-Cadherin phosphorylation.

(a) Ex vivo retinae were preincubated with SU1498 (10 µM), or BAPTA (20 µM) or

STO (10 µM) and treated with VEGF-A and Bradykinin for 2 min. Retinae were then

fixed with 4% PFA and immunostained for phospho-AMPKα (magenta) and with IB4

(green). **(b)** Ex vivo retinae were preincubated with compound C (10 µM), or L-NAME

(10 µM) or SB202190 (10 µM), treated with VEGF-A and Bradykinin for 2 min and

immunostained for phosphorylated VE-cad (using anti-pY685-VEC) (magenta) and

with IB4 (green). **(c)** Proposed signalling networks in the ex-vivo retinal

microvasculature downstream of VEGF-A and Bradykinin. Scale bars, 10 µm.

Supplemental Figure 1. Properties of the ex vivo retina model.

(a-b) Selected frames, pseudocoloured in (b), of time-resolved recordings of a

Sulforhodamine-B-filled rat capillary before and after the addition of 50 ng/ml VEGF-

A, illustrating the rapid loss of fluorophore from the lumen of the vessel. The red box

exemplifies a typical r.o.i used for intensity measurement. Scale bars, 10 µm. **(c-g)**

CLDN5 siRNA was injected into mouse eyes. 48 h later CLDN5 levels were analysed

by immunoblots of retinal lysates (c-d) or by wholemount immunochemical staining

(e). Scale bar, 100 µm. Alternatively, permeability of 4 KDa Rhodamine was measured

in ex vivo retinae from CLDN5 siRNA or control injected eyes (f-g). **$p < 0.01$, ***$p <$

0.001.

Figure 1. VEGF and BK induced permeability in cultured brain MVECs.
(a, b) Primary rat brain microvascular ECs were grown on permeable Transwell inserts to confluence and until they reached full electrical barrier (500-800 Ω.cm2). VEGF-A (a) or BK (b) were added at time 0. Shown are means ± SD of normalized resistance changes (n=3). Significant changes were detected in the short- and long-term, as well as overall responses. **(c)** Changes in the distribution of VE-Cad and Occludin in response to basal (corresponding to abluminal) stimulation with VEGF-A or BK were analysed by confocal microscopy in post-confluent primary rat brain microvascular ECs. Whites arrows indicate the broadening of the VE-Cad staining. Scale bar: 10 µM. **(d-f)** Cryo-immuno-EM of VE-Cad distribution in control (d) and VEGF-A (e) or BK (f) stimulated human hCMEC/D3 cells. Shown are interendothelial junction areas with the two adjacent membranes (red arrowheads). White arrowheads point out gold labelled VE-Cad, which in control cells was found predominantly associated with abutting plasma membranes (within 20 nm; i.e. the distance expected by the primary and the secondary bridging Ab). Scale bar: 100 nm. **(g)** Distances of VE-Cad gold particles from cell-cell junctions for determined from three independent preparations as shown in (d-f). **p < 0.01, ***p < 0.001.

Figure 2. Validation of the ex vivo retina model in rats.
(a, b) Control retinae were from animals directly perfused fixed with 4% PFA. Ex vivo retinae were isolated as described, flat mounted and left submerged with Krebs solutions for 1 h before PFA fixation. Whole mounts were stained for Isolectin B4 (IB4), claudin-5 (CLDN5) and VE-cad as indicated. **(c-f)** Sulforhodamine-B fluorescent intensities were recorded in single retinal capillaries. 50 ng/ml VEGF-A (c, d) or 10 μM Bradykinin (BK) (e, f) were applied on top of the retina (abluminal side) at times indicated. Optionally retinae were preincubated with the VEGFR2-selective antagonist SU1498 (10 μM) for 15 min prior to recording. Mean (± SD) permeability changes recorded from at least four retinae are shown in (d) and (f). **(g, h)** Ex vivo retinal explants were incubated with VEGF-A (g) or BK (h) for 2 min, fixed using 4% PFA and then stained using IB4 (green) and for phospho-p38 (pT180/Y182), phospho-Hsp27 (pS82) or phospho-eNOS (pS1177) (magenta). **(i)** Microvessel permeability changes were recorded in mouse retinae as described in (e, f). **(j)** Mouse retinal explants were stimulated and stained as described in (g) using phospho-specific antibodies to p38, HSP27 and eNOS. ns non-significant, ***p < 0.001. Scale bars, 10 μm.

Figure 3. VEGF-A and Bradykinin induced AMPK phosphorylation.
(a-c) Primary brain microvascular ECs were stimulated with VEGF-A from the apical or basal side for 5 min triggering apically (group I), basally specific (group III) or mixed (group II) responses (see text or ref 12 for more details) (a). Cells were lysed and phosphorylation of indicated molecules assessed by phosphoantibody array analysis (b-c). **(d-g)** brain microvascular ECs (d, f) or ex-vivo rat retinae (e, g) were stimulated with 50 ng/ml VEGF-A (d, e) or 10 μM BK (f, g) for the indicated length of time and AMPKα phosphorylation (pT172) analysed. Representative results and quantification of AMPKα activation from three independent experiments are shown as normalised means ± SD. *p < 0.05, **p < 0.01, ***p < 0.001.

Figure 4. AMPK mediated VEGF-A/Bradykinin-induced permeability.
(a-f) Rat ex-vivo retinae were preincubated with or without Compound-C (CC, 10 µM) for 15 minutes. 10 ng/ml VEGF-A (a, b) or 10 µM BK (c, d) were then applied to the top of the retina and changes in permeability were recorded. Alternatively (e, f), retinae were immunostained using IB4 (green) and anti-phospho-p38, -HSP27 or -eNOS (magenta) as detailed in Figure 2. **(g-l)** AMPKα specific siRNA or scrambled control was injected into the vitreous of mouse eyes. After 48 h retinae were isolated, lysed and subjected to immunoblotting as indicated (g). Shown in (h) is the densitometric quantification of 3 independent experiments as shown in (g). Alternatively, after 48 h retinae were prepared for ex vivo permeability measurements and stimulated using VEGF-A (50 ng/ml) (i, j) or BK (10 µM) (k, l). Note that neither PIF induced any permeability in the knocked down ex-vivo retina. Representative results and quantifications (normalized mean ± SD) from three independent experiments are shown. **p < 0.01, ***p < 0.001. Scale bars, 10 µm.

Figure 5. Stimulation of AMPK induced permeability in the ex-vivo retina.
(a-d) Ex vivo preparations were stimulated with the two different AMPK activators A769662 (10 µM) and AICAR (10 µM). Both agonists induced strong and immediate permeability in the ex vivo retinal microvessels. Mean (± SD) permeability changes recorded from at least four retinae are shown in (b, d). **(e)** Ex vivo retinae were stimulated as in (a-d) and after 2 min fixed using 4% PFA and then immunostained using IB4 (green) and or anti-phospho-p38, -HSP27 or -eNOS (magenta) as detailed in Figure 2 (magenta). *p < 0.05. Scale bars, 10 µm.

Figure 6. VEGF-A- and Bradykinin-induced permeability requires Ca^{2+}, CaMKK, p38 and eNOS.
(a-e) Ex vivo retinae were preincubated with 20 µM BAPTA (Ca^{2+} chelator) (a, e as indicated), 10 µM STO-609 (STO, CaMKK inhibitor) (b, e as indicated), 10 µM L-NAME (NOS inhibitor) (c, e as indicated), or 10 µM SB202190 (SB, p38 inhibitor) (d, e as indicated) for 15 minutes. Then VEGF-A (50 ng/ml) was applied to the top of the retina (abluminal side) and changes in microvessel permeability were recorded as described in Figure 2. Alternatively (e), retinae were fixed after 2 min using 4 % PFA and then immunostained with IB4 (green) and for phosphorylation of p38, HSP27 and eNOS (magenta). **(f-j)** As in panels (a-e), except that ex vivo retinae were stimulated with BK (10 µM). Representative results and quantifications (mean ± SD) from four independent experiments are shown. *p < 0.05, **p < 0.01, ***p < 0.001. Scale bars, 10 µm.

Figure 7. VEGF-A and Bradykinin induced VE-Cadherin phosphorylation.
(a) Ex vivo retinae were preincubated with SU1498 (10 μM), or BAPTA (20 μM) or STO (10 μM) and treated with VEGF-A and Bradykinin for 2 min. Retinae were then fixed with 4% PFA and immunostained for phospho-AMPKα (magenta) and with IB4 (green). **(b)** Ex vivo retinae were preincubated with compound C (10 μM), or L-NAME (10 μM) or SB202190 (10 μM), treated with VEGF-A and Bradykinin for 2 min and immunostained for phosphorylated VE-cad (using anti-pY685-VEC) (magenta) and with IB4 (green). **(c)** Proposed signalling networks in the ex-vivo retinal microvasculature downstream of VEGF-A and Bradykinin. Scale bars, 10 μm.

Supplemental Figure 1. Properties of the ex-vivo retina model.
(a-b) Selected frames, pseudocoloured in (b), of time-resolved recordings of a Sulforhodamine-B-filled rat capillary before and after the addition of 50 ng/ml VEGF-A, illustrating the rapid loss of fluorophore from the lumen of the vessel. The red box exemplifies a typical r.o.i used for intensity measurement. Scale bars, 10 μm. **(c-g)** CLDN5 siRNA was injected into mouse eyes. 48 h later CLDN5 levels were analysed by immunoblots of retinal lysates (c-d) or by wholemount immunochemical staining (e). Scale bar, 100 μm. Alternatively, permeability of 4 KDa Rhodamine was measured in ex vivo retinae from CLDN5 siRNA or control injected eyes (f-g). **p < 0.01, ***p < 0.001.