Stress induces cytoprotective TDP-43 nuclear bodies through lncRNA

***NEAT1*-promoted phase separation**

Authors: Chen Wang[1,2,*], Yongjia Duan[1,2,*], Gang Duan[1,2,*], Zhiwei Ma[1,†], Kai Zhang[1,2], Xue

Deng[1,2], Beituo Qian[1,2], Jinge Gu[1,2], Qiangqiang Wang[1], Shuang Zhang[1], Lin Guo[3], Cong Liu[1,2,§],

and Yanshan Fang[1,2,6,§]

Affiliations:

[1]Interdisciplinary Research Center on Biology and Chemistry, Shanghai Institute of Organic

Chemistry, Chinese Academy of Sciences, Shanghai 201210, China

[2]University of Chinese Academy of Sciences, Beijing, 100049, China,

[3]Department of Biochemistry and Molecular Biology, Thomas Jefferson University, Philadelphia,

PA 19107, USA

*These authors contributed equally

[†]Current address: West China School of Basic Medical Sciences and Forensic Medicine,

Sichuan University, Chengdu, 610041, China

[§]Correspondence to:

Cong Liu: liulab@sioc.ac.cn

Yanshan Fang: fangys@sioc.ac.cn

Tel: +86-21-6858.2510

Fax: +86-21-6858.2502

Short title (<50 characters): Phase-separated TDP-43 NBs mitigate stress

Keywords: TDP-43, nuclear body, RNP granule, phase separation, lncRNA *NEAT1*

Word count: ~6500 words (excluding the title page, abstract, references and figure legends)

Graphic Abstract

Highlights *(Up to four bullet points. The length of each highlight cannot exceed 85 characters,*

including spaces)

• TDP-43 forms reversible, LD-like NBs alleviating cytotoxicity and neurodegeneration

• The two RRMs act antagonistically to maintain a core-shell architecture of TDP-43 NBs

• LncRNA *NEAT1* is upregulated in stressed neurons, which promotes TDP-43 LLPS

• ALS-causing mutation D169G abolishes *NEAT1*-induced TDP-43 LLPS and NBs

SUMMARY (< 150 words)

Despite the prominent role of TDP-43 in neurodegeneration, its physiological and pathological

functions are not fully understood. Here, we report an unexpected function of TDP-43, which

forms dynamic, reversible, droplet-like nuclear bodies (NBs) in response to stress, and the

formation of NBs alleviates TDP-43-mediatd cytotoxicity and neurodegeneration in cell and

Drosophila models of amyotrophic lateral sclerosis (ALS). Super-resolution microscopy reveals a

"core-shell" architecture of TDP-43 NBs, which is maintained by the two RRMs antagonistically.

Further, TDP-43 NBs are partially colocalized with nuclear paraspeckles and the paraspeckle

scaffolding lncRNA *NEAT1* is dramatically upregulated in stressed neurons. Moreover, increase

of *NEAT1* promotes TDP-43 liquid-liquid phase separation (LLPS) *in vitro*. Finally, we uncover

that the ALS-associated mutation TDP-43^{D169G} impairs *NEAT1*-induced phase separation,

causing a specific defect in the assembly of TDP-43 NBs but not stress granules. Together, our

findings propose a stress-mitigating role and mechanism for TDP-43 NBs, whose dysfunction

may contribute to ALS pathogenesis.

INTRODUCTION

Amyotrophic lateral sclerosis (ALS) is a progressive motor neuron disease characterized by the degeneration of motor neurons in the brain and spinal cord, which leads to fatal paralysis generally within 2–5 years after diagnosis (*1,2*). Missense mutations in the gene *TARDBP* encoding the trans-activation response element (TAR) DNA-binding protein 43 (TDP-43) predispose to familial ALS, and protein inclusions containing TDP-43 in the affected motor neurons are a pathological hallmark of ALS (*3*, *4*).   TDP-43 is a nuclear protein but can shuttle between the nucleus and the cytoplasm. It has a nuclear localization signal (NLS), a nuclear export signal (NES), two canonical RNA recognition motifs (RRMs) that bind to nucleic acids, and a C-terminal low complexity domain (LCD) that mediates protein-protein interactions and is enriched of disease-associated mutations (*4*, *5*). TDP-43 is engaged in a variety of ribonucleoprotein (RNP) complexes and plays an important role in RNA processing and homeostasis (*6-10*). In addition, TDP-43 is known to participate in cytoplasmic stress granules (SGs), which may undergo aberrant phase transition and promote the formation of solid protein aggregates in diseased conditions (*11-14*).

Nuclear bodies (NBs) are dynamic, membraneless nuclear structures that concentrate specific nuclear proteins and RNAs and play an important role in maintaining nuclear homeostasis and RNA processing (*15*, *16*). Increasing evidence suggests the association of TDP-43 with a variety of NBs, such as paraspeckles (*17-19*), gemini of coiled bodies (GEMs) (*20*), nuclear stress bodies (*21*), interleukin-6 and -10 splicing activating compartment bodies (InSAC bodies) (*22*), and omega speckles (*23*). Notably, in the nucleus of spinal motor neurons of sporadic ALS patients, TDP-43 is found colocalized with paraspeckles and the occurrence of paraspeckles is specifically increased in the early phase of the disease (*18*). In addition, excessive formation of dysfunctional paraspeckles is also observed in the spinal neurons and glia of ALS-FUS patients (*24*). Although it is clear that TDP-43 is present in NBs that sometimes accompanies neurodegeneration, our understanding of the function of TDP-43 in NBs and the contribution of NBs to ALS pathogenesis is still rudimentary.

Recent studies indicate that liquid-liquid phase separation (LLPS) of RNA-binding protein

(RBPs) drives the assembly of liquid droplet (LD)-like, membraneless RNP granules in the

cytoplasm and nucleoplasm (*25-29*). Several ALS-related RBPs including TDP-43, FUS, hnRNP

A1, hnRNP A2 and TIA1 are shown to phase separate *in vitro* (*14, 30-35*). The intrinsically

disordered LCD domains of the RBPs are thought to mediate the LLPS (*28, 31, 36-38*) and

posttranslational protein modifications play an important role in determining the biophysical and

biological properties of the phase behavior of the RBPs (*34, 39-46*). In addition to the LCD,

recent studies reveal that RNA is of vital importance in regulating the LLPS (*47, 48*), which can

either suppress or promote phase separation depending on the contents and concentrations of

the RNAs (*25, 49-51*).

In this study, we find that various cellular stresses induce TDP-43 to form distinct, highly

dynamic and reversible NBs, which not only attenuate the cytotoxicity of TDP-43 in mammalian

cells but also ameliorate neurodegeneration and behavioral deficits in a *Drosophila* model of ALS.

Further investigation with super-resolution microscopy reveals the unexpected opposing

functions of the two RRMs in maintaining the core-shell architecture of TDP-43 NBs.

Furthermore, we demonstrate a crucial role of the paraspeckle scaffolding RNA nuclear-enriched

abundant transcript 1 (*NEAT1*) in promoting TDP-43 LLPS *in vitro*, which is compromised by the

ALS-causing mutation D169G, leading to a defect in forming TDP-43 NBs. Together, our findings

demonstrate how stress induces the assembly of TDP-43 NBs in the generally "suppressive"

nucleoplasm environment and suggest the involvement of dysfunctional TDP-43 NBs in ALS

pathogenesis.

RESULTS

Arsenic stress induces dynamic and reversible TDP-43 NBs

TDP-43 protein is predominantly localized to the nucleus but can shuttle between the nucleus

and the cytoplasm. In response to cellular stress, TDP-43 is recruited to cytoplasmic SGs (*11*).

Interestingly, in an earlier related work of our group (*46*), we noticed that although arsenite, a

commonly used experimental reagent to raise cellular stress (*52*), induced TDP-43$^+$ cytoplasmic

SGs, the majority of TDP-43 signal remained in the nucleus. Furthermore, in a great portion of

the cells, the nuclear TDP-43 lost the normal diffused pattern and instead exhibited a distinct

granular appearance (Movie S1 and Fig. 1, A and B).

A membraneless nuclear structure fulfills the requirements of NBs if it is: (1) microscopically

visible, (2) enriched with specific nuclear factors, and (3) continuously exchanging the contents

with the surrounding nucleoplasm (*15*). Indeed, arsenite-induced TDP-43 nuclear granules were

microscopically visible (Fig. 1). Further examination by immunocytochemistry indicated that

these arsenite-induced nuclear granules were colocalized or partially colocalized with several

known types of NBs, especially paraspeckles marked by splicing factor proline-glutamine rich

(SFPQ) (Fig. S1). These results suggest that cellular stress may be involved in the translocation

of TDP-43 into NBs such as paraspeckles, similar to that found in the spinal motor neurons of

ALS patients (*18*).

To examine if stress-induced TDP-43 nuclear granules were liquid-like and could exchange

with surrounding nucleoplasm, we performed the fluorescence recovery after photobleaching

(FRAP) analysis in *live* cells using a green fluorescent protein (GFP)-tagged TDP-43

(GFP-TDP-43). To induce TDP-43 nuclear granules, cells were pre-treated with arsenite (250 μM)

for 30 minutes before photobleaching. The fluorescence of GFP-TDP-43 nuclear granules was

rapidly recovered after photobleaching, reaching ~50% of the pre-bleaching intensity in ~65

seconds (Movie S2 and Fig. 1, C and D). Thus, these data indicate that stress-induced TDP-43

nuclear granules are liquid-like and highly dynamic. Together, they fulfilled all the above three

requirements and are thereafter called TDP-43 NBs in this study.

Next, we determined whether stress-induced TDP-43 NBs were reversible using arsenite

washout assay. After arsenite removal, the stress-induced TDP-43 NBs gradually disappeared

and eventually recovered to nearly diffused pattern like that before the arsenite treatment (Fig.

1E). Our quantitative analyses indicated that both the percentages of cells with TDP-43 NBs and

the number of TDP-43 NBs per cell decreased in a time-dependent manner after arsenite

washout (Fig. 1, F to H). Of note, there were considerably more cells with TDP-43 NBs than

TDP-43-associated (TDP-43$^+$) SGs (Fig. 1B) and overwhelmingly more TDP-43 NBs than

TDP-43$^+$ SGs per cell before the washout started (Fig. 1E, 0 min). Hence, although the

percentages of cells with visible TDP-43 NBs were not markedly reduced when TDP-43$^+$ SGs

became largely undetectable at 30 min after arsenite removal (Fig. 1, F and G), the average

number of TDP-43 NBs per cell did significantly decrease (Fig. 1H). Together, our data indicate

that arsenic stress-induced TDP-43 NBs are reversible as that of SGs upon the relief of stress.

Formation of TDP-43 NBs as a general cellular stress mechanism

Next, we examined if endogenous TDP-43 could also form NBs upon arsenic stress. We found

that in most cells, endogenous TDP-43 was present as evenly distributed while some cells

showed spontaneous small granules (Fig. S2A). The arsenite treatment led to the formation of

larger and more NBs in a significantly increased number of cells (Fig. S2, A and B). In addition,

we observed stress-induced TDP-43 NBs also in *in vitro* cultured mouse primary neurons (Fig.

S2, C and D). Thus, the formation of TDP-43 NBs is not an artificial effect due to TDP-43

overexpression or the use of non-neuronal cell lines. Moreover, it was previously reported that

heat shock-induced reversible nuclear TDP-43 "aggregations" were colocalized with the marker

of nuclear stress bodies (*21*). These observations raised the possibility that the formation of

TDP-43 NBs might be a general stress response mechanism in the nucleus. Indeed, we found

that disturbance of the nuclear homeostasis by inhibiting nuclear export with leptomycin B (LMB)

also effectively induced endogenous TDP-43 to form NBs (Fig. 2, A and B).

To further determine whether blocking nuclear export of TDP-43 itself was sufficient to

induce the formation of TDP-43 NBs, we generated mutations in the NLS and the NES of

TDP-43, respectively (NLS^{mut} and NES^{mut}; Fig. 2C). Under normal conditions, wild-type (WT)

TDP-43 was mostly present in diffused pattern in the nucleus with rare occurrence of

spontaneous TDP-43 NBs (Fig. 2, D to F). As expected, the NLS^{mut} TDP-43 was predominantly

cytoplasmic and thus the chance of TDP-43 NBs with the NLS^{mut} was even lower than that of the

WT (Fig. 2, D to F). In contrast, the NES^{mut} was exclusively nuclear and formed TDP-43 NBs

even in the absence of arsenite, heat shock or LMB treatment (Fig. 2, D to F). Furthermore, the

solubility of TDP-43-NES^{mut} protein was significantly reduced, as more than half of the protein

was insoluble in radioimmunoprecipitation assay buffer (RIPA) (Fig. 2, G and H).

The marked reduction in the solubility of the NES^{mu} protein raised the question whether the

TDP-43-NES^{mut} NBs were still in a dynamic, liquid-like state or became solid "aggregates". To

address this question, we examined the stability of WT and mutant TDP-43 proteins by treating

cells with protein synthesis inhibitor cycloheximide (CHX) (53). We found that the RIPA-soluble

TDP-43 protein was rather stable, as none of the WT, NLS^{mut} or NES^{mut} TDP-43 in the soluble

fractions showed significant turnover in 24 h after CHX treatment (Fig. S3, A to D). In contrast,

the RIPA-insoluble fraction of TDP-43-NES^{mut} decreased rapidly upon CHX inhibition and

became almost undetectable within 24 h (Fig. S3, A to D). Meanwhile, we found that the

treatment with CHX led to rapid disassembly of TDP-43-NES^{mut} NBs and reduction in the number

of TDP-43-NES^{mut} NBs per cell (Fig. S3, E and F). These results suggest that the RIPA-insoluble

TDP-43-NES^{mut} NBs are in a reversible, liquid-like state rather than solid "aggregates". A similar

observation of misfolded proteins entering the liquid-like, granular nucleolus under stress was

recently reported, which may function as a phase-separated protein quality control compartment

(54).

Next, we further examined how disturbance of the cellular proteostasis by blocking protein

turnover impacted on this process. We inhibited proteasome-mediated protein degradation by

MG132, which led to a significant increase in the RIPA-insoluble fraction of the NES^{mut} as well as

that of the WT TDP-43 protein (Fig. S4, A to D). In contrast, blocking autophagic flux by

chloroquine (CQ) did not significantly affect the levels of insoluble TDP-43 (Fig. S4, E to H). Thus,

the turnover of TDP-43 NBs did not rely on the autophagy-lysosomal pathway, which was more

commonly used to clear misfolded protein aggregates (*55, 56*). Together, our data indicate that

TDP-43 NBs are dynamic, reversible and sensitive to the environment of surrounding

nucleoplasm, which may act as a general cellular mechanism to various stresses.

Formation of TDP-43 NBs alleviates the cytotoxicity

We were keen to understand the functional significance of forming TDP-43 NBs. Was it a

"protective" mechanism for cells, like that of SGs? Indeed, we found that overexpression (OE) of

WT or NLS^{mut} TDP-43 but not the NB-forming NES^{mut} in human 293T cells led to decreased cell

viability (Fig. 3A). To further confirm that the formation of TDP-43 NBs was required for the

cytotoxicity-antagonizing effect rather than simply restricting TDP-43 in the nucleus, we sought

for alternative approaches to promote or suppress TDP-43 NBs without disrupting its

nucleocytoplasmic transport. It was previously shown that an acetylation-mimic K145/192Q

mutation of TDP-43 impaired the RNA-binding and promoted the formation of TDP-43 protein

inclusions in both the nucleus and the cytoplasm, whereas the acetylation-deficient K145/192R

mutation suppressed that (*40*). Hence, we generated the K145/192Q mutation in WT TDP-43

(Fig. 3B) and the K145/192R mutation in TDP-43-NES^{mut} (Fig. 3C).

Indeed, the K145/192Q mutation led to the formation of TDP-43 NBs without affecting the

nucleocytoplasmic distribution of TDP-43 and that was in the absence of LMB or arsenite

treatment (Fig. 3, D, D', F, and H). And importantly, the NB-promoting K145/192Q mutation

alleviated the cytotoxicity of WT TDP-43 (Fig. 3J), similar to the effect of the NES^{mut} (Fig. 3A). On

the other hand, abolishing NB formation of the NES^{mut} by the K145/192R mutation (Fig. 3, E, E',

G, and I) made the originally non-toxic TDP-43-NES^{mut} to exhibit marked cytotoxicity (Fig. 3K),

like that of the WT TDP-43 (Fig. 3A). These data demonstrate that the formation of TDP-43 NBs

reduces the cytotoxicity and increases the cell viability, therefore potentially acting as a

"protective" mechanism to help cells survive stressed or diseased conditions.

TDP-43-NES^{mut} forms neuronal NBs *in vivo* and mitigates neurodegeneration in a

***Drosophila* model of ALS**

To validate our findings in an *in vivo* model, we utilized the binary GAL4-UAS system (*57*) to

generate transgenic flies in order to express WT, NLS^{mut} or NES^{mut} human TDP-43 (hTDP-43) in

a tissue-specific manner. The transgenes of WT and mutant UAS-*hTDP-43* were inserted in the

same locus of the fly genome by ΦC31 integrase-mediated, site-specific integration (*58-60*). This

approach avoided the location effect on transgene expression across different fly lines and

allowed direct comparison between the WT and mutant TDP-43 flies. Immunohistochemistry

analysis of the whole mount fly brains confirmed the predominant nuclear localization of the WT

and NES^{mut} hTDP-43 in the fly neurons (*elav*-Gal4), whereas the NLS^{mut} was largely cytoplasmic

(Fig. 4A). Consistent with the data in 293T cells (Fig. 2, D to F), the NES^{mut} hTDP-43 formed

striking NBs in the nucleus of fly neurons (Fig. 4, A and B). Furthermore, the solubility of the

hTDP-43-NES^{mut} protein extracted from the fly brains was significantly decreased (Fig. 4, C and

D), which was also consistent with the results in the mammalian cells (Fig. 2, G and H).

We then examined the consequence of expressing WT, NLS^{mut} or NES^{mut} hTDP-43 in the fly

eye (GMR-Gal4). WT and NLS^{mut} hTDP-43 flies showed slight eye degeneration at Day 5, as

evidenced by the rough surface of the compound eyes and loss of pigment cells. At the same

time point, however, the eyes of the NES^{mut} flies were largely unaffected and the degeneration

score was much lower (0.4 ± 0.34) than that of WT hTDP-43 flies (1.5 ± 0.41, *P* < 0.001) (Fig. 4E).

By Day 25, the eyes of WT and NLS^{mut} hTDP-43 flies further degenerated as a lot more pigment

cells were lost and some eyes showed swelling especially in the NLS^{mut} hTDP-43 flies.

Compared to WT hTDP-43 flies (2.5 ± 0.13), the eye degeneration of the NLS^{mut} flies was

significantly worse (3.8 ± 0.17, *P* < 0.001), whereas that of the NES^{mut} flies was substantially

milder (0.8 ± 0.14, *P* < 0.001) (Fig. 4E). Similar conclusions were obtained from the climbing

assay, which also showed a striking difference in the severity of neurotoxicity among WT, NLS[mut]

and NES[mut] hTDP-43 flies. Expression of WT hTDP-43 in the fly motor neurons (D42-Gal4)

caused an age-dependent decline of the climbing capability, which deteriorated much faster in

the NLS[mut] flies (Fig. 4F). The behavioral phenotype of NES[mut] flies was the mildest, whose

climbing capability was not statistically different from that of the control flies (UAS-*Luciferase*) at

all time points examined (Fig. 4F). Together, these data indicate that the formation of TDP-43

NBs by the NES[mut] alleviates the TDP-43-mediated neurodegeneration in an *in vivo Drosophila*

model of ALS.

The role of the major functional domains of TDP-43 in the assembly of NBs

To understand how TDP-43 NBs were assembled, we generated truncated TDP-43 proteins,

namely ΔLCD, ΔRRM1 and ΔRRM2, to determine how each major domain of TDP-43 is involved

in regulating this process (Fig. S5A). It was previously shown that the intrinsically disordered

LCD domain mediated protein-protein interaction and was critical for the formation of cytoplasmic

SGs and protein aggregations (*31*, *61*). We found that both the WT and ΔLCD TDP-43 were

mostly soluble (Fig. S5, B and C), however, the ΔLCD was unable to form NBs in response to

LMB (Fig. S5, F to G'), indicating that the formation of TDP-43 NBs in cells requires the LCD

domain.

Next, we examined the impact of deleting RRM1 or RRM2 on the assembly of TDP-43 NBs.

The ΔRRM1 TDP-43 was largely soluble (Fig. S5D), while the ΔRRM2 showed a greatly

increased insolubility (Fig. S5E). Interestingly, the ΔRRM1 TDP-43 formed large, ring-like

structures in the nucleus even in the absence of LMB treatment (Fig. S5H). And, treatment with

LMB did not promote further formation of the ring-like structures or trigger the formation of

WT-like TDP-43 NBs (Fig. S5H'). ΔRRM2 TDP-43 looked similar to WT in cells without stress

(Fig. S5I), whereas treating with LMB gave rise to a grainy appearance (Fig. S5I').

Super-resolution microscopy reveals a "core-shell" architecture of TDP-43 NBs

maintained by the opposing functions of the two RRMs

To further characterize the nuclear structures formed by WT, ΔRRM1 and ΔRRM2 TDP-43, we

utilized the Leica LIGHTNING SP8 confocal microscope to capture multicolor images in

super-resolution down to 120 nm. Interestingly, we found that WT TDP-43 formed solid, mid-size

NBs; ΔRRM1 was much larger, ring-shaped; and ΔRRM2 displayed a mesh-like structure with

many smaller NBs (Fig. 5, A and B). Three-dimensional (3D) rendering revealed that WT TDP-43

NBs were not perfect "spheres" as ones might infer from the 2D images; rather, they were in oval

or cylinder shapes in the cell. Unlike WT TDP-43, ΔRRM1 was mostly large, hollow "pillars";

whereas ΔRRM2 NBs looked similar to those of WT in 3D rendering. Further quantification

confirmed that the average diameter of ΔRRM1 "rings" was drastically larger whereas that of

ΔRRM2 NBs was noticeably smaller than that of WT TDP-43 (Fig. 5C).

Given the distinct morphology of the nuclear structures formed by ΔRRM1 and ΔRRM2

TDP-43, it was tempting to propose a "core-shell" architecture of TDP-43 NBs and the two RRMs

might function antagonistically to maintain such a structure – the RRM1 pushed TDP-43 toward

the core of NBs whereas the RRM2 pulled TDP-43 toward the shell (Fig. 5D). If this hypothesis

were correct, ΔRRM1 TDP-43 that lacked the "centripetal force" of the RRM1 would be mainly

distributed to the shell of NBs, while ΔRRM2 TDP-43 that lacked the "centrifugal force" would be

mainly located in the core. To test this hypothesis, we co-expressed HA-Tagged WT TDP-43

with Myc-tagged WT, ΔRRM1 or ΔRRM2 TDP-43 and induced NB formation in 293T cells by

LMB (Fig. 5, E to H). Indeed, ΔRRM1 formed a ring-like shell wrapping the co-expressed WT

TDP-43 NBs (Fig. 5, F and J) and the diameter of the NBs was dramatically larger (Fig. 5N). The

increased diameter was not simply due to overexpression of two folds of TDP-43 protein, as

co-expression of WT TDP-43-HA and WT myc-TDP-43 did not show such effect (Fig. 5, E, I, and

M). Consistent with our hypothesis, we found that ΔRRM2 co-expressed with the WT TDP-43

was mainly located in the core of NBs (Fig. 5, G and K) and the diameter was significantly

smaller (Fig. 5O). The core-shell architecture could also be clearly seen when ΔRRM1 and

ΔRRM2 TDP-43 were co-expressed (Fig. 5, H, L, and P). Therefore, the data suggest that RRM1

and RRM2 play different or even opposing functions in the assembly of TDP-43 NBs, possibly by

interacting with different RNAs and other RNA-binding NB proteins via the different RRMs.

RNA suppresses TDP-43 *in vitro* de-mixing via the RRMs

Recent studies indicate that the de-mixing process by LLPS condenses RBPs into LD-like

compartments, allowing for the formation of intracellular membraneless RNP granules (*25-29*).

Several ALS-related RBPs including TDP-43, FUS, hnRNP A1, hnRNP A2 and TIA1 have been

shown to phase separate *in vitro* (*14*, *30-35*). The dynamic, reversible and membraneless

characteristics of TDP-43 NBs suggested that LLPS might be involved in the formation of

TDP-43 NBs. Indeed, we found that purified full-length (FL) TDP-43 underwent LLPS and formed

LDs *in vitro* in a dose-dependent manner (Fig. S6, A to C). Similar to the observation in cells (Fig.

S5, G and G'), the C-terminal LCD truncation (TDP-43$^{1-274}$) showed greatly reduced LLPS

capability (Fig. S6D), suggesting a major role of the LCD in mediating TDP-43 de-mixing *in vitro*

and *in vivo*. Nevertheless, the formation of TDP-43$^{1-274}$ LDs was evident at a higher protein

concentration (50 μM) (Fig. S6D), confirming the previous observation that the N-terminal

domains of TDP-43 could drive LLPS *in vivo* (*32*).

To gain insights into the mechanism involved in the regulation of TDP-43 NBs, we

determined how RNA impacts on the LLPS of TDP-43 by adding total RNAs extracted from HeLa

cells into the *in vitro* de-mixing system (Fig. S6, E and F). The addition of total RNAs markedly

reduced the LLPS of FL TDP-43 at a concentration of 500 ng/μl (Fig. S6E), whereas a much

lower concentration of total RNAs (100 ng/μl) was sufficient to suppress the LLPS of TDP-43$^{1-274}$

(Fig. S6F). These results confirm the predominant LLPS-promoting role of the C-terminal LCD

and suggest an LLPS-suppressing function for the N-terminal RNA-binding regions. We then

moved on to test how the two RRMs were involved in the RNA suppression of TDP-43 LLPS.

Since the LLPS-promoting effect of the LCD in TDP-43 was overwhelming and made the function

of the RRMs in RNA suppression difficult to assess in the *in vitro* de-mixing assay (Fig. S6, E and

F), we instead generated and tested ΔRRM1 and ΔRRM2 TDP-43$^{1-274}$ proteins (Fig. 6A and Fig.

S6G). We confirmed that the RNA-binding affinity of ΔRRM1 and ΔRRM2 TDP-43$^{1-274}$ was

significantly reduced (Fig. S6, H and I). WT as well as ΔRRM1 and ΔRRM2 TDP-43$^{1-274}$ formed

LDs by LLPS *in vitro* in a dose-dependent manner, though the de-mixing of ΔRRM1 TDP-43$^{1-274}$

appeared less robust than that of WT or ΔRRM2 (Fig. S6, J to L).

To better compare the difference in RNA suppression, we lowered the NaCl concentration

and increased the crowding agent PEG, which allowed WT TDP-43$^{1-274}$ to form larger LDs (Fig.

6B). We found that total RNAs at a concentration of 100 ng/μl markedly reduced the size and

number of LDs of WT TDP-43$^{1-274}$. ΔRRM1 TDP-43$^{1-274}$ did not form large LDs as WT at the

concentration of 50 μM and we initially thought its LLPS would be more easily suppressed by

total RNAs. To our surprise, a much higher concentration (500 ng/μl) of total RNAs was required

to see effective suppression (Fig. 6C), suggesting that deletion of the RRM1 greatly reduced the

ability of total RNAs to suppress the formation of TDP-43 droplets, which was consistent with the

finding that ΔRRM1 TDP-43 spontaneously phase separated into ring-like nuclear structures in

cells even in the absence of cellular stress (Fig. S5H). We also tested ΔRRM1 TDP-43$^{1-274}$ at 100

μM to allow large ΔRRM1 LDs to form and the result confirmed that the suppression of TDP-43

LLPS by total RNAs was dramatically reduced with ΔRRM1, requiring total RNAs of about 250

ng/μl (Fig. 6C'). The suppression of the LLPS of ΔRRM2 TDP-43$^{1-274}$ by total RNAs (100~250

ng/μl) was slightly reduced compared to that of WT (Fig. 6, B and D), but to a less extent than

that of ΔRRM1 (Fig. 6, C and C'). It was worth noting that ΔRRM1 TDP-43$^{1-274}$ showed both

decreased LLPS capacity and reduced sensitivity to RNA suppression, indicating that a RBP

with a lower tendency to phase separate on its own does not necessarily makes its LLPS more

sensitive to RNA suppression. Nevertheless, these data indicate that both RRMs are involved in

RNA suppression of TDP-43 LLPS.

Upregulation of *NEAT1* RNA promotes TDP-43 LLPS

Recent works have revealed both positive and negative regulation of LLPS of prion-like

RBPs by RNA (*62*). For example, tRNA was shown to reduce the LLPS of FUS *in vitro* (*50*).

Similarly, tRNA also potently suppressed the *in vitro* LLPS of TDP-43 (Fig. 6E). We further

examined the function of the two RRMs in mediating tRNA suppression. We found that ΔRRM1

TDP-43$^{1-274}$ showed similar sensitivity to tRNA suppression, as 1.25 μg/μl of tRNA was sufficient

to suppress the LLPS of both WT and ΔRRM1 TDP-43$^{1-274}$ (Fig. 6, E and F). At this condition,

however, ΔRRM2 LDs could still be observed and the effective suppression required a higher

concentration of tRNA (2.5 μg/μl). It should be pointed out that, unlike the response to total RNAs,

ΔRRM1 TDP-43$^{1-274}$ did not exhibit a greatly decreased sensitivity to tRNA suppression, despite

that again only small ΔRRM1 LDs were present at the beginning of the suppression test. Thus,

the lack of a strong suppression of ΔRRM1 LLPS by total RNAs (Fig. 6C) was not simply

because suppression of small LDs was difficult to detect.

Since stress-induced TDP-43 NBs were co-localized with paraspeckles (Fig. S1, A and B)

and TDP-43 bound directly to the paraspeckle scaffolding RNA *NEAT1* (*18*, *63*), we next

examined whether *NEAT1* RNA impacted on the phase behavior of TDP-43. Interestingly, we

found that *NEAT1* promoted the LLPS of TDP-43 in a dose-dependent manner (Fig. 6H), and

this regulation by *NEAT1* required both the RRM1 and RRM2 of TDP-43 (Fig. 6, I and J).

Furthermore, we showed that increasing concentrations of *NEAT1* antagonized the suppressing

effect of tRNA, inducing the formation of TDP-43 droplets *in vitro* (Fig. 6K). In this suppressive

environment, the *NEAT1*-mediated induction of the LLPS of ΔRRM1 TDP-43$^{1-274}$ was markedly

reduced (Fig. 6L), which was consistent with the observations that ΔRRM1 TDP-43 remained

soluble and LMB did not promote the formation of NBs or increase the number of the ring-like

nuclear structures in cells (Fig. S5, D, H and H'). Under the same condition, *NEAT1* RNA more

readily induced the *in vitro* de-mixing of ΔRRM2 TDP-43$^{1-274}$ (Fig. 6M), likely because tRNA

suppression of ΔRRM2 was less effective (Fig. 6G). The overall increased tendency to form LDs

and to remain phase separated might underlie the formation of more condensed, insoluble

TDP-43 NBs by ΔRRM2 (Fig. S5, E, I and I').

Given that *NEAT1* RNA could act as scaffolds to promote the nucleation of paraspeckles (*50,*

64-67) and that stress-induced TDP-43 NBs were partially co-localized with paraspeckles (Fig.

S1, A and B), our data strongly suggest that *NEAT1* plays an important role in triggering the

formation of stress-induced TDP-43 NBs in cells. Supporting this hypothesis, we found that the

levels of both the total *NEAT1* RNA and the long non-coding RNA (lncRNA) isoform *NEAT1_2*

were dramatically increased in mouse primary neurons treated with arsenite (Fig. 6N) or LMB

(Fig. 6O), consistent with the observation earlier in this study that TDP-43 formed NBs in

stressed neurons (Fig. S2, C and D). Together, we propose that cellular stress increases NEAT1

RNA levels, which antagonize the suppressive environment of the nucleoplasm by providing

nucleation scaffolds that condensate TDP-43 and other NB components, triggering the formation

of TDP-43 NBs in stressed cells.

ALS-associated D169G mutation of TDP-43 shows a specific defect in NB formation

This study was launched by the discovery that TDP-43 formed cytotoxicity-antagonizing NBs in

mammalian cells and the formation of NBs alleviated TDP-43-mediated neurodegeneration in

flies. To further investigate the relevance of TDP-43 NBs in the pathogenesis of human diseases,

we examined several known ALS-causing TDP-43 mutants. Interestingly, we found that the

mutation of D169G (640A>T) (*4*; and Fig. 7A) dramatically reduced the capability of TDP-43 to

form NBs in response to stress induced by arsenite (Fig. 7, B to D) or LMB (Fig. S7, A to C). In

contrast, the nucleocytoplasmic transport and recruitment of D169G TDP-43-D169G to SGs was

unaffected (Fig. 7, B and E), which was consistent with the previous assessments of D169G in

SG formation (*68, 69*). The D169G mutation is in the RRM1, which is thought to cause a local

conformational change that increases the thermal stability of the RRM1 (*70, 71*). It is worth

noting that most known disease-associated mutations such as M337V are enriched in the LCD

region of TDP-43 (Fig. S7, D and E), which are thought to promote aberrant phase transition of

SGs to irreversible protein aggregates in disease (*11*). Unlike D169G, we did not observe a

significant effect of M337V mutation on the assembly of TDP-43 NBs (Fig. S7, F and G). Thus,

TDP-43^{D169G} has a specific defect in the formation of stress-induced NBs. In addition, these

findings indicate that although TDP-43 NBs and SGs are both phase-separated RNP granules,

the exact mechanisms regulating their assembly are different.

Next, we examined if the phase separation behavior of TDP-43-D169G was altered, which

might contribute to the defect in TDP-43 NB formation. We found that purified D169G TDP-43$^{1-274}$

protein formed LDs in a dose-dependent manner just like that of WT in the *in vitro* LLPS assay

(Fig. 7, F, G and H), suggesting that the ability of D169G to phase separate was not dramatically

changed. Further, the suppression of TDP-43 LLPS by total RNAs showed no difference in

D169G compared to that of WT TDP-43$^{1-274}$ (Fig. 7, I and J), suggesting the normal regulation of

TDP-43 RNP granules by total RNAs remained intact in D169G. This was consistent with the

result that D169G did not affect SG assembly or TDP-43 translocation to SGs (Fig. 7B and 7E).

In striking contrast, the induction of TDP-43 LLPS by *NEAT1* RNA was drastically reduced in

D169G (Fig. 7, K and L). Thus, these data not only well cohered with the observation that

TDP-43-D169G was defective in NB formation but also further confirmed the crucial role of

NEAT1 in triggering the assembly of TDP-43 NBs. Together with the cytotoxicity-antagonizing

function of TDP-43 NBs (Fig. 3 and Fig. 4), the findings of D169G (Fig. 7) suggest that a

compromise of stress-induced TDP-43 NBs may be involved in ALS pathogenesis.

DISCUSSION

This study was initiated by the finding that TDP-43 formed distinct nuclear granules in stressed

cells. We then characterized these subnuclear structures and demonstrated that they meet the

three criteria of NBs (*15*) – (1) microscopically distinctive; (2) enriched of TDP-43 and partially

colocalized with paraspeckles; and (3) liquid or hydrogel-like, highly dynamic and sensitive to the

surrounding nucleoplasm. Further, we show that various cellular stresses such as arsenic stress,

nuclear export blockage and proteasomal inhibition induce the formation of TDP-43 NBs. And

TDP-43 NBs are detected in human cells, mouse primary neurons, and fly brains *in vivo*. Thus,

the assembly of TDP-43 NBs may be a general stress response that occurs in multiple cell types

and different organisms.

The cytotoxicity-antagonizing effect of TDP-43 NBs

Although TDP-43 insoluble aggregates are found in the nucleus and cytoplasm of affected

neurons in ALS and FTD patients, reducing TDP-43 aggregates did not reduce or prevent cell

death (*72*), suggesting that TDP-43 aggregation is associated with rather than a primary cause of

neurodegeneration. Interestingly, we find in this study that NB-forming TDP-43 such as the

NES^{mut} and the K145/192Q mutation are much less cytotoxic than diffused WT TDP-43. More

importantly, abolishing the capability of TDP-43-NES^{mut} to form NBs makes the originally

non-toxic NES^{mut} exhibit cytotoxicity. Furthermore, transgenic flies expressing TDP-43-NES^{mut}

form remarkable NBs in the fly neurons and the neurodegeneration is considerably less severe

than that of the TDP-43-WT flies. In line of the cytoprotective role, TDP-43^{+} paraspeckles are

observed in the spinal motor neurons of ALS patients especially in the early phases of the

disease (*18*), suggesting that TDP-43 NBs may be called up and engaged in the first line of

defense against cellular stress and disease conditions.

Paraspeckles are a class of subnuclear RNP granules formed by the scaffolding lncRNA

NEAT1 associated with SFPQ and other paraspeckle proteins (*64, 66, 73-75*), which are

engaged in regulating different cellular functions through nuclear retention of specific mRNAs

and proteins (*76-78*). SGs sequester mRNAs and temporarily arrest protein synthesis to help

cells survive stress (*79*). Given that TDP-43 is presented in both paraspeckles and SGs and

plays an important role in regulating gene expression and RNA metabolism (*63, 80-82*), it is

reasonable to speculate that a similar underpinning mechanism may contribute to the

cytoprotective function of TDP-43 NBs. For example, assembly of TDP-43 NBs may stall DNA

transcription and/or arrest RNA processing; when stress is relieved, TDP-43 NBs disassemble

and release the RNAs and nuclear proteins required for normal cell functions. In addition, since

intranuclear inclusions of TDP-43 are also found in ALS patients (*83*), our data do not exclude

the possibility that TDP-43 NBs may become irreversible protein aggregates under extended

stress or diseased conditions.

The architecture of TDP-43 NBs and the functions of the two RRMs

The super-resolution 3D rendering reveals that TDP-43 NBs are ovals or cylinders in the nucleus

rather than perfect spheres as inferred from the 2D images. The "roundness" has been described

as one of the basic features for the notion that RNP granules are membraneless, intracellular

compartments formed by LCD-mediated phase separation. Our data suggests that substantial

additional forces, possibly provided by protein-protein and/or protein-RNA interactions,

participate in assembling and shaping RNP granules, at least for the case of TDP-43 NBs.

TDP-43 contains two canonical RRMs in tandem. RRM1 has a longer Loop3 region than

RRM2 and is thought to have a higher affinity for RNA targets (*84-85*). Although we did not

detect a significant difference in the binding affinity between ΔRRM1 and ΔRRM2 to total RNAs

in the dot-blot assay, they do show divergent sensitivities to RNA suppression in the *in vitro*

LLPS assays, likely because that the two RRMs have different RNA preferences (*84*). It needs to

be noted that the effects of RNA on ΔRRM1 and ΔRRM2 TDP-43 were compared without the

LCD in the *in vitro* LLPS assays. This is because the LCD presents an overwhelming

LLPS-driving force (*25, 31, 37, 61*), whereas the *in vitro* LLPS system appears to be much less

suppressive than the conditions in the nucleoplasm (see below).

In cells, TDP-43 is diffused in the nucleus and does not form NBs spontaneously under

normal conditions. And, the partition of TDP-43 into phase-separated intracellular compartments

requires the LCD (*31*, *61*, and also see Fig. S5). Thus, ΔRRM1 and ΔRRM2 are examined

inclusive of the LCD in cell-based experiments in this study. ΔRRM1 lacks the RRM1-mediated

suppression but contains the forceful LLPS-promoting LCD, which leads to spontaneous phase

separation of ΔRRM1 in the nucleus. Meanwhile, since the RRM1 is also used by TDP-43 to

anchor itself to the core of the NBs, the phase-separated ΔRRM1 forms large, ring-like structures.

The seemingly conflicting effects of RRM1 – inhibiting TDP-43 phase separation under normal

conditions whereas pushing TDP-43 toward the core of the NBs when the LLPS occurs – may be

achieved by interacting with different RNAs. In contrast, since ΔRRM2 retains the main RNA

suppression force mediated by RRM1, it does not spontaneously phase separate in the nucleus.

When stressed, ΔRRM2 forms small, condensed NBs possibly because the RRM2-mediated

"centrifugal force" is no longer available to counteract the RRM1-mediated "centripetal force".

Together, we propose a "push-and-pull" model for the opposing effects of the two RRMs in

maintaining the "core-shell" architecture of TDP-43 NBs.

The role of RNAs in regulating TDP-43 LDs and NBs

RNA can either promote or suppress the assembly of RNP granules depending on the

concentrations and the types of the RNA (15, 16, *25*, *26*). In addition, the ALS/FTD-related

C9ORF72 repeat RNA promotes phase transition and assembly of RNP granules in cells (*86*). In

this study, we show that tRNA suppresses TDP-43 LLPS, similar to the effect on FUS (*50*);

however, tRNA was previously shown to enhance Tau LLPS (*87*). Thus, even for the same RNA,

its role may vary depending on the protein that it interacts with.

lncRNAs are thought to provide the scaffold for the assembly of paraspeckles (*64*, *65*, *67*)

and nuclear stress bodies (*88*, *89*). In particular, the paraspeckle lncRNA *NEAT1* are increased

in patients and/or animal models of several neurodegenerative diseases including ALS (*18*),

Alzheimer's disease (*90*), Parkinson's disease (*91*), Huntington's disease (*92*), and multiple

sclerosis (*93*). In this study, we find that *NEAT1* lncRNA is substantially upregulated in stressed

neurons and TDP-43 NBs are partially colocalized with paraspeckles, cohering with the similar

observations in the spinal motor neurons of early ALS patients (*18*). Further *in vitro* experiments

indicate that *NEAT1* RNA not only promotes TDP-43 LLPS, but also antagonizes the generally

"suppressive" environment modeled using tRNA. Thus, stress-induced upregulation of *NEAT1*

may provide nucleation scaffolds to concentrate TDP-43 and other NB proteins, which overturns

the suppressive environment in the nucleoplasm, leading to the emergence of TDP-43 NBs. It is

worth noting that lncRNAs are potential targets of TDP-43 (*94-96*), implying complicated

cross-regulation between TDP-43 and *NEAT1* in this cellular stress response mechanism.

The D169G mutation and the relevance of TDP-43 NBs in ALS pathogenesis

More than 30 disease-causing mutations have been identified in the gene encoding TDP-43, of

which most are in the LCD and the adjacent regions (*5*, *97*). D169G is the only known

ALS-causing mutation within the RRM of TDP-43 (*4*) and its pathogenic mechanism has been

elusive. Initially, it was presumed to abrogate RNA binding due to its location within the RRM (*4*).

However, subsequent work indicates that the binding affinity of D169G to RNA or DNA is not

decreased (*85*). And, although TDP-43^{D169G} exhibits slightly increased thermal stability and may

be more susceptible to proteolytic cleavage (*70*, *71*), it does not engender aberrant

oligomerization but rather increases the resistance of TDP-43 to aggregation (*70*). Further, the

D169G mutation does not affect TDP-43 subcellular distribution or SG recruitment (*68*, *69*; also

see Fig. 7). The only reported cellular alteration by D169G is that it reduces polyubiquitination

and co-aggregation of TDP-43 with Ubiquilin 1 (98). However, given the long-standing notion that

ubiquitinated cytoplasmic inclusions containing TDP-43 is a hallmark and major cause of the

disease (*6*, *10*, *11*), it is difficult to understand how decreased ubiquitination and aggregation of

TDP-43^{D169G} leads to disease pathogenesis.

In this study, we discover that TDP-43^{D169G} has a striking and specific defect in the formation

of stress-induced TDP-43 NBs but not SGs. D169G is thought to cause a small local

conformational change in the Turn6 of the β-sheets within the RRM1 of TDP-43 (*70*, *71*), but

does not reduce the overall binding affinity of TDP-43 to nucleic acids (*85*). Indeed, we find that

the suppression of TDP-43 LLPS by total RNAs is unaffected by D169G, whereas the induction

of TDP-43 LLPS by *NEAT1* is dramatically diminished. Together with the cytoprotective role of

TDP-43 NBs, D169G provides a paradigm for compromise of TDP-43 NBs involved in ALS

pathogenesis. Given that cytoplasmic mislocalization and nuclear depletion of TDP-43 are

common in diseased neurons, it is likely that loss of the TDP-43 NB-mediated stress-mitigating

function may also underlie other cases of ALS and related diseases.

REFERENCES AND NOTES

1. J. P. Taylor, R. H. Brown, Jr., D. W. Cleveland, Decoding ALS: from genes to mechanism. *Nature* **539**, 197-206 (2016).

2. M. A. van Es *et al.*, Amyotrophic lateral sclerosis. *The Lancet* **390**, 2084-2098 (2017).

3. M. Neumann *et al.*, Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Science* **314**, 130-133 (2006).

4. E. Kabashi *et al.*, TARDBP mutations in individuals with sporadic and familial amyotrophic lateral sclerosis. *Nat Genet* **40**, 572-574 (2008).

5. E. B. Lee, V. M. Lee, J. Q. Trojanowski, Gains or losses: molecular mechanisms of TDP43-mediated neurodegeneration. *Nat Rev Neurosci* **13**, 38-50 (2011).

6. A. Jovicic, J. W. Paul, 3rd, A. D. Gitler, Nuclear transport dysfunction: a common theme in amyotrophic lateral sclerosis and frontotemporal dementia. *J Neurochem* **138 Suppl 1**, 134-144 (2016).

7. A. Ratti, E. Buratti, Physiological functions and pathobiology of TDP-43 and FUS/TLS proteins. *J Neurochem* **138 Suppl 1**, 95-111 (2016).

8. H. Ederle, D. Dormann, TDP-43 and FUS en route from the nucleus to the cytoplasm. *FEBS Letters* **591**, 1489-1507 (2017).

9. B. Fahrenkrog, A. Harel, Perturbations in Traffic: Aberrant Nucleocytoplasmic Transport at the Heart of Neurodegeneration. *Cells* **7**, 232 (2018).

10. M. Zhao, J. R. Kim, R. van Bruggen, J. Park, RNA-Binding Proteins in Amyotrophic Lateral Sclerosis. *Mol Cells* **41**, 818-829 (2018).

11. Y. R. Li, O. D. King, J. Shorter, A. D. Gitler, Stress granules as crucibles of ALS pathogenesis. *J Cell Biol* **201**, 361-372 (2013).

12. M. Ramaswami, J. P. Taylor, R. Parker, Altered ribostasis: RNA-protein granules in degenerative disorders. *Cell* **154**, 727-736 (2013).

13. T. Murakami *et al.*, ALS/FTD Mutation-Induced Phase Transition of FUS Liquid Droplets and Reversible Hydrogels into Irreversible Hydrogels Impairs RNP Granule Function. *Neuron* **88**, 678-690 (2015).

14. A. Patel *et al.*, A Liquid-to-Solid Phase Transition of the ALS Protein FUS Accelerated by Disease Mutation. *Cell* **162**, 1066-1077 (2015).

15. D. Stanek, A. H. Fox, Nuclear bodies: news insights into structure and function. *Curr Opin Cell Biol* **46**, 94-101 (2017).

16. M. Wegener, M. Muller-McNicoll, Nuclear retention of mRNAs - quality control, gene regulation and human disease. *Semin Cell Dev Biol* **79**, 131-142 (2018).

17. E. B. Dammer *et al*., Coaggregation of RNA-binding proteins in a model of TDP-43
proteinopathy with selective RGG motif methylation and a role for RRM1 ubiquitination.
PLoS One **7**, e38658 (2012).

18. Y. Nishimoto *et al*., The long non-coding RNA nuclear-enriched abundant transcript 1_2
induces paraspeckle formation in the motor neuron during the early phase of amyotrophic
lateral sclerosis. *Mol Brain* **6**, 31 (2013).

19. J. A. West *et al*., Structural, super-resolution microscopy analysis of paraspeckle nuclear
body organization. *J Cell Biol* **214**, 817-830 (2016).

20. T. Ishihara *et al*., Decreased number of Gemini of coiled bodies and U12 snRNA level in
amyotrophic lateral sclerosis. *Hum Mol Genet* **22**, 4136-4147 (2013).

21. M. Udan-Johns *et al*., Prion-like nuclear aggregation of TDP-43 during heat shock is
regulated by HSP40/70 chaperones. *Hum Mol Genet* **23**, 157-170 (2014).

22. S. Lee *et al*., Identification of a subnuclear body involved in sequence-specific cytokine RNA
processing. *Nat Commun* **6**, 5791 (2015).

23. L. Lo Piccolo *et al*., Loss of ISWI Function in Drosophila Nuclear Bodies Drives Cytoplasmic
Redistribution of Drosophila TDP-43. *Int J Mol Sci* **19**, 1082 (2018).

24. H. An *et al*., ALS-linked FUS mutations confer loss and gain of function in the nucleus by
promoting excessive formation of dysfunctional paraspeckles. *Acta Neuropathol Commun* **7**,
7 (2019).

25. Y. Lin, D. S. Protter, M. K. Rosen, R. Parker, Formation and Maturation of Phase-Separated
Liquid Droplets by RNA-Binding Proteins. *Mol Cell* **60**, 208-219 (2015).

26. M. Feric *et al*., Coexisting Liquid Phases Underlie Nucleolar Subcompartments. *Cell* **165**,
1686-1697 (2016).

27. S. F. Banani, H. O. Lee, A. A. Hyman, M. K. Rosen, Biomolecular condensates: organizers
of cellular biochemistry. *Nat Rev Mol Cell Biol* **18**, 285-298 (2017).

28. V. N. Uversky, Intrinsically disordered proteins in overcrowded milieu: Membrane-less
organelles, phase separation, and intrinsic disorder. *Curr Opin Struct Biol* **44**, 18-30 (2017).

29. S. Nakagawa, T. Yamazaki, T. Hirose, Molecular dissection of nuclear paraspeckles:
towards understanding the emerging world of the RNP milieu. *Open Biol* **8**, pii: 180150
(2018).

30. A. A. Hyman, C. A. Weber, F. Julicher, Liquid-liquid phase separation in biology. *Annu Rev
Cell Dev Biol* **30**, 39-58 (2014).

31. A. Molliex *et al*., Phase separation by low complexity domains promotes stress granule
assembly and drives pathological fibrillization. *Cell* **163**, 123-133 (2015).

32. H. B. Schmidt, R. Rohatgi, In Vivo Formation of Vacuolated Multi-phase Compartments

Lacking Membranes. *Cell Rep* **16**, 1228-1236 (2016).

33. I. R. Mackenzie *et al.*, TIA1 Mutations in Amyotrophic Lateral Sclerosis and Frontotemporal
Dementia Promote Phase Separation and Alter Stress Granule Dynamics. *Neuron* **95**,
808-816.e809 (2017).

34. V. H. Ryan *et al.*, Mechanistic View of hnRNPA2 Low-Complexity Domain Structure,
Interactions, and Phase Separation Altered by Mutation and Arginine Methylation. *Molecular*
Cell **69**, 465-479.e467 (2018).

35. L. Wang, J. Kang, L. Lim, Y. Wei, J. Song, TDP-43 NTD can be induced while CTD is
significantly enhanced by ssDNA to undergo liquid-liquid phase separation. *Biochem*
Biophys Res Commun **499**, 189-195 (2018).

36. S. Hennig *et al.*, Prion-like domains in RNA binding proteins are essential for building
subnuclear paraspeckles. *J Cell Biol* **210**, 529-539 (2015).

37. S. Xiang *et al.*, The LC Domain of hnRNPA2 Adopts Similar Conformations in Hydrogel
Polymers, Liquid-like Droplets, and Nuclei. *Cell* **163**, 829-839 (2015).

38. D. T. Murray *et al.*, Structure of FUS Protein Fibrils and Its Relevance to Self-Assembly and
Phase Separation of Low-Complexity Domains. *Cell* **171**, 615-627 e616 (2017).

39. O. A. Brady, P. Meng, Y. Zheng, Y. Mao, F. Hu, Regulation of TDP-43 aggregation by
phosphorylation and p62/SQSTM1. *J Neurochem* **116**, 248-259 (2011).

40. T. J. Cohen *et al.*, An acetylation switch controls TDP-43 function and aggregation
propensity. *Nat Commun* **6**, 5845 (2015).

41. W. Li *et al.*, Heat Shock-induced Phosphorylation of TAR DNA-binding Protein 43 (TDP-43)
by MAPK/ERK Kinase Regulates TDP-43 Function. *J Biol Chem* **292**, 5089-5100 (2017).

42. M. Hofweber *et al.*, Phase Separation of FUS Is Suppressed by Its Nuclear Import Receptor
and Arginine Methylation. *Cell* **173**, 706-719.e713 (2018).

43. F. Luo *et al.*, Atomic structures of FUS LC domain segments reveal bases for reversible
amyloid fibril formation. *Nat Struct Mol Biol* **25**, 341-346 (2018).

44. L. McGurk *et al.*, Poly(ADP-Ribose) Prevents Pathological Phase Separation of TDP-43 by
Promoting Liquid Demixing and Stress Granule Localization. Molecular *Cell* **71**,
703-717.e709 (2018).

45. S. Qamar *et al.*, FUS Phase Separation Is Modulated by a Molecular Chaperone and
Methylation of Arginine Cation-pi Interactions. *Cell* **173**, 720-734 e715 (2018).

46. Y. Duan *et al.*, PARylation regulates stress granule dynamics, phase separation, and
neurotoxicity of disease-related RNA-binding proteins. *Cell Res* **29**, 233-247 (2019).

47. D. Dominguez *et al.*, Sequence, Structure, and Context Preferences of Human RNA Binding
Proteins. *Mol Cell* **70**, 854-867 e859 (2018).

48. S. Namkoong, A. Ho, Y. M. Woo, H. Kwak, J. H. Lee, Systematic Characterization of
Stress-Induced RNA Granulation. *Mol Cell* **70**, 175-187 e178 (2018).

49. S. P. Shevtsov, M. Dundr, Nucleation of nuclear bodies by RNA. *Nat Cell Biol* **13**, 167-173
(2011).

50. S. Maharana *et al.*, RNA buffers the phase separation behavior of prion-like RNA binding
proteins. *Science* **360**, 918-921 (2018).

51. J. R. Mann *et al.*, RNA Binding Antagonizes Neurotoxic Phase Transitions of TDP-43.
Neuron **102**, 1-18 (2019).

52. L. Bernstam, J. Nriagu, Molecular aspects of arsenic stress. *J Toxicol Environ Health B Crit*
Rev **3**, 293-322 (2000).

53. L. A. Kominek, Cycloheximide Production by Streptomyces griseus Control Mechanisms of
Cycloheximide Biosynthesis. *Antimicrob Agents Chemother* **7**, 856-860 (1975).

54. F. Frottin *et al.*, The nucleolus functions as a phase-separated protein quality control
compartment. *Science* **365**, 342-347 (2019).

55. D. Glick, S. Barth, K. F. Macleod, Autophagy: cellular and molecular mechanisms. *J Pathol*
221, 3-12 (2010).

56. S. J. Barmada *et al.*, Autophagy induction enhances TDP43 turnover and survival in
neuronal ALS models. *Nature Chemical Biology* **10**, 677-685 (2014).

57. A. H. Brand, N. Perrimon, Targeted gene expression as a means of altering cell fates and
generating dominant phenotypes. *Development* **118**, 401-415 (1993).

58. A. C. Groth, M. Fish, R. Nusse, M. P. Calos, Construction of transgenic Drosophila by using
the site-specific integrase from phage PhiC31. *Genetics* **166**, 1775-1782 (2003).

59. J. R. Bateman, A. M. Lee, C. T. Wu, Site-specific transformation of Drosophila via phiC31
integrase-mediated cassette exchange. *Genetics* **173**, 769-777 (2006).

60. J. Bischof, R. K. Maeda, M. Hediger, F. Karch, K. Basler, An optimized transgenesis system
for Drosophila using germ-line-specific phiC31 integrases. *Proc Natl Acad Sci U S A* **104**,
3312-3317 (2007).

61. A. E. Conicella, G. H. Zerze, J. Mittal, N. L. Fawzi, ALS Mutations Disrupt Phase Separation
Mediated by alpha-Helical Structure in the TDP-43 Low-Complexity C-Terminal Domain.
Structure **24**, 1537-1549 (2016).

62. L. Guo, J. Shorter, It's Raining Liquids: RNA Tunes Viscoelasticity and Dynamics of
Membraneless Organelles. *Mol Cell* **60**, 189-192 (2015).

63. J. R. Tollervey *et al.*, Characterizing the RNA targets and position-dependent splicing
regulation by TDP-43. *Nat Neurosci* **14**, 452-458 (2011).

64. C. M. Clemson *et al.*, An architectural role for a nuclear noncoding RNA: NEAT1 RNA is

essential for the structure of paraspeckles. *Mol Cell* **33**, 717-726 (2009).

65. L. L. Chen, G. G. Carmichael, Altered nuclear retention of mRNAs containing inverted
repeats in human embryonic stem cells: functional role of a nuclear noncoding RNA. *Mol*
Cell **35**, 467-478 (2009).

66. Y. T. Sasaki, T. Ideue, M. Sano, T. Mituyama, T. Hirose, MENepsilon/beta noncoding RNAs
are essential for structural integrity of nuclear paraspeckles. *Proc Natl Acad Sci U S A* **106**,
2525-2530 (2009).

67. T. Yamazaki *et al.*, Functional Domains of NEAT1 Architectural lncRNA Induce Paraspeckle
Assembly through Phase Separation. *Molecular Cell* **70**, 1038-1053.e1037 (2018).

68. K. K. McDonald *et al.*, TAR DNA-binding protein 43 (TDP-43) regulates stress granule
dynamics via differential regulation of G3BP and TIA-1. *Hum Mol Genet* **20**, 1400-1410
(2011).

69. E. Bentmann *et al.*, Requirements for stress granule recruitment of fused in sarcoma (FUS)
and TAR DNA-binding protein of 43 kDa (TDP-43). *J Biol Chem* **287**, 23079-23094 (2012).

70. J. A. Austin *et al.*, Disease causing mutants of TDP-43 nucleic acid binding domains are
resistant to aggregation and have increased stability and half-life. *Proc Natl Acad Sci U S A*
111, 4309-4314 (2014).

71. C. H. Chiang *et al.*, Structural analysis of disease-related TDP-43 D169G mutation: linking
enhanced stability and caspase cleavage efficiency to protein accumulation. *Sci Rep* **6**,
21581 (2016).

72. Y. C. Liu, P. M. Chiang, K. J. Tsai, Disease animal models of TDP-43 proteinopathy and
their pre-clinical applications. *Int J Mol Sci* **14**, 20079-20111 (2013).

73. A. H. Fox *et al.*, Paraspeckles: A Novel Nuclear Domain. *Current Biology* **12**, 13-25 (2002)

74. K. V. Prasanth *et al.*, Regulating gene expression through RNA nuclear retention. *Cell* **123**,
249-263 (2005).

75. T. Naganuma *et al.*, Alternative 3'-end processing of long noncoding RNA initiates
construction of nuclear paraspeckles. *EMBO J* **31**, 4020-4034 (2012).

76. C. S. Bond, A. H. Fox, Paraspeckles: nuclear bodies built on long noncoding RNA. *J Cell*
Biol **186**, 637-644 (2009).

77. A. H. Fox, A. I. Lamond, Paraspeckles. *Cold Spring Harb Perspect Biol* **2**, a000687 (2010).

78. A. H. Fox, S. Nakagawa, T. Hirose, C. S. Bond, Paraspeckles: Where Long Noncoding RNA
Meets Phase Separation. *Trends Biochem Sci* **43**, 124-135 (2018).

79. L. Liu-Yesucevitz *et al.*, Tar DNA binding protein-43 (TDP-43) associates with stress
granules: analysis of cultured cells and pathological brain tissue. *PLoS One* **5**, e13250
(2010).

80. J. K. Bose, I. F. Wang, L. Hung, W. Y. Tarn, C. K. Shen, TDP-43 overexpression enhances
exon 7 inclusion during the survival of motor neuron pre-mRNA splicing. *J Biol Chem* **283**,
28852-28859 (2008).

81. T. W. Han *et al*., Cell-free formation of RNA granules: bound RNAs identify features and
components of cellular assemblies. *Cell* **149**, 768-779 (2012).

82. J. Seo *et al*., Oxidative Stress Triggers Body-Wide Skipping of Multiple Exons of the Spinal
Muscular Atrophy Gene. *PLoS One* **11**, e0154390 (2016).

83. M. S. Forman, J. Q. Trojanowski, V. M. Lee, TDP-43: a novel neurodegenerative
proteinopathy. *Curr Opin Neurobiol* **17**, 548-555 (2007).

84. E. Buratti, F. E. Baralle, Characterization and functional implications of the RNA binding
properties of nuclear factor TDP-43, a novel splicing regulator of CFTR exon 9. *J Biol Chem*
276, 36337-36343 (2001).

85. P. H. Kuo, C. H. Chiang, Y. T. Wang, L. G. Doudeva, H. S. Yuan, The crystal structure of
TDP-43 RRM1-DNA complex reveals the specific recognition for UG- and TG-rich nucleic
acids. *Nucleic Acids Res* **42**, 4712-4722 (2014).

86. M. M. Fay, P. J. Anderson, P. Ivanov, ALS/FTD-Associated C9ORF72 Repeat RNA
Promotes Phase Transitions In Vitro and in Cells. *Cell Rep* **21**, 3573-3584 (2017).

87. T. Ukmar-Godec *et al*., Lysine/RNA-interactions drive and regulate biomolecular
condensation. *Nat Commun* **10**, 2909 (2019).

88. C. Jolly et al., Stress-induced transcription of satellite III repeats. *J Cell Biol* **164**, 25-33
(2004).

89. R. Valgardsdottir et al., Structural and Functional Characterization of Noncoding Repetitive
RNAs Transcribed in Stressed Human Cells. *Molecular Biology of the Cell* **16**, 2597-2604
(2005).

90. N. Puthiyedth, C. Riveros, R. Berretta, P. Moscato, Identification of Differentially Expressed
Genes through Integrated Study of Alzheimer's Disease Affected Brain Regions. *PLoS One*
11, e0152342 (2016).

91. E. Mariani *et al*., Meta-Analysis of Parkinson's Disease Transcriptome Data Using TRAM
Software: Whole Substantia Nigra Tissue and Single Dopamine Neuron Differential Gene
Expression. *PLoS One* **11**, e0161567 (2016).

92. C. Cheng *et al*., The long non-coding RNA NEAT1 is elevated in polyglutamine repeat
expansion diseases and protects from disease gene-dependent toxicities. *Hum Mol Genet*
27, 4303-4314 (2018).

93. L. Schirmer *et al*., Neuronal vulnerability and multilineage diversity in multiple sclerosis.
Nature [Epub ahead of print] (2019).

94. M. Polymenidou *et al*., Long pre-mRNA depletion and RNA missplicing contribute to

neuronal vulnerability from loss of TDP-43. *Nat Neurosci* **14**, 459-468 (2011).

95. C. Y. Chung *et al.*, Aberrant activation of non-coding RNA targets of transcriptional
elongation complexes contributes to TDP-43 toxicity. *Nat Commun* **9**, 4406 (2018).

96. M. Modic *et al.*, Cross-Regulation between TDP-43 and Paraspeckles Promotes
Pluripotency-Differentiation Transition. *Mol Cell* **74**, 951-965 e913 (2019).

97. A. S. Chen-Plotkin, V. M. Lee, J. Q. Trojanowski, TAR DNA-binding protein 43 in
neurodegenerative disease. *Nat Rev Neurol* **6**, 211-220 (2010).

98. S. H. Kim *et al.*, Potentiation of amyotrophic lateral sclerosis (ALS)-associated TDP-43
aggregation by the proteasome-targeting factor, ubiquilin 1. *J Biol Chem* **284**, 8083-8092
(2009).

99. X. Sun *et al.*, Distinct multilevel misregulations of Parkin and PINK1 revealed in cell and
animal models of TDP-43 proteinopathy. *Cell Death Dis* **9**, 953 (2018).

100.X. Cao *et al.*, In vivo imaging reveals mitophagy independence in the maintenance of axonal
mitochondria during normal aging. *Aging Cell* **16**, 1180-1190 (2017).

101.M. H. M. Koken *et al.*, The t(15;17) translocation alters a nuclear body in a retinoic
acid-reversible fashion. *The EMBO Journal* **13**, 1073-1083 (1994).

102.N. Santama *et al.*, Characterization of hCINAP, a novel coilin-interacting protein encoded by
a transcript from the transcription factor TAFIID32 locus. *J Biol Chem* **280**, 36429-36441
(2005).

103.X-D. Fu, T. Maniatis, Factor required for mammalian spliceosome assembly is localized to
discrete regions in the nucleus. *Nature* **343**, 437–444 (1990).

ACKNOWLEDGMENTS

We thank the BDSC for providing the fly strains, the SIBCB *Drosophila* Center for the fly embryo injection services, L. Pan, J. Zhou, and Z. Zhang for the cloning vectors and plasmids, Z. Zhang for help in the pulse-chase assay, S. Qiu for technical supports, J. Yuan, A. Li and Y. Chen for critical reading of the manuscript, and the members of the Fang lab for helpful discussion. **Funding**: Supported by grants from the National Key R&D Program of China (No. 2016YFA0501902) to Y.F., the National Natural Science Foundation of China (No. 81671254 and No. 31471017) to Y.F and (No. 91853112 and No. 31470748) to C.L, and the Alzheimer's AssOCiation (AARF-16-441196) and Target ALS Springboard Fellowship to L.G. **Author contributions:** G.D., Z.M. and Y.F. conceived the research; C.W., Y.D., G.D., C.L., L.G. and Y.F. designed the experiments; C.W., Y.D. G.D., K.Z., X.D., B.Q., J.G. and S.Z. performed the experiments; C.W., Y.D., G.D., Z.M., J.G. and C.L contributed important new reagents; C.W., Y.D. G.D., and Y.F. analyzed the data and interpreted the results; C.W., Y.D., G.D. and Y.F. prepared the figures; and C.W., Y.D. G.D., and Y.F. wrote the paper. All authors read and approved the final manuscript. **Competing interests:** None. **Data and materials availability:** All data are available in the manuscript or the supplementary materials.

SUPPLEMENTARY INFORMATION

Online Methods

Figures S1-S7

Movies S1-S2

References (99-103)

FIGURES AND FIGURE LEGENDS

Figure 1. TDP-43 forms reversible, dynamic NBs in response to arsenic stress.

(**A**) Representative confocal images of HeLa cells in the absence (PBS) or presence of arsenite

(250 μM, 30 min) forming TDP-43 NBs (arrows) and TDP-43$^+$ SGs (arrowheads) (also see Movie

S1). All cells are transfected with TDP-43-HA (anti-HA) and stained for SGs with anti-TIAR;

merged images with DAPI staining for DNA (blue) are shown. Arrows, TDP-43 NBs; arrowheads,

SGs associated with TDP-43 (TDP-43$^+$ SGs). (**B**) Quantification of the percentage of cells that

show TDP-43 NBs or TDP-43$^+$ SGs before and after arsenite the treatment in (A). Data are

shown as mean ± SEM, n = ~100 cells for each group from pooled results of 3 independent

repeats; *p < 0.05, **p < 0.01, ***p < 0.001; Student's t-test for comparison between stress and

no stress within the same group, two-way ANOVA for comparison of the stress-induced changes

between different groups. (**C**) Representative images of the FRAP analysis of GFP-TDP-43 NBs

in *live* cells (also see Movie S2). (**D**) The FRAP recovery curve is quantified by averaging the

relative fluorescence intensity (FI) of 9 TDP-43 NBs of similar size at indicated time after

photobleaching. The relative FI of each TDP-43 NB prior to photobleaching is set to 100%. Time

0 refers to the time point right after photobleaching. (**E**) Representative images showing the

disappearance of TDP-43 NBs and TDP-43$^+$ SGs after arsenite washout. (**F-H**) Quantification of

the percentage of cells showing TDP-43 NBs (F), TDP-43$^+$ SGs (G) or the average count of

TDP-43 NBs per cell (H) at indicated time points after arsenite washout in (E). Mean ± SEM,

except for (H) box-and-whisker plots. n = ~100 cells for each group in (F-G) and ~15 cells for

each group in (H), pooled results from 3 independent repeats; statistic significance is determined

by comparing to time 0; *p < 0.05, **p < 0.01, ***p < 0.001; one-way ANOVA. Scale bars, 10 μm

in (A) and 5 μm in (C, E).

Figure 2. Inhibition of nuclear export induces TDP-43 NBs.

(**A-B**) Representative images of endogenous TDP-43 (endo-TDP-43) forming NBs upon LMB

treatment (25 nM, 24 h). Percentage of cells showing distinct TDP-43 NBs in (A) is quantified in

(B). (**C**) A diagram showing the major domains of human TDP-43 and the residues mutated in

the localization signals. NLS, nuclear localization signal; RRM, RNA recognition motif; NES,

nuclear export signal; LCD, low complexity domain. (**D**) Representative images of 293T cells

transfected with WT, NLS^{mut} or NES^{mut} TDP-43. (**E-F**) The percentage of cells with NBs (E) and

the average number of NBs per cell (F) are quantified. (**G-H**) Representative Western blot

images (G) and quantifications (H) of WT, NLS^{mut} and NES^{mut} TDP-43 proteins in the soluble (S,

supernatants in RIPA) and insoluble fractions (I, pellets resuspended in 9 M of urea). Mean ±

SEM, except for (F) box-and-whisker plots. n >120 cells each condition in (B, E) and ~ 10 cells

each condition in (F) from pooled results of 3 independent repeats, n = 3 in (G-H); *p < 0.05, **p

< 0.01, ***p < 0.001; ns, not significant; Student's t-test in (B), one-way ANOVA in (E-F, H). Scale

bar: 5 μm.

Figure 3_CWang

Figure 3. Formation of NBs mitigates TDP-43-mediated cytotoxicity.

(**A**) OE of the WT or NLS^{mut} but not NES^{mut} TDP-43 in 293T cells exhibits cytotoxicity and

decreases the cell viability, which is measured by the Cell Counting Kit-8 (CCK-8) assay. (**B-C**)

Diagrams showing the K145/192Q mutation in the WT TDP-43 and the K145/192R mutation in

the NES^{mut} TDP-43, respectively. (**D-E'**) Representative confocal images of the WT (D) and "WT

+ K145/192Q" TDP-43 (D'), or the NES^{mut} (E) and "NES^{mut} + K145/192R" TDP-43 (E') in 293T

cells. (**F-G**) Quantifications of the average numbers of TDP-43 NBs per cell formed in each group

in (D-E') as indicated. (**H-I**) Percentages of cells with TDP-43 exclusively in the nucleus (nuclear)

or in both the nucleus and the cytoplasm (cyto) in (D-E') are quantified. (**J-K**) The viability of cells

expressing the WT or NES^{mut} TDP-43 with the indicated NB-promoting or NB-suppressing

mutation is assessed using the CCK-8 assay. Mean ± SEM, except for (F-G) box-and-whisker

plots; n = 3 in (A, J-K), n = ~12 cells each group in (F-G), n >100 cells per group in (H-I). *p <

0.05, **p < 0.01, ***p < 0.001; ns, not significant; one-way ANOVA in (A), Student's *t*-test in (F-G,

J-K), two-way ANOVA in (H-I). Scale bar: 5 μm.

Figure 4_CWang

Figure 4. Transgenic flies expressing WT, NLS^mut or NES^mut hTDP-43.

(**A**) Representative confocal images of the fly central brain neurons (*elav*-Gal4) expressing the

WT or mutant *hTDP-43* as indicated. The whole-mount fly brains were dissected and

immunostained for hTDP-43 (green) and DAPI (blue, to show the nucleus). (**B**) The average

percentages of fly neurons with or without TDP-43 NBs are quantified. Mean ± SEM, n = ~100

neurons each group. (**C-D**) Representative image (C) and quantification (D) of Western blot

analysis of the WT or mutant hTDP-43 protein extracted from the fly head in RIPA-soluble (S) or

RIPA-insoluble (I, resuspended in 9 M of urea) fractions. All protein levels are normalized to

GAPDH. The relative level of soluble TDP-43-WT is set to 100%. Mean ± SEM, n = 4. (**E**)

Representative z-stack images of the fly eye (GMR-Gal4) expressing the WT or mutant hTDP-43

transgenes as indicated at Day 5 or Day 25. The average degeneration score (mean ± SEM) and

the statistic significance (compared to hTDP-43-WT flies) of each group are indicated at the

bottom. n = 8 eyes each group. (**F**) The climbing capability of the flies expressing the WT or

mutant hTDP-43 in motor neurons (D42-Gal4) is evaluated as percentage of flies climbing over 3

cm in 10 seconds. Control, UAS-*luciferase*. Mean ± SEM, the number of flies tested for each

genotype is shown. *$p < 0.05$, ***$p < 0.001$; ns, not significant; two-way ANOVA in (B, F),

one-way ANOVA in (D-E). Scale bars: 2 μm in (A) and 100 μm in (E).

Figure 5_CWang

Figure 5. The role of RRM1 and RRM2 in maintaining the architecture of TDP-43 NBs.

(**A**) A diagram showing the major functional domains of WT TDP-43 and the ΔRRM1 and

ΔRRM2 mutants. (**B**) LIGHTNING super-resolution microscopy of 293T cells transfected with

Myc-TDP-43-WT, ΔRRM1 or ΔRRM2 and treated with LMB. Higher magnification images

(zoom-ins) and 3D renderings of the boxed areas are shown. (**C**) The average diameters of the

WT, ΔRRM1 and ΔRRM2 TDP-43 nuclear structures in (B) are quantified. (**D**) A schematic

model of the core-shell architecture of TDP-43 NBs (red) and the distinct distribution of ΔRRM1

(green) and ΔRRM2 (blue). (**E-H**) LIGHTNING multicolor super-resolution microscopy and higher

magnification images of representative LMB-induced TDP-43 NBs. Cells are co-transfected with

a combination of HA- or Myc-tagged WT, ΔRRM1 or ΔRRM2 TDP-43 as indicated. (**I-P**) The

intensity profiles (I-L) along the indicated lines and the average diameters (M-P) of TDP-43 NBs

in each group are shown. The box-and-whisker plots with the value of each NB measured are

shown. n = ~900 NBs in (C, M-P) for each group from pooled results of at least 3 independent

repeats; $^{\#}p$ < 0.0001; one-way ANOVA in (C) and Student's *t*-test in (M-P). Scale bars: 5 μm in

(B), 2 μm in (E-H) and 200 nm in the zoom-ins.

Figure 6_CWang

Figure 6. The effects of different RNAs on TDP-43 LLPS *in vitro*.

(**A**) A diagram showing the WT, ΔRRM1 and ΔRRM2 of TDP-43$^{1-274}$. (**B-D**) The total RNA

extracts, (**E-G**) tRNA, and (**H-J**) lncRNA *NEAT1* are added into the *in vitro* LLPS assay of WT,

ΔRRM1 and ΔRRM2 TDP-43$^{1-273}$ as indicated. C', a higher concentration of ΔRRM1 is also

included in order to start the suppression assay with large LDs as those of WT TDP-43. (**K-M**)

tRNA-mediated suppression of TDP-43 LLPS can be antagonized by increasing *NEAT1* RNA

levels. The concentrations of NaCl, the crowding agent PEG, TDP-43 proteins and RNAs used in

the *in vitro* assays are as indicated. (**N-O**) Cellular stress induced by arsenite (N) or LMB (O)

increases *NEAT1* RNA levels in mouse primary neurons compared to the vehicle control PBS

and ethanol, respectively. Mean ± SEM, n = 3; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$; Student's *t*-test.

Scale bars: 2 μm.

Figure 7_CWang

Figure 7. The ALS-associated D169G mutation impairs the ability of TDP-43 to form

stress-induced NBs and *NEAT1*-promoted LDs.

(**A**) A diagram showing WT and D169G mutant TDP-43. (**B**) Representative confocal images of

HeLa cells transfected with WT or D169G TDP-43 in the absence (PBS) or presence of arsenite.

Green, TDP-43-HA (anti-HA); Red, anti-TIAR for SGs; Blue, DAPI staining for DNA (nucleus).

Arrows, TDP-43 NBs; arrowheads, TDP-43[+] SGs. (**C-E**) The percentage of cells with TDP-43

NBs (C), the average count of TDP-43 NBs per cell (D) and the percentage of cells with TDP-43[+]

SGs (E) in (B) are quantified. Mean ± SEM, except for (D) box-and-whisker plots; n = ~100 cells

for each group in (C, E) and n = ~15 cells for each group in (D) from pooled results of 3

independent repeats; ***$p < 0.001$; ns, not significant; Student's *t*-test. (**F**) Coomassie blue

staining confirming the purified WT and D169G TDP-43$^{1-274}$ proteins used in the *in vitro* LLPS

assays. (**G-H**) Both WT and D169G TDP-43 form LDs by LLPS in a dose-dependent manner.

(**I-J**) Suppression of the *in vitro* LLPS of WT and D169G TDP-43 by total RNA extracts. (**K-L**)

Promotion of TDP-43 LLPS by *NEAT1* RNA is dramatically reduced in the D169G TDP-43. The

concentrations of NaCl, TDP-43, PEG and RNA used in the *in vitro* assays are as indicated.

Scale bars: 5 μm in (B) and 2 μm in (G-L).

Supplementary Information for

4 Stress induces cytoprotective TDP-43 nuclear bodies through lncRNA

5 *NEAT1*-promoted phase separation

6 Chen Wang, Yongjia Duan, Gang Duan, Zhiwei Ma, Kai Zhang, Xue Deng, Beituo Qian,

7 Jinge Gu, Shuang Zhang, Lin Guo, Cong Liu, and Yanshan Fang

8 Correspondence to: fangys@sioc.ac.cn

11 This PDF file includes:

12   Online Methods

13   Figure S1 to S7

14   Captions for Movies S1 to S2

16 Other Supplementary Data for this manuscript include the following:

17   Movies S1 to S2

 Online Methods

Plasmids and constructs

The pcDNA3.1-TDP-43-HA and pCMV-myc-TDP-43 plasmids were as previously described (*99*)

and used as the templates to generate the following plasmids. The pcDNA3.1-TDP-43-NES^{mut},

NLS^{mut}, D169G and M337V-HA plasmids were generated by site-directed mutagenesis using the

Fast Mutagenesis Kit II (Vazyme). The pcDNA3.1-TDP-43-K145/192Q-HA and

pcDNA3.1-TDP-43-NES^{mut}-K145/192R-HA plasmids were generated by site-directed

mutagenesis using the Fast Mutagenesis Kit MultiS (Vazyme). The pCMV-myc-TDP-43$^{1-274}$

pCMV-myc-TDP-43-ΔRRM1, and pCMV-myc-TDP-43-ΔRRM2 plasmids were generated by

PCR using the pCMV-myc-TDP-43 plasmid as a template. To generate the pCAG-GFP-TDP-43

plasmid, the GFP coding sequence was amplified from pcDNA3.1-GFP-AXR3 (a gift from Dr. Z.

Zhang) by PCR and then subcloned into a pCAG-hTDP-43 plasmid (*99*) using the ClonExpress

MultiS One Step Cloning Kit (Vazyme).

For lentivirus infection of mouse primary neurons, the pLenti-hSyn-TDP-43-HA plasmid was

generated by PCR using the pcDNA3.1-TDP-43-HA as a template and sub-cloned into the

pLenti-hSyn vector (a gift from Dr. Y. Chen).

For generation of transgenic fly lines, the WT and mutant UAS-hTDP-43 constructs were

generated by PCR using the above plasmids as a template and sub-cloned into the

pBID-UASC-6*Myc vector (100) between the KpnI and ApaI sites.

For *Escherichia coli* (*E. coli*) expression, the pET28a-6×His-sumo-TDP-43,

sumo-TDP-43-$^{1-274}$, His-TDP-43$^{1-274}$, His-TDP-43$^{1-274}$-ΔRRM1, His-TDP-43$^{1-274}$-ΔRRM2 and

His-TDP-43$^{1-274}$-D169G constructs were generated by PCR using the above WT or mutant

TDP-43 plasmid as a template and were sub-cloned into a pET28a-6×His (a gift from Dr. L. Pan)

or pET28a-6×His-sumo (a gift from Dr. J. Zhou) vector.

For *in vitro* transcription of NEAT1 RNA, the pcDNA3.1-NEAT1 was generated by PCR

using the pEGFP-C1-mNEAT1 plasmid (a gift from Dr. L. Chen) as a template and sub-cloned

into the pcDNA3.1 vector.

The primers used for PCR to generate the expression plasmids are summarized below. All

constructs were verified by sequencing to ensure the integrity of the cloned open reading frames.

pcDNA3.1-TDP-43-HA:

5'- CGTTTAAACGGGCCCTCTAGAGCCACCATGTCTGAATATATTCGG -3'

5'-CACAGTGGCGGCCGCCTAAGCGTAGTCTGGGACGTCGTATGGGTACATTCCCCAG

CCAGAAG -3'

pcDNA3.1-TDP-43-NLS^{mut}-HA:

5'- TAACGCAGCAGCAATGGATGAGACAGATGCTTCATCA -3

5'- CCATTGCTGCTGCGTTATCTTTTGGATAGTTGACAACATACA -3

pcDNA3.1-TDP-43-NES^{mut}-HA:

5'- GGCAGCGCAGTCTGCATGTGGAGAGGACTTGATCATTAAAGG -3

5'- ATGCAGACTGCGCTGCCTGATCATCTGCAAATGTAACAAAGG -3

pLenti-hSyn-TDP-43-HA:

5'- AGAGCGCAGTCGAGAGGATCCGCCACCATGTCTGAATATATTCGG -3'

5'- GATAAGCTTGATATCGAATTCTCATTAAGCGTAGTCTGGGACGTCG -3'

pBID-UASC-TDP-43-Myc:

5'- CGGCCGCGGCTCGAGGGTACCATGTCTGAATATATTCGGGTAACCG -3'

5'- GAGTTTTTGTTCGAAGGGCCCTCTAGACTCGAGCATTCCCCAGCCAGAAGACTT-3'

pBID-UASC-TDP-43-NLS^{mut}-Myc:

5'- CGGCCGCGGCTCGAGGGTACCATGTCTGAATATATTCGGGTAACCG -3'

5'- GAGTTTTTGTTCGAAGGGCCCTCTAGACTCGAGCATTCCCCAGCCAGAAGACTT-3'

pBID-UASC-TDP-43-NES^{mut}-Myc:

5'- CGGCCGCGGCTCGAGGGTACCATGTCTGAATATATTCGGGTAACCG -3'

5'- GAGTTTTTGTTCGAAGGGCCCTCTAGACTCGAGCATTCCCCAGCCAGAAGACTT-3'

pcDNA3.1-TDP-43-K145/192Q-HA:

5'- TCATTCACAGGGGTTTGGCTTTGTTCGTTTT -3

5'- CAAACCCCTGTGAATGACCAGTCTTAAGATCTTTCTTG -3

5'- GCAGACAAGTGTTTGTGGGGCGCTGTAC -3

5'- CACAAACACTTGTCTGCTTCTCAAAGGCTCATCTT -3

pcDNA3.1-TDP-43-NES^{mut}-K145/192R-HA:

5'- TCATTCAAGGGGGTTTGGCTTTGTTCGTTTT -3

5'- CAAACCCCCTTGAATGACCAGTCTTAAGATCTTTCTTG -3

5'- GCAGAAGAGTGTTTGTGGGGCGCTGTAC -3

5'- CACAAACACTCTTCTGCTTCTCAAAGGCTCATCTT -3

pcDNA3.1-TDP-43-D169G-HA:

5'- ATGATAGGTGGACGATGGTGTGACTGCAAACT -3'

5'- CATCGTCCACCTATCATATGTCGCTGTGACATTACTTTC -3'

pcDNA3.1-TDP-43-M337V-HA:

5'- GGTATGGTGGGCATGTTAGCCAGCCAGCAGAA -3'

5'- AACATGCCCACCATACCCCAACTGCTCTGTAGTGC -3'

pCMV-Myc-TDP-43:

5'- ATGGCCATGGAGGCCCGAATTCATGTCTGAATATATT-3'

5'- CCGCGGCCGCGGTACCTCGAGCTACATTCCCCAGCCAGAAGAC -3'

pCMV-Myc-TDP-43$^{1–274}$:

5'- ATGGCCATGGAGGCCCGAATTCATGTCTGAATATATT -3'

5'- CCGCGGCCGCGGTACCTCGAGCTATCCACTTCTTTCTAACTGTCTATTGC -3'

pCMV-Myc-TDP-43-ΔRRM1:

5'- ATGGCCATGGAGGCCCGAATTCATGTCTGAATATATT -3'

5'- CCGCGGCCGCGGTACCTCGAGCTACATTCCCCAGCCAGAAGAC -3'

pCMV-Myc-TDP-43-ΔRRM2:

5'- ATGGCCATGGAGGCCCGAATTCATGTCTGAATATATT -3'

5'- CCGCGGCCGCGGTACCTCGAGCTACATTCCCCAGCCAGAAGAC -3'

pCAG-GFP-TDP-43:

5'- CATCATTTTGGCAAAGAATTCCACCATGGTGAGCAAGGGCGAGG -3'

5'- CAGACATGCTTCCGCCCTTGTACAGCTCGTCCATGCC -3'

5'- CGAGCTGTACAAGGGCGGAAGCATGTCTGAATATATTCGGGTAACCG -3'

5'- GCTCCCCGGGGGTACCTCGAGCTATTACATTCCCCAGCCAGAAGACTT -3'

pET28a-6×His-TDP-43$^{1–274}$:

5'- CAGCAAATGGGTCGCGGATCCATGTCTGAATATATTCGGGTAACCG -3'

5'- GTGGTGGTGGTGGTGCTCGAGTTAACTTCTTTCTAACTGTCTATTGCTATTG -3'

pET28a-6×His-TDP-43$^{1–274}$-ΔRRM1:

5'- CAGCAAATGGGTCGCGGATCCATGTCTGAATATATTCGGGTAACCG -3'

5'- GTGGTGGTGGTGGTGCTCGAGTTAACTTCTTTCTAACTGTCTATTGCTATTG -3'

pET28a-6×His-TDP-43$^{1–274}$-ΔRRM2:

5'- CAGCAAATGGGTCGCGGATCCATGTCTGAATATATTCGGGTAACCG -3'

5'- GTGGTGGTGGTGGTGCTCGAGTTAACTTCTTTCTAACTGTCTATTGCTATTG -3'

pET28a-6×His-TDP-43$^{1–274}$-D169G:

5'- CAGCAAATGGGTCGCGGATCCATGTCTGAATATATTCGGGTAACCG -3'

5'- GTGGTGGTGGTGGTGCTCGAGTTAACTTCTTTCTAACTGTCTATTGCTATTG -3'

pET28a-6×His-sumo-TDP-43:

5'- AGAGAACAGATTGGTGGATCCATGTCTGAATATATTCGGGTAACCG -3'

5'-TTGTCGACGGAGCTCGAATTCTTACATTCCCCAGCCAGAAGAC -3'

pET28a-6×His-sumo-TDP-43-$^{1–274}$:

5'- AGAGAACAGATTGGTGGATCCATGTCTGAATATATTCGGGTAACCG -3'

5'- TTGTCGACGGAGCTCGAATTCTTATCCACTTCTTTCTAACTGTCTATTGC -3'

pcDNA3.1-NEAT1:

5'- CGTTTAAACGGGCCCTCTAGAGAGTTAGTGACAAGGAGGGCTCG -3'

5'- ATATCCAGCACAGTGGCGGCCGCTTCAATCTCAAACCTTTATTTTGCTG-3'

Cell cultures and transfection

293T and HeLa cells were cultured in Dulbecco's Modified Eagle Medium (Sigma, D0819)

supplemented with 10% (v/v) fetal bovine serum (FBS, BioWest) and 1% penicillin/streptomycin

at 37°C in 5% CO_2. Transient transfection was performed using Lipofectamine™ 3000

(Invitrogen) in Opti-MEM (Invitrogen). Cells were transfected for at least 24h before the

subsequent drug treatments or examinations.

Pharmacological experiments:

Arsenite treatment: HeLa or 293T cells were grown on coverslips in a 24-well plate and

transfected with the indicated plasmids for 24 h. Cells were then treated with 250 µM of $NaAsO_2$

or PBS for 30 min, prior to fixation with 4% paraformaldehyde. For the recovery experiments, the

culture medium containing $NaAsO_2$ was removed and the cells were incubated in fresh medium

for indicated time prior to fixation.

LMB treatment: For the nuclear export inhibition assays, LMB was added into the culture

medium after 6h transfection at indicated final concentrations.

CHX treatment: For the pulse-chase assays, CHX was added into the medium at each time

point at a final concentration of 25 ng/ml.

MG132 treatment: For the proteasomal inhibition assays, MG132 was added into the

medium at each time at a final concentration of 25 µM.

CQ treatment: For autophagy inhibition assays, CQ was added into the medium at each time

at a final concentration of 25 mM.

The cells incubated in the culture medium with the above drugs for indicated time before

fixation for immunocytochemistry or western blotting analysis.

Immunocytochemistry and immunohistochemistry assays

HeLa or 293T cells grown on coverslips pre-coated with PLL (Sigma) in a 24-well plate were

transfected and treated as described above. The cells were then fixed in 4% paraformaldehyde

in PBS for 15 min at room temperature (RT), permeabilized in 0.5% Triton X-100 (Sigma) in PBS

for 15 min and blocked with 3% goat serum in PBST (0.1% Triton X-100 in PBS) for 1h at RT.

For immunostaining of the whole-mount *Drosophila* brain, fly brains were dissected on ice

and immediately fixed in 4% paraformaldehyde in PBS for 30 min at RT and permeabilized with

0.5% Triton X-100 in PBS overnight at 4°C overnight.

The above primary and secondary antibodies were then incubated in the blocking buffer at

4°C overnight or at RT for 1-2 h. After 3 washes with PBST, cells were mounted on glass slides

using the VECTASHIELD Antifade Mounting Medium with DAPI (Vector Laboratories).

Confocal and super-resolution imaging

Fluorescent confocal images were taken with Leica TCS SP8 confocal microscopy system using

a 63X or 100X oil objective (NA=1.4). Super-resolution images were captured using the Leica

SP8 LIGHTNING confocal microscope, which allowed simultaneous multicolor imaging in

super-resolution down to 120 nm. Confocal or super-resolution images were then processed in

LAS X (Leica) and assembled into figures using Adobe Photoshop CS6.

Live cell imaging

HeLa cells transfected with pCAG-GFP-TDP-43 were grown on Nunc™ Lab-Tek™ Chambered

Coverglass (Thermo) for 24 h before the cells were treated with 250 µM of $NaAsO_2$. For live cell

imaging, the chambered coverglasses were placed in the Incubation System for Microscopes

(Tokai Hit) and maintained at 37°C in 5% CO_2 for the duration of the experiment. Time-lapse

images were captured using the Leica TCS SP8 confocal microscopy system every 10 min for

about 1 h.

Fluorescence recovery after photobleaching (FRAP) assay

The PFRP assay was performed using the FRAP module of the Leica SP8 confocal microscopy

system. In brief, each GFP-TDP-43 NB was bleached using a 488 nm laser at 100% laser power

for approximately 5 s. After photobleaching, time-lapse images were captured every 10 s for the

about 5 min. For each indicated time point (t), the fluorescence intensity within the bleached NB

was normalized to the fluorescence intensity of a nearby, unbleached NB (to control for

photobleaching during prolonged live imaging). The normalized fluorescence intensity of

pre-bleaching was set to 100% and the normalized fluorescence intensity at each time point (I_t)

was used to calculate the fluorescence recovery according to the following formula: FR(t) =

$I_t/I_{pre-bleaching}$. Image J was used for quantification and GraphPad Prism to plot and analyze the

FRAP experiments.

Antibodies

The following antibodies were used for Western blotting, immunoprecipitation and

immunofluorescence assays: mouse anti-FLAG (Sigma, F3165), mouse anti-HA (Proteintech,

66006-1), rabbit anti-c-Myc (Sigma, c3956), rabbit anti-TDP-43 (Proteintech, 10782-2-AP),

anti-β-Tubulin III (Sigma, T2200), rabbit anti-TIAR (Cell Signaling Technology, 8509S), rabbit

anti-SC35 (Abcam, ab204916), rabbit anti-SFPQ (Abcam, ab177149), rabbit anti-TAF9 (Abcam,

ab169784), rabbit anti-PML (Abcam, ab179466), and chicken anti-MAP2 (Abcam, ab5392). HRP

conjugated secondary antibodies: goat anti-mouse (Sigma, A4416) and goat anti-rabbit (Sigma,

A9169). Fluorescent secondary antibodies: goat anti-mouse-Alexa Fluor 488 (Life Technologies,

A11029), goat anti-rabbit-Alexa Fluor 568 (Life Technologies, A11036) and goat

anti-Chicken-Alexa Fluor 568 (Life Technologies, A11041).

Protein extraction and Western blotting

Total protein was extracted from cells in a 2% SDS extraction buffer (50 mM Tris pH 6.8, 2%

SDS, 1% mercaptoethanol, 12.5% glycerol and 0.04% bromophenol blue) containing the

protease inhibitor cocktail (Roche, 04693132001). For separation of soluble and insoluble

proteins, cells or fly heads were lysed on ice in a RIPA buffer (50 mM Tris pH 8.0, 150 mM NaCl,

1% NP-40, 5 mM EDTA, 0.5% sodium deoxycholate, 0.1% SDS) supplemented with protease

and phosphatase inhibitors (Roche). After sonication, the homogenates were centrifuged at

16,000 g for 15min at 4°C. The supernatant was used as the soluble fraction and the pellets

containing the insoluble fraction were dissolved in a urea buffer (9 M urea, 50 mM Tris buffer, pH

8.0) after wash.

All protein samples were then boiled at 100°C for 5 min and separated using a 10% Bis-Tris

SDS-PAGE (Invitrogen). Detection was performed using the High-sig ECL Western Blotting

Substrate (Tanon). Images were captured using an Amersham Imager 600 (GE Healthcare) and

densitometry was measured using ImageQuant TL Software (GE Healthcare). The contrast and

brightness were optimized equally using Adobe Photoshop CS6. All experiments were

normalized to tubulin or GAPDH as indicated in the figures.

Cell viability assay

Transfected 293T cells were seeded in 96-well plates (Corning) at the density of 9×10^3 cells/well

and cultured in 100 µL of culture medium. Otherwise, cell viability was examined 48-72 h after

transfection using the Cell Counting Kit-8 (CCK-8) (Dojindo), according to the manufacturer's

instructions. Briefly, 10 µL of the CCK-8 solution were added to each well and incubated at 37°C

for 2.5 h. Finally, the absorbance at 450 nm was measured with a Synergy2 microplate reader

(BioTek Instruments).

Mouse care and surgical procedures

All mouse procedures were performed in compliance with the institutional guidelines on the

scientific use of living animals at Interdisciplinary Research Center on Biology and Chemistry,

the Chinese Academy of Sciences (CAS). "Principles of laboratory animal care" (NIH publication

No. 86-23, revised 1985) were followed. Animal distress and conditions requiring euthanasia

were addressed and the number of animals used was minimized.

Lentivirus production and primary neuron culture

To generate lentivirus for infecting primary neurons, 293T cells were co-transfected with

pLenti-hSyn-TDP-43-HA, psPAX2 and pMD2.G with a ratio of 4:2:1 in Opti-MEM medium using

Lipofectamine™ 2000. Culture supernatant was collected at 48 h after transfection and passed

through a 0.45-μm filter. Viral particles were concentrated from culture supernatants by Lenti-X™

Concentrator (Clontech). Viral pellets used for neuronal infection were resuspended in

Neurobasal medium (Invitrogen).

Primary cortical neuron were isolated from C57BL/6 mouse cortex at embryonic day 17 (E17)

and cultured in serum-free Neurobasal medium (Invitrogen) supplemented with 2% B27,

GlutaMax, and penicillin-streptomycin (Invitrogen). At 7 days in vitro (DIV), neurons were

infected with pLenti-hSyn-TDP-43-HA for 5 days before extraction for RNA or

immunofluorescence.

RNA extraction and real-time quantitative PCR

For quantitative PCR (qPCR), total RNA was isolated from mouse primary neuron using TRIzol

(Invitrogen) according to the manufacturer's instruction. After DNase (Promega) treatment, the

reverse transcription reactions were performed using All-in-One cDNA Synthesis SuperMix kit

(Bimake). The cDNA was then used for real-time qPCR using the SYBR Green qPCR Master Mix

(Bimake) with the QuantStudio™ 6 Flex Real-Time PCR system (Life Technologies). The mRNA

levels of GAPDH were used as an internal control to normalize the mRNA levels of NEAT1. The

qPCR primers used in this study are listed below:

Total mNEAT1:

5'- ACTCTTGCCCCTCACTCTGA -3'

5'- CAGGGTGTCCTCCACCTTTA -3';

mNEAT1_2:

5'- CCCACACCTCAGTGGTTTCT -3'

5'- ACAGAACCAAGGCACAATCC -3';

mGAPDH:

5'- CACCATCTTCCAGGAGCGAG -3'

5'- CCTTCTCCATGGTGGTGAAGAC -3';

Purification of TDP-43 proteins

WT or mutant TDP-43 protein was expressed in BL21 (DE3) *E. coli* (TransGenBiotech, CD601-03) at 19°C for 16 h after induction by adding 100 μM of IPTG as previously described (*46*). In brief, cells were harvested by centrifugation at 4000 rpm for 20 min at 4 °C and lysed in 50 mL of lysis buffer (50 mM Tris-HCl, 500 mM NaCl, pH 8.0, 10 mM imidazole, 4 mM β-mercaptoethanol, 1 mM PMSF, and 0.1 mg/mL RNase A). After cell lysates were filtered with a 0.22 μm filter, the protein was purified using Ni columns (GE Healthcare, USA) and then eluted in an elution buffer (50 mM Tris-HCl, 500 mM NaCl, pH 8.0, 250 mM imidazole and 4 mM β-mercaptoethanol). The proteins were further purified using the Superdex 200 16/600 columns (GE Healthcare) in a buffer containing 50 mM Tris-HCl pH 7.5, 300 mM NaCl and 2 mM DTT, and freshly frozen in liquid nitrogen and stored at -80 °C. RNase A was routinely added in cell lysates and administrated again before chromatography during the protein purification procedure. All purified proteins were confirmed by Coomassie brilliant blue staining and Western blotting before use.

RNA-binding dot-blot assay

Purified WT or mutant TDP-43 protein was diluted in a blotting buffer (50 mM Tris-HCl, pH 7.5, 300 mM NaCl, and 5% glycerol) and blotted onto a 0.45 μm nitrocellulose membrane. The membranes were left to dry at room temperature for 30 min and then stained with Ponceau S for 10s. Images were captured using an Amersham Imager 600 (GE Healthcare). After the images were captured using an Amersham Imager 600 (GE Healthcare), the membranes were washed with PBST (0.05% TWEEN 20 in PBS) for 30 min and then incubated in the PBST containing 25 ng/μl total RNAs or NEAT1 RNA for 1h with gentle rocking and rotation at room temperature. The membranes were then washed in PBST and incubated in the SYBR™ Gold Nucleic Acid Gel Stain (Invitrogen) for 10 min. RNAs bound to the membranes were imaged and examined using the Gel Image System (Tanon).

In vitro phase separation and RNA buffering assay

For the *in vitro* LLPS experiments, purified WT or mutant TDP-43 protein was mixed with NaCl at

indicated concentrations in a phase separation buffer (50 mM Tris-HCl, pH 7.5 and 5-10% *(w/v)*

PEG 8000 (Sigma)) and incubated for 1 min at room temperature. For the RNA buffering assay,

the TDP-43 proteins were incubated with total RNAs, tRNA or NEAT1 RNA in the above phase

separation buffer with NaCl at indicated concentrations as shown in the figures. Finally, 5 μL of

each sample was pipetted onto a coverslip and imaged using a Leica microscope with differential

interference contrast (DIC). The total RNAs were extracted from HeLa cells using TRIzol

(Invitrogen) according to the manufacturer's instructions. Yeast tRNA was purchased from

Invitrogen. The NEAT1 RNA was *in vitro* transcribed and purified using HiScribe™ T7 Quick High

Yield RNA Synthesis Kit (NEB). Total RNAs and NEAT1 RNA were used within one day of

production.

Drosophila strains

Flies tested in this study were raised on standard cornmeal media and maintained at 25 °C and

60% relative humidity. The following strains were obtained from the Bloomington *Drosophila*

Stock Center (BDSC): *D42*-Gal4 (#8816), *elav*-Gal4 (#8760) and UAS-*LacZ* (#8529). The

transgenic fly strains of WT and mutant UAS-hTDP-43 were generated by ΦC31

integrase-mediated, site-specific integration, which allowed uniform transgene expression across

different lines. The attP2 landing site stock used for the fly embryo injection and transformation in

this study was y[1] M{vas-int.Dm}ZH-2A w[*]; P{CaryP}attP40 (25C6), and the transgenic

pBID-UASC-Luciferase (UAS-Luc) fly strain that was generated using the same approach at the

same landing site (*100*) was used as a control.

Fly climbing assay

For the climbing assay, ~20 flies per vial, ~10 vials per group were tested. All flies were

transferred into an empty polystyrene vial and allowed 10 min for flies to recover. The flies were

then gently tapped down to the bottom of the vial and the number of flies that climbed over 3cm

within 10 seconds was recorded. The test was repeated three times for each via.

Statistical Analysis

Statistical significance in this study is determined by one-way analysis of variance (ANOVA) with

Tukey's HSD post-hoc test, two-way ANOVA with Bonferroni's post-hoc test, or unpaired,

two-tailed Student's t-test with unequal variance at $*p < 0.05$, $**p < 0.01$, and $***p < 0.001$ as

indicated in each figure legend. Error bars represent the standard error of the mean (SEM).

Supplementary Figures

Figure S1_CWang

Figure S1. Co-immunostaining of TDP-43 with several known NB markers

Representative confocal images of the nucleus of 293T cells transfected with TDP-43-HA and treated with 250 μM NaAsO$_2$ for 30 min. All cells are examined by immunocytochemistry with anti-HA for TDP-43-HA (green) and the indicated antibodies (red): (**A-B**) SFPQ (splicing factor proline-glutamine rich), for paraspeckles (*74*). (**C-D**) PML (promyelocytic leukaemia) protein, for PML bodies (*101*). (**E-F**) TAF9 (TATA-box binding protein associated factor 9) for Cajal bodies (*102*). (**G-H**) SC35 (serine/arginine-rich splicing factor-like protein), for nuclear speckles (*103*). Merged images with DAPI staining (for DNA) are shown. Arrows indicate the examples of co-localization of TDP-43 with the indicated NB markers. The co-localization of TDP-43 with each type of NBs is further evaluated by the line scanning analysis shown in (B), (D), (F) and (H), Scale bar: 5 μm. The results indicate that TDP-43 NBs are well colocalized with paraspeckles, and in a few rare cases partial colocalization with PML bodies, Cajal bodies or nuclear speckles are also spotted.

Figure S2. Stress-induced NBs are observed with endogenous TDP-43 and in primary neurons

(**A-B**) Representative images of endogenous TDP-43 (endo-TDP-43) forming NBs upon arsenite treatment (500 μM, 30 min). DAPI staining to show the nucleus and merged images are also shown. Percentage of cells showing distinct TDP-43 NBs in (A) is quantified in (B). Arrows indicate TDP-43 NBs. (**C-D**) Representative images (C) and quantification (D) of TDP-43-HA NBs in primary mouse cortical neurons induced by arsenite treatment. TDP-43-HA (green), microtubule associated protein 2 (MAP2) to mark neuronal perikarya and dendrites (red), and merged images with DAPI staining to label the nucleus (blue) are shown. Zoom-in images of the boxed areas are also shown. Mean ± SEM, n = 100 cells each group from pooled results of 3 independent repeats; **$p < 0.01$, ***$p < 0.001$; Student's t-test. Scale bars: 5 μm in (A and C), and 2 μm in the zoom-ins.

356 **Figure S3. TDP-43-NES^{mut} NBs are dynamic and reversible**

(**A-D**) Representative Western blot images (A) and quantifications (B-D) of the pulse-chase
experiments of WT, NLS^{mut} and NES^{mut} TDP-43 protein in the soluble (S, supernatants in RIPA)
and the insoluble (I, pellets resuspended in 9 M of urea) fractions at indicated time after
Cycloheximide (CHX) treatment. All proteins are normalized to tubulin in the soluble fraction and
the relative level at 0 h is set to 100%. Mean ± SEM, n = 3 independent repeats; compared to the
level at 0 h; *P < 0.05, ***P < 0.001; ud, undetected; one-way ANOVA. (**E**) Representative
images showing TDP-43-NES^{mut} NBs in 293T cells at indicated time after CHX treatment. (**F**)
Quantification of the average number of TDP-43 NBs per cell in (E). Mean ± SEM, n = 10~11
cells per time point, from pooled results of 3 independent repeats; ***p < 0.001; one-way ANOVA.
Scale bar: 5 μm.

Figure S4. The turnover of insoluble TDP-43 is sensitive to proteasomal but not autophagic inhibition

(**A-D**) Representative images (A) and quantifications (B-D) of Western blotting analyses of WT, NLS^{mut} and NES^{mut} TDP-43 in the soluble (S) and insoluble (I) fractions at indicated time after treating the cells with MG132 to inhibit the proteasome-mediated protein degradation. (**E-H**) Representative images (E) and quantifications (F-H) Western blot analyses of WT, NLS^{mut} and NES^{mut} TDP-43 in the soluble (S) and insoluble (I) fractions at indicated time after treating the cells with chloroquine (CQ) to block the autophagic flux. S, supernatants in RIPA; I, precipitates in RIPA that are re-suspended in 9 M of urea. All proteins are normalized to tubulin in the soluble fraction and the relative level of each fraction at 0 h is set to 1. Mean ± SEM, n = 3 independent repeats; *P < 0.05, ***p < 0.001; ud, undetected; one-way ANOVA.

Figure S5. The impact of the LCD and RRM domains on the assembly of TDP-43 NBs

(**A**) A diagram showing the major functional domains and the truncated ΔLCD, ΔRRM1 and ΔRRM2 TDP-43. (**B-E**) Western blot analysis of TDP-43-WT (B), ΔLCD (C), ΔRRM1 (D) and ΔRRM2 (E) in the RIPA-soluble and RIPA-insoluble (resuspended in 9 M of urea) fractions. All TDP-43 proteins are normalized to tubulin in the soluble fraction and the relative level is quantified as percentage to the soluble fraction of each protein. Mean ± SEM, n = 3 independent repeats. (**F-I'**) Representative confocal images of TDP-43-WT, ΔLCD, ΔRRM1 and ΔRRM2 in 293T cells treated with the vehicle control ethanol (F-I) or LMB (F'-I'). Green, anti-Myc for Myc-TDP-43; blue, DAPI for DNA. Scale bar: 5 μm.

Figure S6_CWang

Figure S6. Total RNAs suppress TDP-43 LLPS *in vitro*

(**A**) A diagram showing the full-length (FL) and truncated TDP-43 protein without the LCD (TDP-43$^{1-274}$). (**B**) Coomassie blue staining confirming the purified FL and truncated sumo-TDP-43 proteins. (**C-D**) The *in vitro* LLPS of FL (C) and TDP-43$^{1-274}$ (D) is dose-dependent. (**E-F**) Suppression of the LLPS of the FL and the TDP-43$^{1-274}$ by total RNAs extracted from HeLa. (**G**) Coomassie blue staining confirming purified WT, ΔRRM1 and ΔRRM2 TDP-43$^{1-274}$ proteins used in the *in vitro* assays below and in Figure 6. (**H**) Representative images of the *in vitro* dot-blot assay confirming the reduced RNA binding affinity of the ΔRRM1 and ΔRRM2 TDP-43$^{1-273}$ compared to the WT. The bound RNA is visualized using the SYBR® Gold Nucleic Acid Gel Stain kit. Bovine serum albumin (BSA) is used as a negative binding control. (**I**) The RNA intensity is normalized to Ponceau S staining (protein loading control) and shown as the percentage to that of WT TDP-43$^{1-273}$ in (H). Mean ± SEM, n = 3; ***$p < 0.001$; Student's *t*-test. (**J-L**) WT (J), ΔRRM1 (K) and ΔRRM2 (L) TDP-43$^{1-273}$ proteins forms LDs *in vitro* by LLPS in a dose-dependent manner. The concentrations of NaCl, the crowding agent PEG, TDP-43 protein and total RNAs used in the above assays are as indicated. Scale bars: 2 μm.

Figure S7. The D169G mutant TDP-43 has a specific defect in NB formation

(**A**) Representative confocal images of HeLa cells expressing with WT or D169G TDP-43 treated
with LMB for indicated times. (**B-C**) Quantifications of the percentage of cells showing TDP-43
NBs (B) and the average number of NBs per cell (C) at the end of the experiment in (A).
LMB-induced NB formation was delayed in D169G TDP-43. (**D**) A diagram showing WT TDP-43
and the M337V mutation in the LCD. (**E**) Representative confocal images of cells transfected
with WT or M337V TDP-43 in the absence (PBS) or presence of arsenite. Green, TDP-43-HA
(anti-HA); Blue, DAPI staining for DNA (blue); arrows, TDP-43 NBs. (**F-G**) The percentage of
cells with TDP-43 NBs (F) and the average count of TDP-43 NBs per cell (G) in (E) are quantified.
Mean ± SEM, n = ~100 cells in total for each group in (B, F) and 11~15 cells per group in (C, G),
pooled results from 3 independent repeats; *$p < 0.05$, ***$p < 0.001$; ns, not significant; Student's
t-test. Scale bars: 5 μm.

Supplementary movies

Movie S1. A representative time-lapse video of *live* HeLa cells showing the formation of
GFP-TDP-43 NBs upon the arsenite treatment

Movie S2. A representative time-lapse video of GFP-TDP-43-GFP NBs in *live* HeLa cells
before and after photobleaching