Bacteria primed by antimicrobial peptides develop tolerance and persist

Authors: Alexandro Rodríguez-Rojas[1§], Desiree Y. Baeder[2§], Paul Johnston[3,4], Roland R. Regoes[2]* and Jens Rolff[1,5]*.

Affiliations:

[1]Freie Universität Berlin, Institut für Biologie, Evolutionary Biology, Königin-Luise-Strasse 1-3, 14195 Berlin, Germany.

[2]Institute of Integrative Biology, Universitätsstrasse 16 ETH Zürich, 8092 Zürich, Switzerland

3. Berlin Center for Genomics in Biodiversity Research, Berlin, Germany;

4. Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin, Germany;

5. Berlin-Brandenburg Institute of Advanced Biodiversity Research (BBIB), Berlin, Germany

*Correspondence: roland.regoes@env.ethz.ch, jens.rolff@fu-berlin.de,

§These authors contributed equally to this work

SUMMARY

Antimicrobial peptides (AMPs) are key components of innate immune defenses. Because of the antibiotic crisis, AMPs have also come into focus as new drugs. Here, we explore whether prior exposure to sublethal doses of AMPs increases bacterial survival and abets the evolution of resistance. We show that *Escherichia coli* primed by sublethal doses of AMPs develop tolerance and increase persistence by producing curli or colanic acid. We develop a population dynamic model that predicts that priming delays the clearance of infections and fuels the evolution of resistance. The effects we describe should apply to many AMPs and other drugs that target the cell surface. The optimal strategy to tackle tolerant or persistent cells requires high concentrations of AMPs and fast and long-lasting expression. Our findings also offer a new understanding of phenotypic drug resistance as an adaptive response and could lead to measures that slow the evolution of resistance.

INTRODUCTION

Antimicrobial peptides - short, usually cationic peptides - are key effectors of innate immune defences of all multicellular life (Zasloff, 2002) and are also important players at the host microbiota interface (Bevins and Salzman, 2011; Mergaert, 2018). Because of their evolutionary success and diversity, AMPs are considered as new antimicrobial drugs to alleviate the antibiotic resistance crisis (Fjell et al., 2012) with currently more than two dozen under clinical trial (Koo and Seo, 2019). *Bona fide* genetic resistance in AMPs has been studied (Joo et al., 2016, Perron et al., 2006, Yu et al., 2018), but not to the extent of antibiotic resistance. Resistance against AMPs evolves usually with a low probability and the levels of resistance are not as high as against antibiotics (Spohn et al., 2019; Yu et al., 2018).

Notably, phenotypic resistance (Levin and Rozen, 2006), the ability of bacteria to survive lethal concentrations of antimicrobials without a genetically encoded resistance mechanism, has hardly been studied for AMPs. One of the very rare studies that has addressed this in a natural host-microbe interaction is the example of the bobtail squid and its symbiont *Vibrio fischeri*. Here, phenotypic resistance is elicited by a low pH and primes the *Vibrios* to colonize the light-emitting organ of the squid (Chen et al., 2017) in the presence of high concentrations of AMPs. Microbes have evolved adaptive physiological alterations to predictable environmental changes (Brown et al., 2019; Mitchell et al., 2009). This has been studied in the context of available carbon sources during gut passage (Mitchell et al., 2009), and a meta-analysis found that priming, the phenotypic response to a low level stressor, provides a fitness benefit for microbes against a variety of stressors including pH, temperature and oxidative stress (Andrade-Linares et al., 2016).

Here we study if the previous encounter with sub-lethal dosages of AMPs induces phenotypic resistance against AMPs. Such low concentrations are common. For example, after infection the induction of AMPs results in sublethal concentrations before killing concentrations are reached. This usually takes a few hours (Makarova et al., 2016). Such pharmacokinetic profiles are mirrored in many antimicrobial drug treatments. During medical application of antibiotics, the pharmacokinetics start at zero and the killing concentrations are building up over time.

Phenotypic resistance can be induced by sublethal levels of antimicrobials (Andersson and Hughes, 2014). Phenotypic resistance, resulting in either drug tolerance or persister cell formation (see Fig. S 1, Balaban et al., 2019; Brauner et al., 2016) has been shown to be of great importance to understand antibiotic resistance (Andersson and Hughes, 2014; Balaban et al., 2019; Levin and Rozen, 2006; Levin-Reisman et al., 2017). It can also facilitate *bona fide* resistance evolution against antibiotics (Levin-Reisman et al., 2017). We adopt these findings and concepts from antibiotic research to understand phenotypic resistance against AMP, an insight which should equally inform our understanding of host-microbe interactions as well as resistance evolution against AMPs as drugs. It is noteworthy that AMPs differ significantly from conventional antibiotics in several aspects including their pharmacodynamics, resulting in narrower mutant selection windows (Yu et al., 2018). Also, the speed at which they kill cells is fast, as they kill within minutes (Fantner et al., 2010) rather than within hours, as it is the case for antibiotics (Yao et al., 2012).

In this study, we specifically investigate whether prior exposure to sublethal doses of AMPs increases bacterial survival via either tolerance or persistence, and the risk of resistance evolution. We use two antimicrobial peptides that are well characterised as a

case study. Melittin is a 26 amino acid residue linear peptide from the honeybee, which kills bacterial cells by permeabilization of the membrane (Raghuraman and Chattopadhyay, 2007). Melittin is also active against eukaryotic parasites such as Leishmania and cancer cells. The other AMPs used here, pexiganan, is the first eukaryotic AMP developed as a drug, a synthetic 22 amino acid residue peptide closely related to magainin from the African clawed frog (Ge et al., 1999). It shows a broad activity against both, gram$^+$ and gram$^-$ bacteria and kills by forming toroidal pores (Gottler and Ramamoorthy, 2009). We combine *in vitro* experiments with a modelling approach to study priming and resistance emergence. We find that a sublethal dose of certain antimicrobial peptides can induce increased tolerance and/or persistence in bacteria and hence prime (Hilker et al., 2016) them for the exposure to a subsequent lethal dose. We identify candidate underlying molecular mechanisms and capture the population dynamics by adapting a classic mathematical model of persistence (Balaban et al. 2004). With computer simulations, we then predict that increasing tolerance and persistence will have a positive effect on bacterial survival and the emergence of resistance.

RESULTS

Primed cells are more resistant

We primed *E. coli* K-12 in vitro by exposing them to a fraction of the minimal inhibitory concentration of an AMP (0.1xMIC, table S1). Subsequently, we exposed the primed bacterial populations to lethal concentrations of the respective AMPs (10 x MIC, table S1) and then monitored bacterial survival over time. We found that the priming treatment resulted in much higher *E. coli* survival (Fig. 1 A, B).

Figure 1: Bacterial tolerance and persistence determine the shape of time-kill curves.

Time-kill experiments with *E. coli* K-12, primed (orange) and naïve (blue) bacteria were

exposed to 10xMIC of (A) melittin and (B) pexiganan. At each timepoint indicated, we

measured the bacterial population size 5 times. We tested with an ANOVA by means of

contrasts if priming influenced tolerance and persistence. For both antimicrobials, priming

significantly increased the slope of the first phase, the measure of tolerance, and the bacterial

level in the second phase, the measure of persistence, (significance level: $p < 0.05$). We

corrected for multiple testing with the Bonferroni-method. The line in the plots indicates the

best fit of a biphasic function (table S2), on which our statistical analysis is based.

Killing is biphasic

The decline of the time kill curves is biphasic suggesting two subpopulations (Abel zur

Wiesch et al., 2015, Balaban et al., 2002, Hedges, 1966, Wiuff et al., 2005). We

excluded that deviations from monophasic decline arise because of decreasing

antimicrobial concentrations over time (Fig. S 2) and fitted the time-kill curves to a

biphasic linear function. For both AMPs, bacterial populations declined faster during

the first than the second phase (Fig. 1, Table S 2). Tolerance, the decline of bacterial

populations in the first phase (Balaban et al., 2019), was significantly higher in primed

than in naïve bacteria for both AMPs. Primed bacteria showed higher survival in the

second phase, indicating higher numbers of persisters (Balaban et. al., 2019). The change in population size in the second phase, however, was not significantly different between primed and naïve populations, indicating that the population dynamics as such in the second phase are not influenced by priming (Fig. 1). In short, priming with AMPs allow bacteria to survive better by increasing both bacterial tolerance and persistence.

Priming is mediated by either curli or colanic acid

To understand how priming leads to tolerance and persistence, we used RNAseq of cells exposed to priming concentrations of AMPs (Fig. 2 A, B). Exposure to sublethal concentrations of pexiganan (0.1xMIC, as above) induced colanic acid synthesis (Fig. 3 A). Colanic acid capsules have been shown to protect against AMPs and antibiotics (Miajlovic and Smith, 2014). The removal of an essential gene for colonic acid production completely abolished the priming effect by pexiganan (Fig 3 B). By phase contrast imaging, we observed the formation of a characteristic colonic acid capsule in pexiganan-primed but not in naïve cells (Fig. 3 C). The priming response was homogenous, with all observed cells producing a colanic acid capsule under priming conditions. Exposure to a sublethal concentration of melittin (0.1xMIC, as above) induced up-regulation of curli fimbriae (Fig 3 D). Curli is an important virulence factor (Barnhart and Chapman, 2006) and a component of extra-cellular matrix that protects against AMPs (Maria-Neto et al., 2015). A curli-deficient mutant showed a decrease of the priming effect induced by melittin (Fig. 3 B). For melittin-primed bacteria, curli induction was documented with a specific curli binding chemical (Fig. 3D). Both AMPs also induced significant overlap in gene expression related to osmotic shock (Fig. 2).

2 **Figure 2. Gene expression in primed *E. coli*.** (A) Venn diagrams showing specific and

3 overlapping responses of *E. coli* MG1655 to priming concentrations of melittin and pexiganan

4 (B) Principal component 1 separates the control from the peptide priming, PC 2 separates the

5 melittin induced response from the pexiganan response.

7 Figure 3 A

1   Figure 3B

3   Figure 3C

5   Figure 3D

8   **Figure 3. Physiological response to AMP priming.** (A) SEM of E. coli MG1655 treated with

9   1/10xMIC (priming concentration) of pexiganan (top row) and melittin (bottom row) and non-

10   treated bacteria (middle row, control). No apparent difference was noticed between melittin-

11   treated cells and control. In the case of pexiganan, the treated cells tend to aggregate, a phenotype

12   that consistent with the presence of colanic acid. Red arrows represent potential areas altered by

the capsule of colanic acid that collapse under drying processing necessary for the preparation.

Bacteria were observed with different magnifications ranging from 3000X to 40000X.(B).

Bacterial mutants in *csgA* (curli mutant) and *wza* (colonic acid mutant) were exposed to the AMPs

melittin and pexiganan respectively (10xMIC) after priming. Colony forming units (CFU) were

determined after two hours of exposure. Boxplot show data tested by repeated-measures of one-

way ANOVA and Dunnetts' tests. For each case, equal letter represents no statistical differences

while differing letter detonates significant ones (for $p<0.05$). (C). *E. coli* MG1655 treated with

1/10xMIC (priming concentration) of pexiganan (left) and non-treated bacteria (right, control)

observed under phase contrast optical microscopy. The specimens consisted of cells suspended in

a 0.1 % solution of nigrosin to create a strong contract to visualize the colonic acid capsules.

Bacteria were observed with magnifications 1000X after exposure to a trigger stimulus. (D).

Detection of curli production by primed cells after exposure to a killing dose of melittin

(10xMIC). Curli production is only detected in small proportion of primed cells (survival

fraction). After treatment, we exposed the cells to ECtracer™680 (Ebba Biotech, Sweden), a red

fluorescent tracer molecule for staining of curli.

After lethal exposure to melittin and pexiganan we observed a high degree of

heterogeneity regarding the killing of individual cells in primed bacteria, with many cells

surviving the killing concentrations (Fig. 4 A). By contrast, almost all the control bacteria

were killed (Fig. 4 A). Primed cells also seem to aggregate, with a stronger effect in

pexiganan-treated cells compared to melittin-treated cells (Fig. 4 A).

Figure 4 A

Figure 4 B

Figure 4. Cell viability and persister formation after AMP priming. (A) Cell viability after treating with priming and trigger doses of melittin and pexiganan were determined using the live/dead BacLight Bacterial Viability Kit (Thermo Scientific, Germany) on chip as described in M&M. After priming during 30 minutes, the treatment was removed by perfusing fresh MHB. The fluorescence signal was analyzed via a using excitation at 485 nm and emission at 530 nm for green fluorescence (Syto9) and using excitation at 485 nm and emission at 630 nm for red fluorescence (propidium iodide). (B) The priming concentrations of pexiganan and melittin, respectively, increased the persister number as determined by treated primed and naïve populations (10^8 CFU/ml) with ciprofloxacin (10x MIC, 2μg/ml). This increase in persistence co-occurs with ATP leakage (determined in the supernatant of the culture medium).

To determine if priming stimulates persistence we used an assay based on the observation

that a decrease of intracellular ATP increases persistence under the antibiotic ciprofloxacin (Shan et al., 2017a). Because ATP leakage is a hallmark of AMP-treated bacteria (Brogden, 2005), we exposed melittin and pexiganan primed populations and controls to ciprofloxacin (Shan et al., 2017a). This resulted in a highly significant increase in the number of persisters (Fig. 4 B). The level of leaked ATP in the culture supernatant was significantly higher in primed bacteria for both AMPs as compared to controls (Fig. 4 B). The pre-treatment with melittin or pexiganan does not change the MIC of *E. coli* to melittin, pexiganan or ciprofloxacin consistent with the definition of persisters (Balaban et al., 2019) (Table S 3).

A mathematical model predicts resistance evolution of primed cells

To quantify the influence of priming on bacterial tolerance and persistence, phenomena inherently linked to the growth dynamics and subpopulation structure, population dynamical models that capture tolerance and persistence are required. We build on a two-state population model previously developed (Balaban et al., 2004) to describe bacterial antibiotic persistence (Fig. 5 A, B). This model assumes that bacteria exist in two phenotypic states, normal cells (N) and persisters (P). The two subpopulations N and P differ in their susceptibility to AMPs, a difference that is implemented as differing net growth rates for the given amount of AMPs (A), with $r_N(A)$ and $r_P(A)$, respectively. Bacteria switch from subpopulation N to P with the rate s_N and back with the rate s_P.

We quantified the model parameters by fitting the analytical solution of the model equations (Fig. 5 B) to the time-kill datasets of melittin and pexiganan. In a preliminary analysis with fitting all four parameters (rN, sN, rP, and sP), the net replication rate of persister cells, rp, was not significantly different from 0 (Fig. S 4). We therefore fixed

rP to 0. The resulting fit described our data well (Fig. 5 C, D, and Table S 5). We found

that priming affected two of the three estimated model parameters (Fig. S 4): rN

increased, which translates into increase in tolerance, and sP decreased. Together, the

increase in rN and decrease in sP result in higher persistence levels (Fig. 5 C, D).To

assess the influence of priming on possible treatment success and resistance evolution,

we extended a previously developed modeling framework to predict resistance

evolution (Yu et al., 2018) by a persistent subpopulation (Fig. S 5). We then explored

the effects of priming on tolerance and persistence, individually and in combination,

which would be challenging empirically. Using our predictive framework, we

investigated the effect of priming on survival based on a zero-order pharmacokinetic

profile (Regoes et al., 2004; Yu et al., 2018) (Fig. S 6) and with parameterized

pharmacodynamics functions (Fig. S 7).

Figure 5

A

B

$$\frac{dN}{dt} = r_N(A)N(t) - s_N N(t) + s_P P(t)$$

$$\frac{dP}{dt} = r_P(A)P(t) - s_P P(t) + s_N N(t)$$

Figure 5: The two-state model describes time-kill data. (A) The two-state model (adapted from

17) consists of two subpopulations, normal cells (N) and persister cells (P) and is parameterized

with the growth rates $r_N(A)$ and $r_P(A)$, which are depended on the concentration of AMPs (A) in

the system, and the switching rates s_N, and s_P. Each subpopulation is described with an ordinary

differential equation (B), each of which describes the change of the respective subpopulation over

time. For each antimicrobial, melittin (C) and pexiganan (D), we fitted the model to the data of

naive (blue) and primed (orange) bacterial populations (see also Fig. 1) individually. The

continuous lines represent the total bacterial population B(t), with B(t) = N(t) + P(t), and the

dashed and dotted lines represent the subpopulations N(t) and P(t), respectively. Bacteria primed

with melittin have an increased net growth rate r_N and decreased s_P compared to the naive

populations. In the case of pexiganan, the parameter r_N is significantly higher in primed compared

to naive populations. For an overview of the fitted parameters, see table S 5.

We found that survival of the population was highly dependent on tolerance (Fig. 6 A,

B, S 8). The presence of persistent cells alone only marginally increased time until

clearance. When we implemented the observed decrease in switching rate s_P, the time

until clearance of the bacterial population is extended at high treatment intensities (high concentrations A_{max}). The results do not qualitatively change for larger pharmacokinetic decay rates (k), typical for AMPs (Fig. S 9). Taken together, an increase in tolerance alone resulted in higher survival independent of persister cells. An increase in persister cells further increased survival at high antimicrobial concentrations.

Next, we assessed how priming affects resistance emergence. Generally, resistance evolution depends on the population size, mutation rate and the replication rate. While priming increases bacterial survival, hence the population size, it also increases the number of persisters that do not replicate and hence are a poor source of resistant mutations (Balaban et al., 2004) (Fig 1, Fig. S 10). Our simulations revealed that the beneficial effects of priming on survival due to increased persistence did not translate into an increased probability of resistance emergence and establishment (Fig. 6 B, Fig. S 8, 9). The probability of resistance emergence was mainly influenced by the effect of priming on tolerance.

Figure 6

Figure 6: Influence of priming on time until clearance and resistance evolution. We extended

a previously developed PKPD framework (*7*) to include persistence (Fig. S 5). With this

framework, we estimated PD curves (Fig. S 7) for primed and naïve bacteria and the

estimated switching rates (Fig. 5) to predict (A) time until clearance and (B) probability

of resistance evolution. We simulated primed bacteria (primed with heterogeneous

population consisting of N and P subpopulations, indicated as "primed, heterogeneous,

smaller s_P") and naïve bacteria (naïve, heterogeneous). In addition, we simulated dynamics

without persistence for both primed and naïve bacteria ("primed homogeneous" and

"naïve homogeneous") and dynamics of primed populations with switching rates of the

naïve population ("primed, heterogeneous"). In (A), each dot is an individual run. In both

(A) and (B), the line denotes the average of all runs. No clearance means that simulated treatment could not reduce bacteria population < 1 cell within 7 days of treatment. Here, we only show the results of melittin (results of pexiganan are qualitatively highly similar, see Fig. S 8). All parameter values used in the mathematical model are listed in table S 5.

DISCUSSION

We find that sublethal dosing of the AMPs melittin and pexiganan primes bacterial cells to increase both tolerance and persistence. As sublethal concentrations of antimicrobials (AMPs and antibiotics) are common in the environment and in hosts/patients, a priming response, if it is a widespread phenomenon, would have significant consequences. Antimicrobials that prime inevitably would induce the formation of tolerant and persisting cells. This should apply to situations of induced immune responses, but equally to drug treatments. In short, generating or increasing populations of phenotypic resistant bacteria, would be inevitable in many situations, and would represent a serious obstacle to clearing bacterial infections.

The priming response we report is mediated by rather general physiological resistance mechanisms. The molecular basis of the induction of AMP tolerance and persistence is based on modifications of bacterial envelopes involving either curli production under melittin treatment, or colonic acid production after pexiganan exposure. Both these mechanisms protect the cells by shielding them from AMPs, either making the access to the outer membrane more difficult and/or capturing AMPs, giving scope for cross-resistance. These physiological resistance mechanisms are not antimicrobial specific and hence likely to be also induced by other antimicrobials.

Interestingly, the activation of both pathways shows different dynamics in biofilm formation (Prigent-Combaret et al., 2000). Curli and colanic acid are important components of the biofilm matrix, curli for the primary adhesion and colanic acid for the biofilm structure (Prigent-Combaret et al., 2000). Triggering their expression by sublethal levels of AMPs, could potentially catalyse biofilm formation. Within a host, if the immune system fails to clear the pathogens, the AMP-priming effect may thereby favour the transition from acute to chronic bacterial infections, where biofilms prevail.

In natural systems of host-microbe interactions, priming plays a role in facilitating colonization of AMP rich environments (Chen et al. 2017). Priming by AMPs also plays a role in infection vectors: in the flea gut the PhoQ-PhoP system is induced in *Yersinia pestis,* the infective agent causing plague, by AMPs leading to biofilm formation that enhances transmission to the final host (Vadyvaloo et al., 2015). It is not clear as yet if phenotypic AMP-resistance will facilitate opportunistic infections in a way similar to genetic AMP-resistance, as has been shown for genetic AMP resistance in *S. aureus* (Cheung et al., 2018), but in the light of our results it seems likely.

Phenotypic resistance can drive the evolution of genetic resistance (Levin and Rozen, 2006). We find that, while priming *E. coli* with AMPs results in increased tolerance and persistence, the main driver of resistance evolution in our model is increased tolerance. Resistance evolution is a product of mutation supply and strength of selection. The selection in our scenarios is strong, but mutations are only supplied from the subpopulation of tolerant cells, not from the persister cells, as they are metabolically inactive. So, while the persisters do not contribute to resistance evolution, they do provide a source for secondary infection, once the immune system overexpression ceases or alternatively, when the antimicrobial drug is removed. In antibiotic resistance evolution, by contrast, persisters have shown to contribute to resistance evolution (Windels et al., 2019). Part of this seems to be explained by increased mutagenesis caused by the

antibiotics (Windels et al 2019), an aspect we explicitly did not model as AMPs do not increase mutagenesis (Rodríguez-Rojas et al., 2014).

Our combined theoretical and empirical results suggest that in hosts the optimal strategy of AMP-deployment would be a very fast increase in concentration to avoid priming and the subsequent development of phenotypic resistance through both, increased persistence and increased tolerance. Such fast increases of AMPs in specific locations are realized in some important natural situations. During insect metamorphosis, when the gut is renewed in the pupa, a cocktail of AMPs and lysozymes is discharged into the gut (Johnston et al., 2019) and references therein), resulting in a quick reduction of bacterial numbers by orders of magnitude. When infections persist, one possible solution is a long-lasting immune response (Haine et al., 2008) that deals with a recurrent infection that could potentially be caused by persisting cells. These natural situations also have parallels in the medical application of antimicrobials. We propose that a fast increase of antimicrobial concentration, as for example in the intra-venous application of antibiotics, should also reduce the probability of persister formation via priming.

Long-lasting antimicrobial exposures are prevalent in natural systems. This is at odds with the observation that long-lasting drug treatments, at least in the case of single drug applications, select for drug resistance. Therefore, at first glance extended treatments do not seem to be a good strategy to manage persisters. In the natural context of AMPs, as they are always expressed as cocktails, this is very likely not an issue and would suggest the use of combination therapies.

Acknowledgments: We are grateful to Sophie Armitage, Charlotte Rafaluk-Mohr and Olivia Judson for comments on the manuscript.

METHODS

Bacteria and growth conditions

The *E. coli* MG1655 was used as bacterial models for all experiments. All cultures related to antimicrobial tests were carried out in Mueller-Hinton I Broth (Sigma). For genetic manipulations (*E. coli* BW25113). All the strains and their derivatives were routinely cultured in Lysogeny Broth (LB medium) or SOB, supplemented with antibiotics when appropriate. Other constructed and strains used in this study are listed in table S 6 of this section.

Minimal inhibitory concentration (MIC)

MICs were determined according to CLSI recommendations by a microdilution method in MHB with some modifications. Inoculum size that was adjusted to 2×10^7 CFU/ml from a regrowth of overnight cultures to be consistent with the downstream experiments. The MIC was defined as the antimicrobial concentration that inhibited growth after 24 h of incubation in liquid MH medium at 37°C. Polypropylene non-binding plates (Th. Geyer, Germany) were used for all experiments. The MIC was considered as the antimicrobial concentration that inhibited growth after 24 hours of incubation in liquid medium at 37°C.

MIC was also determined similarly as above for primed bacteria after two and a half hours of exposure to priming concentrations. The values were compared with naive controls for both antimicrobial peptides, melittin and pexiganan, and for ciprofloxacin.

Priming experiments

Starting from 1×10^8 CFU/ml, where bacteria were exposed (stimulus) to 1/10 MIC of melittin or pexiganan during 30 minutes at 37°C with soft shaking. The tubes were centrifuged at 4000 x g for 10 minutes and allowed to recover for 60 minutes. The cells were challenged (trigger of priming response) with a concentration equivalent to an 10X MIC. The cultures were diluted and plated to determine cell viability. Five replicas per culture were used, and every experiment was repeated twice. Non treated cells were used as a control.

Activity of melittin and pexiganan from the supernatant of challenged cells.

Similarly, to the priming experiments, 1×10^8 CFU/ml of bacteria were exposed 10x MIC (supernatant I) of melittin or pexiganan during 150 minutes at 37°C with soft shaking. The tubes were centrifuged at 4000 x g for 10 minutes, the supernatant were collected and centrifuged again for 20 000 x g for 30 minutes. The new supernatant was filtrated using 0.22 μm sterile filters (Sigma Aldrich, Germany) and used immediately (supernatant II). In parallel, exponentially growing cultures containing 1×10^8 CFU/ml per tube were centrifuged at 4000 x g for 10 minutes, the supernatants were removed by sterile aspiration and the pellets were re-suspended in equal volumes of the supernatant II and incubated during 150 minutes at 37°C with soft shaking. Samples from each tube were from both, supernatant I and supernatant II, taken every 30 minutes and diluted and plated to determine cell viability. Five replicas per culture were used. Data points from time-kill experiments from bacteria treated with the supernatant I (first round) and the supernatant II (second round) were compared to determine changes in the activity of the AMPs either by degradation or adsorption.

Persister antibiotic survival assay

Bacterial cultures were inoculated at 1:100 from a 16-hour overnight culture into MHB medium. Cells were grown for 2 h to reach approximately 2×10^8 CFU/ml). The cultures were treated with priming concentrations (1/10 MIC) of melittin and pexiganan for 30 minutes. Non-treated cultures were used as control. All cultures were washed and centrifuged twice to remove the treatment. The supernatants were used to determine ATP concentration using a Molecular Probes ATP Determination Kit (Thermo Fisher Scientific, Germany). Bacteria were resuspended in equal volume of fresh medium and an aliquot from each culture was taken to determine the number of bacteria at t=0 by diluting and plating in MHB agar. Following, ciprofloxacin was added for a final concentration of 2 µg/ml to treated tubes and to non-treated AMPs control. The cultures were incubated for four hours. Then, bacteria were washed twice with 0.9 % NaCl and plated on MHB agar to determine the counts of survival fractions. The percent survival was calculated as the ratio of cells before and after the treatment as described previously (Shan et al., 2017b). Briefly final CFU/CFU at 0 h) × 100. The results are presented as the average from 5 independent replicates.

Construction and verification of deletion mutants

We inactivated the major curli subunit protein gene *csgA* and the colonic acid precursor gene *wza*. Although both pathways involve many genes, the removal of these two components impair the production of both substances respectively. These mutants were generated in *E. coli* K-12 strain MG1655 following the methodology described elsewhere (Datsenko and Wanner, 2000) with some modifications because we used as template the genomic DNA of the Keio collection (Baba et al., 2006). Briefly, we extracted genomic DNA from the mutants *csgA::Kn* and *wza::Kn* of the Keio collection

(*E. coli* BW25113) and amplified by PRC the flanking regions of the kanamycin

resistance cassette disrupting both genes and including an appropriate homology

sequence. For the *csgA* mutant we used the primers 5'-

GATGCCAGTATTTCGCAAGGTG-3' and 5'-GGTTATCTGACTGGAAAGTGCC-

3' while primers 5'-TAGCGTGTCTGGATGCCTG-3' and 5'-

CCACTTTCAGCTCCGGGT-3' were used for *wza*. The PCR products were purified

and electroporated in the *E. coli* MG1655 carrying a red recombinase helper plasmid,

pKD46. The strain was grown in 10 ml SOB medium with ampicillin (100 μg/ml) and

L-arabinose at 30°C to an OD_{600} of ~0.5 and then made electrocompetent by washing

and centrifuge them with a cold solution of glycerol 10%. Competent cells in 80 μl

aliquots were electroporated with 200 ng of PCR product. Cells were added

immediately 0.9 ml of SOC, incubated during 2 h at 37°C, and then 100 μl aliquots

spread onto LB agar with kanamycin (30 μg/ml). The correct inactivation of genes was

verified by PCR. The antibiotic resistant cassette (Kn) was removed for both mutants

using the flippase plasmid pCP20.

Transcriptome sequencing

The transcriptome sequencing of primed cells was determined on samples treated

identically as described for the priming experiments. Total RNA from 10^8 cell per sample

was isolated using the RNAeasy Isolation kit (Qiagen, Germany). Traces of genomic

DNA were removed from 10 μg of RNA by digestion in a total volume of 500 μl

containing 20 units of TURBO DNase, incubated for 30 minutes at 37°C, immediately

followed by RNeasy (Qiagen) clean-up and elution in 30 μl of RNase-free water.

Following DNase treatment, RNA integrity was assessed using Agilent RNA 6000 Nano

kit and 2100 Bioanalyzer instrument (both from Agilent Technologies). Total RNA was

depleted from ribosomal RNA using the Ribo-Zero Depletion Kit for Gram-negative

bacteria (Illumina, USA). Libraries were prepared using a TruSeq Stranded Total RNA library preparation kit (Illumina, USA) and were sequenced on a MiSeq platform. Transcript abundances were derived from pseudo-alignment of reads to the cDNA sequences from the ASM584v2 assembly of *Escherichia coli* MG1655 (ENA accession GCA_000005845.2) using Salmon version 0.7.2 with default parameters (Patro et al., 2017). Differential gene expression was analyzed using the R package DESeq2 (Love et al., 2014) in conjunction with tximport (Soneson et al., 2015). Pairwise contrasts were performed between the control and each AMP treatment with empirical bayesian shrinkage of both dispersion parameters and fold-change estimation. We defined genes as being significantly differentially expressed when the absolute fold-change in expression was greater than 2, at an FDR-adjusted p-value of less than 0.05. The variance-stabilizing transformation was used to remove the dependence of the variance on the mean and to transform data to the log2 scale prior to ordination using principal component analysis. Quality of RNAseq data were contrasted by Euclidian distance and symmetry of data reads distribution (Fig. S 11).

Observation at single cell level

To observe cell reaction at single cell level during priming experiments, we used an ad hoc microfluidic device developed for this project. It consisted in a main channel for bacterial inoculation and medium perfusion and with several lateral compartments which dimensions are around 1.5 μm height (ensuring all bacteria are kept in focus) and square 200 μm width corresponding to a field size of the used microscope at 1000x magnification (Fig. S 12). The chip was designed in Autocad. We started the replication of our microfluidic chips from a custom made (Sigatec SA) silicon (SiO) master. This silicon master itself was first replicated in Smooth-Cast 310 (Bentley advanced material). Soft lithography produced the chips in PDMS (Sylgard Silicone Elastomer Base and Curing

Agent mixed in 10:1 ratio). The PDMS chips were cured overnight at 75°C in an

incubator. We punched an inlet and outlet hole for the laminar flow in each chip using a

biopsy puncher of 0.5 mm (outer diameter). The chips were bonded to a glass cover slide

(24×60 mm) after a 30-second air plasma treatment (PDC-002, Harrick Plasma). Before

use, the assembled chip was treated for 15 seconds in air plasma and immediately injected

it with filtered MHB medium for passivation. We left the activated chip to incubate for a

least 1 hour before loading the bacteria. The devices were loaded to full capacity with a

bacterial suspension containing nearly $2x10^8$ CFU/ml (exponentially growing bacteria,

0.5 OD_{600}). Cell suspension was injected into the main channel of the chip using a blunt-

end 23G needle attached to a 1ml syringe. We centrifuged the loaded chip at 200 x g for

10 min using in-house adapters, checked the loading.

After we loaded the bacteria cells in the blind ending side channels, we connected

the chip to a syringe pump (AL-6000, WPI, Germany) and placed the chip under an

inverted microscope. A continuous laminar flow (100µl/h) of MHB through the central

channel was maintained throughout the experiment (Fig. S 12). For the life cell imaging,

after infusion with priming or triggering concentration of AMPs, we injected MHB

supplemented with bacterial Live/Dead stain kit solutions (Thermo Scientific, Germany)

for a final concentration of 0.1 µl/ml of MHB. We took pictures of at least 20 fields per

treatment from independent side channels. Fluorescent images were taken of each field

of view with simultaneous acquisition in red and green fluorescent channels during a time

interval of no more than 2 minutes per treatment with a Nikon Ti-2 inverted microscope

(Nikon, Japan). Cells were observed with the 100× objective and controlled by Nis

Element AR software. The chip holder is temperature controlled at 37°C.

Determination of melittin-induced curli

The production of curli due to melittin treatment was determined by using the fluorescent

dye Ectracer^{TM680} (Ebba Biotech, Sweden) that stain extracellular curli. Ectracer^{TM680}

was used according to the instructions of manufacturers. Bacterial cultures were treated

on chip with priming and triggering concentrations of melittin as described for single cell

observation section omitting the live/dead staining. After priming and triggering, the

channels were perfused MHB supplemented with Ectracer^{TM680} in a proportion of 1/1000

related to the medium. Cells were observed with the red channel fluorescence for Cy5

dye using a Nikon Eclipse Ti2 inverted optical microscope using the 100X oil objective.

Two independent samples were prepared for each group (primed and naïve cells).

SEM of *E. coli* treated antimicrobial peptides

Approximately 2×10^7 CFU/ml *E. coli* MG1655 were treated with 1/10 MIC of pexiganan

and melittin during 30 minutes. The cultures were concentrated 10 times by a quick

centrifugation step of 1 minute at 8 000 g and resuspended in 1/10 of its own supernatant.

and resuspension 10 µl drops were placed on a circular glass cover slip (1.5 cm of

diameter). The drops were fixed with osmium tetroxide vapors during one minute and

allow to dry in a laminar flow cabinet. The cover glasses were mounted on aluminum

stubs using double-sided adhesive tape and coated with gold in a sputter coater (SCD-

040; Balzers, Union, Liechtenstein). The specimens were examined with a FEI Quanta

200 SEM (FEI Co., Hillsboro, OR) operating at an accelerating voltage of 15 kV under

high vacuum mode at different magnifications. At least 5 fields from two independent

replicas.

Statistical analysis

To analyze the priming data, we first tested if the dynamics of the depicted in the time-kill curves are biphasic.

We fitted the function

$$f(m_1, m_2, t_{kink}, t) = \begin{cases} log_{10}\big(CFU(t=0)\big) + m_1 t, & t < t_{kink} \\ log_{10}\big(CFU(t=0)\big) + m_1 t_{kink} + m_2(t - t_{kink}), & t \geq t_{kink} \end{cases}$$

to the time-kill data of each AMP and for primed and naïve populations individually using a least square algorithm. Here, t_{kink} is the time point, at which the population dynamics switch from the first phase to the second phase and m_1 and m_2 are the slopes of the first and the second decline, respectively. Note that m_1 is a direct measure of tolerance. The standard error (SE) was calculated as

$$SE(\theta) = \sqrt{\frac{1}{I(\theta)}}$$

The parameter θ denotes to the estimated parameter values of m_1, m_2, and t_{kink} and $I(\theta)$ is the expected Fisher information. The 95% confidence interval was calculated as $\theta \pm 1.96 * SE(\theta)$.

We used an ANOVA by means of contrasts to assess if priming changes tolerance and persistence. For this, we tested if the decrease of the first phase (all data points with $t < t_{kink}$) differed between primed and naive populations (test for differences in tolerance) and if the population size in the second phase (all data points with $t > t_{kink}$) differed between primed and naive populations (test for differences in persistence). Both tests showed significant differences between primed and naive populations for both melittin

1 and pexiganan (significance level: p < 0.05). We corrected for multiple testing with the

2 Bonferroni-method.

4 **Population models**

5 To describe bacterial population dynamics, we used the two-state model by Balaban

6 *et al.* (Balaban et al., 2004) (Fig. 5 A, B):

$$\frac{dN(t)}{dt} = (r_N(A) - s_N)N(t) + s_P P(t)$$

$$\frac{dP(t)}{dt} = (r_P(A) - s_P)P(t) + s_N N(t)$$

In this model, the bacterial population consists of two subpopulations, one with a

normal phenotype, *N(t),* and a second with a persister phenotype,*P(t)*. We denote the total

bacterial population size by *B(t),*with *B(t) = N(t)+P(t)*. The rate of change of the

population is determined by the net growth rate of N and P, r_N and r_P, and the switching

rate from N to P, s_N, and the switching rate from P to N, s_P. The parameter estimates of

the net growth rates are dependent on the AMP concentration A, i.e. *$r_N(A)$* and *$r_P(A)$*.

analytical solution of this ODE system (Balaban et al., 2004; Komarova and Wodarz,

2007) is

$B(t) = N(t) + P(t) = c_1 u_1 e^{\lambda_1 t} + c_1 u_2 e^{\lambda_1 t} + c_2 v_1 e^{\lambda_2 t} + c_2 v_2 e^{\lambda_2 t}$ Eq. S2

with

λ_1

$= \dfrac{r_N + r_P - s_N - s_P - \sqrt{(-r_N - r_P + s_N + s_P)^2 - 4(r_N r_P - r_P s_N - r_N s_P)}}{2}$

$$\lambda_2 = \frac{r_N + r_P - s_N - s_P + \sqrt{(-r_N - r_P + s_N + s_P)^2 - 4(r_N r_P - r_P s_N - r_N s_P)}}{2}$$

$$\vec{u} = \begin{pmatrix} \dfrac{\lambda_1 - r_P + s_P}{s_N} \\ 1 \end{pmatrix}$$

$$\vec{v} = \begin{pmatrix} \dfrac{\lambda_2 - r_P + s_P}{s_N} \\ 1 \end{pmatrix}$$

$$c_1 = \frac{v_1 P(t = 0) - N(t = 0)}{v_1 - u_1}$$

$$c_2 = \frac{-u_1 P(t = 0) + N(t = 0)}{v_1 - u_1}$$

The model was fitted by a least square algorithm, minimizing the residual sum of

squares of the data to the prediction. For the starting conditions ($N(t\text{=}0)$, $P(t\text{=}0)$), we

assumed that the ratio of N/P was at the equilibrium predicted by the model without

antimicrobials when the exposure to lethal concentrations of AMPs started. $N(t\text{=}0)$ and

$P(t\text{=}0)$ were therefore calculated using the eigenvector $\vec{v}$ that corresponds to the largest

eigenvalue of a system without antimicrobials. Here, we assumed that the parameter r_N is

equal the net growth rate in absence of antimicrobials, $r_N = \psi_{max}$. The parameter ψ_{max}

was estimated based on the time-kill curve of bacterial population that grow in absence

of antimicrobials (see below and Fig. S 11). The eigenvector contains information about

the ratio of N and P for $t \rightarrow \infty$: $\frac{N}{P} = \frac{v_1}{v_2}$. Resulting, $P(t = 0) = \frac{B(t=0)}{1+v_1}$ and $N(t = 0) =$

$B(t = 0) - P(t = 0)$. $B(t = 0)$ was estimated from the data. Confidence intervals were

calculated as described above. In a pre-analysis, we used 4 free model parameters that

were fitted: r_N, r_P, s_N and s_P. The parameter r_P was not significant from 0 (Fig. S 8).

Therefore, we set the parameter r_P to 0 and fitted the remaining 3 parameters to the data

(Fig. S 9, table S 5).

Tolerance and persistence in terms of model parameters

The measure of tolerance is the slope m_1. Komarova and Wodarz (Komarova and Wodarz, 2007) showed that the slope can directly be linked to the population model parameters. In our notation,

$$m_1 = log_{10}(c_1(u_1+v_1)e^{\lambda_1 t}) \quad \text{Eq. S1}$$

Note that the first phase is mainly influenced by r_N (Fig. S9), therefore, $m_1 \approx r_N$.

Persistent cell numbers at time t were calculated with the analytic solution:

$$P(t) = c_1 u_2 e^{\lambda_1 t} + c_2 v_2 e^{\lambda_2 t} \quad \text{Eq. S2}$$

Pharmacokinetic and pharmacodynamic function

We used the pharmacokinetic function $A(t) = A_{max}e^{-k(t-n)}$, with $8h * n \le t \le 8h * (n+1)$ and $n = 0,1,2 \ldots$ described previously elsewhere (Regoes et al., 2004). In our simulations, we fixed the decay parameter k and varied the drug input A_{max}. To describe the effect of the AMPs on the bacterial population, we used the pharmacodynamic (PD) function $\psi(A)$ (Regoes et al., 2004; Zhi et al., 1986), with

$$\psi(A) = \psi_{max} - e(A) = \psi_{max} - \frac{(\psi_{max} - \psi_{min})\left(\frac{A}{MIC}\right)^{\kappa}}{\left(\frac{A}{MIC}\right)^{\kappa} - \frac{\psi_{min}}{\psi_{max}}}$$

The parameter ψ_{max} describes the net growth rate in absence of antimicrobials ($\psi_{max} = \psi(A = 0)$). The antimicrobial effect $e(A)$ is dependent on the antimicrobial concentration and is the defined with ψ_{max}, ψ_{min}, the net growth rate in presence of large amounts of antimicrobials ($\psi(A \to \infty)$), with the MIC, the antimicrobial concentration that results in no growth ($\psi(A = MIC) = 0$) and with κ, which determines the steepness of the PD curve.

The PD function was fitted to the time-kill curves (Fig. S6), as described by Regoes

et al. (Regoes et al., 2004). In short, we used log-linear regressions of the time kill curves

within the time-points 0h and 1h to estimate the change of the bacterial population over

time, i.e. the slopes of the log-linear regression. We fixed the parameter ψ_{max} and fitted

the 3 remaining parameters of the PD function with the *Markov-Chain-Monte-Carlo*

method.

Stochastic simulations

To simulate resistance evolution with stochastic simulations, we expanded a

previously developed framework for bacterial population dynamics (Yu et al., 2018). The

framework models bacterial population dynamics exposed to changing levels of

antimicrobials and allows for resistance evolution. In the simulations, the change in

population size of a sensitive strain *S*, with *S = N+P*, and of a resistant strain *R* were

described with the following ODE system:

$$\frac{dN(t)}{dt} = \psi_{max}(1-\mu)N(t)\left(1 - \frac{N(t)+P(t)+R(t)}{K}\right) - s_N N(t) + s_P P(t)$$

$$- e_N(A(t))N(t)$$

$$\frac{dP(t)}{dt} = s_N N(t) - s_P P(t)$$

$$\frac{dR(t)}{dt} = \psi_{max}\,\mu\,N(t)\left(1 - \frac{N(t)+P(t)+R(t)}{K}\right) + \psi_{max}(1-c)R\left(1\right.$$

$$\left. - \frac{N(t)+P(t)+R(t)}{K}\right) - e_R(A(t))R(t)$$

Here, the replication rate is assumed to be equal to the maximum net growth rate ψ_{max}. The effect of the antimicrobial $e_N(A)$ is explained above. Note that we assume that bacteria in class P do not grow and are not affected by antimicrobials. We also assumed that the switching rates are constant. To describe the effect of an antimicrobial on the strain R, $e_R(A)$, we use the same parameter set than with $e_N(A)$, except for the *MIC*: $\frac{MIC_R}{MIC_N} = 10$. In the simulations, we differentiated between the following cases: (i) naïve bacteria, homogeneous population (no persistence), (ii) naïve bacteria, heterogeneous population, (iii) primed bacteria (increase in r_N), homogeneous population, (iv) primed bacteria (increase in r_N), heterogeneous population (v) primed bacteria (increase in r_N and decrease in s_P). The stochastic simulations were run 1000 times for each antimicrobial, for each case, and for a variety of input antimicrobial concentrations A_{max}. All parameter values are listed in table S 6. The intensity is the input antimicrobial concentrations A_{max}. The simulations were run for t=7d. Time until clearance and probability of resistance evolution were calculated as the mean of the value over the 1000 simulations.

Implementation

Statistical testing, simulations and plots were done in R version 3.3.2 (R Core Team, 2015), using Rstudio version 1.0.143 (R Development Core Team, 2015). We used the following R-packages: (i) for plotting: sfsmisc (Maechler), (ii) for fitting the PD function: rjags(iii) for stochastic simulations: adaptivetau (Cao et al., 2007). We used Mathematica version 11.0 (Wolfram Research, 2016) to determine the analytical solutions of the population models.

REFERENCES

Abel zur Wiesch, P., Abel, S., Gkotzis, S., Ocampo, P., Engelstädter, J., Hinkley, T., Magnus, C., Waldor,

M.K., Udekwu, K., and Cohen, T. (2015). Classic reaction kinetics can explain complex patterns of

antibiotic action. Sci. Transl. Med. *7*, 287ra73-287ra73.

Andersson, D.I., and Hughes, D. (2014). Microbiological effects of sublethal levels of antibiotics. Nat.

Rev. Microbiol. *12*, 465–478.

Andrade-Linares, D.R., Lehmann, A., and Rillig, M.C. (2016). Microbial stress priming: a meta-analysis:

Microbial priming to stress. Environ. Microbiol. *18*, 1277–1288.

Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K.A., Tomita, M.,

Wanner, B.L., and Mori, H. (2006). Construction of Escherichia coli K-12 in-frame, single-gene knockout

mutants: the Keio collection. Mol Syst Biol *2*, 2006 0008.

Balaban, N.Q., Merrin, J., Chait, R., Kowalik, L., and Leibler, S. (2004). Bacterial Persistence as a

Phenotypic Switch. Science *305*, 1622–1625.

Balaban, N.Q., Helaine, S., Lewis, K., Ackermann, M., Aldridge, B., Andersson, D.I., Brynildsen, M.P.,

Bumann, D., Camilli, A., Collins, J.J., et al. (2019). Definitions and guidelines for research on antibiotic

persistence. Nat. Rev. Microbiol. *17*, 441–448.

Barnhart, M.M., and Chapman, M.R. (2006). Curli biogenesis and function. Annu. Rev. Microbiol. *60*,

131–147.

Bevins, C.L., and Salzman, N.H. (2011). Paneth cells, antimicrobial peptides and maintenance of

intestinal homeostasis. Nat. Rev. Microbiol. *9*, 356–368.

Brauner, A., Fridman, O., Gefen, O., and Balaban, N.Q. (2016). Distinguishing between resistance,

tolerance and persistence to antibiotic treatment. Nat. Rev. Microbiol. *14*, 320–330.

Brogden, K.A. (2005). Antimicrobial peptides: pore formers or metabolic inhibitors in bacteria? Nat. Rev.

Microbiol. *3*, 238–250.

Brown, A.J.P., Gow, N.A.R., Warris, A., and Brown, G.D. (2019). Memory in Fungal Pathogens

Promotes Immune Evasion, Colonisation, and Infection. Trends Microbiol. *27*, 219–230.

Cao, Y., Gillespie, D.T., and Petzold, L.R. (2007). Adaptive explicit-implicit tau-leaping method with

automatic tau selection. J. Chem. Phys. *126*, 224101.

Chen, F., Krasity, B.C., Peyer, S.M., Koehler, S., Ruby, E.G., Zhang, X., and McFall-Ngai, M.J. (2017).

Bactericidal Permeability-Increasing Proteins Shape Host-Microbe Interactions. MBio *8*, e00040-17,

/mbio/8/2/e00040-17.atom.

Cheung, G., Fisher, E., McCausland, J., Choi, J., Collins, J., Dickey, S., and Otto, M. (2018).

Antimicrobial peptide resistance mechanism contributes to Staphylococcus aureus infection Accepted. J.

Of 1–34.

Datsenko, K.A., and Wanner, B.L. (2000). One-step inactivation of chromosomal genes in Escherichia

coli K-12 using PCR products. Proc Natl Acad Sci U A *97*, 6640–6645.

Fantner, G.E., Barbero, R.J., Gray, D.S., and Belcher, A.M. (2010). Kinetics of antimicrobial peptide

activity measured on individual bacterial cells using high-speed atomic force microscopy. 1–6.

Fjell, C.D., Hiss, J.A., Hancock, R.E.W., and Schneider, G. (2012). Designing antimicrobial peptides:

form follows function. Nat. Rev. Drug Discov. *11*, 37–51.

Ge, Y., Macdonald, D.L., Holroyd, K.J., Thornsberry, C., Wexler, H., and Zasloff, M. (1999). In Vitro

Antibacterial Properties of Pexiganan, an Analog of Magainin. ANTIMICROB AGENTS CHEMOTHER

43, 7.

Gottler, L.M., and Ramamoorthy, A. (2009). Structure, membrane orientation, mechanism, and function

of pexiganan — A highly potent antimicrobial peptide designed from magainin. Biochim. Biophys. Acta

BBA - Biomembr. *1788*, 1680–1686.

Haine, E.R., Moret, Y., Siva-Jothy, M.T., and Rolff, J. (2008). Antimicrobial Defense and Persistent

Infection in Insects. Science *322*, 1257–1259.

Hedges, A.J. (1966). An examination of single-hit and multi-hit hypotheses in relation to the possible

kinetics of colicin adsorption. J. Theor. Biol. *11*, 383–410.

Hilker, M., Schwachtje, J., Baier, M., Balazadeh, S., Bäurle, I., Geiselhardt, S., Hincha, D.K., Kunze, R.,

Mueller-Roeber, B., Rillig, M.C., et al. (2016). Priming and memory of stress responses in organisms

lacking a nervous system: Priming and memory of stress responses. Biol. Rev. *91*, 1118–1133.

Johnston, P.R., Paris, V., and Rolff, J. (2019). Immune gene regulation in the gut during metamorphosis

in a holo- versus a hemimetabolous insect. Philos. Trans. R. Soc. B Biol. Sci. *374*, 20190073.

Joo, H.-S., Fu, C.-I., and Otto, M. (2016). Bacterial strategies of resistance to antimicrobial peptides.

Philos. Trans. R. Soc. B Biol. Sci. *371*, 20150292.

Komarova, N.L., and Wodarz, D. (2007). Effect of Cellular Quiescence on the Success of Targeted CML

Therapy. PLoS ONE *2*, e990.

Koo, H.B., and Seo, J. (2019). Antimicrobial peptides under clinical investigation. Pept. Sci. *111*.

Levin, B.R., and Rozen, D.E. (2006). Non-inherited antibiotic resistance. Nat. Rev. Microbiol. *4*, 556–

562.

Levin-Reisman, I., Ronin, I., Gefen, O., Braniss, I., Shoresh, N., and Balaban, N.Q. (2017). Antibiotic

tolerance facilitates the evolution of resistance. Science *355*, 826–830.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. *15*, 550.

Maechler, M. Utilities from "Seminar fuer Statistik" ETH Zurich [R package sfsmisc version 1.1-4].

Makarova, O., Rodriguez-Rojas, A., Eravci, M., Weise, C., Dobson, A., Johnston, P., and Rolff, J. (2016).

Antimicrobial defence and persistent infection in insects revisited. Philos. Trans. R. Soc. B Biol. Sci. *371*,

20150296.

Maria-Neto, S., de Almeida, K.C., Macedo, M.L.R., and Franco, O.L. (2015). Understanding bacterial

resistance to antimicrobial peptides: From the surface to deep inside. Biochim. Biophys. Acta BBA -

Biomembr. *1848*, 3078–3088.

Mergaert, P. (2018). Role of antimicrobial peptides in controlling symbiotic bacterial populations. Nat.

Prod. Rep. *35*, 336–356.

Miajlovic, H., and Smith, S.G. (2014). Bacterial self-defence: how Escherichia coli evades serum killing.

FEMS Microbiol. Lett. *354*, 1–9.

Mitchell, A., Romano, G.H., Groisman, B., Yona, A., Dekel, E., Kupiec, M., Dahan, O., and Pilpel, Y.

(2009). Adaptive prediction of environmental changes by microorganisms. Nature *460*, 220–224.

Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. (2017). Salmon provides fast and

bias-aware quantification of transcript expression. Nat. Methods *14*, 417–419.

Perron, G.G., Zasloff, M., and Bell, G. (2006). Experimental evolution of resistance to an antimicrobial

peptide. Proc. R. Soc. B Biol. Sci. *273*, 251–256.

Prigent-Combaret, C., Prensier, G., Le Thi, T.T., Vidal, O., Lejeune, P., and Dorel, C. (2000).

Developmental pathway for biofilm formation in curli-producing Escherichia coli strains: Role of

flagella, curli and colanic acid. Environ. Microbiol. *2*, 450–464.

R Core Team (2015). R: A language and environment for statistical computing. (Vienna, Austria: R

Foundation for Statistical Computing).

R Development Core Team (2015). RStudio Team (2015). RStudio: Integrated Development for R.

Raghuraman, H., and Chattopadhyay, A. (2007). Melittin: a Membrane-active Peptide with Diverse

Functions. Biosci. Rep. *27*, 189–223.

Regoes, R.R., Wiuff, C., Zappala, R.M., Garner, K.N., Baquero, F., and Levin, B.R. (2004).

Pharmacodynamic Functions: a Multiparameter Approach to the Design of Antibiotic Treatment

Regimens. Antimicrob. Agents Chemother. *48*, 3670.

Rodríguez-Rojas, A., Makarova, O., and Rolff, J. (2014). Antimicrobials, Stress and Mutagenesis. PLoS

Pathog. *10*, e1004445.

Shan, Y., Brown Gandt, A., Rowe, S.E., Deisinger, J.P., Conlon, B.P., and Lewis, K. (2017a). ATP-

Dependent Persister Formation in Escherichia coli. MBio *8*, e02267-16.

Shan, Y., Brown Gandt, A., Rowe, S.E., Deisinger, J.P., Conlon, B.P., and Lewis, K. (2017b). ATP-

Dependent Persister Formation in *Escherichia coli*. MBio *8*, e02267-16, /mbio/8/1/e02267-16.atom.

Soneson, C., Love, M.I., and Robinson, M.D. (2015). Differential analyses for RNA-seq: transcript-level

estimates improve gene-level inferences. F1000Research *4*, 1521.

Spohn, R., Daruka, L., Lázár, V., Martins, A., Vidovics, F., Grézal, G., Méhi, O., Kintses, B., Számel, M.,

Jangir, P.K., et al. (2019). Integrated evolutionary analysis reveals antimicrobial peptides with limited

resistance. Nat. Commun. *10*, 4538.

Vadyvaloo, V., Viall, A.K., Jarrett, C.O., Hinz, A.K., Sturdevant, D.E., and Hinnebusch, B.J. (2015).

Role of the Phop-PhoQ gene regulatory system in adaptation of yersinia pestis to environmental stress in

the flea digestive tract. Microbiol. U. K. *161*, 1198–1210.

Windels, E.M., Michiels, J.E., Fauvart, M., Wenseleers, T., Van den Bergh, B., and Michiels, J. (2019).

Bacterial persistence promotes the evolution of antibiotic resistance by increasing survival and mutation

rates. ISME J. *13*, 1239–1251.

Wiuff, C., Zappala, R.M., Regoes, R.R., Garner, K.N., Baquero, F., and Levin, B.R. (2005). Phenotypic

Tolerance: Antibiotic Enrichment of Noninherited Resistance in Bacterial Populations. Antimicrob.

Agents Chemother. *49*, 1483–1494.

Wolfram Research, Inc. (2016). Mathematica, Version 11.0.

Yao, Z., Kahne, D., and Kishony, R. (2012). Distinct Single-Cell Morphological Dynamics under Beta-

Lactam Antibiotics. Mol. Cell *48*, 705–712.

Yu, G., Baeder, D., Regoes, R., and Rolff, J. (2018). Predicting Drug Resistance Evolution: Antimicrobial

Peptides Vs. Antibiotics. Proc R Soc Lond B 138107.

Zasloff, M. (2002). Antimcirobial peptides of multicellular Organisms. Nature *415*, 389–395.

Zhi, J., Nightingale, C.H., and Quintiliani, R. (1986). A pharmacodynamic model for the activity of

antibiotics against microorganisms under nonsaturable conditions. J. Pharm. Sci. *75*, 1063–1067.

Supplement

Supplementary tables S1-S6 page 1-6

Supplementary references 7-8

Supplementary figures S1-S12 9-21

Supplementary tables

Table S 1. Minimal inhibitory concentration of the two antimicrobial peptides melittin

and pexiganan and the antibiotic ciprofloxacin used in this work. The MIC was

determined to be used as a concentration reference but it was also determined after

priming treatment to show that the observed enhanced survival is based on a phenotypic

changes and not mutations.

Antimicrobials	MIC (µg/ml) pre-priming	MIC (µg/ml) post-priming
melittin	2	2
pexiganan	1	1
ciprofloxacin	0.125	0.125

1 **Table S 2**. Slopes of fitting the biphasic function with the slopes m_1 and m_2 to the data on

2 a $\log_{10}$ scale. Significant differences between slopes of primed and naïve population

3 counts are indicated with asterisks for each antimicrobial.

Antimicrobial	Treatment	Slope			
		m_1 (± SE)		m_2 (± SE)	
melittin	Primed	-3.9 (± 0.6)	*	-0.3 (± 0.4)	
	Naïve	-6.3 (± 0.6)		-1.0 (± 0.4)	
pexiganan	Primed	-2.9 (±0.3)	*	-0.6 (± 0.894)	
	Naïve	-4.1 (± 0.3)		-0.3 (± 0.9)	

2   Table S 3. Priming response to pexiganan (a) and melittin (b) (0.1xMIC, 30-minute

3   treatment)

5   *See excel files.*

1 **Table S 4.** Results of fitting the two-state model to bacterial population dynamics in the

2 presence of melittin and pexiganan and the classic exponential population growth model

3 to dynamics in the presence of ciprofloxacin and ampicillin (rounded values).

Antimicrobial	Treatment	r_N ($\pm$ SE)	r_P[1]	s_N ($\pm$ SE)	s_P ($\pm$ SE)
melittin	Primed	-8.96 (0.72)	0	0.00085 (0.00098)	0.27 (0.61)
	Naïve	-15.06 (0.86)	0	0.00128 (0.00095)	2.25 (0.45)
pexiganan	Primed	-6.68 (0.32)	0	0.00027 (0.00059)	0.40 (1.10)
	Naïve	-9.61 (0.40)	0	0.00021 (0.00025)	1.26 (0.66)

5 [1] Parameter r_P is fixed at 0 due to results of the preliminary analysis, in which the

6 parameter was not significantly different from 0 (see Figure S 8).

 Table S 5. Parameter values used as input for the stochastic simulations.

Parameter	Description	AMP	Value
k	AMP degradation rate		0.3
μ	Mutation rate		$1*10^{-9}$
K	Carrying capacity		$5*10^9$
ψ_{max}	Maximal net growth rate in absence of antimicrobials	melittin	2.13
		pexiganan	2.06
ψ_{min}	Minimal net growth rate in presence of large amounts of antimicrobials	melittin	-8.86/ -9.89[§]
		pexiganan	-36.87/-66.80[§]
κ	Hill parameter	melittin	1.3/1.1[§]
		pexiganan	0.6/0.7[§]
MIC	Dose of antimicrobials for which the net growth rate is 0	melittin	0.48/1.83[§]
		pexiganan	0.54/2.14[§]
c	Cost of resistance		0.5
$B(t{=}0)$	Population size at time t=0		10^8
t_{max}	Maximal run of simulation		7d
s_N	Switching rate from N to P	melittin	0.0012
		pexiganan	0.00021
s_P	Switching rate from P to N	melittin	2.26/0.28[§]
		pexiganan	1.26/0.40[§]

3   [§] Naïve/Primed

23   Table S 6. Strains and plasmids used in this work and their relevant phenotypes
24

Strain	Relevant phenotype	Reference

E. coli MG1655 WT	F⁻ λ⁻ *ilvG⁻ rfb-50 rph-1*	Lab stock
E. coli MG1655 *csgA*::Kn	*csgA*::Kn F⁻ λ⁻ *ilvG⁻ rfb-50 rph-1*	this work
E. coli MG1655 *csgA*::scar	*csgA*::scar* F⁻ λ⁻ *ilvG⁻ rfb-50 rph-1*	this work
E. coli MG1655 *wza*::Kn	*wza*::Kn F⁻ λ⁻ *ilvG⁻ rfb-50 rph-1*	this work
E. coli MG1655 *wza*::scar*	*wza*::scar* F⁻ λ⁻ *ilvG⁻ rfb-50 rph-1*	this work
E. coli BW25113 *csgA*::Kn	*csgA*::Kn lacI⁺rrnB$_{T14}$ ΔlacZ$_{WJ16}$ hsdR514 ΔaraBAD$_{AH33}$ ΔrhaBAD$_{LD78}$ rph-1 Δ(araB–D)567 Δ(rhaD–B)568 ΔlacZ4787(::rrnB-3) hsdR514 rph-1	Baba et al., 2006
E. coli BW25113 *wza*::Kn	*wza*::Kn lacI⁺rrnB$_{T14}$ ΔlacZ$_{WJ16}$ hsdR514 ΔaraBAD$_{AH33}$ ΔrhaBAD$_{LD78}$ rph-1 Δ(araB–D)567 Δ(rhaD–B)568 ΔlacZ4787(::rrnB-3) hsdR514 rph-1	Baba et al., 2006
Plasmids	**Relevant phenotype**	**Reference**
pKD46	oriR101 w/repA101ts Amp^R	Datsenko and Wanner, 2000
pCP20	oriR101 w/repA101ts Amp^R	Datsenko and Wanner, 2000

1   * asterisk indicates the removal of the antibiotic resistant cassette kanamycin (Kn) by

2   the action of the flippase expressed by the plasmid pCP20 (Datsenko and Wanner,

3   2000).

Supplementary References

1. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware quantification of transcript expression. Nat Methods [Internet]. Nature Publishing Group; 2017 [cited 2019 Jul 3];14:417–9.

2. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol [Internet]. 2014 [cited 2019 Jul 3];15:550.

3. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. F1000Research [Internet]. 2015 [cited 2019 Jul 3];4:1521.

4. Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S. Bacterial Persistence as a Phenotypic Switch. Science (80-) [Internet]. 2004 [cited 2018 Jun 9];305:1622–5.

5. Komarova NL, Wodarz D. Effect of Cellular Quiescence on the Success of Targeted CML Therapy. Agur Z, editor. PLoS One [Internet]. 2007 [cited 2018 Jun 9];2:e990.

6. Regoes RR, Wiuff C, Zappala RM, Garner KN, Baquero F, Levin BR. Pharmacodynamic Functions: a Multiparameter Approach to the Design of Antibiotic Treatment Regimens. Antimicrob Agents Chemother [Internet]. American Society for Microbiology (ASM); 2004 [cited 2019 May 30];48:3670. Available from: http://www.ncbi.nlm.nih.gov/pubmed/15388418

7. Yu G, Baeder D, Regoes R, Rolff J. Predicting Drug Resistance Evolution: Antimicrobial Peptides Vs. Antibiotics. Proc R Soc Lond B. 2018;138107.

8. R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2015.

9. R Development Core Team. RStudio Team (2015). RStudio: Integrated Development for R. [Internet]. Boston: MA URL http://www.rstudio.com/; 2015. Available from: http://www.rstudio.com

10. Maechler M. Utilities from "Seminar fuer Statistik" ETH Zurich [R package sfsmisc version 1.1-4]. Comprehensive R Archive Network (CRAN); [cited 2019 Jul 3]; Available from: https://cran.r-project.org/web/packages/sfsmisc/index.html

11. Cao Y, Gillespie DT, Petzold LR. Adaptive explicit-implicit tau-leaping method with automatic tau selection. J Chem Phys [Internet]. 2007 [cited 2019 Jul 3];126:224101. Available from: http://aip.scitation.org/doi/10.1063/1.2745299

12. Wolfram Research I. Mathematica, Version 11.0. Champaign, Illinois: Wolfram Research, Inc.; 2016.

13. Wiuff C, Zappala RM, Regoes RR, Garner KN, Baquero F, Levin BR. Phenotypic

tolerance: antibiotic enrichment of noninherited resistance in bacterial populations.

Antimicrob Agents Chemother [Internet]. 2005 [cited 2016 Jan 6];49:1483–94.

Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1068602&tool=pmcentrez&

rendertype=abstract

14. Nielsen EI, Friberg LE. Pharmacokinetic-Pharmacodynamic Modeling of

Antibacterial Drugs. Pharmacol Rev [Internet]. 2013 [cited 2019 Jul 4];65:1053–90.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/23803529

Supplementary Figures

Figure S 1. Bacterial tolerance and persistence change the shape of time-kill curves (figure adapted from Brauner et al. (*44*). (A) Tolerance is the ability of bacteria to longer survive exposure to antimicrobials due to decrease in susceptibility. Tolerance is quantified as increase in the slope of the time kill curve. (B) Persistence is the phenomenon of a subpopulation being less susceptible to the antimicrobial than the rest of the population. A persistent subpopulation manifests as a biphasic decline of bacterial population when exposed to lethal concentrations of antimicrobials. Here, the population consists predominantly of the less susceptible persistent subpopulation. (C) Together, tolerance and persistence result in biphasic time-kill curves with decreased susceptibility in the first phase.

Figure S 2. A possible explanation for a nonlinear decline in populations would be a

decrease in active AMP concentration over time due to degradation (*45*, *46*), We tested

this alternative explanation. As in Figure 1 (main text), we measured the population

dynamics of bacteria exposed to 10 x MIC (round 1). At the end of the experiment, we

sampled the supernatant. In round 2, a fresh bacterial population was exposed to the

sampled supernatant. (A) and (B) show survival data of both rounds for melittin and

pexiganan, respectively. The trend line depicts the median of the population size at each

time-point. We tested differences in population size over time and differences between

round one and two for each antimicrobial peptide with an ANOVA and adjusted with the

Bonferroni method for multiple testing. For both, melittin and pexiganan, the population

size changed significantly over time, while the differences in population size between

round one and two and the interaction between time point and round were not significant

($p < 0.05$).

A

melittin

B

pexiganan

2 **Figure S 3:** Net growth rates resulting from the fit of the two-state model with 4 free

3 parameters. For both (A) melittin and (B) pexiganan, r_P is not significantly different from

4 0. Significant differences between naive and primed treatment are indicated with

5 asterisks.

Figure S 4. Parameter (A) r_N, (B) s_N, and (C) s_P of the two-state model fitted to the data

of primed (orange) and naïve (blue) bacteria. The parameter r_P was set to 0. Significant

differences between naïve (blue) and primed (orange) parameter values are indicated with

asterisks. For parameter values, see Table S 5.

Figure S 5. Diagrammatic representation of (A) the two-state model (*17*) (see also Fig.

5A) and (B) our previously developed framework (*38*), which we extended by a persistent

class according to *(17)*. Here, bacteria that replicate mutate with a mutation rate **μ** to a

more resistant subpopulation R.

Figure S 6. Pharmacokinetic profile used in our stochastic simulations.

A

B

C

D

F**igure S 7.** Time-kill curves and PD curves of bacterial population dynamics exposed to

AMPs. Time-kill experiments in which naïve (blue) and primed (orange) bacteria were

exposed to (A) melittin and (B) pexiganan. The AMP dose used in each time-kill

experiment is listed in the legend in each plot. PD functions of melittin (C) and pexiganan

(D) fitted to the data in (A) and (B), respectively. Note that we excluded data points of

experiments in which naïve bacteria were exposed to melittin (40 µg/ml) from the

analysis to ensure the best fit. Parameter values are listed in table S5.

2 **Figure S 8.** Time until clearance and probability of resistance evolution for bacteria

3 exposed to pexiganan. In (A) each dot is an individual run.

—— naive, homogeneous
—— naive, heterogeneous
—— primed, homogeneous
—— primed, heterogeneous
—— primed, heterogeneous, smaller s_P

Figure S 9. Predictions for different parameter values. (A) Time until clearance and (B) probability of resistance evolution is affected by the AMP decay rate k (dashed lines: k = 0.1, solid lines: k = 0.3). In (C) and (D) the mutation rate is varied (dashed lines: μ = 10^{-11}, solid lines: μ = 10^{-9}). All simulations are based on melittin data.

A

B

C

D

Figure S 10. Two-state model predicts biphasic decline depending on the model parameter values. If not varied, $r_N = -10$, $r_P = 0$, $s_N = 0.001$, and $s_P = 1$. Tolerance, i.e. the slope m_1 is mainly influenced by r_N, while the levels of persistence is influenced by all four parameters.

Figure S 11. Quality control of RNA sequencing by evaluating symmetry and distribution of the transcriptome counts, volcano plots showing different degrees of significance (A) and assessing dissimilarities of sample-based Euclidian hierarchical clustering for cells treated with priming concentrations of melitting and pexiganan (B). Datasets are based on RNAseq of three independent biological replicas.

3 Figure S 12. Microfluidic device design used for live imaging of priming and killing of
4 *E. coli* MG1655 by AMPs. Z-values refers to the depth of chip features while X and Y
5 represent the dimensions of 2D axes. Each chamber square compartment was designed to
6 be similar in size (200 µm) to a microscope field with a magnification of 1000X.
7