Bacteria primed by antimicrobial peptides develop tolerance and persist

Authors: Alexandro Rodríguez-Rojas[1§], Desiree Y. Baeder[2§], Paul Johnston[3,4], Roland R. Regoes[2*] and Jens Rolff[1,5*].

Affiliations:

[1]Freie Universität Berlin, Institut für Biologie, Evolutionary Biology, Königin-Luise-Strasse 1-3, 14195 Berlin, Germany.

[2]Institute of Integrative Biology, Universitätsstrasse 16 ETH Zürich, 8092 Zürich, Switzerland

3. Berlin Center for Genomics in Biodiversity Research, Berlin, Germany;

4. Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin, Germany;

5. Berlin-Brandenburg Institute of Advanced Biodiversity Research (BBIB), Berlin, Germany

*Correspondence: roland.regoes@env.ethz.ch, jens.rolff@fu-berlin.de,

§These authors contributed equally to this work

SUMMARY

Antimicrobial peptides (AMPs) are key components of innate immune defenses. Because of the antibiotic crisis, AMPs have also come into focus as new drugs. Here, we explore whether prior exposure to sublethal doses of AMPs increases bacterial survival and abets the evolution of resistance. We show that *Escherichia coli* primed by sublethal doses of AMPs develop tolerance and increase persistence by producing curli or colanic acid. We develop a population dynamic model that predicts that priming delays the clearance of infections and fuels the evolution of resistance. The effects we describe should apply to many AMPs and other drugs that target the cell surface. The optimal strategy to tackle tolerant or persistent cells requires high concentrations of AMPs and fast and long-lasting expression. Our findings also offer a new understanding of non-inherited drug resistance as an adaptive response and could lead to measures that slow the evolution of resistance.

INTRODUCTION

Antimicrobial peptides - short, usually cationic peptides - are key effectors of innate immune defences of all multicellular life (*1*) and are also important players at the host microbiota interface (*2, 3*). Because of their evolutionary success and diversity, AMPs are considered as new antimicrobial drugs to alleviate the antibiotic resistance crisis (*4*) with currently more than two dozen under clinical trial (*5*). *Bona fide* genetic resistance in AMPs has been studied (*6, 7*), but not to the extent of antibiotic resistance. Resistance against AMPs evolves usually with a low probability and the levels of resistance are not as high as against antibiotics (*8, 9*).

Notably, non-inherited resistance (*10*), the ability of bacteria to survive lethal concentrations of antimicrobials without a genetically encoded resistance mechanism, has hardly been studied for AMPs. One of the very rare studies that has addressed this in a natural host-microbe interaction is the example of the bobtail squid and its symbiont *Vibrio fischeri*. Here, phenotypic resistance is elicited by a low pH and primes the *Vibrios* to colonize the light-emitting organ of the squid (*11*) in the presence of high concentrations of AMPs. Microbes have evolved adaptive physiological alterations to predictable environmental changes (*12, 13*). This has been studied in the context of available carbon sources during gut passage (*12*), and a meta-analysis found that priming, the phenotypic response to a low level stressor, provides a fitness benefit for microbes against a variety of stressors including pH, temperature and oxidative stress (*14*).

Here we study if the previous encounter with sub-lethal dosages of AMPs induces tolerance or persistence. Such low concentrations are common. For example, after infection the induction of AMPs results in sublethal concentrations before killing concentrations are reached. This usually takes a few hours (*15*). Such pharmacokinetic profiles are mirrored in many antimicrobial drug treatments. During medical application

of antibiotics, the pharmacokinetics start at zero and the killing concentrations are building up over time.

Resistance can be induced by sublethal levels of antimicrobials (*16*). Non-inherited resistance, resulting in either drug tolerance or persister cell formation (see Fig. S 1, (*17*) has been shown to be of great importance to understand antibiotic resistance (*10, 16–18*). It can also facilitate *bona fide* resistance evolution against antibiotics (*18*). We adopt these findings and concepts from antibiotic research to understand resistance against AMPs, an insight which should equally inform our understanding of host-microbe interactions as well as resistance evolution against AMPs as drugs. It is noteworthy that AMPs differ significantly from conventional antibiotics in several aspects including their pharmacodynamics, resulting in narrower mutant selection windows (*9*). Also, the speed at which they kill cells is fast, as they kill within minutes (*19*) rather than within hours, as it is the case for antibiotics (*20*).

In this study, we specifically investigate whether prior exposure to sublethal doses of AMPs increases bacterial survival via either tolerance or persistence, and the risk of resistance evolution. We use two antimicrobial peptides that are well characterised as a case study. Melittin is a 26 amino acid residue linear peptide from the honeybee, which kills bacterial cells by permeabilization of the membrane (*21*). Melittin is also active against eukaryotic parasites such as Leishmania and cancer cells. The other AMP used here, pexiganan, is the first eukaryotic AMP developed as a drug, a synthetic 22 amino acid residue peptide closely related to magainin from the African clawed frog (*22*). It shows a broad activity against both, gram$^+$ and gram$^-$ bacteria and kills by forming toroidal pores (*23*). We combine *in vitro* experiments with a modelling approach to study priming and resistance emergence. We find that a sublethal dose of certain antimicrobial

peptides can induce increased tolerance and/or persistence in bacteria and hence prime

(24) them for the exposure to a subsequent lethal dose. We identify candidate underlying

molecular mechanisms and capture the population dynamics by adapting a classic

mathematical model of persistence **(25)**. With computer simulations, we then predict that

increasing tolerance and persistence will have a positive effect on bacterial survival and

the emergence of resistance.

RESULTS

Primed cells are more resistant

We primed *E. coli* K-12 in vitro by exposing them to a fraction of the minimal inhibitory

concentration of an AMP (0.1xMIC, table S1). Subsequently, we exposed the primed

bacterial populations to lethal concentrations of the respective AMPs (10 x MIC, table

S1) and then monitored bacterial survival over time. We found that the priming treatment

resulted in much higher *E. coli* survival (Fig. 1 A, B).

A   B

Figure 1: Bacterial tolerance and persistence determine the shape of time-kill curves. Time-kill

experiments with *E. coli* K-12, primed (orange) and naïve (blue) bacteria were exposed to 10xMIC of (A)

melittin and (B) pexiganan. At each timepoint indicated, we measured the bacterial population size 5 times. We tested with an ANOVA by means of contrasts if priming influenced tolerance and persistence. For both antimicrobials, priming significantly increased the slope of the first phase, the measure of tolerance, and the bacterial level in the second phase, the measure of persistence, (significance level: $p < 0.05$). We corrected for multiple testing with the Bonferroni-method. The line in the plots indicates the best fit of a biphasic function (table S2), on which our statistical analysis is based.

Killing is biphasic

The decline of the time kill curves is biphasic suggesting two subpopulations (*25–28*). We excluded that deviations from monophasic decline arise because of decreasing antimicrobial concentrations over time (Fig. S 2) and fitted the time-kill curves to a biphasic linear function. For both AMPs, bacterial populations declined faster during the first than the second phase (Fig. 1, Table S 2). Tolerance, the decline of bacterial populations in the first phase (*17*), was significantly higher in primed than in naïve bacteria for both AMPs. Primed bacteria showed higher survival in the second phase, indicating higher numbers of persisters. The change in population size in the second phase, however, was not significantly different between primed and naïve populations, indicating that the population dynamics as such in the second phase are not influenced by priming (Fig. 1). In short, priming with AMPs allow bacteria to survive better by increasing both bacterial tolerance and persistence.

Priming is mediated by either curli or colanic acid

To understand how priming leads to tolerance and persistence, we used RNAseq of cells exposed to priming concentrations of AMPs (Fig. 2 A, B). Exposure to sublethal concentrations of pexiganan (0.1xMIC, as above) induced colanic acid synthesis (Fig. 3 A). Colanic acid capsules have been shown to protect against AMPs and antibiotics (**29**). The removal of an essential gene for colanic acid production completely abolished the

priming effect by pexiganan (Fig 3 B). By phase contrast imaging, we observed the

formation of a characteristic colanic acid capsule in pexiganan-primed but not in naïve

cells (Fig. 3 C). The priming response was homogenous, with all observed cells

producing a colanic acid capsule under priming conditions. Exposure to a sublethal

concentration of melittin (0.1xMIC, as above) induced up-regulation of curli fimbriae

(Fig 3 D). Curli is an important virulence factor **(30)** and a component of extra-cellular

matrix that protects against AMPs **(31)**. A curli-deficient mutant showed a decrease of the

priming effect induced by melittin (Fig. 3 B). For melittin-primed bacteria, curli induction

was documented with a specific curli binding chemical (Fig. 3D). Both AMPs also

induced significant overlap in gene expression related to osmotic shock (Fig. 2).

Figure 2. Gene expression in primed *E. coli*. (A) Venn diagrams showing specific and overlapping

responses of *E. coli* MG1655 to priming concentrations of melittin and pexiganan   (B) Principal

component 1 separates the control from the peptide priming, PC 2 separates the melittin induced response

from the pexiganan response.

18
19 Figure 3 A

6 Figure 3B

1 Figure 3C

3 Figure 3D

naïve cells

primed cells

10 μm

Figure 3. Physiological response to AMP priming. (A) SEM of E. coli MG1655 treated with 1/10xMIC

(priming concentration) of pexiganan (top row) and melittin (bottom row) and non-treated bacteria (middle

row, control). No apparent difference was noticed between melittin-treated cells and control. In the case of

pexiganan, the treated cells tend to aggregate, a phenotype that consistent with the presence of colanic acid.

Red arrows represent potential areas altered by the capsule of colanic acid that collapse under drying

processing necessary for the preparation. Bacteria were observed with different magnifications ranging

from 3000X to 40000X.(B). Bacterial mutants in *csgA* (curli mutant) and *wza* (colanic acid mutant) were

exposed to the AMPs melittin and pexiganan respectively (10xMIC) after priming. Colony forming units

(CFU) were determined after two hours of exposure. Boxplot show data tested by repeated-measures of

one-way ANOVA and Dunnetts' tests. For each case, equal letter represents no statistical differences while

differing letter detonates significant ones (for $p<0.05$). (C). *E. coli* MG1655 treated with 1/10xMIC

(priming concentration) of pexiganan (left) and non-treated bacteria (right, control) observed under phase

contrast optical microscopy. The specimens consisted of cells suspended in a 0.1 % solution of nigrosin to

create a strong contract to visualize the colanic acid capsules. Bacteria were observed with magnifications 1000X after exposure to a trigger stimulus. (D). Detection of curli production by primed cells after exposure to a killing dose of melittin (10xMIC). Curli production is only detected in small proportion of primed cells (survival fraction). After treatment, we exposed the cells to ECtracer™680 (Ebba Biotech, Sweden), a red fluorescent tracer molecule for staining of curli.

After lethal exposure to melittin and pexiganan we observed a high degree of heterogeneity regarding the killing of individual cells in primed bacteria, with many cells surviving the killing concentrations (Fig. 4 A). By contrast, almost all the control bacteria were killed (Fig. 4 A). As mellitin and pexiganan both damage the membrane, which ultimately leads to cell death, the live-dead stain despite it's limitations, seems to be highly appropriate here. Primed cells also seem to aggregate, with a stronger effect in pexiganan-treated cells compared to melittin-treated cells (Fig. 4 A).

Figure 4 A

1   Figure 4 B

Figure 4. Cell viability and persister formation after AMP priming. (A) Cell viability after treating

with priming and trigger doses of melittin and pexiganan were determined using the live/dead BacLight

Bacterial Viability Kit (Thermo Scientific, Germany) on chip as described in M&M. After priming during

30 minutes, the treatment was removed by perfusing fresh MHB. The fluorescence signal was analyzed via

a using excitation at 485 nm and emission at 530 nm for green fluorescence (Syto9) and using excitation at

485 nm and emission at 630 nm for red fluorescence (propidium iodide). (B) The priming concentrations

of pexiganan and melittin, respectively, increased the persister number as determined by treated primed and

naïve populations (10^8 CFU/ml) with ciprofloxacin (10x MIC, 2µg/ml). This increase in persistence co-

occurs with ATP leakage (determined in the supernatant of the culture medium).

To determine if priming stimulates persistence we used an assay based on the observation

that a decrease of intracellular ATP increases persistence under the antibiotic

ciprofloxacin (*32*). Because ATP leakage is a hallmark of AMP-treated bacteria (*33*), we

exposed melittin and pexiganan primed populations and controls to ciprofloxacin (*34*).

This resulted in a highly significant increase in the number of persisters (Fig. 4 B). The

level of leaked ATP in the culture supernatant was significantly higher in primed bacteria

for both AMPs as compared to controls (Fig. 4 B). The pre-treatment with melittin or

pexiganan does not change the MIC of *E. coli* to melittin, pexiganan or ciprofloxacin

consistent with the definition of persisters (*17*).

A mathematical model predicts resistance evolution of primed cells

To quantify the influence of priming on bacterial tolerance and persistence, phenomena

inherently linked to the growth dynamics and subpopulation structure, population

dynamical models that capture tolerance and persistence are required. We build on a

two-state population model previously developed (*25*) to describe bacterial antibiotic

persistence (Fig. 5 A, B). This model assumes that bacteria exist in two phenotypic

states, normal cells (*N*) and persisters (*P*). The two subpopulations *N* and *P* differ in

their susceptibility to AMPs, a difference that is implemented as differing net growth

rates for the given amount of AMPs (A), with $r_N(A)$ and $r_P(A)$, respectively. Bacteria

switch from subpopulation *N* to *P* with the rate s_N and back with the rate s_P.

We quantified the model parameters by fitting the analytical solution of the model

equations (Fig. 5 B) to the time-kill datasets of melittin and pexiganan. In a preliminary

analysis with fitting all four parameters (rN, sN, rP, and sP), the net replication rate of

persister cells, rp, was not significantly different from 0 (Fig. S 4). We therefore fixed

rP to 0. The resulting fit described our data well (Fig. 5 C, D, and Table S 5). We found

that priming affected two of the three estimated model parameters (Fig. S 4): rN

increased, which translates into increase in tolerance, and sP decreased. Together, the

increase in rN and decrease in sP result in higher persistence levels (Fig. 5 C, D).To

assess the influence of priming on possible treatment success and resistance evolution,

we extended a previously developed modeling framework to predict resistance

evolution (*9*) by a persistent subpopulation (Fig. S 5). We then explored the effects of

priming on tolerance and persistence, individually and in combination, which would be

challenging empirically. Using our predictive framework, we investigated the effect of

priming on survival based on a zero-order pharmacokinetic profile (*9, 26*) (Fig. S 6) and

with parameterized pharmacodynamics functions (Fig. S 7).

Figure 5

A

growth $r_N(A)$ switching s_N growth $r_P(.$

N P

switching s_P

B

$$\frac{dN}{dt} = r_N(A)N(t) - s_N N(t) + s_P P(t)$$

$$\frac{dP}{dt} = r_P(A)P(t) - s_P P(t) + s_N N(t)$$

C

D

Figure 5: The two-state model describes time-kill data. (A) The two-state model (adapted from *17*)
consists of two subpopulations, normal cells (N) and persister cells (P) and is parameterized with the growth
rates $r_N(A)$ and $r_P(A)$, which are depended on the concentration of AMPs (A) in the system, and the
switching rates s_N, and s_P. Each subpopulation is described with an ordinary differential equation (B), each
of which describes the change of the respective subpopulation over time. For each antimicrobial, melittin
(C) and pexiganan (D), we fitted the model to the data of naive (blue) and primed (orange) bacterial
populations (see also Fig. 1) individually. The continuous lines represent the total bacterial population B(t),
with B(t) = N(t) + P(t), and the dashed and dotted lines represent the subpopulations N(t) and P(t),
respectively. Bacteria primed with melittin have an increased net growth rate r_N and decreased s_P compared
to the naive populations. In the case of pexiganan, the parameter r_N is significantly higher in primed
compared to naive populations. For an overview of the fitted parameters, see table S 5.

We found that survival of the population was highly dependent on tolerance (Fig. 6 A,
B, S 8). The presence of persistent cells alone only marginally increased time until
clearance. When we implemented the observed decrease in switching rate s_P, the time
until clearance of the bacterial population is extended at high treatment intensities (high
concentrations A_{max}). The results do not qualitatively change for larger pharmacokinetic
decay rates (k), typical for AMPs (Fig. S 9). Taken together, an increase in tolerance

alone resulted in higher survival independent of persister cells. An increase in persister

cells further increased survival at high antimicrobial concentrations.

Next, we assessed how priming affects resistance emergence. Generally, resistance

evolution depends on the population size, mutation rate and the replication rate. While

priming increases bacterial survival, hence the population size, it also increases the

number of persisters that do not replicate and hence are a poor source of resistant

mutations (Fig 1, Fig. S 10). Our simulations revealed that the beneficial effects of

priming on survival due to increased persistence did not translate into an increased

probability of resistance emergence and establishment (Fig. 6 B, Fig. S 8, 9). The

probability of resistance emergence was mainly influenced by the effect of priming on

tolerance.

23   Figure 6

A	B

Figure 6: Influence of priming on time until clearance and resistance evolution. We extended a

previously developed PKPD framework (*9*) to include persistence (Fig. S 5). With this framework,

we estimated PD curves (Fig. S 7) for primed and naïve bacteria and the estimated switching rates

(Fig. 5) to predict (A) time until clearance and (B) probability of resistance evolution. We simulated

primed bacteria (primed with heterogeneous population consisting of N and P subpopulations,

indicated as "primed, heterogeneous, smaller s_P") and naïve bacteria (naïve, heterogeneous). In

addition, we simulated dynamics without persistence for both primed and naïve bacteria ("primed

homogeneous" and "naïve homogeneous") and dynamics of primed populations with switching

rates of the naïve population ("primed, heterogeneous"). In (A), each dot is an individual run. In

both (A) and (B), the line denotes the average of all runs. No clearance means that simulated

treatment could not reduce bacteria population < 1 cell within 7 days of treatment. Here, we only

show the results of melittin (results of pexiganan are qualitatively highly similar, see Fig. S 8). All

parameter values used in the mathematical model are listed in table S 5.

DISCUSSION

We find that sublethal dosing of the AMPs melittin and pexiganan primes bacterial cells to increase both tolerance and persistence. As sublethal concentrations of antimicrobials (AMPs and antibiotics) are common in the environment and in hosts/patients, a priming response, if it is a widespread phenomenon, would have significant consequences. Antimicrobials that prime inevitably would induce the formation of tolerant and persisting cells. This should apply to situations of induced immune responses, but equally to drug treatments. In short, generating or increasing populations of phenotypic resistant bacteria, would be inevitable in many situations, and would represent a serious obstacle to clearing bacterial infections.

The priming response we report is mediated by rather general physiological resistance mechanisms. The molecular basis of the induction of AMP tolerance and persistence is based on modifications of bacterial envelopes involving either curli production under melittin treatment, or colanic acid production after pexiganan exposure. Both these mechanisms protect the cells by shielding them from AMPs, either making the access to the outer membrane more difficult and/or capturing AMPs, giving scope for cross-resistance. These physiological resistance mechanisms are not antimicrobial specific and hence likely to be also induced by other antimicrobials.

Interestingly, the activation of both pathways shows different dynamics in biofilm formation (*35*). Curli and colanic acid are important components of the biofilm matrix, curli for the primary adhesion and colanic acid for the biofilm structure (*35*). Triggering

their expression by sublethal levels of AMPs, could potentially catalyse biofilm formation. Within a host, if the immune system fails to clear the pathogens, the AMP-priming effect may thereby favour the transition from acute to chronic bacterial infections, where biofilms prevail.

In natural systems of host-microbe interactions, priming plays a role in facilitating colonization of AMP rich environments (*11*). Priming by AMPs also plays a role in infection vectors: in the flea gut the PhoQ-PhoP system is induced in *Yersinia pestis,* the infective agent causing plague, by AMPs leading to biofilm formation that enhances transmission to the final host (*36*). It is not clear as yet if phenotypic AMP-resistance will facilitate opportunistic infections in a way similar to genetic AMP-resistance, as has been shown for genetic AMP resistance in *S. aureus* (*37*), but in the light of our results it seems likely.

Toleranc and persistence can drive the evolution of genetic resistance (*10*)(*18*). We find that, while priming *E. coli* with AMPs results in increased tolerance and persistence, the main driver of resistance evolution in our model is increased tolerance. Resistance evolution is a product of mutation supply and strength of selection. The selection in our scenarios is strong, but mutations are only supplied from the subpopulation of tolerant cells, not from the persister cells, as they are metabolically inactive. So, while the persisters do not contribute to resistance evolution, they do provide a source for secondary infection, once the immune system overexpression ceases or alternatively, when the antimicrobial drug is removed. In antibiotic resistance evolution, by contrast, persisters have shown to contribute to resistance evolution (*38*). Part of this seems to be explained by increased mutagenesis caused by the antibiotics (*38*), an aspect we explicitly did not model as AMPs do not increase mutagenesis (*39*).

Our combined theoretical and empirical results suggest that in hosts the optimal strategy of AMP-deployment would be a very fast increase in concentration to avoid priming and

the subsequent development of phenotypic resistance through both, increased persistence and increased tolerance. Such fast increases of AMPs in specific locations are realized in some important natural situations. During insect metamorphosis, when the gut is renewed in the pupa, a cocktail of AMPs and lysozymes is discharged into the gut (*40*) and references therein), resulting in a quick reduction of bacterial numbers by orders of magnitude. When infections persist, one possible solution is a long-lasting immune response (*41*) that deals with a recurrent infection that could potentially be caused by persisting cells. These natural situations also have parallels in the medical application of antimicrobials. We propose that a fast increase of antimicrobial concentration, as for example in the intra-venous application of antibiotics, should also reduce the probability of persister formation via priming.

Long-lasting antimicrobial exposures are prevalent in natural systems. This is at odds with the observation that long-lasting drug treatments, at least in the case of single drug applications, select for drug resistance. Therefore, at first glance extended treatments do not seem to be a good strategy to manage persisters. In the natural context of AMPs, as they are always expressed as cocktails, this is very likely not an issue and would suggest the use of combination therapies.

Acknowledgments: We are grateful to Sophie Armitage, Charlotte Rafaluk-Mohr and Olivia Judson for comments on the manuscript.

Declaration of interests
The authors declare no conflict of interest

METHODS

Bacteria and growth conditions

The *E. coli* MG1655 was used as bacterial models for all experiments. All cultures related to antimicrobial tests were carried out in Mueller-Hinton I Broth (Sigma). For genetic manipulations (*E. coli* BW25113). All the strains and their derivatives were routinely cultured in Lysogeny Broth (LB medium) or SOB, supplemented with antibiotics when appropriate. Other constructed and strains used in this study are listed in table S 6 of this section.

Minimal inhibitory concentration (MIC)

MICs were determined according to CLSI recommendations by a microdilution method in MHB with some modifications. Inoculum size that was adjusted to 2×10^7 CFU/ml from a regrowth of overnight cultures to be consistent with the downstream experiments. The MIC was defined as the antimicrobial concentration that inhibited growth after 24 h of incubation in liquid MH medium at 37°C. Polypropylene non-binding plates (Th. Geyer, Germany) were used for all experiments. The MIC was considered as the antimicrobial concentration that inhibited growth after 24 hours of incubation in liquid medium at 37°C.

MIC was also determined similarly as above for primed bacteria after two and a half hours of exposure to priming concentrations. The values were compared with naive controls for both antimicrobial peptides, melittin and pexiganan, and for ciprofloxacin.

Priming experiments

Starting from 1×10^8 CFU/ml in mid exponential growth, where bacteria were exposed (stimulus) to 1/10 MIC of melittin or pexiganan during 30 minutes at 37°C with

soft shaking. The tubes were centrifuged at 4000 x g for 10 minutes and allowed to

recover for 60 minutes. The cells were challenged (trigger of priming response) with a

concentration equivalent to an 10X MIC. The cultures were diluted and plated to

determine cell viability. Five biological replicates were generated. Non-treated cells were

used as a control and also harvested during mid exponential growth.

Activity of melittin and pexiganan from the supernatant of challenged cells.

Similarly, to the priming experiments, $1x10^8$ CFU/ml of bacteria were exposed 10x

MIC (supernatant I) of melittin or pexiganan during 150 minutes at 37°C with soft

shaking. The tubes were centrifuged at 4000 x g for 10 minutes, the supernatant were

collected and centrifuged again for 20 000 x g for 30 minutes. The new supernatant was

filtrated using 0.22 µm sterile filters (Sigma Aldrich, Germany) and used immediately

(supernatant II). In parallel, exponentially growing cultures containing $1x10^8$ CFU/ml per

tube were centrifuged at 4000 x g for 10 minutes, the supernatants were removed by

sterile aspiration and the pellets were re-suspended in equal volumes of the supernatant

II and incubated during 150 minutes at 37°C with soft shaking. Samples from each tube

were from both, supernatant I and supernatant II, taken every 30 minutes and diluted and

plated to determine cell viability. Five replicas per culture were used. Data points from

time-kill experiments from bacteria treated with the supernatant I (first round) and the

supernatant II (second round) were compared to determine changes in the activity of the

AMPs either by degradation or adsorption.

Persister antibiotic survival assay

Bacterial cultures were inoculated at 1:100 from a 16-hour overnight culture into

MHB medium. Cells were grown for 2 h to reach approximately 2×10^8 CFU/ml). The

cultures were treated with priming concentrations (1/10 MIC) of melittin and pexiganan

for 30 minutes. Non-treated cultures were used as control. All cultures were washed and

centrifuged twice to remove the treatment. The supernatants were used to determine ATP

concentration using a Molecular Probes ATP Determination Kit (Thermo Fisher

Scientific, Germany). Bacteria were resuspended in equal volume of fresh medium and

an aliquot from each culture was taken to determine the number of bacteria at t=0 by

diluting and plating in MHB agar. Following, ciprofloxacin was added for a final

concentration of 2 µg/ml to treated tubes and to non-treated AMPs control. The cultures

were incubated for four hours. Then, bacteria were washed twice with 0.9 % NaCl and

plated on MHB agar to determine the counts of survival fractions. The percent survival

was calculated as the ratio of cells before and after the treatment as described previously

(*34*). Briefly final CFU/CFU at 0 h) × 100. The results are presented as the average from

5 independent replicates.

Construction and verification of deletion mutants

We inactivated the major curli subunit protein gene *csgA* and the colanic acid

precursor gene *wza*. Although both pathways involve many genes, the removal of these

two components impair the production of both substances respectively. These mutants

were generated in *E. coli* K-12 strain MG1655 following the methodology described

elsewhere (*42*) with some modifications because we used as template the genomic DNA

of the Keio collection (*43*). Briefly, we extracted genomic DNA from the mutants

csgA::Kn and *wza::Kn* of the Keio collection (*E. coli* BW25113) and amplified by PRC

the flanking regions of the kanamycin resistance cassette disrupting both genes and

including an appropriate homology sequence. For the *csgA* mutant we used the primers

5'-GATGCCAGTATTTCGCAAGGTG-3' and 5'-

GGTTATCTGACTGGAAAGTGCC-3' while primers 5'-

TAGCGTGTCTGGATGCCTG-3' and 5'-CCACTTTCAGCTCCGGGT-3' were used

for *wza*. The PCR products were purified and electroporated in the *E. coli* MG1655

carrying a red recombinase helper plasmid, pKD46. The strain was grown in 10 ml SOB

medium with ampicillin (100 µg/ml) and L-arabinose at 30°C to an OD_{600} of ~0.5 and

then made electrocompetent by washing and centrifuge them with a cold solution of

glycerol 10%. Competent cells in 80 µl aliquots were electroporated with 200 ng of

PCR product. Cells were added immediately 0.9 ml of SOC, incubated during 2 h at

37°C, and then 100 µl aliquots spread onto LB agar with kanamycin (30 µg/ml). The

correct inactivation of genes was verified by PCR. The antibiotic resistant cassette (Kn)

was removed for both mutants using the flippase plasmid pCP20.

Transcriptome sequencing

The transcriptome sequencing of primed cells was determined on samples treated

identically as described for the priming experiments. Total RNA from 10^8 cell per sample

was isolated using the RNAeasy Isolation kit (Qiagen, Germany). Traces of genomic

DNA were removed from 10 µg of RNA by digestion in a total volume of 500 µl

containing 20 units of TURBO DNase, incubated for 30 minutes at 37°C, immediately

followed by RNeasy (Qiagen) clean-up and elution in 30 µl of RNase-free water.

Following DNase treatment, RNA integrity was assessed using Agilent RNA 6000 Nano

kit and 2100 Bioanalyzer instrument (both from Agilent Technologies). Total RNA was

depleted from ribosomal RNA using the Ribo-Zero Depletion Kit for Gram-negative

bacteria (Illumina, USA). Libraries were prepared using a TruSeq Stranded Total RNA

library preparation kit (Illumina, USA) and were sequenced on a MiSeq platform.

Transcript abundances were derived from pseudo-alignment of reads to the cDNA

sequences from the ASM584v2 assembly of *Escherichia coli* MG1655 (ENA accession

GCA_000005845.2) using Salmon version 0.7.2 with default parameters (*44*).

Differential gene expression was analyzed using the R package DESeq2 (*45*) in

conjunction with tximport (*46*). Pairwise contrasts were performed between the control

and each AMP treatment with empirical bayesian shrinkage of both dispersion parameters

and fold-change estimation. We defined genes as being significantly differentially

expressed when the absolute fold-change in expression was greater than 2, at an FDR-

adjusted p-value of less than 0.05. The variance-stabilizing transformation was used to

remove the dependence of the variance on the mean and to transform data to the log2

scale prior to ordination using principal component analysis. Quality of RNAseq data

were contrasted by Euclidian distance and symmetry of data reads distribution (Fig. S

11).

Observation at single cell level

To observe cell reaction at single cell level during priming experiments, we used an

ad hoc microfluidic device developed for this project. It consisted in a main channel for

bacterial inoculation and medium perfusion and with several lateral compartments which

dimensions are around 1.5 μm height (ensuring all bacteria are kept in focus) and square

200 μm width corresponding to a field size of the used microscope at 1000x magnification

(Fig. S 12). The chip was designed in Autocad. We started the replication of our

microfluidic chips from a custom made (Sigatec SA) silicon (SiO) master. This silicon

master itself was first replicated in Smooth-Cast 310 (Bentley advanced material). Soft

lithography produced the chips in PDMS (Sylgard Silicone Elastomer Base and Curing

Agent mixed in 10:1 ratio). The PDMS chips were cured overnight at 75°C in an

incubator. We punched an inlet and outlet hole for the laminar flow in each chip using a

biopsy puncher of 0.5 mm (outer diameter). The chips were bonded to a glass cover slide

(24×60 mm) after a 30-second air plasma treatment (PDC-002, Harrick Plasma). Before

use, the assembled chip was treated for 15 seconds in air plasma and immediately injected

it with filtered MHB medium for passivation. We left the activated chip to incubate for a

least 1 hour before loading the bacteria. The devices were loaded to full capacity with a

bacterial suspension containing nearly 2×10^8 CFU/ml (exponentially growing bacteria,

0.5 OD_{600}). Cell suspension was injected into the main channel of the chip using a blunt-

end 23G needle attached to a 1ml syringe. We centrifuged the loaded chip at 200 x g for

10 min using in-house adapters, checked the loading.

After we loaded the bacteria cells in the blind ending side channels, we connected

the chip to a syringe pump (AL-6000, WPI, Germany) and placed the chip under an

inverted microscope. A continuous laminar flow (100µl/h) of MHB through the central

channel was maintained throughout the experiment (Fig. S 12). For the life cell imaging,

after infusion with priming or triggering concentration of AMPs, we injected MHB

supplemented with bacterial Live/Dead stain kit solutions (Thermo Scientific, Germany)

for a final concentration of 0.1 µl/ml of MHB. We took pictures of at least 20 fields per

treatment from independent side channels. Fluorescent images were taken of each field

of view with simultaneous acquisition in red and green fluorescent channels during a time

interval of no more than 2 minutes per treatment with a Nikon Ti-2 inverted microscope

(Nikon, Japan). Cells were observed with the 100× objective and controlled by Nis

Element AR software. The chip holder is temperature controlled at 37°C.

Determination of melittin-induced curli

The production of curli due to melittin treatment was determined by using the fluorescent

dye Ectracer[TM680] (Ebba Biotech, Sweden) that stain extracellular curli. Ectracer[TM680]

was used according to the instructions of manufacturers. Bacterial cultures were treated

on chip with priming and triggering concentrations of melittin as described for single cell

observation section omitting the live/dead staining. After priming and triggering, the

channels were perfused MHB supplemented with Ectracer^TM680 in a proportion of 1/1000

related to the medium. Cells were observed with the red channel fluorescence for Cy5

dye using a Nikon Eclipse Ti2 inverted optical microscope using the 100X oil objective.

Two independent samples were prepared for each group (primed and naïve cells).

SEM of *E. coli* treated antimicrobial peptides

Approximately 2×10^7 CFU/ml *E. coli* MG1655 were treated with 1/10 MIC of pexiganan

and melittin during 30 minutes. The cultures were concentrated 10 times by a quick

centrifugation step of 1 minute at 8 000 g and resuspended in 1/10 of its own supernatant.

and resuspension 10 μl drops were placed on a circular glass cover slip (1.5 cm of

diameter). The drops were fixed with osmium tetroxide vapors during one minute and

allow to dry in a laminar flow cabinet. The cover glasses were mounted on aluminum

stubs using double-sided adhesive tape and coated with gold in a sputter coater (SCD-

040; Balzers, Union, Liechtenstein). The specimens were examined with a FEI Quanta

200 SEM (FEI Co., Hillsboro, OR) operating at an accelerating voltage of 15 kV under

high vacuum mode at different magnifications. At least 5 fields from two independent

replicas.

Statistical analysis

To analyze the priming data, we first tested if the dynamics of the depicted in the time-

kill curves are biphasic.

We fitted the function

$$f(m_1, m_2, t_{kink}, t) = \begin{cases} log_{10}\big(CFU(t=0)\big) + m_1 t, & t < t_{kink} \\ log_{10}\big(CFU(t=0)\big) + m_1 t_{kink} + m_2(t - t_{kink}), & t \geq t_{kink} \end{cases}$$

to the time-kill data of each AMP and for primed and naïve populations individually using a least square algorithm. Here, t_{kink} is the time point, at which the population dynamics switch from the first phase to the second phase and m_1 and m_2 are the slopes of the first and the second decline, respectively. Note that m_1 is a direct measure of tolerance. The standard error (SE) was calculated as

$$SE(\theta) = \sqrt{\frac{1}{I(\theta)}}$$

The parameter θ denotes to the estimated parameter values of m_1, m_2, and t_{kink} and $I(\theta)$ is the expected Fisher information. The 95% confidence interval was calculated as $\theta \pm 1.96 * \mathrm{SE}(\theta)$.

We used an ANOVA by means of contrasts to assess if priming changes tolerance and persistence. For this, we tested if the decrease of the first phase (all data points with $t < t_{kink}$) differed between primed and naive populations (test for differences in tolerance) and if the population size in the second phase (all data points with $t > t_{kink}$) differed between primed and naive populations (test for differences in persistence). Both tests showed significant differences between primed and naive populations for both melittin and pexiganan (significance level: $p < 0.05$). We corrected for multiple testing with the Bonferroni-method.

Population models

To describe bacterial population dynamics, we used the two-state model by Balaban *et al.* (*25*) (Fig. 5 A, B):

$$\frac{dN(t)}{dt} = (r_N(A) - s_N)N(t) + s_P P(t)$$

$$\frac{dP(t)}{dt} = (r_P(A) - s_P)P(t) + s_N N(t)$$

In this model, the bacterial population consists of two subpopulations, one with a

normal phenotype, *N(t)*, and a second with a persister phenotype,*P(t)*. We denote the total

bacterial population size by *B(t)*,with *B(t)* = *N(t)+P(t)*. The rate of change of the

population is determined by the net growth rate of N and P, r_N and r_P, and the switching

rate from N to P, s_N, and the switching rate from P to N, s_P. The parameter estimates of

the net growth rates are dependent on the AMP concentration A, i.e. *$r_N(A)$* and *$r_P(A)$*.

analytical solution of this ODE system (*25, 47*) is

$B(t) = N(t) + P(t) = c_1 u_1 e^{\lambda_1 t} + c_1 u_2 e^{\lambda_1 t} + c_2 v_1 e^{\lambda_2 t} + c_2 v_2 e^{\lambda_2 t}$ Eq. S2

with

λ_1

$$= \frac{r_N + r_P - s_N - s_P - \sqrt{(-r_N - r_P + s_N + s_P)^2 - 4(r_N r_P - r_P s_N - r_N s_P)}}{2}$$

$$\lambda_2 = \frac{r_N + r_P - s_N - s_P + \sqrt{(-r_N - r_P + s_N + s_P)^2 - 4(r_N r_P - r_P s_N - r_N s_P)}}{2}$$

$$\vec{u} = \begin{pmatrix} \dfrac{\lambda_1 - r_P + s_P}{s_N} \\ 1 \end{pmatrix}$$

$$\vec{v} = \begin{pmatrix} \dfrac{\lambda_2 - r_P + s_P}{s_N} \\ 1 \end{pmatrix}$$

$$c_1 = \frac{v_1 P(t = 0) - N(t = 0)}{v_1 - u_1}$$

1
$$c_2 = \frac{-u_1 P(t=0) + N(t=0)}{v_1 - u_1}$$

The model was fitted by a least square algorithm, minimizing the residual sum of

squares of the data to the prediction. For the starting conditions ($N(t=0)$, $P(t=0)$), we

assumed that the ratio of N/P was at the equilibrium predicted by the model without

antimicrobials when the exposure to lethal concentrations of AMPs started. $N(t=0)$ and

$P(t=0)$ were therefore calculated using the eigenvector $\vec{v}$ that corresponds to the largest

eigenvalue of a system without antimicrobials. Here, we assumed that the parameter r_N is

equal the net growth rate in absence of antimicrobials, $r_N = \psi_{max}$. The parameter ψ_{max}

was estimated based on the time-kill curve of bacterial population that grow in absence

of antimicrobials (see below and Fig. S 11). The eigenvector contains information about

the ratio of N and P for $t \rightarrow \infty$: $\frac{N}{P} = \frac{v_1}{v_2}$. Resulting, $P(t=0) = \frac{B(t=0)}{1+v_1}$ and $N(t=0) =$

$B(t=0) - P(t=0)$. $B(t=0)$ was estimated from the data. Confidence intervals were

calculated as described above. In a pre-analysis, we used 4 free model parameters that

were fitted: r_N, r_P, s_N and s_P. The parameter r_P was not significant from 0 (Fig. S 8).

Therefore, we set the parameter r_P to 0 and fitted the remaining 3 parameters to the data

(Fig. S 9, table S 5).

Tolerance and persistence in terms of model parameters

The measure of tolerance is the slope m_1. Komarova and Wodarz (*47*) showed that

the slope can directly be linked to the population model parameters. In our notation,

$m_1 = log_{10}\big(c_1(u_1+v_1)e^{\lambda_1 t}\big)$ Eq. S1

Note that the first phase is mainly influenced by r_N (Fig. S9), therefore, $m_1 \approx r_N$.

Persistent cell numbers at time t were calculated with the analytic solution:

$P(t) = c_1 u_2 e^{\lambda_1 t} + c_2 v_2 e^{\lambda_2 t}$ Eq. S2

Pharmacokinetic and pharmacodynamic function

We used the pharmacokinetic function $A(t) = A_{max}e^{-k(t-n)}$, with $8h * n \leq t \leq$ $8h * (n + 1)$ and $n = 0,1,2 ...$ described previously elsewhere (*26*). In our simulations, we fixed the decay parameter k and varied the drug input A_{max}. To describe the effect of the AMPs on the bacterial population, we used the pharmacodynamic (PD) function $\psi(A)$ (*26, 48*), with

$$\psi(A) = \psi_{max} - e(A) = \psi_{max} - \frac{(\psi_{max} - \psi_{min})\left(\frac{A}{MIC}\right)^{\kappa}}{\left(\frac{A}{MIC}\right)^{\kappa} - \frac{\psi_{min}}{\psi_{max}}}$$

The parameter ψ_{max} describes the net growth rate in absence of antimicrobials ($\psi_{max} = \psi(A = 0)$). The antimicrobial effect $e(A)$ is dependent on the antimicrobial concentration and is the defined with ψ_{max}, ψ_{min}, the net growth rate in presence of large amounts of antimicrobials ($\psi(A \to \infty)$), with the MIC, the antimicrobial concentration that results in no growth ($\psi(A = MIC) = 0$) and with κ, which determines the steepness of the PD curve.

The PD function was fitted to the time-kill curves (Fig. S6), as described by Regoes *et al.* (*26*). In short, we used log-linear regressions of the time kill curves within the time-points 0h and 1h to estimate the change of the bacterial population over time, i.e. the slopes of the log-linear regression. We fixed the parameter ψ_{max} and fitted the 3 remaining parameters of the PD function with the *Markov-Chain-Monte-Carlo method.*

Stochastic simulations

To simulate resistance evolution with stochastic simulations, we expanded a previously developed framework for bacterial population dynamics (*9*). The framework models bacterial population dynamics exposed to changing levels of antimicrobials and

allows for resistance evolution. In the simulations, the change in population size of a sensitive strain S, with $S = N+P$, and of a resistant strain R were described with the following ODE system:

$$\frac{dN(t)}{dt} = \psi_{max}(1 - \mu)N(t)\left(1 - \frac{N(t) + P(t) + R(t)}{K}\right) - s_N N(t) + s_P P(t)$$

$$- e_N(A(t))N(t)$$

$$\frac{dP(t)}{dt} = s_N N(t) - s_P P(t)$$

$$\frac{dR(t)}{dt} = \psi_{max} \mu N(t)\left(1 - \frac{N(t) + P(t) + R(t)}{K}\right) + \psi_{max}(1 - c)R\left(1\right.$$

$$\left. - \frac{N(t) + P(t) + R(t)}{K}\right) - e_R(A(t))R(t)$$

Here, the replication rate is assumed to be equal to the maximum net growth rate ψ_{max}. The effect of the antimicrobial $e_N(A)$ is explained above. Note that we assume that bacteria in class P do not grow and are not affected by antimicrobials. We also assumed that the switching rates are constant. To describe the effect of an antimicrobial on the strain R, $e_R(A)$, we use the same parameter set than with $e_N(A)$, except for the *MIC*: $\frac{MIC_R}{MIC_N} = 10$. In the simulations, we differentiated between the following cases: (i) naïve bacteria, homogeneous population (no persistence), (ii) naïve bacteria, heterogeneous population, (iii) primed bacteria (increase in r_N), homogeneous population, (iv) primed bacteria (increase in r_N), heterogeneous population (v) primed bacteria (increase in r_N and decrease in s_P). The stochastic simulations were run 1000 times for each antimicrobial, for each case, and for a variety of input antimicrobial concentrations A_{max}. All parameter values are listed in table S 6. The intensity is the input antimicrobial concentrations A_{max}.

The simulations were run for t=7d. Time until clearance and probability of resistance evolution were calculated as the mean of the value over the 1000 simulations.

Implementation

Statistical testing, simulations and plots were done in R version 3.3.2 (*49*), using Rstudio version 1.0.143 (*50*). We used the following R-packages: (i) for plotting: sfsmisc (*51*), (ii) for fitting the PD function: rjags(iii) for stochastic simulations: adaptivetau (*52*). We used Mathematica version 11.0 (*53*) to determine the analytical solutions of the population models.

REFERENCES

1.   B. P. Lazzaro, M. Zasloff, J. Rolff, Antimicrobial peptides: Application informed by evolution. *Science* **368**, eaau5480 (2020).

2.   C. L. Bevins, N. H. Salzman, Paneth cells, antimicrobial peptides and maintenance of intestinal homeostasis. *Nat. Rev. Microbiol.* **9**, 356–368 (2011).

3.   P. Mergaert, Role of antimicrobial peptides in controlling symbiotic bacterial populations. *Nat. Prod. Rep.* **35**, 336–356 (2018).

4.   N. Mookherjee, M. A. Anderson, H. P. Haagsman, D. J. Davidson, Antimicrobial host defence peptides: functions and clinical potential. *Nat. Rev. Drug Discov.* **19**, 311–332 (2020).

5.   H. B. Koo, J. Seo, Antimicrobial peptides under clinical investigation. *Pept. Sci.* **111** (2019), doi:10.1002/pep2.24122.

6.   H.-S. Joo, C.-I. Fu, M. Otto, Bacterial strategies of resistance to antimicrobial peptides. *Philos. Trans. R. Soc. B Biol. Sci.* **371**, 20150292 (2016).

7.   G. G. Perron, M. Zasloff, G. Bell, Experimental evolution of resistance to an antimicrobial peptide. *Proc. R. Soc. B Biol. Sci.* **273**, 251–256 (2006).

8.   R. Spohn, L. Daruka, V. Lázár, A. Martins, F. Vidovics, G. Grézal, O. Méhi, B. Kintses, M. Számel, P. K. Jangir, B. Csörgő, Á. Györkei, Z. Bódi, A. Faragó, L. Bodai, I. Földesi, D. Kata, G. Maróti, B. Pap, R. Wirth, B. Papp, C. Pál, Integrated evolutionary analysis reveals antimicrobial peptides with limited resistance. *Nat. Commun.* **10**, 4538 (2019).

9.   G. Yu, D. Baeder, R. Regoes, J. Rolff, Predicting Drug Resistance Evolution:
Antimicrobial Peptides Vs. Antibiotics. *Proc R Soc Lond B*, 138107 (2018).

10. B. R. Levin, D. E. Rozen, Non-inherited antibiotic resistance. *Nat. Rev. Microbiol.*
4, 556–562 (2006).

11. F. Chen, B. C. Krasity, S. M. Peyer, S. Koehler, E. G. Ruby, X. Zhang, M. J. McFall-
Ngai, Bactericidal Permeability-Increasing Proteins Shape Host-Microbe
Interactions. *mBio*. **8**, e00040-17, /mbio/8/2/e00040-17.atom (2017).

12. A. Mitchell, G. H. Romano, B. Groisman, A. Yona, E. Dekel, M. Kupiec, O. Dahan,
Y. Pilpel, Adaptive prediction of environmental changes by microorganisms.
Nature. **460**, 220–224 (2009).

13. A. J. P. Brown, N. A. R. Gow, A. Warris, G. D. Brown, Memory in Fungal
Pathogens Promotes Immune Evasion, Colonisation, and Infection. *Trends*
Microbiol. **27**, 219–230 (2019).

14. D. R. Andrade-Linares, A. Lehmann, M. C. Rillig, Microbial stress priming: a meta-
analysis: Microbial priming to stress. *Environ. Microbiol.* **18**, 1277–1288 (2016).

15. O. Makarova, A. Rodriguez-Rojas, M. Eravci, C. Weise, A. Dobson, P. Johnston, J.
Rolff, Antimicrobial defence and persistent infection in insects revisited. *Philos.*
Trans. R. Soc. B Biol. Sci. **371**, 20150296 (2016).

16. D. I. Andersson, D. Hughes, Microbiological effects of sublethal levels of
antibiotics. *Nat. Rev. Microbiol.* **12**, 465–478 (2014).

17. N. Q. Balaban, S. Helaine, K. Lewis, M. Ackermann, B. Aldridge, D. I. Andersson,
M. P. Brynildsen, D. Bumann, A. Camilli, J. J. Collins, C. Dehio, S. Fortune, J.-M.
Ghigo, W.-D. Hardt, A. Harms, M. Heinemann, D. T. Hung, U. Jenal, B. R. Levin,
J. Michiels, G. Storz, M.-W. Tan, T. Tenson, L. Van Melderen, A. Zinkernagel,
Definitions and guidelines for research on antibiotic persistence. *Nat. Rev.*
Microbiol. **17**, 441–448 (2019).

18. I. Levin-Reisman, I. Ronin, O. Gefen, I. Braniss, N. Shoresh, N. Q. Balaban,
Antibiotic tolerance facilitates the evolution of resistance. *Science*. **355**, 826–830
(2017).

19. G. E. Fantner, R. J. Barbero, D. S. Gray, A. M. Belcher, Kinetics of antimicrobial
peptide activity measured on individual bacterial cells using high-speed atomic force
microscopy, 1–6 (2010).

20. Z. Yao, D. Kahne, R. Kishony, Distinct Single-Cell Morphological Dynamics under
Beta-Lactam Antibiotics. *Mol. Cell*. **48**, 705–712 (2012).

21. H. Raghuraman, A. Chattopadhyay, Melittin: a Membrane-active Peptide with
Diverse Functions. *Biosci. Rep.* **27**, 189–223 (2007).

22. Y. Ge, D. L. Macdonald, K. J. Holroyd, C. Thornsberry, H. Wexler, M. Zasloff, In
Vitro Antibacterial Properties of Pexiganan, an Analog of Magainin. *ANTIMICROB*
AGENTS CHEMOTHER. **43**, 7 (1999).

23. L. M. Gottler, A. Ramamoorthy, Structure, membrane orientation, mechanism, and function of pexiganan — A highly potent antimicrobial peptide designed from magainin. *Biochim. Biophys. Acta BBA - Biomembr.* **1788**, 1680–1686 (2009).

24. M. Hilker, J. Schwachtje, M. Baier, S. Balazadeh, I. Bäurle, S. Geiselhardt, D. K. Hincha, R. Kunze, B. Mueller-Roeber, M. C. Rillig, J. Rolff, T. Romeis, T. Schmülling, A. Steppuhn, J. van Dongen, S. J. Whitcomb, S. Wurst, E. Zuther, J. Kopka, Priming and memory of stress responses in organisms lacking a nervous system: Priming and memory of stress responses. *Biol. Rev.* **91**, 1118–1133 (2016).

25. N. Q. Balaban, J. Merrin, R. Chait, L. Kowalik, S. Leibler, Bacterial Persistence as a Phenotypic Switch. *Science*. **305**, 1622–1625 (2004).

26. R. R. Regoes, C. Wiuff, R. M. Zappala, K. N. Garner, F. Baquero, B. R. Levin, Pharmacodynamic Functions: a Multiparameter Approach to the Design of Antibiotic Treatment Regimens. *Antimicrob. Agents Chemother.* **48**, 3670 (2004).

27. Hedges, A, An examination of single-hit and multi-hit hypotheses in relation to the possible kinetics of colicin adsorption. *J. Theor. Biol.* **11**, 383–410.

28. P. Abel zur Wiesch, S. Abel, S. Gkotzis, P. Ocampo, J. Engelstädter, T. Hinkley, C. Magnus, M. K. Waldor, K. Udekwu, T. Cohen, Classic reaction kinetics can explain complex patterns of antibiotic action. *Sci. Transl. Med.* **7**, 287ra73-287ra73 (2015).

29. H. Miajlovic, S. G. Smith, Bacterial self-defence: how Escherichia coli evades serum killing. *FEMS Microbiol. Lett.* **354**, 1–9 (2014).

30. M. M. Barnhart, M. R. Chapman, Curli biogenesis and function. *Annu. Rev. Microbiol.* **60**, 131–47 (2006).

31. S. Maria-Neto, K. C. de Almeida, M. L. R. Macedo, O. L. Franco, Understanding bacterial resistance to antimicrobial peptides: From the surface to deep inside. *Biochim. Biophys. Acta BBA - Biomembr.* **1848**, 3078–3088 (2015).

32. Y. Shan, A. Brown Gandt, S. E. Rowe, J. P. Deisinger, B. P. Conlon, K. Lewis, ATP-Dependent Persister Formation in Escherichia coli. *mBio*. **8**, e02267-16 (2017).

33. K. A. Brogden, Antimicrobial peptides: pore formers or metabolic inhibitors in bacteria? *Nat. Rev. Microbiol.* **3**, 238–50 (2005).

34. Y. Shan, A. Brown Gandt, S. E. Rowe, J. P. Deisinger, B. P. Conlon, K. Lewis, ATP-Dependent Persister Formation in *Escherichia coli*. *mBio*. **8**, e02267-16, /mbio/8/1/e02267-16.atom (2017).

35. C. Prigent-Combaret, G. Prensier, T. T. Le Thi, O. Vidal, P. Lejeune, C. Dorel, Developmental pathway for biofilm formation in curli-producing Escherichia coli strains: Role of flagella, curli and colanic acid. *Environ. Microbiol.* **2**, 450–464 (2000).

36. V. Vadyvaloo, A. K. Viall, C. O. Jarrett, A. K. Hinz, D. E. Sturdevant, B. J. Hinnebusch, Role of the Phop-PhoQ gene regulatory system in adaptation of

yersinia pestis to environmental stress in the flea digestive tract. *Microbiol. U. K.* **161**, 1198–1210 (2015).

37. G. Cheung, E. Fisher, J. McCausland, J. Choi, J. Collins, S. Dickey, M. Otto, Antimicrobial peptide resistance mechanism contributes to Staphylococcus aureus infection Accepted. *J. Of*, 1–34 (2018).

38. E. M. Windels, J. E. Michiels, M. Fauvart, T. Wenseleers, B. Van den Bergh, J. Michiels, Bacterial persistence promotes the evolution of antibiotic resistance by increasing survival and mutation rates. *ISME J.* **13**, 1239–1251 (2019).

39. A. Rodríguez-Rojas, O. Makarova, J. Rolff, Antimicrobials, Stress and Mutagenesis. *PLoS Pathog.* **10**, e1004445 (2014).

40. P. R. Johnston, V. Paris, J. Rolff, Immune gene regulation in the gut during metamorphosis in a holo- versus a hemimetabolous insect. *Philos. Trans. R. Soc. B Biol. Sci.* **374**, 20190073 (2019).

41. E. R. Haine, Y. Moret, M. T. Siva-Jothy, J. Rolff, Antimicrobial Defense and Persistent Infection in Insects. *Science*. **322**, 1257–1259 (2008).

42. K. A. Datsenko, B. L. Wanner, One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U A*. **97**, 6640–6645 (2000).

43. T. Baba, T. Ara, M. Hasegawa, Y. Takai, Y. Okumura, M. Baba, K. A. Datsenko, M. Tomita, B. L. Wanner, H. Mori, *Mol Syst Biol*, in press, doi:msb4100050 [pii] 10.1038/msb4100050.

44. R. Patro, G. Duggal, M. I. Love, R. A. Irizarry, C. Kingsford, Salmon provides fast and bias-aware quantification of transcript expression. *Nat. Methods*. **14**, 417–419 (2017).

45. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

46. C. Soneson, M. I. Love, M. D. Robinson, Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. *F1000Research*. **4**, 1521 (2015).

47. N. L. Komarova, D. Wodarz, Effect of Cellular Quiescence on the Success of Targeted CML Therapy. *PLoS ONE*. **2**, e990 (2007).

48. J. Zhi, C. H. Nightingale, R. Quintiliani, A pharmacodynamic model for the activity of antibiotics against microorganisms under nonsaturable conditions. *J. Pharm. Sci.* **75**, 1063–7 (1986).

49. R Core Team, *R: A language and environment for statistical computing.* (R Foundation for Statistical Computing, Vienna, Austria, 2015).

50. R Development Core Team, RStudio Team (2015). RStudio: Integrated Development for R. (2015).

1   51. M. Maechler, Utilities from "Seminar fuer Statistik" ETH Zurich [R package
2   sfsmisc version 1.1-4].

3   52. Y. Cao, D. T. Gillespie, L. R. Petzold, Adaptive explicit-implicit tau-leaping method
4   with automatic tau selection. *J. Chem. Phys.* **126**, 224101 (2007).

5   53. Inc. Wolfram Research, Mathematica, Version 11.0 (2016).

Supplement

Supplementary tables S1-S6

Supplementary figures S1-S12

                Supplementary tables

Table S 1. Minimal inhibitory concentration of the two antimicrobial peptides melittin and pexiganan and the antibiotic ciprofloxacin used in this work. The MIC was determined to be used as a concentration reference but it was also determined after priming treatment to show that the observed enhanced survival is based on a phenotypic changes and not mutations.

Antimicrobials	MIC (µg/ml) pre-priming	MIC (µg/ml) post-priming
melittin	2	2
pexiganan	1	1
ciprofloxacin	0.125	0.125

Table S 2. Slopes of fitting the biphasic function with the slopes m_1 and m_2 to the data on a $\log_{10}$ scale. Significant differences between slopes of primed and naïve population counts are indicated with asterisks for each antimicrobial.

Antimicrobial	Treatment	Slope			
		m_1 (± SE)		m_2 (± SE)	
melittin	Primed	-3.9 (± 0.6)	*	-0.3 (± 0.4)	
	Naïve	-6.3 (± 0.6)		-1.0 (± 0.4)	
pexiganan	Primed	-2.9 (±0.3)	*	-0.6 (± 0.894)	
	Naïve	-4.1 (± 0.3)		-0.3 (± 0.9)	

Table S 3. Priming response to pexiganan (a) and melittin (b) (0.1xMIC, 30-minute treatment)

See excel files.

Table S 4. Results of fitting the two-state model to bacterial population dynamics in the presence of melittin and pexiganan and the classic exponential population growth model to dynamics in the presence of ciprofloxacin and ampicillin (rounded values).

Antimicrobial	Treatment	r_N (± SE)	$r_P{}^1$	s_N (± SE)	s_P (± SE)
melittin	Primed	-8.96 (0.72)	0	0.00085 (0.00098)	0.27 (0.61)
	Naïve	-15.06 (0.86)	0	0.00128 (0.00095)	2.25 (0.45)
pexiganan	Primed	-6.68 (0.32)	0	0.00027 (0.00059)	0.40 (1.10)
	Naïve	-9.61 (0.40)	0	0.00021 (0.00025)	1.26 (0.66)

[1] Parameter r_P is fixed at 0 due to results of the preliminary analysis, in which the parameter was not significantly different from 0 (see Figure S 8).

Table S 5. Parameter values used as input for the stochastic simulations.

Parameter	Description	AMP	Value
k	AMP degradation rate		0.3
μ	Mutation rate		$1*10^{-9}$
K	Carrying capacity		$5*10^9$
ψ_{max}	Maximal net growth rate in absence of antimicrobials	melittin	2.13
		pexiganan	2.06
ψ_{min}	Minimal net growth rate in presence of large amounts of antimicrobials	melittin	-8.86/ -9.89[§]
		pexiganan	-36.87/-66.80[§]
κ	Hill parameter	melittin	1.3/1.1[§]
		pexiganan	0.6/0.7[§]
MIC	Dose of antimicrobials for which the net growth rate is 0	melittin	0.48/1.83[§]
		pexiganan	0.54/2.14[§]
c	Cost of resistance		0.5
$B(t=0)$	Population size at time t=0		10^8
t_{max}	Maximal run of simulation		7d
s_N	Switching rate from N to P	melittin	0.0012
		pexiganan	0.00021
s_P	Switching rate from P to N	melittin	2.26/0.28[§]
		pexiganan	1.26/0.40[§]

[§] Naïve/Primed

Table S 6. Strains and plasmids used in this work and their relevant phenotypes

Strain	Relevant phenotype	Reference
E. coli MG1655 WT	F⁻ λ⁻ *ilvG⁻ rfb-50 rph-1*	Lab stock
E. coli MG1655 *csgA*::Kn	*csgA*::Kn F⁻ λ⁻ *ilvG⁻ rfb-50 rph-1*	this work
E. coli MG1655 *csgA*::scar	*csgA*::scar* F⁻ λ⁻ *ilvG⁻ rfb-50 rph-1*	this work
E. coli MG1655 *wza*::Kn	*wza*::Kn F⁻ λ⁻ *ilvG⁻ rfb-50 rph-1*	this work
E. coli MG1655 *wza*::scar*	*wza*::scar* F⁻ λ⁻ *ilvG⁻ rfb-50 rph-1*	this work
E. coli BW25113 *csgA*::Kn	*csgA*::Kn *lacI⁺rrnB*$_{T14}$ Δ*lacZ*$_{WJ16}$ *hsdR*514 Δ*araBAD*$_{AH33}$ Δ*rhaBAD*$_{LD78}$ *rph-1* Δ*(araB–D)567* Δ*(rhaD–B)568* Δ*lacZ4787(::rrnB-3) hsdR514 rph-1*	Baba et al., 2006
E. coli BW25113 *wza*::Kn	*wza*::Kn *lacI⁺rrnB*$_{T14}$ Δ*lacZ*$_{WJ16}$ *hsdR*514 Δ*araBAD*$_{AH33}$ Δ*rhaBAD*$_{LD78}$ *rph-1* Δ*(araB–D)567* Δ*(rhaD–B)568* Δ*lacZ4787(::rrnB-3) hsdR514 rph-1*	Baba et al., 2006
Plasmids	**Relevant phenotype**	**Reference**
pKD46	oriR101 w/repA101ts Amp^R	Datsenko and Wanner, 2000
pCP20	oriR101 w/repA101ts Amp^R	Datsenko and Wanner, 2000

* asterisk indicates the removal of the antibiotic resistant cassette kanamycin (Kn) by the action of the flippase expressed by the plasmid pCP20 (Datsenko and Wanner, 2000).

Supplementary References

1. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware quantification of transcript expression. Nat Methods [Internet]. Nature Publishing Group; 2017 [cited 2019 Jul 3];14:417–9.

2. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol [Internet]. 2014 [cited 2019 Jul 3];15:550.

3. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. F1000Research [Internet]. 2015 [cited 2019 Jul 3];4:1521.

4. Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S. Bacterial Persistence as a Phenotypic Switch. Science (80-) [Internet]. 2004 [cited 2018 Jun 9];305:1622–5.

5. Komarova NL, Wodarz D. Effect of Cellular Quiescence on the Success of Targeted CML Therapy. Agur Z, editor. PLoS One [Internet]. 2007 [cited 2018 Jun 9];2:e990.

6. Regoes RR, Wiuff C, Zappala RM, Garner KN, Baquero F, Levin BR. Pharmacodynamic Functions: a Multiparameter Approach to the Design of Antibiotic Treatment Regimens. Antimicrob Agents Chemother [Internet]. American Society for Microbiology (ASM); 2004 [cited 2019 May 30];48:3670. Available from: http://www.ncbi.nlm.nih.gov/pubmed/15388418

7. Yu G, Baeder D, Regoes R, Rolff J. Predicting Drug Resistance Evolution: Antimicrobial Peptides Vs. Antibiotics. Proc R Soc Lond B. 2018;138107.

8. R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2015.

9. R Development Core Team. RStudio Team (2015). RStudio: Integrated Development for R. [Internet]. Boston: MA URL http://www.rstudio.com/; 2015. Available from: http://www.rstudio.com

10. Maechler M. Utilities from "Seminar fuer Statistik" ETH Zurich [R package sfsmisc version 1.1-4]. Comprehensive R Archive Network (CRAN); [cited 2019 Jul 3]; Available from: https://cran.r-project.org/web/packages/sfsmisc/index.html

11. Cao Y, Gillespie DT, Petzold LR. Adaptive explicit-implicit tau-leaping method with automatic tau selection. J Chem Phys [Internet]. 2007 [cited 2019 Jul 3];126:224101. Available from: http://aip.scitation.org/doi/10.1063/1.2745299

12. Wolfram Research I. Mathematica, Version 11.0. Champaign, Illinois: Wolfram Research, Inc.; 2016.

13. Wiuff C, Zappala RM, Regoes RR, Garner KN, Baquero F, Levin BR. Phenotypic tolerance: antibiotic enrichment of noninherited resistance in bacterial populations. Antimicrob Agents Chemother [Internet]. 2005 [cited 2016 Jan 6];49:1483–94. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1068602&tool=pmcentrez&rendertype=abstract

14. Nielsen EI, Friberg LE. Pharmacokinetic-Pharmacodynamic Modeling of Antibacterial Drugs. Pharmacol Rev [Internet]. 2013 [cited 2019 Jul 4];65:1053–90. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23803529

Supplementary Figures

Figure S 1. Bacterial tolerance and persistence change the shape of time-kill curves (figure adapted

from Brauner et al. (*44*). (A) Tolerance is the ability of bacteria to longer survive exposure to

antimicrobials due to decrease in susceptibility. Tolerance is quantified as increase in the slope of

the time kill curve. (B) Persistence is the phenomenon of a subpopulation being less susceptible to

the antimicrobial than the rest of the population. A persistent subpopulation manifests as a biphasic

decline of bacterial population when exposed to lethal concentrations of antimicrobials. Here, the

population consists predominantly of the less susceptible persistent subpopulation. (C) Together,

tolerance and persistence result in biphasic time-kill curves with decreased susceptibility in the

first phase.

Figure S 2. A possible explanation for a nonlinear decline in populations would be a decrease in active AMP concentration over time due to degradation (*45*, *46*), We tested this alternative explanation. As in Figure 1 (main text), we measured the population dynamics of bacteria exposed to 10 x MIC (round 1). At the end of the experiment, we sampled the supernatant. In round 2, a fresh bacterial population was exposed to the sampled supernatant. (A) and (B) show survival data of both rounds for melittin and pexiganan, respectively. The trend line depicts the median of the population size at each time-point. We tested differences in population size over time and differences between round one and two for each antimicrobial peptide with an ANOVA and adjusted with the Bonferroni method for multiple testing. For both, melittin and pexiganan, the population size changed significantly over time, while the differences in population size between round one and two and the interaction between time point and round were not significant ($p < 0.05$).

Figure S 3: Net growth rates resulting from the fit of the two-state model with 4 free parameters. For both (A) melittin and (B) pexiganan, r_P is not significantly different from 0. Significant differences between naive and primed treatment are indicated with asterisks.

Figure S 4. Parameter (A) r_N, (B) s_N, and (C) s_P of the two-state model fitted to the data of primed (orange) and naïve (blue) bacteria. The parameter r_P was set to 0. Significant differences between naïve (blue) and primed (orange) parameter values are indicated with asterisks. For parameter values, see Table S 5.

Figure S 5. Diagrammatic representation of (A) the two-state model (*17*) (see also Fig. 5A) and (B) our previously developed framework (*38*), which we extended by a persistent class according to *(17)*. Here, bacteria that replicate mutate with a mutation rate **μ** to a more resistant subpopulation R.

Figure S 6. Pharmacokinetic profile used in our stochastic simulations.

A

B

C   D

Fi**gure S 7.** Time-kill curves and PD curves of bacterial population dynamics exposed to AMPs. Time-kill experiments in which naïve (blue) and primed (orange) bacteria were exposed to (A) melittin and (B) pexiganan. The AMP dose used in each time-kill experiment is listed in the legend in each plot. PD functions of melittin (C) and pexiganan (D) fitted to the data in (A) and (B), respectively. Note that we excluded data points of experiments in which naïve bacteria were exposed to melittin (40 µg/ml) from the analysis to ensure the best fit. Parameter values are listed in table S5.

Figure S 8. Time until clearance and probability of resistance evolution for bacteria exposed to

pexiganan. In (A) each dot is an individual run.

Figure S 9. Predictions for different parameter values. (A) Time until clearance and (B) probability of resistance evolution is affected by the AMP decay rate k (dashed lines: k = 0.1, solid lines: k = 0.3). In (C) and (D) the mutation rate is varied (dashed lines: $\mu = 10^{-11}$, solid lines: $\mu = 10^{-9}$). All simulations are based on melittin data.

Figure S 10. Two-state model predicts biphasic decline depending on the model parameter values. If not varied, $r_N = -10$, $r_P = 0$, $s_N = 0.001$, and $s_P = 1$. Tolerance, i.e. the slope m_1 is mainly influenced by r_N, while the levels of persistence is influenced by all four parameters.

Figure S 11. Quality control of RNA sequencing by evaluating symmetry and distribution of the transcriptome counts, volcano plots showing different degrees of significance (A) and assessing dissimilarities of sample-based Euclidian hierarchical clustering for cells treated with priming concentrations of melitting and pexiganan (B). Datasets are based on RNAseq of three independent biological replicas.

Figure S 12. Microfluidic device design used for live imaging of priming and killing of *E. coli* MG1655 by AMPs. Z-values refers to the depth of chip features while X and Y represent the dimensions of 2D axes. Each chamber square compartment was designed to be similar in size (200 µm) to a microscope field with a magnification of 1000X.