1
Under-ice mesocosm experiments reveal the primacy of light but the importance of

zooplankton in winter phytoplankton dynamics

Allison R. Hrycik[1,2] and Jason D. Stockwell[1]

[1]University of Vermont, Rubenstein Ecosystem Science Laboratory, 3 College Street,

Burlington, Vermont, USA 05401

[2]University of Vermont Biology Department, 120A Marsh Life Science, 109 Carrigan Drive,

Burlington, Vermont 05405

Keywords: Winter limnology, food webs, phytoplankton, zooplankton, rotifers

Abstract

Factors that regulate planktonic communities under lake ice may be vastly different than the open-water season. However, under-ice food webs in temperate lakes are poorly understood, despite expected changes in light availability, ice cover, and snowfall associated with climate change. We hypothesized that light limitation (bottom-up control) outweighs zooplankton grazing (top-down control) on phytoplankton biovolume and community structure under ice in a north temperate lake. Using *in situ* under-ice mesocosms, we found that light had stronger effects on phytoplankton abundance than zooplankton, as expected. Specifically, low light limited growth of diatoms, cryptophytes, chrysophytes, and chlorophytes. Zooplankton, however, also significantly affected phytoplankton by decreasing diatoms and cryptophytes, in contrast to the common assumption that zooplankton grazing has negligible effects under ice. Ammonia and soluble reactive phosphorus decreased in high light treatments presumably through uptake by phytoplankton, whereas ammonia and soluble reactive phosphorus increased in high zooplankton treatments, likely through excretion. *In situ* experimental studies are commonly applied to understand food web dynamics in open-water conditions, but are extremely rare under ice. Our results suggest that changes in the light environment under ice have significant, rapid effects on phytoplankton growth and community structure and that zooplankton may play a more active role in winter food webs than previously thought. Changes in snow and ice dynamics associated with climate change may alter the light environment in ice-covered systems and significantly influence community structure.

                                            Introduction

The relatively recent and rapid changes in winter conditions in temperate zones have led

to declining ice cover in temperate lakes (Magnuson 2000, O'Reilly et al. 2015, Sharma et al.

2019) and altered snow cover and snowmelt dynamics (Adam et al. 2009, Musselman et al.

2017). However, the impact of such changes in winter conditions on lake food web dynamics

under ice is poorly understood (Salonen et al. 2009, Sommer et al. 2012) because winter

limnology has been under-studied compared to the open water season (Salonen et al. 2009,

Hampton et al. 2015). Experimental studies crucial to understand lake processes during the open-

water season (e.g., Schindler 1977, Lampert 1978, Huisman et al. 1999) are missing during the

ice-covered season, despite the potential for winter plankton community dynamics to set

inoculum conditions for the open water season. For example, phytoplankton can reach bloom

densities under ice (Twiss et al. 2012, Katz et al. 2015) and support abundant zooplankton

populations (Grosbois and Rautio 2018). A study of 101 lakes worldwide found that winter

chlorophyll-*a* (a proxy for phytoplankton abundance) reached an average of 43% of summer

chlorophyll-*a* (Hampton et al. 2017). Winter phytoplankton populations may also be important as

inoculum for spring blooms (Feuchtmayr et al. 2012). To better understand if and how changes

in snow and ice cover will affect biotic communities under ice and potentially into the open

water season, we must first disentangle the drivers of food web dynamics under ice.

Light is widely accepted as the major driver of phytoplankton production under ice

because it limits photosynthetic activity and can be highly variable depending on winter

conditions (Salonen et al. 2009, Hampton et al. 2015), including day length, ice thickness, ice

clarity, and especially snow cover (Bolsenga and Vanderploeg 1992, Eilertsen and Degerlund

2010, Lei et al. 2011). Physical factors such as light and temperature are considered the main

drivers of winter phytoplankton biovolume in the Plankton Ecology Group (PEG) Model that

describes planktonic community succession in temperate lakes (Sommer et al. 1986, 2012).

Predictably, total phytoplankton production is highest when light transmission is highest during

winter (Maeda and Ichimura 1973, Wojciechowska and Lenard 2014). Additionally, community

structure may be highly sensitive to changes in light levels because different taxa may be better

equipped to deal with different light conditions during winter. For example, diatoms have been

found at bloom densities in conditions with clear ice and minimal snowpack (Katz et al. 2015).

Some phytoplankton taxa with adaptations that allow them to succeed during light-limited

conditions, such as mixotrophic or mobile flagellated taxa, are often found in high proportions

under ice (McKnight et al. 2000, Özkundakci et al. 2016). To this end, we may expect a higher

proportion of known mixotrophic taxa, such as chrysophytes (Sanders et al. 1990, Caron et al.

1993), when light is limited, and higher total phytoplankton biovolume with high light

transmission.

Zooplankton can control phytoplankton biovolume and community structure during the

open water season through grazing, including selective feeding on specific phytoplankton groups

(Bergquist et al. 1985, Böing et al. 1998, Cyr and Curtis 1999). However, less is known about

top-down effects of zooplankton under ice, which makes interpretation and prediction of food

web interactions under ice and entering the spring phytoplankton bloom difficult (Sommer et al.

2012). Zooplankton actively feed under the ice (Vanderploeg et al. 1992, Grosbois and Rautio

2018), although they may be heavily dependent on accumulated lipid stores (Mariash et al. 2017,

Grosbois et al. 2017). Similar to the open water season, zooplankton grazing rates and

phytoplankton response during winter are expected to be dependent on the zooplankton and

phytoplankton species present, and their interactions. For example, winter zooplankton

communities are often dominated by copepods and rotifers (Primicerio and Klemetsen 1999,

Blank et al. 2009, Virro et al. 2009). A higher ratio of herbivorous to predatory rotifers may be

expected under ice if *Daphnia* are limited (Obertegger et al. 2011). Copepods may have

particularly strong impacts on phytoplankton community structure through selective raptorial

feeding (Ramakrishna Rao and Kumar 2002, Sommer et al. 2003), suggesting that winter

zooplankton communities that are actively feeding may influence under-ice phytoplankton

community structure and biovolume.

Changes in nutrient concentrations under ice may be closely linked to changes in

phytoplankton and zooplankton communities. Nutrients are generally not expected to limit

phytoplankton growth during winter, especially in eutrophic systems (Sommer et al. 2012).

However, we may still expect a signal from phytoplankton communities in nutrient

concentrations. Higher phytoplankton growth generally corresponds with reductions in forms of

nitrogen and phosphorus that can be assimilated quickly, such as soluble reactive phosphorus

(SRP), ammonia, and sometimes nitrate (Tamminen 1995, Glibert et al. 2016). Manipulation of

zooplankton biomass may also alter nutrient levels through excretion (Corner and Newell 1967,

Oliver et al. 2015). Other sources of nutrient inputs that are important in the open-water season,

including phosphorus release from sediment (Penn et al. 2000, Søndergaard et al. 2003), may

also be a significant source of phosphorus under ice if oxygen is limited (Joung et al. 2017). We

expected that nutrient concentrations would respond to changes in plankton communities but not

to the extent that they affected phytoplankton growth (Sommer et al. 2012).

In this study, we used an *in situ* under ice carboy experiment to test the relative

importance of zooplankton grazing versus light limitation on winter phytoplankton biovolume,

community structure, and nutrient concentrations in a north temperate lake. We hypothesized

that both low light and high zooplankton grazing would decrease phytoplankton biovolume, but

that light would be quantitatively more important than zooplankton to drive total phytoplankton

biovolume and have a larger impact on phytoplankton community structure under ice. Our

hypothesis follows the PEG Model (Sommer et al. 1986) and its recent update (Sommer et al.

2012), in which physical factors are thought to shape winter phytoplankton communities

compared to higher influence of zooplankton grazing and nutrient limitation during the open

water season. To our knowledge, our under-ice carboy experiment is the first application of this

type of mesocosm experiment under ice, despite wide application of carboy experiments during

the open water season (e.g., Havens et al. 1996, Nayar et al. 2004).

Methods

The experiment took place in Shelburne Pond, Vermont, a small, hypereutrophic system

(Ferber et al. 2004) with a mean depth of 3.4 m and maximum depth of 7.6 m (Vermont

Department of Environmental Conservation 2018). We initiated the experiment over 2

consecutive days with 12 carboys on 25 January and 12 carboys on 26 January 2018. The

transparent carboys were deployed on the north end of Shelburne Pond (44.39388°N,

73.16278°W) in an area with 0-2 cm of patchy snow on top of 30 cm of secondary ice

(black/clear ice; Block et al. 2018) and a water column depth of 4.6 m. We set carboys in a

randomized grid pattern of six carboys by four carboys spaced 5 m apart. Each carboy was

suspended by steel cable approximately 50 cm below the ice (Fig. 1A).

We tested four replicates for each of six treatments: (1) no zooplankton (0x)/unshaded,

(2) no zooplankton/shaded, (3) ambient zooplankton (1x)/unshaded, (4) ambient

zooplankton/shaded, (5) high zooplankton (10x)/unshaded, and (6) high zooplankton/shaded

(Fig. 1B/C). We mixed ambient water for all treatments in a 208-L plastic barrel. The barrel was

filled by raising and lowering the intake of a hand pump throughout the top 4.0 m of the water

column and filtering through a 350-µm sieve to remove large zooplankton. Pilot work indicated

350-µm was the best mesh size to remove large grazing zooplankton but kept most colonial

phytoplankton. The sieved ambient lake water in the barrel was mixed constantly as each group

of six 22.7-L carboys (i.e., one replicate for each treatment) was filled.

We then added zooplankton treatments directly to carboys. Two of the six carboys had no

zooplankton added, two had ambient zooplankton added, and two had 10X natural density of

large (i.e., sieved) zooplankton added. Zooplankton for the treatments were collected with a 13-

cm diameter, 64-µm Wisconsin net through the upper 4.0 m of the water column and then

retained on a 350-µm sieve. Desired densities were achieved for the ambient zooplankton

treatment by using a plankton splitter and adding half of the sieved net haul to each of the two

ambient density treatments because half of the volume strained for a 4-m tow was approximately

equal to the carboy volume. The 10X treatments had sieved zooplankton from five zooplankton

net tows added to each carboy. Finally, we covered one carboy from each zooplankton treatment

with greenhouse shade cloth that blocked 85% of incoming light to simulate the light-limiting

effects of snow cover. The set-up process was repeated twice on 25 January and twice on 26

January for a total of 4 replicates per treatment. We affixed a MK-9 light and temperature sensor

(Wildlife Computers Inc., Redmond, WA, USA) to the outside of one carboy without shade cloth

and between another carboy and its shade cloth on 25 January. We also added a HOBO

temperature sensor (Onset Computer Corporation, Bourne, MA, USA) to the inside of one

shaded and one unshaded carboy on 26 January. Light values from MK-9 sensors were converted

from relative units to $\mu E \cdot m^{-2} \cdot s^{-1}$ (Kotwicki et al. 2009). We did not leave any head space in the

carboys to simulate the sealed conditions of an ice-covered lake. The ice over the carboys re-

froze within one day of deployment.

We sampled phytoplankton, zooplankton, and nutrients while filling carboys to measure

initial experimental conditions. We collected three 100-mL water samples and preserved them in

1% Lugol's solution for phytoplankton enumeration. Three replicate 10-L water samples from

the barrel that was previously sieved with 350-µm mesh were filtered through a 20-µm sieve for

microzooplankton and rotifer enumeration. We also preserved the zooplankton filtered out of

each barrel with the 350 µm sieve. All zooplankton and rotifers were anesthetized with Alka

Seltzer before preservation with 70% ethanol. Finally, three replicate 500-mL water samples

were collected from each barrel for nutrient analyses. Once back at the lab, each nutrient sample

was split into several different portions for nutrient analyses. A 100-mL sample was preserved

with three drops of sulfuric acid to achieve a pH of 2.0 for later analysis of total phosphorus

(TP), total nitrogen (TN), and total organic carbon (TOC). Two 45-mL samples were filtered

through 0.45-µm syringe filters and frozen: one for soluble reactive phosphorus (SRP), and one

for ammonia and nitrate + nitrite (NO_x) quantification.

Carboys were extracted from the lake 14 days after deployment. Although most summer

carboy experiments are much shorter in duration (e.g., Havens et al. 1996, Miller et al. 1997,

Griniene et al. 2016), we expected that slow phytoplankton growth rates at low temperatures

(Cloern 1977, Richardson et al. 2000) would necessitate a longer incubation time. Each carboy

was inverted several times to homogenize contents before opening. We collected three

phytoplankton samples by filling 125-mL bottles with water and its associated phytoplankton

community, which was later measured out to 100 mL and preserved in Lugol's solution. One

500-mL nutrient sample was taken per carboy, and split into portions for different analyses, as

described above. We strained the remaining 21.84 L of water through a 20-µm Wisconsin net to

sample crustacean zooplankton and rotifers. The organisms were anesthetized with Alka Seltzer

then preserved in 70% ethanol.

We identified phytoplankton to genus and counted full fields of view at 400x until

reaching at least 300 natural units (cells for single-celled species or colonies for colonial

species). We measured ten natural units per genus for each sample to calculate biovolume using

Spot Basic software (Spot Imaging, Sterling Heights, MI). Dimensions measured were dependent

on phytoplankton taxa present, e.g., we measured diameter for spherical cells and diameter and

length for ellipsoid cells (Hillebrand et al. 1999). Within colonies, we measured ten individual

cells per colony when possible. If fewer than ten cells were present or clearly visible, we

measured all cells. We used those median measurements to calculate taxon-specific biovolume

for each sample (Hillebrand et al. 1999), then taxon-specific biovolume was multiplied by cell

density to estimate total biovolume per sample for each taxon. We only processed one out of the

three phytoplankton samples collected per carboy because replicates within each carboy were

very similar and would not have added to statistical power due to pseudoreplication. Analysis of

three pairs of phytoplankton samples from the same carboys showed an average of 4.2%

difference in cell counts for each genus.

We processed rotifers and crustacean zooplankton by measuring and counting at least 200

individuals per sample (200 rotifers and 200 zooplankton). Rotifers were counted and measured

on a Nikon Eclipse Ni-U compound microscope (Nikon, Tokyo, Japan) with Spot Basic software

(Spot Imaging, Sterling Heights, Michigan, USA), while zooplankton were identified and

enumerated on an Olympus SZX12 dissecting microscope (Olympus Corporation, Tokyo, Japan)

interfaced with a GTCO CalComp digitizer for measurements (Turning Technologies, Inc.,

Youngstown, Ohio, USA). Rotifer and crustacean zooplankton biomass were calculated using

length-to-biomass conversions (all crustacean zooplankton and most rotifers) or length/width-to-

biomass conversions (*Filinia* rotifers) (Watkins et al. 2011, United States Environmental

Protection Agency 2016).

We measured nutrient concentrations primarily to ensure that we did not artificially limit

nutrients in our study. Nutrient samples were either stored frozen (SRP, ammonia, and NO_x) or

acidified and refrigerated (TN, TOC, and TP) until analysis. We measured TN and TOC on a

TOC-L total organic carbon analyzer with a TNM-L TN measuring unit (Shimadzu Corporation,

Kyoto, Japan). We analyzed TP and SRP using the molybdenum colorimetry method (USEPA

1993) with ascorbic acid modification and a persulfate digestion for TP on a Shimadzu UV-VIS

2600 spectrophotometer (Shimadzu Corporation, Kyoto, Japan). Ammonia and NO_x were

measured on a SEAL AA3 continuous flow autoanalyzer (SEAL Analytical, Mequon,

Wisconsin, USA) using Method No. G-171-96 Rev. 15 with salicylate for ammonia and Method

No. G-172-96 Rev. 18 for NO_x.

All response variables (phytoplankton density and biovolume, rotifer density and

biomass, and nutrient concentrations) were analyzed using ANOVAs ($\alpha=0.05$) with zooplankton

and light as predictor variables. We then used the ANOVA output for variance partitioning to

quantify the contribution (partial R^2) of zooplankton and light separately on response variables.

The values are reported as R^2_{zoop} and R^2_{light}. We originally ran ANOVAs with an interaction term

(zooplankton*light), but this term was removed for final ANOVAs used in variance partitioning

because we found only one significant interaction effect out of 22 tests. Samples taken during

mesocosm setup were not included in statistical analyses because they do not represent treatment

effects, but are included in visualizations for demonstrative purposes.

We performed non-metric multidimensional scaling (nMDS) with Bray-Curtis distance

on phytoplankton species composition to visualize differences in phytoplankton communities

between treatments using the R package "vegan" (version 2.4.3; Oksanen et al. 2013). We

removed the small number of phytoplankton records where phytoplankton could not be identified

to genus, which should not significantly alter nMDS interpretation (Pos et al. 2014). We used

perMANOVA with Bray-Curtis distance and 999 permutations to test significance of light and

zooplankton treatments on overall phytoplankton community composition (Anderson 2001,

Oksanen et al. 2013).

243   **Results**

Light and temperature behaved as expected throughout the experiments. Light was

limited by greenhouse shade cloth (Supporting Information Fig. S1). Temperature was virtually

indistinguishable between light and shaded treatments, although internal carboy temperatures

were slightly higher than external water temperatures. However, the difference between internal

and external temperature (<0.3°C) was small compared to the overall increase in water

temperature over the course of the experiment (Fig. S1).

Zooplankton density and biomass were different between treatments, but differences

were not as great as intended (Fig. 2). Our "0x" zooplankton treatments averaged (± SD) 64.5 ±

14.21 µg dry/L, "1x" treatments averaged 99.8 ± 5.36 µg dry/L, and "10x" zooplankton averaged

338.6 ± 56.93 µg dry/L. That is, our "10x" zooplankton level had 3.4x the zooplankton biomass

as our "1x" treatment. Consequently, we refer to zooplankton levels as low, medium, and high.

Zooplankton biomass was dominated by *Diacyclops thomasi* and zooplankton density was

dominated by both *D. thomasi* and copepod nauplii (Fig. S2). Crustacean zooplankton body size

followed a bimodal distribution with a smaller peak that represented copepod nauplii and

Chydorus spp. and the larger peak represented adult zooplankton (Fig. S3). The smaller

zooplankton were depleted in treatments with higher densities of large-bodied zooplankton (Fig.

S3).

Differences in phytoplankton community composition among treatments was primarily

driven by light, although some groups were also significantly affected by zooplankton levels.

The biovolume of total phytoplankton and chlorophytes, and density and biovolume of diatoms,

cryptomonads, and chrysophytes significantly increased with light (Table 1; Fig. 3; Fig. 4). Light

most strongly increased diatoms ($R^2_{light} = 0.73$ for density and $R^2_{light} = 0.62$ for biovolume) and

chrysophytes ($R^2_{light} = 0.76$ for density and $R^2_{light} = 0.71$ for biovolume) (Table 1). Zooplankton

significantly decreased diatom density and cryptomonad biovolume but explained less variation

overall than light treatments ($R^2_{zoop} = 0.07$ for diatom density and $R^2_{zoop} = 0.33$ for cryptophyte

biovolume) (Table 1). Cyanobacteria and haptophytes were also found in all samples but did not

vary by treatment (Table 1). Phytoplankton samples comprised diatoms, chlorophytes,

cyanobacteria, cryptomonads, dinoflagellates, chrysophytes, haptophytes, synurophytes,

desmids, and euglenoids (Table 2).

Non-metric multidimensional scaling resulted in a clear distinction in phytoplankton

communities between shaded and unshaded treatments for all axes, but little effect from

zooplankton (Fig. 5). Treatments at the beginning of experiments were more similar to shaded

treatments than unshaded treatments. Our final ordination had three axes to reduce stress from

0.22 with two axes to 0.15 with three axes (Fig. 5). Phytoplankton genera that drove separations

along nMDS axes were mostly rare species that were only found in some treatments such as

Crucigenia, Diatoma, Aphanocapsa, and *Dictyosphaerium* for the first axis and *Oocystis,*

Stephanodiscus, Selenastrum, and *Gonium* for the second axis (Fig. S4). More common genera

such as *Chrysochromulina* and *Woronichinia* were found across treatments, so contributed little

to separations in nMDS axes (Fig. S4). PerMANOVA indicated a significant effect of light

treatment (pseudoF$_{1,20}$ = 12.3; p=0.001) but no effect of zooplankton treatment (pseudoF$_{2,20}$ =

1.36; p=0.184).

Rotifer density was not dependent on experimental conditions, but rotifer biomass was

significantly affected by zooplankton levels. Rotifer biomass increased with increased

zooplankton (Fig. 2; Table 1). Changes in zooplankton treatments accounted for 33% of the

variation in rotifer biomass (i.e., R^2_{zoop} = 0.33). *Keratella cochlearis* was the dominant rotifer

species found in mesocosms and made up > 90% of individuals, but only 39% of biomass due to

their small size (Fig. S5). Other rotifers found in mesocosms were *Keratella hiemalis*,

Brachionus angularis, *Asplanchna spp.*, and *Filinia spp*.

Some nutrient levels were significantly different between treatments (Fig. 6; Table 1).

SRP and ammonia significantly decreased with light and significantly increased with

zooplankton. Light explained 73% of the variation in SRP and 72% of the variation in ammonia,

while zooplankton explained 15% of the variation in SRP and 21% in ammonia (Table 1).

Zooplankton also significantly increased TN and explained 20% of the variance between

treatments (Table 1). Neither light nor zooplankton significantly affected TP, NO$_x$, or TOC (Fig.

6; Table 1).

Discussion

We found that light had a stronger influence on phytoplankton biovolume and community

composition than zooplankton grazing in our under-ice carboy experiment. The results support

our hypothesis that bottom-up control exceeds top-down control on phytoplankton under ice.

Specifically, the greatest contrast in chrysophytes and diatoms occurred when light was

manipulated. However, zooplankton grazing had non-negligible effects on both phytoplankton

and nutrient concentrations. The significance of zooplankton in our experiment contrasts with the

PEG Model assumption that the effects of zooplankton grazing are low during winter (Sommer

et al. 1986, 2012). Actively overwintering zooplankton have the potential to impact

phytoplankton biovolume under ice, likely through selective feeding and nutrient cycling. Thus,

the role of winter zooplankton may be more nuanced than primarily functioning as a standing

stock to graze the spring phytoplankton bloom after ice-out as proposed in the "overwintering

zooplankton" scenario of the updated PEG Model (Sommer et al. 2012). Changes in

phytoplankton community dynamics with zooplankton biomass mediated through light

conditions could have direct consequences for which phytoplankton communities are present at

ice-out.

Light had a stronger effect on phytoplankton community composition than zooplankton,

supporting the hypothesis that light is the main driver of phytoplankton biovolume under ice.

The difference was apparent in both nMDS visualizations and ANOVA analysis of specific

phytoplankton groups. Changes in community composition contributed to differences in overall

phytoplankton biovolume, despite relatively constant total phytoplankton density across

treatments. In particular, we observed higher proportion of chrysophytes (mostly *Chrysococcus*)

and diatoms when more light was available. *Chrysococcus* can be successful under the ice in

north temperate lakes (Phillips and Fawley 2002) so *Chrysococcus* as a dominant phytoplankton

in Shelburne Pond during winter is not surprising. Although *Chrysococcus* is a known mixotroph

that can withstand low light conditions (Olrik 1998), it still responded strongly to high light in

our experiment. Diatoms are also often abundant under clear ice with high light transmission

similar to our unshaded experimental conditions (Katz et al. 2015). Interestingly, the changes we

observed in phytoplankton community composition took place over just two weeks, indicating

that highly variable or rapidly changing light environments (e.g., patchy snow, rapid snowfall on

top of clear ice, or snow that is abruptly windswept off ice) could have large impacts on

phytoplankton community structure over relatively short time scales.

Light treatments significantly altered ammonia and SRP presumably through increased

phytoplankton production. Nitrate decreased in treatments with high phytoplankton levels, but

ammonia remained constant across treatments. Many phytoplankton can take up nitrate faster

than ammonia, and in some cases, ammonia uptake may inhibit nitrate uptake, although these

relationships can be highly variable (Dortch 1990, Glibert et al. 2016). SRP and nitrate most

likely did not limit phytoplankton growth in mesocosms because concentrations of both were

higher than concentrations found in Shelburne Pond during summer when phytoplankton

production is highest (Ferber et al. 2004).

Despite discrepancies between expected and actual zooplankton biomass, zooplankton

still had a major effect in carboys through consumption of certain phytoplankton groups and

alteration of nutrient concentrations. Zooplankton significantly decreased total phytoplankton

biovolume, cryptomonad biovolume, and diatom density, suggesting that zooplankton selectively

grazed larger cryptophytes (specifically *Cryptomonas*) and smaller or non-colonial diatoms.

Diatoms and cryptophytes are known food sources for both rotifers and crustacean zooplankton

(Hansen 2000, Mohr and Adrian 2002, Zhou et al. 2011, Tõnno et al. 2016). Furthermore,

nutrients showed signals of zooplankton activity through increases in SRP, ammonia, and TN

with increased zooplankton. Zooplankton increase ammonia and SRP through excretion

(Moegenburg and Vanni 1991), whereas higher TN was likely the direct result of higher

zooplankton biomass. Although top-down control from zooplankton grazing was minimal in

comparison to bottom-up control from light limitation in this experiment, zooplankton have the

potential to indirectly affect phytoplankton though mobilization of nutrients during a period

when external nutrient inputs are low.

Rotifers unexpectedly increased in biomass as crustacean zooplankton increased. We

would expect rotifers to decrease when crustacean zooplankton are abundant because crustacean

zooplankton typically outcompete rotifers when phytoplankton resources are limited (Gilbert

1985, Fussmann 1996), and copepods such as *Diacyclops* may directly consume rotifers (Ciros-

Pérez et al. 2004). One possibility is that the increased nutrients from adding zooplankton

increased primary production that was then consumed by zooplankton and rotifers. However, we

cannot evaluate whether nutrient cycling and phytoplankton regeneration rates increased because

we only measured standing stock of phytoplankton at the end of experiments. Another potential

explanation is that additional rotifers were added with the zooplankton treatments. However, this

possibility is unlikely because samples from the 350-µm mesh sieve that were prevented from

going into the barrels indicated extremely low rotifer densities of 0.26 individuals/L. Another

possibility is that rotifers experienced competitive release if large crustacean zooplankton

outcompeted small crustacean zooplankton that may consume phytoplankton in the same size

range as rotifers. In this case, having a high prevalence of large crustacean zooplankton would

release rotifers from competition. This possibility seems most likely because small-bodied

zooplankton were found in lower proportions in treatments with more large-bodied zooplankton.

The guild ratio theory (Obertegger et al. 2011) that predicts that *Daphnia* would outcompete

microphagous (herbivorous) rotifers was potentially supported because we found low *Daphnia*

abundances and high microphagous rotifer abundance in all treatments.

Mesocosms were effective in maintaining physical parameters such as light and

temperature but were less predictable in maintaining zooplankton levels. Water temperatures

increased throughout the course of the experiment similarly between shaded and unshaded

treatments, so temperature was not an influential factor in differences between treatments. The

shade cloth covers maintained differences in light readings between treatments, including during

a significant snowfall the night before we began extracting carboys at the end of the experiment

(February 8). Zooplankton maintained differences between treatments, but at lower magnitudes

than expected. The most likely explanation is that zooplankton experienced mortality as they

were collected, sieved, and added to carboys. Alternatively, the highest zooplankton densities

may have exceeded the carrying capacity of the carboys and experienced mortality during the

experiment.

Mesocosm studies are necessarily limited in their scope to manipulate only the factors

under study. In this experiment, we suspended our sealed carboys just below the ice. The limited

mixing with the rest of the water column negated the potential for phytoplankton to settle to the

bottom of the lake. However, our experiments were conducted during a period when the water

was warming before ice-out and was likely in a convective mixing state (Bruesewitz et al. 2015).

Convective mixing could re-suspend phytoplankton such as diatoms (Vehmaa and Salonen

2009), and thus, we would not expect phytoplankton to settle out as quickly as they would in a

winter stratified state. Additionally, our sealed systems may have relaxed selective pressures for

flagellated phytoplankton taxa by limiting their ability to migrate in the water column.

In this experiment, we demonstrated that light is the main driving factor of phytoplankton

biovolume and community structure under ice. Variations in light can also lead to significant

changes in nutrient cycling. However, the role of zooplankton under ice should not be

overlooked. Zooplankton decreased some phytoplankton taxa and altered nutrient concentrations

in mesocosms, which suggests that we may miss important contributions of zooplankton in

shaping phytoplankton communities and nutrient cycling under ice if we assume that

overwintering zooplankton have negligible effects. Furthermore, high prevalence of copepod

nauplii suggests that some crustacean zooplankton reproduced under ice. Winter copepod

reproduction is often overlooked in temperate lakes, despite its occurrence in multiple systems

(this study; Vanderploeg et al. 1992). Additional application of open water experimental

techniques to ice-covered ecosystems is an important step in disentangling food web drivers

under ice.

The results of this experiment are relevant to understand what may happen with changing

winter conditions associated with climate change. As climate change progresses, some regions

are expected to experience increased snowpack and other areas are expected to see decreased

snowpack (Räisänen 2008, Trujillo and Molotch 2014). Most regions are also predicted to have

earlier snowmelt (Stewart et al. 2004, Klein et al. 2016) that is potentially more protracted

(Musselman et al. 2017). Our study which simulated a change in snow cover of just a few

centimeters (Bolsenga and Vanderploeg 1992) for two weeks was enough to significantly alter

phytoplankton community structure, indicating that minor events such as rain-on-snow events

that melt a layer of snow or slightly altered snowfall totals may have disproportionately large

effects on phytoplankton communities. Ice cover duration is expected to shorten in most regions

as a result of climate change (Magnuson 2000, Magee and Wu 2017, Sharma et al. 2019), and

while outside the scope of this study, we would expect that shorter ice duration would alter

spring mixing regimes (Roberts et al. 2018) and potentially increase zooplankton survival

(Preston and Rusak 2010). As indicated in this study, higher zooplankton biomass may increase

concentrations of nutrients relevant for phytoplankton growth. Increases in SRP and ammonia

could increase nutrient cycling rates if light levels are sufficiently high to sustain phytoplankton

growth. The interplay of light transmission and zooplankton grazing and their effects on

phytoplankton communities will likely change rapidly with climate change.

425   **Acknowledgements**

We thank Brad Roy, Thomas Hamling, Ben Block, Jessica Griffin, Wilton Burns, Posy

Labombard, Taylor Stewart, Brianna Wilkins, Hannah Lister, Natalie Flores, Lisa Izzo, Frances

Ianucci, Hannah Lachance, and Brian O'Malley for field work assistance and Jonathan

Rickwood for lab assistance. Nicholas Gotelli, Ana Morales-Williams, and Andrew Schroth were

instrumental in project design. We also thank the Marsden and Stockwell laboratories for

comments on earlier drafts of the manuscript. This work was supported by the University of

Vermont Biology Department, the Morse Fund, the UVM Roberto Fabri Fialho Research Award

to ARH, the Vermont Space Grant Consortium under NASA Cooperative Agreement

NNX15AP86H, and a NASA Earth and Space Science Fellowship to ARH under NASA

Cooperative Agreement 80NSSC18K1394 P00001.

Literature Cited

Adam, J. C., A. F. Hamlet, and D. P. Lettenmaier. 2009. Implications of global climate change

for snowmelt hydrology in the twenty-first century. Hydrological Processes 23:962–972.

Anderson, M. J. 2001. A new method for non-parametric multivariate analysis of variance.

Austral Ecology 26:32–46.

Bergquist, A. M., S. R. Carpenter, and J. C. Latino. 1985. Shifts in phytoplankton size structure

and community composition during grazing by contrasting zooplankton assemblages.

Limnology and Oceanography 30:1037–1045.

Blank, K., J. Haberman, M. Haldna, and R. Laugaste. 2009. Effect of winter conditions on spring

nutrient concentrations and plankton in a large shallow Lake Peipsi (Estonia/Russia).

Aquatic Ecology 43:745–753.

Block, B. D., B. A. Denfeld, J. D. Stockwell, G. Flaim, H. F. Grossart, L. B. Knoll, D. B. Maier,

R. L. North, M. Rautio, J. A. Rusak, S. Sadro, G. A. Weyhenmeyer, A. J. Bramburger, D.

K. Branstrator, K. Salonen, and S. E. Hampton. 2018. The unique methodological

challenges of winter limnology. Limnology and Oceanography: Methods 17:lom3.10295.

Böing, W. J., A. Wagner, H. Voigt, T. Deppe, and J. Benndorf. 1998. Phytoplankton responses to

grazing by *Daphnia galeata* in the biomanipulated Bautzen reservoir. Hydrobiologia

389:101–114.

Bolsenga, S. J., and H. A. Vanderploeg. 1992. Estimating photosynthetically available radiation

into open and ice-covered freshwater lakes from surface characteristics; a high

transmittance case study. Hydrobiologia 243–244:95–104.

Bruesewitz, D. A., C. C. Carey, D. C. Richardson, and K. C. Weathers. 2015. Under-ice thermal

stratification dynamics of a large, deep lake revealed by high-frequency data. Limnology

and Oceanography 60:347–359.

Caron, D., R. Sanders, E. Lim, C. Marras, L. Amaral, S. Whitney, R. Aoki, and K. Porters. 1993.

Light-dependent phagotrophy in the freshwater mixotrophic chrysophyte *Dinobryon*

cylindricum. Microbial Ecology 25:93–111.

Ciros-Pérez, J., M. J. Carmona, S. Lapesa, and M. Serra. 2004. Predation as a factor mediating

resource competition among rotifer sibling species. Limnology and Oceanography 49:40–

50.

Cloern, J. E. 1977. Effects of light intensity and temperature on *Cryptomonas ovata*

(Cryptophyceae) growth and nutrient uptake rates. Journal of Phycology 13:389–395.

Corner, E. D. S., and B. S. Newell. 1967. On the Nutrition and metabolism of zooplankton IV.

The forms of nitrogen excreted By *Calanus*. Journal of the Marine Biological Association

of the United Kingdom 47:113–120.

Cyr, H., and J. M. Curtis. 1999. Zooplankton community size structure and taxonomic

composition affects size-selective grazing in natural communities. Oecologia 118:306–315.

Dortch, Q. 1990. The interaction between ammonium and nitrate uptake in phytoplankton.

Marine Ecology Progress Series 61:183–201.

Eilertsen, H. C., and M. Degerlund. 2010. Phytoplankton and light during the northern high-

latitude winter. Journal of Plankton Research 32:899–912.

Ferber, L. R., S. N. Levine, A. Lini, and G. P. Livingston. 2004. Do cyanobacteria dominate in

eutrophic lakes because they fix atmospheric nitrogen? Freshwater Biology 49:690–708.

Feuchtmayr, H., S. J. Thackeray, I. D. Jones, M. De Ville, J. Fletcher, B. James, and J. Kelly.

2012. Spring phytoplankton phenology - are patterns and drivers of change consistent

among lakes in the same climatological region? Freshwater Biology 57:331–344.

Fussmann, G. 1996. The importance of crustacean zooplankton in structuring rotifer and

phytoplankton communities; an enclosure study. Journal of Plankton Research 18:1897–

1915.

Gilbert, J. J. 1985. Competition between Rotifers and *Daphnia*. Ecology 66:1943–1950.

Glibert, P. M., F. P. Wilkerson, R. C. Dugdale, J. A. Raven, C. L. Dupont, P. R. Leavitt, A. E.

Parker, J. M. Burkholder, and T. M. Kana. 2016. Pluses and minuses of ammonium and

nitrate uptake and assimilation by phytoplankton and implications for productivity and

community composition, with emphasis on nitrogen-enriched conditions. Limnology and

Oceanography 61:165–197.

Griniene, E., S. Šulčius, and H. Kuosa. 2016. Size-selective microzooplankton grazing on the

phytoplankton in the Curonian Lagoon (SE Baltic Sea). Oceanologia 58:292–301.

Grosbois, G., H. Mariash, T. Schneider, and M. Rautio. 2017. Under-ice availability of

phytoplankton lipids is key to freshwater zooplankton winter survival. Scientific Reports

7:11543.

Grosbois, G., and M. Rautio. 2018. Active and colorful life under lake ice. Ecology 99:752–754.

Hampton, S. E., A. W. E. Galloway, S. M. Powers, T. Ozersky, K. H. Woo, R. D. Batt, S. G.

Labou, C. M. O'Reilly, S. Sharma, N. R. Lottig, E. H. Stanley, R. L. North, J. D. Stockwell,

R. Adrian, G. A. Weyhenmeyer, L. Arvola, H. M. Baulch, I. Bertani, L. L. Bowman, C. C.

Carey, J. Catalan, W. Colom-Montero, L. M. Domine, M. Felip, I. Granados, C. Gries, H.-

P. Grossart, J. Haberman, M. Haldna, B. Hayden, S. N. Higgins, J. C. Jolley, K. K.

Kahilainen, E. Kaup, M. J. Kehoe, S. MacIntyre, A. W. Mackay, H. L. Mariash, R. M.

McKay, B. Nixdorf, P. Nõges, T. Nõges, M. Palmer, D. C. Pierson, D. M. Post, M. J. Pruett,

M. Rautio, J. S. Read, S. L. Roberts, J. Rücker, S. Sadro, E. A. Silow, D. E. Smith, R. W.

Sterner, G. E. A. Swann, M. A. Timofeyev, M. Toro, M. R. Twiss, R. J. Vogt, S. B.

Watson, E. J. Whiteford, and M. A. Xenopoulos. 2017. Ecology under lake ice. Ecology

Letters 20:98–111.

Hampton, S. E., M. V. Moore, T. Ozersky, E. H. Stanley, C. M. Polashenski, and A. W. E.

Galloway. 2015. Heating up a cold subject: prospects for under-ice plankton research in

lakes. Journal of Plankton Research 37:277–284.

Hansen, A.-M. 2000. Response of ciliates and *Cryptomonas* to the spring cohort of a cyclopoid

copepod in a shallow hypereutrophic lake. Journal of Plankton Research 22:185–203.

Havens, K., T. East, and J. Beaver. 1996. Experimental studies of zooplankton-phytoplankton-

nutrient interactions in a large subtropical lake (Lake Okeechobee, Florida, U.S.A.).

Freshwater Biology 36:579–597.

Hillebrand, H., C.-D. Dürselen, D. Kirschtel, U. Pollingher, and T. Zohary. 1999. Biovolume

Calculation for Pelagic and Benthic Microalgae. Journal of Phycology 35:403–424.

Huisman, J., R. R. Jonker, C. Zonneveld, and F. J. Weissing. 1999. Competition for light

between phytoplankton species: Experimental tests of mechanistic theory. Ecology 80:211–

222.

Joung, D., M. Leduc, B. Ramcharitar, Y. Xu, P. D. F. Isles, J. D. Stockwell, G. K. Druschel, T.

Manley, and A. W. Schroth. 2017. Winter weather and lake-watershed physical

configuration drive phosphorus, iron, and manganese dynamics in water and sediment of

ice-covered lakes. Limnology and Oceanography 62:1620–1635.

Katz, S. L., L. R. Izmest'eva, S. E. Hampton, T. Ozersky, K. Shchapov, M. V. Moore, S. V.

Shimaraeva, and E. A. Silow. 2015. The "Melosira years" of Lake Baikal: Winter

environmental conditions at ice onset predict under-ice algal blooms in spring. Limnology

and Oceanography 60:1950–1964.

Klein, G., Y. Vitasse, C. Rixen, C. Marty, and M. Rebetez. 2016. Shorter snow cover duration

since 1970 in the Swiss Alps due to earlier snowmelt more than to later snow onset.

Climatic Change 139:637–649.

Kotwicki, S., A. De Robertis, P. von Szalay, and R. Towler. 2009. The effect of light intensity on

the availability of walleye pollock (*Theragra chalcogramma*) to bottom trawl and acoustic

surveys. Canadian Journal of Fisheries and Aquatic Sciences 66:983–994.

Lampert, W. 1978. Release of dissolved organic carbon by grazing zooplankton. Limnology and

Oceanography 23:831–834.

Lei, R., M. Leppäranta, A. Erm, E. Jaatinen, and O. Pärn. 2011. Field investigations of apparent

optical properties of ice cover in Finnish and Estonian lakes in winter 2009. Estonian

Journal of Earth Sciences 60:50.

Maeda, O., and S.-E. Ichimura. 1973. On the High Density of a Phytoplankton Population Found

in a Lake under Ice. Internationale Revue der gesamten Hydrobiologie und Hydrographie

58:673–689.

Magee, M. R., and C. H. Wu. 2017. Effects of changing climate on ice cover in three

morphometrically different lakes. Hydrological Processes 31:308–323.

Magnuson, J. J. 2000. Historical Trends in Lake and River Ice Cover in the Northern

Hemisphere. Science 289:1743–1746.

Mariash, H. L., M. Cusson, and M. Rautio. 2017. Fall Composition of Storage Lipids is

Associated with the Overwintering Strategy of Daphnia. Lipids 52:83–91.

McKnight, D. M., B. L. Howes, C. D. Taylor, and D. D. Goehringer. 2000. Phytoplankton

dynamics in a stably stratified Antarctic lake during winter darkness. Journal of Phycology 36:852–861.

Miller, C., P. Glibert, G. Berg, and M. Mulholland. 1997. Effects of grazer and substrate amendments on nutrient and plankton dynamics in estuarine enclosures. Aquatic Microbial Ecology 12:251–261.

Moegenburg, S. M., and M. J. Vanni. 1991. Nutrient regeneration by zooplankton: effects on nutrient limitation of phytoplankton in a eutrophic lake. Journal of Plankton Research 13:573–588.

Mohr, S., and R. Adrian. 2002. Reproductive success of the rotifer *Brachionus calyciflorus* feeding on ciliates and flagellates of different trophic modes. Freshwater Biology 47:1832–1839.

Musselman, K. N., M. P. Clark, C. Liu, K. Ikeda, and R. Rasmussen. 2017. Slower snowmelt in a warmer world. Nature Climate Change 7:214–219.

Nayar, S., B. P. L. Goh, and L. M. Chou. 2004. Environmental impact of heavy metals from dredged and resuspended sediments on phytoplankton and bacteria assessed in *in situ* mesocosms. Ecotoxicology and Environmental Safety 59:349–369.

O'Reilly, C. M., S. Sharma, D. K. Gray, S. E. Hampton, J. S. Read, R. J. Rowley, P. Schneider, J. D. Lenters, P. B. McIntyre, B. M. Kraemer, G. A. Weyhenmeyer, D. Straile, B. Dong, R. Adrian, M. G. Allan, O. Anneville, L. Arvola, J. Austin, J. L. Bailey, J. S. Baron, J. D. Brookes, E. de Eyto, M. T. Dokulil, D. P. Hamilton, K. Havens, A. L. Hetherington, S. N. Higgins, S. Hook, L. R. Izmest'eva, K. D. Joehnk, K. Kangur, P. Kasprzak, M. Kumagai, E. Kuusisto, G. Leshkevich, D. M. Livingstone, S. MacIntyre, L. May, J. M. Melack, D. C. Mueller-Navarra, M. Naumenko, P. Noges, T. Noges, R. P. North, P.-D. Plisnier, A. Rigosi,

A. Rimmer, M. Rogora, L. G. Rudstam, J. A. Rusak, N. Salmaso, N. R. Samal, D. E.

Schindler, S. G. Schladow, M. Schmid, S. R. Schmidt, E. Silow, M. E. Soylu, K. Teubner,

P. Verburg, A. Voutilainen, A. Watkinson, C. E. Williamson, and G. Zhang. 2015. Rapid

and highly variable warming of lake surface waters around the globe. Geophysical Research

Letters 42:10,773-10,781.

Obertegger, U., H. A. Smith, G. Flaim, and R. L. Wallace. 2011. Using the guild ratio to

characterize pelagic rotifer communities. Hydrobiologia 662:157–162.

Oksanen, J., F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. B. O'Hara, G. L. Simpson,

P. Solymos, M. H. H. Stevens, and H. Wagner. 2013. Vegan: Community ecology package.

Oliver, S. K., D. K. Branstrator, T. R. Hrabik, S. J. Guildford, and R. E. Hecky. 2015. Nutrient

excretion by crustacean zooplankton in the deep chlorophyll layer of Lake Superior.

Canadian Journal of Fisheries and Aquatic Sciences 72:390–399.

Olrik, K. 1998. Ecology of mixotrophic flagellates with special reference to Chrysophyceae in

Danish lakes. Pages 329–338Phytoplankton and Trophic Gradients. Springer Netherlands,

Dordrecht.

Özkundakci, D., A. S. Gsell, T. Hintze, H. Täuscher, and R. Adrian. 2016. Winter severity

determines functional-trait composition of phytoplankton in seasonally ice-covered lakes.

Global change biology 22:284–298.

Penn, M. R., M. T. Auer, S. M. Doerr, C. T. Driscoll, C. M. Brooks, and S. W. Effler. 2000.

Seasonality in phosphorus release rates from the sediments of a hypereutrophic lake under a

matrix of pH and redox conditions. Canadian Journal of Fisheries and Aquatic Sciences

57:1033–1041.

Phillips, K. A., and M. W. Fawley. 2002. Winter phytoplankton community structure in three

shallow temperate lakes during ice cover. Hydrobiologia 470:97–113.

Pos, E., J. E. Guevara Andino, D. Sabatier, J.-F. Molino, N. Pitman, H. Mogollón, D. Neill, C.

Cerón, G. Rivas, A. Di Fiore, R. Thomas, M. Tirado, K. R. Young, O. Wang, R. Sierra, R.

García-Villacorta, R. Zagt, W. Palacios, M. Aulestia, and H. ter Steege. 2014. Are all

species necessary to reveal ecologically important patterns? Ecology and Evolution 4:4626–

4636.

Preston, N. D., and J. A. Rusak. 2010. Homage to Hutchinson: does inter-annual climate

variability affect zooplankton density and diversity? Pages 165–177Fifty years after the

"Homage to Santa Rosalia": Old and new paradigms on biodiversity in aquatic ecosystems.

Springer Netherlands, Dordrecht.

Primicerio, R., and A. Klemetsen. 1999. Zooplankton seasonal dynamics in the neighbouring

lakes Takvatn and Lombola (Northern Norway). Hydrobiologia 411:19–29.

Räisänen, J. 2008. Warmer climate: less or more snow? Climate Dynamics 30:307–319.

Ramakrishna Rao, T., and R. Kumar. 2002. Patterns of prey selectivity in the cyclopoid copepod

Mesocyclops thermocyclopoides. Aquatic Ecology 36:411–424.

Richardson, T. L., C. E. Gibson, and S. I. Heaney. 2000. Temperature, growth and seasonal

succession of phytoplankton in Lake Baikal, Siberia. Freshwater Biology 44:431–440.

Roberts, D. C., A. L. Forrest, G. B. Sahoo, S. J. Hook, and S. G. Schladow. 2018. Snowmelt

Timing as a Determinant of Lake Inflow Mixing. Water Resources Research 54:1237–1251.

Salonen, K., M. Leppäranta, M. Viljanen, and R. D. Gulati. 2009. Perspectives in winter

limnology: closing the annual cycle of freezing lakes. Aquatic Ecology 43:609–616.

Sanders, R. W., K. G. Porter, and S. J. Bennett. 1990. Heterotrophic, autotrophic, and

mixotrophic nanoflagellates: Seasonal abundances and bacterivory in a eutrophic lake.

Limnology and Oceanography 35:1821–1832.

Schindler, D. W. 1977. Evolution of phosphorus limitation in lakes. Science.

Sharma, S., K. Blagrave, J. J. Magnuson, C. M. O'Reilly, S. Oliver, R. D. Batt, M. R. Magee, D.

Straile, G. A. Weyhenmeyer, L. Winslow, and R. I. Woolway. 2019. Widespread loss of

lake ice around the Northern Hemisphere in a warming world. Nature Climate Change

9:227–231.

Sommer, U., R. Adrian, L. De Senerpont Domis, J. J. Elser, U. Gaedke, B. Ibelings, E. Jeppesen,

M. Lürling, J. C. Molinero, W. M. Mooij, E. van Donk, and M. Winder. 2012. Beyond the

Plankton Ecology Group (PEG) Model: Mechanisms Driving Plankton Succession. Annual

Review of Ecology, Evolution, and Systematics 43:429–448.

Sommer, U., Z. M. Gliwicz, W. Lampert, and A. Duncan. 1986. PEG-model of Seasonal

Succession of Planktonic Events in Fresh Waters. Archiv fur Hydrobiologie 106:433–471.

Sommer, U., F. Sommer, B. Santer, E. Zöllner, K. Jürgens, C. Jamieson, M. Boersma, and K.

Gocke. 2003. *Daphnia* versus copepod impact on summer phytoplankton: functional

compensation at both trophic levels. Oecologia 135:639–647.

Søndergaard, M., J. P. Jensen, and E. Jeppesen. 2003. Role of sediment and internal loading of

phosphorus in shallow lakes. Hydrobiologia 506–509:135–145.

Stewart, I. T., D. R. Cayan, and M. D. Dettinger. 2004. Changes in Snowmelt Runoff Timing in

Western North America under a `Business as Usual' Climate Change Scenario. Climatic

Change 62:217–232.

Tamminen, T. 1995. Nitrate and ammonium depletion rates and preferences during a Baltic

spring bloom. Marine Ecology Progress Series 120:123–133.

Tõnno, I., H. Agasild, T. Kõiv, R. Freiberg, P. Nõges, and T. Nõges. 2016. Algal Diet of Small-

Bodied Crustacean Zooplankton in a Cyanobacteria-Dominated Eutrophic Lake. PLOS

ONE 11:e0154526.

Trujillo, E., and N. P. Molotch. 2014. Snowpack regimes of the Western United States. Water

Resources Research 50:5611–5623.

Twiss, M. R., R. M. L. McKay, R. A. Bourbonniere, G. S. Bullerjahn, H. J. Carrick, R. E. H.

Smith, J. G. Winter, N. A. D'souza, P. C. Furey, A. R. Lashaway, M. A. Saxton, and S. W.

Wilhelm. 2012. Diatoms abound in ice-covered Lake Erie: An investigation of offshore

winter limnology in Lake Erie over the period 2007 to 2010. Journal of Great Lakes

Research 38:18–30.

United States Environmental Protection Agency. 2016. Standard Operating Procedure for

Zooplankton Analysis LG403.

USEPA. 1993. Method 365.1, Revision 2.0: Determination of Phosphorus by Semi-Automated

Colorimetry. Cincinnati, Ohio, USA.

Vanderploeg, H. A., S. J. Bolsenga, G. L. Fahnenstiel, J. R. Liebig, and W. S. Gardner. 1992.

Plankton ecology in an ice-covered bay of Lake Michigan: utilization of a winter

phytoplankton bloom by reproducing copepods. Hydrobiologia 243–244:175–183.

Vehmaa, A., and K. Salonen. 2009. Development of phytoplankton in Lake Pääjärvi (Finland)

during under-ice convective mixing period. Aquatic Ecology 43:693–705.

Vermont Department of Environmental Conservation. 2018. Spring Phosphorus Monitoring

Lake Water Quality Data. https://anrweb.vermont.gov/dec/_dec/SpringPhosphorus.aspx.

Virro, T., J. Haberman, M. Haldna, and K. Blank. 2009. Diversity and structure of the winter

rotifer assemblage in a shallow eutrophic northern temperate Lake Võrtsjärv. Aquatic

Ecology 43:755–764.

Watkins, J., L. Rudstam, and K. Holeck. 2011. Length-weight regressions for zooplankton

biomass calculations – A review and a suggestion for standard equations.

Wojciechowska, W., and T. Lenard. 2014. Effect of extremely severe winters on under-ice

phytoplankton development in a mesotrophic lake (Eastern Poland). Oceanological and

Hydrobiological Studies 43:147–153.

Zhou, S., Q. Cai, Y. Xu, and X. Han. 2011. Daily and vertical dynamics of rotifers under the

impact of diatom blooms in the Three Gorges Reservoir, China. Hydrobiologia 675:29–40.

Tables and Figures

Table 1. ANOVA and variance partitioning results for each response variable. N=24 for each test (4 replicates per treatment). Dinoflagellate, desmid, euglenoid, and synurophyte phytoplankton density and biovolume were not analyzed statistically due to low sample size but are included in total phytoplankton calculations. Significant p-values (α=0.05) are bolded with their respective R^2s.

Response variable	R^2_{light}	R^2_{zoop}	p-value$_{light}$	p-value$_{zoop}$
Total phytoplankton density	0.07	0.08	0.2124	0.4303
Chlorophyte density	0.01	0.02	0.7630	0.8258
Diatom density	**0.73**	**0.07**	**3.564*10^{-8}**	**0.0470**
Cyanobacteria density	0.00	0.05	0.7646	0.6303
Cryptomonad density	**0.37**	0.12	**0.001074**	0.1121
Chrysophyte density	**0.76**	0.05	**2.103*10^{-8}**	0.1081
Haptophyte density	0.00	0.00	0.9224	0.9853
Total phytoplankton biovolume	**0.51**	0.09	**6.265*10^{-5}**	0.1143
Chlorophyte biovolume	**0.21**	0.08	**0.0246**	0.3256
Diatom biovolume	**0.62**	0.00	**1.239*10^{-5}**	0.9172
Cyanobacteria biovolume	0.02	0.06	0.4801	0.5605

Cryptomonad biovolume	**0.23**	**0.33**	**0.0044**	**0.0038**
Chrysophyte biovolume	**0.71**	0.05	$2.210*10^{-7}$	0.1618
Haptophyte biovolume	0.00	0.05	0.8303	0.6268
Rotifer density	0.03	0.03	0.4514	0.4105
Rotifer biomass	0.04	**0.33**	0.2768	**0.0034**
Total phosphorus	0.11	0.01	0.1261	0.5462
Soluble reactive phosphorus	**0.73**	**0.15**	$3.230*10^{-10}$	$6.577*10^{-5}$
Total organic carbon	0.00	0.00	0.7834	0.8293
Total nitrogen	0.08	**0.20**	0.15004	**0.02577**
Ammonia	**0.72**	**0.21**	$6.526*10^{-13}$	$4.084*10^{-8}$
Nitrate	0.01	0.00	0.7065	0.8240

Table 2. Phytoplankton genera found in mesocosms.

Taxonomic Group	Genera	Taxonomic Group	Genera
Diatoms	*Asterionella*	**Cyanobacteria**	*Aphanocapsa*
	Diatoma		*Dolichospermum*
	Fragilaria		*Microcystis*
	Navicula		*Pseudanabaena*
	Stephanodiscus		*Woronichinia*
Chlorophytes	*Ankistrodesmus*	**Cryptophytes**	*Chroomonas*
	Chlamydomonas		*Cryptomonas*
	Crucigenia		*Komma*
	Dictyosphaerium	**Dinoflagellates**	*Peridinium*
	Eudorina	**Chrysophytes**	*Chrysococcus*
	Franceia		*Dinobryon*
	Gonium	**Haptophytes**	*Chrysochromulina*
	Kirchneriella	**Synurophytes**	*Mallomonas*
	Micractinium	**Desmids**	*Cosmarium*
	Oocystis		*Staurastrum*
	Scenedesmus	**Euglenoids**	*Euglena*
	Schroederia		
	Selenastrum		
	Tetrabaena		
	Tetraspora		

Fig. 1. Carboy experimental design. A.) Each carboy was suspended in a randomized grid approximately 0.5 m below the ice by a steel cable harness connected to a PVC anchor in the shape of an "X." PVC was placed on loose wood blocks to prevent it from freezing into the ice. B.) Greenhouse shade cloth covers blocked 85% of incoming light to simulate snow cover (photo credit: Hannah Lachance). C.) We crossed two light levels (unshaded and shaded) with three zooplankton levels with four replicates per treatment. Our target for "low" zooplankton was complete removal of large-bodied zooplankton, "medium" zooplankton mimicked ambient zooplankton densities, and "high" zooplankton targeted ten times ambient zooplankton densities (see Results for actual densities).

Fig. 2. Crustacean zooplankton and rotifer density and biomass. "Before" indicates samples taken from barrels at the start of experiments. "Z" indicates zooplankton level. Center lines indicate medians, boxes indicate first and third quartiles of data, and whiskers indicate minimum and maximum of data.

Fig. 3. Total phytoplankton biovolume and density. "Before" indicates samples taken from barrels at the start of experiments. "Z" indicates zooplankton level. Center lines indicate medians, boxes indicate first and third quartiles of data, and whiskers indicate minimum and maximum of data.

Fig. 4. Phytoplankton proportional density and biovolume by taxonomic group. "Before"

indicates samples taken from barrels at the start of experiments. "Z" indicates zooplankton level.

Fig. 5. Results from nMDS analysis on phytoplankton community composition with three axes (stress = 0.15). Each point represents one replicate. "Before" indicates samples taken from barrels at the start of experiments. "Z" indicates zooplankton level.

Fig. 6. Nutrient results from experimental treatments. "Before" indicates samples taken from barrels at the start of experiments. "Z" indicates zooplankton level. Center lines indicate medians, boxes indicate first and third quartiles of data, and whiskers indicate minimum and maximum of data.