TITLE

Active zone compaction in presynaptic homeostatic potentiation

AUTHORS

Achmed Mrestani[1], Philip Kollmannsberger[2], Martin Pauli[1,6], Felix Repp[1,2,6], Robert J. Kittel[1,3,4], Jens Eilers[4], Sören Doose[5], Markus Sauer[5], Anna-Leena Sirén[6], Manfred Heckmann[1*], Mila M. Paul[1*]

[1] Institute for Physiology, Department of Neurophysiology, Julius-Maximilians-University Würzburg, D-97070 Würzburg, Germany

[2] Center for Computational and Theoretical Biology, Julius-Maximilians-University Würzburg, D-97074 Würzburg, Germany

[3] Institute of Biology, Department of Animal Physiology, Leipzig University, D-04103 Leipzig, Germany

[4] Carl-Ludwig-Insitute for Physiology, Leipzig University, D-04103 Leipzig, Germany

[5] Department of Biotechnology and Biophysics, Julius-Maximilians-University Würzburg, D-97074 Würzburg, Germany

[6] Department of Neurosurgery, University Hospital of Würzburg, D-97080 Würzburg, Germany

[*] Correspondence should be adressed to M.H. (heckmann@uni-wuerzburg.de) and M.M.P. (mila.paul@uni-wuerzburg.de)

CLASSIFICATION

major category: Biological Sciences, minor category: Neuroscience

KEYWORDS

active zone, Bruchpilot, compaction, homeostasis, presynaptic plasticity, super-resolution microscopy

AUTHOR CONTRIBUTIONS

A.M., J.E., A.L.S., M.S., M.H. and M.M.P designed experiments. A.M., M.P., M.H. and M.M.P. performed experiments. A.M., P.K., F.R., M.P., R.J.K., J.E., S.D., M.H. and M.M.P analyzed the data. A.M., M.H. and M.M.P. wrote the manuscript with the help

of all co-authors. A.L.S., M.S., M.H. and M.M.P. coordinated the study and provided

funding.

This PDF file includes:

Main Text

Figures 1-5

Supplementary Figures 1-5

Supplementary Table 1

ABSTRACT

Brain function relies on neurotransmission which is stabilized by presynaptic

homeostatic potentiation (PHP). PHP operates on time scales ranging from minute- to

life-long adaptations and likely involves reorganization of presynaptic active zones

(AZs). At *Drosophila melanogaster* neuromuscular junctions, earlier work ascribed AZ

enlargement by incorporating more Bruchpilot (Brp) scaffold protein a central

mechanistic role in PHP.

We used localization microscopy (*direct* stochastic optical reconstruction microscopy,

*d*STORM) and hierarchical density-based spatial clustering of applications with noise

(HDBSCAN) to study AZ plasticity during PHP. We found that both acute,

philanthotoxin (PhTx)-induced and chronic, genetically-induced PHP lead to

compaction of individual AZs without altering Brp copy numbers per AZ. This

compaction even occurs within Brp subclusters of the AZ scaffold which also move

towards AZ centers. Furthermore, lowering imaging resolution revealed how AZ

compaction in PHP translates into apparent increases in AZ area and Brp protein

content as implied earlier. Our results suggest AZ compaction in PHP as an effective

mechanism to raise presynaptic protein density and transmitter release.

SIGNIFICANCE STATEMENT

Homeostatic plasticity stabilizes chemical synaptic transmission in multiple organisms

ranging from insects to humans. Changes in active zones (AZs), membrane

specializations of the presynapse where synaptic vesicles are discharged, are thought

to be crucial in homeostatic adaptations. AZ growth by protein incorporation was

proposed as a core mechanism in presynaptic homeostatic potentiation (PHP).

Localization microscopy of an abundant AZ scaffold protein uncovered that instead of

growing, AZs are compacted in acute and chronic PHP. At lower imaging resolution,

however, AZs appear larger and brighter although protein numbers are not increased.

In summary, our findings suggest AZ compaction as new and effective mechanism to

raise presynaptic protein density and transmitter release in PHP.

INTRODUCTION

Chemical synapses are complex nanomachines that integrate and process information. Synapses are optimized for fast and reliable performance in combination with miniaturization and plasticity (Atwood and Karunanithi, 2002; Chua et al., 2010; Kittel and Heckmann, 2016; Neher and Brose, 2018). An intriguing form of synaptic plasticity is presynaptic homeostatic potentiation (PHP, Turrigiano, 2008; Davis and Müller, 2015). Homeostatic plasticity has been studied at vertebrate synapses (Turrigiano et al., 1998; Murthy et al., 2001; Thiagarajan et al., 2005; Lindskog et al., 2010; Jeans et al., 2017; Delvendahl et al., 2019) as well as at *Drosophila melanogaster* neuromuscular junctions (NMJs, Davis and Goodman, 1998; DiAntonio et al., 1999; Frank et al., 2006; Müller et al., 2012; Younger et al., 2013; Orr et al., 2017; Gratz et al., 2019). While the structural correlates of PHP remain elusive, several mechanisms are conceivable including incorporation, removal, modification and/or rearrangement of key proteins at active zones (AZs), presynaptic membrane specializations where synaptic vesicles are discharged.

The cytomatrix at the active zone (CAZ) consists of an organized set of proteins (Südhof 2012). Large α-helical coiled-coil proteins of the ELKS/CAST family are crucial for the regulation of synaptic transmission (Held et al., 2016; Dong et al., 2018). In *Drosophila,* the ELKS/CAST homolog Bruchpilot (Brp) is essential for synchronous glutamate release and its amount was shown to correlate with release probability and structural synaptic differentiation (Kittel et al., 2006; Ehmann et al., 2014; Peled et al., 2014; Paul et al., 2015; Akbergenova et al., 2018). Brp is a major AZ scaffold protein responsible for clustering of presynaptic calcium channels close to release sites creating a proper molecular environment for fast and precise neurotransmitter release (Kittel et al., 2006; Held and Kaeser, 2018). Whereas the Brp N-term was mapped in membrane-proximity, its C-term localizes about 155 nm above the postsynaptic receptors (Fouquet et al., 2009; Liu et al., 2011) and is important for tethering synaptic vesicles (Hallermann et al., 2010). Remarkably, Brp is distributed heterogeneously at *Drosophila* NMJs forming about 15 subclusters within an AZ (Ehmann et al., 2014). Subclusters with high protein concentration should promote vesicle tethering as shown for Syntaxin (Sieber et al., 2007; Barg et al., 2010; van den Bogaart et al., 2011). Early work at *Drosophila* NMJs showed no evidence for protein synthesis during PHP (Frank et al., 2006). However, confocal and STED microscopy suggested an increase in Brp

amount during acute and chronic PHP (Weyhersmüller et al., 2011; Goel et al., 2017;

Böhme et al., 2019).

Here, we reason that localization microscopy in terms of *direct* stochastic optical

reconstruction microscopy (*d*STORM, Heilemann et al., 2008; van de Linde et al.,

2011; Löschberger et al., 2012) permits precise quantification of Brp in PHP (Ehmann

et al., 2014; Paul et al., 2015). For unbiased data analysis we perform hierarchical

density-based spatial clustering of applications with noise (HDBSCAN, Campello et al.,

2013). We address the following fundamental questions: How robust are localization-

based results relying on density-based clustering algorithms? Does the amount of

protein within individual AZs increase during PHP and/or does the nanoarchitecture of

the AZ scaffold change? Do AZs grow or shrink? Using data simulations and correlative

confocal–*d*STORM imaging we analyze the relation between thresholding- and

localization-based fluorescence quantification of small structures below the diffraction

limit. We discover striking differences of Brp organization during acute,

pharmacologically-induced and chronic, genetically-induced homeostasis and reason

that AZ compaction and not AZ growth is the structural correlate of release

strengthening in PHP.

RESULTS

Analysis of super-resolution data

We imaged type Ib boutons on abdominal muscles 6 and 7 of male *Drosophila melanogaster* 3[rd] instar larvae (Figure 1A) using *d*STORM (Heilemann et al., 2008; van de Linde et al., 2011; Löschberger et al., 2012; Ehmann et al., 2014; Paul et al., 2015) and a highly specific monoclonal antibody Brp[Nc82], that maps to the C-terminal region of Brp (Kittel et al., 2006; Fouquet et al., 2009). Since Brp is an abundant protein and our epitope in Brp covers the spatial extent of individual AZs, we interpret 'Brp area' as 'AZ area'. Density-based clustering algorithms allow to study synaptic nanotopology (Bar-On et al., 2012; Ehmann et al., 2014; Szoboszlay et al., 2017; Rahbek-Clemmensen et al., 2017; Rebola et al., 2019). We used a Python implementation of hierarchical density-based spatial clustering of applications with noise (HDBSCAN), which robustly extracts clusters in data with varying density (Campello et al., 2013; McInnes et al., 2017). To test the algorithm's robustness, we varied its free parameters 'minimum cluster size' and 'minimum samples' and analyzed their influence on the number of detected Brp clusters, the number of Brp localizations and the area per cluster (Figure 1B-E, Supplementary Figure 1). A wide range of parameters delivered robust results and all Brp clusters in the following analyses were extracted with the combination 100 and 25 for minimum cluster size and minimum samples, respectively.

To probe the existence of Brp subclusters in our data we computed H functions (derivative of Ripley's K function) and calculated an averaged curve (Figure 1F). Positive values for H(r) indicate clustering, negative dispersion or edge effects, and maximum positive values roughly correspond to the radius of putative clusters (Kiskowski et al., 2009). The maximum of the curve at 22 nm is in agreement with the dimensions of Brp subclusters reported in earlier work (Ehmann et al., 2014). Subsequently, we performed a second HDBSCAN on individual AZs with adjusted parameters that yielded similar subcluster radii as the H function (Figure 1G, H). In addition to alpha shape generation for area determination, localization-based analysis yields the possibility to define a center of mass (c.o.m.) of an individual subcluster and of the entire AZ and to measure their specific interspaces (Figure 1G). The distance between AZ c.o.m. and subcluster c.o.m. will be referred to as radial distance.

Compaction of the Bruchpilot scaffold in acute and chronic homeostasis

The wasp venom Philanthotoxin (PhTx) induces PHP within minutes and increases

quantal content and release-ready vesicles in our preparation (Frank et al., 2006;

Weyhersmüller et al., 2011; Davis and Müller, 2015). We compared AZs of type Ib

boutons incubated in PhTx (phtx) or DMSO (control, Figure 2A). Brp localization

numbers as a measure for the amount of Brp protein per AZ (Ehmann et al., 2014)

were not changed in phtx, however, AZ area was reduced compared to controls and,

hence, Brp density increased in phtx (Figure 2B, see Supplementary Table 1 for all

data presented here). Thus, PHP induces a rearrangement of Brp without changing

the number of Brp molecules within individual AZs of type Ib boutons.

Previous work discovered a proximo-distal functional and structural gradient with distal

type Ib boutons containing more AZs, more Brp per AZ and releasing more glutamate

than proximal boutons (Peled and Isacoff, 2011; Ehmann et al., 2014; Paul et al.,

2015). To test whether the above described changes in phtx occur in a distinct spatial

pattern within the NMJ we performed a subgroup analysis of AZs in boutons number

1-6 (Paul et al., 2015). Spearman correlation coefficients displayed no correlation

between bouton number and Brp localization density in both groups (Supplementary

Figure 2A). Thus, structural plasticity in acute homeostasis appears to occur

homogeneously within the NMJ. Furthermore, there was neither a change in the total

number of localizations nor the amount of unclustered Brp localizations outside AZs

per bouton (Supplementary Figure 2B).

To clarify whether AZ compaction only appears after an acute homeostatic challenge

or whether it is a general phenomenon in PHP, we used a knockout of the postsynaptic

glutamate receptor subunit GluRIIA as a genetic, 'chronic' PHP model (GluRIIAko,

Petersen et al., 1997; DiAntonio et al., 1999; Frank et al., 2006). We performed

dSTORM imaging and HDBSCAN analysis in GluRIIAko and wildtype larvae as

described before (Figure 2C). Remarkably, we found comparable effects to PhTx

treatment with unchanged Brp localization numbers per AZ, decreased AZ area and

higher Brp density (Figure 2D). Thus, Brp scaffolds are also compacted in chronic PHP.

AZ subcluster compaction

Next, we tested whether AZ reorganization during PHP also takes place within AZ

subclusters (Figure 3A, C). Brp subcluster number per AZ was unchanged in phtx and

GluRIIAko versus controls (Figure 3B, D). While Brp localization numbers in

subclusters were unchanged, subcluster area was decreased (Figure 3B, D). Thus,

mean Brp localization density in subclusters increased in acute and chronic PHP
(Figure 3B, D). We also examined the radial distances between the c.o.m. of an
individual AZ and the c.o.m.s of its subclusters and found that they decreased in phtx
and GluRIIAko compared to controls (Figure 3B, D). In addition, the minimum distance
between subcluster c.o.m.s was decreased in both forms of PHP, whereas the space
between subclusters was only reduced in chronic PHP (Supplementary Figure 5).
Taken together, these data show that Brp is compacted within AZ subclusters in acute
and chronic homeostasis but the number of Brp molecules does not increase.
Next, we quantified the circularity of Brp clusters as the ratio of their Eigenvalues of
the covariance matrix (ratio between 0 and 1 with 1 indicating a perfect circle, i.e.
presumably in top view, see Material and Methods). AZ circularity was similar in phtx
and control, thus, analysis of the entire dataset without orientation selection appears
appropriate. We then investigated the relation between AZ area and circularity.
Interestingly, this uncovered an inverse correlation (Supplementary Figure 2C). We
interpret that low circularity is indicative of AZs in side view or of large Brp spots that
partially arise from merged AZs lying nearby in 2D projection. The latter have been
referred to as double ring structures, grouped CAZ units, or cluster AZs (Kittel et al.,
2006; Ehmann et al., 2014; Akbergenova et al., 2018). Assuming that some structural
parameters depend strongly on AZ orientation, we analyzed AZ area and radial
distance in a subsample of AZs with circularity ≥ 0.5 and more planar orientation
(Supplementary Figure 2D). Again, we found a pronounced decrease in AZ area and
a similar decrease of radial distance in both groups. This indicates that AZ compaction
in PHP appears regardless of AZ orientation.
In earlier studies abdominal muscle 4 which exclusively forms type Ib boutons was
used to analyze structural effects of PhTx on AZs (Goel et al., 2017; Böhme et al.,
2019). Therefore, we imaged PhTx treated AZs on abdominal muscle 4 and applied
the described cluster analysis. This yielded similar results as shown for AZs on
abdominal muscles 6/7 including subcluster parameters (Supplementary Figure 3).
223
Correlative confocal-*d*STORM imaging of AZs
Earlier confocal and STED microscopy reported an increased fluorescence intensity in
PHP and interpreted this as evidence for protein recruitment to the AZ (Weyhersmüller
et al., 2011; Goel et al., 2017; Böhme et al., 2019). Since our results are in sharp
contrast to these findings we wondered whether the difference in molecular density

before and after PHP could bias AZ area quantification either in our localization-based
approach or in intensity-based approaches (confocal and STED microscopy). To test
this, we compared both methods using simulated circular clusters without noise and
subclustering (Supplementary Figure 4). Our results indicate that the area of large
objects with low molecular density (i.e. lower fluorescence intensity) can be grossly
misjudged by pixel- and intensity-based quantification, which could explain the
discrepancy between our and earlier findings. To further examine this experimentally,
we perfomed sequential confocal-*d*STORM imaging of Brp in type Ib boutons (Figure
4A, B). Confocal data were analyzed with an intensity threshold of 100 a. u. (Figure
4Aii). Figure 4C shows enlarged AZs overlaid in confocal and *d*STORM resolution from
Figure 4A and B in green, magenta and blue to mark AZ identity. The examples
demonstrate area overestimation due to high localization density (Figure 4Ci) and
show that on the other hand thresholding of confocal images can lead to complete loss
of signal that is included in *d*STORM quantification (Figure 4Cii). Additionally,
segmentation of closely spaced AZs is clearly superior using HDBSCAN-based
quantification of localization data compared to thresholding-based quantification of
confocal data (Figure 4Ciii). In a next step, we quantified AZs in the complete dataset
after correlative confocal-*d*STORM imaging. Distribution of AZ area was broader in
confocal microscopy than in *d*STORM (Figure 4D). As expected, we obtained positive
correlations between the localization density per AZ measured with *d*STORM and the
mean pixel intensity per AZ measured with confocal (Figure 4E). Interestingly,
localization density correlated stronger with confocally measured than *d*STORM-
measured AZ area (Figure 4F) matching our simulation results (Supplementary Figure
4). Confocal microscopy mostly overestimates AZ area, whereas underestimation only
occurs at low AZ localization density (Figure 4G). This indicates that higher localization
density in *d*STORM leads to an apparent enlargement of AZ area reported by
thresholded confocal microscopy.
256

Homeostatic compaction is translated into an apparent area increase
In contrast to earlier studies (Weyhersmüller et al., 2011; Goel et al., 2017; Böhme et
al., 2019) we found smaller AZ areas but increased localization density following PHP.
To gain a better understanding of this discrepancy we performed simulations of
confocal spatial resolution applying a Gaussian filter of 150 nm SD (Figure 5A, B). 3D
histograms of two example AZs in control and phtx illustrate the applied thresholding

(Figure 5C, compare Figure 4B in Weyhersmüller et al., 2011). In phtx, AZ area is

decreased and localization density increased in original *d*STORM resolution (x-axes,

Figure 5D, E), whereas both AZ area and mean pixel intensity increased in the same

images in simulated confocal resolution (y-axes). We conclude that intensity-based

quantification with confocal resolution may lead to inverse results concerning AZ area

compared to localization-based analysis. In a last step, we simulated STED resolution

as a less dramatic resolution decrease applying a Gaussian filter of 25 nm SD (Figure

5F). AZ area was measured similarly as mentioned above, AZ diameter was estimated

from AZ area and intensity maxima per AZ were detected in selected planar-oriented

AZs with a peak finding algorithm in FIJI after thresholding (Material and Methods;

Böhme et al., 2019). This indicated an increase in AZ area, AZ diameter and number

of intensity maxima per AZ after induction of PHP (Figure 5G), apparent findings

contrary to our localization-based data evaluation. In summary, simulation of lower

resolution gives inverse results for the number of clusters per AZ matching

aforementioned findings.

DISCUSSION

We introduce AZ compaction as a mechanism for presynaptic structural adaptation in acute and chronic PHP. Compaction encompasses reduced AZ area and enhanced molecular density without increasing protein amount. Brp subclusters are likewise compacted and located closer to AZ centers during PHP. Notably, AZ compaction appears as an increase of AZ area in thresholding-based analysis, explaining the apparent contradiction with earlier results.

Protein proximity promotes vesicle traffic

Despite substantial progress in single-molecule based microscopy, resolving the spatial relations between individual subcellular molecules remains challenging (Nahmani et al., 2017). In this study, we reason that dSTORM and a localization-based clustering algorithm are a powerful experimental conjunction to decipher protein dense networks at single presynaptic AZs. The effects reported here are small in absolute terms, but very reproducible under different conditions (Figure 2 and 3, Supplementary Figure 2, 3 and 5). In the light of mechanistic considerations, it is plausible that changes of molecular proximities at AZs in the nanometer range lead to considerable functional consequences (Neher 1998). For example, it is established for the t-SNARE protein Syntaxin that the formation of protein clusters enhances local protein concentration, which in turn may raise the probability of protein interactions (Sieber et al., 2007; van den Bogaart et al., 2011). Furthermore, in hippocampal neurons RIM clusters with diameters of ~80 nm were described to show remarkable reorganization during synaptic plasticity (Tang et al., 2016). Molecular concentrations can be estimated from our data and Brp molecule numbers per AZ reported previously by Ehmann et al., 2014 assuming a distribution of the Brp[Nc82] epitope in a hemisphere. For radii of 185 and 175 nm (Figure 2D, Supplementary Table 1) the hemisphere volumes before and after PHP are ~0.013 and 0.011 µm^3. 170 Brp molecules (Ehmann et al., 2014) yield concentrations of 21 and 25 µM before and after PHP, respectively. Thus, the ~7 % density increase translates into a 19 % concentration increase. Brp subclusters are assumed to contain bundles of ~7 Brp molecules creating a platform for a single synaptic vesicle as described below (Ehmann et al., 2014). For hemispheres with subcluster radii of 22.5 and 21.7 nm (Figure 3D) this yields protein concentrations of 490 and 540 µM before and after PHP. While the above-mentioned estimates for Brp concentrations in the whole AZ are probably reasonably accurate, the estimates for

the concentration in subclusters provide only a lower limit. Here, subcluster radii are most certainly overestimated due to the linkage error introduced by the combination of a primary antibody with the secondary F(ab')$_2$ fragment leading to a 'labeling complex diameter' of 26 nm (Ehmann et al., 2014). Thus, local protein concentrations could be even higher and, likewise, the difference in the two scenarios. This could be clarified in future studies using smaller labeling approaches, e.g. directly fluorophore-conjugated Brp^{Nc82} F(ab')$_2$ fragments. Brp was shown to directly or indirectly tether synaptic vesicles at AZs via its C-term (Hallermann et al., 2010; Scholz et al., 2019). Other coiled-coil proteins like Golgins tether and traffic vesicles at the Golgi network by guidance via multiple vesicle-binding sites (Gillingham and Munro, 2016; Cheung and Pfeffer, 2016). Likewise, Brp could provide a meshwork guiding vesicles via multiple vesicle-interaction sites represented by subclusters. The Brp^{Nc82} epitope is located close to the C-term and an increased protein concentration, achieved by AZ compaction, could enhance the probability of these interactions and facilitate vesicle supply. Taking the ~40.5 nm vesicle diameter from Karunanithi et al., 2002 and adding ~10 nm of protein trimmings (Takamori et al., 2006) resulting in 50-60 nm veritable vesicle extent, the distance between Brp clusters should fit approximately these dimensions. This is the case and since we observe compaction of Brp subclusters and a reduced radial distance this should further enhance trafficking of vesicles over the surface generated by Brp C-terms (Supplementary Figure 5). This structural adaption is in line with evidence for increased vesicle traffic during PHP (Weyhersmüller et al., 2011; Delvendahl et al., 2019).

Vesicle cluster formation depends on liquid-liquid phase separation, achieved by intrinsically disordered regions (IDRs, Milovanovic et al., 2018). Glutamine-rich IDRs were implicated in the formation of coiled-coils and protein agglomerates (Fiumara et al., 2010). The vesicle tethering Brp C-term comprises a stretch of glutamine residues (Hallermann et al., 2010). Vesicle traffic in liquid-liquid phase separation with Brp C-terms could profit from AZ compaction. While the molecular binding partners of the Brp C-term are still unclear, a functional interaction with Complexin was reported (Scholz et al., 2019). We assume that all Brp molecules forming one subcluster create a platform for tethering a single synaptic vesicle. This hypothesis matches reasonably well with EM studies describing on average 12 vesicles tethered to a single CAZ

346 (Böhme et al., 2016) and with reconstructions of EM tomography data (Zhan et al.,
347 2016).

Compaction of the AZ scaffold without recruitment of Brp molecules

Localization data can be used to estimate protein numbers (Ehmann et al., 2014;
Löschberger et al., 2012). In this study, the number of Brp localizations (i.e. Brp
molecules) did not change after induction of homeostasis, which is in line with previous
work ruling out protein synthesis during PHP (Frank et al., 2006). Confocal and STED
microscopy provided data which were interpreted as evidence for protein incorporation
into AZs during acute and chronic homeostasis (Weyhersmüller et al., 2011; Goel et
al., 2017; Böhme et al., 2019). However, our data support an alternative interpretation
(Figure 2B, D, Supplementary Figure 2B). Confocal and STED data were analyzed
using pixel- and thresholding-based approaches that depend on signal intensity
(Weyhersmüller et al., 2011; Goel et al., 2017; Böhme et al., 2019). Here, we show
that large structures with low intensity appear smaller than small structures with higher
intensity (Supplementary Figure 4C). We performed correlative confocal-*d*STORM
imaging and demonstrated that a higher protein density in *d*STORM translates into an
apparent area increase in confocal microscopy (Figure 4F). In confocal microscopy
usually a stack of multiple slices is obtained, whereas in 2D-*d*STORM we sampled only
signal from a single focal plane. However, we found no evidence for signal that is
included in confocal images but is missing in corresponding *d*STORM data (compare
Figure 4Ai and B). We measured that the z-range of boutons in confocal stacks ranges
from 800 nm to 2 μm, roughly corresponding to the axial capture range of our *d*STORM
measurements (Tokunaga et al., 2008). The correlation between localization density
and AZ area in confocal microscopy is corroborated by simulating confocal resolution
(Figure 5D, E).

A molecular concept of homeostatic plasticity

Diverse cellular plasticity mechanisms enable the remarkable capacity of the nervous
system to adapt, refine and increase in strength. Among them, the unique form of
homeostatic plasticity stabilizes brain function on time scales of minutes to decades
proceeding via different molecular pathways (Turrigiano et al., 2008; Younger et al.,
2013; Orr et al., 2017; Ortega et al., 2018; Frank et al., 2020). We here report that
compaction of the AZ scaffold is a crucial hallmark of homeostasis, due to the increase

of protein density and therefore promotion of vesicle traffic. In contrast, disuse hypersensitivity, another form of synaptic plasticity induced by blockade of the AMPA glutamate receptor, was linked to AZ enlargement reversable by application of PhTx (Murthy et al., 2001; Thiagarajan et al., 2005). This illustrates that distinct forms of plasticity can exhibit varying structural expressions. At the *Drosophila* NMJ a proximo-distal gradient in release strength is accompanied by (ultra-)structural adaptations of AZs (Peled and Isacoff, 2011; Ehmann et al., 2014; Paul et al., 2015). To test whether this form of intraneuronal differentiation influences the ability of an AZ to undergo homeostatic plasticity, we analyzed our *d*STORM data according to this aspect. Interestingly, we find that PHP increases Brp density regardless of the bouton position (Supplementary Figure 2A).

We describe compaction on the level of individual AZs (i.e. clusters of Brp epitopes) and of Brp subclusters. However, the molecular mechanism of compaction for release strengthening might be extended to other AZ molecules. Orr et al., 2017 demonstrated that the Semaphorin2b-PlexinB complex regulates PHP via control of presynaptic actin. Whereas the role of actin for synaptic efficacy has been described unambiguously (Cingolani and Goda, 2008), the significance for compaction of AZs remains to be determined. Furthermore, AZ molecules located in membrane-proximity like ENaCs (Younger et al., 2013), presynaptic calcium channels (Turrigiano 2008), RIM and RIM-binding protein should be examined using superresolution microscopy to determine whether compaction is a universal plasticity concept during PHP. Finally, it appears promising to correlate transmission properties of individual AZs with its nanoarchitecture using transgenically expressed GCaMP Ca^{2+} sensors (Akbergenova et al., 2018; Gratz et al., 2019) to link synaptic activity with ultrastructure.

MATERIAL AND METHODS

Fly stocks

Flies were raised on standard cornmeal and molasses medium at 25 °C. *Drosophila melanogaster* male 3rd instar larvae of the following strains were used for experiments: Wildtype: *w1118* (Bloomington *Drosophila* Stock Center). GluRIIA knockout: *DGluRIIAAD9/Df(2L)clh4* (kindly provided by Mathias A. Böhme).

Philanthotoxin treatment and and larval preparation

Philanthotoxin 433 tris (trifluoroacetate) salt (PhTx, P207 Sigma) was dissolved in dimethyl sulfoxide (DMSO) to obtain a stock solution of 4 mM and stored at -20 °C. For each experiment, the respective volume was further diluted with freshly prepared haemolymph-like solution (HL-3, Stewart et al., 1994) to a final PhTx concentration of 20 µM in 0.5 % DMSO. Control experiments were performed with the same DMSO concentration in HL-3. PhTx treatment of semi-intact preparations was performed essentially as described previously (Frank et al., 2006). In brief, larvae were pinned down in calcium-free, ice-cold HL-3 at the anterior and posterior endings, followed by a dorsal incision along the longitudinal axis. Larvae were incubated in 10 µl of 20 µM PhTx in DMSO for 10 minutes at room temperature (22 °C). Following this incubation time, PhTx was replaced by HL-3 and larval preparations were completed, followed by fixation and staining.

Fixation, staining and immunofluorescence

After PhTx treatment and dissection, larvae were fixed with 4 % paraformaldehyde in phosphate buffered saline (PBS) for 10 minutes and blocked for 30 minutes with PBT (PBS containing 0.05 % Triton X-100, Sigma) including 5 % natural goat serum (Dianova). Primary antibodies were added for overnight staining at 4 °C. After two short and three long (20 min each) washing steps with PBT, preparations were incubated with secondary antibodies for 3 hours at room temperature, followed by two short and three long washing steps with PBT. Preparations were kept in PBS at 4 °C until imaging. All data were obtained from NMJs formed on abdominal muscles 6/7 in segments A2 and A3, except data for Supplementary Figure 3, which were obtained from NMJs formed on abdominal muscle 4 in segments A2-A4. Directly compared data (e.g. Figure 2B) were obtained from larvae stained in the same vial and measured in one imaging session.

_d_STORM (_direct_ stochastic optical reconstruction microscopy)

Super-resolution imaging of the specimen was performed essentially as previously reported (Ehmann et al., 2014; Paul et al., 2015). Preparations were incubated with monoclonal antibody (mAb) Brp^{Nc82} (1:100, Antibody Registry ID: AB_2314866, Developmental Studies Hybridoma Bank) and secondary antibody goat α-mouse F(ab')$_2$ fragments labelled with Alexa Fluor647 (1:500, A21237, Thermofisher). Boutons were visualized using Alexa Fluor488 conjugated goat α-horseradish-peroxidase antibody (α-hrp, 1:250, Jackson Immuno Research). After staining, larval preparations were incubated in 100 mM mercaptoethylamine (MEA, Sigma-Aldrich) in a 0.2 M sodium phosphate buffer, pH 7.8 to 7.9, to allow reversible switching of single fluorophores during data acquisiton (van de Linde et al., 2008). The buffer additionally included an oxygen-scavenging system (10% (wt/vol) glucose, 10 U/ml glucose oxidase and 200 U/ml catalase). In all experiments, images were acquired using an inverted microscope (Olympus IX-71, 60x, NA 1.49, oil immersion) equipped with a nosepiece-stage (IX2-NPS, Olympus). 647 nm (F-04306-113, MBP Communications Inc.) and 488 nm (iBEAM-SMART-488-S, Toptica) lasers were used for excitation of Alexa Fluor647 and Alexa Fluor488, respectively. Laser beams were passed through clean-up filters (BrightLine HC 642/10, Semrock and ZET 488/10, Chroma, respectively), combined by a dichroic mirror (LaserMUX BS 473-491R, 1064R, F38-M03, AHF Analysentechnik), and directed onto the probe by an excitation dichroic mirror (HC Quadband BS R405/488/532/635, F73-832, AHF Analysentechnik). The emitted fluorescence was filtered with a quadband-filter (HC-quadband 446/523/600/677, Semrock) and a longpass- (Edge Basic 635, Semrock) or bandpass-filter (Brightline HC 582/75, Semrock) for the red and green channel, respectively, and divided onto two cameras (iXon Ultra DU-897-U, Andor) using a dichroic mirror (HC-BS 640 imaging, Semrock). The green channel was used for visualizing individual presynaptic boutons in normal fluorescence microscopy. For the red channel, image resolution was 126 nm x 126 nm per pixel to obtain super-resolution of Brp. Single fluorophores were localized and high resolution-images were reconstructed with rapi_d_STORM (Heilemann et al., 2008; van de Linde et al., 2011; Wolter et al., 2010; Wolter et al., 2012; www.super-resolution.de). Only fluorescence spots with more than 12000 photons were analyzed and subpixel binning of 10 nm px^{-1} was applied. For visualization of representative _d_STORM measurements (Figure 4B, Figure 5A, B, F), reconstructed images from rapi_d_STORM with 5 nm binning were opened in FIJI

(Schindelin et al., 2012) and where contrast enhanced for clarity. Localization precision was determined with the NeNa algorithm (nearest neighbor based analysis, Endesfelder et al. 2014), implemented in the LAMA software package (LocAlization Microscopy Analyzer; Malkusch and Heilemann, 2016) in 108 dSTORM measurements of Brp and was 6.3 ± 0.6 nm (mean ± SD) in this study. To check for homogenous illumination of the samples we analyzed and compared localization intensity of all dSTORM measurements, (i) for the whole image, and (ii) only for analyzed regions and found no differences in A/D counts between the groups.

Confocal microscopy

For confocal imaging larvae were mounted in PBS and imaged using a commercial confocal laser scanning microscope (Zeiss LSM 700) equipped with an oil-immersion objective (Plan-Apochromat 63x/1.40 Oil M27). Alexa Fluor647 and Alexa Fluor488 were excited with the 639 nm and the 488 nm diode laser lines, repectively. The microscope was controlled via ZEN 12 software (black edition, Zeiss AG). For each z-stack 7-13 slices with 200 nm axial spacing were obtained. The gain of the photomultiplier tubes was adjusted to 700 and 600 V and laser power was set to 1.5 % and 1 % in the red and green channel, respectively, to obtain a good signal with little photobleaching allowing subsequent dSTORM imaging. The pinhole was set to 70.6 µm, corresponding to 1.32 Airy units in the red and 1.73 Airy units in the green channel. Images were recorded in 1024 x 1024 (lines x pixels) format with 16-bit data depth. Given an approximate resolution of 200 nm in the red channel, the pixel size, adjusting the zoom factor to 1.2, was set to about 83 nm for sufficient data sampling. Settings resulted in a pixel dwell time of 3.15 µs.

Data evaluation

Localization microscopy data were analyzed with custom written Python code (https://www.python.org/, language version 3.6) and the web-based Python interface Jupyter (https://jupyter.org/index.html). Localization tables from rapidSTORM were directly loaded and analyzed. Prior to the Python-based analysis the regions of interest (ROI) were masked in the reconstructed, binned images from rapidSTORM using FIJI (1.440, Schindelin et al., 2012). These ROIs corresponded to the terminal 6 boutons according to the α-hrp staining. In earlier work on super-resolved Brp-data, density-based spatial clustering of applications with noise (DBSCAN) was used to identify AZ

subclusters (Ehmann et al., 2014). Given the parameters k and ε, DBSCAN considers

a group of localizations as a cluster if there are at least k residues in a circle with radius

ε around a certain localization (Ester et al., 1996). Here, we used an improved

approach called hierarchical density-based spatial clustering of applications with noise

(HDBSCAN) that, in contrast, extracts the most robust clusters from a cluster hierarchy

over varying ε environments that are least sensible to ε variation, i.e. have the longest

lifetime in the cluster tree (Campello et al., 2013). The algorithm is thus more powerful

for cluster detection in data with variable density. We used the Python implementation

of HDBSCAN (McInnes et al., 2017; https://github.com/scikit-learn-contrib/hdbscan),

which takes 'minimum cluster size' and 'minimum samples' as the main free

parameters, and performed two steps of clustering with different parameters on our

localization data: (i) to identify the AZs, and (ii) to extract the subclusters from the AZs.

Setting minimum samples to smaller values than minimum cluster size allows the

algorithm to be less conservative, i.e. extract clusters that might be smaller than

minimum cluster size but very robust in the cluster hierarchy. We explored optimal

clustering parameters for the detection of active zones and varied minimum cluster

size and minimum samples in a certain range (Figure 1B, Supplementary Figure 1).

Too high values falsely merge adjacent AZs together (Figure 1C) whereas too low

values lead to fragmenting AZs into smaller clusters (Figure 1E). A wide range of

parameters delivered robust results and all Brp clusters in this study were extracted

with the combination 100 and 25 for minimum cluster size and minimum samples. A

second HDBSCAN on the individual AZ clusters was performed with minimum cluster

size 24 and minimum samples 6, and a similar number of subclusters with a

comparable spatial extent compared to the DBSCAN-based subcluster analysis of

Ehmann et al., 2014 was found (compare Figure 1H for subcluster radii and Figure 3B,

D for subcluster areas). For subcluster analysis, the cluster selection method was

changed to 'leaf' clustering, which comprises a tendency to more homogenous clusters

by extracting those that lie on leaf nodes of the cluster tree rather than the most stable

clusters. To quantify cluster area, we computed 2D alpha shapes using CGAL

(Computational Geometry Algorithms Library, https://www.cgal.org) in Python. The

geometrical concept of alpha shapes can be used to calculate the shape and thus the

area of a set of points. Given a finite set of points and an alpha value α, an edge of the

shape is drawn between two points whenever they lie on the boundary of a disk with

radius $\alpha^{1/2}$ that does not contain any other point (Edelsbrunner and Mücke, 1994). To

get the alpha shapes of the AZ clusters and AZ subclusters we choose α-values of 800 nm^2 and 300 nm^2, respectively. The subcluster center of mass (c.o.m.) was calculated as mean of the x- and y-values of all localizations of the subcluster, and the AZ c.o.m. as mean of its subcluster c.o.m.s. To estimate the distance of subcluster c.o.m. from AZ c.o.m. the Euclidean distance of these points was computed. The distance to the nearest subcluster c.o.m. was determined for each individual subcluster c.o.m. and the mean value per AZ was obtained (Supplementary Figure 5A). To get the distance between subclusters the mean subcluster radius per AZ was subtracted from the subcluster center distance (Supplementary Figure 5B). For evaluation of Brp cluster circularity, the ratio of the Eigenvalues of each cluster was computed, where 1 indicates a perfect circle and values < 1 indicate decreasing circularity (Supplementary Figure 2C). Exclusion criteria for outliers in all dSTORM data evaluations of Brp were AZ area < 0.03 μm^2 (Ehmann et al., 2014) and > 0.3 μm^2, absolute localization counts per AZ > 8000 and mean AZ localization density > 60000 localizations per μm^2 (about 3-5-fold median). Additional exclusion criteria for type Ib neuron recordings of Brp were average AZ localization count < 1000 and at the same time average AZ area < 0.095 μm^2 per image, indicative of insufficient data quality. For visualization of the localization-based results (Figure 1A, C-E, G, Figure 2A, C, Figure 3A, C, Figure 4C, Supplementary Figure 3A) scatter plots were created in Python. The H function (Figure 1F) as derivative of Ripley's K function was computed using Python package Astropy (Robitaille et al., 2013) for each individual AZ and for the random Poisson distribution. Curves for display were averaged (mean ± SD). The function was evaluated in nm steps for radii from 0 to 120 nm and without correction for edge effects. Simulated localization data (Supplementary Figure 4) for a given radius r were created in Python using the function numpy.random.uniform() passing the arguments -r and r. Thus, random numbers >= -r and < r were created and used as x- and y-coordinates of random points. Simulation was further constrained by computing the Euclidean distance of the simulated point and the center of the simulated point cloud (x = 0, y = 0) and discarding all points with a distance > r. This gave rise to point clouds consisting of 1500 points with circular distributions in a circle with radius r around the center. For HDBSCAN analysis nine random point clouds per given radius in the same coordinate system were used as input spacing the center points of adjacent point clouds three radii apart. To simulate confocal resolution the point coordinates were binned into 5 nm pixels. Binned images were imported as text images in FIJI and automatically

converted to 32-bit images were raw integrated density is equal to the absolute number

of localizations in a pixel. A Gaussian filter with 150 nm standard deviation (SD) was

applied to blur the images. The maximally possible pixel intensity after blurring was

0.265 arbitrary units (a. u.) corresponding to 1500 points in one pixel before blurring.

To achieve a somewhat realistic gain all images were equally scaled to a maximum of

0.35 a. u. Thresholding was performed in 8-bit with 50 a. u.

Confocal images were processed and evaluated in FIJI. Z-stacks were maximum

projected and Brp spots segmented with a pixel intensity threshold of 100 a. u. in 8-bit

using the 'Analyze Particles' function (Figure 4Aii). Mean pixel intensity of resulting

masks was measured in the original images. Corresponding Brp spots, extracted with

100 a. u. threshold, and Brp localization clusters were assigned manually. Only

clusters that were clearly distinguishable in both confocal and *d*STORM without

confluence to neighboring signal were used. Bouton height ranged from 0.8 to 2 µm in

confocal imaging. To ensure comparability to earlier quantification approaches we

created maximum projections of confocal data. The capture range of our *d*STORM

measurements is at least 1 µm, but probably even higher (Tokunaga et al., 2008). To

minimize bias arising from analysis of different focal planes we inspected all correlative

images and found no evidence for signal that was included in confocal stacks but

missing in *d*STORM images (compare Figure 4Ai and B). Additionally, only AZs were

correlated that could be clearly assigned in both imaging techniques.

To simulate confocal resolution (Figure 5A, B) localization tables from rapi*d*STORM

with 5 nm binning were converted to FIJI-readable density matrices where raw

integrated density corresponds to the localization count in a pixel. Images were

contrast enhanced to obtain 0.1 % saturated pixels, a Gaussian filter with 150 nm SD

was applied and the images were scaled to a pixel size of 80 nm. The same images

as for the localization-based analysis were included and only pixels in the same ROIs

as described above were used for subsequent processing and analysis. Images were

converted to 8-bit and the gain was adjusted by scaling all pixels in both images to the

same value by setting the brightest of all analyzed pixels in all images to the maximum

value of 255 a. u.. Following a standard protocol for the quantification of confocal data

of the *Drosophila* NMJ (Schmid et al., 2008; Weyhersmüller et al., 2011) a minimal

threshold of 50 a. u. was applied, and the 'Analyze Particles' function in FIJI was used

to create individual masks from the thresholded images. The resulting masks were

applied to the non-thresholded simulated confocal images to quantify AZ area as well

as the mean intensity of AZs. STED resolution in Figure 5F was simulated similarly, but a Gaussian filter with 25 nm SD was applied to the density matrices. The gain of all images was adjusted by setting 175 a. u. to the maximum. AZ area and mean pixel intensity per AZ were measured with a threshold of 50 a. u. (Weyhersmüller et al., 2011) and only Brp spots with an area > 0.03 μm^2 were analyzed. The diameter of Brp spots was calculated under the assumption of a circular area. To quantify intensity maxima per AZ planar-oriented, ring-like AZs were selected. Selection was performed blinded with respect to the experimental groups. Single AZs were cut out from the whole images. Thresholding was performed with 100 a. u. and all pixels below the threshold were removed. The maxima were extracted using the 'Find Maxima' function in FIJI (Prominence = 0) and quantified per AZ.

Statistics

Statistical analyses were performed with Sigma Plot 13 (Systat Software). Shapiro-Wilk was used to test normality. If data were not normally distributed, we used the non-parametric Mann-Whitney rank sum test for statistical analysis and reported data as median (25^{th}-75^{th} percentile). If data were normally distributed they were reported as mean ± SD unless indicated otherwise (Figure 1F). Asterisks indicate the significance level (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$) and n denotes sample number. In box plots, horizontal lines represent median, boxes quartiles and whiskers 10^{th} and 90^{th} percentiles. Scatter plots show individual data points unless indicated otherwise. Bin counts in histograms were normalized to the total number of observed events which was set to 1. Color codes in contour plots (Figure 1B, Supplementary Figure 1) were logarithmically scaled. Linear regression curves were fitted and Spearman correlation coefficient r and statistical significance p of correlations were evaluated in Sigma Plot. All plots were produced with Sigma Plot software and figures assembled using Adobe Illustrator (Adobe, 2015.1.1 release). Supplementary Table 1 contains all numerical values not stated in text and figure legends including p-values and samples sizes.

Code and data availability

The authors declare that custom written Python code and all data sets supporting the findings of this work are available from the corresponding authors.

ACKNOWLEDGEMENTS

645 This work was supported by grants from the German Research Foundation (DFG)

646 TRR166/B06 to A.L.S. and M.H., TRR166/B02 to S.D., TRR166/B04 to M.S.,

647 KI1460/5-1 and KI1460/4-1 to R.J.K. and from the IZKF Würzburg to A.L.S. and M.H.

648 (N229). The authors thank T. Langenhan for scientific discussions and M. Oppmann

649 and F. Köhler for technical assistance. The authors declare no conflict of interest.

REFERENCES

1. H. L. Atwood, S. Karunanithi, Diversification of synaptic strength: presynaptic elements. *Nat. Rev. Neurosci.* **3**, 497–516 (2002).

2. J. J. Chua, S. Kindler, J. Boyken, R. Jahn, The architecture of an excitatory synapse. *J. Cell Sci.* **123**, 819–823 (2010).

3. R. J. Kittel, M. Heckmann, Synaptic vesicle proteins and active zone plasticity. *Front. Synaptic Neurosci.* **8,** 8 (2016).

4. E. Neher, N. Brose, Dynamically primed synaptic vesicle states: Key to understand synaptic short-term plasticity. *Neuron* **100**, 1283–1291 (2018).

5. G. G. Turrigiano, The self-tuning neuron: Synaptic scaling of excitatory synapses. *Cell* **135**, 422–435 (2008).

6. G. W. Davis, M. Müller, Homeostatic control of presynaptic neurotransmitter release. *Annu. Rev. Physiol.* **77**, 251–270 (2015).

7. G. G. Turrigiano, K. R. Leslie, N. S. Desai, L. C. Rutherford, S. B. Nelson, Activity-dependent scaling of quantal amplitude in neocortical neurons. *Nature* **391**, 892–896 (1998).

8. V. N. Murthy, T. Schikorski, C. F. Stevens, Y. Zhu, Inactivity produces increases in neurotransmitter release and synapse size. *Neuron* **32**, 673–682 (2001).

9. T. C. Thiagarajan, M. Lindskog, R. W. Tsien, Adaptation to synaptic inactivity in hippocampal neurons. *Neuron* **47**, 725–737 (2005).

10. M. Lindskog *et al.*, Postsynaptic GluA1 enables acute retrograde enhancement of presynaptic function to coordinate adaptation to synaptic inactivity. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 21806–21811 (2010).

11. A. F. Jeans, F. C. van Heusden, B. Al-Mubarak, Z. Padamsey, N. J. Emptage, Homeostatic presynaptic plasticity is specifically regulated by P/Q-type Ca^{2+} channels at mammalian hippocampal synapses. *Cell Rep.* **21**, 341–350 (2017).

12. I. Delvendahl, K. Kita, M. Müller, Rapid and sustained homeostatic control of presynaptic exocytosis at a central synapse. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 23783–23789 (2019).

13. G. W. Davis, C. S. Goodman, Synapse-specific control of synaptic efficacy at the terminals of a single neuron. *Nature* **392**, 82–86 (1998).

14. A. DiAntonio, S. A. Petersen, M. Heckmann, C. S. Goodman, Glutamate receptor expression regulates quantal size and quantal content at the Drosophila neuromuscular junction. *J. Neurosci.* **19**, 3023–3032 (1999).

15. C. A. Frank, M. J. Kennedy, C. P. Goold, K. W. Marek, G. W. Davis, Mechanisms underlying the rapid induction and sustained expression of synaptic homeostasis. *Neuron* **52**, 663–677 (2006).

16. M. Müller, K. S. Y. Liu, S. J. Sigrist, G. W. Davis, RIM controls homeostatic plasticity through modulation of the readily-releasable vesicle pool. *J. Neurosci.* **32**, 16574–16585 (2012).

17. M. A. Younger, M. Müller, A. Tong, E. C. Pym, G. W. Davis, A presynaptic ENaC channel drives homeostatic plasticity. *Neuron* **79**, 1183–1196 (2013).

18. B. O. Orr, R. D. Fetter, G. W. Davis, Retrograde semaphorin-plexin signalling drives homeostatic synaptic plasticity. *Nature* **550**, 109–113 (2017).

19. S. J. Gratz *et al.*, Endogenous tagging reveals differential regulation of Ca^{2+} channels at single AZs during presynaptic homeostatic potentiation and depression. *J. Neurosci.* **39**, 2416–2429 (2019).

20. T. C. Südhof, The presynaptic active zone. *Neuron* **75**, 11–25 (2012).

21. R. G. Held, C. Liu, P. S. Kaeser, ELKS controls the pool of readily releasable vesicles at excitatory synapses through its N-terminal coiled-coil domains. *Elife* **5**, e14862 (2016).

22. W. Dong *et al.*, CAST/ELKS proteins control voltage-gated Ca^{2+} channel density and synaptic release probability at a mammalian central synapse. *Cell Rep.* **24**, 284–293 (2018).

23. R. J. Kittel *et al.*, Bruchpilot promotes active zone assembly, Ca^{2+} channel clustering, and vesicle release. *Science* **312**, 1051–1054 (2006).

24. N. Ehmann *et al.*, Quantitative super-resolution imaging of Bruchpilot distinguishes active zone states. *Nat. Commun.* **5**, 4650 (2014).

25. E. S. Peled, Z. L. Newman, E. Y. Isacoff, Evoked and spontaneous transmission favored by distinct sets of synapses. *Curr. Biol.* **24**, 484–493 (2014).

26. M. M. Paul *et al.*, Bruchpilot and Synaptotagmin collaborate to drive rapid glutamate release and active zone differentiation. *Front. Cell. Neurosci.* **9**, 29 (2015).

27. Y. Akbergenova, K. L. Cunningham, Y. V. Zhang, S. Weiss, J. T. Littleton, Characterization of developmental and molecular factors underlying release heterogeneity at Drosophila synapses. *Elife* **7**, e38268 (2018).

28. R. G. Held, P. S. Kaeser, ELKS active zone proteins as multitasking scaffolds for secretion. *Open Biol.* **8**, 170258 (2018).

29. W. Fouquet *et al.*, Maturation of active zone assembly by Drosophila Bruchpilot. *J. Cell Biol.* **186**, 129–145 (2009).

30. K. S. Liu *et al.*, RIM-binding protein, a central part of the active zone, is essential for neurotransmitter release. *Science* **334**, 1565–1569 (2011).

31. S. Hallermann *et al.*, Naked dense bodies provoke depression. *J. Neurosci.* **30**, 14340–14345 (2010).

32. J. J. Sieber *et al.*, Anatomy and dynamics of a supramolecular membrane protein cluster. *Science* **317**, 1072–1076 (2007).

33. S. Barg, M. K. Knowles, X. Chen, M. Midorikawa, W. Almers, Syntaxin clusters assemble reversibly at sites of secretory granules in live cells. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 20804–20809 (2010).

34. G. van den Bogaart *et al.*, Membrane protein sequestering by ionic protein-lipid interactions. *Nature* **479**, 552–555 (2011).

35. A. Weyhersmüller, S. Hallermann, N. Wagner, J. Eilers, Rapid active zone remodeling during synaptic plasticity. *J. Neurosci.* **31**, 6041–6052 (2011).

36. P. Goel, X. Li, D. Dickman, Disparate postsynaptic induction mechanisms ultimately converge to drive the retrograde enhancement of presynaptic efficacy. *Cell Rep.* **21**, 2339–2347 (2017).

37. M. A. Böhme *et al.*, Rapid active zone remodeling consolidates presynaptic potentiation. *Nat. Commun.* **10**, 1085 (2019).

38. M. Heilemann *et al.*, Subdiffraction-resolution fluorescence imaging with conventional fluorescent probes. *Angew. Chem. Int. Ed. Engl.* **47**, 6172–6176 (2008).

39. S. van de Linde *et al.*, Direct stochastic optical reconstruction microscopy with standard fluorescent probes. *Nat. Protoc.* **6**, 991–1009 (2011).

40. A. Löschberger *et al.*, Super-resolution imaging visualizes the eightfold symmetry of gp210 proteins around the nuclear pore complex and resolves the central channel with nanometer resolution. *J. Cell Sci.* **125**, 570–575 (2012).

41. R. J. G. B. Campello, D. Moulavi, J. Sander, "Density-based clustering based on hierarchical density estimates" in *Proceedings of the 17th Pacific-Asia*

Conference on Knowledge Discovery and Data Mining, J. Pei, V. S. Tseng, L. Cao, H. Motoda, G. Xu, Eds. (Springer, 2013), pp. 160–172.

42. D. Bar-On *et al.*, Super-resolution imaging reveals the internal architecture of nano-sized syntaxin clusters. *J. Biol. Chem.* **287**, 27158–27167 (2012).

43. M. Szoboszlay, T. Kirizs, Z. Nusser, Objective quantification of nanoscale protein distributions. *Sci. Rep.* **7**, 15240 (2017).

44. T. Rahbek-Clemmensen *et al.*, Super-resolution microscopy reveals functional organization of dopamine transporters into cholesterol and neuronal activity-dependent nanodomains. *Nat. Commun.* **8**, 740 (2017).

45. N. Rebola *et al.*, Distinct nanoscale calcium channel and synaptic vesicle topographies contribute to the diversity of synaptic function. *Neuron* **104**, 693–710 (2019).

46. L. McInnes, J. Healy, S. Astels, hdbscan: Hierarchical density based clustering. *Journal of Open Source Software* **2**, 205 (2017).

47. M. A. Kiskowski, J. F. Hancock, A. K. Kenworthy, On the use of Ripley's K-function and its derivatives to analyze domain size. *Biophys. J.* **97**, 1095–1103 (2009).

48. E. S. Peled, E. Y. Isacoff, Optical quantal analysis of synaptic transmission in wild-type and rab3-mutant Drosophila motor axons. *Nat. Neurosci.* **14,** 519–526 (2011).

49. S. A. Petersen, R. D. Fetter, J. N. Noordermeer, C. S. Goodman, A. DiAntonio, Genetic analysis of glutamate receptors in Drosophila reveals a retrograde signal regulating presynaptic transmitter release. *Neuron* **19**, 1237–1248 (1997).

50. M. Nahmani, C. Lanahan, D. DeRosier, G. G. Turrigiano, High-numerical-aperture cryogenic light microscopy for increased precision of superresolution reconstructions. *Proc. Natl. Acad. Sci. U.S.A.* **114**, 3832–3836 (2017).

51. E. Neher, Vesicle pools and Ca^{2+} microdomains: New tools for understanding their roles in neurotransmitter release. *Neuron* **20**, 389–399 (1998).

52. A. H. Tang *et al.*, A trans-synaptic nanocolumn aligns neurotransmitter release to receptors. *Nature* **536**, 210–214 (2016).

53. N. Scholz *et al.*, Complexin cooperates with Bruchpilot to tether synaptic vesicles to the active zone cytomatrix. *J. Cell Biol.* **218**, 1011–1026 (2019).

54. A. K. Gillingham, S. Munro, Finding the Golgi: Golgin coiled-coil proteins show the way. *Trends Cell. Biol.* **26**, 399–408 (2016).

55. P. Y. Cheung, S. R. Pfeffer, Transport vesicle tethering at the trans Golgi network: Coiled coil proteins in action. *Front. Cell. Dev. Biol.* **4**, 18 (2016).

56. S. Karunanithi, L. Marin, K. Wong, H. L. Atwood, Quantal size and variation determined by vesicle size in normal and mutant Drosophila glutamatergic synapses. *J. Neurosci.* **22**, 10267–10276 (2002).

57. S. Takamori *et al.*, Molecular anatomy of a trafficking organelle. *Cell* **127**, 831–846 (2006).

58. D. Milovanovic, Y. Wu, X. Bian, P. De Camilli, A liquid phase of synapsin and lipid vesicles. *Science* **361**, 604–607 (2018).

59. F. Fiumara, L. Fioriti, E. R. Kandel, W. A. Hendrickson, Essential role of coiled coils for aggregation and activity of Q/N-rich prions and PolyQ proteins. *Cell* **143**, 1121–1135 (2010).

60. M. A. Böhme *et al.*, Active zone scaffolds differentially accumulate Unc13 isoforms to tune Ca^{2+} channel-vesicle coupling. *Nat. Neurosci.* **19**, 1311–1320 (2016).

61. H. Zhan, J. Bruckner, Z. Zhang, K. O'Connor-Giles, Three-dimensional imaging
of Drosophila motor synapses reveals ultrastructural organizational patterns. *J.*
Neurogenet. **30**, 237–246 (2016).
62. M. Tokunaga, N. Imamoto, K. Sakata-Sogawa, Highly inclined thin illumination
enables clear single-molecule imaging in cells. *Nat. Methods* **5**, 159–161 (2008).
63. J. M. Ortega, Ö. Genç, G. W. Davis, Molecular mechanisms that stabilize short
term synaptic plasticity during presynaptic homeostatic plasticity. *Elife* **7**,
e40385 (2018).
64. C. A. Frank, T. D. James, M. Müller, Homeostatic control of Drosophila
neuromuscular junction function. *Synapse* **74**, e22133 (2020).
65. L. A. Cingolani, Y. Goda, Actin in action: the interplay between the actin
cytoskeleton and synaptic efficacy. *Nat. Rev. Neurosci.* **9**, 344–356 (2008).
66. B. A. Stewart, H. L. Atwood, J. J. Renger, J. Wang, C. F. Wu, Improved stability
of Drosophila larval neuromuscular preparations in haemolymph-like
physiological solutions. *J. Comp. Physiol. A.* **175**, 179–191 (1994).
67. S. van de Linde, M. Sauer, M. Heilemann, Subdiffraction-resolution
fluorescence imaging of proteins in the mitochondrial inner membrane with
photoswitchable fluorophores. *J. Struct. Biol.* **164**, 250–254 (2008).
68. S. Wolter *et al.*, Real-time computation of subdiffraction-resolution fluorescence
images. *J. Microsc.* **237**, 12–22 (2010).
69. S. Wolter *et al.*, rapi*d*STORM: accurate, fast open-source software for
localization microscopy. *Nat. Methods* **9**, 1040–1041 (2012).
70. J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis.
Nat. Methods **9**, 676–682 (2012).
71. U. Endesfelder, S. Malkusch, F. Fricke, M. Heilemann, A simple method to
estimate the average localization precision of a single-molecule localization
microscopy experiment. *Histochem. Cell Biol.* **141**, 629–638 (2014).
72. S. Malkusch, M. Heilemann, Extracting quantitative information from single-
molecule super-resolution imaging data with LAMA - LocAlization Microscopy
Analyzer. *Sci. Rep.* **6**, 34486 (2016).
73. M. Ester, H.-P. Kriegel, J. Sander, X. Xu, "A density-based algorithm for
discovering clusters in large spatial databases with noise" in *KDD'96:*
Proceedings of the Second International Conference on Knowledge Discovery
and Data Mining, E. Simoudis, J. Han, U. Fayyad, Eds. (AAAI Press, 1996), pp.
226–231.
74. H. Edelsbrunner, E. P. Mücke, Three-dimensional alpha shapes. *ACM*
Transactions on Graphics **13**, 43–72 (1994).
75. T. P. Robitaille *et al.*, Astropy: A community Python package for astronomy.
Astronomy and Astrophysics **558**, A33 (2013).
76. A. Schmid *et al.*, Activity-dependent site-specific changes of glutamate receptor
composition in vivo. *Nat. Neurosci.* **11**, 659–666 (2008).
841

Figure 1. Cluster analysis of presynaptic Brp localizations.

(A) Scatter plot of 15470 localizations from *d*STORM of a representative type Ib bouton

of a wildtype *Drosophila* NMJ on abdominal muscles 6/7 stained with Brp[Nc82] antibody

labelled with Alexa Fluor647 conjugated F(ab')$_2$ fragments. Arrow indicates an

individual cluster of Brp localizations, defined as an AZ throughout the manuscript.

(B) Contour plot of the number of detected AZs in the data from (A) depending on the

HDBSCAN clustering parameters 'minimum samples' and 'minimum cluster size'.

Parameter combinations for clustering in (C-E) are indicated.

(C-E) Data from (A) after HDBSCAN analysis. Colors indicate cluster identity,

unclustered localizations are displayed in black. Parameter combinations for minimum

cluster size and minimum samples were 2000 and 600 (C), 100 and 25 (D) and 15 and

2 (E). Note that multiple AZs are merged into two large clusters in (C) and that

individual AZs are split into multiple small clusters in (E). Minimum cluster size 100 and

minimum samples 25 are used for AZ detection throughout this manuscript. Asterisk

in (D) highlights the AZ cluster shown in (G).

(F) Averaged H function (grey, mean ± SD, derivative of Ripley's K function) from n = 568 AZs obtained from 13 NMJs from 6 wildtype animals. The maximum of the curve indicates a mean subcluster (sc.) radius of 22 nm. Dashed black line indicates the prediction for a random Poisson distribution.

(G) HDBSCAN for subcluster detection applied to an individual AZ (asterisk in D). Alpha shapes used for AZ area quantification are indicated by black lines. Left: Colored Brp subclusters surrounded by colored lines indicating alpha shapes. Unclustered Brp localizations are shown as grey dots. Right: The centers of mass (c.o.m.) of this AZ (cross) and of each subcluster (colored dots) are indicated. A dashed line shows the Euclidean distance between the AZ c.o.m and an individual sc. c.o.m., referred to as 'radial distance'.

(H) Histogram of mean Brp subcluster radius per AZ assuming a circular area, median (25^{th}-75^{th} percentile): 23 (21-26) nm) for AZs in (F). Scale bars 1 μm in (A) and 100 nm in (G).

Figure 2. Acute and chronic homeostatic potentiation decrease AZ area and increase Brp density.

(A) Scatter plots of type Ib AZs from a control (ctrl, left) and a Philanthotoxin treated animal (phtx, right) for induction of acute presynaptic homeostatic potentiation (PHP). Localizations outside alpha shape are not shown for clarity.

(B) Number of Brp localizations per AZ (p = 0.236), AZ area (p = 0.009) and Brp localization density (p = 0.009) for ctrl (grey, n = 568 AZs from 13 NMJs from 6 animals) and phtx larvae (red, n = 792 AZs from 14 NMJs from 5 animals) shown as box plots where horizontal lines represent median, boxes quartiles and whiskers 10th and 90th percentiles. Asterisks indicate significance level (* p < 0.05, ** p < 0.01, *** p < 0.001).

(C) Scatter plots of type Ib AZs from a control animal (ctrl, left) and a GluRIIA receptor null mutant (GluRIIAko, right) representing chronic PHP. Localizations outside alpha shape are deleted for clarity.

(D) Number of Brp localizations per AZ (p = 0.391), AZ area (p < 0.001) and Brp localization density (p < 0.001) for ctrl (grey, n = 872 AZs from 19 NMJs from 6 animals) and GluRIIAko larvae (blue, n = 1020 AZs from 19 NMJs from 6 animals). Scale bars 100 nm in (A) and (C).

Figure 3. Homeostatic potentiation compacts AZ subclusters.

(A) Scatter plots of type Ib AZs from a ctrl and a phtx animal. Brp subclusters are colored individually, unclustered localizations are displayed as grey dots. Localizations outside alpha shape are deleted for clarity.

(B) Number of subclusters per AZ (p = 0.084), sc. area (p = 0.025), sc. localization density (p = 0.022) and radial distance between the AZ c.o.m. and the c.o.m. of individual subclusters (p = 0.003) for ctrl (grey, n = 568 AZs from 13 NMJs from 6 animals) and phtx larvae (red, n = 792 AZs from 14 NMJs from 5 animals).

(C) Scatter plots of type Ib AZs from ctrl and GluRIIAko as shown in (A).

(D) Number of subclusters per AZ (p = 0.118), sc. area (p < 0.001), sc. localization density (p < 0.001) and radial distance between the AZ c.o.m. and the c.o.m. of individual subclusters (p = 0.002) for ctrl (grey, n = 872 AZs from 19 NMJs from 6 animals) and GluRIIAko larvae (blue, n = 1020 AZs from 19 NMJs from 6 animals). Scale bars 100 nm in (A) and (C).

Figure 4. Correlative confocal-*d*STORM microscopy links increased localization density to increased confocal intensity and AZ area.

(A) Confocal image of a type Ib bouton. (i) unprocessed original image and (ii) after applying a 100 a. u. threshold to the original image. Symbols mark corresponding regions in (B) enlarged in (C).

(B) *d*STORM image of the same bouton as shown in (A). Boxes correspond to enlarged regions in (C).

(C) Enlarged confocal AZs thresholded at 100 a. u. in grey corresponding to boxed regions in (B) and overlaid scatter plots of Brp localizations from *d*STORM (green, magenta, blue for cluster identity). Localization clusters at the edges are not displayed. Note that the green and magenta clusters in (Ci) are not separated in confocal microscopy after thresholding, that the magenta cluster in (Cii) is lost after confocal

thresholding, and that the two clusters in (Ciii) are not separated by confocal microscopy.

(D) AZ area in confocal (grey) and *d*STORM imaging (black, for D-F n = 343 corresponding AZs from 12 NMJs from 7 animals).

(E) Spearman correlation coefficient shows a positive correlation between the normalized mean pixel intensity in confocal imaging and *d*STORM localization density.

(F) Spearman correlation coefficients show a stronger correlation between AZ area in confocal imaging and localization density (grey) compared to AZ area measured with *d*STORM and localization density (black).

(G) Ratio of confocal and *d*STORM area of corresponding AZs plotted against their localization density. Values below 1 (dashed black line) indicate an underestimation of confocal area quantification with respect to *d*STORM, whereas values larger than 1 indicate overestimation. Scale bars 1 µm in (A) and (B), 330 nm in (C).

Figure 5. Compaction is reported as increased area in confocal simulation and

as higher AZ diameter and more clusters in STED simulation.

(A) Sections of *d*STORM images in ctrl and phtx (5 nm binning). Boxes highlight

enlarged regions in (B).

(B) Enlarged regions from (A) with *d*STORM resolution (i / iii) and simulated confocal

resolution (ii / iv, 80 nm pixels, 150 nm Gaussian blur). Boxes highlight selected AZs

in (C).

(C) 3D bar plots of mean pixel intensity in confocal resolution of boxed AZs in (B). Grey

plane indicates thresholding level of 50 a. u. which was used for quantification in (D,

E).

(D) AZ area (median (25th-75th percentile)) quantified as shown in (C), i.e. by assuming

confocal resolution, plotted against AZ area from original *d*STORM analysis (data from

Figure 2B) in ctrl (grey, n = 345 AZs from 13 NMJs from 6 animals in confocal resolution

for (D, E)) and phtx (red, n = 457 AZs from 14 NMJs from 5 animals in confocal

resolution for (D, E), p = 0.026). Note the inverse, significant results for AZ area

change.

(E) Mean pixel intensity in simulated 'confocal' resolution (right, p = 0.016) plotted

against dSTORM localization density (data from Figure 2B) in ctrl and phtx. Also note

the inverse, significant results for this analysis.

(F) Example type Ib AZs of ctrl (upper panels) and phtx (lower panels) in dSTORM

resolution (left) and simulated STED resolution (25 nm Gaussian blur, right). White

crosses indicate maxima detected with FIJI peak finding algorithm.

(G) AZ area and AZ diameter, estimated under the assumption of a circular area,

obtained from simulated STED analysis in ctrl (n = 591 AZs from 13 NMJs from 6

animals) and phtx (n = 820 AZs from 14 NMJs from 5 animals, p = 0.044 and 0.044)

and number of intensity peaks, i.e. Brp modules, per AZ in ctrl (n = 122 AZs from 12

NMJs from 6 animals) and phtx (n = 186 AZs from 14 NMJs from 5 animals, p = 0.002).

Scale bars 4 µm in (A), 1.2 µm in (B) and 100 nm in (F).

SUPPLEMENTARY MATERIAL

Supplementary Figure 1. Exploring HDBSCAN parameters. *Related to Figure 1.*
Contour plots of the mean number of localizations per AZ (A) and the mean AZ area
(B) for the data shown in Figure 1A depending on the HDBSCAN clustering parameters
'minimum samples' and 'minimum cluster size'. The parameter combination used for
analysis throughout this study is indicated (minimum cluster size 100, minimum
samples 25).

Supplementary Figure 2. Spatial differentiation, bouton analysis and circularity.

Related to Figures 2 and 3.

(A) Brp localization density shown as boxplots for individual boutons number 1-6 from the end of the bouton string. Spearman correlation coefficients show no significant correlation between AZ localization density and bouton position in ctrl (grey, n = 240, 140, 102, 71, 54, 19 AZs for boutons 1-6) and phtx (red, n = 251, 134, 146, 117, 78, 51 AZs for boutons 1-6, respectively).

(B) Number of all Brp localizations per bouton and number of unclustered (unclust.) localizations per bouton in ctrl (n = 13 distal Ib boutons from 6 animals) and phtx (n = 14 distal Ib boutons from 5 animals, p = 0.577 and 0.903).

(C) Scatter plots illustrating correlation of AZ circularity and area in ctrl and phtx and linear regression lines. Spearman coefficients r and p-values show negative correlations in both groups.

(D) AZs with circularity ≥ 0.5 (indicative for an AZ in top view) were used to compare AZ organization in ctrl and phtx. Box plots of AZ area and of the radial distance between the AZ c.o.m. and the c.o.m. of individual subclusters (n = 442 AZs from 13 NMJs from 6 animals and n = 621 AZs from 14 NMJs from 5 animals in ctrl and phtx, respectively, p = 0.002 and < 0.001).

Supplementary Figure 3. AZ nanoarchitecture in type Ib boutons from NMJs on abdominal muscle 4 of *Drosophila* 3[rd] instar larvae. *Related to Figures 2 and 3.*

(A) Scatter plots of a ctrl and a phtx type Ib AZ on muscle 4.

(B, C) AZ area (B, p = 0.018) and Brp localization density (C, p < 0.001) in ctrl (grey, n = 1183 AZs from 20 NMJs from 6 animals for B-G) and phtx (red, n = 1445 AZs from 22 NMJs from 8 animals for B-G).

(D-G) Subcluster numbers per individual type Ib AZ (D, p = 0.803), subcluster area (E, p = 0.020), subcluster localization density (F, p = 0.003) and radial distance between the AZ c.o.m. and the c.o.m. of individual subclusters (G, p = 0.022) in both groups.

Scale bar 200 nm in (A).

Supplementary Figure 4. Limits of area quantification.

(A) Simulation of 1500 random localizations (blue, compare ctrl value in Figure 2B, D) randomly distributed in circular areas with radius (i) 400 nm and (ii) 300 nm. Black lines indicate alpha shapes for area quantification. Holes in Ai exemplify the local failure of alpha shape-based area quantification.

(B) Radius of simulated localization clusters (magenta, mean ± SD, n = 9 simulated clusters per given radius) estimated from alpha shape area under the assumption of a circular area and line for real radius (grey, equal to given radius) plotted against the given radius for simulation. Symbols (i, ii) correspond to the given radii used in (A). This revealed a robust area quantification and allowed us to define a critical density of about 2700 localizations per µm^2 above which the estimated radius was always at least 95 % of the given radius. The first given radius below the critical density was 430 nm which exceeds the observed maximal AZ radius. We conclude that localization density is not limiting for area quantification given the high densities of AZs and of AZ subclusters in our data.

(C) Data from (A) after binning in 5 nm pixels, application of a Gaussian filter of 150 nm SD and intensity scaling to impose confocal resolution. White lines indicate a 50 a. u. threshold used for area quantification (compare standard procedures for quantification of confocal data: Schmidt et al., 2008; Weyhersmüller et al., 2011; Goel et al., 2017; Böhme et al., 2019). Radii estimated from the area under the assumption of a circular area are displayed. Strikingly, the radius measured in confocal resolution by thresholding shows an inverse effect compared to the given radii and the radii computed by alpha shapes.

(D) Radii after imposing confocal resolution and thresholding on simulated localization

clusters (magenta, mean ± SD, n = 10 simulated clusters per given radius) and line for

real radius (grey) plotted against the given radius for simulation. Symbols (i, ii)

correspond to the given radii used in (C). Note the drop of the estimated radius above

a given radius of 250 nm (corresponding to a localization density roughly below 7600

locs. per μm^2). This implies that the area of large objects with low molecular density

(i.e. lower fluorescence intensity) is systematically underestimated by pixel- and

intensity-based quantification offering an explanation for the apparent discrepancy

between our data and earlier findings. Scale bars 200 nm in (A) and (C).

Supplementary Figure 5. Homeostatic potentiation reduces subcluster spacing.
(A, B) Minimum distance between subcluster centers (mean per AZ). Sc. center distance is decreased in acute PHP (ctrl: grey, n = 568 AZs from 13 NMJs from 6 animals for (A, C); phtx: red, n = 790 AZs from 14 NMJs and 5 animals for (A, C); p = 0.024) and in chronic PHP (ctrl: grey, n = 872 AZs from 19 NMJs from 6 animals for (B, D); GluRIIAko: blue, n = 1020 AZs from 19 NMJs from 6 animals for (B, D); p < 0.001).

(C, D) Distance between subclusters (mean sc. diameter per AZ substracted from sc. center distance) in acute (C) and chronic (D) PHP. The distance between subclusters (inter sc. distance) is unchanged in acute PHP (p = 0.354) but reduced in chronic PHP (p = 0.007).

Supplementary Table 1. Data summary and statistical information.

Numerical values not stated in text and figure legends including p-values and sample sizes. Data were not normally distributed and Mann-Whitney rank sum test was used to evaluate statistical significance. The largest number stated for n indicates the sample size used for the statistical test.

parameter	median (25th-75th percentile)		p-value	figure
Brp ctrl vs. phtx				
locs. per AZ	1532 (1007-2303)	1453 (926-2346) (792)	0.236	2B
AZ area [µm^2]	0.109 (0.074-0.169)	0.101 (0.068-0.148)	0.009	2B
loc. density [locs./µm^2]	13555 (11308-16236)	14082 (11756-17204)	0.009	2B
sc. per AZ	15 (10-21)	13 (9-21)	0.084	3B
locs. per sc.	57 (51-64)	56 (51-64)	0.480	
sc. area [nm^2]	1677 (1342-2149)	1617 (1301-1994)	0.025	3B
sc. loc. density [locs./µm^2]	33914 (28276-40553)	34290 (29464-42454)	0.022	3B
radial distance [nm]	125 (102-163)	116 (96-153)	0.003	3B
circularity [a.u.]	0.64 (0.52-0.75)	0.65 (0.52-0.77)	0.161	
n (AZs, NMJs, animals)	568, 13, 6	792, 14, 5		
locs. per bouton	40352 (19442-46915)	33922 (16721-44260)	0.577	S2B
unclustered locs. per bouton	1383 (428-2309)	1065 (593-1493)	0.903	S2B
n (distal Ib boutons, animals)	13, 6	14, 5		
AZ with circularity ≥ 0.5				
AZ area [µm^2]	0.103 (0.075- 0.149)	0.094 (0.068- 0.135)	0.002	S2D
radial distance [nm]	119 (101-148)	112 (95-139)	< 0.001	S2D
n (AZs, NMJs, animals)	442, 13, 6	621, 14, 5		
Brp ctrl vs. GluRIIAko				
locs. per AZ	1546 (1067-2270)	1480 (1059-2325)	0.391	2D
AZ area [µm^2]	0.107 (0.081-0.156)	0.096 (0.069-0.144)	< 0.001	2D
loc. density [locs./µm^2]	14203 (12091-16481)	15414 (13522-17294)	< 0.001	2D
sc. per AZ	15 (10-21)	13 (10-21)	0.118	3D
locs. per sc.	55 (50-62)	55 (50-62)	0.860	
sc. area [nm^2]	1588 (1291-1960)	1483 (1228-1750)	< 0.001	3D
sc. loc. density [locs./µm^2]	35053 (29821-40440)	37622 (33003-42921)	< 0.001	3D
radial distance [nm]	116 (98-148)	113 (93-146)	0.002	3D
n (AZs, NMJs, animals)	872, 19, 6	1020, 19, 6		
Brp ctrl vs. phtx muscle 4				
locs. per AZ	1591 (1022-2321)	1519 (995-2378)	0.705	
AZ area [µm^2]	0.115 (0.081-0.166)	0.108 (0.075-0.161)	0.018	S3B
loc. density [locs./µm^2]	13197 (11146-15712)	13757 (11744-16197)	< 0.001	S3C
sc. per AZ	14 (9-20)	14 (9-20)	0.803	S3D
locs. per sc.	54 (49-61)	54 (49-61)	0.834	
sc. area [nm^2]	1724 (1374-2138)	1679 (1331-2090)	0.020	S3E
sc. loc. density [locs./µm^2]	31505 (26557-37713)	32550 (27702-38679)	0.003	S3F
radial distance [nm]	123 (99-158)	119 (96-155)	0.022	S3G
n (AZs, NMJs, animals)	1183, 20, 6	1445, 22, 8		
simulated localization clusters, given radius 400 and 300 nm				

alpha shape area [µm^2]	0.463	0.270		S4A
measured radius [nm]	384	293		S4A
thresholding area [µm^2]	0.090	0.202		S4C
measured radius [nm]	170	253		S4C
correlative confocal-*d*STORM microscopy				
AZ area [µm^2]	0.164 (0.103-0.246)	0.101 (0.0768-0.147)		4D
n (AZs, NMJs, animals)	343, 12, 7			
Brp ctrl vs. phtx in confocal and STED simulation				
AZ area [µm^2]	0.115 (0.051-0.204)	0.134 (0.057-0.256)	0.026	5D
mean pixel intensity [a.u.]	58.6 (54.0-65.4)	60.3 (54.4-68.9)	0.016	5E
n (AZs, NMJs, animals)	345, 13, 6	457, 14, 5		
AZ area [µm^2]	0.081 (0.052-0.125)	0.085 (0.058-0.131)	0.044	5G
AZ diameter [nm]	322 (257-399)	329 (273-408)	0.044	5G
n (AZs, NMJs, animals)	591, 13, 6	820, 14, 5		
intensity peaks per AZ	3 (1-4)	4 (2-5)	0.002	5G
n (AZs, NMJs, animals)	122, 12, 6	186, 14, 5		
Brp subcluster distances ctrl vs. phtx				
sc. center distance [nm]	55 (48-64)	54 (47-62)	0.024	S5A
inter sc. distance [nm]	9 (5-13)	9 (4-14)	0.354	S5C
n (AZs, NMJs, animals)	568, 13, 6	790, 14, 5		
Brp subcluster distances ctrl vs. GluRIIAko				
sc. center distance [nm]	53 (47-60)	51 (46-57)	< 0.001	S5B
inter sc. distance [nm]	8 (4-12)	7 (4-11)	0.007	S5D
n (AZs, NMJs, animals)	872, 19, 6	1020, 19, 6		

1056