Title Page

Significant improvement of reprogramming efficiency by transient overexpression of ZGA inducer Dux but not Dppa2/4

Lei Yang[1], Xuefei Liu[1], Lishuang Song[1,2], Anqi Di[1,2], Guanghua Su[1,2], Chunling Bai[1,2], Zhuying Wei[1,2], and Guangpeng Li[1,2]

1 State Key Laboratory of Reproductive Regulation and Breeding of Grassland Livestock (R2BGL), Inner Mongolia University, Hohhot 010070, China.

2 College of Life Sciences, Inner Mongolia University, Hohhot 010070, China.

Correspondence: Guangpeng Li (gpengli@imu.edu.cn)

Running Title: Dux promotes somatic cell reprogramming

Abstract

Cloned animal have been reported by somatic cell nuclear transfer (SCNT) for many years. However, the SCNT biotechnology is extremely inefficient, and zygotic genome activation (ZGA) is required for SCNT and chemical mediated somatic cell reprogramming. To identify candidate factors that facilitate ZGA in SCNT reprogramming, we carried out a siRNA-repressor and mRNA-inducer screening in SCNT to identify epigenetic and chromatin regulators of the ZGA transcriptional program. Through a series of experiments, we demonstrate that transient overexpression of Dux not only improved SCNT efficiency, but also increased the efficiency of chemical reprogramming. Furthermore, transient overexpression of Dux combined with inactivation of DNA methyltransferases (Dnmts) further promote the overall development of SCNT-derived animals. These findings enhance our understanding of ZGA-regulators in SCNT and chemical mediated somatic cell reprogramming, and suggest that transient overexpression of Dux may serve as a useful tool to enhances the somatic reprogramming efficiency.

Introduction

The scientists at the Institute of Neuroscience (Chinese Academy of Sciences), recently cloned the world's first monkeys Zhong Zhong and Hua Hua (the Zhong Hua is a mandarin term for china) [1]. After the first mammal Dolly sheep to be cloned [2], the created non-human primate has once again attracted worldwide attention. Indeed, successful cloning of the sheep and monkey not only made reproductive cloning possible, but also raised the possibility of therapeutic cloning. Somatic cell nuclear transfer (SCNT) is the technical term for cloning in which terminal differentiated somatic cell is transferred into enucleated oocyte, in which the cytoplasmic factors reprogram the nucleus to become a zygote-like state [3]. Both SCNT derived and Yamanaka factors induced pluripotent stem (iPS) cells have the potential to generate patient-specific pluripotent stem cells for replacement therapy. Compared with iPS, the SCNT reprogramming is transgene-free and oncogene-free methodologies. Thus, SCNT derived pluripotent stem cells would be more suitable for human therapeutic applications.

Following fertilization, the newly formed zygotic genome is activated through a process known as the zygotic genome activation (ZGA), which enables zygotes can subsequently drive the differentiation of all the diverse cell types of the adult animal [4]. A similar mechanism is likely used in ZGA of SCNT embryos [3,5]. The 'erase-and-rebuild' strategy is used to establish the zygotic transcription program during SCNT reprogramming [5]. Consistently, our and others recent results have revealed that incomplete ZGA is a major SCNT reprogramming barrier [6-8]. Although many advances have been made in SCNT technology via different epigenetic regulators, the SCNT for producing cloned animals still inefficient.

Materials and Methods

Animals and Chemicals

Specific pathogen-free (SPF)-grade BDF1 (C57BL/6N × DBA/2), CD1, and Kun-Ming (KM) mice mice were purchased from Laboratory Animal Research Center (Inner Mongolia University) or Vital River Laboratories (Beijing, China). All animal experiments were approved by the Animal Care and Use Committee of Inner Mongolia University. All procedures were carried out in strict accordance with the recommendations made in the Guide for the Care and Use of Laboratory Animals of the National Veterinary and Quarantine Service. MERLV:: tdTomato transgenic mice, which carry a transgenic MERLV promoter LTR driven tdTomato reporter [8], were mated with the same positive mice. dDux-MEFs cells were isolated from E12.5 fetus. The MERLV:: tdTomato embryos used in the experiment were produced by MERVL::tdTomato sperm and MII oocytes from the littermates of transgenic mice. All chemicals used in this study were purchased from Sigma (USA), unless otherwise indicated.

SCNT and embryo culture

The SCNT processes followed previously published studies [8,9]. Briefly, groups of ~50 MII oocytes were transferred to a chamber containing oil-covered M2 supplemented with 5 µg/ml cytochalasin B (CB). The spindle-chromosome complex (SCC) was removed by a blunt Piezo-

driven pipette (~10 µm internal diameter) on a 37°C heating stage of an inverted microscope

(Nikon, Japan). The nuclei of donor cells were drawn in and out of the injection pipette until its

plasma membrane was broken and was then injected into enucleated oocytes. The

reconstructed embryos were cultured in aMEM medium (Thermo, USA) containing 10% fetal

calf serum (FCS; Hyclone, USA) for 1 h before activation treatment. The reconstructed

embryos were activated in Ca_{2+} free KSOM medium containing 10 mM strontium and 5 µg/ml

CB for 6 h. Activated embryos were thoroughly washed and cultured in G1/G2 medium (1:1,

vol / vol; Vitrolife, Sweden). Induction of Dux was performed by administration of doxycycline

(2 µg/ml) in the G1/G2 medium.

Immunofluorescence staining

Embryos and CiPS cells were rinsed three times in phosphate-buffered saline (PBS) with 0.3%

BSA, fixed with 4% paraformaldehyde (PFA) overnight at 4°C, and then permeabilized with

0.2% (vol / vol) Triton X-100 for 15 min at room temperature, followed by washing thoroughly

in PBS containing 0.3% BSA. Fixed samples were blocked in 0.05% Tween 20 in PBS

containing 3% BSA (PBST) at 37°C for 1 h and then incubated with the primary antibodies

overnight at 4°C. Samples were incubated with primary antibodies against HA (Santa Cruz,

sc-7392, USA), Oct4 (Santa Cruz, sc-8629, USA), and Ssea1 (Santa Cruz, sc-21702, USA).

After incubating, the samples were needed to wash several times in PBST and then incubated

with appropriate secondary antibodies conjugated with Alexa Fluor 594 and Alexa Fluor 488

(Thermo, USA) for 1 h at 37°C. For imaging, the embryos were mounted in 10 µl anti-fade

solution with DAPI (Thermo, USA) and compressed with a cover-slip. All samples were

observed by laser scanning microscope (Nikon, Japan).

***In vitro* mRNA synthesis and microinjection in oocytes**

The *Dux*, *Dppa2*, and *Dppa4* mRNA were cloned into T7-driven vectors, and mRNAs were

synthetized *in vitro* using mMESSAGE mMACHINE T7 Ultra Kit (Thermo, USA) following the

manufacturer's instructions. The integrity of manufactured mRNA was confirmed by

electrophoresis with formaldehyde gels. The final concentration of mRNA was diluted to 900

ng/ml before injection. As previously described [8], 8 pl of mRNA was microinjected into the

cytoplasm of denuded oocytes. These oocytes were obtained by superovulating BDF1 female

mice by intraperitoneal injection of 10 IU pregnant mare serum gonadotropin (PMSG;

Sansheng, China) and 10 IU human chorionic gonadotropin (hCG; Sansheng, China) 48 h

apart. The cumulus-oocyte complexes (COCs) were collected from oviducts 14 h post-hCG,

and the cumulus cells were dispersed by EmbryoMax FHM Mouse Embryo Media (Millipore,

USA). Oocytes were injected using Piezo-operated blunt-end micropipette (3~5 μm internal

diameter). After injection, oocytes were kept at room-temperature for 30 min and then moved

into the incubator. The siRNA information was presented in Table S4.

Generation of CiPS cells

This section was adapted from Deng's lab [10]. Reagents setup: Small molecules: VPA,

CHIR99021, 616452, Tranylcypromine, Forskolin, Ch55, EPZ004777, DZNep, Decitabine,

SGC0946, and PD0325901. Stage I medium: FBS/KSR-based medium supplemented with

the small-molecule cocktail VC6TF5E (100 μM VPA, 40 μM CHIR99021, 10 μM 616452, 5 μM

Tranylcypromine, 10 μM Forskolin, 1 μM Ch55, and 5 μM EPZ004777). N2B27-SII medium:

N2B27-based medium supplemented with 10 ng/ml LIF, 50 μg/ml vitamin C, 25 ng/ml bFGF,

2 mg/ml Albumax-II, and the small-molecule cocktail VC6TF5ZDS (VPA at 1 mM). Stage III

medium: N2B27-based medium with 3 μM CHIR99021, 1 μM PD0325901, 10 ng/ml LIF, and

50 μg/ml vitamin C. On day –1, MEFs were seeded at 50,000 cells per well of 6-well plate with

MEF culture medium. The next day (day 0), change the medium into optimized stage I medium.

From day 6~12, cells were cultured in optimized N2B27-SII medium. During day 12~16, cells

were cultured in stage III medium with 500 μM VPA. On day 16, VPA was removed and cells

were cultured in stage III medium. After another 4~10 days, CiPS colonies emerged. Induction

of Dux was performed by administration of doxycycline (2 μg/ml) in the culture medium.

Transgenic mice generation

The pCW57.1-mDux-CA vector was a gift from Stephen Tapscott (Addgene 99284). The

pronuclear microinjection for the production of transgenic mice followed previously published

studies [11]. Briefly, the vector was injected into the well-recognized pronuclei. Injected zygotes

were transferred into pseudopregnant female mice (~ 30 zygotes per mouse) after 4 h

recovery culture in KSOM-AA medium (Millipore, USA). For founder identification, tail tips were

subjected to standard DNA extraction procedures. The amplified DNA fragments were

subjected to TA cloning and sequencing. The founder mice were crossed to the littermates of

founder mice for four generations to produce homozygous Dux mice.

Alkaline phosphatase staining

For alkaline phosphatase staining, CiPS cells were fixed with 4% PFA in PBS for 2 min, rinsed

once with PBS and detection was performed using a Alkaline Phosphatase Assay Kit

(VECTOR, USA) according to the manufacturer's protocol.

Real-time qPCR analysis

Total RNA was isolated using TRIzol reagent (Thermo, USA) and was immediately reverse-

transcribed using a Prime Script RT reagent kit (Takara, Japan). The reverse transcription

PCR was were amplified using Ex Taq (Takara, Japan). The qPCR was performed using a

SYBR Premix Ex Taq (Takara, Japan) and signals were detected with Applied Biosystems

7500 real-time PCR System (Thermo, USA). Relative mRNA expression was calculated use

the $2^{(-\Delta\Delta Ct)}$ method. The primer information was presented in Table S5.

Chimeric mice and embryo transfer

For chimeric experiments, CiPS cells were used one day before passaging, which showed an

optimal undifferentiated morphology. The CiPS cells were microinjected into CD1/KM

blastocysts using a Piezo (Primetech, Japan) microinjection pipette. After culturing for 3 h, the

embryos were transplanted into the uterus of pseudopregnant mice (~15 embryos per mouse).

The 2-cell stage SCNT embryos were transferred to the oviducts of E0.5 pseudopregnant (~20

embryos per mouse). The cloned pups nursed by lactating CD1/KM females. SSLP analysis
was performed for D6Mit15 and D2Mit102. The primer information is presented in Table S5.
154

Teratoma formation
Teratoma formation analysis was carried out to evaluate the pluripotency of CiPS cells.
Approximately 10^6 CiPS cells were injected subcutaneously into the hind limbs of a 8-week-
old nude mice. After 6 weeks, fully formed teratomas were dissected and fixed with PBS
containing 4% PFA, then embedded in paraffin, sectioned and stained with haematoxylin and
eosin for histological analysis.
161

162 **Bisulfite genomic sequencing**
163 Genomic DNA was extracted by the tissue & blood DNA extraction kit (Qiagen, Germany) and
164 treated with the Methylamp DNA modification kit (Epigentek, USA). The bisulphite conversion
165 was performed as previously published studies [12].
166

RNA-sequencing
The single-cell RNA-seq method followed previously published studies [7,13]. The reverse
transcription was performed directly on the cytoplasmic lysate of individual embryos. The total
cDNA library was then amplified by 18~20 cycles for library construction, which was performed
following manufacturer's instructions (Illumina, USA). Paired-end sequencing was further
performed at Annoroad (Beijing, China). Three biological replicates were analyzed for each
treatment condition. After removing low-quality reads and adapters, the raw reads were
mapped to the mm9 genome using Tophat (v1.3.3) with default parameters [14]. Expression
levels for all RefSeq transcripts were quantified to fragments per kilobase of exon model per
million mapped reads (FPKM) using Cufflinks (v1.2.0) [15]. Gene Ontology (GO) analyses for
differentially expressed genes were performed by using the R package GO.db and Database
for Annotation, Visualization and Integrated Discovery (DAVID) [16].
179

Statistics analysis and data availability

Statistical analyses were done using the univariate analysis of variance (ANOVA) followed by the Student t-test with SPSS 21.0 statistical software (Armonk, USA). $P < 0.05$ was considered significant. Sequencing data have been deposited in the NCBI sequence read archive (SRA) under accession codes: SCNT-1 (SAMN12871322), SCNT-2 (SAMN12871323), SCNT-3 (SAMN12871324); D-SCNT-1 (SAMN12871325), D-SCNT-2 (SAMN12871326), D-SCNT-3 (SAMN12871327); ICSI-1 (SAMN12871328), ICSI-2 (SAMN12871329), ICSI-3 (SAMN12871330).

Results and Discussion

We have recently established a MERVL::tdTomato based ZGA real-time monitor system [8]. To identify candidate factors that facilitate ZGA in SCNT reprogramming, we carried out a siRNA-repressor and mRNA-inducer screening in SCNT (Fig.1A). From the screen, we identified a number of factors that when over-expression or knock-down resulted in reporter activation (Supplementary information, Fig.S1A). Among the identified factors, Dux [17-19], Dppa2 and Dppa4 [20-22] showed the strongest phenotype (Fig.1B, 1C). Intriguingly, we found that most SCNT embryos injected with Dppa2 or Dppa4 mRNA arrested at the 1-cell stage (called '1-cell block' for simplicity; Fig.1D, 1E; Supplementary information, Fig.S1B, Movie S1, S2). Direct injection of Dux mRNAs had no significant impact on the blastocyst formation rate (*vs.* canonical SCNT; 21.2% *vs.* 23.6%; Fig.1D; Supplementary information, Table S1). Indeed, direct injection of Dux will increase the rate of SCNT embryo fragmentation (Fig.1D, 1E; Supplementary information, Movie S2). Nevertheless, when Dux was over-expressed in SCNT embryos, the rate of 2-cell block was significantly reduced (Fig.1D). These results suggest that Dux (but not Dppa2/4) is a key factor whose overexpression can rescue 2-cell arrest in SCNT embryos.

It is known that ZGA is governed by a time-dependent 'zygotic clock' (i.e. 24h post-fertilization), and Dux is expressed as a brief pulse in early 2-cell stage [5, 20,21,23]. Therefore, we

wonder whether Dux is time-depend on improving the efficiency of SCNT embryo development.

To answer this question, we generated transgenic mouse lines containing doxycycline (dox)

inducible Dux gene (Supplementary information, Fig.S2A-B). By switching the culture medium

between dox-containing and -lacking at various time points, the critical period for the dox-Dux

requirement was mapped to a 24-h window (Fig.1F). When the dox was supplied in this

window, the SCNT blastocyst formation rate was significantly increased (Fig.1G;

Supplementary information, Table S2). Subsequently, we performed RNA sequencing (RNA-

seq) to examine the possible changes of the ZGA related gene expressions in Dux-

overexpressed 2-cell SCNT embryos (D-SCNT). Principal component analysis (PCA)

revealed that the ICSI and D-SCNT generated embryos in the same cluster, and ZGA-genes

were upregulated in D-SCNT embryos (Fig.1H, 1I; Supplementary information, Fig.S3A-D;

Dataset S1, S2). These results indicating the improvement of Dux on SCNT embryonic

development depends on its treatment time.

The SCNT blastocyst rate was improved by injecting Kdm4d mRNA (K-SCNT) [6]. We next

compared the developmental potential of SCNT embryos derived by K-SCNT and D-SCNT.

We found that the blastocyst formation rate of D-SCNT was similar to K-SCNT (84.72% *vs.*

86.79%; Supplementary information, Fig.S4A, Table S2). Nonetheless, the cloned pup birth

rate of D-SCNT was higher than K-SCNT (10.71% *vs.* 7.81%; Fig.1J; Supplementary

information, Table S3). Despite D-SCNT or K-SCNT, the abnormally large placentas are still

observed in those cloned pups (Supplementary information, Fig.S4B, Table S3). Previous

studies have indicated that knock-down DNA-methyltransferases Dnmt3a/b led to less

placental abnormalities (Si-SCNT) [24]. We next investigated whether combined D-SCNT and

Si-SCNT could further improve SCNT embryonic development. Compared with canonical

SCNT, combined K-SCNT and Si-SCNT improved the pup birth-rate from 1.01% to 12.24%,

but less than D-SCNT combined with Si-SCNT (18.60%; Fig.1J). Genotyping analysis

confirmed that all the cloned pups were generated from donor cells (Supplementary

information, Fig.S4C). The large placental phenotype was rescued by the D-Si-SCNT

combined approach (Supplementary information, Fig.S4B).

Deng et al. identified a unique embryonic 2-cell like status as the key molecular event that

led to successful chemical reprogramming (CiPS) [10]. Thus, we tried to improve the CiPS by

overexpression of ZGA inducer Dux (D-CiPS). The dox-inducible mouse embryonic fibroblasts

(dDux-MEFs) cells were isolated from the transgenic mice. As shown in Fig.1K, we treated the

dDux-MEFs with dox for various durations, and observed that Dux displayed remarkable effect

at the stage-2 of CiPS. About 36 alkaline phosphatase-positive (AP+) colonies could be

observed in D-CiPS wells (starting from 50,000 MEFs per well of 6-well plate), while only 26

AP+ colonies could be observed in the canonical CiPS well (Fig.1K; Supplementary

information, Fig.S5A, S5B). Consistently, the ZGA related genes were significantly

upregulated after transient expression of Dux (Supplementary information, Fig.S5C). These

D-CiPS induced cells were pluripotent as demonstrated by standard characterization

procedures (Supplementary information, Fig.S5D-F). We found, based on bisulfite sequencing,

that successful demethylation occurred in the promoters of Oct4 and Nanog in the D-CiPS

induced cell lines (Fig.1L).

In summary, we demonstrate here that transient overexpression of Dux not only improved

SCNT efficiency, but also increased the efficiency of chemical reprogramming. The Dux factor

was recently identified as a inducer of ZGA in mouse, human, and likely all mammalian

animals [17,25]. Thus, we reasoned that transient overexpression of Dux could also improves

reprogramming efficiency of all mammals.

Acknowledgements

We thank Prof. Shaorong Gao (Tongji University), Prof. Zhiming Han (Chinese Academy

of Sciences), and Prof. Qing Xia (Peking University) for their technical assistance. We also

thank Prof. Yi Zhang (Harvard Medical School) for sharing the list of RRRs. This study was

supported by the Genetically Modified Organisms Breeding Major Projects (2016ZX08007-

Author contributions

269   L.Y. and G.L. conceived and designed the study. L.Y., X.L., L.S., A.D., G.S., C.B., and Z.W.

270 performed the experiments. L.Y., X.L., L.S., A.D., and G.S. analyzed the data. G.L. supervised

271 the project. L.Y. and G.L. wrote the paper.

Competing interests

274 The authors declare no competing interests.

References

1. Liu, Z. *et al.* Cloning of macaque monkeys by somatic cell nuclear transfer. *Cell*
172, 881-887 (2018).
2. Wilmut, I., Schnieke, A.E., McWhir, J., Kind, A.J. & Campbell, K.H. Viable offspring
derived from fetal and adult mammalian cells. *Nature* **385**, 810-3 (1997).
3. Matoba, S. & Zhang, Y. Somatic cell nuclear transfer reprogramming: mechanisms
and applications. *Cell Stem Cell* **23**, 471-485 (2018).
4. Lu, F. & Zhang, Y. Cell totipotency: molecular features, induction, and
maintenance. *Natl Sci Rev* **2**, 217-225 (2015).
5. Gao, T. *et al.* Nuclear reprogramming: the strategy used in normal development
is also used in somatic cell nuclear transfer and parthenogenesis. *Cell Res* **17**, 135-
50 (2007).
6. Matoba, S. *et al.* Embryonic development following somatic cell nuclear transfer
impeded by persisting histone methylation. *Cell* **159**, 884-95 (2014).
7. Liu, W. *et al.* Identification of key factors conquering developmental arrest of
somatic cell cloned embryos by combining embryo biopsy and single-cell
sequencing. *Cell Discov* **2**, 16010 (2016).
8. Yang, L., Song, L., Liu, X., Bai, L. & Li, G. KDM6A and KDM6B play contrasting roles
in nuclear transfer embryos revealed by MERVL reporter system. *EMBO Rep* **19**,
e46240 (2018).
9. Kishigami, S. *et al.* Production of cloned mice by somatic cell nuclear transfer. *Nat*
Protoc **1**, 125-38 (2006).
10. Zhao, T. *et al.* Single-cell RNA-seq reveals dynamic early embryonic-like
programs during chemical reprogramming. *Cell Stem Cell* **23**, 31-45 (2018).
11. Ittner, L.M. & Gotz, J. Pronuclear injection for the production of transgenic mice.
Nat Protoc **2**, 1206-15 (2007).
12. Cao, S. *et al.* Chromatin accessibility dynamics during chemical induction of
pluripotency. *Cell Stem Cell* **22**, 529-542 (2018).
13. Tang, F. *et al.* RNA-Seq analysis to capture the transcriptome landscape of a single
cell. *Nat Protoc* **5**, 516-35 (2010).
14. Trapnell, C., Pachter, L. & Salzberg, S.L. TopHat: discovering splice junctions with
RNA-Seq. *Bioinformatics* **25**, 1105-11 (2009).

15. Trapnell, C. *et al.* Transcript assembly and quantification by RNA-Seq reveals
unannotated transcripts and isoform switching during cell differentiation. *Nat
Biotechnol* **28**, 511-5 (2010).
16. Huang da, W., Sherman, B.T. & Lempicki, R.A. Systematic and integrative analysis
of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57
(2009).
17. Iturbide, A. & Torres-Padilla, M.E. Starting embryonic transcription for the first
time. *Nat Genet* **49**, 820-821 (2017).
18. Hendrickson, P.G. *et al.* Conserved roles of mouse DUX and human DUX4 in
activating cleavage-stage genes and MERVL/HERVL retrotransposons. *Nat Genet*
49, 925-934 (2017).
19. De Iaco, A. *et al.* DUX-family transcription factors regulate zygotic genome
activation in placental mammals. *Nat Genet* **49**, 941-945 (2017).
20. Eckersley-Maslin, M. *et al.* Dppa2 and Dppa4 directly regulate the Dux-driven
zygotic transcriptional program. *Genes Dev* **33**, 194-208 (2019).
21. De Iaco, A., Coudray, A., Duc, J. & Trono, D. DPPA2 and DPPA4 are necessary to
establish a 2C-like state in mouse embryonic stem cells. *EMBO Rep* **20**, e47382
(2019).
22. Yan, Y.L. *et al.* DPPA2/4 and SUMO E3 ligase PIAS4 opposingly regulate zygotic
transcriptional program. *PLoS Biol* **17**, e3000324 (2019).
23. Rodriguez-Terrones, D. *et al.* A molecular roadmap for the emergence of early-
embryonic-like cells in culture. *Nat Genet* **50**, 106-119 (2018).
24. Gao, R. *et al.* Inhibition of aberrant DNA re-methylation improves post-
implantation development of somatic cell nuclear transfer embryos. *Cell Stem Cell*
23, 426-435 (2018).
25. Whiddon, J.L., Langford, A.T., Wong, C.J., Zhong, J.W. & Tapscott, S.J. Conservation
and innovation in the DUX4-family gene network. *Nat Genet* **49**, 935-940 (2017).
335

Figure legends

Figure 1 Dux act as promote factors for SCNT- and chemical-mediated somatic cell reprogramming.

(A) Schematic illustration of the screen strategy.

(B) Quantification of embryos that expression ZGA reporter (MERVL::tdTomato) after injection with siRNA-repressor or mRNA- inducer. N, total number of embryos analyzed for each condition. Error bars, mean ± S.D.; three independent experiment replicates were performed.

(C) Representative fluorescence image of SCNT embryos derived by different methods. The SCNT embryos produced by transfer of MERVL::tdTomato cumulus cell into WT enucleated oocytes. Scale bar, 100 μm.

(D) Stacked bar plots show fraction of embryos at blastocyst stages after injection with different mRNA as indicated. At least three independent experiment replicates were performed. N, total number of embryos analyzed for each condition.

(E) Representative live-cell images of dynamics ZGA reporter (MERVL::tdTomato) expression during SCNT embryos preimplantation development. The selected images from a series acquired every 15 min. The time after starting observation is shown on the bottom of image. Scale bar, 50 μm.

(F) Schematic transient induction of Dux. All timings reported in this paper is hour post-activation.

(G) The preimplantation development of SCNT embryos derived from D-SCNT or SCNT. Shown is the percentage of embryos that reaches each indicated stage. Scale bar, 100 μm.

(H) Unsupervised hierarchical clustering. The D-SCNT embryos mostly resembled the ICSI derived embryos.

(I) Venn diagram showing the overlap between the genes that failed to be activated in SCNT 2-cell embryos and derepressed in D-SCNT. Heatmap, GO terms showing the expression pattern of 257 overlap genes.

(J) The birth rate of SCNT embryos derived by different methods.

(K) Durations of dox addition at different time points during chemical induction (upper) and

assessment of AP+ colonies (bottom). Error bars, mean ± S.D.; the Dox inducible Dux-

MEFs were used for chemical reprogramming.

(L) Bisulfite sequencing analysis of Oct4 and Nanog demethylation in D-CiPS or canonical

CiPS cells (filled squares are methylated CpGs and empty ones indicate unmethylated

sites).

Fig.1