Short Report

Significant improvement of reprogramming efficiency by transient overexpression of ZGA inducer Dux but not Dppa2/4

Lei Yang[1], Xuefei Liu[1], Lishuang Song[1, 2], Anqi Di[1, 2], Guanghua Su[1, 2], Chunling Bai[1, 2], Zhuying Wei[1, 2], and Guangpeng Li[1, 2, *]

1 State Key Laboratory of Reproductive Regulation and Breeding of Grassland Livestock (R$_2$BGL), Inner Mongolia University, Hohhot 010070, China.

2 College of Life Sciences, Inner Mongolia University, Hohhot 010070, China.

* Correspondence: Guangpeng Li (gpengli@imu.edu.cn)

Abstract

Cloned animal have been reported by somatic cell nuclear transfer (SCNT) for many years.

However, the SCNT is extremely inefficient, and zygotic genome activation (ZGA) is required

for SCNT and chemical mediated reprogramming. To identify candidate factors that facilitate

ZGA in reprogramming, we performed siRNA-repressor and mRNA-inducer screening, which

revealed Dux, Dppa2, and Dppa4 as key factors enhancing the ZGA in SCNT. Direct injection

of ZGA-inducers had no significant effect on the SCNT blastocyst formation, and even

destroyed the ZGA. Through a inducible Dux transgenic mouse model, we demonstrate that

transient overexpression of Dux not only improved SCNT efficiency, but also increased the

efficiency of chemical reprogramming. Transcriptome profiling revealed that Dux treated

SCNT embryos were similar to fertilized embryos. Furthermore, transient overexpression of

Dux combined with inactivation of DNA methyltransferases (Dnmts) further promote the overall

development of SCNT-derived animals. These findings enhance our understanding of ZGA-

regulators in somatic reprogramming.

Introduction

The scientists at the Institute of Neuroscience (Chinese Academy of Sciences), recently cloned the world's first monkeys Zhong Zhong and Hua Hua (the Zhong Hua is a mandarin term for china) (Z. Liu et al., 2018). After the first mammal Dolly sheep to be cloned (Wilmut, Schnieke, McWhir, Kind, & Campbell, 1997), the created non-human primate has once again attracted worldwide attention. Indeed, successful cloning of the sheep and monkey not only made reproductive cloning possible, but also raised the possibility of therapeutic cloning. Somatic cell nuclear transfer (SCNT) is the technical term for cloning in which terminal differentiated somatic cell is transferred into enucleated oocyte, in which the cytoplasmic factors reprogram the nucleus to become a zygote-like state (Matoba & Zhang, 2018). Both SCNT derived and Yamanaka factors induced pluripotent stem (iPS) cells have the potential to generate patient-specific pluripotent stem cells for replacement therapy. Compared with iPS, the SCNT reprogramming is oncogene-free methodologies. Thus, SCNT derived pluripotent stem cells would be more suitable for human therapeutic applications.

Following fertilization, the newly formed zygotic genome is activated through a process known as the zygotic genome activation (ZGA), which enables zygote can subsequently development to the adult animal (Lu & Zhang, 2015). A similar mechanism is likely used in ZGA of SCNT embryos (T. Gao et al., 2007; Matoba & Zhang, 2018). Indeed, our and others recent results have revealed that incomplete ZGA is a major reprogramming barrier (W. Liu et al., 2016; Matoba et al., 2014; Wang et al., 2018; Yang, Song, Liu, Bai, & Li, 2018). Although many advances have been made in SCNT technology via different epigenetic regulators, the SCNT for producing cloned animals still inefficient.

The transcription factor of double homeobox (Dux) was identified as a key inducer of ZGA in normal fertilized embryos (De Iaco et al., 2017; Hendrickson et al., 2017; Whiddon, Langford, Wong, Zhong, & Tapscott, 2017). Very recently, two independent studies found that transcription factors developmental pluripotency associated 2 (Dppa2) and Dppa4 are both

necessary and sufficient for the activation of ZGA (De Iaco, Coudray, Duc, & Trono, 2019;

Eckersley-Maslin et al., 2019). Therefore, as far as we know, at least three factors can be

called master ZGA-inducer in fertilized embryos. Furthermore, overexpression of Dux is

sufficient to drive pluripotent stem cells into a totipotent state by expressing 2-cell embryo

specific transcripts (2-cell-like stem cells) (Macfarlan et al., 2012). In addition, several studies

show totipotent state that the be achieved by depleting ZGA-repressor, such as tripartite motif-

containing 28 (Trim28) (Rodriguez-Terrones et al., 2018), microRNA 34a (miR-34a) (Choi et

al., 2017), chromatin assembly factor (CAF-1) (Ishiuchi et al., 2015), lysine (K)-specific

demethylase 1A (Lsd1/Kdm1a) (Ancelin et al., 2016; Macfarlan et al., 2012; Wasson et al.,

2016), and long interspersed nuclear element (LINE1) (Percharde et al., 2018).

The effect of ZGA-regulators in somatic cell reprograming remains unknown. Thus, the aims

of this study were to explore the relationship between ZGA-regulator and SCNT efficiency,

and to identify a smart factor that could promotes the efficiency of somatic reprogramming.

Results and Discussion

We have recently established a MERVL::tdTomato based ZGA real-time monitor system (Yang

et al., 2018). To identify candidate factors that facilitate ZGA in SCNT reprogramming, we

carried out a siRNA-repressor and mRNA-inducer screening in SCNT (Figure 1A). From the

screen, we identified a number of factors that when over-expression or knock-down resulted

in reporter activation (supplement Figure S1A). Among the identified factors, Dux, Dppa2, and

Dppa4 showed the strongest phenotype (Figure 1B and C). Intriguingly, we found that most

SCNT embryos injected with Dppa2 or Dppa4 mRNA arrested at the 1-cell stage (called '1-

cell block' for simplicity; Figure 1D and E; supplement Figure S1B, Movie S1, S2). Direct

injection of Dux mRNAs had no significant impact on the blastocyst formation rate (*vs.*

canonical SCNT; 21.2% *vs.* 23.6%; Figure 1D; supplement Table S1). Indeed, direct injection

of Dux will increase the rate of SCNT embryo fragmentation (Figure 1D and E; supplement

Movie S2). Nevertheless, when Dux was over-expressed in SCNT embryos, the rate of 2-cell

block was significantly reduced (Figure 1D). These results suggest that Dux (but not Dppa2/4)

is a key factor whose overexpression can rescue 2-cell arrest in SCNT embryos.

It is known that ZGA is governed by a time-dependent 'zygotic clock' (i.e. 24h post-fertilization),

and Dux is expressed as a brief pulse in early 2-cell stage (De Iaco et al., 2019; Eckersley-

Maslin et al., 2019; Rodriguez-Terrones et al., 2018). Therefore, we wonder whether Dux is

time-depend on improving the efficiency of SCNT embryo development. To answer this

question, we generated transgenic mouse lines containing doxycycline (dox) inducible Dux

gene (supplement Figure S2A-B). By switching the culture medium between dox-containing

and -lacking at various time points, the critical period for the dox-Dux requirement was mapped

to a 24-h window (Figure 1F). When the dox was supplied in this window, the SCNT blastocyst

formation rate was significantly increased (Figure 1G; supplement Table S2). Subsequently,

we performed single-cell RNA sequencing (scRNA-seq) to examine the possible changes of

the ZGA related gene expressions in Dux-overexpressed 2-cell SCNT embryos (D-SCNT).

Unsupervised hierarchal clustering revealed that the ICSI and D-SCNT generated embryos in

the same cluster, and ZGA-genes were upregulated in D-SCNT embryos (Figure 1H and I;

supplement Figure S3A-D; Dataset S1, S2). These results indicating the improvement of Dux

on SCNT embryonic development depends on its treatment time.

The SCNT blastocyst rate was improved by injecting Kdm4d mRNA (K-SCNT) (Matoba et al.,

2014). We next compared the developmental potential of SCNT embryos derived by K-SCNT

and D-SCNT. We found that the blastocyst formation rate of D-SCNT was similar to K-SCNT

(84.72% *vs.* 86.79%; supplement Figure S4A, Table S2). Nonetheless, the cloned pup birth

rate of D-SCNT was higher than K-SCNT (10.71% *vs.* 7.81%; Figure 1J; supplement Table

S3). Despite D-SCNT or K-SCNT, the abnormally large placentas are still observed in those

cloned pups (supplement Figure S4B, Table S3). Previous studies have indicated that knock-

down DNA-methyltransferases Dnmt3a/b led to less placental abnormalities (Si-SCNT) (R.

Gao et al., 2018). We next investigated whether combined D-SCNT and Si-SCNT could further

improve SCNT embryonic development. Compared with canonical SCNT, combined K-SCNT

and Si-SCNT improved the pup birth-rate from 1.01% to 12.24%, but less than D-SCNT

combined with Si-SCNT (18.60%; Figure 1J). Genotyping analysis confirmed that all the

cloned pups were generated from donor cells (supplement Figure S4C). The large placental

phenotype was rescued by the D-Si-SCNT combined approach (supplement Figure S4B).

Deng et al. identified a unique embryonic 2-cell like status as the key molecular event that led

to successful chemical reprogramming (CiPS) (Zhao et al., 2018). Thus, we tried to improve

the CiPS by overexpression of ZGA inducer Dux (D-CiPS). The dox-inducible mouse

embryonic fibroblasts (dDux-MEFs) cells were isolated from the transgenic mice. As shown in

Figure 2A, we treated the dDux-MEFs with dox for various durations, and observed that Dux

displayed remarkable effect at the stage-2 of CiPS. About 36 alkaline phosphatase-positive

(AP+) colonies could be observed in D-CiPS wells (starting from 50,000 MEFs per well of 6-

well plate), while only 26 AP+ colonies could be observed in the canonical CiPS well (Figure

2B and C). Consistently, the ZGA related genes were significantly upregulated after transient

expression of Dux (Figure 2D). These D-CiPS induced cells were pluripotent as demonstrated

by standard characterization procedures (Figure 2E-G). We found, based on bisulfite

sequencing, that successful demethylation occurred in the promoters of Oct4 and Nanog in

the D-CiPS induced cell lines (Figure 2H).

When we prepare this manuscript, two recent studies suggested that Dux is important but not

essential for fertilized embryo development by enhancing rather than ZGA inducer (Chen &

Zhang, 2019; Guo et al., 2019). However, we demonstrate here that transient overexpression

of ZGA-inducer Dux not only improved SCNT efficiency, but also increased the efficiency of

chemical reprogramming. Ectopic expression of Dppa2/4 in SCNT embryos resulted in 1-cell

block, which implied that successful timely ZGA is essential for the development of SCNT-

derived embryos. A recent study found that Dppa2/4 interacted with small ubiquitin-like

modifier (SUMO) related proteins (Yan et al., 2019). Therefore, we speculated that

overexpression of Dppa2/4 would accelerate the degradation of reprogramming factors in the

oocyte-cytoplasm, which leads to reprogramming failure. Nonetheless, fertilized embryos

completely differ from SCNT-derived embryos. SCNT embryos are generated by directly

injecting somatic nuclei into enucleated oocytes, whereas the oocyte cytoplasm is

evolutionally designed to reprogram the spermatozoa. Therefore, although Dux and Dppa2/4

as ZGA-inducers, its effects on the reprogramming process likely differ between sperm and

somatic cells. Thus, the relationship among Dux, Dppa2/4, and ZGA in SCNT warrants further

investigation.

Overall, our study revealed that transient overexpression of Dux not only improved SCNT

efficiency, but also increased the efficiency of chemical reprogramming. The Dux was

identified as a ZGA-inducer in mouse, human, and likely all mammalian animals (Iturbide &

Torres-Padilla, 2017; Whiddon et al., 2017). Thus, we reasoned that transient overexpression

of Dux could also improves reprogramming efficiency of all mammals.

Materials and methods

Animals and Chemicals

Specific pathogen-free (SPF)-grade BDF1 (C57BL/6N × DBA/2), CD1, and Kun-Ming (KM) mice mice were purchased from Laboratory Animal Research Center (Inner Mongolia University) or Vital River Laboratories (Beijing, China). All animal experiments were approved by the Animal Care and Use Committee of Inner Mongolia University. All procedures were carried out in strict accordance with the recommendations made in the Guide for the Care and Use of Laboratory Animals of the National Veterinary and Quarantine Service. MERLV:: tdTomato transgenic mice, which carry a transgenic MERLV promoter LTR driven tdTomato reporter (Yang et al., 2018), were mated with the same positive mice. dDux-MEFs cells were isolated from E12.5 fetus. The MERLV:: tdTomato embryos used in the experiment were produced by MERVL::tdTomato sperm and MII oocytes from the littermates of transgenic mice. All chemicals used in this study were purchased from Sigma (USA), unless otherwise indicated.

SCNT and embryo culture

The SCNT processes followed previously published studies (Kishigami et al., 2006; Yang et al., 2018). Briefly, groups of ~50 MII oocytes were transferred to a chamber containing oil-covered M2 supplemented with 5 µg/ml cytochalasin B (CB). The spindle-chromosome complex (SCC) was removed by a blunt Piezo-driven pipette (~10 µm internal diameter) on a 37°C heating stage of an inverted microscope (Nikon, Japan). The nuclei of donor cells were drawn in and out of the injection pipette until its plasma membrane was broken and was then injected into enucleated oocytes. The reconstructed embryos were cultured in aMEM medium (Thermo, USA) containing 10% fetal calf serum (FCS; Hyclone, USA) for 1 h before activation treatment. The reconstructed embryos were activated in Ca_{2+} free KSOM medium containing 10 mM strontium and 5 µg/ml CB for 6 h. Activated embryos were thoroughly washed and cultured in G1/G2 medium (1:1, vol / vol; Vitrolife, Sweden). Induction of Dux was performed by administration of doxycycline (2 µg/ml) in the G1/G2 medium.

Immunofluorescence staining

Embryos and CiPS cells were rinsed three times in phosphate-buffered saline (PBS) with 0.3% BSA, fixed with 4% paraformaldehyde (PFA) overnight at 4°C, and then permeabilized with 0.2% (vol / vol) Triton X-100 for 15 min at room temperature, followed by washing thoroughly in PBS containing 0.3% BSA. Fixed samples were blocked in 0.05% Tween 20 in PBS containing 3% BSA (PBST) at 37°C for 1 h and then incubated with the primary antibodies overnight at 4°C. Samples were incubated with primary antibodies against HA (Santa Cruz, sc-7392, USA), Oct4 (Santa Cruz, sc-8629, USA), and Ssea1 (Santa Cruz, sc-21702, USA). After incubating, the samples were needed to wash several times in PBST and then incubated with appropriate secondary antibodies conjugated with Alexa Fluor 594 and Alexa Fluor 488 (Thermo, USA) for 1 h at 37°C. For imaging, the embryos were mounted in 10 µl anti-fade solution with DAPI (Thermo, USA) and compressed with a cover-slip. All samples were observed by laser scanning microscope (Nikon, Japan).

***In vitro* mRNA synthesis and microinjection in oocytes**

The *Dux*, *Dppa2*, and *Dppa4* mRNA were cloned into T7-driven vectors, and mRNAs were synthetized *in vitro* using mMESSAGE mMACHINE T7 Ultra Kit (Thermo, USA) following the manufacturer's instructions. The integrity of manufactured mRNA was confirmed by electrophoresis with formaldehyde gels. The final concentration of mRNA was diluted to 900 ng/ml before injection. As previously described (Yang et al., 2018), 8 pl of mRNA was microinjected into the cytoplasm of denuded oocytes. These oocytes were obtained by superovulating BDF1 female mice by intraperitoneal injection of 10 IU pregnant mare serum gonadotropin (PMSG; Sansheng, China) and 10 IU human chorionic gonadotropin (hCG; Sansheng, China) 48 h apart. The cumulus-oocyte complexes (COCs) were collected from oviducts 14 h post-hCG, and the cumulus cells were dispersed by EmbryoMax FHM Mouse Embryo Media (Millipore, USA). Oocytes were injected using Piezo-operated blunt-end micropipette (3~5 µm internal diameter). After injection, oocytes were kept at room-

209   temperature for 30 min and then moved into the incubator. The siRNA information was

210   presented in Table S4.

Generation of CiPS cells

This section was adapted from Deng's lab (Zhao et al., 2018). Reagents setup: Small

molecules: VPA, CHIR99021, 616452, Tranylcypromine, Forskolin, Ch55, EPZ004777,

DZNep, Decitabine, SGC0946, and PD0325901. Stage I medium: FBS/KSR-based medium

supplemented with the small-molecule cocktail VC6TF5E (100 µM VPA, 40 µM CHIR99021,

10 µM 616452, 5 µM Tranylcypromine, 10 µM Forskolin, 1 µM Ch55, and 5 µM EPZ004777).

N2B27-SII medium: N2B27-based medium supplemented with 10 ng/ml LIF, 50 µg/ml vitamin

C, 25 ng/ml bFGF, 2 mg/ml Albumax-II, and the small-molecule cocktail VC6TF5ZDS (VPA at

1 mM). Stage III medium: N2B27-based medium with 3 µM CHIR99021, 1 µM PD0325901,

10 ng/ml LIF, and 50 µg/ml vitamin C. On day –1, MEFs were seeded at 50,000 cells per well

of 6-well plate with MEF culture medium. The next day (day 0), change the medium into

optimized stage I medium. From day 6~12, cells were cultured in optimized N2B27-SII medium.

During day 12~16, cells were cultured in stage III medium with 500 µM VPA. On day 16, VPA

was removed and cells were cultured in stage III medium. After another 4~10 days, CiPS

colonies emerged. Induction of Dux was performed by administration of doxycycline (2 µg/ml)

in the culture medium.

Transgenic mice generation

The pCW57.1-mDux-CA vector was a gift from Stephen Tapscott (Addgene 99284). The

pronuclear microinjection for the production of transgenic mice followed previously published

studies (Ittner & Gotz, 2007). Briefly, the vector was injected into the well-recognized pronuclei.

Injected zygotes were transferred into pseudopregnant female mice (~ 30 zygotes per mouse)

after 4 h recovery culture in KSOM-AA medium (Millipore, USA). For founder identification, tail

tips were subjected to standard DNA extraction procedures. The amplified DNA fragments

were subjected to TA cloning and sequencing. The founder mice were crossed to the

littermates of founder mice for four generations to produce homozygous Dux mice.

Alkaline phosphatase staining

For alkaline phosphatase staining, CiPS cells were fixed with 4% PFA in PBS for 2 min, rinsed

once with PBS and detection was performed using a Alkaline Phosphatase Assay Kit

(VECTOR, USA) according to the manufacturer's protocol.

Real-time qPCR analysis

Total RNA was isolated using TRIzol reagent (Thermo, USA) and was immediately reverse-

transcribed using a Prime Script RT reagent kit (Takara, Japan). The reverse transcription

PCR was were amplified using Ex Taq (Takara, Japan). The qPCR was performed using a

SYBR Premix Ex Taq (Takara, Japan) and signals were detected with Applied Biosystems

7500 real-time PCR System (Thermo, USA). Relative mRNA expression was calculated use

the $2_{(-\Delta\Delta Ct)}$ method. The primer information was presented in Table S5.

Chimeric mice and embryo transfer

For chimeric experiments, CiPS cells were used one day before passaging, which showed an

optimal undifferentiated morphology. The CiPS cells were microinjected into CD1/KM

blastocysts using a Piezo (Primetech, Japan) microinjection pipette. After culturing for 3 h, the

embryos were transplanted into the uterus of pseudopregnant mice (~15 embryos per mouse).

The 2-cell stage SCNT embryos were transferred to the oviducts of E0.5 pseudopregnant (~20

embryos per mouse). The cloned pups nursed by lactating CD1/KM females. SSLP analysis

was performed for D6Mit15 and D2Mit102. The primer information is presented in Table S5.

Teratoma formation

Teratoma formation analysis was carried out to evaluate the pluripotency of CiPS cells.

Approximately 10_6 CiPS cells were injected subcutaneously into the hind limbs of a 8-week-

old nude mice. After 6 weeks, fully formed teratomas were dissected and fixed with PBS

containing 4% PFA, then embedded in paraffin, sectioned and stained with haematoxylin and

eosin for histological analysis.

Bisulfite genomic sequencing

Genomic DNA was extracted by the tissue & blood DNA extraction kit (Qiagen, Germany) and

treated with the Methylamp DNA modification kit (Epigentek, USA). The bisulphite conversion

was performed as previously published studies (Cao et al., 2018).

RNA-sequencing

The single-cell RNA-seq method followed previously published studies (W. Liu et al., 2016;

Tang et al., 2010). The reverse transcription was performed directly on the cytoplasmic lysate

of individual embryos. The total cDNA library was then amplified by 18~20 cycles for library

construction, which was performed following manufacturer's instructions (Illumina, USA).

Paired-end sequencing was further performed at Annoroad (Beijing, China). Three biological

replicates were analyzed for each treatment condition. After removing low-quality reads and

adapters, the raw reads were mapped to the mm9 genome using Tophat (v1.3.3) with default

parameters (Trapnell, Pachter, & Salzberg, 2009). Expression levels for all RefSeq transcripts

were quantified to fragments per kilobase of exon model per million mapped reads (FPKM)

using Cufflinks (v1.2.0) (Trapnell et al., 2010). Gene Ontology (GO) analyses for differentially

expressed genes were performed by using the R package GO.db and Database for Annotation,

Visualization and Integrated Discovery (DAVID) (Huang da, Sherman, & Lempicki, 2009).

Statistics analysis and data availability

Statistical analyses were done using the univariate analysis of variance (ANOVA) followed by

the Student t-test with SPSS 21.0 statistical software (Armonk, USA). $P < 0.05$ was considered

significant. Sequencing data have been deposited in the NCBI sequence read archive (SRA)

under accession codes: SCNT-1 (SAMN12871322), SCNT-2 (SAMN12871323), SCNT-3

(SAMN12871324); D-SCNT-1 (SAMN12871325), D-SCNT-2 (SAMN12871326), D-SCNT-3

(SAMN12871327);   ICSI-1   (SAMN12871328),   ICSI-2   (SAMN12871329),   ICSI-3

(SAMN12871330).

Acknowledgements

We thank Prof. Shaorong Gao (Tongji University), Prof. Zhiming Han (Chinese Academy of

Sciences), and Prof. Qing Xia (Peking University) for their technical assistance. We also thank

Prof. Yi Zhang (Harvard Medical School) for sharing the list of RRRs. This study was supported

by the Genetically Modified Organisms Breeding Major Projects (2016ZX08007-002), the

opening project of State Key Laboratory of R2BGL (to L.Y.), the Inner Mongolia University

Chief Scientist Program (to G.L.), the Inner Mongolia Autonomous Region Basic Research

Project (to G.L.).

Author contributions

L.Y. and G.L. conceived and designed the study. L.Y., X.L., L.S., A.D., G.S., C.B., and Z.W.

performed the experiments. L.Y., X.L., L.S., A.D., and G.S. analyzed the data. G.L. supervised

the project. L.Y. and G.L. wrote the paper.

Competing interests

The authors declare no competing interests.

References

Ancelin, K., Syx, L., Borensztein, M., Ranisavljevic, N., Vassilev, I., Briseno-Roa, L., . . . Heard, E. (2016). Maternal LSD1/KDM1A is an essential regulator of chromatin and transcription landscapes during zygotic genome activation. *Elife, 5*, e08851. doi:10.7554/eLife.08851

Cao, S., Yu, S., Li, D., Ye, J., Yang, X., Li, C., . . . Pei, D. (2018). Chromatin accessibility dynamics during chemical induction of pluripotency. *Cell Stem Cell, 22*(4), 529-542. doi:10.1016/j.stem.2018.03.005

Chen, Z., & Zhang, Y. (2019). Loss of DUX causes minor defects in zygotic genome activation and is compatible with mouse development. *Nat Genet, 51*(6), 947-951. doi:10.1038/s41588-019-0418-7

Choi, Y. J., Lin, C. P., Risso, D., Chen, S., Kim, T. A., Tan, M. H., . . . He, L. (2017). Deficiency of microRNA miR-34a expands cell fate potential in pluripotent stem cells. *Science, 355*(6325), eaag1927. doi:10.1126/science.aag1927

De Iaco, A., Coudray, A., Duc, J., & Trono, D. (2019). DPPA2 and DPPA4 are necessary to establish a 2C-like state in mouse embryonic stem cells. *EMBO Rep, 20*(5), e47382. doi:10.15252/embr.201847382

De Iaco, A., Planet, E., Coluccio, A., Verp, S., Duc, J., & Trono, D. (2017). DUX-family transcription factors regulate zygotic genome activation in placental mammals. *Nat Genet, 49*(6), 941-945. doi:10.1038/ng.3858

Eckersley-Maslin, M., Alda-Catalinas, C., Blotenburg, M., Kreibich, E., Krueger, C., & Reik, W. (2019). Dppa2 and Dppa4 directly regulate the Dux-driven zygotic transcriptional program. *Genes Dev, 33*(3-4), 194-208. doi:10.1101/gad.321174.118

Gao, R., Wang, C., Gao, Y., Xiu, W., Chen, J., Kou, X., . . . Gao, S. (2018). Inhibition of aberrant DNA re-methylation improves post-implantation development of somatic cell nuclear transfer embryos. *Cell Stem Cell, 23*(3), 426-435. doi:10.1016/j.stem.2018.07.017

Gao, T., Zheng, J., Xing, F., Fang, H., Sun, F., Yan, A., . . . Sheng, H. Z. (2007). Nuclear reprogramming: the strategy used in normal development is also used in somatic cell nuclear transfer and parthenogenesis. *Cell Res, 17*(2), 135-150. doi:10.1038/cr.2007.2

Guo, M., Zhang, Y., Zhou, J., Bi, Y., Xu, J., Xu, C., . . . Wang, Y. (2019). Precise temporal regulation of Dux is important for embryo development. *Cell Res*, Published online 2019-Oct-7. doi:10.1038/s41422-019-0238-4

Hendrickson, P. G., Dorais, J. A., Grow, E. J., Whiddon, J. L., Lim, J. W., Wike, C. L., . . .

Cairns, B. R. (2017). Conserved roles of mouse DUX and human DUX4 in activating cleavage-stage genes and MERVL/HERVL retrotransposons. *Nat Genet, 49*(6), 925-934. doi:10.1038/ng.3844

Huang da, W., Sherman, B. T., & Lempicki, R. A. (2009). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc, 4*(1), 44-57. doi:10.1038/nprot.2008.211

Ishiuchi, T., Enriquez-Gasca, R., Mizutani, E., Boskovic, A., Ziegler-Birling, C., Rodriguez-Terrones, D., . . . Torres-Padilla, M. E. (2015). Early embryonic-like cells are induced by downregulating replication-dependent chromatin assembly. *Nat Struct Mol Biol, 22*(9), 662-671. doi:10.1038/nsmb.3066

Ittner, L. M., & Gotz, J. (2007). Pronuclear injection for the production of transgenic mice. *Nat Protoc, 2*(5), 1206-1215. doi:10.1038/nprot.2007.145

Iturbide, A., & Torres-Padilla, M. E. (2017). Starting embryonic transcription for the first time. *Nat Genet, 49*(6), 820-821. doi:10.1038/ng.3880

Kishigami, S., Wakayama, S., Thuan, N. V., Ohta, H., Mizutani, E., Hikichi, T., . . . Wakayama, T. (2006). Production of cloned mice by somatic cell nuclear transfer. *Nat Protoc, 1*(1), 125-138. doi:10.1038/nprot.2006.21

Liu, W., Liu, X., Wang, C., Gao, Y., Gao, R., Kou, X., . . . Gao, S. (2016). Identification of key factors conquering developmental arrest of somatic cell cloned embryos by combining embryo biopsy and single-cell sequencing. *Cell Discov, 2*, 16010. doi:10.1038/celldisc.2016.10

Liu, Z., Cai, Y., Wang, Y., Nie, Y., Zhang, C., Xu, Y., . . . Sun, Q. (2018). Cloning of macaque monkeys by somatic cell nuclear transfer. *Cell, 172*(4), 881-887. doi:10.1016/j.cell.2018.01.020

Lu, F., & Zhang, Y. (2015). Cell totipotency: molecular features, induction, and maintenance. *Natl Sci Rev, 2*(2), 217-225. doi:10.1093/nsr/nwv009

Macfarlan, T. S., Gifford, W. D., Driscoll, S., Lettieri, K., Rowe, H. M., Bonanomi, D., . . . Pfaff, S. L. (2012). Embryonic stem cell potency fluctuates with endogenous retrovirus activity. *Nature, 487*(7405), 57-63. doi:10.1038/nature11244

Matoba, S., Liu, Y., Lu, F., Iwabuchi, K. A., Shen, L., Inoue, A., & Zhang, Y. (2014). Embryonic development following somatic cell nuclear transfer impeded by persisting histone methylation. *Cell, 159*(4), 884-895. doi:10.1016/j.cell.2014.09.055

Matoba, S., & Zhang, Y. (2018). Somatic cell nuclear transfer reprogramming: mechanisms

and applications. *Cell Stem Cell, 23*(4), 471-485. doi:10.1016/j.stem.2018.06.018

Percharde, M., Lin, C. J., Yin, Y., Guan, J., Peixoto, G. A., Bulut-Karslioglu, A., . . . Ramalho-

Santos, M. (2018). A LINE1-nucleolin partnership regulates early development and

ESC identity. *Cell, 174*(2), 391-405. doi:10.1016/j.cell.2018.05.043

Rodriguez-Terrones, D., Gaume, X., Ishiuchi, T., Weiss, A., Kopp, A., Kruse, K., . . . Torres-

Padilla, M. E. (2018). A molecular roadmap for the emergence of early-embryonic-like

cells in culture. *Nat Genet, 50*(1), 106-119. doi:10.1038/s41588-017-0016-5

Tang, F., Barbacioru, C., Nordman, E., Li, B., Xu, N., Bashkirov, V. I., . . . Surani, M. A. (2010).

RNA-Seq analysis to capture the transcriptome landscape of a single cell. *Nat Protoc,*

5(3), 516-535. doi:10.1038/nprot.2009.236

Trapnell, C., Pachter, L., & Salzberg, S. L. (2009). TopHat: discovering splice junctions with

RNA-seq. *Bioinformatics, 25*(9), 1105-1111. doi:10.1093/bioinformatics/btp120

Trapnell, C., Williams, B. A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M. J., . . . Pachter,

L. (2010). Transcript assembly and quantification by RNA-seq reveals unannotated

transcripts and isoform switching during cell differentiation. *Nat Biotechnol, 28*(5), 511-

515. doi:10.1038/nbt.1621

Wang, Y., Zhao, C., Hou, Z., Yang, Y., Bi, Y., Wang, H., . . . Gao, S. (2018). Unique molecular

events during reprogramming of human somatic cells to induced pluripotent stem cells

(iPSCs) at naive state. *Elife, 7*, e29518. doi:10.7554/eLife.29518

Wasson, J. A., Simon, A. K., Myrick, D. A., Wolf, G., Driscoll, S., Pfaff, S. L., . . . Katz, D. J.

(2016). Maternally provided LSD1/KDM1A enables the maternal-to-zygotic transition

and prevents defects that manifest postnatally. *Elife, 5*, e08848.

doi:10.7554/eLife.08848

Whiddon, J. L., Langford, A. T., Wong, C. J., Zhong, J. W., & Tapscott, S. J. (2017).

Conservation and innovation in the DUX4-family gene network. *Nat Genet, 49*(6), 935-

940. doi:10.1038/ng.3846

Wilmut, I., Schnieke, A. E., McWhir, J., Kind, A. J., & Campbell, K. H. (1997). Viable offspring

derived from fetal and adult mammalian cells. *Nature, 385*(6619), 810-813.

doi:10.1038/385810a0

Yan, Y. L., Zhang, C., Hao, J., Wang, X. L., Ming, J., Mi, L., . . . Wang, Y. (2019). DPPA2/4 and

SUMO E3 ligase PIAS4 opposingly regulate zygotic transcriptional program. *PLoS Biol,*

17(6), e3000324. doi:10.1371/journal.pbio.3000324

Yang, L., Song, L., Liu, X., Bai, L., & Li, G. (2018). KDM6A and KDM6B play contrasting roles

in nuclear transfer embryos revealed by MERVL reporter system. *EMBO Rep, 19*(12),

e46240. doi:10.15252/embr.201846240

Zhao, T., Fu, Y., Zhu, J., Liu, Y., Zhang, Q., Yi, Z., . . . Deng, H. (2018). Single-cell RNA-seq

reveals dynamic early embryonic-like programs during chemical reprogramming. *Cell*

Stem Cell, 23(1), 31-45. doi:10.1016/j.stem.2018.05.025

Figure legends

Figure 1. Dux act as promote factors for SCNT-mediated somatic cell reprogramming.

(A) Schematic illustration of the screen strategy.

(B) Quantification of embryos that expression ZGA reporter (MERVL::tdTomato) after injection with siRNA-repressor or mRNA- inducer. N, total number of embryos analyzed for each condition. Error bars, mean ± S.D.; three independent experiment replicates were performed.

(C) Representative fluorescence image of SCNT embryos derived by different methods. The SCNT embryos produced by transfer of MERVL::tdTomato cumulus cell into WT enucleated oocytes. Scale bar, 100 µm.

(D) Stacked bar plots show fraction of embryos at blastocyst stages after injection with different mRNA as indicated. N, total number of embryos analyzed for each condition.

(E) Representative live-cell images of dynamics ZGA reporter expression during SCNT embryos development. The time after starting observation is shown on the bottom of image. Scale bar, 50 µm.

(F) Schematic transient induction of Dux. All timings reported in this paper is hour post-activation.

(G) The preimplantation development of SCNT embryos. Shown is the percentage of embryos that reaches each indicated stage. Scale bar, 100 µm.

(H) Unsupervised hierarchical clustering.

(I) Venn diagram showing the overlap between the genes that failed to be activated in SCNT 2-cell embryos and derepressed in D-SCNT. Heatmap, GO terms showing the expression pattern of 257 overlap genes.

(J) The birth rate of SCNT embryos derived by different methods.

Figure 2. Dux facilitates chemical-mediated somatic cell reprogramming.

(A) Durations of dox addition at different time points during chemical induction (left) and assessment of AP+ colonies (right). Error bars, mean ± S.D.

(B) Morphological changes at distinct time points during D-CiPS. Scale bar, 100 μm.

(C) Representative images of the number of AP+ positive colonies.

(D) Relative expression levels by qPCR of select ZGA genes in D-CiPS, compared with

canonical CiPS. Error bars, S.E.M.; n = 3; ***P < 0.001 by Student's t-test.

(E) Immunostaining images of D-CiPS expressed pluripotency markers. Scale bar, 50 μm.

(F) Histology by H&E staining of teratoma tissues derived by D-CiPS cells. Scale bar, 200 μm.

(G) Chimeric mouse generated from D-CiPS cells.

(H) Bisulfite sequencing analysis of Oct4 and Nanog demethylation in D-CiPS or canonical

CiPS cells (filled squares are methylated CpGs and empty ones indicate unmethylated).

Fig.1

Fig.2