A CellAge epigenetic clock for expedited discovery of anti-ageing compounds *in vitro*

Celia Lujan[1,*], Eleanor J. Tyler[2,*], Amy P. Webster[3,*], Eleanor R. Stead[1], Victoria E. Martinez

Miguel[1], Simone Ecker[3], Deborah Milligan[2], James C. Garbe[4], Martha R. Stampfer[4], Stephan Beck[3,#],

Robert Lowe[5,#], Cleo L. Bishop[2,#], Ivana Bjedov[1,#]

Affiliations

[1]: UCL Cancer Institute, Paul O'Gorman Building, University College London, 72 Huntley Street London, London WC1E 6DD, UK
[2]: Centre for Cell Biology and Cutaneous Research, Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, 4 Newark Street, London E1 2AT
[3]: Medical Genomics Group, UCL Cancer Institute, University College London, London WC1E 6BT, UK
[4]: Biological Systems and Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA, United States
[5]: Centre for Genomics and Child Health, Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, 4 Newark Street, London E1 2AT

*Equal contribution
#corresponding authors :
 i.bjedov@ucl.ac.uk c.l.bishop@qmul.ac.uk rlowe.compbio@gmail.com s.beck@ucl.ac.uk

Abstract

We aim to improve anti-ageing drug discovery, currently achieved through laborious and lengthy longevity analysis. Recent studies demonstrated that the most accurate molecular method to measure human age is based on CpG methylation profiles, as exemplified by several epigenetics clocks that can accurately predict an individual's age. Here, we developed CellAge, a new epigenetic clock that measures subtle ageing changes in primary human cells *in vitro*. As such, it provides a unique tool to measure the effects of relatively short pharmacological treatments on ageing. We validated our CellAge clock against known longevity drugs such as rapamycin and trametinib. Moreover, we uncovered novel anti-ageing drugs, torin2 and Dactolisib (BEZ-235), demonstrating the value of our approach as a screening and discovery platform for anti-ageing strategies. CellAge outperforms other epigenetic clocks in measuring subtle ageing changes in primary human cells in culture. The tested drug treatments reduced senescence and other ageing markers, further consolidating our approach as a screening platform. Finally, we showed that the novel anti-ageing drugs we uncovered *in vitro,* indeed increased longevity *in vivo*. Our method expands the scope of CpG methylation profiling from measuring human chronological and biological age from human samples in years, to accurately and rapidly detecting anti-ageing potential of drugs using human cells *in vitro*, providing a novel accelerated discovery platform to test sought after geroprotectors.

One of the remarkable achievements of developed countries is a continuous increase in life expectancy at birth, leading to greater longevity. However, a higher proportion of elderly in modern societies is accompanied by a steep increase in people suffering from age-related diseases. For example, cancer incidence rates, currently at 17 million worldwide, are expected to increase to 26 million in 2040[1], and a similar rise is expected for Alzheimer's and Parkinson's disease[2]. Compression of late-life morbidity is, therefore, a priority to alleviate suffering in the elderly[3] and to reduce a growing economic burden to society[4].

Critically, seminal discoveries in the biology of ageing showed that ageing is a malleable process and that down-regulation of major cellular nutrient signalling pathways, either glucose-sensing insulin signalling or amino acid-sensing target-of-rapamycin signalling, results in longevity and health improvement in all model organisms tested from yeast to mammals[5]. For instance, the long-lived mutants in *C. elegans* are protected from tumorous cell proliferation[6] and have reduced toxic protein aggregation[7], while *Drosophila* show less deterioration in their hearts[8]. Long-lived mouse mutants are protected from osteoporosis, cataracts and skin pathology, as well as decline in glucose homeostasis, immune and motor function[9]. The effect of these mutations is conserved from yeast to mammals, and it is, therefore, expected that if drugs replicate the biological impact of these changes, this could improve health in the elderly and prevent age-related disease. It is increasingly recognised that directly targeting ageing through pharmacological interventions, as opposed to specific age-related diseases, is a highly promising strategy for broad-spectrum disease protection[10]. However, at present, there are only a handful of reliable anti-ageing drugs whose effects have been confirmed in mammals, such as rapamycin[11] and metformin[12]. Crucially, there are currently no sufficiently reliable ageing biomarkers for testing drugs on human cells *in vitro,* and the development of a specialised epigenetic clock seems the most promising current approach[13,14,15].

To accelerate the discovery workflow for anti-ageing drugs, we took advantage of the breakthrough in the ageing field which showed that epigenetic clocks provide the most accurate measurements of human age, for instance, the approximate error rate for the Skin and Blood clock is ±2.5 years (maximal correlation coefficient 0,98)[13]. Epigenetic clocks surpass the accuracy of other ageing biomarkers such as telomere length and those based on transcriptomic, metabolomics or proteomic approaches, potentially because the latter approaches detect more transient and less stable cellular changes[16]. Ageing is accompanied by overall CpG hypomethylation, whilst some CpG islands and gene regions become hypermethylated[17]. Remarkably, only a small selection of the 56 million CpG sites in the diploid human genome, coupled with computational algorithms, is sufficient to provide an accurate readout of human age. One of the first epigenetic clocks was developed by Hannum using just 71 CpG sites to estimate age from blood samples[18], while Horvath's multi-tissue age estimator[16] and Skin and Blood clock[13] use 353 and 391 CpG sites, respectively[19,20]. Even a single CpG site in the ELOVL2 gene is sufficient to determine age[21], albeit clocks using only a few CpG sites are less accurate and less applicable to different tissues[19]. The epigenetic clocks measure the ageing

process inherent to all our cells and tissues, irrespective of their proliferation rate[14]. As the human
epigenome reflects physiological changes, epigenetic clocks cannot only predict chronological age
from a human sample but also give an estimate of biological age as has widely been demonstrated
by the associations of epigenetic age with morbidity and mortality[19,22]. Recently, valuable predictors
focussing on this aspect have been developed: PhenoAge[23] and GrimAge[24], which form the best
epigenetic morbidity and mortality predictors available to date.

DNA methylation also captures information on the approximate number of cell divisions a stem cell
has been through, as has been shown by epiTOC[25], a mitotic clock that approximates stem cell
divisions and correlates with cancer risk[26]. The biology underlying CpG methylation alterations at the
sites linked to ageing clocks is not well understood. Horvath suggests that it is linked to epigenetic
maintenance programmes being reflected in DNA methylation alterations[16,19]. Some recent findings
implicate loss of H3K36 histone methyltransferase NSD1 in epigenetic ageing clock acceleration [27].
Despite the enigma regarding epigenetic clock mechanism, these clocks are extremely useful and
reliable predictors of age. However, little is known so far about the performance of these clocks in *in*
vitro ageing experiments. It has recently been shown that the rate of epigenetic ageing in cultured
cells is significantly faster than in the human body[14,28] and that epigenetic age is retarded by
rapamycin *in vitro*[14], but neither of the clocks specialised for *in vitro* drug discovery nor were they
tested on multiple anti-ageing drugs.

Therefore, we aimed to exploit the exceptional accuracy of CpG methylation clocks to uncover new
anti-ageing pharmacological treatments. The current gold standard for discovering novel anti-ageing
drugs are longevity experiments, which are laborious, lengthy and expensive. For instance, in mice,
they take three years, thereby precluding any large scale drug screens. Existing screens in *C.*
elegans commonly use live *E. coli* as food[29,30], which is a disadvantage as drugs are metabolised
first by the bacteria making their effect on worms secondary, which may lead to confounded
results[31,32]. Yeast drug screens lack the crucial aspect of tissue toxicity[33]. In addition, all longevity
assays require constant supply of the drug, making them highly expensive. Other attempts to
uncover anti-ageing effects of drugs are based on computational analysis using existing
transcriptomic information on the ageing process combined with drug characteristics[34]. However,
transcriptomic changes are more transient and noisy when compared to DNA methylation and are,
therefore, a less consistent ageing marker[19].

We tested if existing epigenetic clocks could be used to measure anti-ageing drug potential in human
primary cells *in vitro* and if we could build a new clock specialised for this purpose. Senescence is
tightly associated with ageing of the organism, and because of the pronounced resemblance of
ageing in primary cells *in vitro* to ageing *in vivo,* together with the evidence that human DNA
methylation signatures are conserved and accelerated in cultured fibroblasts[28], we used cultured
human cells as a proxy for human ageing[14,35]. The ability to test anti-ageing drug properties directly
on human cells *in vitro* could considerably accelerate the discovery of new compounds promoting

healthy ageing. To this end, we used normal human mammary fibroblasts (HMFs) from a healthy 21-
year old donor that we cultured from passage 10 to passage 20, which is before these cells reach
senescence at passage 29 (Supplementary Fig. 1a-d). To measure CpG methylation, we used EPIC
Arrays (Illumina) that measure methylation at 850,000 sites.

First, we tested the three most suitable existing epigenetic clocks, to determine if they could detect
weekly and monthly ageing differences occurring during serial passaging of HMFs (Fig. 1a). The
Multi-tissue clock[16] consistently predicted a higher epigenetic age, and at passage ten this was
43.6±1.0 years (Fig. 1a), consistent with what was recently reported[28]. This increased age estimate,
compared to 16 years of the donor, is in accordance with published data demonstrating that this
epigenetic clock overestimates the age of mammary tissue samples[16]. The PhenoAge clock[23],
developed to predict mortality and morbidity risks, reported the epigenetic age of the donor to be
3.5±1.1 years younger (Fig. 1a). The most accurate age estimate, predicting the age of the donor at
23.2±0.87 years, was obtained using the Skin and Blood clock, which is specialised for determining
donor age of the cells in culture and of easily accessible human tissues (Fig. 1a). The Multi-tissue
clock and Skin and Blood clock showed a small increase in age with progressive passaging (from
passage 10 to 20, age estimate increased from 43.6±1.0 to 53.9±1.7 and from 23.2±0.87 to 31.6±1.2
years, respectively), whilst this increase was greater for the PhenoAge clock (from 3.5±1.1 to
26.6±9.7 years). This suggests that, of the tested clocks, the PhenoAge clock is most suitable to
measure ageing *in vitro* (Fig. 1a). However, the PhenoAge clock showed substantial variability in
predictions for higher passages, which would obstruct the detection of subtle ageing differences
upon anti-ageing drug treatments. In conclusion, while the Skin and Blood clock[13] measures
fibroblast ageing in culture, none of the existing clocks was ideally suited to accurately measure
subtle anti-ageing drug potential in human primary cells *in vitro,* and similar comparisons have
recently been reported by others[14,28].

This prompted us to develop a new clock that, rather than predicting donor age in years, specialises
in measuring methylation changes occurring during ageing of primary cells in culture and could
differentiate DNA methylation state between each passage. To this end, we developed a clock using
two different cell types, the above-mentioned HMFs and human dermal fibroblasts (HDFs), which
were obtained from a different donor, have a different proliferative lifespan *in vitro*, and a different
rate of DNA methylation change. Like the HMFs, the HDFs were serially passaged and sampled
every other passage for DNA methylation analysis (Fig. 1b). We filtered for significant probes using a
p-value threshold of 1×10^{-11}, which gave the lowest error under our experimental setup, leaving
2,543 probes to build the clock using lasso regression, similar to the method used by Horvath[16]. We
then tested our novel epigenetic clock, which we named the CellAge clock, using an entirely different
set of samples, and we observed accurate prediction of passage number for both HMFs and HDFs,
with a Root Mean Square Error (RMSE) of 0.37 (Fig. 1c and Supplementary Fig. 2a).

Having built a precise epigenetic clock that measures methylation changes during replicative ageing

of human primary cells *in vitro*, we tested if anti-ageing drug treatment of HMFs and HDFs

decelerated the CellAge clock. We chose an mTOR inhibitor, rapamycin, which is one of the most

robust and evolutionarily conserved anti-ageing drug targets[36], and which mediates its effect through

down-regulation of S6K and Pol III, and up-regulation of autophagy[37,38]. We chose relatively low

rapamycin concentration of 5nM that did not inhibit cell growth (Supplementary Fig. 1a) but

moderately downregulated mTOR signalling, as evidenced by decreased pS6K and p4E-BP

phosphorylation (Supplementary Fig. 3). This setup mimics the pro-longevity effects of rapamycin *in*

vivo where it is well accepted that only mild nutrient sensing pathway inhibition increases life- and

healthspan[5,39]. DNA methylation profiles from HMFs collected following four, six and eight weeks of

rapamycin treatment (passage 16, 18 and 20; Fig. 2) were analysed using the CellAge clock and

clearly demonstrated that rapamycin slows down methylation changes associated with replicative

ageing. Interestingly, this clock deceleration was more pronounced upon longer treatment as shown

by the gradual decrease of predicted-actual passage from 16 to 20 weeks. The low dose rapamycin

treatment did not affect population doublings, confirming that the methylation changes were not a

reflection of proliferation inhibition or slowing of the cell cycle (Supplementary Fig. 1). This is further

evidenced by comparing the predicted passage from the CellAge clock against cumulative

population doubling, showing rapamycin samples lie on a separate line to that of the control samples

(Supplementary Fig. 2b,c). Contrarily, rapamycin samples and controls differed to considerably

lesser extent when actual passage and cumulative population doublings are compared

(Supplementary Fig. 2b,c). Importantly, we observed a similar pattern for HMFs and HDFs (Fig. 2),

suggesting that the CellAge clock could be applicable to different cells, albeit calibration is required

for cells that reach senescence at different rates.

We then focused on HMFs to test another anti-ageing drug, trametinib[40], an inhibitor of the

MEK/ERK signalling pathway, which we also applied in low concentration to avoid any effect on

growth and population doubling (Supplementary Fig. 1 and Fig. 3). The CellAge clock analysis of

trametinib treatment showed clock deceleration for all three passages tested (Fig. 2), thereby

confirming previous results in *Drosophila in vivo* that trametinib extends lifespan[40]. Next, we

examined the effect of two other inhibitors of nutrient-sensing pathways as mutations in these

pathways in model organisms represent the most evolutionary conserved anti-ageing interventions[5].

We tested Dactolisib/BEZ235, a dual ATP competitive PI3K and mTOR inhibitor, for which we again

optimised the dose of the treatment to obtain a reduction in signalling without significant proliferation

impairment, as shown by pS6K downstream target 4E-BP (Supplementary Fig. 3).

Dactolisib/BEZ235 slowed down the DNA methylation changes similar to rapamycin, suggesting that

Dactolisib/BEZ235 could be a new anti-ageing drug according to the output of the CellAge clock (Fig.

2). We also tested torin2, which is a selective inhibitor of the mTOR pathway that inhibits both

mTORC1 and mTORC2, unlike rapamycin, which targets solely mTORC1. Owing to its more

complete inhibition of the mTOR pathway, we were interested in examining its effect on replicative ageing, especially as the role of mTORC2 in ageing is less well established. The impact of mTORC2 inhibition on lifespan can be positive or negative depending on which of the mTORC2 downstream effectors is affected, in which tissue, and whether females or male mice are used for the experiment[41]. Some of the negative effects of mTOR pathway inhibition, such as insulin resistance and hyperlipidemia, are attributed to the mTORC2 branch of the pathway and may arise under certain conditions of prolonged and/or high dose rapamycin treatment[41]. Interestingly, while our CellAge clock suggests that torin2 is indeed a novel anti-ageing drug (Fig. 2), its effect on ageing in mammalian cell culture appears to be less pronounced than that of rapamycin. This is in line with literature suggesting that a promising strategy to improve healthy ageing is the development of inhibitors that are highly specific for mTORC1 or that target mTORC1 downstream effectors separately[41].

Next, we compared our anti-ageing drug screening results obtained by the CellAge clock with analyses using Horvath's Multi-tissue and Skin and Blood clock, as well as the PhenoAge clock. The clocks did not detect any significant effect of anti-ageing drug treatment (Supplementary Fig. 4). The Skin and Blood clock[26] was used recently to measure deceleration of ageing in primary fibroblasts[14,28], however the concentration of rapamycin used in our conditions was five times lower without effect on cell growth, highlighting the sensitivity of our epigenetic clock to detect age-related methylation changes at very low drug concentrations. Under our conditions, the only epigenetic clock that detected gradual methylation changes from passage 10 to passage 20 was the PhenoAge clock (Supplementary Fig. 4). However, its output was more variable between samples and inconsistent for anti-ageing drug treatments, reporting both clock acceleration and deceleration. For instance, rapamycin, Dactolisib/BEZ235 and torin2 treated cells appeared slightly younger compared to controls, whereas trametinib treated cells were estimated older to some extent (Supplementary Fig. 4), unlike the results we obtained with our CellAge clock (Fig. 2). Overall, the CellAge clock that we developed here was more consistent and performed significantly better on ageing cells in culture and following known anti-ageing drug treatments compared to existing clocks, as evidenced by its ability to detect subtle ageing differences. Our results are supportive of clocks being highly specialised for a certain task, and suggests that while other popular epigenetic clocks perform remarkably on determining donor's age in years and their health status, they were not able to robustly detect slight ageing changes in human primary cells induced by drug treatment over a short period of time *in vitro*.

Next, we assessed if the CellAge clock is suitable for the screening of novel anti-ageing drugs. To this aim, we examined if drugs that decelerate the CellAge clock also reduce features associated with senescence, such as morphological changes and expression of ageing biomarkers[42]. Rapamycin, Dactolisib/BEZ235 and Trametinib treatment slowed down morphological alteration in cells that gradually occur during replicative ageing, namely cell elongation, increased nuclear area

and cell area, and the treated cells appeared particularly 'youthful' (Fig. 3). Another characteristic of
senescence is increased expression of the cyclin-dependent kinase inhibitors p21$^{CIP1/Waf1}$ and
p16^{INK4a}. p21$^{CIP1/Waf1}$ triggers G1 cycle arrest upon damage and can lead to senescence or
apoptosis[43,44]. Expression of p16^{INK4a}, which is produced from the CDKN2A gene together with
p19^{ARF} (p14^{ARF} in humans) increases exponentially during ageing and was suggested to stabilise the
senescent state[45]. p16^{INK4a} expression was the marker of choice for senescent cell clearance leading
to prolonged lifespan in mice[46]. Our results demonstrate that drugs which decelerate the CellAge
clock at the same time reduce expression of both nuclear p21$^{CIP1/Waf1}$ and p16^{INK4a} compared to non-
treated cells, showing their efficacy in delaying the senescence programme (Fig. 3b,c). In addition,
the most frequently used senescent marker, senescent-associated β-galactosidase activity (SA-
βgal), was significantly decreased upon anti-ageing drugs treatment with rapamycin and
Dactolisib/BEZ235, but not in cells treated with trametinib (Fig. 3b,c). Another difference in
senescent markers was observed with interleukin-6 (IL-6), which is one of the most important
inflammatory cytokines and part of the senescent-associated secretory phenotype. IL-6 was
significantly reduced in aged cells upon rapamycin and Dactolisib/BEZ235 treatment but not in
trametinib treated cells (Fig. 3b,c). This difference possibly stems from the overactivated RAS/ERK
pathway being a more prominent inducer of senescence than the overactivated mTOR/PI3K
pathway[47], and hence corresponding inhibitors have different potency in inhibiting senescence.
Finally, we examined the nucleolus, an organelle dedicated to rRNA production and ribosomal
assembly, as it has recently emerged that maintenance of its structure, and low levels of nucleolar
methyltransferase fibrillarin, is a common denominator for major anti-ageing intervention from worms
to mice[48]. We observed that as a consequence of ageing, nucleoli in aged HMFs lose their defined
round shape, are more diffused, and stain less well. For rapamycin and Dactolisib/BEZ235, we
observed clearly defined and 'younger' looking nucleoli in aged cells. However, trametinib treated
cells resembled the nucleoli of controls. In summary, a panel of the most frequently used markers for
cell senescence confirmed that drugs which decelerate the CellAge clock also make the cells appear
more youthful. This strongly suggests that the CellAge clock can be used as a robust and sensitive
detector of novel anti-ageing treatments.

Finally, having discovered two novel potential anti-ageing drug treatments using the CellAge clock,
Dactolisib/BEZ235 and torin2, we tested and validated them *in vivo* using the fruit fly *Drosophila*
melanogaster as a model organism. This is important as tissue-specific drug toxicity, which can be
missed in cell culture, is one of the major reasons for drug failure in clinical trials. For *in vivo*
longevity studies we used the outbred wild-type w^{Dah} strain which is particularly suitable for ageing
studies, Drosoflipper device for fast fly transfer, and specially formulated holidic medium[49] to
increase drug bioavailability compared to standard sugar-yeast-agar fly food. We used rapamycin as
a positive control for longevity experiments in flies and showed that median lifespan extension on
holidic media varied from 7% to 9% compared to ethanol solvent control, depending on 1μM or 5μM
concentration, respectively (p<0.001, log-rank test), which is comparable to published literature[50]

(Fig. 4a). Importantly, both Dactolisib/BEZ235 and torin2 significantly extended lifespan in
Drosophila for 7% (p<0.001, log-rank test) (Fig. 4b,c). This firmly demonstrates that drugs that
decelerate the CellAge clock have similarly favourable output on major anti-ageing biomarkers *in*
vitro and extend longevity *in vivo*.

For the first time, we have a robust epigenetic clock for the rapid discovery of anti-ageing drugs
directly in human cells, bypassing lower model organisms and significantly shortening discovery time
compared to 3-year long mice longevity analysis. Testing different compounds for ageing using the
CellAge clock could potentially reveal new anti-ageing pathways and help us to improve our
knowledge base of not only ageing biology but of molecular pathways underpinning the epigenetic
clocks as well, understanding of which is limited. Other available epigenetic clocks could not detect
anti-ageing drugs so accurately. The CellAge clock, however, does not predict the chronological age
of the sample, demonstrating that epigenetic clocks are highly specialised to the purpose for which
they were trained. We expect many biological outputs to be extracted by different epigenetic clock
algorithms in the future, given the wealth of information stored in our epigenome.

Better experimental systems to test anti-ageing drugs are very much needed, given a rising
proportion of the elderly in modern societies and, as a consequence, larger numbers of people
suffering from age-related diseases. Our results show that by using the CellAge clock, cultured
primary human cells can be used as a proxy to measure human ageing and can reliably detect anti-
ageing effects upon a relatively brief treatment. By doing so, this fast and accurate method is
expected to accelerate the discovery of novel preventive treatments for age-related disease, directly
using human cells. Importantly, further research will be focused on expanding our findings on
different types of primary cells from donors with different ages, as well as on testing various
compounds. While ageing itself is not a disease, potential anti-ageing drugs could be FDA approved
separately for different conditions. The first study to test broad-spectrum protection capacity of
metformin, the TAME study, is underway[51]. In addition, it was shown that rapamycin/everolimus pre-
treatment dramatically improves flu vaccination and immune response in the elderly[52]. In mice, it
also lowers the incidence of tumours[53], and it shows promising results in the field of
neurodegeneration[54]. This supports the idea that targeting healthy ageing might have multiple
beneficial outputs.

Our novel drug discovery platform will inform on new anti-ageing mechanisms, currently dominated
by IIS and mTOR signalling pathways as well as dietary restriction regimes, and will thereby
advance our understanding of the ageing process. Many drugs targeting growth pathways are
already available from cancer research where they are used in very high doses. With our CellAge
clock, it could be examined which of these compounds can be disease preventative at very low
concentrations. Our experimental setup is also suitable for nutraceutical approaches whereby dietary
supplements could be rigorously tested for their effect on ageing. Overall, we expect our accelerated

discovery platform to be valuable for the discovery of strongly sought after anti-ageing drugs and
geroprotective strategies to improve healthy human ageing.
309

Methods

Cell culture and reagents. Normal finite lifespan human mammary fibroblasts (HMFs) were
obtained from reduction mammoplasty tissue of a 16-year-old individual, donor 48 by Dr Martha
Stampfer (University Berkeley) who has all required IRB approvals to distribute these cell
samples and MTA agreement set in place with Dr Cleo Bishop laboratory. Independent cultures
from these cells were serially passaged from passage 9 through to passage 20 and aliquots
taken upon each passage for Illumina Infinium Methylation EPIC analysis. HMFs were
maintained in Dulbecco's Modified Eagles Medium (DMEM) (Life Technologies, UK)
supplemented with 10% foetal bovine serum (FBS) (Labtech.com, UK), 2mM L-glutamine (Life
Technologies, UK) and 10 μg/mL insulin from bovine pancreas (Sigma).
Normal finite lifespan human dermal fibroblasts (HDFs) were obtained from face lift dermis
following a kind donation from an anonymous healthy patient under standard ethical practice,
reference LREC No. 09/HO704/69. HDFs were grown in DMEM with 4 mM L-glutamine (Life
Technologies), supplemented with 10% FBS.
Cells were plated at 7,500 cells/cm^2 in T25 cell culture flask in 5ml of media to which 5μl of
appropriate drug or vehicle control was added. Media was changed every two days and cells
were passaged every 7 days and trypsinisation was used to detach the cells. All cells were
routinely tested for mycoplasma and shown to be negative.
328

Immunofluorescence microscopy and high content analysis. Cells were washed in
phosphate buffered saline (PBS), fixed for 15 minutes with 3.7% paraformaldehyde with 5%
sucrose, washed and permeabilised for 15 minutes using 0.1% Triton X in PBS (30 minutes
for anti-nucleolin antibody) then washed and blocked in 0.25% bovine serum albumin (BSA)
in PBS before primary antibody incubations. Primary antibodies used were anti-IL6 (R&D
Systems, 1:100; overnight 4°C), anti-nucleolin (Santa Cruz, 1:2000, overnight room
temperature), anti-p16 (Proteintech, 1:500, overnight 4°C), anti-p21 (12D1, Cell Signalling,
1:2000, overnight 4°C). Cells were incubated for 2 hours at room temperature with the
appropriate AlexaFluor-488, AlexaFluor-546 or AlexaFluor-647 conjugated antibody (1:500,
Invitrogen), DAPI (1:1,000 from 1mg/mL stock) and CellMask Orange or Deep Red
(1:200,000, Invitrogen). Images were acquired using the IN Cell 2200 or 6000 automated
microscope (GE) and HCA was performed using the IN Cell Developer software (GE).
342

Z score generation. For each of the parameters analysed, significance was defined as one Z score from the negative control mean. Z scores were generated according to the formula below: Z score = (mean value of experimental condition – mean value of vehicle control/standard deviation (SD) for vehicle control.

Senescence-associated beta-galactosidase (SA-β-Gal) assay. Cells were washed in PBS, fixed for 5 minutes with 0.2% glutaraldehyde, washed and incubated for 24 hour at 37°C (no CO_2) with fresh senescence-associated beta-galactosidase (SA-β-Gal) solution: 1mg of 5-bromo-4-chloro-3-indoyl β-D-galactosidase (X-Gal) per mL (stock = 20mg of dimethylsulfoxide per ml) / 40mM citric acid/sodium phosphate, pH 6.0 / 5mM potassium ferrocyanide / 5mM potassium ferricyanide / 150mM NaCl / 2mM $MgCl_2$). Cells were stained with Hoechst 33342 (1:10,000 from 10mg/mL stock) for 30 minutes. Images were acquired using the IN Cell 2200 automated microscope and HCA was performed using the IN Cell Developer software.

Genomic DNA extraction. For isolation of Genomic DNA from primary human fibroblasts we used QIAamp DNA micro kit (56304) and we followed manufacturers protocol, with an additional washing steps with 500μl AW2 buffer and 500μl 80% ethanol to improve purity. DNA quantification and purity was determined by Nanodrop and QuBit. For bisulfite conversion EZ DNA methylation kit was used (D5001).

Preparation of methylation array data. For each sample, 500ng high-quality DNA was bisulphite converted using the EZ DNA methylation kit (Zymo Research), using the alternative incubation conditions recommended for use with Illumina methylation arrays. Bisulphite converted DNA was eluted in 12ul elution buffer. Methylation was analysed using the Infinium Human Methylation EPIC array (Illumina) using standard operating procedures at the UCL Genomics facility. The EPIC array data have been deposited into ArrayExpress at the European Bioinformatics Institute (https://www.ebi.ac.uk/arrayexpress/) under accession number E-MTAB-8327.

.

Pre-processing of methylation array data. DNA methylation array data was processed using the minfi package[55] within R (http://www.R-project.org/). Initial QC metrics from this package were used to remove low-quality samples. Probes were filtered using a detection p-value cut-off >0.01 and normalised using the Noob procedure. Cross-hybridising probes

were removed from analysis based on the list published in McCartney et al.[56] The training

and test sets were pre-processed separately to obtain a fair estimate of the performance of

CellAge clock.

Estimation of sample age using existing epigenetic clocks. Following pre-processing

of data, the epigenetic age of all samples was predicted using three epigenetic clocks; the

Multi-tissue clock[16], the Skin and Blood clock[13] and the PhenoAge clock[23].

Development of the CellAge clock. The clock was built using a total of 39 samples, with

6 samples at each of the following passages; 10,12, and 14 and 7 samples at each of the

following passages; 16, 18, and 20. This included both dermal fibroblasts (n=12) and

mammary fibroblasts (n=27). Probes were initially filtered from the 730,453 available using

the dmpFinder function within minfi[55]. A p-value threshold was used to determine which

probes to use in the model building and this was selected using a leave one out validation,

yielding the threshold of 1×10^{-11} and a total of 2,543 probes. We built the model using

elastic net regression (setting alpha=0.5) from the glmnet package[57] within R and

determining the lambda parameter using internal cross validation function provided.

Lifespan measurements. We used *white Dahomey* (w^{Dah}) wild-type flies that were

maintained and all experiments were conducted at 25ºC. Flies were kept on a 12 h light:12

h dark cycle at constant humidity using standard sugar/yeast/agar (SYA) medium. For all

experiments, flies were reared at standard larval density by transferring 18 μl of egg

suspension into SYA bottles. Eclosing adults were collected over a 12-h period and allowed

to mate for 48 h before sorting into single sexes and placed in vials containing either control

or experimental drug food. For lifespan assays, flies were reared at standard density and

maintained at 15 flies per vial and we used holidic media recipe food for all longevity

assays[49]. Flies were transferred to fresh food vials every 2-3 days and scored for deaths. At

least 150 flies were used for each lifespan experiment.

Western blot measurements. Whole flies or human primary cell pellet was homogenised

in 2x Laemmli loading sample buffer (100 mM Tris pH 6.8, 20% glycerol, 4% SDS; Bio-Rad)

containing 50 mM DTT. Extracts were cleared by centrifugation and approximately 20 µg of

protein extract was loaded per lane on a polyacrylamide gel. Proteins were separated and

transferred to nitrocellulose membrane. The following antibodies were used at the indicated

dilutions: H3 (Cell Signaling Technology; 1:2000; 4499S), pS6K (Cell Signaling Technology;

1:1000; 9206S), total S6K (Santa Cruz; 1:1000; 8418), p4EBP (Cell Signalling Technology,

1:500; 2855S), non-phospho4E-BP (Cell Signalling Technology; 1:500; 4923S), pAkt (Cell

Signalling; 1:1000; 4060), pAkt (Cell Signalling; 1:1000; 4056), total Akt (Cell Signalling;

1:1000; 9272), pERK (Cell Signalling; 1:1000; 4370), total (Cell Signalling; 1:1000; 4692).

Blots were developed using the ECL detection system (GE, Amersham), and analysed

using FIJI software (US National Institutes of Health). We used precasted TGX stain-free

gels from Bio-Rad (567-8123 or 567-8124) according to the manufacturer's instructions.

Statistical analysis. Statistical analysis was performed using JMP software (version 4.0.5;

SAS Institute) and R. Log-rank tests were performed on lifespan curves. Probes were

selected to build CellAge using an F-test and CellAge was built using lasso regression.

Online content

The code for CellAge clock is available from GitHub. All methylation microarray data reported

in   this   study   have   been   deposited   in   the   ArrayExpress

(https://www.ebi.ac.uk/arrayexpress/) public repository and they are accessible under

accession number E-MTAB-8327.

References

1   Wilson, B. E. *et al.* Estimates of global chemotherapy demands and corresponding

physician workforce requirements for 2018 and 2040: a population-based study.

Lancet Oncol **20**, 769-780, doi:10.1016/S1470-2045(19)30163-9 (2019).

2   Reeve, A., Simcox, E. & Turnbull, D. Ageing and Parkinson's disease: why is

advancing age the biggest risk factor? *Ageing Res Rev* **14**, 19-30,

doi:10.1016/j.arr.2014.01.004 (2014).

3   Partridge, L., Deelen, J. & Slagboom, P. E. Facing up to the global challenges of

ageing. *Nature* **561**, 45-56, doi:10.1038/s41586-018-0457-8 (2018).

4   Rae, M. J. *et al.* The demographic and biomedical case for late-life interventions in

aging. *Sci Transl Med* **2**, 40cm21, doi:2/40/40cm21 [pii]

10.1126/scitranslmed.3000822 (2010).

5   Lopez-Otin, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The

hallmarks of aging. *Cell* **153**, 1194-1217, doi:S0092-8674(13)00645-4 [pii]

10.1016/j.cell.2013.05.039 (2013).

6   Pinkston, J. M., Garigan, D., Hansen, M. & Kenyon, C. Mutations that increase the
life span of C. elegans inhibit tumor growth. *Science* **313**, 971-975 (2006).
7   Cohen, E., Bieschke, J., Perciavalle, R. M., Kelly, J. W. & Dillin, A. Opposing
activities protect against age-onset proteotoxicity. *Science* **313**, 1604-1610 (2006).
8   Wessells, R. J., Fitzgerald, E., Cypser, J. R., Tatar, M. & Bodmer, R. Insulin
regulation of heart function in aging fruit flies. *Nat Genet* **36**, 1275-1281 (2004).
9   Selman, C. *et al.* Evidence for lifespan extension and delayed age-related
biomarkers in insulin receptor substrate 1 null mice. *Faseb J* **22**, 807-818 (2008).
10   Niccoli, T. & Partridge, L. Ageing as a risk factor for disease. *Curr Biol* **22**, R741-
752, doi:S0960-9822(12)00815-9 [pii]
10.1016/j.cub.2012.07.024 (2012).
11   Harrison, D. E. *et al.* Rapamycin fed late in life extends lifespan in genetically
heterogeneous mice. *Nature* **460**, 392-395, doi:nature08221 [pii]
10.1038/nature08221 (2009).
12   Martin-Montalvo, A. *et al.* Metformin improves healthspan and lifespan in mice. *Nat
Commun* **4**, 2192, doi:10.1038/ncomms3192 (2013).
13   Horvath, S. *et al.* Epigenetic clock for skin and blood cells applied to Hutchinson
Gilford Progeria Syndrome and ex vivo studies. *Aging (Albany NY)* **10**, 1758-1775,
doi:10.18632/aging.101508 (2018).
14   Horvath, S., Lu, A. T., Cohen, H. & Raj, K. Rapamycin retards epigenetic ageing of
keratinocytes independently of its effects on replicative senescence, proliferation
and differentiation. *Aging (Albany NY)* **11**, 3238-3249, doi:10.18632/aging.101976
(2019).
15   Castillo-Quan, J. I., Kinghorn, K. J. & Bjedov, I. Genetics and pharmacology of
longevity: the road to therapeutics for healthy aging. *Adv Genet* **90**, 1-101,
doi:S0065-2660(15)00005-X [pii]
10.1016/bs.adgen.2015.06.002 (2015).
16   Horvath, S. DNA methylation age of human tissues and cell types. *Genome Biol* **14**,
R115, doi:gb-2013-14-10-r115 [pii]
10.1186/gb-2013-14-10-r115 (2013).
17   Booth, L. N. & Brunet, A. The Aging Epigenome. *Mol Cell* **62**, 728-744,
doi:10.1016/j.molcel.2016.05.013 (2016).
18   Hannum, G. *et al.* Genome-wide methylation profiles reveal quantitative views of
human aging rates. *Mol Cell* **49**, 359-367, doi:S1097-2765(12)00893-3 [pii]
10.1016/j.molcel.2012.10.016 (2013).

19   Horvath, S. & Raj, K. DNA methylation-based biomarkers and the epigenetic clock

theory of ageing. *Nat Rev Genet* **19**, 371-384, doi:10.1038/s41576-018-0004-3

(2018).

20   Field, A. E. *et al.* DNA Methylation Clocks in Aging: Categories, Causes, and

Consequences. *Mol Cell* **71**, 882-895, doi:10.1016/j.molcel.2018.08.008 (2018).

21   Garagnani, P. *et al.* Methylation of ELOVL2 gene as a new epigenetic marker of

age. *Aging Cell* **11**, 1132-1134, doi:10.1111/acel.12005 (2012).

22   Marioni, R. E. *et al.* DNA methylation age of blood predicts all-cause mortality in

later life. *Genome Biol* **16**, 25, doi:10.1186/s13059-015-0584-6 (2015).

23   Levine, M. E. *et al.* An epigenetic biomarker of aging for lifespan and healthspan.

Aging (Albany NY) **10**, 573-591, doi:10.18632/aging.101414 (2018).

24   Lu, A. T. *et al.* DNA methylation GrimAge strongly predicts lifespan and healthspan.

Aging (Albany NY) **11**, 303-327, doi:10.18632/aging.101684 (2019).

25   Yang, Z. *et al.* Correlation of an epigenetic mitotic clock with cancer risk. *Genome*

Biol **17**, 205, doi:10.1186/s13059-016-1064-3

10.1186/s13059-016-1064-3 [pii] (2016).

26   Tomasetti, C., Li, L. & Vogelstein, B. Stem cell divisions, somatic mutations, cancer

etiology, and cancer prevention. *Science* **355**, 1330-1334, doi:355/6331/1330 [pii]

10.1126/science.aaf9011 (2017).

27   Martin-Herranz, D. E. *et al.* Screening for genes that accelerate the epigenetic aging

clock in humans reveals a role for the H3K36 methyltransferase NSD1. *Genome*

Biol **20**, 146, doi:10.1186/s13059-019-1753-9 (2019).

28   Sturm, G. *et al.* Human aging DNA methylation signatures are conserved but

accelerated in cultured fibroblasts. *Epigenetics*, 1-16,

doi:10.1080/15592294.2019.1626651 (2019).

29   Ye, X., Linton, J. M., Schork, N. J., Buck, L. B. & Petrascheck, M. A

pharmacological network for lifespan extension in Caenorhabditis elegans. *Aging*

Cell **13**, 206-215, doi:10.1111/acel.12163 (2014).

30   Lucanic, M., Lithgow, G. J. & Alavez, S. Pharmacological lifespan extension of

invertebrates. *Ageing Res Rev* **12**, 445-458, doi:10.1016/j.arr.2012.06.006 (2013).

31   Cabreiro, F. *et al.* Metformin retards aging in C. elegans by altering microbial folate

and methionine metabolism. *Cell* **153**, 228-239, doi:10.1016/j.cell.2013.02.035

(2013).

32 Pryor, R. *et al.* Host-Microbe-Drug-Nutrient Screen Identifies Bacterial Effectors of
Metformin Therapy. *Cell* **178**, 1299-1312 e1229, doi:10.1016/j.cell.2019.08.003
(2019).
33 Zimmermann, A. *et al.* Yeast as a tool to identify anti-aging compounds. *FEMS*
Yeast Res **18**, doi:10.1093/femsyr/foy020 (2018).
34 Donertas, H. M., Fuentealba Valenzuela, M., Partridge, L. & Thornton, J. M. Gene
expression-based drug repurposing to target aging. *Aging Cell* **17**, e12819,
doi:10.1111/acel.12819 (2018).
35 Lowe, R. *et al.* The senescent methylome and its relationship with cancer, ageing
and germline genetic variation in humans. *Genome Biol* **16**, 194,
doi:10.1186/s13059-015-0748-4 (2015).
36 Saxton, R. A. & Sabatini, D. M. mTOR Signaling in Growth, Metabolism, and
Disease. *Cell* **168**, 960-976, doi:10.1016/j.cell.2017.02.004 (2017).
37 Bjedov, I. *et al.* Mechanisms of life span extension by rapamycin in the fruit fly
Drosophila melanogaster. *Cell Metab* **11**, 35-46, doi:S1550-4131(09)00374-X [pii]
10.1016/j.cmet.2009.11.010 (2010).
38 Filer, D. *et al.* RNA polymerase III limits longevity downstream of TORC1. *Nature*
552, 263-267, doi:10.1038/nature25007 (2017).
39 Bjedov, I. & Partridge, L. A longer and healthier life with TOR down-regulation:
genetics and drugs. *Biochem Soc Trans* **39**, 460-465, doi:BST0390460 [pii]
10.1042/BST0390460 (2011).
40 Slack, C. *et al.* The Ras-Erk-ETS-Signaling Pathway Is a Drug Target for Longevity.
Cell **162**, 72-83, doi:S0092-8674(15)00707-2 [pii]
10.1016/j.cell.2015.06.023 (2015).
41 Kennedy, B. K. & Lamming, D. W. The Mechanistic Target of Rapamycin: The
Grand ConducTOR of Metabolism and Aging. *Cell Metab* **23**, 990-1003,
doi:10.1016/j.cmet.2016.05.009 (2016).
42 Hanzelmann, S. *et al.* Replicative senescence is associated with nuclear
reorganization and with DNA methylation at specific transcription factor binding
sites. *Clin Epigenetics* **7**, 19, doi:10.1186/s13148-015-0057-5 (2015).
43 He, S. & Sharpless, N. E. Senescence in Health and Disease. *Cell* **169**, 1000-1011,
doi:10.1016/j.cell.2017.05.015 (2017).
44 McHugh, D. & Gil, J. Senescence and aging: Causes, consequences, and
therapeutic avenues. *J Cell Biol* **217**, 65-77, doi:10.1083/jcb.201708092 (2018).

45   Gire, V. & Dulic, V. Senescence from G2 arrest, revisited. *Cell Cycle* **14**, 297-304,
doi:10.1080/15384101.2014.1000134 (2015).

46   Baker, D. J. *et al.* Naturally occurring p16(Ink4a)-positive cells shorten healthy
lifespan. *Nature* **530**, 184-189, doi:10.1038/nature16932 (2016).

47   Kennedy, A. L. *et al.* Activation of the PIK3CA/AKT pathway suppresses
senescence induced by an activated RAS oncogene to promote tumorigenesis. *Mol*
Cell **42**, 36-49, doi:10.1016/j.molcel.2011.02.020 (2011).

48   Tiku, V. & Antebi, A. Nucleolar Function in Lifespan Regulation. *Trends Cell Biol* **28**,
662-672, doi:10.1016/j.tcb.2018.03.007 (2018).

49   Piper, M. D. *et al.* A holidic medium for Drosophila melanogaster. *Nat Methods* **11**,
100-105, doi:10.1038/nmeth.2731 (2014).

50   Fan, X. *et al.* Rapamycin preserves gut homeostasis during Drosophila aging.
Oncotarget **6**, 35274-35283, doi:10.18632/oncotarget.5895 (2015).

51   Barzilai, N., Crandall, J. P., Kritchevsky, S. B. & Espeland, M. A. Metformin as a
Tool to Target Aging. *Cell Metab* **23**, 1060-1065, doi:S1550-4131(16)30229-7 [pii]
10.1016/j.cmet.2016.05.011 (2016).

52   Mannick, J. B. *et al.* TORC1 inhibition enhances immune function and reduces
infections in the elderly. *Sci Transl Med* **10**, doi:10.1126/scitranslmed.aaq1564
(2018).

53   Anisimov, V. N. *et al.* Rapamycin increases lifespan and inhibits spontaneous
tumorigenesis in inbred female mice. *Cell Cycle* **10**, 4230-4236, doi:18486 [pii]
10.4161/cc.10.24.18486 (2011).

54   Bove, J., Martinez-Vicente, M. & Vila, M. Fighting neurodegeneration with
rapamycin: mechanistic insights. *Nat Rev Neurosci* **12**, 437-452,
doi:10.1038/nrn3068 (2011).

55   Fortin, J. P., Triche, T. J., Jr. & Hansen, K. D. Preprocessing, normalization and
integration of the Illumina HumanMethylationEPIC array with minfi. *Bioinformatics*
33, 558-560, doi:10.1093/bioinformatics/btw691 (2017).

56   McCartney, D. L. *et al.* Identification of polymorphic and off-target probe binding
sites on the Illumina Infinium MethylationEPIC BeadChip. *Genom Data* **9**, 22-24,
doi:10.1016/j.gdata.2016.05.012 (2016).

57   Friedman, J., Hastie, T. & Tibshirani, R. Regularization Paths for Generalized Linear
Models via Coordinate Descent. *J Stat Softw* **33**, 1-22 (2010).

Acknowledgements

IB acknowledges funding from ERC StG 311331, ERC PoC 842174, Royal Society Research Grant, The Bill Lyons foundation. This work was supported in part by the CRUK-UCL Centre Award [C416/A25145] awarded to IB and SB. EJT has been funded by MRC (MR/K501372/1) and BBSRC (BB/P002579/1) and DM is funded by the BBSRC (BB/N503629/1). JG, and MS are supported by U.S. Department of Energy under Contract No. DE-AC02-05CH11231 and the Congressionally Directed Medical Research Programs Breast Cancer Research Program Era of Hope Scholar Award BC141351.

Authors contribution

SB, IB developed initial concept. SB, IB, RL and CLB finalised concept developing and designed experiments. RL and APW analysed data. RL developed CellAge clock. CLB and EJT provided cell culture expertise. CL, EJT, ERS, VEMM, DM and IB performed all experiments. SE, SB, IB, RL, CLB wrote the manuscript. JCG and MRS provided critical reagents. All authors discussed results and commented on and approved the manuscript.

Competing interest

The authors declare no competing interest.

Additional information

Correspondence and requests for materials should be addressed to IB, CLB, RL and SB.

Figure Legends

Figure 1. Development of CellAge clock for monitoring subtle ageing difference in cells in culture.

A) Predicted age of control samples using three existing epigenetic clocks. Predicted epigenetic age for control samples across all experiments as estimated by the Multi-tissue clock (green), the Skin and Blood clock (orange) and the PhenoAge clock (yellow). All three clocks show a trend to increase in predicted age with progressing passage, however there is a lot of variability in predictions, particularly for the PhenoAge clock. The Multi-tissue clock consistently predicted cells to have the highest epigenetic age, while the PhenoAge clock consistently predicted cells to

have the lowest epigenetic age, which even reached below zero for several samples

at various passages.

B) Genome-wide methylation changes upon cell passages of primary human

mammary fibroblasts (HMF) and primary human dermal fibroblast (HDF).

C) Testing the CellAge clock on HMF and HDF samples that were not used to train the

clock, demonstrates accurate prediction of the cell passage.

Figure 2. Using the CellAge clock for the detection of anti-ageing drugs.

A) The methylome of Human Dermal Fibroblasts (HDF) and

B) Human Mammary Fibroblasts (HMF) was analysed using the CellAge clock.

Represented is Predicted-Actual Passage for Passage 16, 18 and 20, showing

deceleration of CellAge upon treatment with anti-ageing drugs rapamycin (5nM),

Dactolisib/BEZ235 (10nM), torin2 (5nM) and Trametinib (0.1nM).

Figure 3. Treatment with anti-ageing drugs decreases markers of senescence.

A) Schematic illustrating the experimental set-up conducted in P10 to P22 HMFs,

passaged weekly.

B) Multi-parameter analysis of senescence markers. Colour coding used to illustrate

the number of Z scores of the experimental drug value from the respective control

mean. Scores highlighted in red denote a shift towards a more proliferative

phenotype and scores highlighted in green denote a shift to a more senescent

phenotype.

C) P22 HMFs stained with DAPI (blue) and Cell Mask, p21, p16, IL-6, or nucleolin

(red), or SA-β-Gal (blue) following 96-day treatment with 5nM Rapamycin, 10nM

Dactolisib/BEZ235, 0.1nM Trametinib or their respective controls. Size bar,

100μm.

Figure 4. Drugs that decelerate CellAge extend lifespan *in vivo*.

A) Lifespan analysis on w^{Dah} background wild type flies fed with SYA food containing

different concentration of rapamycin or ethanol as solvent control. For each

condition, 150 flies were used.

B) Lifespan analysis on w^{Dah} background wild type flies fed with SYA food containing

different concentration of Dactolisib/BEZ235 or DMSO as solvent control. For each

condition, 150 flies were used.

C) Lifespan analysis on w^{Dah} background wild type flies fed with SYA food containing

647   different concentration of torin2 or DMSO as solvent control. For each condition,

648   150 flies were used.

Figure 1.

Figure 2.

Figure 3

Figure 4