Indirect versus Direct Effects of Freshwater Browning on Larval Fish Foraging

Dina M. Leech, Troy L. Clift, Jessica L. Littlefield, Nicholas R. Ravagli, and Jacob E. Spain

Department of Biological and Environmental Sciences
Longwood University Farmville, VA 23901

Corresponding Author: Dina M. Leech, Department of Biological and Environmental Sciences, 201 High Street, Longwood University, Farmville, VA, 23909; Office #: 434.395.4966; Email: leechdm@longwood.edu

Keywords: brownification, fish larvae, foraging, predation, zooplankton

Running Title: Browning and larval fish foraging

ABSTRACT

Fish foraging and energy flow are both predicted to decline with freshwater 'browning'

due to reductions in light availability. Studies investigating these predictions have focused on

juveniles and adults; however, the larval stage represents a 'critical period' in fish development.

We investigated the indirect versus direct effects of browning on zooplankton-larval fish

interactions by altering water color with SuperHume (absorbance at 440 nm = 1.6 – 10.8 m^{-1}).

Phytoplankton and zooplankton densities were monitored across experimental tanks in the

laboratory for one month leading up to fish spawning. Larval largemouth bass were then

introduced to assess indirect effects on fish feeding rates and growth. Direct effects on foraging

of largemouth bass and bluegill were determined with separate short-term feeding

experiments. Browning did not directly alter the ability of larval fish to capture prey. However,

significant indirect effects on larval fish foraging, growth, and survival were observed as

phytoplankton and zooplankton decreased with increased browning. Our data suggest lake

browning will reduce energy transfer to larval fish due to a reduction in prey availability but not

visual foraging.

Introduction

In recent decades, many freshwater and coastal ecosystems in the Northern

Hemisphere have become browner in color due to an increased export of chromophoric

dissolved organic matter (cDOM) from the terrestrial watershed (Monteith et al. 2007;

Erlandsson et al. 2008; Haaland et al. 2010; Solomon et al. 2015). The mechanisms underlying

this 'browning' are currently debated and include changes in climate, hydrology, and land use,

reduced atmospheric acid deposition, and increased inputs of dissolved iron (Freeman et al.

2001; Monteith et al. 2007; Erlandsson et al. 2008; Kritzberg and Ekström 2012). Whatever the

mechanism, this phenomenon has far reaching ecological consequences for the structure and

function of aquatic ecosystems, including energy flow.

One concern is the effect of browning on the underwater light environment. As waters

become browner in color, the quantity of light in the water column is reduced, resulting in a

shallower euphotic zone (Bukaveckas and Robbins-Forbes 2000; Einem and Graneli 2010). In

addition, shorter wavelength ultraviolet and visible radiation are more readily absorbed by

cDOM compared to longer wavelengths (Morris et al. 1995; Wetzel 2001). Consequently, the

light environment shifts to the red portion of the visible spectrum. These alterations in the

quantity and quality of light have the potential to affect zooplankton-fish interactions--

indirectly through reduced energy transfer up the food chain and directly through reduced

foraging efficiency.

While recent studies have shown that the initial browning of low productive systems

can stimulate primary production (Ask et al. 2012; Seekell et al. 2015; Williamson et al. 2015),

excessive browning of fresh waters can reduce rates of photosynthesis due to competition for

photons (Kirk 1994). cDOM has been reported to sequester 10 times more photons within the

visible spectrum (400 - 700 nm) compared to photosynthetic pigments (Thrane et al. 2014). As

a consequence, areal primary production often decreases with increased water color (Jones

1992; Carpenter et al. 1998; Ask et al. 2009; Karlsson et al. 2009; Thrane et al. 2014), and

pelagic primary production typically exceeds benthic primary production (Vasconcelos et al.

2018; Vasconcelos et al. 2019). Phytoplankton community composition can also shift to

predominantly cyanobacteria, which are generally less nutritious or inedible to zooplankton

(Ekvall et al. 2013; Robidoux et al. 2015). Overall, aquatic ecosystems often become net

heterotrophic with increased browning as bacterial production exceeds primary production

(Cole et al. 1994; Ask et al. 2012).

Reductions in the quantity and quality of basal resources then 'cascade up' the grazer

food chain to influence zooplankton and fish. For example, Robidoux et al. (2015) observed

decreases in crustacean zooplankton biomass and density with increased water color while

Craig et al. (2017) found that bluegill in lakes of increasing color were smaller in size and had

lower fecundity compared to those living in lakes with less color. Taipale et al. (2018) noted

that zooplankton and fish had poorer nutritional quality in browner systems because the

phytoplankton on which they fed had lower concentrations of essential fatty acids, proteins,

lipids, and carbohydrates. Combined, these results suggest that freshwater browning indirectly

alters energy flow to higher trophic levels.

Trophodynamics between larval fish and zooplankton are particularly important as the

larval stage represents a critical phase in fish development that ultimately affects population

growth and biomass (Fuiman and Werner 2002; Karlsson et al. 2009; Karlsson et al. 2015).

Generally, in temperate climates, increased primary production in the late spring/early summer

leads to increased zooplankton production. Shortly thereafter, many fish species begin to

spawn, matching larval fish hatching with an abundance of zooplankton prey (Mills et al. 1989;

Mehner and Thiel 1999; Hansson et al. 2007). However, as fresh waters continue to brown and

energy flow to zooplankton is reduced, fish larvae may compete for fewer zooplankton prey.

Freshwater browning may also directly affect fish foraging behavior by altering the light

environment to which fish are adapted. Many fish are visually orienting predators, depending

on the quantity (i.e. intensity) and quality (i.e. spectra) of underwater light to locate and

capture prey (Guthrie and Muntz 1993; Leech and Johnsen 2009). Results from studies

investigating the direct effects of browning on fish foraging are varied, ranging from no effect

to enhanced effects (e.g. Stasko et al. 2012; Jönsson et al. 2013; Weidel et al. 2017). However,

this research has focused on juvenile and adult life history stages, with no knowledge of the

effect of browning on fish larvae.

Foraging in early life history stages may be particularly affected by freshwater browning

due the rapid attenuation of shorter wavelength light. Over evolutionary time, the visual

system of fishes spectrally tunes to the intensity and wavelengths of light present in their

environment, varying with age and behavior (Douglas and Djamgoz 1990). Larval fish spend

most of their time foraging in the top few meters of the epilimnion, and many species have

been shown to possess UV photoreceptors during only their early stages of development

(reviewed in Leech and Johnsen 2009). Thus, fish larvae may rely on short-wavelength

ultraviolet and blue radiation to forage (Leech and Johnsen 2009 and references therein).

Alternatively, as light levels decline, fish larvae may rely on other sensory mechanisms to

forage, such as olfactory cues or mechanoreception, as demonstrated in marine fish larvae and

zebrafish (Jones and Janssen 1992; Cobcroft and Pankhurst, 2003; Sampson et al. 2013; Carillo

and McHenry 2016).

Here, we use laboratory experiments to assess the indirect versus direct effects of

browning on larval fish foraging at the time of hatching (i.e., late spring/early summer in central

Virginia, USA). Based on the primary literature, we hypothesized that increased browning will

reduce phytoplankton biomass, leading to a reduction in zooplankton abundance, and

consequently larval fish foraging efficiency, growth, and survival during an early, critical stage in

development. We then used the same laboratory set up to examine the direct effects of

browning on fish feeding rate and prey selectivity when given equal zooplankton prey,

hypothesizing that fish larvae will consume less prey as water color increases due to a reduction

in light availability for foraging.

Methods

Indirect Effects of Browning

Laboratory experiments were conducted in twelve 20 L glass aquaria assembled on

three shelves with four tanks per shelf. The entire shelving unit, including individual shelves,

was covered in black plastic with additional black plastic placed in between each tank.

Experimental tanks were arranged in a randomized block design, with at least one tank from

each treatment on each shelf. Nine tanks were filled with 17 L of artificial lake water made from

the COMBO medium recipe, which provides macro- and micronutrients in relatively high

concentrations to support growth (Kilham et al. 1998). SuperHume, a commercially available

source of humic acid, was then added to the tanks at varying concentrations to adjust the

brown color of the water. Three tanks received 7 µg/L of SuperHume to serve as a light brown

treatment (water color measured as absorbance at 440 nm (a_{440}) = 1.6 m^{-1}), three tanks

received 33 µg/L to serve as a moderate brown treatment (a_{440} = 5.7 m^{-1}), and three tanks

received 66 µg/L to serve as a dark brown treatment (a_{440} = 10.8 m^{-1}). While SuperHume adds

carbon to the system, it does not add nutrients. Thus, despite differences in water color, and

consequently light transparency, our treatments had similar nutrient concentrations. We

adjusted the pH of all tanks to ~7 with approximately 2 µL of 2N HCl.

Although the experimental use of SuperHume has been previous tested (Lennon et al.

2013; Övergaard 2019), some have cited toxicity concerns for zooplankton (Robidoux et al.

2015). We therefore filled the remaining three tanks with 17 L of lake water from Sandy River

Reservoir, Farmville, VA (a_{440} = 1.5 m^{-1}) to serve as a comparison with our light brown

SuperHume treatment. Lake water was passed through a GF/F filter to remove bacterio-,

phyto-, and zooplankton, as best as possible. Sandy River Reservoir water was not used as the

base water for all browning treatments because of the logistics of transporting and filtering the

necessary large volume of water needed for the experiment.

Light was provided by 1.2 m long grow lamps containing two Lumichrome ® Full

Spectrum Plus fluorescent, 32W bulbs suspended approximately 10 cm above the water surface

of the tanks on each shelf and placed on a 14-hr light, 10-hr dark cycle. While lamps cannot

exactly reproduce the solar spectrum, these specific bulbs were selected based on their broad

coverage of the ultraviolet and visible spectra. An Ocean Optics Red Tide USB650 UV

spectrophotometer was used to measure the light spectrum, based on photon counts, from 250

to 800 nm in each treatment (Figure 1). After the experiment had concluded, we acquired a

LiCor LI-192 quantum sensor to estimate photosynthetically active radiation (PAR) across our

treatments. The sensor was placed in the center of the tank at a depth of 0.09 m. PAR ranged

from approximately 15 µmol m^{-2} s^{-1} in the dark brown treatment, 22 µmol m^{-2} s^{-1} in the

moderate brown treatment, and 31 – 35 µmol m^{-2} s^{-1} in the lake water and light brown

treatments. These light levels are comparable to those experienced during crepuscular periods,

when larvae often actively forage (Keast and Welsh 1968; Leech and Johnsen 2009). They are

also representative of light levels experienced at midday in the summer between approximately

1.0 – 2.5 m depth in local, natural systems (i.e. ~ 1800 µmol m^{-2} s^{-1} surface irradiance, Figure

S1). All tanks stabilized to a room temperature of 22 - 23 ^{o}C under both light and dark

conditions.

At the beginning of the experiment, each tank received a dense mixture of

phytoplankton containing equal parts of *Ankistrodesmus* sp., *Chlorella* sp*.*, *Scenedesmus* sp*.*,

and *Selenastrum* sp., which resulted in an initial chlorophyll-*a* concentration of approximately

10 µg/L in all tanks. Algae were purchased from Carolina Biological and cultured in COMBO

medium prior to the experiment. A mixed assemblage of zooplankton was then added to each

tank. Zooplankton were collected from Sandy River Reservoir the day before the experiment by

towing a 64 µm mesh bongo net from 0 - 6 m several times. Prior to introduction, zooplankton

were concentrated into a single 4 L container, mixed, and then 50 mL aliquots were introduced

into each tank to provide a starting density of approximately 26 zooplankton per liter in each

tank. This initial relatively low density permitted observations of population growth over time

and reflected natural zooplankton concentrations in the early spring in local systems.

Proportionally, the zooplankton assemblage consisted of 80% copepods (mostly cyclopoids),

15% cladocerans (*Daphnia* sp. and *Bosmina* sp.), and 5% rotifers (*Keratella* sp.). We removed all

Chaoborus before introducing the zooplankton to the tanks to minimize zooplankton mortality

due to predation.

Over the course of the next month, phytoplankton biomass and zooplankton abundance

were measured at 0, 3, 7, 14, 21 and 28 days. Before each sampling, tanks were gently mixed

with a broad, plastic spoon. For phytoplankton, two replicate 50 mL water samples were

removed from each tank and filtered through Whatman GF/F filters to measure chlorophyll-*a*

concentration as a proxy for phytoplankton biomass. Filters were placed in 90% acetone

overnight in the freezer and then chlorophyll-*a* concentration was measured on a Shimadzu

Trilogy Fluorometer using the non-acidification module. Chlorophyll-*a* concentrations are

reported as the average of these two replicates. One hundred milliliters of DI water were added

to each tank to replace the 100 mL removed for chlorophyll-*a* analysis, keeping the volume at

~17 L.

For zooplankton, 3 L of water from each tank was passed through a 64 μm mesh cup,

and then the water was immediately returned to the tank to maintain a constant volume.

Zooplankton collected on the mesh were rinsed into a sample cup and preserved with 70%

ethanol. Zooplankton were identified and enumerated under a dissecting microscope at 40x in

a Ward counting wheel. Zooplankton density (individuals per liter) was determined by counting

the total number of rotifers, copepods, and cladocerans in each sample collection and then

dividing by the total volume of water sampled (i.e. 3 L).

Because bacteria can serve as an alternate food source for zooplankton, either directly

or through the microbial loop (Sanders and Porter 1990; Wylie and Currie 1991), we measured

bacterial abundance at Days 0 and 28. Five milliliters of water were collected from each tank

with a serological pipet, placed in a sterile culture tube, and preserved with glutaraldehyde.

Samples were refrigerated until analysis. Bacterial abundance was determined by counting

DAPI stained cells under an epifluorescent microscope at a magnification of 100x, based on the

methods of Porter and Feig (1980). A minimum of 400 cells were counted per sample to

determine cell density per milliliter.

After 28 days, two larval largemouth bass (*Micropterus salmonoides*), 10-12 mm in

standard length (i.e. body length excluding tail), were introduced to each tank. Preliminary

experiments determined the fish behaved better in pairs compared to single introductions. We

chose to introduce the larvae after one month to simulate the timing of fish hatching in nature

following an increase in algae and zooplankton in late spring (Mills et al. 1989; Mehner and

Thiel 1999; Hansson et al. 2007). Because we were unable to successfully obtain fish larvae

from Sandy River Reservoir prior to the experiment, we used larvae from a local pond with

similar optical characteristics. Fish larvae were collected using light traps (Aquatic Instruments,

Inc., Hope, ID) anchored in the littoral zone of the pond overnight. Fish were immediately

transported back to the lab and housed for 24 hours without food. Fish collection and care

followed approved institutional animal care and use protocols.

Once introduced to the experimental tanks, the fish larvae were allowed to feed for 24

hours. After which, zooplankton were sampled as described above. This provided an estimate

of daily zooplankton consumption in each treatment, assuming that total zooplankton

consumed was represented by the difference in zooplankton density at the beginning and end

of fish feeding. We also assumed that the two fish in each tank fed equally, such that the

difference in zooplankton density was divided by 2 to estimate daily consumption rates per fish

larva.

Fish were then allowed to feed in the tanks for another 5 days with survival monitored

daily. After which, fish larvae were euthanized in MS-222, and final measurements were taken

of their body length under a dissecting microscope to estimate growth. Because we could not

clearly distinguish the two fish in each tank, growth was estimated as the average size of the

two fish at the beginning versus the end of the experiment. A final collection was also made of

zooplankton density in the tanks. However, zooplankton abundance was too low in all the tanks

to make accurate counts.

Throughout the experiment, a YSI 600 XLM sonde was used to measure water

temperature, conductivity, pH, and dissolved oxygen concentration in each tank. To monitor

potential changes in water color over time, a Shimadzu UV/Vis spectrophotometer was used to

measure absorbance at 440 nm of water filtered through a GF/F filter. Dissolved organic carbon

(DOC) concentration was also monitored during the experiment using GF/F filtrate. Samples

were run on a Shimadzu TOC-L analyzer.

Direct Effects of Browning

The same experimental setup was used to investigate direct effects of browning on fish

foraging efficiency. However, we did not use a lake water control. Light to dark brown water

color treatments consisted of four replicate tanks assembled in a randomized, block design on

the three shelves. After adding SuperHume to the tanks, we noted that water color ranged

from a_{440} = 1.6 to 13.1 m^{-1}. In each tank, the pH was adjusted to approximately 7, and the

water was allowed to equilibrate to room temperature (22 - 23°C). A YSI 600 XLM sonde was

used to determine the water temperature, conductivity, dissolved oxygen concentration, and

pH of each tank prior to the beginning of each feeding experiment.

262   Both zooplankton and larval *Micropterus salmonoides* were collected from Sandy River

Reservoir. Fish larvae were starved for 24 hours prior to the experiment. For each experimental

tank, two larval fish (~13 mm in length) were introduced and allowed to acclimate for a

minimum of 1 hour. After which, a known concentration of zooplankton prey (i.e. ~ 20

zooplankton/L) was added, and the fish were allowed to feed for 30 minutes. We staggered the

introduction of the zooplankton every 10-15 minutes to allow time for disassembling each tank

at the end of the timed feeding trial. The zooplankton community consisted of approximately

40% cladocearans (mostly *Daphnia* sp.), 27% adult calanoid copepods, 24% *Chaoborus*, 7%

copepodids, and 2% adult cyclopoid copepods.

271   Fish were removed from the tank and placed in MS-222 for euthanization. The

remaining zooplankton were collected by filtering the water in each tank through a 64 μm

mesh. Zooplankton were rinsed off the mesh and into a sample cup with 70% ethanol.

Zooplankton were identified and enumerated under a dissecting microscope at 10 - 40x in a

Ward counting wheel. Zooplankton density per liter was calculated as the total number of

counted zooplankton divided by the total volume of water in the tank (i.e. 17 L). The total

zooplankton consumed per minute by each fish larvae was then calculated as the difference in

zooplankton density at the beginning versus the end of an experiment divided by 30 minutes

and then divided by 2 larvae. We assumed each fish in each tank fed equally during the

experiment. We chose this method rather than examining gut contents to be consistent with

previous experiments. Preliminary studies filling and emptying tanks with zooplankton resulted

in ~99% recovery, providing confidence in our methodology (unpublished data).

To further explore the direct effects of browning on larval fish foraging, we performed

three additional experiments using the same experimental setup but with bluegill *Lepomis*

machrochirus (~13-15 mm in standard length) collected from a local pond. The goal of these

experiments was to test the limits of larval fish foraging under increasing water color by

reducing the foraging time and number of prey. For each experiment, fish larvae were collected

with light traps 24 - 48 hours before the experiment. Two larvae were placed in each tank and

allowed to acclimate for 24 hours prior to the introduction of zooplankton prey. A YSI sonde

was used to confirm that temperature, pH, conductivity, dissolved oxygen concentrations were

similar across all tanks. For the first experiment, fish larvae fed for 24-hours with a relatively

high density of prey (i.e., 130 zooplankton per liter) to estimate daily feeding rates under ideal

conditions. We then conducted two experiments with reduced time and prey concentration to

test the limits of larval fish foraging: 1) a 10-minute feeding trial with 4 zooplankton per liter

and 2) a 5-minute feeding trial with 2 zooplankton per liter. Zooplankton prey remaining in each

tank were collected at the end of each experiment to assess fish feeding rates and prey

selectivity. For all three experiments, zooplankton were collected from the same pond as the

fish larvae and consisted of approximately 80% cladocearans (mostly *Daphnia* sp.), 15%

cyclopoid copepods, 3% *Chaoborus*, and 2% rotifers (*Keratella* sp. and *Asplanchna* sp.).

Statistical Analyses

All statistical tests were performed using the R Statistical Environment (R Core Team

2018). For the indirect effects experiment, we used the *nlme* package (Pinheiro et al. 2020) to

conduct repeated measures ANOVAs in combination with post hoc Tukey tests to assess

differences in water temperature, conductivity, pH, and dissolved oxygen with browning

treatment, time, and the interaction between time and treatment. Temperature and

conductivity data were log transformed for normality. ANOVAs were computed as linear mixed

models using the *lme* function, including terms for random effects associated with tank number

(i.e. *random* function) and, in some cases, autoregressive effects associated with time points

being unequally spaced (i.e. *corAR1* function). The *anova* function from the *car* package was

used to report the results of the models (Fox and Weisberg 2019). If significant, we used the *cld*

function in the *lsmeans* package (Lenth 2016) to summarize the Tukey results. Residuals from

each test were calculated using the *residuals* function and then plotted with the

plotNormalHistorgram function from the *rcompanion* package (Mangiafico 2020). Akaike

Information Criteria were used to select the best model, particularly if autoregressive

correlations improved the model.

Repeated measures ANOVA with post hoc Tukey tests were also performed on log

transformed chlorophyll-*a* concentrations to assess differences in algal biomass with treatment,

time, and interactions between treatment and time. The test was performed as described

above. In addition, linear regressions of chlorophyll-*a* concentrations in each treatment over

the first week (Days 0, 3, and 7) were performed to determine initial phytoplankton growth

rates, using the slope as an estimate of added algal biomass per day.

For the zooplankton, we observed similar densities across treatments except on the last

sampling date. We therefore assessed differences in zooplankton density only on Day 28 with a

Welch's ANOVA. This test is recommended when the data have high heteroscedasticity and is

paired with a post hoc Games-Howell test to compute pairwise comparisons of treatments. We

used the *welch_anova_test* function from the *rstatix* package (Kassambara 2020). Results from

the Welch's ANOVAs were used to compute effect sizes based on omega squared values.

A two-way ANOVA combined with a post hoc Tukey test was used to compare bacterial

abundance across treatments between Day 0 and Day 28. A Shapiro-Wilk's test confirmed data

normality prior to running the ANOVA. The *lme* function in the *nlme* package was used to run

the model, using tank number as a random variable. Summary results were presented with the

anova function and residuals were checked with the *residuals* and *plotNormalHistogram*

functions. Results of the Tukey test were observed using the *lsmeans* and *cld* functions as

described above.

For the fish data in the indirect effects experiment, we again observed unequal variance

across treatments, and therefore, used Welch's ANOVAs with post hoc Game-Howell tests to

assess potential differences in zooplankton consumption and growth with water color. Because

we noted variability in zooplankton abundance within our treatments, we also calculated

Pearson correlation coefficients between zooplankton prey availability at the time of fish

introduction and zooplankton consumption rates after 24 hours as well as larval fish growth

after 6 days. E*Indices were calculated to determine potential differences in prey selectivity

with browning (Lechowicz 1982). Values were checked for normality and then a two-way

ANOVA was performed to test for significant differences in prey selectivity between treatment,

zooplankton taxa, and interactions between treatment and taxa. Tukey pairwise comparisons

were performed for significant results.

For the direct effect experiment with largemouth bass, we observed differences in the

absorbance coefficients within treatments. We therefore used linear regression analysis to

assess the relationship between zooplankton consumption and water color (i.e. absorbance at

440 nm). For the bluegill experiments, absorbance coefficients were similar within treatments,

and thus, we assessed differences in zooplankton consumption across water color treatments

with Welch's ANOVAs, using omega squared values to estimate effect sizes. E*Indices were

again calculated, and two-way ANOVAs were performed to assess significant differences in prey

selectivity with treatment, zooplankton taxa, and their interaction. For the largemouth bass

experiment, this required us to bin tanks into light, moderate, and dark brown treatments

based on similarities in a_{440}, with 4 replicates per treatment.

Results

Indirect Effects of Browning

Over the course of the 28-day experiment, there were no significant differences in

water temperature, pH, or dissolved oxygen concentration among the 12 tanks with time or

treatment (Table 1). Conductivity did not significantly vary with time but was significantly lower

in the lake water treatment compared to the SuperHume treatments (Table 1). Absorbance at

440 nm was similar in the light brown and lake water treatments but significantly differed in the

moderate and dark brown treatments (Table 1). Water color decreased during the first week in

all treatments and then remained relatively constant (Figure S2). On average, absorbance

coefficients at 440 nm were 1.3 m^{-1} in the light brown and lake water treatments, 5.0 m^{-1} in the

moderate brown treatment, and 9.7 m^{-1} in the dark brown treatment. DOC concentration

significantly differed among the four treatments, with the light brown treatment having the

lowest DOC concentration (2.6 mg/L) and the lake water treatment having the highest DOC

concentration (5.1 mg/L) (Table 1). DOC concentration initially increased by 0.3 – 0.8 mg/L over

the first week of the experiment and then declined by ~0.5 mg/L over the next 3 weeks, except

in the lake water treatment, which did not significantly differ over the next three weeks (Figure

S2). Interestingly, only marginal increases in DOC concentration were observed with

SuperHume additions despite an approximate 10 times increase in water color. Higher DOC

concentrations in the lake water treatment, compared to the SuperHume treatments, were due

to high inputs of non-chromophoric, algal-derived organic carbon in the eutrophic reservoir.

Bacterial abundance significantly differed with time ($F_{(1)}$=32.95, p=0.0004, n=12) and

treatment ($F_{(3)}$=24.11, p < 0.0001, n=12). Based on the results of the Tukey test, bacterial

abundance in the lake water treatment (i.e., 2.86 x 10^6 cells per mL $\pm$ 1.4 x 10^5 S.E.) was

significantly greater than all other treatments (i.e., approximately 7.1 x 10^5 cells per mL $\pm$ 1.5 x

10^5 S.E. in the light brown treatment and approximately 1.3 x 10^6 cells per mL $\pm$ 9 x 10^5 S.E. in

the moderate and dark brown treatments). This suggests that not all bacteria were removed

from the lake water treatment during the initial set-up, and bacteria were introduced with the

addition of SuperHume. By the end of the experiment, bacterial abundance did not significantly

differ across the light, moderate, and dark brown treatments but was significantly higher in the

lake water treatment (i.e., ~ 3.19 x 10^6 cells per mL in the lake water treatment compared to ~ 2

x 10^6 cells per mL in the SuperHume treatments). Interestingly, bacterial abundance differed

by only 10 $\pm$ 9% between Day 0 and Day 28 in the lake water treatment while in the low,

moderate, and dark brown treatments, bacterial abundance was 64 $\pm$ 5%, 35 $\pm$ 5%, and 23 $\pm$ 3%

higher on Day 28, respectively.

Phytoplankton biomass, as estimated by chlorophyll-*a* concentration, significantly

differed with treatment ($F_{(3)}$= 33.79, p < 0.0001, n=12), time ($F_{(1)}$= 8.07, p=0.004, n=12), and the

interaction between treatment and time ($F_{(3)}$=20.18, p=0.0002, n=12) (Table 1; Figure 2). The

highest chlorophyll-*a* concentrations throughout the experiment were observed in the light

brown treatment (i.e., 46.1 µg/L by Day 28, Figure 2) while there was no significant difference

in the moderate brown, dark brown and lake water treatments (i.e., declining to 2 -7 µg/L by

Day 28, Figure 2). Based on slopes from regression analyses, algal growth during the first week

was fastest in the light brown treatment (5.61 µg/L per day $\pm$ 0.25 S.E.) followed by the lake

water (4.15 µg/L per day $\pm$ 0.22 S.E), moderate brown (2.36 µg/L per day $\pm$ 0.28 S.E), and dark

brown treatments (1.51 µg/L per day $\pm$ 0.2 S.E).

After 28 days, total zooplankton abundance was significantly different between

treatments ($F_{(3,3.35)}$=34.4, p=0.005, n=12, est ω^2= 0.89: 95% CI [0.0, 0.97]). Based on post hoc

pairwise comparisons, zooplankton in the lake water (mean = 190 individuals per L $\pm$ 25 S.E.)

and light brown (mean = 120 individuals per L $\pm$ 35 S.E.) treatments were not significantly

different (p>0.05), but the lake water treatment was significantly greater than the moderate

brown (mean = 72 individuals per L $\pm$ 7 S.E., p=0.02) and dark brown treatments (mean = 47

individuals per L $\pm$ 10 S.E., p = 0.003) (Figure 3A). Removing one outlier from the light treatment

resulted in significant differences between all three SuperHume treatments at the p $\leq$ 0.01

level. Overall, the cladoceran *Bosmina* sp. was the most abundant zooplankton in all treatments

by Day 28 (Figure 3B). Despite being the dominant zooplanktors at the beginning of the

experiment, copepods represented only 12% of the zooplankton community in the lake water

treatment and 5% of the zooplankton community in the SuperHume treatments by Day 28

(Figure 3D & 3E). *Daphnia* sp. abundance was low in all treatments by Day 28, except for the

light brown treatment (Figure 3C). While counting the zooplankton under the microscope,

flocculant SuperHume was observed, but not quantified, in the guts of cladocerans (i.e.,

Daphnia and *Bosmina*) but not calanoid or cyclopoid copepods (Figure 4). Rotifers rapidly

decreased in abundance and were not observed in any of the treatments after Day 7.

All fish survived the 6-day incubation in the lake water and light brown treatments.

However, there was a 33% mortality rate in both the moderate and dark brown treatments.

One dead fish was found in two replicate tanks of each treatment after 5-6 days, at which point

most zooplankton were visually depleted in the tanks. In the first 24 hours, fish consumed twice

as many prey in the lake water treatment (i.e. ~610 zooplankton per larva) compared to the

light brown treatment (~323 zooplankton per larva) and four times as many prey than in the

dark brown treatment (~143 zooplankton per larva). However, because of the variability in

zooplankton abundance within treatments, zooplankton consumption by larval fish did not

significantly differ with water color ($F_{(3,3.49)}$=4.79, p=0.09, n=12, est ω^2= 0.48: 95% CI [0.0, 0.78]).

Rather, zooplankton consumption in the first 24 hours correlated highly with zooplankton

densities at the time fish were introduced to the experimental tanks (Pearson Correlation

Coefficient =0.88, P=0.0002; Figure 5A). Proportionally, fish larvae primarily consumed *Bosmina*

in all treatments as they were the most numerous species in the tanks. However, *Daphnia* had

a higher electivity value (E*= 0.24 $\pm$ 0.0004), followed by *Bosmina* (E*= -0.12 $\pm$ 0.02) and then

copepods (E*= -0.64 $\pm$ 0.02). These differences in selectivity of zooplankton taxa were

statistically significant based on a two-way ANOVA ($F_{(2,20)}$=250.9, p < 0.0001, n=12). No

statistical differences in electivity were observed across treatment ($F_{(2,20)}$= 0.44, p = 0.78, n=12)

or the interaction between treatment and taxa ($F_{(8,20)}$=0.64, p = 0.73, n =12). Note that one dark

brown tank had to be eliminated from the two-way ANOVA because of a lack of *Daphnia* at the

time the fish were introduced.

Fish growth over the 6-day feeding experiment was generally greater in the lake water

and light brown treatments compared to the moderate and dark brown treatments; however,

these differences were not significantly different ($F_{(3,4.02)}$=3.17, p=0.15, n=12, est ω^2= 0.35: 95%

CI [0.0, 0.68]). On average, fish grew ~1.2 – 1.5 mm in the lake water and light brown

treatments, ~0.66 mm in the moderate brown treatment, and ~0.33 mm in the dark brown

treatment. Growth was moderately correlated with initial zooplankton abundance in the tank

(Pearson Correlation Coefficient =0.62, P=0.03; Figure 5B). After 6 days, the surviving fish larvae

in the moderate and dark brown treatments displayed empty and concave stomachs while

those in the lake water and light brown treatments were full with zooplankton prey (Figure 5C).

Note that previous studies have reported that it takes 4 – 6 hours for larvae to fully evacuate

their guts (Werner 1969).

Direct Effects of Browning

In the experiment with largemouth bass, we observed no significant differences in fish

foraging efficiency with increased browning over the 30-minute feeding trial based on linear

regression (Figure 6). Fish consumed on average 2.5 zooplankton per minute $\pm$ 0.3 S.E. across all

treatments, consuming mostly *Daphnia* sp. (percent consumed = 61% $\pm$ 0.03 S.E.) and

Chaoborus sp. (percent consumed = 57% $\pm$ 0.07 S.E.) followed by copepods (percent consumed

= 0.07% $\pm$ 0.02 S.E.). E*Index values significantly differ across zooplankton taxa ($F_{(2,27)}$= 71.14, p

< 0.0001, n=12), with *Daphnia* sp. (E* Index= 0.19 $\pm$ 0.02 S.E.) and *Chaoborus* sp. (E* Index =

0.15 $\pm$ 0.03 S.E.) prey preferred over copepods (E* Index= -0.7 $\pm$ 0.07 S.E.). However, prey

selectivity did not significantly differ with treatment ($F_{(2,27)}$= 0.07, p = 0.93, n=12) or the

interaction between treatment and zooplankton taxa ($F_{(4,27)}$= 0.38, p = 0.82, n=12).

Similar patterns were observed in the experiments with bluegill (Tables 2 and 3).

Despite reducing zooplankton prey concentration and foraging time, no significant differences

in foraging rate were detected with increasing water color (Table 2). Similar to largemouth

bass, larval bluegill primarily consumed *Daphnia* sp. and *Chaoborus* sp. prey compared to

copepods in all treatments (Table 3). In the 24-hour experiment, larval bluegill in all treatments

consumed ~80% of the zooplankton in their respective tanks, with low consumption rates for

rotifers (Table 2).

Discussion

The transition from endogenous to exogenous feeding is a 'critical period' in larval fish

development, characterized by high mortality rates that influence longer-term population

growth (Fuiman and Werner 2002). Here, we demonstrate that browning may add stress to this

critical stage by decreasing zooplankton prey availability. Browning did not, however, directly

alter larval fish feeding efficiency or prey selectivity, suggesting plasticity in foraging behavior

under varying light conditions. To our knowledge, this is the first study examining the effects of

freshwater browning on larval fishes. Understanding the balance between direct versus indirect

effects of browning on early life history stages will improve fish conservation and management

strategies in response to continued organic matter loading with changes in climate and land

use.

Indirect Effects of Browning

The transfer of energy through the traditional grazer food chain depends on basal

resources to support higher trophic levels. Therefore, one of the most profound effects of

freshwater browning is the reduction in primary production, due to increased light attenuation

(Wetzel 2001; Thrane et al. 2014; Solomon et al. 2015), that subsequently reduces energy flow

to zooplankton and fish (Jones et al. 2012; Solomon et al. 2015; Creed et al. 2018). Our study

supports these general observations and raises concerns for larval fish. With increasing brown

color, we observed a reduction in the quantity of light, decreased phytoplankton biomass, and

decreased zooplankton densities. In turn, the foraging rate, growth, and survival of larval

largemouth bass declined.

Starvation affects larval fish more than juvenile or adult stages because of their high

metabolic demands coupled with low energy reserves in their tissues (Fuiman 2002). Within 4 -

5 days at 25 - 30 °C, larvae will likely starve to death given low to no food rations (Fuiman

2002). In the present study, after 6 days at 22 - 23°C, fish larvae in the moderate and dark

brown treatments exhibited a 33% mortality rate while the surviving larvae in these treatments

displayed concave-shaped, empty stomachs. Although we did not quantify gut fullness, these

observations suggest a positive relationship between browning and starvation risk. Fish larvae

were found dead in the darker brown treatments near the end of the experiment (Day 5 or 6)

when zooplankton densities were already visibly depleted.

At the time fish larvae were introduced to our experimental tanks, zooplankton

densities in the moderate and dark brown treatments were 2 to 4 times less than in the light

brown and lake water treatments (i.e. 46 - 72 individuals per liter compared to 120 - 190

individuals per liter). Daily consumption rates for larval fish are often greater than a larvae's

own biomass (Post 1990). We did not directly measure zooplankton or larval fish biomass in

our study. However, based on published length-weight regressions (larval fish: Brecker 1993;

zooplankton: reviewed in Watkins et al. 2011), we roughly estimate that larval fish biomass was

~2020 μg dry weight per fish at the time of their introduction to the experimental tanks while

total zooplankton biomass was ~1577 $\pm$ 447 μg dry weight in the dark brown treatment, ~2965

$\pm$ 361 μg dry weight in the moderate brown treatment, ~6300 $\pm$ 1582 μg dry weight in the light

brown treatment, and ~4962 $\pm$ 228 μg dry weight in the lake water treatment. This resulted in a

ratio of zooplankton:fish biomass of 0.8, 1.5, 3.1, and 2.5 in the dark brown, moderate brown,

light brown, and lake water treatments, respectively. After 24 hours, fish larvae consumed

approximately 0.26, 0.54, 1.0, and 1.5 grams of zooplankton per gram of fish in the dark brown,

moderate brown, light brown, and lake water treatments, respectively. Based on data from

Houde and Zastrow (1990), freshwater fish should consume a minimum of ~0.5 - 0.7 grams of

prey per gram of fish per day at 23 ^{o}C to meet mean weight-specific growth rates. This suggests

that larvae in the moderate and dark brown treatments were food limited.

We placed two larvae in each tank to encourage routine behavior, which resulted in a

fish density of ~118 larvae m^{-3}. This is relatively high but not uncommon in nature, particularly

during the early months of fish hatching (Santucci et al. 2003). Previous studies have shown

that when zooplankton consumption outweighs zooplankton reproduction, fish larvae may

rapidly deplete their food source, affecting future growth and survival (Mehner and Thiel 1999;

Santucci et al. 2003; Hansson et al. 2007). This may occur more often in brown systems if

zooplankton abundances are low at the time of fish hatching. Furthermore, as surface water

temperatures rise as a result of global climate change, fish feeding rates are predicted to

increase as a consequence of higher metabolic rates (van Dorst et al. 2019). Brown waters are

likely to increase in temperature more rapidly due to the increased absorption of solar

radiation (Solomon et al. 2015), further exacerbating competition for limited resources in these

systems.

Unlike prey abundance, we did not see significant differences in zooplankton

community composition with increased water color. Others studies have also noted minimal

changes in zooplankton community composition with freshwater browning, and there does not

seem to be a clear pattern in positive versus negative effects on specific zooplankton groups

(Nicolle et al. 2012; Ekvall and Hansson 2012; Kelly et al. 2014; Robidoux et al. 2015; Lebret et

al. 2018; Leech et al. 2018). In the present study, the zooplankton community shifted from

copepod to cladoceran dominated across all treatments, with smaller-bodied *Bosmina* the

dominant zooplankton prior to larval fish introduction. Abundances of large-bodied *Daphnia*

were similar in all treatments during the first two weeks of the experiment, but for unknown

reasons, continued to increase only in the light brown treatment (i.e., ~ 30 individuals per L at

Day 28). It is possible that the light brown treatment provided the ideal diet for *Daphnia*, with

sufficient carbon and nutrients from both algal- and terrestrially-derived resources (Lennon et

al. 2013; Gall et al. 2017; Tang et al. 2018).

Overall, the zooplankton community that developed in our experiment provided a

standard diet for fish larvae. Typically, larger crustacean zooplankton promote foraging success,

growth, and survival (Crowder et al. 1987; Mills et al. 1989). However, for younger larvae that

are gape-limited, smaller prey, like *Bosmina* spp., can be more beneficial (Mehner and Thiel

1999). Nevertheless, zooplankton prey abundance in the moderate and dark brown treatments

was too low to support larval development.

Similar to previous studies (Batt et al. 2015; Karlsson et al. 2015; Solomon et al. 2015),

reduced zooplankton abundance coincided with reduced phytoplankton biomass as browning

increased. Interestingly, despite having similar water color, the light brown and lake water

treatments displayed opposite trends in phytoplankton biomass after Day 14. In the lake water

treatment, increased zooplankton abundance likely resulted in greater consumption of

phytoplankton biomass. However, we are uncertain why the same pattern did not occur in the

light brown treatment. It is possible that the light brown treatment, made with the COMBO

medium, provided phytoplankton with more nutrients for growth to keep up with zooplankton

consumption. We were unable to measure nutrient concentrations at the time of these

experiments; however, water quality monitoring data collected by the Virginia Department of

Environmental Quality for Sandy River Reservoir indicate generally lower nitrogen and

phosphorus concentrations than present in the COMBO medium.

Phytoplankton species composition and nutritional quality were not assessed as part of

our study; however, we recognize that these factors can also influence energy and nutrient

availability for zooplankton, and consequently larval fish (Taipale et al. 2018; Creed et al. 2018).

Because of logistics, we chose to use an artificial assemblage of green algal species that are

generally favorable foods. In nature, phytoplankton communities can shift towards

cyanobacteria with increased water color, which could further reduce zooplankton densities

(Ekvall et al. 2013; Robidoux et al. 2015). It is also possible that the algae used in our

experiment may not have been adapted to the low light conditions of brown water systems

given that they were purchased from Carolina Biological. However, our results are comparable

to field studies examining natural communities along a water color gradient (Ask et al. 2009;

Karlsson et al. 2009; Thrane et al. 2014).

Reductions in prey consumption, and possibly less nutritious prey, ultimately led to

reductions in fish growth. Fish larvae in the lake water and light brown treatments, on average,

grew twice as fast as the larvae in the moderate brown treatment and four times faster than

the larvae in the dark brown treatment. Although, we recognize that there was relatively high

unexplained variability in growth within our treatments (e.g. fish in two replicate tanks of the

moderate brown treatment exhibited no growth while the third replicate exhibited relatively

high growth). Growth rates are important for several reasons. Larval fish become better at

avoiding predators and detecting prey as swimming strength and visual acuity increases with

body size (Fuiman 2002). In addition, larger larvae consume bigger, more energy-rich prey.

Growth in juvenile and adult fish is often slower in brown compared to blue lakes, resulting in a

smaller length-at-age (Estlander et al. 2010; Horppila et al. 2010; Benoit et al. 2016; van Dorst

et al. 2018), and our data suggest similar patterns for larval fish. These reductions in individual

growth lead to further reductions in population growth and fish biomass as lakes darken in

color (Karlsson et al. 2009, 2015; Finstad et al. 2014), which can alter food web structure and

lower fisheries yields.

Importantly, the results of our indirect effects experiment were influenced by the timing

of fish introduction at Day 28. If the fish had been introduced at earlier time points, when

zooplankton densities were similar in all treatments, we likely would have observed no effect of

browning on larval fish. Indeed, in systems with low prey availability, fish foraging, growth, and

survival are reduced regardless of water color (Mehner and Thiel 1999; Santucci et al. 2003;

Hansson et al. 2007). Nonetheless, similar to our experimental results, browning has been

demonstrated in field and laboratory studies to reduce zooplankton densities (e.g. Ekvall et al.

2013; Robidoux et al. 2015; Leech et al. 2018). Moreover, it is often stated that these

differences in prey availability affect fish growth and survival, but experimental evidence is

limited. Here, we provide quantitative data on how alterations in prey densities, due to

increased browning, may affect larval fish foraging, growth, and survival in nature. We argue

that these valuable data can served as a starting point to design larger-scaled mesocosm

experiments or observational studies.

Direct Effects of Browning

When given an equal abundance of zooplankton prey, neither the foraging efficiency

nor prey selectivity of larval largemouth bass or bluegill were affected by increased browning.

Our results are similar to Stasko et al. (2012), which found no significant effect of browning on

juvenile roach feeding rates but are contrary to Jönsson et al. (2013), which reported decreases

in reactive distance and capture success of piscivorous Northern pike (*Esox lucius*) feeding on

roach (*Rutilus rutilus*) with increased water color. Weidel et al. (2017) reported significant

effects of browning on juvenile largemouth bass and bluegill foraging, but water color explained

only ~25-28% of the variation in foraging rates. Combined, results from our study and others

(Estlander et al. 2012; Ranaker et al. 2012; Nurminen et al. 2014) suggest that effects of

browning on fish foraging rates may be age- or species-specific.

Light intensities across all our treatments (i.e. ~15 – 35 μmol m^{-2} s^{-1} PAR) may have been

adequate for larval fish foraging, resulting in the lack of direct effects. For example, Miner and

Stein (1993) reported that larval bluegill successfully feed on zooplankton at light intensities

above 450 lux (i.e., ~8 µmol m^{-2} s^{-1} based on the conversion factor for sunlight). Some have

suggested that reductions in light levels have a greater effect on later life history stages of fish

because of the positive relationship between sighting distance and body size (Askne and Giske

1993; Fiksen et al. 2002). If browning does directly affect visual foraging, it may do so by

limiting the thickness and daily duration of the photic zone, such that there is a spatiotemporal

contraction of foraging habitat critical to growth and survival. Additionally, the metabolic costs

associated with searching for prey may increase with reductions in light levels, particularly if

prey availability declines with increased browning.

Larval fish used in our experiments may have been pre-adapted to the low-light

environment of brown water systems. In general, the man-made reservoirs and ponds of

central Virginia, USA have relatively high light attenuation due to increased cDOM inputs

and/or increased phytoplankton growth (Figure S1). Compared to local lakes and ponds in the

region, our dark brown treatment increased water color approximately 10 times. However, we

may have seen direct effects of browning on fish feeding had we continued to add organic

matter to the tanks. Moreover, fish inhabiting clear, blue lakes may show a greater response to

freshwater browning than those in our study (Stasko et al. 2015), and we encourage further

experimentation with larvae from these systems.

Because foraging success plays a critical role in survival and reproduction, there is likely

strong selection pressure to adapt to changing environmental conditions. As light availability for

visual foraging declines with browning, fish may shift from vision to other sensory mechanisms,

such as mechanoreception or olfactory cues, to detect zooplankton prey. For example, previous

research ablating the function of superficial neuromasts with neomycin or streptomycin has

shown reduced feeding rates in marine fish larvae (Jones and Janssen 1992; Cobcroft and

Pankhurst, 2003; Sampson et al. 2013). Larval zebrafish have been shown to learn to use

mechanoreception to feed in the dark (Carillo and McHenry 2016). There is also recent

evidence that juvenile bluegill may actually feed more in the open pelagic water column at

night and horizontally migrate towards the littoral zone during the day to avoid piscivorous

predators (Shoup et al. 2014). Future research should investigate the potential for alternative

feeding strategies as waters brown in color and be cautious about our biases as human

researchers relying on light and vision (Cumming et al. 2018).

Small Enclosures versus Natural Systems

We recognize that our experiment was conducted in relatively small containers and that

caution must be applied when scaling up results to natural systems. Nevertheless, our

observations of phytoplankton, zooplankton, and fish responses to browning are similar to

those reported by recent field-based studies (Karlsson et al. 2009, 2015; Finstad et al. 2014 and

others mentioned above), providing confidence in our results. Working with fragile fish larvae is

challenging, and smaller containers minimize some of the logistical constraints. Yet, the small,

confined tank could have influenced feeding rates. Given the size of larvae used in our

experiments (10 - 13 mm), their visual acuity, or reaction distance, is approximately one body

length (Werner 1969). However, we do not know how long it took larvae to search the tank.

Larger containers may have revealed greater differences in feeding rates across treatments,

similar to Weidel et al. (2017).

We also did not quantify the spatial distribution of zooplankton in the tanks, which

could have influenced feeding rates. Experimental tanks were covered during the feeding trials

to avoid distracting the fish. Zooplankton are known to swim downward in the presence of light

and fish, and it is possible that the zooplankton could have clumped at the bottom of the tanks,

making them easier to find and capture. However, this behavioral pattern in the zooplankton

was not obviously apparent to us while breaking down the tanks.

Interestingly, Seekell et al. (2018) reported that the relationship between fish biomass

and water color was more negative in deep lakes compared to shallow lakes within the boreal

region of Sweden. The authors state that the negative effects of browning associated with light

extinction and decreased primary production are minimized in shallow lakes because light often

reaches the lake bottom. Moreover, most fish inhabiting deeper lakes were observed in the

littoral zone. Moving into shallow waters may provide fish adequate light to forage in brown

waters, as supported by our results in shallow, experimental tanks (i.e., ~0.2 m depth).

Implications and Applications

Given increasing trends in temperature and precipitation with global climate change,

the browning of inland and coastal waters is predicted to continue, with far reaching

consequences for aquatic ecosystems (de Wit et al. 2016; Weyhenmeyer et al. 2016). Here, we

demonstrate that freshwater browning reduces energy flow to higher trophic levels, negatively

affecting larval fish growth and survival during a 'critical period'. Because of their ecological and

economic value, fish population dynamics are closely monitored and predicted by natural

resource managers and conservationists. While current population growth models focus on the

importance of temperature and nutrients (e.g., Deslauriers et al. 2017), our study and others,

highlight the need to incorporate water color (i.e. cDOM) to more accurately predict

recruitment strength, sustainable yields, and food web stability in systems affected by

browning.

Acknowledgements

Acknowledgements

We thank David and Navona Hart for granting access to their pond and Mark Rolfing for his

assistance in counting bacteria samples. Funding for this research came from the Longwood

University PRISM Summer Research Program and a faculty development grant to DML from

Longwood University. Collection of larval bluegill and largemouth bass was approved by the

Virginia Department of Game and Inland Fisheries, Permit# 051397. This manuscript was

improved by thoughtful comments from Sönke Johnsen, Tessa DeWalt, and two anonymous

reviewers.

References

Aksnes, D. L. and Giske, J. 1993. A theoretical model of aquatic visual feeding. Ecol. Model. **67**: 233-250.

Ask, J., Karlsson, J. and Jansson M. 2012. Net ecosystem production in clear-water and brown-water lakes. Glob. Biogeochem. Cyc. **26** 7. doi:10.1029/2010GB003951.

Bartels, P., Hirsch, P. E., Svanbäck, R. and Eklöv, P. 2016. Dissolved organic carbon reduces habitat coupling by top predators in lake ecosystems. Ecosystems **19**: 955-967.

Batt, R. D., Carpenter, S.R., Cole, J.J., Pace, M.L. Johnson, R.A., Kurtzweil, J.T. and Wilkinson, G.M. 2015. Altered energy flow in the food web of an experimentally darkened lake. Ecosphere **6**(3): 33. http://dx.doi.org/10.1890/ES14-00241.1.

Benoît, P.O., Beisner, B.E, and Solomon, C.T. 2016. Growth rate and abundance of common fishes is negatively related to dissolved organic carbon concentration in lakes. Can. J. Fish. Aq. Sci. **73**(8): 1230-1236. https://doi.org/10.1139/cjfas-2015-0340.

Breck, J. Hurry up and wait: Growth of young bluegill in ponds and in simulations with an individual-based model. Trans Amer Fish Soc 122: 467-480.

Bukaveckas, P.A. & Robbins-Forbes, M. 2000. Role of dissolved organic carbon in the attenuation of photosynthetically active and ultraviolet radiation in Adirondack lakes. Freshw. Biol. **43**(3): 339–354.

Carrillo, A, and McHenry, M.J. 2016. Zebrafish learn to forage in the dark. J. Exper. Biol. **219**: 582-**589**. doi: 10.1242/jeb.128918.

Carpenter, S.R., Cole, J.J., Kitchell, J.F. & Pace, M.L. 1998. Impact of Dissolved Organic Carbon, Phosphorus, and Grazing on Phytoplankton Biomass and Production in Experimental Lakes. Limnol. Oceanogr. **43**(1): 73–80.

Cole, J. J., Caraco, N.F., Kling, G.W. and Kratz, T.K. 1994. Carbon dioxide supersaturation in the surface waters of lakes. Science **265**: 1568–1570.

Cobcroft, J. M. and Pankhurst, P. M. 2003. Sensory organ development in cultured striped trumpeter larvae *Latris lineata*: implications for feeding behaviour. Mar. Freshw. Res. **54**: 669-682.

Craig, N., Jones, S.E., Weidel, B.C., and Solomon, C.T. 2017. Life history constraints explain negative relationship between fish productivity and dissolved organic carbon in lakes. Ecol. Evol. **7**(16): 6201–6209. doi:10.1002/ece3. 3108.

785 Creed, IF, Bergström, A-K, Trick, C.G., Grimm, N.B., Hessen, D.O., Karlsson, J. … Weyhenmeyer,
786   G.A. 2018. Global change-driven effects on dissolved organic matter composition:
787   Implications for food webs of northern lakes. Glob. Chan. Biol. **24**: 3692– 3714.
788   https://doi.org/10.1111/gcb.14129.
789
790 Crowder, L.B., McDonald, M.E., and Rice, J.A. 1987. Understanding recruitment of Lake
791   Michigan fishes: the importance of size-based interactions between fish and
792   zooplankton. Can. J. Fish. Aq. Sci. **44** (2): 141-147.
793
794 Cummings ME, Endler JA, Rebecca C. Fuller. 2018. 25 Years of sensory drive: the evidence and
795   its watery bias. Curr Zool. 64: 471-484.
796
797 Deslauriers, D., Rosburg, A.J., and Chipps, S.R. 2017. Development of a foraging model
798   framework to reliably estimate daily food consumption by young fishes Can. J. Fish. Aq.
799   Sci. **74**: 1668-1681. https://doi.org/10.1139/cjfas-2016-0331.
800
801 de Wit, H.A., Valinia, S., Weyhenmeyer, G.A., Futter, M.N., Kortelainen, P., Austnes, K., Hessen,
802   D.O., Räike, A., Laudon, H. and Vuorenmaa, H. 2016. Current Browning of Surface
803   Waters Will Be Further Promoted by Wetter Climate. Environ. Sci. Tech. Let. **3** (12): 430-
804   435. DOI: 10.1021/acs.estlett.6b00396.
805
806 Douglas, R.H. and Djamgoz, M. 1990. The Visual system of fish. Chapman and Hall.
807
808 Einem, J. von and Granéli, W. 2010. Effects of fetch and dissolved organic carbon on epilimnion
809   depth and light climate in small forest lakes in southern Sweden. Limnol. Oceanogr. **55**:
810   920–930.
811
812 Ekvall, M.K. and Hansson L-A. 2012. Differences in Recruitment and Life-History Strategy Alter
813   Zooplankton Spring Dynamics Under Climate-Change Conditions. *PLoS ONE* **7**(9):
814   e44614. https://doi.org/10.1371/journal.pone.0044614
815
816 Ekvall, M.K., De la Calle M.J., Faassen, E.J., Gustafsson, S., Lürling, M., and Hansson, L-A. 2013.
817   Synergistic and species-specific effects of climate change and water colour on
818   cyanobacterial toxicity and bloom formation. Freshw. Biol. **58**: 2414–2422.
819
820 Erlandsson, M., Buffam, I., Folster, J., Laudon, H., Temnerud, J., Weyhenmeyer, G.A., and
821   Bishop, K. 2008. Thirty-five years of synchrony in the organic matter concentrations of
822   Swedish rivers explained by variation in flow and sulphate. *Global Change Biol.* **14**:
823   1191–1198.
824
Estlander, S., Nurminen, L., Olin, M., Vinni, M., Immonen, S., Rask, M., and Lehtonen, H. 2010.
Diet shifts and food selection of perch *Perca fluviatilis* and roach *Rutilus rutilus* in humic
lakes of varying water colour. J. Fish Biol. **77**(1): 241– 256.
https://doi.org/10.1111/j.1095-8649.2010.02682.x.

Fiksen, Ø., Aksnes, D.L., Flyum, M.H. and Giske, J. 2002. The influence of turbidity on growth
and survival of fish larvae: a numerical analysis. Hydrobiologia **484**: 49–59.
https://doi.org/10.1023/A:1021396719733

Finstad, A. G., Helland, I.P., Ugedal, O., Hesthagen, T., and Hessen, D.O. 2014. Unimodal
response of fish yield to dissolved organic carbon. Ecol. Let. **17**: 36–43.

Fox J, Weisberg S. 2019. *An R Companion to Applied Regression*, Third edition. Sage, Thousand
Oaks CA. https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

Freeman, C., Evans, C.D, Monteith, D.T., Reynolds, B., and Fenner, N. 2001. Export of organic
carbon from peat soils. *Nature* **412**: 785–786. doi:10.1038/35090628.

Fuiman L.A. 2002. Special considerations of fish eggs and larvae, pp. 1-32. *In* L.A. Fuiman and
G.W. Robert (eds.), Fishery Science, The unique contributions of early life stages.
Blackwell publishing, 326 pp.

Fuiman, L.A. and R.G. Werner, eds. 2002. Fishery Science: The Unique Contributions of Early
Life Stages. Blackwell Science, Oxford.

Gall, A., Kainz, M., and Rasconi, S. 2017. *Daphnia magna* fitness during low food supply under
different water temperature and brownification scenarios. *J. Limnol.* **76**: 161-169. DOI:
10.4081/jlimnol.2016.1450.

Guthrie, D. M. and Muntz, W. R. A. 1993. Role of vision in fish behavior. *In* Behavior of Teleost
Fishes, 2nd edn (ed Pitcher, T. P.), pp. 89-121. London: Chapman and Hall.

Haaland, S., Hongve, D., Laudon, H., Riise, G., and Vogt, R.D. 2010. Quantifying the drivers of the
increasing colored organic matter in boreal surface waters. Environ. Sci. Tech. 44: 2975–
2980.

Hansson, L-A., Nicolle, A., Brodersen, J., Romare, P., Anders Nilsson, P., Brönmark, C., & Skov, C.
2007. Consequences of fish predation, migration and juvenile ontogeny on zooplankton
spring dynamics. *Limnol. Oceanogr.* **52**: 696–706.

Houde, E.D. and Zastrow, C.E. 1993, Ecosystem- and taxon-specific dynamic and energetics
properties of larval fish assemblages. *Bulletin of Marine Science* 53: 290-335.

Horppila, J., Olin, M., Vinni, M., Estlander, S., Nurminen, L., Rask, M., Ruuhijärvi, J. and
Lehtonen, H. 2010. Perch production in forest lakes: the contribution of abiotic and
biotic factors. Ecol. Freshw. Fish **19**: 257-266.

Jones, R.I. 1992. The influence of humic substances on lacustrine planktonic food chains.
Hydrobiolog. **229**: 73–91.
875
876 Jones, W. R. and Janssen, J. 1992. Lateral line development and feeding behavior in the mottled
877   sculpin, *Cottus bairdi* (Scorpaeniformes: Cottidae). *Copeia* **1992**: 485-492.
878
879 Kassambara, A. 2020. Rstatix: Pipe-Friendly framework for basic statistical tests.
880   https://rpkgs.datanovia.com/rstatix/
881
882 Karlsson, J., Byström, P., Ask, J., Ask, P., Persson, L. and Jansson, M. 2009. Light limitation of
883   nutrient-poor lake ecosystems. *Nature* **460**(7254): 506–509.
884
885 Karlsson, J., Bergstrom, A.K., Bystrom, P., Gudasz, C., Rodriguez, P., and Hein, C. 2015.
886   Terrestrial organic matter input suppresses biomass production in lake ecosystems.
887   Ecology 96: 2870– 2876. doi:10.1890/15-0515.1.
888
889 Keast, A., and Welsh L. 1968. Daily feeding periodicities, food uptake rates, and dietary changes
890   with hour of day in some lake fishes. *J. Fish. Res. Board. Can.* 25: 1133-1144.
891
892 Kilham, S.S., Kreeger, D.A., Lynn, S.G., Goulden, C.E., and Herrera, L. 1998. COMBO: a defined
893   freshwater culture medium for algae and zooplankton. *Hydrobiolog.* **377**: 147-159.
894
895 Kirk, J.T. O. 1994. Light and photosynthesis in aquatic ecosystems. Melbourne: Cambridge
896   University Press. http://www.loc.gov/catdir/toc/cam025/93037395.html
897
898 Koizumi S., Craig, N., Zwart, J.A., Kelly, P.T., Ziegler, J.P., Weidel, B.C., Jones, S.E., & Solomon,
899   C.T. 2018. Experimental whole-lake dissolved organic carbon increase alters fish diet and
900   density but not growth or productivity. Can. J. Fish. Aq. Sci. **75**: 1859-1867.
901   https://doi.org/10.1139/cjfas-2017-0283.
902
903 Kritzberg, E.S. and Ekström, S.M. 2012. Increasing iron concentrations in surface waters – a
904   factor behind brownification? *Biogeosciences* **9**(4): 1465–1478.
905
906 Lebret, K., Langenheder, S., Colinas, N., Östman, Ö., & Lindström, E.S. 2018. Increased water
907   colour affects freshwater plankton communities in a mesocosm study. Aq. Microb. Ecol*.*
908   **81**: 1–17. https://doi.org/10.3354/ame01858
909
910 Lechowicz, M.J. 1982. The sampling characteristics of electivity indices. *Oecologia* **52**: 22–30.
911
912 Leech, D.M. and Johnsen, S. 2009. Light and biological receptors. *In* Encyclopedia of Inland
913   Waters. Editor, G. Likens. Pp. 671-681.
914

Leech, D.M., Pollard, A.I., Labou, S.G., and Hampton, S.E. 2018. Fewer blue lakes and more
murky lakes in the U.S.: Implications for planktonic food webs. Limnol. Oceanogr. **63**:
2661-2680. doi.org/10.1002/lno.10967.

Lenth, R.V. 2016. Least-Squares Means: The R Package lsmeans. Journal of Statistical Software,
69(1), 1-33.doi:10.18637/jss.v069.i01

Mangiafico, S. 2020. Rcompanion: Functions to Support Extension Education Program
Evaluation. http://rcompanion.org

Mehner, T. and Thiel, R. 1999. A review of predation impact by 0+ fish on zooplankton in fresh
and brackish waters of the temperate northern hemisphere. *Environ. Biol. Fish.* **56**:
169–181.

Mills, E. L., Sherman, R. and Robson, D.S. 1989. Effect of zooplankton abundance and body size
on growth of age-0 yellow perch (*Perca flavescens*) in Oneida Lake, New York, 1975-
1986. Can. J. Fish. Aq. Sci. **46**: 880- 886.

Miner, J.G., and R.A. Stein. 1993. Interactive influence of turbidity and light on larval bluegill
(*Lepomis macrochirus*) foraging. Can. j. Fish. Aquat. Sci. **50**: 704-788.

Monteith, D.T., Stoddard, J.L., Evans, C.D., de Wit, Heleen A., Forsius, M., Høgåsen, T., Wilander,
A., Skjelkvåle, B.L., Jeffries, D.S., Vuorenmaa, J., Keller, B., Kopácek, J. and Vesely, J.
2007. Dissolved organic carbon trends resulting from changes in atmospheric deposition
chemistry. *Nature* **450**(7169): 537–540.

Morris D.P., Zagarese, H., Williamson, C.E., Balseiro, E.G., Hargreaves, B.R., Modenutti, B.,
Moeller, R., and Queimalinos, C. 1995. The attenuation of solar UV radiation in lakes and
the role of dissolved organic carbon. *Limnol. Oceanogr.* **40**: 1381–1391.

Nicolle, A., Hallgren P., Von Einem J., Kritzberg E.S., Graneli W., Persson A. … Hansson, L-A.
2012. Predicted warming and browning affect timing and magnitude of plankton
phenological events in lakes: an enclosure study. Freshw. Biol. 57: 684–695.

Nurminen, L, Estlander S, Olin M, and Lehtonen H. 2014. Feeding efficiency of planktivores
under disturbance, the effect of water colour, predation threat and shoal composition:
feeding of planktivores under disturbance. J. Fish Biol. **84**: 1195–1201.

Övergaard, R. 2019. The effects of different browning agents SuperHume, HuminFeed, and
Reverse Osmosis Concentrate on the fitness and survival of *Daphnia magna*
(Dissertation). Retrieved from http://urn.kb.se/resolve?urn=urn:nbn:se:uu:diva-392705

Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team (2020). *nlme: Linear and Nonlinear Mixed*
Effects Models. R package version 3.1-149. https://CRAN.R-project.org/package=nlme.
960

961 Porter, K.G. and Feig, Y. S. 1980. The use of DAPI for identifying and counting aquatic
962 microflora. Limnol. Oceanogr. **25**: doi: 10.4319/lo.1980.25.5.0943.
963

964 R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for
965 Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.
966

967 Ranåker L, Jönsson M, Nilsson PA, and Brönmark C. 2012. Effects of brown and turbid water on
968 piscivore–prey fish interactions along a visibility gradient. Freshw. Biol. **57**(9): 1761-
969 1768.
970

971 Sampson, J.A., Duston, J., and Croll, R.P. 2013. Superficial neuromasts facilitate non-visual
972 feeding by larval striped bass (*Morone saxatilis*). J. Exper. Biol. **216**: 3522-3530. doi:
973 10.1242/jeb.087395.
974

975 Santucci, V.J., Jr., and Wahl, D.H. 2003. The effects of growth, predation, and first-winter
976 mortality on recruitment of bluegill cohorts. Trans. Amer. Fish. Soc. **132**: 346-360.
977

978 Seekell, D. A., Lapierre, J.-F., Ask, J., Bergström, A.-K., Deininger, A., Rodríguez, P., & Karlsson, J.
979 2015. The influence of dissolved organic carbon on primary production in northern
980 lakes. Limnol. Oceanogr. **60**: 1276–1285. doi:10.1002/lno.10096.
981

982 Seekell, D. A., Byström, P. and Karlsson, J. 2018. Lake morphometry moderates the relationship
983 between water color and fish biomass in small boreal lakes. Limnol. Oceanogr. **63**: 2171-
984 2178. doi:10.1002/lno.10931
985

986 Shoup, D.E., Boswell, K.M. and Wahl, D.H. 2014. Diel Littoral–Pelagic Movements by Juvenile
987 Bluegills in a Small Lake. Trans. Amer. Fish. Soc. **143**(3): 796-801.
988 Solomon, C. T., S. E. Jones, B. C. Weidel, and others. 2015. Ecosystem Consequences of
989 Changing Inputs of Terrestrial Dissolved Organic Matter to Lakes: Current Knowledge
990 and Future Challenges. Ecosystems 18: 376–389. doi:10.1007/s10021-015-9848-y
991

Stasko, A.D., Gunn, J.M., and Johnston, T.A. 2012. Role of ambient light in structuring north
temperate fish communities: potential effects of increasing dissolved organic carbon
concentration with a changing climate. *Environ. Rev.* **20**(3): 173-190. doi:10.1139/a2012-
010.
Stasko AD, Johnston TA, and Gunn JM. 2015. Effects of water clarity and other environmental
factors on trophic niches of two sympatric piscivores. Freshw. Biol. 60: 1459–1474.
998

999 Taipale, S. J., Kahilainen, K. K., Holtgrieve, G. W., and Peltomaa, E. T. 2018. Simulated
1000 eutrophication and browning alters zooplankton nutritional quality and determines
1001 juvenile fish growth and survival. Ecol. Evol. **8**(5): 2671–2687. doi:10.1002/ece3.3832.

1002
1003   Tang, Y., Yang, X., Xu, R., Zhang, X., Liu, Z., Zhang, Y. and Dumont, H. J. 2019. Heterotrophic
1004   microbes upgrade food value of a terrestrial carbon resource for *Daphnia magna*. Limol.
1005   Oceanogr. 64: 474-482. doi:10.1002/lno.11052.
1006
1007   Thrane, J.-E., Hessen, D.O. and Andersen, T. 2014. The Absorption of Light in Lakes: Negative
1008   Impact of Dissolved Organic Carbon on Primary Productivity. *Ecosystems* 17: 1040–1052.
1009
1010   van Dorst, R.M., Gårdmark, A., Svanbäck, R., Beier, U., Weyhenmeyer, G.A., and Huss, M. 2019.
1011   Warmer and browner waters decrease fish biomass production. Global Change Biol. 25:
1012   1395– 1408. https://doi.org/10.1111/gcb.14551.
1013
1014   Vasconcelos, F.R., Diehl, S., Rodríguez, P., Karlsson, J., and Byström, P. 2018. Effects of
1015   terrestrial organic matter on aquatic primary production as mediated by pelagic–benthic
1016   resource fluxes. Ecosystems 21: 1255. https://doi.org/10.1007/s10021-017-0217-x.
1017
1018   Vasconcelos, FR, Diehl, S, Rodríguez, P, Hedström, P, Karlsson, J, and Byström, P. 2019. Bottom-
1019   up and top-down effects of browning and warming on shallow lake food webs. Global
1020   Change Biol. 25: 504– 521. https://doi.org/10.1111/gcb.14521.
1021
1022   Watkins, J., Rudstam, L., and Holeck, K. 2011. Length-weight regressions for zooplankton
1023   biomass calculations – A review and a suggestion for standard equations. Technical
1024   Report. Cornell University. Available at https://hdl.handle.net/1813/24566.
1025
1026   Weidel, B.C., Baglini, K., Jones, S.E., Kelly, P.T., Solomon, C.T., and Zwart, J.A. 2017. Light climate
1027   and dissolved organic carbon concentration influence species-specific changes in fish
1028   zooplanktivory. *Inland Waters* 7(2): 210-217. doi:10.1080/20442041.2017.1329121.
1029
1030   Werner, R.G. 1969. Ecology of Limnetic Bluegill (*Lepomis macrochirus*) Fry in Crane Lake,
1031   Indiana. *The American Midland Nat*uralist **81**(1): 164-181.
1032   Wetzel, R. G. 2001. Limnology: Lake and river ecosystems, 3rd ed. Academic Press.
1033
1034   Weyhenmeyer, G.A., Müller, R.A., Norman, M., and Tranvik, L.J. 2016. Sensitivity of freshwaters
1035   to browning in response to future climate change. Climatic Change 134: 225-239.
1036   https://doi.org/10.1007/s10584-015-1514-z.
1037
1038   Williamson, C. E., Overholt, E.P., Pilla, R.M., Leach, T.H., Brentrup, J.A., Knoll, L.B., Mette, E.M.
1039   and Moeller, R.E. 2015. Ecological consequences of long-term browning in lakes.
1040   Scientific Reports **5**: srep18666. doi:10.1038/srep18666.
1041
1042   Wylie, J.L. and Currie, D.J. 1991. The relative importance of bacteria and algae as food sources
1043   for crustacean zooplankton. Limnol. Oceanogr. **36**: 708-728. doi:
1044   10.4319/lo.1991.36.4.0708.
1045

1046 **Table 1**. Mean $\pm$ S.E. of major physical and chemical parameters across water color treatments.
1047 Significance is denoted in the last column, with, T = treatment effect, D =Day, I = Treatment:Day
1048 interaction, and NS = not significant. Lowercase letters indicate which treatments were
1049 significantly different from one another based on treatment effect.
1050

	Lake Water	Light Brown	Moderate Brown	Dark Brown	Significance
Temperature (ºC)	22.1 $\pm$ 0.18	22.09 $\pm$ 0.16	22.04 $\pm$ 0.16	22.03 $\pm$ 0.15	(T) NS (D) NS (I) NS
Dissolved Oxygen (mg/L)	9.05 $\pm$ 0.09	9.12 $\pm$ 0.09	8.89 $\pm$ 0.04	8.83 $\pm$ 0.03	(T) NS (D) <0.0001 (I) NS
Conductivity (mS/cm)	0.08 $\pm$ 0 a	0.25 $\pm$ 0.01 b	0.25 $\pm$ 0 b	0.25 $\pm$ 0.01 b	(T) <0.0001 (D) 0.01 (I) NS
pH	7.36 $\pm$ 0.04	7.25 $\pm$ 0.04	7.15 $\pm$ 0.22	7.14 $\pm$ 0.02	(T) NS (D) NS (I) NS
a440	1.77 $\pm$ 0.65 a	0.99 $\pm$ 0.08 a	4.84 $\pm$ 0.12 b	9.83 $\pm$ 0.17 c	(T) <0.0001 (D) 0.003 (I) NS
Chl-*a* (ug/L)	16.34 $\pm$ 2.69 ab	29.65 $\pm$ 3.83 b	12.4 $\pm$ 2.12 a	8.66 $\pm$ 1.57 a	(T) <0.0001 (D) 0.004 (I) 0.0002
DOC (mg/L)	5.22 $\pm$ 0.05 a	2.58 $\pm$ 0.07 b	3.40 $\pm$ 0.06 c	4.66 $\pm$ 0.07 d	(T) <0.0001 (D) 0.02 (I) NS

1059 **Table 2**. Results from the three foraging experiments with larval bluegill (*Lepomis machrochirus*), including the
1060 mean $\pm$ S.E. consumption rates. The approximate total number of each zooplankton type in the tanks at the start
1061 of a feeding trial is provided in parentheses in the first column. Welch's ANOVA results and omega-squared
1062 estimates of effect size were calculated for total zooplankton consumed only.
1063

	Consumption Rates (day^{-1} $larva^{-1}$ or min^{-1} $larva^{-1}$)		
	Light	Moderate	Dark
Experiment 1: 24 hours, ~130 zooplankton L^{-1}			
Large Cladocerans (210)	99 $\pm$ 4	101 $\pm$ 2	102 $\pm$ 1
Small Cladocerans (1817)	884 $\pm$ 7	876 $\pm$ 8	876 $\pm$ 11
Chaoborus (33)	23 $\pm$ 0	23 $\pm$ 0	23 $\pm$ 0
Copepods (47)	16 $\pm$ 0.1	16 $\pm$ 0.2	16 $\pm$ 0.2
Rotifers (77)	33 $\pm$ 1	34 $\pm$ 1	35 $\pm$ 0.1
Total (2183)	1055 $\pm$ 6	1051 $\pm$ 9	1052 $\pm$ 10
Welch's ANOVA	$F_{(2, 5.7)}$ = 0.081, p = 0.92, n = 12		
Omega Squared Estimate	est ω^2= 0.18: 95% CI [0.0, 0.68]		
Experiment 2: 10 minutes, ~4 zooplankton L^{-1}			
Large Cladocerans (39)	2 $\pm$ 0.03	2 $\pm$ 0.1	2 $\pm$ 0.03
Chaoborus (5)	0.3 $\pm$ 0	0.3 $\pm$ 0	0.25 $\pm$ 0
Copepods (23)	1 $\pm$ 0.02	1 $\pm$ 0.04	0.8 $\pm$ 0.1
Total (67)	3 $\pm$ 0.01	3 $\pm$ 0.2	2.8 $\pm$ 0.4
Welch's ANOVA	$F_{(2, 4.0)}$ = 1.53, p = 0.32, n = 12		
Omega Squared Estimate	est ω^2= 0.08: 95% CI [0.0, 0.47]		
Experiment 3: 5 minutes, ~2 zooplankton L^{-1}			
Large Cladocerans (22)	1.3 $\pm$ 0.2	1.5 + 0.08	1.4 $\pm$ 0.2
Small Cladoceran (7)	0.3 $\pm$ 0.1	0.5 + 0.1	0.38 $\pm$ 0.1
Copepods (11)	0.5 $\pm$ 0.2	0.7 + 0.2	0.5 $\pm$ 0.1
Total (40)	2 $\pm$ 0.3	2 $\pm$ 0.2	2 $\pm$ 0.5
Welch's ANOVA	$F_{(2, 5.3)}$ = 1.45, p = 0.31, n = 12		
Omega Squared Estimate	est ω^2= 0.08: 95% CI [0.0, 0.43]		

1064 **Table 3**. E* Index values from the three foraging experiments with larval bluegill (*Lepomis*
1065 *machrochirus*), including the mean $\pm$ S.E. for prey selectivity. The E* Index ranges from 1 to -1,
1066 where 1 represents a strong selection for a prey type, -1 equals strong avoidance of a prey type,
1067 and 0 equals random feeding. Two-way ANOVAs were performed to assess significant
1068 differences with treatment, zooplankton taxa, and their interaction.

	E* Index		
	Light	**Moderate**	**Dark**
Experiment 1: 24 hours, ~130 zooplankton L^{-1}			
Large Cladocerans	-0.007 $\pm$ 0.02	0.004 $\pm$ 0.004	0.007 $\pm$ 0.006
Small Cladocerans	0.01 $\pm$ 0.005	0.003 $\pm$ 0.005	0.003 $\pm$ 0.005
Chaoborus	0.02 $\pm$ 0.002	0.02 $\pm$ 0.005	0.02 $\pm$ 0.002
Copepods	0.01 + 0.003	0.006 $\pm$ 0.003	-0.01 $\pm$ 0.005
Rotifers	-0.04 $\pm$ 0.02	-0.04 $\pm$ 0.01	-0.03 $\pm$ 0.004
ANOVA Results:	Treatment: $F_{(2,45)}$ = 0.09; p = 0.91, n= 12		
	Zooplankton Taxa: $F_{(2,45)}$ = 20.62; p < 0.0001, n= 12		
	Treatment:Zooplankton: Taxa: $F_{(8,45)}$ = 0.84; p = 0.57, n= 12		
Experiment 2: 10 minutes, ~4 zooplankton L^{-1}			
Large Cladocerans	0.004 $\pm$ 0.008	0.006 $\pm$ 0.02	-0.03 $\pm$ 0.06
Chaoborus	0.01 $\pm$ 0.002	0.04 $\pm$ 0.02	0.06 $\pm$ 0.06
Copepods	- 0.02 $\pm$ 0.009	-0.05 $\pm$ 0.007	-0.06 $\pm$ 0.03
ANOVA Results:	Treatment: $F_{(2,27)}$ = 0.10; p = 0.91, n= 12		
	Zooplankton Taxa: $F_{(2,27)}$ = 4.81; p = 0.02, n= 12		
	Treatment:Zooplankton: Taxa: $F_{(4,27)}$ = 0.72; p = 0.58, n= 12		
Experiment 3: 5 minutes, ~2 zooplankton L^{-1}			
Large Cladocerans	0.16 $\pm$ 0.07	-0.01 $\pm$ 0.07	0.10 $\pm$ 0.06
Small Cladocerans	-0.23 $\pm$ 0.26	-0.05 $\pm$ 0.08	-0.2 $\pm$ 0.3
Copepods	-0.25 $\pm$ 0.29	-0.02 $\pm$ 0.03	-0.08 $\pm$ 0.07
ANOVA Results:	Treatment: $F_{(2,27)}$ = 0.15; p = 0.86, n= 12		
	Zooplankton Taxa: $F_{(2,27)}$ = 1.96; p = 0.17, n= 12		
	Treatment:Zooplankton: Taxa: $F_{(4,27)}$ = 0.48; p = 0.75, n= 12		

1069 **Figure Legends**

1070

1071 **Figure 1**. Spectral characteristics of the four treatments measured in photon counts per
1072 wavelength. Data were collected by an Ocean Optics Red Tide USB650 UV spectrophotometer
1073 from 250 to 800 nm with the sensor placed at the bottom of the tank pointing straight up
1074 toward the overhead grow lamps.

1075

1076 **Figure 2**. Mean phytoplankton biomass, as estimated by chlorophyll-*a* concentration, in the
1077 four treatments during the 28-day experiment. Error bars represent the standard error of three
1078 replicate tanks per treatment. Phytoplankton biomass increased in all treatments during the
1079 first 7 days but was comparatively lower in treatments with increased water color throughout
1080 the experiment.

1081

Figure 3. Zooplankton densities (individuals per Liter) in the four treatments during the 28-
day experiment. Error bars represent the standard error of three replicate tanks per treatment. The
top panel displays changes in total zooplankton densities (A) while the bottom panels display
individual zooplankton genera or groups, including the cladocerans *Bosmina* (B) and *Daphnia*
(C) as well as cyclopoid (D) and calanoid (E) copepods. In general, the zooplankton community
shifted from copepod to cladoceran dominated and abundance was significantly lower in
treatments with increased water color.

1089

1090 **Figure 4**. Flocculant organic matter was observed in the guts of cladocerans, including *Bosmina*
1091 and *Daphnia*, but not copepods in the moderate and dark brown treatments. Brown arrows
1092 highlight cladoceran gut contents. Image taken at 40x magnification.

1093

Figure 5. Daily prey consumption (A) and total growth over 6 days (B) in relation to zooplankton
prey availability at the time fish were introduced to the experimental tanks. Note that there are
2 points for the moderate brown treatment with ~1400 zooplankton prey and 0 mm growth.
After 6-days, surviving larval fish in the moderate and dark brown treatments displayed
concave empty stomachs while those in the light brown and lake water treatments had full
stomachs. Photos show example larvae from the light brown and dark brown treatments (C).
Red tick marks below fish are in millimeters.

1101

1102 **Figure 6**. Zooplankton consumption rates for larval largemouth bass with increasing water
1103 color, as measured by absorbance at 440 nm. When given equal prey concentrations in all
1104 treatments (i.e., 20 individuals per liter), there was no significant difference in the number of
1105 prey consumed during the 30-minute feeding trial. This suggests no direct effect of browning
1106 on visual foraging.

1113
1114   **Supplementary Materials**
1115
1116   **Figure S1**. Representative downwelling irradiance curves for photosynthetically active radiation
1117   (PAR) in Sandy River Reservoir and Hart Pond. Both are located near Farmville, VA, USA and
1118   were used as our study systems to collect larval fish and zooplankton. Measurements were
1119   taken with a Biospherical Instruments, Inc. BIC profiling radiometer in June 2014.
1120
1121   **Figure S2**. Mean $\pm$ S.E. of (A) a_{440} and (B) dissolved organic carbon concentrations over time
1122   during the Indirect Effects experiment. Data were collected at Day 0, 3, 7, 14, 21, and 28 as
1123   described in the Methods.

Figure 1

Figure 2

Figure 3

Copepods

Cladocerans

Figure 4

A — Total Zooplankton Consumed in 24 hrs vs. Zooplankton Prey Availability (Total # Individuals)

Legend: Light, Moderate, Dark, Lake

B — Growth (mm) vs. Zooplankton Prey Availability (Total # Individuals)

C

Light Brown

Dark Brown

Figure 5

Figure 6

Figure S1

Figure S2