1 **Indirect versus Direct Effects of Freshwater Browning on**
2 **Larval Fish Foraging**
3

4 Dina M. Leech, Troy L. Clift, Jessica L. Littlefield, Nicholas R. Ravagli, and Jacob E.
5 Spain
6
7 Department of Biological and Environmental Sciences
8 Longwood University Farmville, VA 23901
9
10 **Corresponding Author:** Dina M. Leech, Department of Biological and Environmental
11 Sciences, 201 High Street, Longwood University, Farmville, VA, 23909; Office #: 434.395.4966;
12 Email: leechdm@longwood.edu

13
14 **Keywords:** brownification, fish larvae, foraging, predation, zooplankton
15
16 **Running Title:** Browning and larval fish foraging

ABSTRACT

1) The larval stage in fish development is characterized as a 'critical period' with unusually high mortality rates. Additional stress may be imposed on fish larvae as fresh waters 'brown' in color due to increased inputs of dissolved organic matter from the terrestrial watershed. Both energy transfer and foraging efficacy are predicted to decrease as light availability declines with browning.

2) Here, we investigated the balance between indirect versus direct effects of browning on zooplankton-larval fish interactions. Experimental tanks were established in the laboratory where water color was altered with increasing concentrations of SuperHume (water color measured as absorbance at 440 nm = $1.6 - 10.8$ m^{-1}). Phytoplankton and zooplankton densities were monitored across treatments for one month to assess possible changes in community structure leading up to fish spawning (i.e., late Spring in central Virginia, USA). We then introduced larval largemouth bass to the tanks to assess indirect effects on larval fish feeding rate and growth. Additionally, direct effects on the foraging efficiency and selectivity of largemouth bass and bluegill were determined with separate short-term feeding experiments using the same experimental setup.

3) Browning did not directly alter the ability of larval fish to see and capture prey across our brown water treatments. However, significant indirect effects on larval fish foraging, growth, and survival were observed as phytoplankton and zooplankton densities decreased with increasing water color.

4) Our data suggest lake browning will reduce energy transfer to larval fish, during a critical stage of development, due to a reduction in prey availability but not visual foraging.

61   **1. INTRODUCTION**
62
In recent decades, many freshwater and coastal ecosystems in the Northern

Hemisphere have become browner in color due to an increased export of chromophoric

dissolved organic matter (cDOM) from the terrestrial watershed (Monteith et al. 2007;

Erlandsson et al. 2008; Haaland et al. 2010; Solomon et al. 2015). The mechanisms underlying

this 'browning' are currently debated and include changes in hydrology and land use, increased

global temperatures, reduced atmospheric acid deposition, and increased inputs of dissolved

iron (Freeman et al. 2001; Monteith et al. 2007; Erlandsson et al. 2008; Kritzberg and Ekström

2012). Whatever the mechanism, this phenomenon has far reaching ecological consequences

for the structure and function of aquatic ecosystems, including energy flow.

One concern is the effect of browning on the underwater light environment. As waters

become browner in color, the quantity of light in the water column is reduced, resulting in a

shallower euphotic zone (Bukaveckas and Robbins-Forbes 2000; Einem and Graneli 2010). In

addition, shorter wavelength ultraviolet and visible radiation are more readily absorbed by

cDOM compared to longer wavelengths (Morris et al. 1995; Wetzel 2001). Consequently, the

light environment shifts to the red portion of the visible spectrum. These alterations in the

quantity and quality of light have the potential to affect zooplankton-fish interactions--

indirectly through reduced energy transfer up the food chain and directly through reduced

foraging efficiency.

While recent studies have shown that the initial browning of low productive systems

can stimulate primary production (Ask et al. 2012; Seekell et al. 2015; Williamson et al. 2015),

excessive browning of fresh waters can reduce rates of photosynthesis due to competition for

photons (Kirk 1994). cDOM has been reported to sequester 10 times more photons within the

visible spectrum (400 - 700 nm) compared to photosynthetic pigments (Thrane et al. 2014). As

a consequence, areal primary production often decreases with increased water color (Jones

1992; Carpenter et al. 1998; Ask et al. 2009; Karlsson et al. 2009; Thrane et al. 2014), and

pelagic primary production typically exceeds benthic primary production (Vasconcelos et al.

2018; Vasconcelos et al. 2019). Phytoplankton community composition can also shift to

predominantly cyanobacteria, which are generally less nutritious or inedible to zooplankton

consumers (Ekvall et al. 2013; Robidoux et al. 2015). Overall, aquatic ecosystems often become

net heterotrophic with increased browning as bacterial production exceeds primary production

(Cole et al. 1994; Ask et al. 2012).

Reductions in the quantity and quality of basal resources then 'cascade up' the grazer

food chain to influence zooplankton and fish. For example, Robidoux et al. (2015) observed

decreases in crustacean zooplankton biomass and density with increased water color while

Craig et al. (2017) found that bluegill in lakes of increasing color were smaller in size and had

lower fecundity compared to those living in lakes with less color. Taipale et al. (2018) noted

that zooplankton and fish had poorer nutritional quality in browner systems because the

phytoplankton on which they fed had lower concentrations of essential fatty acids, proteins,

lipids, and carbohydrates. Combined, these results suggest that freshwater browning indirectly

alters energy flow to higher trophic levels, reducing ecosystem productivity.

Freshwater browning may also directly affect fish foraging behavior by altering the light

environment to which fish are adapted. Many fish are visually orienting predators, depending

on the quantity (i.e. intensity) and quality (i.e. spectra) of underwater light to locate and

capture prey (Guthrie and Muntz 1993; Leech and Johnsen 2009). However, results from

studies investigating direct effects of browning on fish foraging are varied. Some have noted

reductions in foraging efficiency as water color darkens (Estlander et al. 2010; Ranaker et al.

2012; Stasko et al. 2012; Jönsson et al. 2013; Weidel et al. 2017) while others have found no

effect of browning (Jönsson et al. 2103; Nurminen et al. 2014) or enhanced effects (Jönsson et

al. 2103). This suggests that the effects of browning on fish foraging may be species-specific. It

is also important to note that most studies have focused on juvenile or adult life history stages.

The larval stage represents an especially critical phase in fish development that

ultimately affects population growth and fish biomass (Hjort 1914; Fuiman and Werner 2002;

Karlsson et al. 2009; Karlsson et al. 2015). Because of their dependence on shorter wavelengths

of light for foraging, early life history stages may be particularly affected by freshwater

browning. Over evolutionary time, the visual system of fishes spectrally tunes to the intensity

and wavelengths of light present in their environment (Douglas and Djamgoz 1990). This

spectral tuning often varies with age and a fish's behavior. Larval fish spend most of their time

foraging in top few meters of the eplimnion, relying on short-wavelength ultraviolet and blue

radiation, while older fish predominantly forage at deeper depths on larger prey, relying on

longer wavelength green, yellow, and red light (Leech and Johnsen 2009).

Here, we use laboratory experiments to assess the indirect versus direct effects of

browning on larval fish foraging at the time of hatching (i.e., late spring/early summer in central

Virginia, USA). We hypothesized that increased browning will reduce phytoplankton biomass,

leading to a reduction in zooplankton abundance, and consequently larval fish foraging

efficiency, growth, and survival during an early, critical stage in development. We then used

the same laboratory set up to examine the direct effects of browning on fish feeding rate and

prey selectivity when given equal zooplankton prey, hypothesizing that fish larvae will consume

less prey as water color increases due to a reduction in light availability for foraging.

2. METHODS

2.1 Indirect Effects of Browning

Laboratory experiments were conducted in twelve 20 L glass aquaria assembled on

three shelves with four tanks per shelf. The entire shelving unit, including individual shelves,

was covered in black plastic with additional black plastic placed in between each tank. Light

was provided by 1.2 m long grow lamps containing two Lumichrome ® Full Spectrum Plus

fluorescent, 32W bulbs suspended approximately 10 cm above the water surface of each tank

on each shelf. While lamps cannot exactly reproduce the solar spectrum, these specific bulbs

were selected based on their broad coverage of the ultraviolet and visible specta. The relative

light intensity (i.e., photon count) in each tank was measured with an Ocean Optics Red Tide

USB650 UV spectrophotometer from 250 to 800 nm (Figure 1). Lights were placed on a 14-hr

light, 10-hr dark cycle.

Experimental tanks were arranged in a randomized block design, with at least one tank

from each treatment on each shelf. Nine tanks were filled with 17 L of artificial lake water made

from the COMBO medium recipe (Kilham et al. 1998). SuperHume, a commercially available

source of humic acid, was then added to the tanks at varying concentrations to adjust the

brown color of the water (Lennon et al. 2013). Three tanks received 7 µg/L of SuperHume to

serve as a light brown treatment (water color measured as absorbance at 440 nm (a_{440}) = 1.6 m$^-$

^{1}), three tanks received 33 µg/L to serve as a moderate brown treatment (a_{440} = 5.7 m^{-1}), and

three tanks received 66 µg/L to serve as a dark brown treatment (a_{440} = 10.8 m^{-1}). The pH of

each tank was then adjusted to 7 with approximately 2 µL of 2N HCl. The remaining three tanks

were filled with 17 L of lake water from Sandy River Reservoir, Farmville, VA to compare with

the artificial SuperHume treatments (a_{440} = 1.5 m^{-1}). Lake water was passed through a GF/F

filter to remove bacterio-, phyto-, and zooplankton, as best as possible. All tanks were allowed

to stabilize at room temperature, 22 - 23 ^{o}C.

At the beginning of the experiment, each tank received a dense mixture of

phytoplankton containing equal parts of *Ankistrodesmus* sp., *Chlorella* sp., *Scenedesmus* sp.,

and *Selenastrum* sp., which resulted in an initial chlorophyll-*a* concentration of approximately

10 µg/L in all tanks. Algae were purchased from Carolina Biological and cultured in COMBO

medium prior to the experiment. A mixed assemblage of zooplankton was then added to each

tank. Zooplankton were collected from Sandy River Reservoir the day before the experiment by

towing a 64 µm mesh bongo net from 0 - 6 m several times. Prior to introduction, zooplankton

were concentrated into a single 4 L container, mixed, and then 50 mL aliquots were introduced

into each tank to provide a starting density of approximately 26 zooplankton per liter in each

tank. This initial relatively low density permitted observations of population growth over time.

Proportionally, the zooplankton assemblage consisted of 80% copepods (mostly cyclopoids),

15% cladocerans (*Daphnia* sp. and *Bosmina* sp.), and 5% rotifers (*Keratella* sp.). We removed all

Chaoborus before introducing the zooplankton to the tanks to minimize zooplankton mortality

due to predation.

Over the course of the next month, phytoplankton biomass and zooplankton abundance

were measured at 0, 3, 7, 14, 21 and 28 days. Tanks were gently mixed with a broad, plastic

spoon before each sampling. For phytoplankton, two replicate 50 mL water samples were

removed from each tank and filtered through Whatman GF/F filters to measure chlorophyll-*a*

concentration as a proxy for phytoplankton biomass. Filters were placed in 90% acetone

overnight in the freezer and then chlorophyll-*a* concentration was measured on a Shimadzu

Trilogy Fluorometer using the non-acidification module. Chlorophyll-*a* concentrations are

reported as the average of these two replicates. One hundred milliliters of DI water were added

to each tank to replace the 100 mL removed for chlorophyll-*a* analysis, keeping the volume at

~17 L.

Because bacteria can serve as an alternate food source for zooplankton, either directly

or through the microbial loop (Sanders and Porter 1990; Wylie and Currie 1991), we measured

bacterial abundance at Days 0 and 28. Five milliliters of water were collected from each tank

with a serological pipet, placed in a sterile culture tube, and preserved with glutaraldehyde.

Samples were refrigerated until analysis. Bacterial abundance was determined by counting

DAPI stained cells under an epifluorescent microscope at a magnification of 100x, based on the

methods of Porter and Feig (1980). A minimum of 400 cells were counted per sample to

determine cell density per milliliter.

For zooplankton, 3 L of water from each tank was passed through a 64 μm mesh cup

and then poured back into the tank. Zooplankton collected on the mesh were rinsed into a

sample cup and preserved with 70% ethanol. Zooplankton were identified and enumerated

under a dissecting microscope at 40x in a Ward counting wheel. Zooplankton density

(individuals per liter) was determined by counting the total number of rotifers, copepods, and

cladocerans in each sample collection and then dividing by 3, or the total volume of water

sampled.

After 28 days, two larval largemouth bass (*Micropterus salmonoides*), 10-12 mm in

length, were introduced to each tank. Preliminary experiments determined the fish behaved

better in pairs compared to single introductions. Because we were unable to successfully obtain

fish larvae from Sandy River Reservoir prior to the experiment, we used larvae from a local

pond with similar optical characteristics. Fish larvae were collected using light traps (Aquatic

Instruments, Inc., Hope, ID) anchored in the littoral zone of the pond overnight. Fish were

immediately transported back to the lab and housed for 24 hours without food. Fish collection

and care followed approved institutional animal care and use protocols.

Once introduced to the experimental tanks, the fish larvae were allowed to feed for 24

hours. After which, zooplankton were sampled as described above. This provided an estimate

of daily zooplankton consumption in each treatment, assuming that total zooplankton

consumed was represented by the difference in zooplankton density at the beginning and end

of fish feeding. We also assumed that the two fish in each tank fed equally, such that the

difference in zooplankton density was divided by 2 to estimate daily consumption rates per fish

larvae.

Fish were then allowed to feed in the tanks for another 5 days. After which, fish larvae

were euthanized in MS-222, and final measurements were taken of their body length under a

dissecting microscope to estimate growth. Because we could not clearly distinguish the two fish

in each tank, growth was estimated by first adding together the length of the two fish in each

tank at the beginning and end of the experiment. We then took the difference of these two

numbers and divided by 2 to estimate growth per larvae. A final collection was also made of

zooplankton density in the tanks. However, zooplankton abundance was too low in all the tanks

to make accurate counts.

Throughout the experiment, a YSI 600 XLM sonde was used to measure water

temperature, conductivity, pH, and dissolved oxygen concentration in each tank. To monitor

potential changes in water color over time, a Shimadzu UV/Vis spectrophotometer was used to

measure absorbance at 440 nm of water filtered through a GF/F filter. Dissolved organic carbon

(DOC) concentration was also monitored during the experiment using GF/F filtrate. Samples

were run on a Shimadzu TOC-L analyzer.

2.2 Direct Effects of Browning

The same experimental setup was used to investigate direct effects of browning on fish

foraging efficiency. However, we did not use a lake water control. Light to dark brown water

color treatments consisted of four replicate tanks assembled in a randomized, block design on

the three shelves. After adding SuperHume to the tanks, we noted that water color ranged

from a_{440} = 1.6 to 13.1 m^{-1}. In each tank, the pH was adjusted to approximately 7, and the

water was allowed to equilibrate to room temperature (22 - 23^{o}C). A YSI 600 XLM sonde was

used to determine the water temperature, conductivity, dissolved oxygen concentration, and

pH of each tank prior to the beginning of each feeding experiment.

Both zooplankton and larval *Micropterus salmonoides* were collected from Sandy River

Reservoir. Fish larvae were starved for 24 hours prior to the experiment. For each experimental

tank, two larval fish (~13 mm in length) were introduced and allowed to acclimate for a

minimum of 1 hour. After which, a known concentration of zooplankton prey (i.e. ~ 20

zooplankton/L) was added, and the fish were allowed to feed for 30 minutes. We staggered the

introduction of the zooplankton every 10-15 minutes to allow time for disassembling each tank

at the end of the timed feeding trial. The zooplankton community consisted of approximately

40% cladocearans (mostly *Daphnia* sp.), 27% adult calanoid copepods, 24% *Chaoborus*, 7%

copepodids, and 2% adult cyclopoid copepods.

Fish were removed from the tank and placed in MS-222 for euthanization. The

remaining zooplankton were collected by filtering the water in each tank through a 64 μm

mesh. Zooplankton were rinsed off the mesh and into a sample cup with 70% ethanol.

Zooplankton were identified and enumerated under a dissecting microscope at 10 - 40x in a

Ward counting wheel. Zooplankton density per liter was calculated as the total number of

counted zooplankton divided by 17, or the total volume of water in the tank. The total

zooplankton consumed per minute by each fish larvae was then calculated as the difference in

zooplankton density at the beginning versus the end of an experiment divided by 30, or the

number of minutes the fish fed, and then divided by 2, or per larvae. We assumed each fish in

each tank fed equally during the experiment. We chose this method rather than examining gut

contents to be consistent with previous experiments. Preliminary studies filling and emptying

tanks with zooplankton resulted in ~99% recovery, providing confidence in our methodology

(unpublished data).

To further explore the direct effects of browning on larval fish foraging, we performed

three additional experiments using the same experimental setup but with bluegill *Lepomis*

machrochirus (~13-15 mm in length) collected from a local pond. The goal of these experiments

was to test the limits of larval fish foraging under increasing water color by reducing the

foraging time and number of prey. For each experiment, fish larvae were collected with light

traps 24 - 48 hours before the experiment. Two larvae were placed in each tank and allowed to

acclimate for 24 hours prior to the introduction of zooplankton prey. A YSI sonde was used to

confirm that temperature, pH, conductivity, dissolved oxygen concentrations were similar

across all tanks. For the first experiment, fish larvae fed for 24-hours with a relatively high

density of prey (i.e., 130 zooplankton per liter) to estimate daily feeding rates under ideal

conditions. We then conducted two experiments with reduced time and prey concentration to

test the limits of larval fish foraging: 1) a 10-minute feeding trial with 4 zooplankton per liter

and 2) a 5-minute feeding trial with 2 zooplankton per liter. Zooplankton prey remaining in each

tank were collected at the end of each experiment to assess fish feeding rates and prey

selectivity. For all three experiments, zooplankton were collected from the same pond as the

fish larvae and consisted of approximately 80% cladocearans (mostly *Daphnia* sp.), 15%

cyclopoid copepods, 3% *Chaoborus*, and 2% rotifers (*Keratella* sp. and *Asplanchna* sp.).

2.3 Statistical Analyses

For the first experiment investigating indirect effects, repeated measures one-way

ANOVAs were used to assess potential differences in water temperature, conductivity, pH,

dissolved oxygen, and chlorophyll-*a* concentration over the course of the experiment in each

treatment. A two-way ANOVA was used to compare bacterial abundance across treatments

between Day 0 and Day 28. Total zooplankton density was assessed only on Day 28 with a one-

way ANOVA as densities were similar on all other dates.

For both the indirect and direct effects experiments, linear regressions were performed

to determine differences in fish feeding rates with increasing water color. Zooplankton

consumption served as the dependent variable and absorbance at 440 nm the independent

variable. We also used a linear regression to calculate differences in fish growth with increasing

water color during the 6-day feeding trial in the indirect effects experiment. Pearson correlation

coefficients were calculated to estimate the effect of zooplankton densities on larval fish

consumption rates and growth. E*Indices were calculated to determine potential differences in

prey selectivity with browning (Lechowicz 1982). All statistical tests were performed using the R

Statistical Environment (R Core Team 2018).

3. RESULTS

3.1 Indirect Effects of Browning

Over the course of the 28-day experiment, there were no significant differences in

water temperature, pH, or dissolved oxygen concentration among the 12 tanks with time or

treatment (Table 1). Conductivity did not significantly vary with time but was significantly lower

in the lake water treatment compared to the SuperHume treatments (Table 1). Absorbance at

440 nm was similar in the light brown and lakewater treatments but significantly different from

the moderate and dark brown treatments (Table 1). Water color decreased during the first

week in all treatments and then remained relatively constant. On average, absorbance

coefficients at 440 nm were 1.3 m^{-1} in the light brown and lake water treatments, 5.0 m^{-1} in the

moderate brown treatment, and 9.7 m^{-1} in the dark brown treatment. DOC concentration

significantly differed among the four treatments, with the light brown treatment having the

lowest DOC concentration (2.6 mg/L) and the lake water treatment having the highest DOC

concentration (5.1 mg/L) (Table 1). However, DOC concentration did not significantly vary over

time. Interestingly, only marginal increases in DOC concentration were observed with

SuperHume additions despite an approximate 10 times increase in water color. Higher DOC

concentrations in the lake water treatment, compared to the SuperHume treatments, were due

to high inputs of non-chromophoric, algal-derived organic carbon in the eutrophic reservoir.

Bacterial abundance was significantly higher on Day 28 compared to Day 0 across all

treatments (F=20.34, DF=1, P= 0.0002). The lake water treatment had an initial bacterial

abundance of 2.86 x 10^6 cells per mL $\pm$ 1.4 x 10^5 S.E. compared to approximately 7.1 x 10^5 cells

per mL $\pm$ 1.5 x 10^5 S.E. in the light brown treatment and approximately 1.3 x 10^6 cells per mL $\pm$ 9

x 10^5 S.E. in the moderate and dark brown treatments. This suggests that not all bacteria were

removed from the lake water treatment during the initial set-up, and bacteria were introduced

with the addition of SuperHume. By the end of the experiment, bacterial abundance did not

significantly differ across the light, moderate, and dark brown treatments but was significantly

higher in the lake water treatment (i.e., ~ 3.19 x 10^6 cells per mL in the lake water treatment

compared to ~ 2 x 10^6 cells per mL in the SuperHume treatments (F=24.61, DF=3, P>0.0001)).

Interestingly, bacterial abundance differed by only 10 $\pm$ 9% between Day 0 and Day 28 in the

lake water treatment while in the low, moderate, and dark brown treatments, bacterial

abundance was 64 $\pm$ 5%, 35 $\pm$ 5%, and 23 $\pm$ 3% higher on Day 28, respectively.

Phytoplankton biomass, as estimated by chlorophyll-*a* concentration, significantly

differed with treatment but not time (Figure 2). The highest chlorophyll-*a* concentrations

throughout the experiment were observed in the light brown treatment (i.e., 46.1 µg/L by Day

28, Figure 2) while there was no significant difference in the moderate brown, dark brown and

lake water treatments (i.e., declining to 2 -7 µg/L by Day 28, Figure 2). Based on slopes from

regression analyses, algal growth during the first week was fastest in the light brown treatment

(5.61 µg/L per day $\pm$ 0.25 S.E.) followed by the lake water (4.15 µg/L per day $\pm$ 0.22 S.E),

moderate brown (2.36 µg/L per day $\pm$ 0.28 S.E), and dark brown treatments (1.51 µg/L per day

$\pm$ 0.2 S.E).

After 28 days, total zooplankton density was significantly greater in the lake water

treatment (mean = 190 individuals per L $\pm$ 25 S.E.) followed by the light brown (mean = 120

individuals per L $\pm$ 35 S.E.), moderate brown (mean = 72 individuals per L $\pm$ 7 S.E.), and dark

brown treatments (mean = 47 individuals per L $\pm$ 10 S.E.) (Figure 3A). Overall, the cladoceran

Bosmina sp. was the most abundant zooplankton in all treatments by Day 28 (Figure 3B).

Despite being the dominant zooplanktors at the beginning of the experiment, copepods

represented only 12% of the zooplankton community in the lake water treatment and 5% of the

zooplankton community in the SuperHume treatments by Day 28 (Figure 3D & 3E). *Daphnia* sp.

abundance was low in all treatments by Day 28, except for the light brown treatment (Figure

3C). While counting the zooplankton under the microscope, flocculant SuperHume was

observed, but not quantified, in the guts of cladocerans (i.e., *Daphnia* and *Bosmina*) but not

calanoid or cyclopoid copepods (Figure 4). Rotifers rapidly decreased in abundance and were

not observed in any of the treatments after Day 7.

All fish survived the 6 day incubation in the lake water and light brown treatments.

However, there was a 33% mortality rate in both the moderate and dark brown treatments,

with one dead fish found in two replicate tanks of each treatment after 5-6 days. As predicted,

fish foraging rate decreased with increasing water color (Figure 5A). In the first 24 hours, fish

consumed twice as many prey in the lake water treatment (i.e. ~610 zooplankton per larva)

compared to the light brown treatment (~323 zooplankton per larva) and four times as many

prey than in the dark brown treatment (~143 zooplankton per larva). Daily zooplankton

consumption correlated highly with zooplankton densities at the time fish were introduced to

the experimental tanks (Pearson Correlation Coefficient =0.88, P=0.0002). Proportionally, fish

larvae primarily consumed *Bosmina* in all treatments as they were the most numerous species

in the tanks. However, *Daphnia* had a higher electivity value (E*= 0.24 $\pm$ 0.0004), followed by

Bosmina (E*= -0.12 $\pm$ 0.02) and then copepods (E*= -0.64 $\pm$ 0.02). No statistical differences in

electivity were observed across treatments.

Fish growth over the 6-day feeding experiment was greatest in the lake water and light

brown treatments followed by the moderate and dark brown treatments (Figure 5B). On

average, fish grew ~1.2 – 1.5 mm in the lake water and light brown treatments, ~0.66 mm in

the moderate brown treatment, and ~0.33 mm in the dark brown treatment. However, based

on linear regression, water color explained only 33% of the variation in larval fish growth (y= -

0.13x + 1.51, P=0.05), and growth was not significantly correlated with total zooplankton

consumption (Pearson Correlation Coefficient =0.45, P=0.15). After 6 days, the surviving fish

larvae in the moderate and dark brown treatments displayed empty and concave stomachs

while those in the lake water and light brown treatments were full with zooplankton prey

(Figure 5C). Note that previous studies have reported that it takes 4 – 6 hours for larvae to fully

evacuate their guts (Werner 1969).

3.2 Direct Effects of Browning

In the experiment with largemouth bass, we observed no significant effect of increased

browning on fish foraging efficiency or prey selectivity over the 30-minute feeding trial (Figure

6). Fish consumed on average 2.5 zooplankton per minute $\pm$ 0.3 S.E. across all treatments,

consuming mostly *Daphnia* sp. (percent consumed = 61% $\pm$ 0.03 S.E.) and *Chaoborus* sp.

(percent consumed = 57% $\pm$ 0.07 S.E.) followed by copepods (percent consumed = 0.07% $\pm$ 0.02

S.E.). E*Index values did not significantly differ across treatments, with *Daphnia* sp. (E* Index=

0.19 $\pm$ 0.02 S.E.) and *Chaoborus* sp. (E* Index = 0.15 $\pm$ 0.03 S.E.) prey preferred over copepods

(E* Index= -0.7 $\pm$ 0.07 S.E.).

Similar patterns were observed in the experiments with bluegill. Despite reducing

zooplankton prey concentration and foraging time, no significant differences in foraging rate or

prey selectivity were detected with increasing water color (Table 2). Similar to largemouth

bass, larval bluegill primarily consumed *Daphnia* sp. and *Chaoborus* sp. prey compared to

copepods in all treatments (Table 2). In the 24-hour experiment, larval bluegill in all treatments

consumed ~80% of the zooplankton in their respective tanks, with low consumption rates for

rotifers (Table 2).

4. DISCUSSION

The transition from endogenous to exogenous feeding is a 'critical period' in larval fish

development characterized by unusually high mortality rates that influence longer-term

population growth (Hjort 1914; Fuiman and Werner 2002). Here, we demonstrate that

browning will likely add stress to this critical stage by decreasing zooplankton prey abundance.

Browning did not, however, directly alter larval fish feeding efficiency or prey selectivity,

suggesting plasticity in foraging behavior under varying light conditions. Understanding the

balance between direct versus indirect effects of browning on the early life history of fishes will

improve conservation and management strategies in response to continued organic matter

loading to aquatic ecosystems with future changes in climate and land use.

4.1 Indirect Effects of Browning

The transfer of energy through the traditional grazer food chain depends on basal

resources to support higher trophic levels. Therefore, one of the most profound effects of

freshwater browning is the reduction in primary production, due to increased light attenuation

(Wetzel 2001; Thrane et al. 2014; Solomon et al. 2015), that subsequently reduces energy flow

to zooplankton and fish (Jones et al. 2012; Solomon et al. 2015; Creed et al. 2018). Our study

supports these general observations and raises concerns for larval fish. With increasing brown

color, we observed a reduction in the quantity of light, decreased phytoplankton biomass, and

decreased zooplankton densities. In turn, the foraging rate, growth, and survival of larval

largemouth bass significantly declined.

Starvation is more serious for larval fish than juvenile or adult stages because of their

high metabolic demands coupled with low energy reserves in their tissues (Fuiman 2002).

Within 4 - 5 days at 25 - 30 ^{o}C, larvae will likely starve to death given low to no food rations

(Fuiman 2002). In the present study, after 6 days at 22 - 23^{o}C, fish larvae in the moderate and

dark brown treatments exhibited a 33% mortality rate while the surviving larvae in these

treatments displayed concave-shaped, empty stomachs. Although we did not quantify gut

fullness, these observations suggest a positive relationship between browning and starvation

risk.

Fish larvae are primarily zooplanktivorous, and therefore, zooplankton abundance and

community composition (e.g. small- vs. large-bodied) are of critical importance to larval fish

survival and growth. Generally, increased primary production in the late Spring and early

Summer leads to increased zooplankton production. Shortly thereafter, many fish species begin

to spawn, matching larval fish hatching with an abundance of zooplankton prey (Mills et al.

1989; Mehner and Thiel 1999; Hansson et al. 2007). However, as fresh waters continue to

brown and energy flow to zooplankton is reduced, planktivorous fishes of all stages may be

forced to compete for fewer zooplankton prey or switch to alternate resources (Batt et al.

2015). Competition may be further intensified as fish density in the epilimnion increases due to

a shallowing of the thermocline with increased browning (Solomon et al. 2015; Koizumi et al.

2018). Fish larvae may be especially negatively affected as they are not as apt at locating and

capturing prey as larger juveniles and adults (Fuiman and Werner 2002).

At the time fish larvae were introduced to our experimental tanks, zooplankton

densities in the moderate and dark brown treatments were 2 to 4 times less than in the light

brown and lake water treatments (i.e. 46 - 72 individuals per liter compared to 120 - 190

individuals per liter). Daily consumption rates for larval fish are often greater than a larvae's

own biomass (Post 1990). We did not directly measure zooplankton or larval fish biomass in our

study. However, based on published length-weight regressions (reviewed in Watkins et al.

2011) and assuming average body length, we roughly estimate that zooplankton biomass was

~93 μg dry weight in the dark brown treatment, ~175 μg dry weight in the moderate brown

treatment, ~373 μg dry weight in the light brown treatment, and ~473 μg dry weight in the lake

water treatment at the time the fish larvae began to forage. Approximately 70-80% of the

zooplankton were consumed in their respective tanks, suggesting that zooplankton prey

availability was potentially low in all of our treatments.

We placed two larvae in each tank to encourage routine behavior, which resulted in a

fish density of ~118 larvae m^{-3}. This is relatively high but not uncommon in nature, particularly

during the early months of fish hatching (Santucci et al. 2003). Previous studies have shown

that when zooplankton consumption outweighs zooplankton reproduction, fish larvae may

rapidly deplete their food source, affecting future growth and survival (Mehner and Thiel 1999;

Santucci et al. 2003; Hansson et al. 2007). This may occur more often in brown systems if

zooplankton abundances are low at the time of fish hatching. Furthermore, as surface water

temperatures rise as a result of global climate change, larval fish feeding rates are predicted to

increase as a consequence of higher metabolic rates (Van Dorst 2017; Van Dorst et al. 2017).

Brown waters are likely to increase in temperature more rapidly due to the increased

absorption of solar radiation (Solomon et al. 2015), further exacerbating competition for limited

resources in these systems.

Unlike prey abundance, we did not see significant differences in zooplankton

community composition with increased water color. Others studies have also noted minimal

changes in zooplankton community composition with freshwater browning, and there does not

seem to be a clear pattern in positive versus negative effects on specific zooplankton groups

(Nicolle et al. 2012; Ekvall and Hansson 2012; Kelly et al. 2014; Robidoux et al. 2015; Lebret et

al. 2018; Leech et al. 2018). In the present study, the zooplankton community shifted from

copepod to cladoceran dominated across all treatments, with smaller-bodied *Bosmina* the

dominant zooplankton prior to larval fish introduction. Abundances of large-bodied *Daphnia*

were similar in all treatments during the first two weeks of the experiment, but for unknown

reasons, continued to increase only in the light brown treatment (i.e., ~ 30 individuals per L at

Day 28). It is possible that the light brown treatment provided the ideal diet for *Daphnia*, with

sufficient carbon and nutrients from both algal- and terrestrially-derived resources (Lennon et

al. 2013; Gall et al. 2017; Tang et al. 2018).

Conversely, copepod abundances declined in all treatments over the 28 days, especially

in treatments with SuperHume. Cladoceran zooplankton may have outcompeted copepods

because of their faster life cycle combined with their consumption of flocculant organic matter

and/or bacteria as phytoplankton biomass declined with browning (Tang et al. 2018).

Alternatively, SuperHume has been suggested to be toxic to zooplankton (Robidoux et al. 2015).

After our experiment, we reared gravid calanoid females and copepod nauplii in beakers with

COMBO medium, algae, and increasing concentrations of SuperHume to investigate possible

inhibitory effects. The results were variable, but we did see some females drop their egg sacs,

which remained unhatched, in the moderate and dark brown treatments (unpublished data).

We therefore cannot rule out potential toxic effects of SuperHume that affected our results.

However, the *in situ* zooplankton community of Sandy River Reservoir was also observed to

shift from copepod to cladoceran-dominated over time. For example, cladocerans were the

dominant zooplankton at the time the direct effect experiment was conducted with larval

largemouth bass.

Overall, the zooplankton community that developed in our experiment provided a

standard diet for fish larvae. Typically, larger crustacean zooplankton promote foraging success,

growth, and survival (Crowder et al. 1987; Mills et al. 1989). However, for younger larvae that

are gape-limited, smaller prey, like *Bosmina* spp., can be more beneficial (Mehner and Thiel

1999). Nevertheless, zooplankton prey abundance in the moderate and dark brown treatments

was too low to support healthy larval development.

Similar to previous studies (Batt et al. 2015; Karlsson et al. 2015; Solomon et al. 2015),

reduced zooplankton abundance coincided with reduced phytoplankton biomass as browning

increased. Interestingly, despite having similar water color, the light brown and lake water

treatments displayed opposite trends in phytoplankton biomass after Day 14. In the lake water

treatment, increased zooplankton abundance likely resulted in greater consumption of

phytoplankton biomass. However, we are uncertain why the same pattern did not occur in the

light brown treatment.

Phytoplankton species composition and nutritional quality were not assessed as part of

our study; however, we recognize that these factors can also influence energy and nutrient

availability for zooplankton, and consequently larval fish (Taipale et al. 2018; Creed et al. 2018).

Because of logistics, we chose to use an artificial assemblage of green algal species that are

generally favorable foods. In nature, phytoplankton communities can shift towards

cyanobacteria with increased water color, which could further reduce zooplankton densities

(Ekvall et al. 2013; Robidoux et al. 2015). It is also possible that the algae used in our

experiment may not have been adapted to the low light conditions of brown water systems

given that they were purchased from Carolina Biological. However, our results are comparable

to field studies examining natural communities along a water color gradient (Ask et al. 2009;

Karlsson et al. 2009; Thrane et al. 2014).

Reductions in prey consumption, and possibly less nutritious prey, ultimately led to

reductions in fish growth. Fish larvae in the lake water and light brown treatments, on average,

grew twice as fast as the larvae in the moderate brown treatment and four times faster than

the larvae in the dark brown treatment. Growth rates are important for several reasons. Larval

fish become better at avoiding predators and detecting prey as swimming strength and visual

acuity increases with body size (Fuiman 2002). In addition, larger larvae consume bigger, more

energy-rich prey. Growth in juvenile and adult fish is often slower in brown compared to blue

lakes, resulting in a smaller length-at-age (Estlander et al. 2010; Horppila et al. 2010; Benoit et

al. 2016; van Dorst et al. 2018), and our data suggest similar patterns for larval fish. These

reductions in individual growth lead to further reductions in population growth and fish

biomass as lakes darken in color (Karlsson et al. 2009, 2015; Finstad et al. 2014), which can alter

food web structure and lower fisheries yields.

4.2 Direct Effects of Browning

When given an equal abundance of zooplankton prey, neither the foraging efficiency

nor prey selectivity of larval largemouth bass or bluegill were affected by increased browning.

Our results are similar to Stasko et al. (2012), which found no significant effect of browning on

juvenile roach feeding rates but are contrary to Jönsson et al. (2013), which reported decreases

in reactive distance and capture success of piscivorous Northern pike (*Esox lucius*) feeding on

roach (*Rutilus rutilus*) with increased water color. Weidel et al. (2017) reported significant

effects of browning on juvenile largemouth bass and bluegill foraging, but water color explained

only ~25-28% of the variation in foraging rates. Combined, results from our study and others

(Estlander et al. 2012; Ranaker et al. 2012; Nurminen et al. 2014) suggest that effects of

browning on fish foraging rates may be age- or species-specific.

It is also possible that larval fish used in our experiments were pre-adapted to the low-

light environment of brown water systems. In general, the man-made reservoirs and ponds of

central Virginia, USA have high light attenuation due to increased cDOM inputs and/or

increased phytoplankton growth. Compared to local lakes and ponds in the region, our dark

brown treatment increased water color approximately 10 times. However, we may have seen

direct effects of browning on fish feeding had we continued to add organic matter to the tanks.

Moreover, fish inhabiting clear, blue lakes may show a greater response to freshwater

browning than those in our study (Stasko et al. 2015), and we encourage further

experimentation with fish larvae from these systems.

Because foraging success plays a critical role in survival and reproduction, there is likely

strong selection pressure to adapt to changing environmental conditions. As light availability for

visual foraging declines with browning, fish may rely on other sensory mechanisms, such as

mechanoreception or olfactory cues, to detect zooplankton prey. Non-visual feeding in fish

larvae is not well studied, but previous research ablating the function of superficial neuromasts

with neomycin or streptomycin has shown reduced feeding rates in marine fish larvae (Jones

and Janssen 1992; Cobcroft and Pankhurst, 2003; Sampson et al. 2013). Larval zebrafish have

been shown to learn to use mechanoreception to feed in the dark (Carillo and McHenry 2016).

There is also recent evidence that juvenile bluegill may actually feed more in the open pelagic

water column at night and horizontally migrate towards the littoral zone during the day to

avoid piscivorous predators (Shoup et al. 2014). Future research should investigate the

potential for alternative feeding strategies as waters brown in color.

4.3 Small Enclosures versus Natural Systems

We recognize that our experiment was conducted in relatively small experimental

containers and that caution must be applied when scaling up results to natural systems.

Nevertheless, our observations of phytoplankton, zooplankton, and fish responses to browning

are supported by recent field-based studies (Karlsson et al. 2009, 2015; Finstad et al. 2014 and

others mentioned above). Working with fragile fish larvae is challenging, and smaller containers

minimize some of the logistical constraints. Yet, the small, confined tank could have influenced

feeding rates. Given the size of larvae used in our experiments (10 - 13 mm), their visual acuity,

or reaction distance, is approximately one body length (Werner 1969). However, we do not

know how long it took larvae to search the tank. Larger containers may have revealed greater

differences in feeding rates across treatments, similar to Weidel et al. (2017). We also did not

quantify the spatial distribution of zooplankton in the tanks, which could have influenced

feeding rates. Experimental tanks were covered during the feeding trials to avoid distracting the

fish. Zooplankton are known to swim downward in the presence of light and fish, and it is

possible that the zooplankton could have clumped at the bottom of the tanks, making them

easier to find and capture. However, this behavioral pattern in the zooplankton was not

obviously apparent to us while breaking down the tanks.

Interestingly, Seekell et al. (2018) recently reported that the relationship between fish

biomass and water color was more negative in deep lakes compared to shallow lakes within the

boreal region of Sweden. The authors state that the negative effects of browning associated

with light extinction and decreased primary production are minimized in shallow lakes because

light often reaches the lake bottom. The authors further noted that most fish were observed in

the littoral zone of lakes. Moving into shallow waters may provide fish adequate light to forage

in brown waters, as supported by our results in shallow experimental tanks (i.e., ~0.2 m depth).

4.4 Implications and Applications

Given increasing trends in temperature and precipitation with global climate change,

the browning of inland and coastal waters is predicted to continue, with far reaching

consequences for aquatic ecosystems (de Wit et al. 2016; Weyhenmeyer et al. 2016). Here, we

demonstrate that freshwater browning reduces energy flow to higher trophic levels, negatively

affecting larval fish growth and survival during a 'critical period'. Because of their ecological and

economic value, fish population dynamics are closely monitored and predicted by natural

resource managers and conservationists. While current population growth models focus on the

importance of temperature and nutrients (e.g., Deslauriers et al. 2017), our study and others,

highlight the need to incorporate water color (i.e. cDOM) to more accurately predict

recruitment strength, sustainable yields, and food web stability in systems affected by

browning.

ACKNOWLEDGEMENTS

We thank David and Navona Hart for granting access to their pond and Mark Rolfing for his

assistance in counting bacteria samples. Funding for this research came from the Longwood

University PRISM Summer Research Program and a faculty development grant to DML from

Longwood University. Collection of larval bluegill and largemouth bass was approved by the

Virginia Department of Game and Inland Fisheries, Permit# 051397. This manuscript was

improved by thoughtful comments from Tessa DeWalt and Sönke Johnsen.

LITERATURE CITED

Ask, J., Karlsson, J. & Jansson M. (2012). Net ecosystem production in clear-water and brown-water lakes. *Global Biogeochemical Cycles*, 26, 7. doi:10.1029/2010GB003951.

Bartels, P., Hirsch, P. E., Svanbäck, R. & Eklöv, P. (2016). Dissolved organic carbon reduces habitat coupling by top predators in lake ecosystems. *Ecosystems*, 19, 955-967.

Batt, R. D., Carpenter, S.R., Cole, J.J., Pace, M.L. Johnson, R.A., Kurtzweil, J.T. & Wilkinson, G.M. (2015). Altered energy flow in the food web of an experimentally darkened lake. *Ecosphere*, 6(3), 33. http://dx.doi.org/10.1890/ES14-00241.1

Benoît, P.O., Beisner, B.E, Solomon, C.T. (2016). Growth rate and abundance of common fishes is negatively related to dissolved organic carbon concentration in lakes. *Canadian Journal of Fisheries and Aquatic Sciences*, 73(8), 1230-1236. https://doi.org/10.1139/cjfas-2015-0340

Bukaveckas, P.A. & Robbins-Forbes, M. (2000). Role of dissolved organic carbon in the attenuation of photosynthetically active and ultraviolet radiation in Adirondack lakes. *Freshwater Biology*, 43(3), 339–354.

Carrillo, A, & McHenry, M.J. (2016). Zebrafish learn to forage in the dark. *Journal of Experimental Biology*, 219, 582-589; doi: 10.1242/jeb.128918.

Carpenter, S.R., Cole, J.J., Kitchell, J.F. & Pace, M.L. (1998). Impact of Dissolved Organic Carbon, Phosphorus, and Grazing on Phytoplankton Biomass and Production in Experimental Lakes. *Limnology and Oceanography*, 43 (1), 73–80.

Cole, J. J., Caraco, N.F., Kling, G.W. & Kratz, T.K. (1994). Carbon dioxide supersaturation in the surface waters of lakes. *Science*, 265, 1568–1570.

Cobcroft, J. M. & Pankhurst, P. M. (2003). Sensory organ development in cultured striped trumpeter larvae *Latris lineata*: implications for feeding behaviour. *Marine and Freshwater Research*, 54, 669-682.

Craig, N., Jones, S.E., Weidel, B.C. & Solomon, C.T. (2017). Life history constraints explain negative relationship between fish productivity and dissolved organic carbon in lakes. *Ecology and Evolution*, 1–9. doi:10.1002/ece3.3108

Creed, IF, Bergström, A-K, Trick, C.G., Grimm, N.B., Hessen, D.O., Karlsson, J. … Weyhenmeyer, G.A. (2018). Global change-driven effects on dissolved organic matter composition: Implications for food webs of northern lakes. *Global Change Biology*, 24, 3692– 3714. https://doi.org/10.1111/gcb.14129.

670 Crowder, L.B., McDonald, M.E., & Rice, J.A. (1987). Understanding recruitment of Lake Michigan
671   fishes: the importance of size-based interactions between fish and zooplankton.
672   *Canadian Journal of Fisheries and Aquatic Sciences*, 44 (2), 141-147.
673

674 Deslauriers, D., Rosburg, A.J., Chipps, S.R. (2017). Development of a foraging model framework
675   to reliably estimate daily food consumption by young fishes. *Canadian Journal of
676   Fisheries and Aquatic Sciences*, 74, 1668-1681. https://doi.org/10.1139/cjfas-2016-0331.
677

678 de Wit, H.A. , Valinia, S., Weyhenmeyer, G.A., Futter, M.N., Kortelainen, P., Austnes, K., Hessen,
679   D.O., Räike, A., Laudon, H. & Vuorenmaa, H. (2016). Current Browning of Surface
680   Waters Will Be Further Promoted by Wetter Climate. *Environmental Science &
681   Technology Letters*, 3 (12), 430-435. DOI: 10.1021/acs.estlett.6b00396.
682

683 Douglas, R.H. & Djamgoz, M. (1990). The Visual system of fish. Chapman and Hall.
684

685 Einem, J. von & Granéli, W. (2010). Effects of fetch and dissolved organic carbon on epilimnion
686   depth and light climate in small forest lakes in southern Sweden. *Limnology and
687   Oceanography*, 55, 920–930.
688

689 Ekvall, M.K. & Hansson L-A. (2012). Differences in Recruitment and Life-History Strategy Alter
690   Zooplankton Spring Dynamics Under Climate-Change Conditions. *PLoS ONE* 7(9),
691   e44614. https://doi.org/10.1371/journal.pone.0044614
692

693 Ekvall, M.K., De la Calle M.J., Faassen, E.J., Gustafsson, S., Lürling, M., Hansson, L-A. (2013).
694   Synergistic and species-specific effects of climate change and water colour on
695   cyanobacterial toxicity and bloom formation. *Freshwater Biology*, 58, 2414–2422.
696

697 Erlandsson, M., Buffam, I., Folster, J., Laudon, H., Temnerud, J., Weyhenmeyer, G.A., & Bishop,
698   K. (2008). Thirty-five years of synchrony in the organic matter concentrations of Swedish
699   rivers explained by variation in flow and sulphate. *Global Change Biology*, 14, 1191–
700   1198.
701

702 Estlander, S., Nurminen, L., Olin, M., Vinni, M., Immonen, S., Rask, M., & Lehtonen, H. (2010).
703   Diet shifts and food selection of perch Perca fluviatilis and roach Rutilus rutilus in humic
704   lakes of varying water colour. *Journal of Fish Biology*, 77(1), 241– 256.
705   https://doi.org/10.1111/j.1095-8649.2010.02682.x.
706

707 Finstad, A. G., Helland, I.P., Ugedal, O., Hesthagen, T., & Hessen, D.O. (2014). Unimodal
708   response of fish yield to dissolved organic carbon. *Ecology Letters*, 17, 36–43.
709

710 Freeman, C., Evans, C.D, Monteith, D.T., Reynolds, B., & Fenner, N. (2001). Export of organic
711   carbon from peat soils. *Nature*, 412, 785–786. doi:10.1038/35090628.
712

713 Fuiman L.A., (2002). Special considerations of fish eggs and larvae, pp. 1-32. *In* L.A. Fuiman and
714   G.W. Robert (eds.), Fishery Science, The unique contributions of early life stages.
715   Blackwell publishing, 326 pp.
716
717 Fuiman, L.A., and R.G. Werner, eds. 2002. Fishery Science: The Unique Contributions of Early
718   Life Stages. Blackwell Science, Oxford.
719
720 Gall, A., Kainz, M., & Rasconi, S. (2017). *Daphnia magna* fitness during low food supply under
721   different water temperature and brownification scenarios. *Journal of Limnology*, 76,
722   161-169. DOI: 10.4081/jlimnol.2016.1450.
723
724 Guthrie, D. M. & Muntz, W. R. A. (1993). Role of vision in fish behavior. *In* Behavior of Teleost
725   Fishes, 2nd edn (ed Pitcher, T. P.), pp. 89-121. London: Chapman and Hall.
726
727 Haaland, S., Hongve, D., Laudon, H., Riise, G., & Vogt, R.D. (2010). Quantifying the drivers of the
728   increasing colored organic matter in boreal surface waters. *Environmental Science &*
729   *Technology*, 44, 2975–2980.
730
731 Hansson, L-A., Nicolle, A., Brodersen, J., Romare, P., Anders Nilsson, P., Brönmark, C., & Skov, C.
732   (2007). Consequences of fish predation, migration and juvenile ontogeny on
733   zooplankton spring dynamics. *Limnology and Oceanography*, 52, 696–706.
734
735 Hjort, J. (1914). Fluctuations in the great fisheries of northern Europe viewed in the light of
736   biological research. Rapports Et Proces-Verbaux De La Commission Internationale Pour
737   L'exploration Scientifique De La Mer 20 (1): 228.
738
739 Horppila, J., Olin, M., Vinni, M., Estlander, S., Nurminen, L., Rask, M., Ruuhijärvi, J. & Lehtonen,
740   H. (2010). Perch production in forest lakes: the contribution of abiotic and biotic factors.
741   *Ecology of Freshwater Fish*, 19, 257-266.
742
743 Jones, R.I. (1992). The influence of humic substances on lacustrine planktonic food chains.
744   *Hydrobiologia*, 229, 73–91.
745
746 Jones, W. R. & Janssen, J. (1992). Lateral line development and feeding behavior in the mottled
747   sculpin, Cottus bairdi (Scorpaeniformes: Cottidae). *Copeia*, 1992, 485-492.
748
749 Karlsson, J., Byström, P., Ask, J., Ask, P., Persson, L. & Jansson, M. (2009). Light limitation of
750   nutrient-poor lake ecosystems. *Nature*, 460(7254), 506–509.
751
752 Karlsson, J., Bergstrom, A.K., Bystrom, P., Gudasz, C., Rodriguez, P., & Hein, C. (2015). Terrestrial
753   organic matter input suppresses biomass production in lake ecosystems. *Ecology*, 96,
754   2870– 2876. doi:10.1890/15-0515.1.
755

756   Kilham, S.S., Kreeger, D.A., Lynn, S.G., Goulden, C.E., & Herrera, L. (1998). COMBO: a defined
757   freshwater culture medium for algae and zooplankton. *Hydrobiologia*, 377, 147-159.
758

759   Kirk, J.T. O. (1994). Light and photosynthesis in aquatic ecosystems. Melbourne: Cambridge
760   University Press. http://www.loc.gov/catdir/toc/cam025/93037395.html
761

762   Koizumi S., Craig, N., Zwart, J.A., Kelly, P.T., Ziegler, J.P., Weidel, B.C., Jones, S.E., & Solomon,
763   C.T. (2018). Experimental whole-lake dissolved organic carbon increase alters fish diet
764   and density but not growth or productivity. *Canadian Journal of Fisheries and Aquatic*
765   *Sciences*, 75, 1859-1867. https://doi.org/10.1139/cjfas-2017-0283.
766

767   Kritzberg, E.S. & Ekström, S.M. (2012). Increasing iron concentrations in surface waters – a
768   factor behind brownification? *Biogeosciences*, 9(4), 1465–1478.
769

770   Lebret, K., Langenheder, S., Colinas, N., Östman, Ö., & Lindström, E.S. (2018). Increased water
771   colour affects freshwater plankton communities in a mesocosm study. *Aquatic Microbial*
772   *Ecology*, 81, 1–17. https://doi.org/10.3354/ame01858
773

774   Lechowicz, M.J. (1982). The sampling characteristics of electivity indices. *Oecologia*, 52, 22–30.
775

776   Leech, D.M. & Johnsen, S. (2009). Light and biological receptors. *In* Encyclopedia of Inland
777   Waters. Editor, G. Likens. Pp. 671-681.
778

779   Leech, D.M., Pollard, A.I., Labou, S.G., & Hampton, S.E. (2018). Fewer blue lakes and more
780   murky lakes in the U.S.: Implications for planktonic food webs. *Limnology and*
781   *Oceanography*, 63, 2661-2680. doi.org/10.1002/lno.10967.
782

783   Mehner, T. & Thiel, R. (1999). A review of predation impact by 0+ fish on zooplankton in fresh
784   and brackish waters of the temperate northern hemisphere. *Environmental*
785   *Biology of Fishes*, 56, 169–181.
786

787   Mills, E. L., Sherman, R. & Robson, D.S. (1989). Effect of zooplankton abundance and body size
788   on growth of age-0 yellow perch (*Perca flavescens*) in Oneida Lake, New York, 1975-
789   1986. *Canadian Journal of Fisheries and Aquatic Sciences*, 46, 880- 886.
790

791   Monteith, D.T., Stoddard, J.L., Evans, C.D., de Wit, Heleen A., Forsius, M., Høgåsen, T., Wilander,
792   A., Skjelkvåle, B.L., Jeffries, D.S., Vuorenmaa, J., Keller, B., Kopácek, J. & Vesely, J. (2007).
793   Dissolved organic carbon trends resulting from changes in atmospheric deposition
794   chemistry. *Nature*, 450(7169), 537–540.
795

796   Morris D.P., Zagarese, H., Williamson, C.E., Balseiro, E.G., Hargreaves, B.R., Modenutti, B.,
797   Moeller, R., and Queimalinos, C. (1995). The attenuation of solar UV radiation in lakes
798   and the role of dissolved organic carbon. *Limnology and Oceanography*, 40, 1381–1391.
799

800   Nicolle A., Hallgren P., Von Einem J., Kritzberg E.S., Graneli W., Persson A. ... Hansson, L-A.
801   (2012). Predicted warming and browning affect timing and magnitude of plankton
802   phenological events in lakes: an enclosure study. *Freshwater Biology*, 57, 684–695.
803
804   Nurminen L, Estlander S, Olin M, & Lehtonen H. (2014). Feeding efficiency of planktivores under
805   disturbance, the effect of water colour, predation threat and shoal composition: feeding
806   of planktivores under disturbance. *Journal of Fish Biology*, 84, 1195–1201.
807
808   Porter, K.G. & Feig, Y. S. (1980). The use of DAPI for identifying and counting aquatic microflora.
809   *Limnology and Oceanography* 25: doi: 10.4319/lo.1980.25.5.0943.
810
811   R Core Team (2018). R: A language and environment for statistical computing. R Foundation for
812   Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.
813
814   Ranåker L, Jönsson M, Nilsson PA, & Brönmark C. (2012). Effects of brown and turbid water on
815   piscivore–prey fish interactions along a visibility gradient. *Freshwater Biology*, 57(9),
816   1761-1768.
817
818   Sampson, J.A., Duston, J., & Croll, R.P. (2013). Superficial neuromasts facilitate non-visual
819   feeding by larval striped bass (*Morone saxatilis*). *Journal of Experimental Biology*, 216,
820   3522-3530. doi: 10.1242/jeb.087395.
821
822   Santucci, V.J., Jr., & Wahl, D.H. (2003). The effects of growth, predation, and first-winter
823   mortality on recruitment of bluegill cohorts. *Transactions of the American Fisheries
824   Society*, 132, 346-360.
825
826   Seekell, D. A., Lapierre, J.-F., Ask, J., Bergström, A.-K., Deininger, A., Rodríguez, P., & Karlsson, J.
827   (2015). The influence of dissolved organic carbon on primary production in northern
828   lakes. *Limnology and Oceanography*, 60, 1276–1285. doi:10.1002/lno.10096.
829
830   Seekell, D. A., Byström, P. and Karlsson, J. (2018), Lake morphometry moderates the
831   relationship between water color and fish biomass in small boreal lakes. *Limnology and
832   Oceanography*, 63, 2171-2178. doi:10.1002/lno.10931
833
834   Shoup, D.E., Boswell, K.M. & Wahl, D.H. (2014) Diel Littoral–Pelagic Movements by Juvenile
835   Bluegills in a Small Lake. *Transactions of the American Fisheries Society* 143(3), 796-801.
836
837   Solomon, C. T., S. E. Jones, B. C. Weidel, and others. 2015. Ecosystem Consequences of
838   Changing Inputs of Terrestrial Dissolved Organic Matter to Lakes: Current Knowledge
839   and Future Challenges. Ecosystems 18: 376–389. doi:10.1007/s10021-015-9848-y
840
841   Stasko, A.D., Gunn, J.M., & Johnston, T.A. (2012). Role of ambient light in structuring north
842   temperate fish communities: potential effects of increasing dissolved organic carbon

843 concentration with a changing climate. *Environmental Reviews*, 20(3), 173-190.
844 doi:10.1139/a2012-010.
845
846 Stasko AD, Johnston TA, & Gunn JM. (2015). Effects of water clarity and other environmental
847 factors on trophic niches of two sympatric piscivores. *Freshwater Biology*, 60, 1459–
848 1474.
849
850 Taipale, S. J., Kahilainen, K. K., Holtgrieve, G. W., & Peltomaa, E. T. (2018). Simulated
851 eutrophication and browning alters zooplankton nutritional quality and determines
852 juvenile fish growth and survival. *Ecology and Evolution*, *8*(5), 2671–2687.
853 doi:10.1002/ece3.3832.
854
855 Tang, Y. , Yang, X. , Xu, R. , Zhang, X. , Liu, Z. , Zhang, Y. & Dumont, H. J. (2019). Heterotrophic
856 microbes upgrade food value of a terrestrial carbon resource for *Daphnia magna*.
857 *Limnology and Oceanography*, 64, 474-482. doi:10.1002/lno.11052.
858
859 Thrane, J.-E., Hessen, D.O. & Andersen, T. (2014). The Absorption of Light in Lakes: Negative
860 Impact of Dissolved Organic Carbon on Primary Productivity. *Ecosystems*, 17, 1040–
861 1052.
862
863 Van Dorst, R. M. (2017). How do warmer and darker waters influence population
864 dynamics in size structured fish communities? Aqua Introductory Research Essay 2017:2.
865 Swedish University of Agricultural Sciences, Department of Aquatic Resources,
866 Drottningholm Lysekil Öregrund. pp 48.
867
868 van Dorst, R.M., Gårdmark, A., Svanbäck, R., Beier, U., Weyhenmeyer, G.A., & Huss, M. (2019).
869 Warmer and browner waters decrease fish biomass production. *Global Change Biology*,
870 25, 1395– 1408. https://doi.org/10.1111/gcb.14551.
871
872 Vasconcelos, F.R., Diehl, S., Rodríguez, P., Karlsson, J., & Byström, P. (2018). Effects of terrestrial
873 organic matter on aquatic primary production as mediated by pelagic–benthic resource
874 fluxes. *Ecosystems*, 21, 1255. https://doi.org/10.1007/s10021-017-0217-x.
875
876 Vasconcelos, FR, Diehl, S, Rodríguez, P, Hedström, P, Karlsson, J, & Byström, P. (2019).
877 Bottom-up and top-down effects of browning and warming on shallow lake food webs.
878 *Global Change Biology*, 25, 504– 521. https://doi.org/10.1111/gcb.14521
879
880 Watkins, J., Rudstam, L., & Holeck, K. (2011). Length-weight regressions for zooplankton
881 biomass calculations – A review and a suggestion for standard equations. Technical
882 Report. Cornell University. Available at https://hdl.handle.net/1813/24566.
883
884 Weidel, B.C., Baglini, K., Jones, S.E., Kelly, P.T., Solomon, C.T., & Zwart, J.A. (2017). Light climate
885 and dissolved organic carbon concentration influence species-specific changes in fish
886 zooplanktivory. *Inland Waters*, 7(2), 210-217. doi:10.1080/20442041.2017.1329121.

887   Werner, R.G. (1969). Ecology of Limnetic Bluegill (*Lepomis macrochirus*) Fry in Crane Lake,
888   Indiana. *The American Midland Naturalist*, 81(1), 164-181.
889
890   Wetzel, R. G. 2001. Limnology: Lake and river ecosystems, 3rd ed. Academic Press.
891
892   Weyhenmeyer, G.A., Müller, R.A., Norman, M., & Tranvik, L.J. (2016). Sensitivity of freshwaters
893   to browning in response to future climate change. *Climatic Change*, 134, 225-239.
894   https://doi.org/10.1007/s10584-015-1514-z
895
896   Williamson, C. E., Overholt, E.P., Pilla, R.M., Leach, T.H., Brentrup, J.A., Knoll, L.B., Mette, E.M. &
897   Moeller, R.E. (2015). Ecological consequences of long-term browning in lakes. *Scientific
898   Reports* 5: srep18666. doi:10.1038/srep18666.
899
900   Wylie, J.L. & Currie, D.J. (1991). The relative importance of bacteria and algae as food sources
901   for crustacean zooplankton. *Limnology and Oceanography*, 36, 708-728. doi:
902   10.4319/lo.1991.36.4.0708.

932 **TABLE 1**. Mean $\pm$ S.E. of major physical and chemical parameters across the water color
933 treatments. Significance is denoted in the last column, with NS = not significant, T = significant
934 treatment effect, and D = significant time effect (i.e. day sampled). Lowercase letters indicate
935 which treatments were significantly different from one another.
936

	Lake Water	Light Brown	Moderate Brown	Dark Brown	Significance
Temperature (°C)	22.1 $\pm$ 0.18	22.09 $\pm$ 0.16	22.04 $\pm$ 0.16	22.03 $\pm$ 0.15	NS
Dissolved Oxygen (mg/L)	9.05 $\pm$ 0.09	9.12 $\pm$ 0.09	8.89 $\pm$ 0.04	8.83 $\pm$ 0.03	NS
Conductivity (mS/cm)	0.08 $\pm$ 0 a	0.25 $\pm$ 0.01 b	0.25 $\pm$ 0 b	0.25 $\pm$ 0.01 b	< 0.0001 (T)
pH	7.36 $\pm$ 0.04	7.25 $\pm$ 0.04	7.15 $\pm$ 0.22	7.14 $\pm$ 0.02	NS
a440	1.77 $\pm$ 0.65 a	0.99 $\pm$ 0.08 a	4.84 $\pm$ 0.12 b	9.83 $\pm$ 0.17 c	< 0.0001 (T) 0.008 (D)
Chl-*a* (ug/L)	16.34 $\pm$ 2.69 a	29.65 $\pm$ 3.83 b	12.4 $\pm$ 2.12 a	8.66 $\pm$ 1.57 a	< 0.0001 (T)
DOC (mg/L)	5.22 $\pm$ 0.05 a	2.58 $\pm$ 0.07 b	3.40 $\pm$ 0.06 c	4.66 $\pm$ 0.07 d	< 0.0001 (T)

TABLE 2. Results from the three foraging experiments with larval bluegill (*Lepomis*
machrochirus), including consumption rates and E* index values for prey selectivity. The E*
Index ranges from 1 to -1, where 1 represents a strong selection for a prey type, -1 equals
strong avoidance of a prey type, and 0 equals random feeding. The approximate total number
of each zooplankton type in the tanks at the start of a feeding trial is provided in parentheses in
the first column.
960

Zooplankton Type	Consumption Rates (day^{-1} larva^{-1} or min^{-1} larva^{-1})			E* Index		
	Light	Moderate	Dark	Light	Moderate	Dark
24 hours, ~130 zooplankton L^{-1}						
Large Cladocerans (210)	71	85	89	-0.05	0.02	0.04
Small Cladocerans (1817)	769	728	726	0.07	0.02	0.01
Chaoborus (33)	17	17	17	0.15	0.13	0.12
Copepods (47)	21	21	19	0.10	0.06	0.005
Rotifers (77)	10	15	18	-0.48	-0.35	-0.25
10 minutes, ~4 zooplankton L^{-1}						
Large Cladocerans (39)	2	2	1.7	0.006	0.18	0.11
Chaoborus (5)	0.25	0.25	2.5	0.01	0.17	0.17
Copepods (23)	1	0.85	1.7	-0.02	0.07	0.15
5 minutes, ~2 zooplankton L^{-1}						
Large Cladocerans (22)	1.4	1.6	1.5	0.15	0.03	0.10
Small Cladoceran (7)	0.25	0.45	0.38	-0.13	-0.02	-0.02
Copepods (11)	0.45	0.73	0.5	-0.07	-0.01	-0.10

961
962

FIGURE LEGENDS

Figure 1. Spectral characteristics of the four treatments measured in photon counts per
wavelength. Data were collected by an Ocean Optics Red Tide USB650 UV spectrophotometer
from 250 to 800 nm with the sensor placed at the bottom of the tank pointing straight up
toward the overhead grow lamps.

Figure 2. Mean phytoplankton biomass, as estimated by chlorophyll-*a* concentration, in the
four treatments during the 28-day experiment. Error bars represent the standard error of three
replicate tanks per treatment. Phytoplankton biomass increased in all treatments during the
first 7 days but was comparatively lower in treatments with increased water color throughout
the experiment.

Figure 3. Zooplankton densities (individuals per Liter) in the four treatments during the 28-day
experiment. Error bars represent the standard error of three replicate tanks per treatment. The
top panel displays changes in total zooplankton densities (A) while the bottom panels display
individual zooplankton genera or groups, including the cladocerans *Bosmina* (B) and *Daphnia*
(C) as well as cyclopoid (D) and calanoid (E) copepods. In general, the zooplankton community
shifted from copepod to cladoceran dominated and abundance was significantly lower in
treatments with increased water color.

Figure 4. Flocculant organic matter was observed in the guts of cladocerans, including *Bosmina*
and *Daphnia*, but not copepods in the moderate and dark brown treatments. Brown arrows
highlight cladoceran gut contents. Image taken at 40x magnification.

Figure 5. Both daily prey consumption (A) and total growth over 6 days (B) were negatively
correlated with water color, as measured by absorbance at 440 nm. After 6-days, surviving
larval fish in the moderate and dark brown treatments displayed concave empty stomachs
while those in the light brown and lake water treatments had full stomachs. Photos show
example larvae from the light brown and dark brown treatments (C). Red tick marks below fish
are in millimeters.

Figure 6. Zooplankton consumption rates for larval largemouth bass with increasing water
color, as measured by absorbance at 440 nm. When given equal prey concentrations in all
treatments (i.e., 20 individuals per liter), there was no significant difference in the number of
prey consumed during the 30-minute feeding trial. This suggests no direct effect of browning
on visual foraging.

Copepods

Cladocerans

A

Daily Zooplankton Consumption Larvae^{-1} vs Water Color (Absorption Coefficient at 440 nm)

y = -86.967x + 1056.9
R² = 0.4775

B

Larval growth (mm) Larvae^{-1} vs Water Color (Absorption Coefficient at 440 nm)

y = -0.1303x + 1.5126
R² = 0.3338

C

Light Brown

Dark Brown