The evolution of competitive ability for essential resources

Joey R. Bernhardt[1*], Pavel Kratina[2], Aaron Pereira[1], Manu Tamminen[3], Mridul K. Thomas[4], Anita Narwani[1]

[1]*Aquatic Ecology Department, Eawag, Überlandstrasse 133, CH-8600 Dübendorf, Switzerland*

[2]*School of Biological and Chemical Sciences, Queen Mary University of London, Mile End Road, London E1 4NS, United Kingdom.*

[3]*Department of Biology, University of Turku, Natura, University Hill, 20014 Turku, Finland*

[4]*Centre for Ocean Life, DTU Aqua, Technical University of Denmark, Kongens Lyngby, Denmark*

*Correspondence to: joanna.bernhardt@eawag.ch.

Keywords: Chlamydomonas reinhardtii, coexistence theory, competition, eco-evolutionary dynamics, phytoplankton, R*, resource competition theory

Data and code accessibility: Data and code are available at https://github.com/JoeyBernhardt/chlamee-r-star and will be deposited in the Dryad digital data repository should the manuscript be accepted. DNA sequences have been deposited in the Sequence Read Archive (SRA) under the BioProject ID PRJNA558172.

Conflicts of interest: The authors have no competing interests.

Funding: Funding was provided by a postdoctoral fellowship from the Nippon Foundation Nereus Program to JRB, European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska Curie grant agreement TROPHY No. 794264 to MKT, an Eawag postdoctoral fellowship, an Eawag Seed Grant and a Swiss National Science Foundation Project Grant (31003A_176069) to AN.

Abstract

Competition for limiting resources is among the most fundamental ecological interactions and has long been considered a key driver of species coexistence and biodiversity. Species' minimum resource requirements, their R^*, are key traits that link individual physiological demands to the outcome of competition. However, a major question remains unanswered -- to what extent are species' competitive traits able to evolve in response to resource limitation? To address this knowledge gap, we performed an evolution experiment in which we exposed *Chlamydomonas reinhardtii* for approximately 285 generations to seven environments in chemostats which differed in resource supply ratios (including nitrogen, phosphorus and light limitation) and salt stress. We then grew the ancestors and descendants in common garden experiments and quantified their competitive abilities for essential resources. We investigated constraints on trait evolution by testing whether changes in resource requirements for different resources were correlated. Competitive abilities for phosphorus improved in all populations, while competitive abilities for nitrogen and light increased in some populations and decreased in others. In contrast to the common assumption that there are trade-offs between competitive abilities for different resources, we found that improvements in competitive ability for a resource came at no detectable cost. Instead, improvements in competitive ability for multiple resources were either positively correlated or not significantly correlated. Using resource competition theory, we then demonstrated that rapid adaptation in competitive traits altered the predicted outcomes of competition. These results highlight the need to incorporate contemporary evolutionary change into predictions of competitive community dynamics over environmental gradients.

Introduction

Resource limitation and competition for limiting resources are among the most important drivers of population growth [1], species distributions [2,3] and biodiversity [4]. Resource competition theory (RCT, [1]) predicts that a few key resource traits, including the minimum resource level a population requires to maintain positive population growth (R^*), determine the outcome of competition over short time scales [5]. However, we still do not know how these resource traits evolve as populations adapt to new environments, especially in the context of organisms competing for essential resources such as light and nitrogen. This is an important gap in knowledge because rapid evolution may be able to alter competitive outcomes among species [6,7]. Understanding how evolutionary processes influence species' traits that are relevant to coexistence is therefore critical to understanding the ecological mechanisms that create and maintain biodiversity [8]. Evolutionary change in one or multiple competing species can increase the likelihood of coexistence by reducing differences in species' competitive abilities for a given resource (i.e. reducing 'fitness differences') and by altering the identity of the resource that each species finds most limiting (i.e. increasing 'niche differences'). Since we do not currently understand the potential constraints on the adaptation of essential resource-use traits, we cannot predict the degree to which evolution contributes to or prevents competitive coexistence.

Resource competition often acts as a strong selective agent that drives patterns of biodiversity and trait change via character divergence [9,10] and adaptive radiation [11]. Competition can select for individuals that are able to consume 'alternative' resources, or those that are not shared with other competitors [12]. Over time this results in adaptive trait divergence and niche differentiation [13]. What is less well appreciated though, is that when resources are essential, or non-substitutable, opportunities for niche differentiation are limited and competition cannot be avoided by character displacement because all competitors require the same limiting resources [7,14]. In this case, selection favours improved competitive ability, or a reduced

population-level R* for the shared limiting resource [14,15]. However, adaptation may

be constrained by physiological limits, genetic correlations between multiple traits [16],

or lack of genetic variation in resource traits [17]. These constraints may be particularly

strong in the case of adaptation to essential resource limitation because there are few

opportunities for divergence in adaptive strategies.

Trade-offs among species in competitive ability have been observed at large

evolutionary scales (i.e. across clades) [18,19]. Turnover in species abundances across

gradients of resource ratios suggests that these trade-offs structure species

distributions and patterns of biodiversity [1,20]. These trade-offs may arise due to

differences in the local conditions in which the traits evolved, or from biophysical or

genetic constraints that prevent individuals from optimizing several resource-use traits

simultaneously. There are at least two types of trade-offs which can govern resource

competition: gleaner-opportunist trade-offs, (Electronic Supplementary Material (ESM)

Figure S1) [21,22], and trade-offs in the ability to acquire different essential limiting

resources (e.g. light versus nitrogen or nitrogen versus phosphorus) [18,19,23,24].

Although the existence of trade-offs in resource-use traits has been demonstrated on a

macroevolutionary scale spanning large swaths of evolutionary time, the

microevolutionary processes by which they may arise and the mechanisms that

maintain them are still poorly understood.

To understand how essential resource competition traits evolve and how adaptation is

constrained, we used experimental evolution with a model organism, *Chlamydomonas*

reinhardtii. Experimental evolution allowed us to control the ecological conditions of

selection in chemostat, to isolate the effect of single limiting resources, and to

minimize confounding selective factors across treatments and replicates. We created

seven distinct selection environments in chemostats that varied either in the supply of

essential resources or salt concentration and quantified how populations' resource-

competition traits and salt tolerances evolved. We replicated the evolutionary

treatments across five ancestral populations in order to quantify heterogeneity in the

responses to selection, and the repeatability of evolutionary outcomes [25]. Using

whole genome resequencing of the ancestors and descendants of the evolution

experiment, we confirmed that the descendants had fixed mutations over the course of

the experiment, and were no longer genetically identical to the ancestors, suggesting

that the observed phenotypic changes have a genetic basis.

We tested three predictions:

1) When populations are exposed to limitation of essential resources, selection on

resource-use traits should reduce R^*, the minimum resource requirement.

Additionally, evolutionary changes in R^* should be larger in the genotypically

diverse population relative to the isoclonal populations [26] because adaptation

from standing genetic variation can occur more rapidly [27] than adaptation

acting on novel mutations [28]. Lastly, we predicted that salt stress, in addition

to resource limitation would lead to greater adaptive trait change, particularly

because stress can increase rates of mutation [29].

2) Adaptive trait change is subject to trade-offs. Trade-offs between competitive

abilities for different resources, gleaner-opportunist trade-offs, or trade-offs

between resisting salt stress and having a high growth rate or low R^* may

constrain or structure adaptive change in resource traits [15,19] and potentially

cause adaptation in one environment to come at a cost to performance in

another environment [30]. Alternatively, positively correlated competitive traits

may cause selection for a lower R^* for one resource to reduce R^* for another

(pleiotropic or correlated fitness benefits in low-resource environments) [30–32].

3) We predicted that if trade-offs in resource-use traits cause traits to diverge

across different selection environments, this would increase the chance that

populations selected in different environments could competitively co-exist.

Methods

Evolution experiment

We obtained a strain of *Chlamydomonas reinhardtii* (cc1690 wild type mt+) from the Chlamydomonas Center (chlamycollection.org). We selected four random colonies derived from single cells (hereafter referred to as Anc 2, Anc 3, Anc 4 and Anc 5), and inoculated them into liquid COMBO freshwater medium [33]. We randomly assigned seven chemostats to each of the four isoclonal ancestral populations (Anc 2-5) and the genotypically diverse population, cc1690. The seven chemostats assigned to each of the ancestral populations were then randomly assigned to one of seven treatments which we maintained for 285 days: COMBO (hereafter referred to as 'C'), nitrogen limitation ('N'), phosphorus limitation ('P'), light limitation ('L'), salt stress ('S'), biotically-depleted medium (i.e. medium previously used to grow seven other species of phytoplankton) ('B'), and a combination of salt stress and biotically-depleted medium ('BS'). Here we used the term 'population' to refer to Anc 2, Anc 3, Anc 4, Anc 5, cc1690 (the 'ancestors') as well as all of their descendant populations ('descendants'). In total there were five ancestral populations, and 32 descendant populations because three were lost to contamination. Detailed information on experimental evolution methods is available in the Supplementary Methods in the ESM (Appendix A).

Determination of R^* and salt tolerance

We determined the minimum resource requirements for positive population growth (R^*) for each population [1] via batch culture experiments. We defined N^* as the minimum nitrogen concentration and P^* as the minimum phosphorus concentration for positive population growth. We define I^* as the minimum light level required for positive population growth (similar to I_c in [34]). We estimated R^* by measuring population growth rates at ten resource levels for each of nitrogen, phosphorus and light for three days (see ESM Appendix A: Supplementary Methods: *'Competitive trait assays'* for more details on the resource levels, acclimation and measurements). We estimated 'consumption vectors' ([1]) for N and P via stoichiometry of exponentially growing populations [3], and cell size by measuring single cell lengths using a high throughput

imager (Biotek®Cytation 5), and calculating cell biovolume assuming cells were

spheres using 4/3*pi*radius^3.

In order to determine populations' R*, we modeled resource-dependent population

growth via a Monod curve [1,35]. We estimated the parameters of the Monod curve

directly from population-level time series of chlorophyll-*a* relative fluorescence units

measured over the resource gradients. We modeled the resource-dependent rate of

population growth, *r*, during the exponential phase as:

$$F(t) = F(0)e^{r(R)t} \tag{1a}$$

where $F(t)$ is the population-level RFU at time *t*, and $r(R)$ is given by:

$$r(R) = \mu max \left(\frac{R}{k_s + R} \right) \tag{1b}$$

using nonlinear least-squares regression with the *nls.LM* function in the *minpack.LM*

package [36] in R. Population growth rate, *r*, is a function of *μmax*, the maximum

population growth rate, *R*, the resource concentration, and k_s, the half-saturation

constant for population growth.

Using the estimated parameters of the Monod curve (i.e. Equation 1b), we estimated R*

as:

$$R^* = \frac{m \, k_s}{\mu max - m} \tag{2}$$

where *m* is the mortality rate, which we set to be 0.56/day to reflect the mortality

caused by dilution in chemostat experiments. To simplify our analyses, we used

Equations 1 and 2 to estimate minimum light requirements (I*), where *R* = irradiance.

We also included ESM Figure S2 with parameters estimated from an Eilers-Peeters

curve [37] for light.

To estimate the uncertainty in the Monod curve (Equation 1) fits, we determined

confidence intervals around the fitted Monod curves using non-parametric

bootstrapping of mean-centered residuals using the *nlsBoot* function with 999

iterations in the *nlstools* [38] package in R. We calculated 95% confidence intervals as

the range between the 2.5th and 97.5th quantiles.

We defined the salt tolerance as salt concentration at which growth rates are half their

maximum (which occurs at a salt concentration of zero). We estimated salt tolerance by

modeling population growth rates during the exponential phase, *r*, as a function of salt

concentration, *S*, using a simplified form of the logistic function:

$$r(S) = \frac{a}{1 + e^{-b(S-c)}}$$ (3)

where *a* (the upper asymptote), is the maximum population growth rate (not salt-

stressed), *b* is the decline in growth rate with increasing salt concentration and *c* is the

salt concentration at which growth rates are half their maximum, in g/L.

Quantifying trait change and testing for trade-offs

We tested for changes in R* between descendant and ancestral populations by

subtracting the ancestral trait value from the descendant trait value and quantifying

whether the 95% on the difference overlapped zero. We tested whether the change in

resource-use traits was greater in the genotypically diverse populations than the

isoclonal populations by comparing the 95% CI of the trait changes.

We tested for trade-offs between:

1. growth rates at high vs low supply of a given resource (ie. μmax vs R*, or a

244   gleaner-opportunist trade-off),

246   2. competitive abilities for different resources, or competitive ability and cell size,

247   and

3. changes in traits in multiple dimensions.

We tested for trade-offs using multiple linear regressions. We quantified competitive

ability for a given resource as $1/R^*$ [19], and tested for trade-offs among competitive

abilities for different resources (trade-off 2). In order to assess trade-offs among

multiple traits and cell size, we centered and scaled the variables using the *scale*

function in R (mean=0, standard deviation=1) so all variables could be compared on

the same scale. In all cases of multiple regression, we included ancestor ID as a fixed

effect to account for relationships among ancestors and descendants.

We tested for differences in multivariate trait change as a function of selection

treatment and ancestor using redundancy analysis (RDA) with the *capscale* function in

the R package *vegan* [39], version 2.5-4. Here we included all of the traits we

measured: R^*s, cell biovolume, consumption vectors (i.e. stoichiometry), and salt

tolerances. We used permutation tests (*anova.cca* in *vegan*) to test the statistical null

hypothesis that selection treatment and ancestor ID had no significant impact on any

independently varying linear combination of traits. We used the same approach to test

the effects of treatment on trait variation along the individual axes. We assessed which

descendant populations had diverged from their ancestors in different environments

using post-hoc Tukey tests using the *TukeyHSD* function in R.

Quantifying genetic changes associated with selection environments

DNA was extracted using a chloroform-methanol extraction and libraries were

prepared using the Bioo Scientific NEXTflex Rapid Illumina DNA-Seq Library Prep Kit.

For details and bioinformatic methods, please refer to the Supplementary Methods in

the ESM (Appendix A).

Testing the potential for altered predicted outcomes of competition

We used resource competition theory [1] to predict the outcome of pairwise

competition for two resources: nitrogen and phosphorus. RCT predicts that two

populations can coexist stably if they meet three conditions: 1) their zero net growth
isoclines (ZNGIs) cross (i.e. populations differ in the identity of the resource that most
limits their growth), 2), they each consume more of the resource which most limits their
growth (i.e. each population has a steeper consumption vector for the resource which
is most limiting to it) and 3), the supply point of resources in the environment falls
above their ZNGIs and between the consumption vectors of the two populations. If the
pair of populations meets criterion 1 and 3 but not 2, theory predicts unstable
coexistence or priority effects. If the pair of populations meets one or none of these
criteria, theory predicts competitive exclusion. We compared all possible combinations
of the five populations in their ancestral state and after selection in the different
resource environments. We then assessed the proportion of these pairwise interactions
that would be expected to lead to unstable coexistence, stable coexistence or
competitive exclusion.
292

Results

Evolutionary changes in R^* and salt tolerance

Relative to their ancestors, P^* declined in all five populations exposed to P limitation (P,
Figure 1A). Declines in P^* ranged from 43% to 85% across the replicate populations. In
response to N limitation, N^* declined in two populations (14%, 34% decline), did not
change in two populations and increased in one population (47% increase) (N, Figure
1B). I^* increased in two populations exposed to low light (L, 12%; 28% increase), and
did not change in the remaining three populations exposed to low light (Figure 1C).
Salt tolerance increased in all populations exposed to high salt (93% - 369% S and BS,
Figure 1D). Consumption vectors, quantified as the P:N molar ratio in the biomass of
populations growing exponentially, decreased in all of the populations subjected to
nitrogen limitation, and increased in four of the five populations exposed to
phosphorus limitation (ESM Figure S4, see also Figures S5-6). This suggests that
populations selected under nitrogen limitation contained more nitrogen relative to
phosphorus whereas populations selected under low phosphorus contained more
phosphorus relative to nitrogen. Contrary to our predictions, the descendants of the

genotypically diverse cc1690 population did not show more trait change than any of the isoclonal populations (triangles vs small dots in Figure 1). However, the genotypically diverse cc1690 population did match our predictions in terms of the direction of adaptive trait change in all selection environments: P* decreased under P-limitation, N* decreased under N-limitation, salt tolerance increased in the high salt environment, and I* decreased under low light, though the change in I* was not statistically significant.

When considering all traits together, descendant populations diverged from their ancestors, and variation in these new phenotypes was associated with selection environment (Redundancy Analysis, Figure 1E). We tested for constraints on adaptive change by assessing whether there was significant separation between ancestors and descendants on the RDA axes. RDA axes 1 and 2, which represent linear combinations of selection environment and ancestor ID (PERMANOVA p < 0.01) explain 85% of the variation associated with selection environment, and 36% of the total variation. On RDA axis 1 (PERMANOVA p < 0.001), which is primarily associated with variation in salt tolerance and P*, populations selected in the P, S, B and BS environments were significantly different (separated) from the ancestors. The salt selected populations (S and BS) were also different from the COMBO (C) and low-light selected treatments (L). On RDA axis 2 (PERMANOVA p = 0.005) which is associated with variation in P:N (consumption vector slope), P is different from ancestors and the C, L, B, S, BS populations. The RDA showed that most of the variation in multivariate phenotypes across selection environments was associated with variation in salt tolerance and P*, and much less independent variation was associated with N* and I* (Figure 1E), suggesting that variation in these traits may be subject to physiological or genetic constraints.

Figure 1. Changes in resource competition traits relative to ancestors in seven different selection environments (A: ancestors, C: COMBO, L: light-limited, P: P-limited, N: N-limited, B: biotically depleted media, S: high salt, BS: biotically depleted and high salt).

Small points correspond to individual populations and large points correspond to the
average change (error bars are $\pm$ 1 SE) of all populations in a given environment.
Populations represented with a triangle are the genotypically diverse populations
(cc1690), circles are isoclonal. Error bars on small points in A-E correspond to 95% CI
from non-parametric bootstrapping. E) Redundancy Analysis of N*, I*, P*, salt tolerance,
consumption vectors and cell biovolume across selection environments.

The structure of trait variance: observed trade-offs

Maximum growth rate across populations increased with minimum resource
requirements (R*) for light (OLS slope = 0.021, 95% CI: 0.0027, 0.039, Adjusted R^2 =
0.11), for nitrogen (OLS slope = 0.019, 95% CI: 0.0078, 0.031, Adjusted R^2 = 0.30), and
for phosphorus (OLS slope = 0.12, 95% CI: 0.0094, 0.23, Adjusted R^2 = 0.13),
indicating a trade-off between growth at high and low resource supplies (because a
lower R* indicates faster growth at minimum resource levels) (Figure 2, A-C). Across
populations, competitive abilities for N and P (CN and CP) were positively associated
(ESM Tables 4-6, Figure 2D, ESM Figures S13-15). After accounting for covariance with
competitive abilities for other resources and ancestor ID, competitive abilities for light
were negatively associated with cell biovolume, while and N and P competitive abilities
were not related to cell size (ESM Tables 4-6, ESM Figures S13-15). Principal
components analysis of cell biovolume, competitive abilities for light, nitrogen and
phosphorus showed that 74% of the variation in cell volume and competitive abilities is
explained by the first two PC axes. The first two PC axes demonstrate a positive
association between competitive abilities for N and P, and a possible trade-off
between biovolume and competitive ability for light (Figure 2D).

Figure 2. High maximum population growth rates (μmax) are positively associated with high minimum resource requirements for phosphorus (A), nitrogen (B) and light (C). Estimates of μmax and R* in A-C are from Monod curves generated over independent gradients of phosphorus, nitrogen and light. D) Loadings on the first two PC axes of a PCA using competitive abilities for phosphorus, nitrogen and light (CP = 1/P*, CN = 1/N*, CI = 1/I*) and cell biovolume. 74.3% of the variation in resource use traits and cell size is explained by the first two PC axes.

Correlations in changes across traits

Though theory often assumes that competitive abilities for different resources are

negatively related [15,40], our results did not support this finding either when

considering absolute variation in competitive abilities (ESM Tables 4-6) or variation in

the change in R* relative to the ancestral populations (Figure 3; ESM Tables 1-3, ESM

Figure S10). The changes in R* for different resources never showed evidence of any

trade-offs, and instead were either positively associated (Figure 3A, B) or showed no

significant relationship (Figure 3C, ESM Figure S10).

Figure 3. Partial regression plots showing how changes in descendants relative to

ancestors in two traits are related to each other, while holding all factors in the

statistical model that are not being displayed constant (complete model results in ESM

Tables 1-3). Positive slopes indicate positively associated trait changes. Grey arrows

indicate the directions in which descendant populations' traits got better or worse

relative to ancestors. Panel A describes the four quadrants delineating areas where

both traits get better, both get worse, or one gets better while the other gets worse

(trade-off).

Genetic changes after selection

Genetic differences between the ancestral and descendant populations were identified

by whole genome re-sequencing. The presence or absence of single nucleotide

polymorphisms (SNPs) identified within the populations were compared between the

ancestors and descendants for each selection treatment. The number of variable SNPs

ranged from 396 to 582 (ESM Table 7), roughly corresponding to mutation fixation

rates of 1.25e-8 to 1.83e-8 mutations/[sites × generations]. Contrary to our

expectations, salt stress did not increase the number of fixed mutations. Selection

treatment had no significant effect on the total number of fixed mutations (ESM Figure

S11 A; ANOVA p = 0.788), but the effect of the ancestor was highly significant (ESM

Figure S11 B; ANOVA p < 1e-7).

Evolutionary adaptation altered the predicted outcomes of competition

Divergence in minimum resource requirements and consumption vectors among

populations of the same ancestor selected in different environments (Figure 1E) was

sufficient in some cases to lead to predicted coexistence. One such example is

illustrated in Figure 4, where descendants of Ancestor 3 selected in low light and high

salt environments have diverged sufficiently in their P^* and N^* such that they could

possibly coexist. While neither descendant population could coexist with Ancestor 3,

the two descendant populations could coexist with one another at a supply point

illustrated as a yellow dot. In their ancestral state, out of all pairwise combinations of

our five ancestor populations, resource competition theory [1] predicts unstable

coexistence in three of 10 cases and competitive exclusion in seven of 10 cases. After

selection across the range of environments in our study, resource ratio theory predicts

unstable coexistence in 19% of all possible pairwise interactions (698 total), stable

coexistence in 28% of all possible pairwise interactions and competitive exclusion in

53% of all possible interactions (ESM Figure S12). Among populations selected in the

same environment (59 total), resource ratio theory predicts competitive exclusion in

49%, stable coexistence in 32%, unstable coexistence in 19%.

Figure 4. Descendants of Ancestor 3 evolved in high salt (red) and low light (orange) environments have diverged in their P* and N* such that they can co-exist. Neither descendant could co-exist with the ancestor. Zero net growth isoclines (solid lines) and consumption vectors (dashed lines) for Ancestor 3 (black) and the descendant of Ancestor 3 selected in low light (orange) and high salt (red).

Discussion

Resource competition is among the most important processes structuring ecological communities [41], but competition theory often assumes that traits underlying competitive abilities remain fixed over ecological time scales [1,42]. Here we found that the traits that underlie competitive abilities for essential resources can adapt rapidly in new resource-limited environments. Populations often adapted to resource limitation by reducing their minimum resource requirements. When exposed to high salt, populations evolved higher salt tolerances. Not only could populations respond adaptively to new environments, but they could also adapt within approximately 285 generations. While we observed gleaner-opportunist trade-offs, we did not find evidence for trade-offs in competitive abilities for different resources. Instead, adaptive

changes in competitive ability for one resource were often positively associated with

improvements in competitive ability for another. Since the ancestral and descendant

populations were maintained under identical conditions when estimating their traits,

the changes we observed were heritable.

Trade-offs in resource-use traits have been invoked to explain changes in dominance

across supply ratio gradients and the coexistence of as many species as resources

[1,18]. Trade-offs in competitive abilities for nitrogen and phosphorus [19], iron and

light [23], and light and nitrogen [24] have been documented among and within

species of phytoplankton. These trade-offs may arise due to local adaptation, or due to

biophysical constraints on the acquisition or metabolism of different resources [18,40].

Individuals may invest resources into two main types of cellular machinery: uptake or

assembly machinery. Uptake machinery is composed of nutrient uptake proteins and

chloroplasts, which are both relatively nitrogen-rich, and both of which may scale with

cell size because uptake and photosynthesis must take place at the cell surface.

Assembly machinery, primarily composed of ribosomes, is relative phosphorus-rich and

may also depend on cell size, as smaller cells tend to grow faster ("growth rate

hypothesis" [43]). Consistent with expectations, competitive abilities for light were

negatively associated with cell size, but in contrast to expectations, N and P

competitive abilities were not. Furthermore, we did not find evidence for a trade-off

between competitive abilities for nitrogen and phosphorus.

Trade-offs between competitive abilities for different resources may not arise if the

same resource is required for multiple metabolic processes. For example, positive

correlations in changes in competitive ability, such as we observed, may arise because

the ability to grow in one resource-limited environments results from multiple

components of metabolism becoming more efficient simultaneously [44–46]. For

example, selection under low light could cause individuals to increase the efficiency

with which nitrogen-intensive chloroplasts harvest light, thereby simultaneously leading

to reduced nitrogen requirements.

Patterns of genotypic variation across populations revealed negative correlations
between fitness at low and high resource supply for a given resource (gleaner-
opportunist trade-offs, Figure 2), and positive correlations between competitive
abilities for different resources (Figure 1E, ESM Tables 4 - 6), suggesting that the
evolution of competitive ability could be constrained by genetic correlations between
multiple resource traits under selection. The genetic correlations between different
resource traits could explain the positively associated trait changes (i.e. improvements
in multiple minimum resources requirements simultaneously). In addition, it is possible
that unmeasured traits can be involved in the trade-off, resulting in a positive genetic
covariance between any two resource traits [17]. When testing for trade-offs, we
accounted for concurrent variation in cell size, but other fitness-correlated traits that we
did not measure, such as resistance to grazers or pathogens [47,48], may be involved in
the trade-offs.

Traits relevant to competitive ability, such as cell size, are known to change as a result
of phenotypic plasticity and evolutionary adaptation [49]. We have demonstrated that
adaptation in response to resource limitation and salt stress can alter competitive traits
sufficiently to change the predicted outcome of competition. Adaptation to different
environments caused competitive traits to diverge and enable coexistence. Contrary to
our expectations, we did not find that coexistence would be more likely among two
populations selected in different environments than two populations selected in the
same environment. This may be explained by the fact that even small differences in the
magnitude of adaptive trait change in the same environment can be sufficient to
enable predicted coexistence (i.e. under P-limitation, P^* for one competitor decreases
slightly more than the P^* for the other competitor). The changes in resource ratios and
salt levels represented in our different selection environments are on the same order of
magnitude as gradients of resource ratios and salinity in natural environments [50]. This
means that predictions of the outcomes of competition should incorporate the
potential for evolutionary changes to influence competitive dynamics [15].

Our results are directly relevant to understanding eco-evolutionary feedbacks in

competitively structured communities. Theory predicts that species will converge in

their resource-use traits when competing for essential resources [14,15]. This

expectation, however, depends on two critical assumptions. These assumptions are

that species' consumption vectors remain fixed and that competitive abilities for

different limiting resources trade off. While we did not grow pairs of populations

together in the evolution experiment, the resource limitation treatments mimicked the

effects of a better competitor for the limiting resource, while avoiding exclusion of the

poorer competitor. Our results do not provide empirical support for either of the

assumptions above, suggesting that theoretical predictions of evolutionary adaptation

under essential resource competition may need to be revised [14,15].

Understanding patterns of biodiversity and coexistence requires accounting for past

and current evolutionary changes in species' competitive traits. While macro-

evolutionary patterns show trade-offs in species' resource-use traits, we found that

positively correlated adaptive trait changes drive within-species responses to resource

limitation, altering the expected outcome of competition. Such micro-

evolutionary changes in species' competitive abilities will need to be considered if we

are to improve our predictions of competitive outcomes and community dynamics in a

changing world.

Acknowledgments:

We thank Jennifer Martiny and James Prosser for the opportunity to participate in this special feature. We thank the following people who helped us maintain cultures, and collect and process the data: C. Carvalho, G. Siegrist, P. Ganesanandamoorthy, L. Rihakova, D. Steiner, E. Burmeister, M. Thali and J-C. Walser. Thanks to J. Jokela for helpful comments on an earlier version of the manuscript.

References

1. Tilman D. 1982 *Resource Competition and Community Structure*. Princeton, New Jersey: Princeton University Press.

2. Edwards KF, Litchman E, Klausmeier CA. 2013 Functional traits explain phytoplankton responses to environmental gradients across lakes of the United States. *Ecology* **94**, 1626–1635.

3. Keddy PA. 2015 Competition. *eLS*. (doi:10.1002/9780470015902.a0003162.pub2)

4. Chesson P. 2000 Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.* **31**, 343–366.

5. Miller *et al.* 2005 A Critical Review of Twenty Years' Use of the Resource- Ratio Theory. *Am. Nat.* **165**, 439–448.

6. Hart SP, Turcotte MM, Levine JM. 2019 Effects of rapid evolution on species coexistence. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 2112–2117.

7. Lankau RA. 2011 Rapid Evolutionary Change and the Coexistence of Species. *Annu. Rev. Ecol. Evol. Syst.* **42**, 335–354.

8. Germain RM, Williams JL, Schluter D, Angert AL. 2018 Moving Character Displacement beyond Characters Using Contemporary Coexistence Theory. *Trends in Ecology & Evolution*. **33**, 74–84. (doi:10.1016/j.tree.2017.11.002)

9. Pfennig DW, Pfennig KS. 2010 Character Displacement and the Origins of Diversity. *The American Naturalist*. **176**, S26–S44. (doi:10.1086/657056)

10. Dayan T, Simberloff D. 2005 Ecological and community-wide character displacement: the next generation. *Ecol. Lett.* **8**, 875–894.

11. Schluter D. 2000 *The ecology of adaptive radiation*. OUP Oxford.

12. Grant PR, Grant BR. 2006 Evolution of character displacement in Darwin's finches.

Science **313**, 224–226.

13. MacArthur R, Levins R. 1967 The limiting similarity, convergence, and divergence
of coexisting species. *Am. Nat.* **101**, 377–385.

14. Abrams PA. 1987 Alternative models of character displacement and niche shift. I.
Adaptive shifts in resource use when there is competition for nutritionally
nonsubstitutable resources. *Evolution* **41**, 651–661.

15. Fox JW, Vasseur DA. 2008 Character convergence under competition for
nutritionally essential resources. *Am. Nat.* **172**, 667–680.

16. Lande R, Arnold SJ. 1983 The measurement of selection on correlated characters.
Evolution **37**, 1210–1226.

17. Blows MW, Hoffmann AA. 2005 A reassessment of genetic limits to evolutionary
change. *Ecology*

18. Litchman E, Klausmeier CA, Schofield OM, Falkowski PG. 2007 The role of
functional traits and trade-offs in structuring phytoplankton communities: scaling
from cellular to ecosystem level. *Ecol. Lett.* **10**, 1170–1181.

19. Edwards KF, Klausmeier CA, Litchman E. 2011 Evidence for a three-way trade-off
between nitrogen and phosphorus competitive abilities and cell size in
phytoplankton. *Ecology* **92**, 2085–2095.

20. Litchman E, Klausmeier CA. 2008 Trait-Based Community Ecology of
Phytoplankton. *Annu. Rev. Ecol. Evol. Syst.* **39**, 615–639.

21. Grover JP, Hudziak J, Grover JD. 1997 *Resource Competition*. Springer Science &
Business Media.

22. Kirk KL. 2002 Competition in variable environments: experiments with planktonic
rotifers. *Freshwater Biology.* **47**, 1089–1096. (doi:10.1046/j.1365-
2427.2002.00841.*x*)

23. Strzepek RF, Harrison PJ. 2004 Photosynthetic architecture differs in coastal and
oceanic diatoms. *Nature* **431**, 689–692.

24. Rhee G--U, Gotham IJ. 1981 The effect of environmental factors on phytoplankton
growth: Light and the interactions of light with nitrate limitation 1. *Limnol.*
Oceanogr. **26**, 649–659.

25. Bell G. 2008 *Selection: the mechanism of evolution.* Oxford University Press on
Demand.

26. Hughes AR, Inouye BD, Johnson MT, Underwood N, Vellend M. 2008 Ecological
consequences of genetic diversity. *Ecol. Lett.* **11**, 609–623.

27. Barrett RDH, Schluter D. 2008 Adaptation from standing genetic variation. *Trends*
Ecol. Evol. **23**, 38–44.

28. Bell G, Collins S. 2008 Adaptation, extinction and global change. *Evol. Appl.* **1**, 3–
16.

29. Goho S, Bell G. 2000 Mild environmental stress elicits mutations affecting fitness in
Chlamydomonas. *Proc. R. Soc. Lond. B Biol. Sci.* **267**, 123–129.

30. Velicer GJ, Lenski RE. 1999 Evolutionary trade-offs under conditions of resource
abundance and scarcity: experiments with bacteria. *Ecology* **80**, 1168–1179.

31. Velicer GJ. 1999 Pleiotropic effects of adaptation to a single carbon source for
growth on alternative substrates. *Appl. Environ. Microbiol.* **65**, 264–269.

32. Lahti DC, Johnson NA, Ajie BC, Otto SP, Hendry AP, Blumstein DT, Coss RG,
Donohue K, Foster SA. 2009 Relaxed selection in the wild. *Trends Ecol. Evol.* **24**,
487–496.

33. Kilham SS, Kreeger DA, Lynn SG, Goulden CE, Herrera L. 1998 COMBO: a defined
freshwater culture medium for algae and zooplankton. *Hydrobiologia* **377**, 147–
159.

34. Huisman J, Weissing FJ. 1994 Light-limited growth and competition for light in
well-mixed aquatic environments: an elementary model. *Ecology* **75**, 507–520.

35. Monod J. 1949 The growth of bacterial cultures. *Annu. Rev. Microbiol.* **3**, 371–394.

36. Elzhov TV, Mullen KM, Spiess A-N, Bolker B. 2013 *minpack.lm: R interface to the*
Levenberg-Marquardt nonlinear least-squares algrothim found in MINPACK, pluss
support for bounds. See https://cran.r-
project.org/web/packages/minpack.lm/minpack.lm.pdf.

37. Eilers PHC, Peeters JCH. 1988 A model for the relationship between light intensity
and the rate of photosynthesis in phytoplankton. *Ecol. Modell.* **42**, 199–215.

38. Baty F, Ritz C, Charles S, Brutsche M, Flandrois J-P, Delignette-Muller M-L. 2015 A
Toolbox for Nonlinear Regression in R : The Package nlstools. *J. Stat. Softw.* **66**, 1–
21.

39. Oksanen J *et al.* 2013 Package 'vegan'. *Community ecology package, version* **2**.

40. Klausmeier CA, Litchman E, Daufresne T, Levin SA. 2004 Optimal nitrogen-to-
phosphorus stoichiometry of phytoplankton. *Nature* **429**, 171–174.

41. Gurevitch J, Morrow LL, Wallace A, Walsh JS. 1992 A Meta-Analysis of
Competition in Field Experiments. *Am. Nat.* **140**, 539–572.

42. Adler PB, Hillerislambers J, Levine JM. 2007 A niche for neutrality. *Ecol. Lett.* **10**,
95–104.

43. Sterner RW, Elser JJ. 2002 *Ecological stoichiometry: the biology of elements from*
molecules to the biosphere. Princeton University Press.

44. Toseland A *et al.* 2013 The impact of temperature on marine phytoplankton
resource allocation and metabolism. *Nat. Clim. Chang.* **3**, 979–984.

45. Tamminen M, Betz A, Pereira AL, Thali M, Matthews B, -F. Suter MJ, Narwani A.
2018 Proteome evolution under non-substitutable resource limitation. *Nature*

Communications. **9**. (doi:10.1038/s41467-018-07106-z)

46. Passarge J, Hol S, Escher M, Huisman J. 2006 Competition for nutrients and light:
stable coexistence, alternative stable states, or competitive exclusion? *Ecol.*
Monogr. **76**, 57–72.

47. Yoshida T, Hairston NG Jr, Ellner SP. 2004 Evolutionary trade-off between defence
against grazing and competitive ability in a simple unicellular alga, Chlorella
vulgaris. *Proceedings of the Royal Society B - Biological Sciences* **271**, 1947–1953.

48. Bowers Roger G., Boots Michael, Begon Michael. 1994 Life-history trade-offs and
the evolution of pathogen resistance: competition between host strains.
Proceedings of the Royal Society of London. Series B: Biological Sciences **257**,
247–253.

49. Malerba ME, Marshall DJ. 2019 Size-abundance rules? Evolution changes scaling
relationships between size, metabolism and demography. *Ecology Letters.* **22**,
1407–1416. (doi:10.1111/ele.13326)

50. Wetzel RG. 2001 *Limnology: Lake and River Ecosystems.* Gulf Professional
Publishing.