Gene-environment interactions characterized by single embryo

transcriptomics

Alfire Sidik[1], Groves B. Dixon[1], Hannah G. Kirby[1], and Johann K. Eberhart[1*]

[1]Department of Molecular Biosciences, Institute for Cellular and Molecular Biology,

Institute for Neuroscience, Waggoner Center for Alcohol and Addiction Research,

University of Texas, Austin, TX 78712, USA

*Correspondence to: Johann K. Eberhart, PAT 522, The University of Texas at Austin,

Molecular Biosciences, College of Natural Sciences, 2401 Speedway, Austin, TX, 78712.

Phone: 512-232-8340.

ABSTRACT

Gene-environment interactions are likely to underlie most human birth defects. The

most common environmental contributor to birth defects is likely prenatal alcohol

exposure. Fetal Alcohol Spectrum Disorders (FASD) describes the full range of defects

that result from prenatal alcohol exposure. Gene-ethanol interactions underlie

susceptibility to FASD but we lack a mechanistic understanding of these interactions. Here,

we leverage the genetic tractability of zebrafish to address this problem. We first show that

vangl2, a member of the Wnt/planar cell polarity (Wnt/PCP) pathway that mediates

convergent extension movements, strongly interacts with ethanol during late blastula and

early gastrula stages. Embryos mutant or heterozygous for *vangl2* are sensitized to ethanol-

induced midfacial hypoplasia. We performed single-embryo RNA-Seq during early

embryonic stages, to assess individual variation to the transcriptional response to ethanol

and determine the mechanism of the *vangl2*-ethanol interaction. To identify the pathway(s)

that are disrupted by ethanol we used these global changes in gene expression to identify

small molecules that mimic the effects of ethanol via the Library of Integrated Network-

based Cellular Signatures (LINCS L1000) dataset. Surprisingly, this dataset predicted that

the Sonic Hedgehog (Shh) pathway inhibitor, cyclopamine, would mimic the effects of

ethanol, despite the fact that ethanol did not alter the expression levels of direct targets of

Shh signaling. Indeed, we found that ethanol and cyclopamine strongly interact to disrupt

midfacial development. Collectively, these results suggest that the midfacial defects in

ethanol-exposed *vangl2* mutants are due to an indirect interaction between ethanol and the

Shh pathway. Vangl2 functions as part of a signaling pathway that regulates coordinated

cell movements during midfacial development. Consistent with an indirect model, a critical

source of Shh signaling that separates the developing eye field into bilateral eyes, allowing

the expansion of the midface, becomes mispositioned in ethanol-exposed *vangl2* mutants.

We demonstrate that ethanol also interacts with another Wnt/PCP pathway member, *gpc4*,

and a chemical inhibitor, blebbistatin. By characterizing membrane protrusions, we

demonstrate that ethanol synergistically interacts with the loss of *vangl2* to disrupt cell

polarity required for convergent extension movements. Collectively, our results shed light

on the mechanism by which the most common teratogen can disrupt development.

INTRODUCTION

Birth defects manifest as structural or functional malformations at birth and are the

leading cause of infant mortality in the United States [1]. Although the multifactorial

etiologies of birth defects are not well understood, many are thought to derive from a

complex interplay between genetic and environmental factors. The adverse effects of

teratogens, or environmental agents that cause irreversible developmental defects, were

first recognized in the 1950s and 1960s [2]. It was then that investigators began to observe

developmental anomalies in infants exposed to methylmercury and thalidomide in utero

[3, 4]. Since then, other environmental teratogens have come to light, including but not

limited to, pollutants, pharmaceuticals, and chemicals.

Prenatal alcohol exposure (PAE) is the most common cause of birth defects, with

the prevalence of fetal alcohol spectrum disorders (FASD) in the US is as high as 2-5%

[5]. Fetal alcohol syndrome (FAS) is the most severe outcome following PAE and is

characterized by midfacial hypoplasia, as well as growth and neural deficits [6]. While

PAE is required for the development of FASD, the teratogenic effects of ethanol are

modulated by genetics [7, 8]. For instance, monozygotic twins were 100% concordant for

FAS, whereas dizygotic twins were only 63% concordant [8, 9]. Furthermore, animal

models of FAS show strain-specific differences after controlling for environmental

variables such as dose and timing [10, 11]. Despite this, the genetic factors that protect or

predispose an individual to FASD are poorly understood. Moreover, we still lack a basic

understanding of the mechanism of ethanol teratogenesis. Zebrafish are well suited to

address this problem due to their genetic tractability and ease of embryological

manipulation [12].

In a screen to identify genetic modifiers of ethanol teratogenicity, *vang-like 2*

(*vangl2*), a member of the non-canonical Wnt/PCP pathway, emerged as an ethanol

sensitive locus [13]. In zebrafish, mutations in the *vangl2* locus disrupt convergent

extension movements during gastrulation, resulting in a shortened, broadened, body axis

[14]. These mutants infrequently present with cyclopia, fusion of the bilateral eyes, but the

phenotypic expressivity of this trait varies with factors such as temperature and genetic

background [15]. In the screen, all untreated *vangl2* mutants displayed proper separation

of the eyes and craniofacial skeletal elements were intact [13]. Upon exposure to a

subteratogenic dose of ethanol, *vangl2* mutants were fully penetrant for cyclopia and

displayed profound defects to the midfacial skeleton [13]. Ethanol-treated *vangl2*

heterozygotes were largely indistinguishable from their wild-type siblings, with the

exception of a single synophthalmic ethanol-treated heterozygote, providing evidence for

latent haploinsufficiency. Together, these data suggest a synergistic interaction between a

mutation in *vangl2* and ethanol.

We know *vangl2* plays a critical role in mediating convergent extension movements

as evidenced by their body axis defect, however because the early effects of ethanol

exposure remain poorly defined, the precise mechanism of the *vangl2*-ethanol interaction

remain elusive. To better understand how ethanol interacts with loss of *vangl2* to alter

phenotypic outcomes, we took an unbiased approach to assess the transcriptional response

to ethanol. we performed single embryo RNA-sequencing (RNA-seq) on control

(untreated) and ethanol-treated wild-type embryos in a time-course spanning gastrulation

and early segmentation stages in zebrafish.

Bioinformatic and functional analyses indicated that midfacial defects in ethanol-

exposed *vangl2* mutants was due to an indirect interaction between ethanol and the Shh

pathway. While there was no alteration in the level of expression of direct Shh targets, a

critical source of Shh signaling that separates the developing eye field into bilateral eyes

becomes mispositioned in ethanol-exposed *vangl2* mutants. We demonstrate that ethanol

and loss of *vangl2*, synergistically interact to disrupt polarized cellular protrusions required

for proper convergent extension movements.

RESULTS

Early embryogenesis is the sensitive time window for *vangl2* mutants

To determine the critical time window of ethanol sensitivity for the *vangl2* mutants,

we first initiated ethanol treatment at various stages comprising late blastula to early

gastrula for 24 hours. The inner lens-to-lens width was used as a measure of synophthalmia

and we calculated the occurrence of cyclopia to characterize ethanol-induced teratogenesis.

In control conditions, the spacing of the eyes in *vangl2* mutants mirrored that of wild-type

zebrafish. Ethanol-exposed *vangl2* mutants exhibited midline defects ranging in severity

from synophthalmia to cyclopia across all time points examined, but these malformations

were fully penetrant for cyclopia (100% fused; n=5/5) when ethanol was applied at shield

stage (6 hours post fertilization, hpf) at the onset of gastrulation (Fig 1A-B). Interestingly,

heterozygotes only displayed cyclopia when ethanol was applied at high stage (3.3 hpf)

(22% fused; n=4/18), a time when treating wild-type embryos with higher concentrations

of ethanol causes similar defects (Fig 1B) [16]. Thus, heterozygotes and homozygotes are

divergent in their time window of greatest ethanol sensitivity. This may be due to a

compensatory genetic mechanism in *vangl2* heterozygotes because zygotic gene
expression initiates after high stage (4 hpf) in zebrafish.

123 The effect of ethanol on the early zebrafish transcriptome is subtle relative to
124 developmental time

To determine if ethanol caused transcriptional changes that could underlie the
interaction with *vangl2*, we designed two RNA-Seq experiments that largely overlapped in
design (Fig 2A-B). Embryos were exposed to a subteratogenic dose of 1% ethanol in
embryo media (171 mM), which equilibrates to approximately 50 mM tissue concentration
[17]. For the first experiment, embryos were treated at the onset of gastrulation (6 hpf) and
collected at mid-gastrulation (8 hpf) and the end of gastrulation (10 hpf) (Fig 2A). A second
experiment was performed to increase power (Fig 2B). Embryos were similarly exposed,
but a 14 hpf time window was included, when the eye fields have completely separated.
Each sample consisted of an individual zebrafish embryo with five replicates per timepoint
and treatment. The 6 hpf control samples were omitted since they lacked ethanol-treated
samples for comparison. The data from both experiments were combined for subsequent
analyses, controlling for batch effects.

To assess the effect of time, batch (i.e. experiment 1 and 2), and ethanol, on the
early zebrafish transcriptome, we performed principal component analysis (PCA).
Individuals from each timepoint clustered tightly along *PC1*, which accounted for 38% of
the transcriptional variation observed across the datasets. This strong effect of time on the
zebrafish transcriptome is in agreement with previous studies [18]. Clustering of samples
by time, regardless of ethanol treatment suggested that control and ethanol-treated samples
were accurately staged, indicating that ethanol did not delay developmentally regulated
transcriptome patterns. There was greater discrimination of 14 hpf samples relative to

earlier timepoints (6, 8, and 10 hpf), which is consistent with greater distinction of this timepoint in terms of developmental time and morphology (Fig 2C). *PC2* largely captured batch effects between experiment 1 and 2 (Fig 2D).

The majority of variation between samples did not appear to be due to treatment, with control and ethanol-treated samples randomly interspersed along *PC1* and *PC2* (Fig 2E). Separation by treatment was observed along *PC8* and *PC9*, which accounts for 3% of the variation in the data (Fig 2F). Hierarchical clustering of samples based on correlation, further corroborated this finding. The 6 and 14 hpf samples showed the greatest dissimilarity, whereas the 8 and 10 hpf samples showed the greatest similarity, irrespective of treatment (S1 Fig). In summary, the transcriptional effect of exposure to a subteratogenic dose of ethanol on the early zebrafish transcriptome was subtle, while time provided the strongest transcriptional fingerprint.

Ethanol has effects on transcription that are largely distinct between different developmental timepoints

Although developmental age was a stronger source of transcriptional variation, we still detected substantial variation in gene expression following subteratogenic ethanol exposure. There were 1,414 differentially expressed genes (DEGs), with a false-discovery rate (FDR) less than 0.1 (Fig 3A; Benjamini–Hochberg procedure). There were more upregulated than downregulated DEGs among ethanol-treated individuals across timepoints (S2 Fig) and these DEGs were distinct between developmental timepoints (Fig 3A,B). In summary, while some genes are generally affected by ethanol across several critical periods, ethanol largely elicits unique responses at distinct developmental stages spanning early embryogenesis.

Ethanol does not affect the Wnt/PCP pathway at the transcriptional level

While *vangl2* mutants do not typically exhibit cyclopia, cyclopia is observed in double mutants between *vangl2* and other Wnt/PCP pathway members [15, 19]. One potential mechanism for the interaction between ethanol and *vangl2* would be the transcriptional misregulation of other Wnt/PCP pathway members. However, transcription of Wnt/PCP pathway members was largely unaffected by ethanol exposure. No Wnt/PCP pathway members were among the 14 shared DEGs across all timepoints (Fig 3C). KEGG pathway enrichment analysis further confirmed that ethanol had little effect on the Wnt/PCP pathway at the level of transcription (Fig 3D). Only two Wnt/PCP pathway members were moderately affected: ethanol exposure moderately decreased expression of the cofactor, *glypican 4* (*gpc4*), (log_2 fold= -0.237; p-value = 0.036) across all timepoints [15, 20] and upregulated expression of *rac3a* (log_2 fold = 0.737; p-value = 8.89E-06), a member of the Rho family of small GTPases [21]. We plotted the normalized read counts across each time point from the RNA-seq experiment to further investigate the dynamics of the potential alteration in *gpc4* levels. The expression of *gpc4* was modestly reduced in ethanol-treated embryos across the RNA-seq dataset (Fig 3E). Ethanol exposure consistently downregulated *gpc4* at 8 and 10 hpf, but the magnitude of the downregulation was relatively modest. To statistically compare ethanol-treated and control embryos, expression of *gpc4* at 10 hpf was investigated using qRT-PCR. This result demonstrated that *gpc4* was not significantly affected by ethanol exposure (Fig 3F). Thus, direct transcriptional alteration to the Wnt/PCP pathway is unlikely to explain the ethanol-induced phenotypes in *vangl2* mutants and heterozygotes.

Modules of co-regulated genes related to ethanol exposure

To determine if ethanol disrupted networks of genes that could explain the *vangl2*-

ethanol interaction, we next performed Weighted Gene Co-expression Network Analysis

(WGCNA) [22]. This unsupervised network analysis identifies groups of genes, termed

modules, based on correlated expression patterns across the samples. Modules are

summarized by the first principal component for the expression estimates of the included

genes, termed the module eigengene, which can be correlated with sample traits to identify

biological significance. The cluster dendrogram generated in this analysis illustrates the

presence of highly distinct and clustered modules (S3A Fig). Merging of similar modules

produced eleven total modules (S3B Fig). Consistent with our PCA analysis, the module

eigengenes were primarily correlated with time. We used these modules to further validate

our ethanol exposure regimen.

Previous work has shown that ethanol delays development in a dose-dependent

manner at concentrations equal to or greater than 1.5% ethanol [23]. For the experiments

herein, ethanol-treated samples were morphologically stage-matched to control samples to

exclude differences due to developmental age or delay. To confirm that the ethanol samples

were indeed age-matched to the control samples at the level of transcription, we compared

expression patterns of developmentally regulated genes between the ethanol and control

samples from each timepoint. For developmentally regulated genes, we used the gene with

the highest module membership (i.e. the hub gene) from each of the WGCNA modules that

was associated with time ($p < 0.05$) (S4 Fig). Consistent with the results from the PCA (Fig

2C), the slope of the expression levels for each gene were similar in control and ethanol-

treated samples across age, most clearly demonstrated in the magenta4, thistle1, and

blueviolet modules (S4B,C,G Fig). Interestingly, we find a time-specific difference in the

expression of hub genes in the honeydew1, greenyellow, and saddlebrown modules at 10

hpf (S4F,H,K Fig). Together, these data indicate samples were accurately age-matched and the observed changes were biologically relevant.

Two modules (mediumpurple4 and darkolivegreen4) correlated with ethanol treatment (S3B-D Fig). The purple and green modules positively and negatively correlated with ethanol treatment, respectively (S3C Fig). A significant differentially expressed gene from the green (S3G Fig) and purple module (S3H Fig) were selected for independent validation on independent biological replicate samples derived from the same wild-type zebrafish line using quantitative real-time RT-PCR (qRT-PCR). These results indicate that our RNA-seq faithfully represents transcript levels. The green module is only weakly downregulated and is enriched for genes encoding zinc finger (ZnF) proteins, which included *znf1015* (S3E Fig). This module is almost entirely composed of genes on chromosome 4, that have been shown to be co-regulated during these stages of development [24, 25]. The purple module revealed GO enrichment of transmembrane transporters, which included *slc16a9a* (S3F Fig). As an upregulated module enriched in transporters, these genes are likely to represent the physiological response to ethanol. Thus, while we did identify two modules associated with ethanol (S1 Table), the module membership did not provide clues to the interaction between ethanol and *vangl2*.

Cyclopamine is predicted to mimic the effects of ethanol

One challenge in RNA-seq analyses is inferring the mechanism underlying a diseased or environmentally-perturbed state from a large set of differentially expressed genes. Individual functional analyses of significant gene-ethanol interactions are time consuming and thus inefficient. To circumvent this problem, we adopted a bioinformatics approach, utilizing the Library of Integrated Network-Based Cellular Signatures (LINCS L1000) toolkit [26]. This transcriptomic dataset includes gene expression data from nine

human cancer cell lines exposed to thousands of small molecule drugs [26]. We queried

the top 100 and 150 up- and down-regulated genes induced by ethanol exposure against

the LINCS L1000 dataset using the clue.io platform (https://clue.io). The query generated

a list of small molecules predicted to have a positive or negative correlation to the input

signature (*i.e.* small molecules predicted to either mimic or antagonize the transcriptional

effects of ethanol). We were particularly interested in those chemicals with positive

correlation to ethanol as they would give insight into the mechanism of ethanol

teratogenicity and highlight potential co-factors that exacerbate ethanol teratogenicity.

Interestingly, cyclopamine, a hallmark Shh pathway inhibitor that inhibits the core

Shh pathway protein Smoothened (Smo), was predicted to positively correlate with the

ethanol signature. The Shh pathway is critical for midfacial development and mice

deficient in *Sonic Hedgehog* (*Shh*) exhibit severe brain and face malformations, including

holoprosencephaly and a single medial eye (cyclopia) [27]. In zebrafish, reduction of *shh*

or null mutations in *smo* similarly results in severe loss of craniofacial midline structures

(i.e. the anterior neurocranium) [28]. Animal models have demonstrated ethanol is an

environmental risk factor for holoprosencephaly, resulting in a characteristic set of

midfacial defects, a hypomorphic forebrain, and in severe cases, cyclopia [16, 29-32].

Thus, attenuation of the Shh pathway could mechanistically explain the *vangl2*-ethanol

interaction.

Ethanol indirectly attenuates Shh signaling

To investigate the interaction of cyclopamine and ethanol, we first exposed wild-

type zebrafish embryos to cyclopamine (50 μM) at shield stage (6 hpf) for 24 hours,

mimicking the ethanol exposure window for the *vangl2* mutants. Embryos were fixed at 4

dpf and the cartilage and bone were stained with Alcian blue and Alizarin red, respectively.

We observed a range of midfacial defects with the most severe phenotype being a complete

loss of the anterior neurocranium and reduced spacing between the eyes (Fig 4A). Since

cyclopamine was predicted to mimic the effects of ethanol, we next combined this low

dose of cyclopamine with the subteratogenic dose of ethanol. Strikingly, all embryos

presented with synophthalmia or cyclopia and significant reductions and defects of the

cartilages of the neuro- and viscerocranium. To quantify this combinatorial effect, we

measured the distance between the lenses and observed a significant reduction in co-

exposed embryos relative to those exposed to either ethanol or cyclopamine alone

(p<0.0001), suggesting a strong synergistic interaction (Fig 4B).

In chick, ethanol exposure during somitogenesis has been proposed to suppress Shh

signaling and induce apoptosis in cranial neural crest cells that make up the craniofacial

skeleton [33]. However, work in zebrafish shows only a modest increase in cell death

within the eye field at a much higher dose of 2% ethanol [34]. To ensure that cells in the

eye field are not simply undergoing apoptosis, we performed a TUNEL cell death assay in

ethanol-treated *vangl2* mutants at 11 hpf, prior to optic vesicle evagination. As expected,

we failed to detect an increase in apoptotic cells within the eye field in *vangl2* mutants or

their siblings (S5 Fig). Thus, Shh-mediated pro-survival signals do not appear to be

significantly disrupted by ethanol.

To test for a direct effect of ethanol on Shh signaling, we quantified the relative

expression of *ptch2*, a canonical read-out of Shh pathway activity, at bud stage (10 hpf).

Subteratogenic doses of ethanol had no effect on the expression levels of *ptch2*, consistent

with RNA-seq results (Fig 4D). While cyclopamine significantly reduced Shh signaling (p

= 0.0006), ethanol did not further reduce *ptch2* levels significantly (p = 0.1115). KEGG

analysis from the RNA-seq confirmed that ethanol does not affect the Shh pathway as a

whole, nor does it reduce the levels of Shh target genes (i.e. *gli*, *ptch*) (Fig 4E). Thus, at

normally subteratogenic doses, ethanol does not appear to directly attenuate Shh signaling.

Ethanol disrupts convergent extension

The ethanol-induced *vangl2* mutant phenotype closely mirrors those in compound

mutants between *vangl2* and other Wnt/PCP pathway members [15, 19]. These double

mutants display a further reduction in convergent extension, as evidenced by a shorter and

broader body axis [15]. Based on these data, we hypothesized that ethanol disrupts

convergent extension, which would mislocalize the Shh signal and result in eye defects.

We performed *in situ* hybridization on untreated and ethanol-treated *vangl2*

embryos to examine convergent extension and to test the hypothesis that the Shh expression

domain is mislocalized in ethanol-exposed *vangl2* mutants. We analyzed expression of *shh*

in the axial mesoderm and *paired box 2a* (*pax2a*) in the midbrain-hindbrain boundary at

bud stage (10 hpf) [35]. Ethanol was initiated at high stage (3.3 hpf), when *vangl2*

heterozygotes and homozygotes are equally sensitive to the effects of ethanol. We observed

a gene and ethanol-dose dependent reduction in the length of the *shh* expression domain

(extension) and an increase in the width of the *pax2a* expression domain (convergence)

(Fig 5A).

To quantify the effect of ethanol on convergent extension, we plotted the

normalized expression values of *shh*/*pax2* (Fig 5B). Post-hoc analyses (Tukey's) revealed

significant differences between untreated embryos across *vangl2* genotypes, indicating that

vangl2 gene dosage affects convergent extension. Similarly, we found significant

differences between ethanol-treated mutants and their heterozygous and wild-type siblings

($p = 0.017$). This observation confirms that loss of *vangl2* results in reduced convergent

extension movements, as evidenced by their short body axis. This phenotype was

exacerbated with ethanol treatment, resulting in reduced convergent extension for *vangl2*

heterozygotes and homozygotes, compared to their untreated counterparts. While there was

a trend, we did not observe a difference in convergent extension between ethanol-treated

vangl2 heterozygotes and their wild-type siblings in their *shh/pax2* ratio. However,

synophthalmia occurred in ethanol-treated heterozygotes but not wild-type embryos,

demonstrating that the nonsignificant reduction in convergent extension in the

heterozygotes can have a phenotypic consequence.

Ethanol alters *six3* and *rx3* expression in the eye field

Transcription factors involved in the specification of the eye field have also been

implicated in the mechanism of eye field separation. The expression of three of these

transcription factors, *six3*, *rx3*, and *rx1*, are altered by ethanol exposure [34]. We examined

the effect of ethanol on *six3a* and *rx3* in ethanol-treated *vangl2* mutants at the initiation of

optic vesicle evagination. *In situ* hybridization of *six3a* in 11 hpf embryos shows a heart-

shaped expression pattern in the prospective forebrain. The caudal indentation marks the

splitting of the eye field into bilateral domains (Fig 5C). This expression pattern becomes

more diffuse with loss of *vangl2*. In untreated homozygous mutants, we observed a

shortening along the anterior-posterior (AP) axis and a broadening along the mediolateral

axis at 11 hpf, consistent with the convergent extension defect. This expression pattern was

further exacerbated in ethanol-treated mutants with complete loss of the caudal indentation.

We observed a similar effect of genotype and ethanol on *rx3* expression at mid-evagination

(12 hpf) (S6 Fig). At this stage, *rx3* is localized to the prospective forebrain and retina [36].

Ethanol-treated *vangl2* homozygous mutants exhibit a compressed expression domain,

clearly displaying a reduction in convergent extension. Our expression analyses suggest

the eye field is specified but mis-localized and fails to separate into bilateral domains, likely

due to defects in mesodermal migration.

Mutation in *gpc4* enhances cyclopia in a dose-dependent manner

Our data support a model in which ethanol interacts with *vangl2* via a combinatorial

disruption of convergent extension. If ethanol disrupts convergent extension, which leads

to interactions with *vangl2*, then further genetic disruption to convergent extension should

exacerbate the ethanol-*vangl2* phenotype. Embryos deficient in *gpc4* similarly have a

defect in convergent extension as evidenced by their shortened, broadened body axis [15].

Previous work in zebrafish has shown a functional interaction between *vangl2* and *gpc4*,

where *vangl2;gpc4* double mutants were invariably cyclopic [15]. We conducted

additional functional analyses to further examine the relationship between these two genes

in the context of ethanol exposure. Consistent with Marlow *et al.*, double mutants were

fully penetrant for cyclopia with or without ethanol. While 1% ethanol exposure altered

the facial morphology of *gpc4* mutants (Fig 6A), it did not cause cyclopia. However,

ethanol concentrations greater than 1%, which did not cause cyclopia in wild-type siblings,

resulted in cyclopia in *gpc4* mutants (S7 Fig) [15]. Additionally, embryos carrying 3

mutant alleles (either *vangl2$^{-/-}$;gpc4$^{+/-}$* or *vangl2$^{+/-}$;gpc4$^{-/-}$*) were greatly sensitized to

cyclopia when exposed to 1% ethanol (Fig 6B). Collectively these data indicate that in

sensitized genotypes, ethanol disrupts convergent extension, thus mispositioning a source

of Shh that is essential for separation of the eye field into bilateral primordia.

Blebbistatin phenocopies the ethanol-induced defect in *vangl2* mutants.

If ethanol interacts with the Wnt/PCP pathway via convergent extension, then

replacing ethanol with an inhibitor that disrupts cell movements, should similarly result in

synophthalmia and/or cyclopia. We extended our analyses to test the effect of blebbistatin,

a myosin II inhibitor that reduces axis elongation, in *vangl2* mutants [37]. Blebbistatin

treatment during gastrulation (6-10 hpf) induced synophthalmia in *vangl2* heterozygotes

and homozygotes (Fig 7A). We then quantified this interaction by measuring the inner lens-

to-lens width and found a significant reduction in blebbistatin-treated *vangl2* homozygotes

(Fig 7B). Furthermore, 26.92% and 100% of blebbistatin-treated heterozygotes and

homozygotes, respectively, were cyclopic (Fig 7C). These data suggest ethanol further

inhibits convergent extension in the sensitized background, *vangl2*, which not only disrupts

their ability to elongate their body axis, but also inhibits the splitting of the eye field.

Ethanol affects *vangl2* filopodia organization on migrating cells

Based on our data, cell dynamics underlying convergent extension are likely

involved in the *vangl2*-ethanol interaction. During gastrulation, mesodermal cells adopt a

bipolar shape and orient their actin-based cytoskeletal projections on their leading and

trailing edge, with respect to the notochord, allowing for proper convergent extension [38].

In *vangl2* mutant embryos, these gastrula cells fail to elongate and align with the primary

axis [39]. Furthermore, *vangl2* mutant ectodermal cells possess a greater number of

filopodial protrusions than wild-type siblings and display projections in an unpolarized

manner [40].

To determine whether ethanol may be exacerbating this phenotype, we analyzed

filopodia protrusion length and orientation around the circumference of migrating live

mesodermal cells. We injected memGFP mRNA into a single blastomere of *vangl2*

embryos at the 8-32 cell stage, to obtain mosaic expression. We then treated a subset of the

injected embryos with 1% ethanol from 6-10 hpf. Homozygous mutants were identified

based on their reduced body axis elongation at 10 hpf. Cells were imaged at 40-60º from

the primary axis. Spike-like filopodia were highly unpolarized in ethanol-treated *vangl2*

mutants compared to their wild-type and heterozygous siblings (Figure 8A). We quantified

the polarity and number of filopodia for each cell, with respect to the head (90º), the

primary axis/notochord (180º), and the tail (270º) (Figure 8B). Rose diagrams suggest

ethanol-treated *vangl2* mutants have more un-polarized projections on the

anterior/posterior edge (red) as opposed to the leading (green) or trailing (blue) edge. We

next reduced the number of bins in the rose diagrams to 90º and noticed a clear over-

representation of filopodia in *vangl2* mutants on their anterior/posterior edge (Figure 8C).

Tukey's post-hoc analyses confirmed ethanol-treated *vangl2* mutants had significantly

more filopodia on their anterior/posterior edge compared to other axes and their wild-type

or heterozygous siblings (Figure 8D). These data confirm that ethanol exposure is

associated with an increase in the number of filopodial projections oriented in the wrong

direction, or perpendicular to the path of dorsal migration, in ethanol-treated *vangl2*

mutants.

DISCUSSION

Molecular staging for the early zebrafish embryo

Despite their importance, the nature and mechanisms of gene-environment

interactions are largely unknown. Here, we performed single embryo bioinformatic

analyses to shed light upon the mechanism behind gene-environment interactions. A

growing number of researchers are using the zebrafish for toxicological and/or gene-

environment studies. One potential confounding variable in these types of studies is

developmental delay. Our WGCNA analyses are rich in gene modules that similarly vary

over developmental time, providing a way to molecularly stage embryos. Thus, our results

will provide important resources to ensure the rigor and reproducibility of these analyses.

The effects of ethanol on the transcriptome is subtle.

Our hierarchical clustering analysis demonstrates that the effect of ethanol on the

transcriptome is minor, within any given time point ethanol-exposed and unexposed

embryos intermingled. PCA results support this, demonstrating that only 3% of the

variance in the transcriptomic data is attributable to ethanol exposure. In sum, we found

1,414 differentially regulated genes across all time points with the majority of these genes

being upregulated. Most differentially expressed genes were specific to a single time point.

At any given time point there were between 92 and 328 differentially expressed genes.

Thus, exposure to a normally subteratogenic dose of ethanol results in subtle and dynamic

alterations to the transcriptome.

Our findings identified dramatically fewer differentially expressed genes than those

observed in transcriptomic analyses of ethanol-exposed chicken and mouse embryos. In

chicken 3,422 differentially regulated genes were observed after 6 hours of ethanol

treatment and the preponderance (1,924 genes) were downregulated [41]. In a mouse

microarray, 2,906 differentially expressed genes were identified 3 hours after ethanol

injection [42]. Species-specific developmental trajectories or the tissue analyzed, cranial

neural fold (chicken), headfold (mouse), versus whole embryo (our study) may account for

some of these differences. However, it is important to note that the chicken and mouse

strains analyzed are highly sensitive to the dose of ethanol used while our zebrafish AB

strain is not (without the addition of a sensitizing mutation). Thus, the chicken and mouse

data likely represent transcriptomic changes representative of pathogenesis. Our zebrafish

data reflect a direct effect of ethanol, which may or may not be pathogenic depending upon

the genetic and environmental milieu.

WGCNA analysis identified two modules (mediumpurple4 and darkolivegreen4)

of genes that are coordinately altered by ethanol exposure. The vast majority of genes that

negatively correlate with ethanol exposure (membership in darkolivegreen4) encode zinc

finger proteins (ZnF) located on chromosome 4. Ethanol has similarly been shown to

downregulate *Zinc finger protein, subfamily 1A*, 4 in mouse fetuses exposed to ethanol

during early development [43]. However, the functional significance of these changes in

ZnF proteins remains unknown and they often have diverse binding affinities and

functions. The long arm of zebrafish chromosome 4 (*Chr4q*) is typically heterochromatic

(condensed) and lacking in protein-coding genes [44]. However, it is not late replicating

until the end of gastrulation or bud stage (10 hpf) [25]. Previous work in zebrafish has

found ZnF proteins on chromosome 4 to undergo robust expression from the initiation of

zygotic transcription until mid-gastrula stage [24]. Since many chromosome 4 genes are

downregulated across ethanol-treated individuals during gastrulation, ethanol may

interfere with replication timing, blocking the early-to-late replication switch, or chromatin

remodeling. Based on these findings, the differentially expressed genes may follow

disrupted expression of ZnF proteins due to chromatin remodeling.

Ethanol interacts with *vangl2* through the indirect effects on the Shh pathway

Because Vangl2 is a core member of the Wnt/PCP pathway, a simple explanation

for ethanol-sensitivity in *vangl2* mutants would be that ethanol transcriptionally

dysregulates the Wnt/PCP pathway. However, our RNA-seq results indicated that this is

not the case. The expression of *gpc4* was the only member of the Wnt/PCP pathway that

was downregulated in our RNA-seq dataset. Cyclopia is observed in *vangl2;gpc4* double

mutants [15] and we show that heterozygosity for *gpc4* further sensitized *vangl2* mutants

to ethanol. However, the downregulation of *gpc4* was quite modest and unlikely to have

any substantial impact on the phenotype of ethanol-exposed *vangl2* mutant.

We used an unbiased approach, the LINCS L1000 dataset to identify pathways that

might be disrupted by ethanol and generate cyclopia in *vangl2* mutants. One of the top

matches to the effects of ethanol was cyclopamine, a Hh pathway inhibitor. Shh signaling

is critically important for development of the midface across species [27, 45-47]. In

zebrafish *shha* and *smo* mutants have cyclopia because Shh signaling to the diencephalon

is critical for separation of the eye fields [48, 49]. Furthermore, cyclopia observed in

Wnt/PCP pathway mutants is due to the mispositioning of this source of Shh [16, 19]. Thus,

disruption of the Shh pathway could mechanistically explain the ethanol sensitivity of

vangl2 mutants.

The Shh pathway has been implicated in ethanol teratogenesis. However, there has

been much controversy regarding whether this interaction is direct or not [8]. As predicted

by LINCS, we find a synergistic interaction between ethanol and the Hh pathway inhibitor

cyclopamine. However, both our RNA-seq and qPCR data demonstrate that, at this timing

and dose, ethanol does not alter the expression of direct readouts of Shh activity. Such an

alteration would be expected if ethanol interacted directly with core components of the

pathway. As supported by our *in situ* analysis, we conclude that ethanol indirectly effects

the Shh pathway by mislocalizing the source of the ligand.

Ethanol disrupts convergent extension by disrupting polarized cellular protrusions.

The phenotypes observed in ethanol-exposed *vangl2* mutants suggest that ethanol

further disrupts convergent extension. Our findings are consistent with previous

observations that high doses of ethanol can generate phenotypes consistent with a

disruption of convergent extension in zebrafish and Xenopus [16, 50, 51]. More recently,

zebrafish microarray analyses demonstrated that disrupted gastrulation movements due to

ethanol exposure were caused by a reduction in *pchd18a* expression [52]. The expression

of *pchd18a* is not altered in our dataset. However, Sarmah and colleagues used the ethanol-

sensitive TL strain and exposed embryos to ethanol from 2 to 8 hpf. This later difference

is particularly intriguing giving our finding that the time window of greatest sensitivity to

ethanol was different for *vangl2* heterozygotes and mutants, with heterozygotes being most sensitive at 3.3 hpf (the earliest time we tested). This raises the intriguing possibility that the earlier time window of teratogenesis is due to a different mechanism.

We propose that ethanol leads to midfacial defects and cyclopia, not by dysregulating Wnt/PCP at the level of transcription, but indirectly through inhibition of convergent extension and mislocalization of important developmental regulators like *shha*. At the start of convergent extension, lateral gastrula cells migrate and converge to the dorsal region of the developing embryo, where they intercalate between neighboring cells to drive extension of the body axis [20, 53, 54]. As these cells begin their dorsal migration, they elongate along their mediolateral axis and polarize their actin-based cytoskeletal processes medially and laterally to drive intercalation [54, 55]. Convergent extension movements during gastrulation also alter the shape of the eye field, whereby diencephalic precursor cells articulate both medially and caudally [56]. We know from previous work that *vangl2* mutant lateral gastrula cells fail to elongate their mediolateral axis and have a slight anterior bias in their dorsal trajectory [39, 57]. In addition to their shape changes, *vangl2* mutant ectodermal cells have more filopodia than wildtype cells and defective polarization (i.e. less filopodia localized on the trailing edge), resulting in an indirect dorsal trajectory [40]. Using live confocal imaging coupled with memGFP expression, we examined the effect of ethanol on filopodia number and polarity in *vangl2* embryos. Interestingly, we did not find more filopodia in *vangl2* lateral mesodermal cells relative to wildtype cells. Despite this, we find that ethanol-treated *vangl2* mutant embryos have more filopodia on their anterior/posterior edge relative to their mediolateral edge. In support of our hypothesis, we demonstrate that ethanol disrupts protrusion polarity, convergent extension, and *shha* localization, with greater disruption observed in *vangl2* mutants.

MATERIALS AND METHODS

Zebrafish (Danio rerio) Care and Use

Zebrafish were cared for using standard IACUC-approved protocols at the University of Texas at Austin. The wild-type AB strain was used for RNA-seq analysis. The *vangl2^{m209}* allele, originally described as *tri^{m209}* [39], was obtained from the Zebrafish International Resource Center (ZIRC) as reported [13]. The gpc4^{fr6} line was provided by Dr. Lila Solnica-Krezel. Adult fish were maintained on a 14h/10h light-dark cycle at 28.5°C. Embryos were collected and staged according to morphology and somite number [58].

Chemical treatments

Embryos were treated with 1% ethanol diluted in embryo media. This subteratogenic dose mimics an acute (binge-like) alcohol exposure roughly equivalent to a blood alcohol concentration of 0.19% in humans. AB embryos were treated with 50 μM cyclopamine (Toronto Research Chemicals C988400), diluted in embryo media. The concentration of ethanol (vehicle) in the embryo media was controlled for between treatment groups and equals a final concentration of 0.5% in the EM. Blebbistatin (Sigma-Aldrich B0560) was dissolved in DMSO and diluted to a concentration of 37.5 μM in 3 mL of embryo media. Following chemical exposure, embryos were washed 3x in embryo media and then grown to 4 dpf before fixation.

mRNA injection

memGFP cloned in the pCS2 vector was linearized by *NotI* restriction endonuclease. Synthetic mRNA was generated using the mMESSAGE mMACHINE™

526   SP6 Transcription Kit (Invitrogen, AM1340). Embryos were microinjected at 100pg

527   between the 8-16 cell stage [40].

Sample collection and RNA extraction

Single embryos were manually dechorionated and collected in a 1.75 mL

microcentrifuge tube with 500 mL of TRIzol reagent (Life Technologies, 15596-026).

Embryos were homogenized with a motorized pestle (VWR, 47747-370) and stored at -

80°C until RNA extraction. Total RNA was processed according to the TRIzol RNA

isolation protocol. Samples were re-suspended with 50 μL of nuclease-free water and

subsequently purified using the RNA Clean & Concentrator kit (Zymo, R1018). The

concentration of each sample was determined using a Nanodrop spectrophotometer. The

quality of total RNA was analyzed with the Agilent BioAnalyzer to ensure that the RNA

Integrity Number (RIN) was ≥ 8. Samples were submitted to the Genomic Sequencing and

Analysis Facility (GSAF) at the University of Texas at Austin. The GSAF performed

standard RNA-Seq library preparations with poly-A mRNA capture.

RNA-seq data processing

Sequencing on the NextSeq 500 platform produced an average of 40.8 million ±

1.4 million (SE) raw paired end reads per sample. Adapter trimming was performed using

Cutadapt with a minimum length of 25 bp [59]. Following adapter trimming, we retained

an average of 40.0 ± 1.3 million (SE) reads per sample. Genome Reference Consortium

Zebrafish Build 10 (GRCz10) for *D. rerio* was downloaded from Ensembl [60]. Trimmed

reads were mapped to the reference using STAR [61]. Mean mapping efficiency was 78.2%

± 0.8% (SE). Following mapping PCR duplicates were removed using Picard

(https://broadinstitute.github.io/picard/). Duplication rate was estimated at 85% ± 0.8%

(SE). Sorting and conversion between SAM and BAM files was performed using samtools

[62]. Reads mapping to annotated genes were counted with HTseq version 0.6.1p1 using

the intersection nonempty mode [63]. The final number of reads mapped to annotated genes

was on average 3.9 ± 0.2 million reads per sample. Detailed instructions and example

commands for implementing the data processing steps described above are available on

Github (https://github.com/grovesdixon/Drerio_early_ethanol_RNAseq).

Differential expression analysis

Normalization and statistical analysis of read counts was performed using DESeq2

[64]. Factors included in the differential expression models were *ethanol treatment*

(control, treated), *developmental timepoint* (8 hpf, 10 hpf, and 14 hpf), and *sequencing*

batch (experiment 1 or experiment 2). Because none of the 6 hpf samples were treated

with ethanol, these samples were not included. We tested for differential expression

associated with ethanol treatment using likelihood ratio tests—comparing the model

including all three factors to a reduced model that did not include ethanol treatment. To

further examine stage-specific ethanol effects we split the samples by developmental

timepoint and tested for ethanol effects within each group.

Weighted Gene Correlation Network Analysis (WGCNA)

Gene expression data were further analyzed with Weighted Gene Correlation

Network Analysis (WGCNA) [22]. For input into the analysis, we used variance stabilized

counts generated using the rlog function in DESeq2 [64]. Genes that were not sequenced

across sufficient samples (4320 in total) were removed using the goodSamplesGenes

function in the WGCNA package. Because there were no ethanol treated samples for the

six hour timepoint, the five samples from this timepoint were removed before further

analysis. To ensure sufficient expression for correlation detection, genes were further

filtered based on a base mean expression cutoff of 5. We controlled for batch effects using

the ComBat function from the R package sva [65]. We selected a soft threshold of 15,

where the scale free topology model fit surpassed 0.8. WGCNA was run with a minimum

module size of 10. Following network analysis, we tested for GO enrichment within

modules using Fisher's exact tests.

GO enrichment analysis

Enrichment of Gene Ontology (GO) terms for ethanol responsiveness was tested

using two-tailed Mann-Whitney U-tests [66] followed by Benjamini-Hochberg procedure

for false discovery correction [67]. The results were plotted as a dendrogram tracing

hierarchical relationships between significant GO terms. The direction of enrichment (for

upregulation or downregulation) was indicated by text color and significance of enrichment

by font type. An advantage of this approach is that it does not require an arbitrary cutoff to

provide counts of "significant" and "non-significant" genes as in typical enrichment tests.

Quantitative Real-Time RT-PCR (qRT-PCR)

To validate our RNA-seq data, we selected two genes to test using qRT-PCR. Total

RNA was reverse transcribed using SuperScript™ First-Strand Synthesis System for RT-

PCR (Invitrogen) with oligo-d(T) primers. qRT-PCR was performed with Power Sybr

Green PCR Master Mix (Thermo Fisher Scientific, 4367659) on the Applied Biosystems

ViiA™ 7 Real-Time PCR System. QuantStudio Real-Time PCR Software was used for

data analysis using the $2^{-\Delta\Delta Ct}$ method. The endogenous control *lsm12b* was selected based

on its stable expression profiles across treatment and stage groups in the RNA-seq datasets.

KEGG enrichment methods and results

We tested for enrichment of significant upregulated and downregulated genes

among KEGG pathways separately using Fisher's exact tests. For each KEGG pathway,

we tested a two-way contingency table with inclusion in the KEGG pathway as columns,

and significant (FDR < 0.05) upregulation or downregulation as rows. P-values were then

adjusted for multiple tests using Benjamini Hochberg Method [67].

Three KEGG pathways were enriched for significant differential expression. Two

for upregulated genes, and one for downregulated genes. The upregulated KEGG pathways

(dre00250 and dre00480) were (1) Alanine, aspartate and glutamate metabolism, and (2)

Glutathione metabolism. The downregulated KEGG pathway (dre03013) was RNA

transport.   The   pathview   figures   for   these   are   in

kegg_pathways/significant_pathway_figures/ in the git repository.

Cartilage and Bone Staining and Measurements

607   Embryos were fixed at 4 days post fertilization (dpf) and stained with Alcian blue

608 for cartilage and Alizarin red for mineralized bone [68]. Whole-mounts of $vangl2^{m209}$

609 embryos were captured using a Zeiss Axio Imager A1 microscope. To assess the degree of

610 cyclopia, the distance between the medial edges of the lenses was measured using the

611 AxiovisionLE software.

In Situ Hybridization

613   Antisense digoxygenin-labeled riboprobes for *shha*, *pax2*, *dlx3*, *six3a* and *rx3*

614 (together as a gift from Dr. Steve Wilson) were used. Whole-mount *in situ* hybridization

615 was performed as described [69]. Images were captured using the Zeiss Axio Imager A1

616 and expression domains were measured using the AxiovisionLE software. An ANOVA

617 and post-hoc Tukey's test were used for statistical analyses.

TUNEL Staining

Whole-mount TUNEL staining was based on previous characterizations [12]. Samples were fixed overnight in 4% paraformaldehyde in PBS (PFA) at 4°C. Samples were dehydrated in methanol and subsequently rehydrated in phosphate-buffered saline containing 0.5% Triton X-100 (PBTx). Samples were permeabilized with 25 ug/mL proteinase K (1mg/ml) in PBT for 30 min. After two, 5 min washes with PBTx, samples were fixed with 4% PFA for 20 min at room temperature. Residual PFA was removed with four, 5 min washes of PBTx. Samples were incubated with 50 µl of 1:10 Enzyme:TUNEL reagent (TdT and fluorescein-dUTP) (Roche, Cat No. 11684795910) at 37°C for 3 h in the dark. The reaction was stopped with two, 5 min washes of PBTx. Confocal images were captured with a Zeiss LSM 710.

Confocal Imaging

Control and ethanol-treated (6-10 hpf) embryos were dechorionated and staged at 10 hpf. Homozygous mutants were distinguished between wild-type and heterozygotes by their body axis elongation at bud stage. Live embryos were mounted in methyl cellulose. Images were acquired using a Zeiss LSM710 confocal microscope using a 10x and 60x objective.

Quantification of filopodia

Number of filopodia was acquired through projections of Z-stacks using the 60x objective on the Zen software. Angle of projections were calculated using Fiji.

Rose diagrams of the mean number of filopodia found were plotted using geom_bar() and the coord_polar() functions from ggplot2. Mean numbers of filopodia for different embryonic regions were compared between groups using ANOVA and Tukey's

'Honest Significant Difference' method. Significantly distinct means were assigned based

on the Tukey's 'Honest Significant Difference' results using the R package multcomp [70].

ACKNOWLEDGEMENTS

We thank Dr. Steve Wilson for his kind contributions of the *six3a* and *rx3* riboprobes and

Dr. John Wallingford for his contributions of the memGFP plasmid. We also thank Dr. Lila

Solnica-Krezel for providing the *gpc4^{fr6}* line. We are grateful to Anna Louise Percy, Angie

Martinez, and Cadianna Cotham, for maintenance and care of all zebrafish lines. Lastly,

we are grateful to our former undergraduate researchers, Jenna Beam, Elaine Avshman,

and Jennyly Nguyen, for their assistance in the lab.
650

Bibliography

1.   Czeizel AE. Birth defects are preventable. Int J Med Sci. 2005;2(3):91-2. Epub
2005/07/12. doi: 10.7150/ijms.2.91. PubMed PMID: 16007259; PubMed Central PMCID:
PMCPMC1168872.

2.   Shepard TH. Detection of human teratogenic agents. The Journal of Pediatrics.
1982;101(5):810-5.

3.   Harada M. Minamata Disease: Methylmercury Poisoning in Japan Caused by
Environmental Pollution. Critical Reviews in Toxicology. 1995;25(1):1-24.

4.   Kim JH, Scialli AR. Thalidomide: the tragedy of birth defects and the effective
treatment   of   disease.   Toxicol   Sci.   2011;122(1):1-6.   Epub   2011/04/22.   doi:
10.1093/toxsci/kfr088. PubMed PMID: 21507989.

5.   May PA, Gossage JP, Kalberg WO, Robinson LK, Buckley D, Manning M, et al.
Prevalence and Epidemiologic Characteristics of FASD from Various Research Methods
with an Emphasis on Recent Inschool Studies Developmental Disabilities 2009;15:176-92.

6.   Jones KL, Smith DW. Recognition of the Fetal Alcohol Syndrome in Early Infancy.
The Lancet. 1973:999-1001.

7.   McCarthy N, Sidik A, Bertrand JY, Eberhart JK. An Fgf-Shh signaling hierarchy
regulates early specification of the zebrafish skull. Dev Biol. 2016;415(2):261-77. Epub
2016/04/10. doi: 10.1016/j.ydbio.2016.04.005. PubMed PMID: 27060628; PubMed
Central PMCID: PMCPMC4967541.

8.   Eberhart JK, Parnell SE. The Genetics of Fetal Alcohol Spectrum Disorders.
Alcohol Clin Exp Res. 2016;40(6):1154-65. Epub 2016/04/29. doi: 10.1111/acer.13066.
PubMed PMID: 27122355; PubMed Central PMCID: PMCPMC5125635.

9.   Streissguth AP, Dehaene P. Fetal alcohol syndrome in twins of alcoholic mothers:
concordance of diagnosis and IQ. Am J Med Genet. 1993;47(6):857-61. Epub 1993/11/01.
doi: 10.1002/ajmg.1320470612. PubMed PMID: 8279483.

10.   Downing C, Balderrama-Durbin C, Broncucia H, Gilliam D, Johnson TE. Ethanol
teratogenesis in five inbred strains of mice. Alcohol Clin Exp Res. 2009;33(7):1238-45.
Epub 2009/04/25. doi: 10.1111/j.1530-0277.2009.00949.x. PubMed PMID: 19389189;
PubMed Central PMCID: PMCPMC2951264.

11.   Loucks E, Carvan MJ, 3rd. Strain-dependent effects of developmental ethanol
exposure in zebrafish. Neurotoxicol Teratol. 2004;26(6):745-55. Epub 2004/09/29. doi:
10.1016/j.ntt.2004.06.017. PubMed PMID: 15451039.

12.   Lovely CB, Fernandes Y, Eberhart JK. Fishing for Fetal Alcohol Spectrum
Disorders: Zebrafish as a Model for Ethanol Teratogenesis. Zebrafish. 2016;13(5):391-8.
Epub 2016/05/18. doi: 10.1089/zeb.2016.1270. PubMed PMID: 27186793; PubMed
Central PMCID: PMCPMC5035362.

13.   Swartz ME, Wells MB, Griffin M, McCarthy N, Lovely CB, McGurk P, et al. A
screen of zebrafish mutants identifies ethanol-sensitive genetic loci. Alcohol Clin Exp Res.
2014;38(3):694-703. Epub 2013/10/30. doi: 10.1111/acer.12286. PubMed PMID:
24164477; PubMed Central PMCID: PMCPMC3959233.

14.   Solnica-Krezel L, Stemple DL, Mountcastle-Shah E, Rangini Z, Neuhauss SCF,
Malicki J, et al. Mutations affecting cell fates and cellular rearrangements during
gastrulation in zebrafish. Development. 1996;123:67-80.

15.   Marlow F, Zwartkruis F, Malicki J, Neuhauss SCF, Abbas L, Weaver M, et al.
Functional Interactions of Genes Mediating Convergent Extension, knypek and trilobite,
during the Partitioning of the Eye Primordium in Zebrafish. Developmental Biology.
1998;203:382-99.

16.   Blader P, Strahle U. Ethanol Impairs Migration of the Prechordal Plate in the
Zebrafish Embryo. Developmental Biology. 1998;201:185-201.

17.   Lovely CB, Nobles RD, Eberhart JK. Developmental age strengthens barriers to
ethanol accumulation in zebrafish. Alcohol. 2014;48(6):595-602. Epub 2014/07/12. doi:
10.1016/j.alcohol.2014.06.003. PubMed PMID: 25012627; PubMed Central PMCID:
PMCPMC4163099.

18.   Farrell JA, Wang Y, Riesenfeld SJ, Shekhar K, Regev A, Schier AF. Single-cell
reconstruction of developmental trajectories during zebrafish embryogenesis. Science.
2018;360(6392).

19.   Heisenberg C-P, Tada M, Rauch G-Jr, Saude L, Concha ML, Geisler R, et al.
Silberblick/Wnt11 mediates convergent extension movements during zebrafish
gastrulation. Nature. 2000;405:76-81.

20.   Topczewski J, Sepich DS, Myers DC, Walker C, Amores A, Lele Z, et al. The
Zebrafish Glypican Knypek Controls Cell Polarity during Gastrulation Movements of
Convergent Extension. Developmental Cell. 2001;1:251-64.

21.   White JJ, Mazzeu JF, Coban-Akdemir Z, Bayram Y, Bahrambeigi V, Hoischen A,
et al. WNT Signaling Perturbations Underlie the Genetic Heterogeneity of Robinow
Syndrome. Am J Hum Genet. 2018;102(1):27-43. Epub 2017/12/26. doi:
10.1016/j.ajhg.2017.10.002. PubMed PMID: 29276006; PubMed Central PMCID:
PMCPMC5777383.

22.   Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network
analysis. BMC Bioinformatics. 2008;9:559. Epub 2008/12/31. doi: 10.1186/1471-2105-9-
559. PubMed PMID: 19114008; PubMed Central PMCID: PMCPMC2631488.

23.   Loucks EJ, Ahlgren SC. Deciphering the role of Shh signaling in axial defects
produced by ethanol exposure. Birth Defects Res A Clin Mol Teratol. 2009;85(6):556-67.
Epub 2009/02/25. doi: 10.1002/bdra.20564. PubMed PMID: 19235835.

24.   White RJ, Collins JE, Sealy IM, Wali N, Dooley CM, Digby Z, et al. A high-
resolution mRNA expression time course of embryonic development in zebrafish. Elife.
2017;6. Epub 2017/11/17. doi: 10.7554/eLife.30860. PubMed PMID: 29144233; PubMed
Central PMCID: PMCPMC5690287.

25.   Siefert JC, Georgescu C, Wren JD, Koren A, Sansam CL. DNA replication timing
during development anticipates transcriptional programs and parallels enhancer activation.
Genome Res. 2017;27(8):1406-16. Epub 2017/05/18. doi: 10.1101/gr.218602.116.
PubMed PMID: 28512193; PubMed Central PMCID: PMCPMC5538556.

26.   Subramanian A, Narayan R, Corsello SM, Peck DD, Natoli TE, Lu X, et al. A Next
Generation Connectivity Map: L1000 Platform and the First 1,000,000 Profiles. Cell.
2017;171(6):1437-52 e17. Epub 2017/12/02. doi: 10.1016/j.cell.2017.10.049. PubMed
PMID: 29195078; PubMed Central PMCID: PMCPMC5990023.

27.   Chiang C, Litingtung Y, Lee E, Young KE, Corden JL, Westphal H, et al. Cyclopia
and defective axial patterning in mice lacking Sonic hedgehog gene function. Nature.
1996;383:407-13.

28.   Eberhart JK, Swartz ME, Crump JG, Kimmel CB. Early Hedgehog signaling from
neural to oral epithelium organizes anterior craniofacial development. Development.
2006;133(6):1069-77. Epub 2006/02/17. doi: 10.1242/dev.02281. PubMed PMID:
16481351.

29.   Sulik KK, Johnston MC, Webb MA. Fetal Alcohol Syndrome: Embryogenesis in a
Mouse Model. Science. 1981;214(4523):936-8.

30.   Cohen MM, Sulik KK. Perspectives on holoprosencephaly: Part II. Central nervous
system, craniofacial anatomy, syndrome commentary, diagnostic approach, and
experimental studies. Journal of craniofacial genetics and developmental biology.
1992;12:196-244.

31.   Hong M, Krauss RS. Ethanol itself is a holoprosencephaly-inducing teratogen.
PLoS One. 2017;12(4):e0176440. Epub 2017/04/26. doi: 10.1371/journal.pone.0176440.
PubMed PMID: 28441416; PubMed Central PMCID: PMCPMC5404885.

32.   Kietzman HW, Everson JL, Sulik KK, Lipinski RJ. The teratogenic effects of
prenatal ethanol exposure are exacerbated by Sonic Hedgehog or GLI2 haploinsufficiency
in the mouse. PLoS One. 2014;9(2):e89448. Epub 2014/03/04. doi:
10.1371/journal.pone.0089448. PubMed PMID: 24586787; PubMed Central PMCID:
PMCPMC3929747.

33.   Ahlgren SC, Thakur V, Bronner-Fraser M. Sonic hedgehog rescues cranial neural
crest from cell death induced by ethanol exposure. Proc Natl Acad Sci U S A.
2002;99(16):10476-81. Epub 2002/07/26. doi: 10.1073/pnas.162356199. PubMed PMID:
12140368; PubMed Central PMCID: PMCPMC124946.

34.   Santos-Ledo A, Cavodeassi F, Carreno H, Aijon J, Arevalo R. Ethanol alters gene
expression and cell organization during optic vesicle evagination. Neuroscience.
2013;250:493-506. Epub 2013/07/31. doi: 10.1016/j.neuroscience.2013.07.036. PubMed
PMID: 23892006; PubMed Central PMCID: PMCPMC3988994.

35.   Krauss S, Johansen T, Korzh V, Fjose A. Expression of the zebrafish paired box
gene pax[zf-b] during early
neurogenesis. Development. 1991;113:1193-206.

36.   Mathers PH, Grinberg A, Mahon KA, Jamrich M. The Rx homeobox gene is
essential for vertebrate eye development. Nature. 1997;387:603-7.

37.   Bénazéraf B, Francois P, Baker RE, Denans N, Little CD, Pourquie O. A random
cell motility gradient downstream of FGF controls elongation of an amniote embryo.
Nature. 2010;466:248–52.

38.   Wallingford JB, Fraser SE, Harland RM. Convergent Extension: The Molecular
Review Control of Polarized Cell Movement during Embryonic Development.
Developmental Cell. 2002;2:695–706.

39.   Jessen JR, Topczewski J, Bingham S, Sepich DS, Marlow F, Chandrasekhar A, et
al. Zebrafish trilobite identifies new roles for Strabismus in gastrulation and neuronal
movements. Nat Cell Biol. 2002;4(8):610-5. Epub 2002/07/10. doi: 10.1038/ncb828.
PubMed PMID: 12105418; PubMed Central PMCID: PMCPMC2219916.

40.   Love AM, Prince DJ, Jessen JR. Vangl2-dependent regulation of membrane
protrusions and directed migration requires a fibronectin extracellular matrix.
Development. 2018;145(22). doi: UNSP dev165472
10.1242/dev.165472. PubMed PMID: WOS:000451191500007.

41.   Berres ME, Garic A, Flentke GR, Smith SM. Transcriptome Profiling Identifies
Ribosome Biogenesis as a Target of Alcohol Teratogenicity and Vulnerability during Early
Embryogenesis. PLoS One. 2017;12(1):e0169351. Epub 2017/01/04. doi:

10.1371/journal.pone.0169351. PubMed PMID: 28046103; PubMed Central PMCID:
PMCPMC5207668.

42.   Green ML, Singh AV, Zhang Y, Nemeth KA, Sulik KK, Knudsen TB.
Reprogramming of genetic networks during initiation of the Fetal Alcohol Syndrome. Dev
Dyn. 2007;236(2):613-31. Epub 2007/01/04. doi: 10.1002/dvdy.21048. PubMed PMID:
17200951.

43.   Da Lee R, Rhee GS, An SM, Kim SS, Kwack SJ, Seok JH, et al. Differential gene
profiles in developing embryo and fetus after in utero exposure to ethanol. J Toxicol
Environ   Health   A.   2004;67(23-24):2073-84.   Epub   2004/10/30.   doi:
10.1080/15287390490515001. PubMed PMID: 15513904.

44.   Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, et al. The
zebrafish reference genome sequence and its relationship to the human genome. Nature.
2013;496(7446):498-503. Epub 2013/04/19. doi: 10.1038/nature12111. PubMed PMID:
23594743; PubMed Central PMCID: PMCPMC3703927.

45.   Marcucio RS, Cordero DR, Hu D, Helms JA. Molecular interactions coordinating
the development of the forebrain and face. Dev Biol. 2005;284(1):48-61. Epub 2005/06/28.
doi: 10.1016/j.ydbio.2005.04.030. PubMed PMID: 15979605.

46.   Aoto K, Shikata Y, Imai H, Matsumaru D, Tokunaga T, Shioda S, et al. Mouse Shh
is required for prechordal plate maintenance during brain and craniofacial morphogenesis.
Dev Biol. 2009;327(1):106-20. Epub 2008/12/24. doi: 10.1016/j.ydbio.2008.11.022.
PubMed PMID: 19103193.

47.   Hong M, Krauss RS. Cdon mutation and fetal ethanol exposure synergize to
produce midline signaling defects and holoprosencephaly spectrum disorders in mice.
PLoS Genet. 2012;8(10):e1002999. Epub 2012/10/17. doi: 10.1371/journal.pgen.1002999.
PubMed PMID: 23071453; PubMed Central PMCID: PMCPMC3469434.

48.   Chen W, Burgess S, Hopkins N. Analysis of the zebrafish smoothened mutant
reveals conserved and divergent functions of hedgehog activity. Development. 2001;128:
2385-96.

49.   Brand M, Heisenberg C-P, Warga RM, Pelegri F, Karlstrom RO, Beuchle D, et al.
Mutations affecting development of the midline and general body shape during zebrafish
embryogenesis. Development. 1996;123: 129-42.

50.   Yelin R, Schyr RB, Kot H, Zins S, Frumkin A, Pillemer G, et al. Ethanol exposure
affects gene expression in the embryonic organizer and reduces retinoic acid levels. Dev
Biol. 2005;279(1):193-204. Epub 2005/02/15. doi: 10.1016/j.ydbio.2004.12.014. PubMed
PMID: 15708568.

51.   Zhang Y, Ding Y, Chen YG, Tao Q. NEDD4L regulates convergent extension
movements in Xenopus embryos via Disheveled-mediated non-canonical Wnt signaling.
Dev Biol. 2014;392(1):15-25. Epub 2014/05/17. doi: 10.1016/j.ydbio.2014.05.003.
PubMed PMID: 24833518.

52.   Sarmah S, Muralidharan P, Curtis CL, McClintick JN, Buente BB, Holdgrafer DJ,
et al. Ethanol exposure disrupts extraembryonic microtubule cytoskeleton and embryonic
blastomere cell adhesion, producing epiboly and gastrulation defects. Biol Open.
2013;2(10):1013-21. Epub 2013/10/30. doi: 10.1242/bio.20135546. PubMed PMID:
24167711; PubMed Central PMCID: PMCPMC3798184.

53.   Solnica-Krezel L, Sepich DS. Gastrulation: making and shaping germ layers. Annu
Rev Cell Dev Biol. 2012;28:687-717. Epub 2012/07/19. doi: 10.1146/annurev-cellbio-
092910-154043. PubMed PMID: 22804578.

54.   Keller R. Shaping the Vertebrate Body Plan by
Polarized Embryonic Cell Movements. Science. 2002;298:1950-4.

55.   Solnica-Krezel L. Conserved patterns of cell movements during vertebrate
gastrulation. Curr Biol. 2005;15(6):R213-28. Epub 2005/03/31. doi:
10.1016/j.cub.2005.03.016. PubMed PMID: 15797016.

56.   Chuang JC, Raymond PA. Zebrafish genes rx1 and rx2 help define the region of
forebrain that gives rise to retina. Dev Biol. 2001;231(1):13-30. Epub 2001/02/22. doi:
10.1006/dbio.2000.0125. PubMed PMID: 11180949.

57.   Roszko I, D SS, Jessen JR, Chandrasekhar A, Solnica-Krezel L. A dynamic
intracellular distribution of Vangl2 accompanies cell polarization during zebrafish
gastrulation. Development. 2015;142(14):2508-20. Epub 2015/06/13. doi:
10.1242/dev.119032. PubMed PMID: 26062934; PubMed Central PMCID:
PMCPMC4510859.

58.   Kimmel CB, Ballard WW, Kimmel SR, Ullman B, Schilling TF. Stages of
Embryonic Development of the Zebrafish. Developmental Dynamics. 1995;203:253-310.

59.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnetjournal. 2011;17(1):10-2. Epub May 2011.

60.   Aken BL, Achuthan P, Akanni W, Amode MR, Bernsdorff F, Bhai J, et al. Ensembl
2017. Nucleic Acids Res. 2017;45(D1):D635-D42. Epub 2016/12/03. doi:
10.1093/nar/gkw1104. PubMed PMID: 27899575; PubMed Central PMCID:
PMCPMC5210575.

61.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:
ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21. Epub 2012/10/30.
doi: 10.1093/bioinformatics/bts635. PubMed PMID: 23104886; PubMed Central PMCID:
PMCPMC3530905.

62.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9. Epub
2009/06/10. doi: 10.1093/bioinformatics/btp352. PubMed PMID: 19505943; PubMed
Central PMCID: PMCPMC2723002.

63.   Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-
throughput sequencing data. Bioinformatics. 2015;31(2):166-9. Epub 2014/09/28. doi:
10.1093/bioinformatics/btu638. PubMed PMID: 25260700; PubMed Central PMCID:
PMCPMC4287950.

64.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550. Epub 2014/12/18. doi:
10.1186/s13059-014-0550-8. PubMed PMID: 25516281; PubMed Central PMCID:
PMCPMC4302049.

65.   Leek JT, Storey JD. Capturing Heterogeneity in Gene Expression Studies by
"Surrogate Variable Analysis". PLoS Genetics. 2005;preprint(2007). doi:
10.1371/journal.pgen.0030161.eor.

66.   Dixon GB, Davies SW, Aglyamova GV, Meyer E, Bay LK, Matz MV. Genomic
determinants of coral heat tolerance across latitudes. Science. 2015;348 (6242).

67.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series B
(Methodological). 1995;57(1):289-300.

68.   Walker MB, Kimmel CB. A two-color acid-free cartilage and bone stain for
zebrafish larvae. Biotech Histochem. 2007;82(1):23-8. Epub 2007/05/19. doi:
10.1080/10520290701333558. PubMed PMID: 17510811.

69.   Miller CT, Schilling TF, Lee K-H, Parker J, Kimmel CB. sucker encodes a
zebrafish Endothelin-1 required for ventral pharyngeal arch development. Development.
2000;127:3815-28.

70.   Hothorn T, Bretz F, Westfall P. Simultaneous inference in general parametric
models. Biom J. 2008;50(3):346-63. Epub 2008/05/16. doi: 10.1002/bimj.200810425.
PubMed PMID: 18481363.

889
890

Figures

892
Fig 1. Ethanol interacts with *vangl2* during early embryogenesis. (A) Treatment of
vangl2 embryos with 1% ethanol at four different stages comprising late blastula to early
gastrula [3.3, 4, 4.5, and 6 hours post fertilization (hpf)] for 24 hours. Homozygous mutants
showed a significant decrease in the inner lens-to-lens width when treated from 6 to 30 hpf.
(B) Heterozygous mutants were most sensitive to ethanol when exposed from 3.3 to 27.3
hpf (n=4/18 cyclopic). Homozygous mutants were most sensitive when exposed from 6 to
30 hpf (n=5/5 cyclopic).
900

901
Fig 2. Developmental age is the dominant driver of variation in the dataset. (A)
Schematic representation of the RNA-seq experimental design. Wild-type AB embryos
were exposed to a subteratogenic dose of 1% ethanol in the embryo media at shield stage
(6 hpf). Embryos were subsequently collected at 8 and 10 hpf for experiment 1 and (B) 8,
10, and 14 hpf, for experiment 2. Each sample consisted of a single zebrafish embryo with
5 control and 5 ethanol-treated samples per timepoint with the exception of the 6 hpf
timepoint, which only had 5 control samples. (C-F) Principal components analysis (PCA)
of top 25,000 most variable genes. The percentage of variance explained is given on each
axis label. (C) *PC1* and *PC2* color coded by developmental timepoint. (D) *PC1* and *PC2*
color coded by RNA-seq experiment (batch). (E) *PC1* and *PC2* color coded by ethanol
treatment group. (F) *PC8* and *PC9* showing separation of ethanol treated and control
samples.
914

915
Fig 3. Effects on transcription are largely distinct between developmental time points.
(A) Volcano plot showing differential expression due to ethanol treatment across all
samples. Significant genes (FDR < 0.1)* are indicated in red. Genes that were significant
at all time points (overlap) are indicated in blue. (B) Venn-diagram showing overlap of
significant genes between the three individual timepoints. (C) Table of genes that were
significant for each timepoint individually. (D) KEGG pathway schematic illustrating
differential expression due to ethanol treatment in the Wnt/planar cell polarity (Wnt/PCP)
pathway. Color coding indicates $\log_2$ fold differences due to ethanol treatment across all
samples, red indicates upregulation due to ethanol, blue indicates downregulation due to
ethanol. Only pathway members with significant changes were color coded. In the event
that multiple genes from the dataset were annotated with the same pathway member, the
$\log_2$ fold difference for the gene with the greatest absolute value for the difference was
used.
929

930
Fig 4. Ethanol indirectly attenuates Shh signaling. (A) Alcian blue and Alizarin red
whole mount staining of untreated (control), cyclopamine-treated (50 μM), and ethanol-
and cyclopamine-treated (1% ethanol plus 50 μM cyclopamine) wild-type embryos from 6
hpf to 4 dpf. Panels represent the spectrum of phenotypes observed for the treatment
groups. Embryos were fixed at 4 dpf. Dorsal view, anterior to the left. (B) Quantification
of the effect of ethanol and cyclopamine on the eye field. Inner lens-to-lens width was used
as a morphometric measure of cyclopia. Both cyclopamine alone and cyclopamine and
ethanol, were significantly different from controls (p<0.0001). (C) qRT-PCR of *ptch2* in
10 hpf wild-type embryos. *slc25a5* (*solute carrier family 25*) was used as a normalization
control. (D) KEGG pathway schematic illustrating differential expression due to ethanol
treatment in the sonic hedgehog (Shh) signaling pathway. Color coding indicates $\log_2$ fold
differences due to ethanol treatment across all samples; red indicates upregulation due to
ethanol, blue indicates downregulation due to ethanol. In the event that multiple genes from
the dataset were annotated with the same pathway member, the $\log_2$ fold difference for the
gene with the greatest absolute value for the difference was used.
946

947
Fig 5. Ethanol disrupts convergent extension. (A) *vangl2* embryos were treated with 1%
ethanol from 3.3 to 10 hpf. Embryos were subsequently harvested at 10 hpf for *in situ*
hybridization with *shha* (midline) and *pax2a* (midbrain-hindbrain boundary) probes.
Dorsal view, anterior to the left. (B) Quantification for the normalized *shh/pax2a*
expression domains were calculated using the AxiovisionLE software. Two-way ANOVA
followed by Tukey's post-hoc test was used to analyze the results. Differences were
observed between control genotypes and ethanol-treated homozygous mutants. (C)
Expression pattern of transcription factor *six3a*, stained using whole mount *in situ*
hybridization at 11 hpf.
957

958
Fig 6. Mutation in *gpc4* exacerbates cyclopia in a dose-dependent manner. (A)
Normalized read counts indicating expression of *gpc4* across different subsets of the
dataset. (B) qRT-PCR of *gpc4* in 10 hpf wild-type embryos. *actb1* (*actin, beta 1*), *lsm12b*
(*like-Sm protein 12 homolog b*), and *slc25a5* (*solute carrier family 25*) were used for
normalization. (C) Alcian blue and Alizarin red whole- and flat-mount staining of untreated
and 1% ethanol-treated (6 hpf – 4 dpf) *gpc4* homozygous mutants. Embryos fixed at 4 dpf.
Dorsal view, anterior to the left. (D) Functional analyses of *vangl2;gpc4* double mutants.
Enhanced cyclopia was observed in ethanol-treated *vangl2;gpc4* double mutants with loss
of at least one copy of either gene.
968

Fig 7. Blebbistatin phenocopies ethanol treatment in *vangl2* mutant embryos. (A)
Alcian blue and Alizarin red whole-mount staining of *vangl2* embryos treated with 37.5µM
blebbistatin (6-10 hpf). Embryos fixed at 4 dpf. Dorsal view, anterior to the left. (B)
Quantification of the inner lens-to-lens width of control and blebbistatin-treated embryos.
Lettering indicates differences in significance based on multiple comparisons test using
Tukey's HSD. (C) Representation of the percentage of control and blebbistatin-treated
embryos exhibiting cyclopia. Genotypes indicated on the x-axes.

979
Fig 8. Membrane protrusions in untreated and ethanol-treated (6-10 hpf) *vangl2*
embryos. (A) Confocal images of untreated and ethanol-treated *vangl2* mesodermal cells
expressing memGFP at 10 hpf. n=6 (B) Rose diagram of filopodial projections representing
length and distribution around the circumference of a cell. 24 bins, divided into 15° angles
(C) Rose diagram of filopodial projections representing length and distribution around the
circumference of a cell. 4 bins, divided into 90° angles. Horizontal plane (0°) represents the
mediolateral axis; vertical plane (90°) represents the anterior-posterior axis.

Supporting Information

993
994 **S1 Fig. Time is the major driver of variation in the dataset.** Heatmap showing overall
995 correlation of gene expression among samples. Samples were hierarchically clustered
996 based on similarity. Treatment [control or ethanol treated (C or E)] and developmental age
997 [hours post fertilization (6, 10, 8, or 14)] are indicated in the row and column labels.
998

999
S2 Fig. There are more upregulated than downregulated genes among ethanol-
treated individuals. Volcano plot showing variation in the transcriptional response to
ethanol treatment across developmental timepoints. Significant genes (FDR < 0.1) are
indicated in red. For each subset, the names of the topmost significantly dysregulated genes
are noted near gene's data point (A) All timepoints combined together (B) 8 hpf only (C)
10 hpf only (D) 14 hpf only.
1006

1007
S3 Fig. WGCNA identifies two modules that are significantly correlated with ethanol
exposure. (A) Dendrogram illustrating the hierarchical clustering of the genes and with
their corresponding modules colors. The top layer of colors indicates the Dynamic Tree
Cutoff, including all assigned modules, before merging by module similarity. The bottom
layer indicates the module colors after merging. These merged modules were used for
further analysis. (B) Heatmap of module-trait correlations. The eigengene for each module
was correlated with ethanol treatment (ethanol), hours post fertilization (age), experimental
batch (seqjob), and as a negative control, a randomly shuffled version of the ethanol
treatments (rand.eth). Intensity of the color in each cell indicates the strength of correlation
between the module (row labels) and the sample trait (column labels). Two modules,
(mediumpurple4 and darkolivegreen4) significantly correlated with ethanol treatment (p <
0.05). (C) Scatterplots of correlation with ethanol treatment against module membership.
Each datapoint is a gene assigned to the indicated module. Ethanol correlation is the
Pearson correlation between the gene's expression level and ethanol treatment. Module

membership is the correlation between the genes expression level and the module eigengene and describes how well the gene matches the overall patterns of the module. (D) Boxplot of log_2 fold differences due to ethanol for the two significant modules. (E) Gene ontology enrichment tree for Molecular Function for the darkolivegreen4 module. (F) Gene ontology enrichment tree for Molecular Function for the mediumpurple4 module. (G) Changes in gene expression from the RNA-seq were validated using wild-type embryos at 10 hpf. *slc16a9a* was selected from the mediumpurple4 module and (H) *znf1015* was selected from the darkolivegreen4 module. Fold change indicates the degree of change between untreated control and ethanol-treated stage-matched embryos. *lsm12b* (*like-Sm protein 12 homolog b*) was used as a normalization control gene.

1033
S4 Fig. Expression of hub genes for development-related modules indicate ethanol did
not retard developmental progression. (A-K) Development-related WGCNA modules
were identified as those with significant relationship to hours post fertilization (Pearson
correlation; $p<0.05$). The hub gene for each of these modules was identified as the gene
assigned to that module with the highest module membership (defined as the correlation of
the gene's expression level with the module eigengene). Normalized expression levels for
the hub genes were plotted against hours post fertilization. The relationship between
expression and time is largely consistent between the ethanol treated (E, teal) and control
(C, pink) samples. Even for the two modules associated with ethanol treatment
(mediumpurple4 and darkolivegreen4), the slopes of the lines are very similar, indicating
subteratogenic ethanol exposure did not cause significant developmental delay.

1045

1046
1047 **S5 Fig. Cell death by TUNEL at 11 hpf in untreated and ethanol-treated** *vangl2*
1048 **mutants.** The number of positive cells in the eye field (indicated by dashed circle) was not
1049 higher in homozygous or ethanol-treated mutants. ant = anterior; nc = notochord
1050

1051
1052 **S6 Fig. Ethanol alters *rx3* expression in the eye field.** Expression pattern of transcription
1053 factor *rx3*, stained using whole mount *in situ* hybridization at 12 hpf. Dorsal view, anterior
1054 to the left.
1055

1056
1057   **S7 Fig. Ethanol interacts with *gpc4*.** (A) Alcian blue and Alizarin red whole-mount
1058   staining of untreated and 1.6% ethanol-treated (6 hpf – 24 hpf) *gpc4* homozygous mutants.
1059   Embryos fixed at 4 dpf. Dorsal view, anterior to the left. (B) Dose-response curve of *gpc4*
1060   mutants treated with 1-1.9% ethanol (6-30 hpf).

1061

1062

1063

1064

1065