# Reference transcriptome data in silkworm

# *Bombyx mori*

Kakeru Yokoi[1,2,*,**], Takuya Tsubota[3,*], Jianqiang Sun[2], Akiya Jouraku[1], Hideki Sezutsu[3], Hidemasa Bono[4]

1 Insect Genome Research and Engineering Unit, Division of Applied Genetics, Institute of Agrobiological Sciences (NIAS), National Agriculture and Food Research Organization (NARO), 1-2 Owashi, Tsukuba, Ibaraki 305-8634, Japan

2 Research Center for Agricultural Information Technology (RCAIT), National Agriculture and Food Research Organization (NARO), Kintetsu Kasumigaseki Building Kasumigaseki 3-5-1 Chiyoda-ku, Tokyo 100-0013, Japan

3 Transgenic Silkworm Research Unit, Division of Biotechnology, Institute of Agrobiological Sciences (NIAS), National Agriculture and Food Research Organization (NARO), 1-2 Owashi, Tsukuba, Ibaraki 305-8634, Japan

4 Database Center for Life Science (DBCLS), Joint Support-Center for Data Science Research, Research Organization of Information and Systems, 1111 Yata, Mishima, Shizuoka 411-8540, Japan

*These authors equally contributed to this work.

**Corresponding author: Kakeru Yokoi

Email adresses

Kakeru Yokoi: yokoi123@affrc.go.jp

Takuya Tsubota: tsubota@affrc.go.jp

Sun Jianqiang: j.sun@affrc.go.jp

Akiya Jouraku: joraku@affrc.go.jp

Hideki Sedutsu: hsezutsu@affrc.go.jp

Hidemasa Bono: bono@dbcls.rois.ac.jp

## Abstract

The silkworm *Bombyx mori* has long been used in the silk industry and utilized in studies of physiology, genetics, molecular biology, and pathology. We recently reported high quality reference genome data for *B. mori.* In the present study, we constructed a

reference transcriptome data set using the reference genome data and RNA-seq data

of 10 tissues from P50T strain larvae. Reference transcriptome data contained 51,926

transcripts, with 39,619 having one or more coding sequence region. The abundance

of each transcript in the 10 tissues as well as 5 tissues from other strain larvae was

estimated, and hierarchical clustering was performed. The results obtained showed

that data on abundance were highly reproducible and there here is a little difference of

transcriptome abundance between the two strain larvae. New isoforms of silk-related

genes were searched for in the reference transcriptomes, and the longest isoform of

*sericin-1* possessing a long exon was identified. We also extracted transcripts that

were strongly expressed in one or more parts of the silk glands. An enrichment

analysis performed using the functional annotation data of the transcripts provided

novel insights into the functions of the silk gland parts. Therefore, the reference

transcriptome data set obtained has extended *B. mori* genomic and transcriptomic

insights and may contribute to advances in entomologic and vertebrate research,

including that on humans.

## 48    Introduction

The silkworm *Bombyx mori* is a lepidopteran insect that has been utilized in

studies of physiology, genetics, molecular biology, and pathology. Functional analyses

of genes related to hormone synthesis/degradation, pheromone reception, larval

marking formation, and virus resistance have been performed using this silkworm (Tan

et al., 2005; Ito et al., 2008; Sakurai et al., 2011; Daimon et al., 2015; Kondo et al.,

2017), and the findings obtained have contributed to the promotion of insect science.

The silkworm has the ability to produce large amounts of silk proteins, which is one of

the most prominent characteristics of this species. Silk proteins are mainly composed

of the fibrous protein Fibroin and aqueous protein Sericin and are produced in the

larval tissue silk gland (SG) (Yukuhiro et al., 2016). A transgenic technique has been

applied to the silkworm (Tamura et al., 2000), and has enabled the production of a

large amount of recombinant proteins through the introduction of transgenes, which are

overexpressed in the SG (Tatematsu et al., 2010). Through this method, it is possible

to utilize the silkworm as a significant bioreactor.

Based on its biological and industrial importance, the whole genome sequence

data of the silkworm was reported in 2004 from two research groups (Mita el al., 2004;

Xia et al., 2004), and was the first lepidopteran whole genome data. Silkworm whole

genome data was updated in 2008 from an international research group (International

Silkworm Genome Consortium 2008). Several data related to the silkworm genome

have since become available (e.g. microarray-based gene expression profiles in
multiple tissues, BAC-based linkage map full-length cDNA data and of *B. mori* in
KaikoBase) (Xia et al., 2007; Yamamoto et al., 2008; Suetsugu et al., 2013). These
findings have strongly promoted studies on *B. mori* and other lepidopteran insects in
the past decade.

In 2019, we reported a new and high-quality genome silkworm reference
genome assembly the silkworm p50T strain using PacBio long-read and Illumina short-
read sequencers (Kawamoto et al., 2019). The new genome assembly consists of 696
scaffolds with N50 of 16.8 M and only 30 gaps, and a new gene model based on this
sequence was predicted. These data are expected to be utilized in silkworm and
Lepidopteran research. Reference transcriptome data using this genome sequence
and the predicted gene set and transcriptome profile of important tissues significantly
contribute to advances in silkworm and Lepidopteran research. In the present study,
we constructed a reference transcript sequence data set using the RNA-seq data of 10
important tissues from silkworm larvae and reference genome data for improving the
predicted gene set data of Kawamoto et al. (2019) (Fig. 1). We also showed
comprehensive expression data for 10 different organs.   These results will contribute
to further advances in silkworm as well as entomological and vertebrate research,
including that on humans.

## 87  Results

<u>Reference transcriptome data</u>

Total RNAs were extracted from the fat body (FB), midgut (MG), Malpighian
tubules (MT), testis (TT), anterior silk gland (ASG), anterior part of the middle silk gland
(MSG-A), middle part of the middle silk gland (MSG-M), posterior part of the middle silk
gland (MSG-P) and posterior silk gland (PSG) of one male P50T larva and from the
ovary (OV) of a female larva. Extraction was repeated three times. Total RNAs were
sequenced and thirty sets of sequenced data were used as RNA-seq data. We
obtained "reference transcriptome data" by using the reference genome, gene model
data (Kawamoto et al., 2019), and RNA-seq data. Transcriptome data contains 51,926
transcripts in 24236 loci (Supporting data 1). The previously constructed gene model
data contained 16,880 genes in 16,845 loci, while our reference transcriptome data
contains new transcripts and loci (Fig. 2A). Among 51,926 transcripts, 7,704 transcripts
belonged to new loci while 27,342 transcripts are newly identified isoforms of which loci
was already identified in Kawamoto et al., (2019) (Fig. 2B). These results suggest that
our reference transcriptome data extend gene model data and contain many

unidentified transcripts and loci. To annotate transcripts, we predicted the coding

sequence region (CDS) and amino acid sequences (Supporting data 2) against all

transcripts. We found that 39,619 transcripts and 16,632 loci had at least one CDS.

The predicted amino acid sequences were used for gene functional annotations by a

homology search against human and *Drosophila* gene sets (Supporting data 3).

<u>Estimating the abundance of the reference transcriptome in multiple tissues</u>

We calculated the abundance of each transcript in ten tissues: FB, MG, MT, TT,

OV, ASG, MSG-A, MSG-M, MSG-P, PSG, TT, and OV. In comparisons of our

calculated results and evaluations of the reliability of our expression analysis, we

additionally quantified transcript expression by using public RNA-seq data that were

sequenced from the FB, MG, MT, TT, and SG of the *B. mori* o751 strain (Kikuchi et al.,

2017; Ichino et al., 2018; Kobayashi et al., 2019). To distinguish between our RNA-seq

data and public data, we referred to the FB, MG, MT, TT, and SG of the o751 strain as

BN_FB, BM_MG, BN_ MT, BM_TT, and BM_SG, respectively. Abundance values are

shown as transcripts per million (tpm) and listed in Supporting data 4.

Hierarchical clustering was performed for comparisons of transcriptome

abundance between multiple tissues (Fig. 3). The results obtained showed that, except

for MSG_P and MSG_M, all samples were clearly grouped by tissue types. Moreover,

clusters of samples that were the same tissues, but extracted from different strain

larvae, namely, FB and BN_FB, MT and BN_MT, MG and BN_MG, and TT and

BN_TT, were placed in adjacent positions, of which samples were obtained from

different studies as well as different strains. These results suggest that the data

obtained on abundance were highly reproducible and there were a little difference in

transcriptome profiles between P50T and o751 larvae. These results also indicated

that there were slight individual genetic differences between the larval samples used

and a marginal artificial effect, demonstrating that our abundance data may be used as

reference expression data for silkworm larvae.

<u>Transcripts of *sericin*, *fibroin*, and *fibrohexamerin* genes</u>

The *sericin*, *fibroin*, and *fibrohexamerin* (*fhx*) genes are known to play

important roles in silk synthesis in *B. mori*. Our transcriptome data contains several

isoforms of *sericin*, *fibroin*, *and fhx* (Table 1). A detailed sequence analysis was

performed to elucidate isoform structures. A previous analysis of *sericin-1* revealed that

this gene is composed of 9 exons, with exons 3-6 being under the selection of

alternative splicing (Garel et al., 1997; Yukuhiro et al., 2016). Within these exons, exon

6 is responsible for the most abundant component of the sericin protein (Sericin M),

which is ~6500 bp in length and encodes a serine-rich repetitive sequence (Yukuhiro et

al., 2016). However, the structure of this exon has not yet been elucidated in detail. We

herein found that exon 6 of MSTRG.2477.1, the longest *sericin-1* isoform identified in

the present study, had a length of 6234 bp (from 2,552,212 to 2,558,445 bp in

chromosome 11, see Supporting data 2 and Fig. 4A) In the present study, we newly

identified a *sericin-1* isoform that contained exon 6 with 6234 bp. We speculate that

this isoform corresponds to the full-length or nearly full-length transcript of *sericin-1*.

The product from this exon is enriched with serine residues and also has abundant

residues of glycine, asparagine and threonine (Supplemental Fig. 1).

Sericin-3 is another major silk protein that has a relatively soft texture (Takasu

et al., 2006; 2007). In gene model data, there is a frame shift in the predicted amino

acid sequence (KWMTBOMO08464), presumably due to the 73-bp deletion present in

exon 3. In the present study, we found that reference transcriptome data

(MSTRG.2595.1) provided an accurate gene structure. Sericin-4 is another recently

identified sericin protein that is composed of 34 exons (Dong et al., 2019). In gene

model data, it is split into three genes (KWMTBOMO06324, KWMTBOMO06325 and

KWMTBOMO06326) whereas our reference transcriptome data represent an exact

structure (MSTRG.2610.1, Fig. 4B).

In contrast, a better structure is provided by gene model data for the fibroin

heavy chain (*h-fib*); this gene encodes a protein with a large and highly repetitive

sequence (Zhou et al., 2000) and although there is a small deletion in the repeat motifs

for both models, the deletion length is shorter for gene model data (32 amino acid

deletion for gene model data and 223 for reference transcriptome data). Regarding

other silk genes (*sericin-2*, *fibroin light chain* (*l-fib*), and *fhx*), both models provide exact

structures (data not shown).

<u>Transcript abundance in the silk gland</u>

As described above, silk is synthesized in the SG. While the role of each SG

part in silk synthesis is known, the underlying molecular and genetic mechanisms

remain unclear. Therefore, the genes or transcripts that are strongly expressed in each

SG part need to be identified in order to elucidate these mechanisms. We searched for

transcripts that showed values of more than 30 transcripts per kilobase million (tpm) in

the five SG parts. The results obtained are shown in Fig. 5. The numbers of transcripts

that were strongly expressed in only ASG, MSG_A, MSG_M, MSG_P, and PSG were

351, 180, 99, 71, and 100, respectively, while more than 1,000 transcripts were

strongly expressed in all parts of the SG.

By using the annotation data of the strongly expressed transcripts, a functional

enrichment analysis (FEA) was performed using the transcripts strongly expressed in

each part of the SG to predict their role. In the enrichment analysis, we utilized the

annotation results against the human gene set. Fig. 6A shows the results of FEA using

the annotation of transcripts strongly expressed in MSG_P, MSG_M specific plus both

in MSG_P and MSG_M. The reason for utilizing MSG_P, MSG_M specific plus MSG_P

and MSG_M classes is that the samples of MSG_P and MSG_M in Fig. 3 did not form

different clusters, suggesting that both tissues have the same functions.   The highly

enriched function of the category (-log(P) > 10) was "Metabolism of RNA", while the

moderately enriched functions (6 < -log(P) < 10) were "ncRNA metabolic process",

"regulation of mRNA processing", "HIV Infection", and "Asparagine N-linked

glycosylation". In MSG_A, the moderately enriched function was "Metabolism of

vitamins and cofactors", while the highly enriched functions of the category were not

found (Fig. 6B). In ASG, the highly enriched functions of the category were

"carbohydrate metabolic process" and "Transport of small molecules" (Fig. 6C). The

moderately enriched functions were "anion transport", "Glycolysis/Gluconeogenesis",

"Ascorbate and aldarate metabolism", and "Metabolism of carbohydrates". In PSG, the

moderately enriched function was "tRNA modification", while the highly enriched

function of the category was not found (Fig. 6D).

## Discussion

In the present study, we obtained RNA-seq data on ten tissues of *B. mori* on the

3$^{rd}$ day of fifth instar larvae from the P50T strain. Using RNA-seq data and new

reference genome data (Kawamoto et al., 2019), we constructed reference

transcriptome data. Our transcriptome data contained new loci and isoforms, thereby

updating the reference genomic and transcriptome data of *B. mori*. The reference

transcriptome consists of 51,926 transcripts in 24,236 loci (16,632 loci have coding

genes), and 39,619 transcripts contain single or multiple CDS. In the mouse reference

data set (GRCm38.p6), there are 52332 loci (22,480 coding genes, 16,324 non-coding

genes, and 13,528 pseudogenes) and 142,333 transcripts

(http://asia.ensembl.org/Mus_musculus/Info/Annotation). In the human reference data

set (GRCh38.p12), there are 63,048 loci (20,454 coding genes, 23,940 non-coding

genes, and 15,204 pseudogenes) and 226,950 transcripts

(http://asia.ensembl.org/Homo_sapiens/Info/Annotation). In *Drosophila melanogaster*

(BDGP6.22), there are 17,753 loci (13,931 coding genes, 3,513 non-coding genes, and

309 pseudogenes) and 34,802 transcripts

(http://asia.ensembl.org/Drosophila_melanogaster/Info/Annotation). In Zebrafish
(GRCz11), there are 32,506 loci (25,592 coding genes, 6,599 non-coding genes, and
315 pseudogenes) and 59,878 transcripts
(http://asia.ensembl.org/Danio_rerio/Info/Annotation). In consideration of the basic
status of the reference data of these model species, our transcriptome data is not
unusual. It suggests that transcriptome data cover the majority of actual transcripts.
We estimated transcriptome abundance in multiple tissues plus several
tissues of other strain larvae. Transcriptome abundance in the tissues MG, TT, MT,
and FB did not markedly differ between the P50T and o751 strains. These results
suggest that these tissues at this stage did not contribute to phenotypic differences
between the two strains. To elucidate the underlying genetic mechanisms for
phenotypic differences, the RNA-seq data and transcriptome data of other stages are
needed. On the other hand, transcriptome abundance in MSG_M and MSG_P samples
was not divided into two independent clusters, suggesting that both parts have similar
roles in this stage, while MSG_A has distinct roles from the other parts.
We searched for new or previously unidentified isoforms of the *sericin, fibroin*,
*and fhx* genes in reference transcriptome data. While new or previously unidentified
isoforms of *sericin-2*, *l-fib*, *h-fib*, and *fhx* were not found, the long or structured isoforms
of *Sericin-1, Sericin-3*, and *Sericin-4* were identified in the reference transcriptome. The
longest isoform of *Sericin-3* (MSTRG.2595.1) possessed slightly more extensive
nucleotide sequences than that of KWMTBOMO08464, in which 73 bases of exon 3
were deleted, resulting in the prediction of incorrect ORF. The predicted amino acid
sequences of KWMTBOMO08464 were not similar to the sericin-3 amino acid
sequence in UniProtKB (ID: A8CEQ1), while that of MSTRG.2595.1 was similar.
Therefore, our transcriptome data provide more precise gene predictions. In the case
of *Sericin-4*, which was recently identified (Dong et al., 2019), we found a longer
transcript in our transcriptome data than the gene model reported by Dong et al.
(2019), which may contribute to the further characterization of sericin-4. The new
isoform of *Sericin-1* contains CDS that code glycine-, asparagine-, and threonine-rich
regions. It was not possible to elucidate the sequences of CDS because they were very
repetitive. Using long-read sequencers, repetitive sequences have been accurately
elucidated in the new reference genome. We consider our reference transcriptome
data to have significantly improved gene model data.
We searched for strongly expressed transcripts in one or more SG parts.
While more than 1,000 transcripts were strongly and ubiquitously expressed in the SG,
801 transcripts were strongly expressed in single parts of the SG. FEA with annotation

data on these transcripts in each part of the SG, except for the categories of MSG_A,
MSG_M specific plus MSG_A and MSG_M, was performed. The FEA results for
MSG_A, MSG_M specific plus MSG_A and MSG_M showed that these parts have
roles in coding or non-coding RNA processing. Some functional descriptions of these
ontologies are related to "splice variant processing". Some isoforms of *sericin-1* (IDs of
MSTRG.2477.1 - MSTRG.2477.16, and KWMTBOMO06216.mrna1) were strongly
expressed in MSG_A and MSG_M. Moreover, the FEA result contained "Asparagine
N-linked glycosylation". These results suggest that the splice variant processing of
*sericin-1* and asparagine processing of the sericin-1 protein, which possesses many
asparagine residues, occurred in MSG_A and MSG_M. The FEA results for ASG
suggested that ASG produced large amounts of energy via carbohydrate metabolic
processes. Silk proteins are mainly synthesized in PSG and MSG. After several
processes, matured silk protein, which is a large complex, is exported and released
through ASG (Takiya et al., 2016). Therefore, the strong expression of "carbohydrate
metabolic"-related transcripts may contribute to the export of silk protein. Since there is
moderate ontology for MSG_A and PSG, we cannot predict the roles of these parts.
In the present study, we performed RNA-seq on multiple tissues of *B. mori* and
constructed reference transcriptome data. The reference transcriptome data
constructed using RNA-seq data and new reference genome data contained
unidentified loci and isoforms, including a long and almost complete *sericin-1* isoform,
which are not present in the gene model data based on a reference new genome
(Kawamoto et al., 2019). Moreover, comprehensive transcriptome abundance and
annotation data will contribute to elucidating the functions of SG parts previously not
proven. The transcript data obtained herein will lead to advances in entomologic and
vertebrate research, including that on humans (Tabunoki et al., 2016).

# Methods

## Silkworm rearing, RNA extraction, and sequencing

The silkworm P50T (*daizo*) strain was reared on an artificial diet (Nihon Nosan
Kogyo, Yokohama, Japan) at 25°C under a 12-hour light/dark photoperiod. Tissues of
the SG, FB, MG, MT, TT, and OV were dissected on the 3$^{rd}$ day of fifth instar larvae.
The SG was further subdivided into ASG, MSG-A, MSG-M, MSG-P, and PSG. Each
tissue was dissected from one individual, except for OV, and three biological replicates
were obtained and analyzed separately. Tissues were homogenized using ISOGEN
(NIPPON GENE, Tokyo, Japan) and the SV Total RNA Isolation System (Promega,

Madison, WI) was used for RNA extraction. Extracted total RNA samples were

sequenced by Illumina Novaseq6000 (Macrogen Japan Corp., Kyoto, Japan).

<u>Construction of reference transcription data and estimation of the expression of</u>

<u>each transcript</u>

The raw RNA-seq data of 30 samples were trimmed by Trimmomatic-version

0.36 (Bolger et al., 2014). The trimmed RNA-seq data of each tissue were mapping to

the new reference genome with the new gene model (Kawamoto et al., 2019) by Hisat2

version 2.1.0. Each mapped data were assembled to transcriptome data by stringtie

version 1.3.3 (Pertea et al., 2016). The 30 transcriptome data sets were merged to one

transcriptome data set referred to as "a reference transcriptome" by the stringtie.

gffcompare v0.10.6 was used ([https://ccb.jhu.edu/software/stringtie/gffcompare.shtml](https://ccb.jhu.edu/software/stringtie/gffcompare.shtml))

for comparisons with the reference transcriptome and gene set of Kawamoto et al.

(2019).

To estimate the expression of the reference transcriptome in 30 samples, the

raw fastq data of each sample and reference transcript data were used with Kallisto

ver0.44.0 (Bray et al., 2016). In comparisons of transcriptome data, the raw RNA-seq

data of multiple tissues in *B. mori* strain o751 from the Sequence Read Archive (SRA)

and reference transcript data were used: the accession numbers of raw RNA-seq data

are DRA005094, DRA005878 and DRA005094 (Kikuchi et al., 2017; Ichino et al.,

2018; Kobayashi et al., 2019).

We used TIBCO Spotfire Desktop (v7.6.0) software with the "Better World"

program license (TIBCO, Inc., Palo Alto, CA; http://spotfire.tibco.com/better-world-

donation-program/) for the classification of differentially expressed samples in silkworm

tissues in hierarchical clustering using Ward's method.

<u>Annotation for the reference transcriptome and functional enrichment analysis</u>

Transcoder (v5.5.0) was used to identify coding regions within transcript

sequences and convert transcript sequences to amino acid sequences

([https://transdecoder.github.io/](https://transdecoder.github.io/)).

Transcriptome sequence sets were compared at the amino acid sequence level by the

successive execution of the blastp program in the NCBI BLAST software package

(v2.9.0+) with default parameters and an E-value cut-off of 1e-10 (Altschul et al.,1997).

Regarding the reference database sets to be blasted, human and fruit fly (*D.*

*melanogaster*) protein datasets in the Ensembl database (v.97) were used because the

sequences of these organisms were functionally well-annotated and amenable to

computational methods, such as a pathway analysis (Tabunoki et al., 2013). The

names of top-hit genes in the human and fruit fly datasets were annotated to *B. mori*

transcripts utilizing Ensembl Biomart (Kinsella et al., 2011) and Spotfire Desktop

software under TIBCO Spotfire's "Better World" program license (TIBCO Software, Inc.,

Palo Alto, CA, USA) (https://spotfire.tibco.com/better-world-donation-program/ ).

Functional enrichment analyses were performed using the metascape portal site

with annotation results against the human gene set (URL:

http://metascape.org/gp/index.html#/main/step1, Zhou et al., 2019).

<u>Investigation of gene structures of *sericin*, *fibroin*, and *fhx*</u>

In investigations on the *sericin*, *fibroin*, and *fhx* gene structures, we visualized

the positions of the new gene set and reference transcript data in the new reference

genome (Kawamoto et al., 2019) using the Integrative genomics viewer (IGV) (James

et al., 2011). In the gene model data set, *sericin-1* corresponded to

KWMTBOMO06216, *sericin-2* KWMTBOMO06334, *sericin-3* KWMTBOMO06311,

*sericin-4* KWMTBOMO06324-06326, fibroin heavy chain (*h-fib*) KWMTBOMO15365,

fibroin light chain *l-fib* KWMTBOMO08464, and *fhx* KWMTBOMO01001. The structures

of these models were compared visually with our new reference transcriptome data.

The several isoforms identified are listed in Table 1. We performed sequence

alignment using gene model sequence data and public sequences deposited in the

NCBI database.

## Data Availability

The RNA-seq reads supporting the conclusions of this study are available in the SRA

with accession number DRA008737 (The accession number of RNA-seq data of each

sample is shown in Table 2A).

Assembled transcriptome sequences are available at the Transcriptome Shotgun

Assembly (TSA) database under accession IDs ICPK01000001-ICPK01051926.

The estimated abundance of transcripts is available from the Gene Expression Archive

(GEA) in DDBJ under accession ID E-GEAD-315.

Supporting data are available in The Life Science Database Archive. The title in the

Archive is "KAIKO - Metadata of reference transcriptome data"

(DOI:10.18908/lsdba.nbdc02443-000.V001).

## References

Altschul, S.F. et al. "Gapped BLAST and PSI-BLAST: a new generation of protein

database search programs." Nucleic Acids Research 25:3389-3402 (1997). DOI:

10.1093/nar/25.17.3389

Bolger, A. M. et al. Trimmomatic: A flexible trimmer for Illumina Sequence Data.
Bioinformatics, btu170 (2014). DOI: 10.1093/bioinformatics/btu170

Bray, N. L. et al. Near-optimal probabilistic RNA-seq quantification. Nature
Biotechnology 34:525–527 (2016). DOI: 10.1038/nbt.3519

Daimon, T. et al. Knockout silkworms reveal a dispensable role for juvenile hormones
in holometabolous life cycle. Proceedings of the National Academy of Sciences of
the United States of America 112 E4226-E4235 (2015). DOI:
10.1073/pnas.1506645112

Dong, Z. et al. Identification of *Bombyx mori* sericin 4 protein as a new biological
adhesive. International Journal of Biological Macromolecules 132:1121-1130 (2019).
DOI: 10.1016/j.ijbiomac.2019.03.166

Garel, A. et al. Structure and organization of the *Bombyx mori* sericin 1 gene and of the
sericins 1 deduced from the sequence of the Ser 1B cDNA. Insect Biochemistry and
Molecular Biology 27:469–477 (1997). DOI: 10.1016/S0965-1748(97)00022-2

Ichino, F. et al. Construction of a simple evaluation system for the intestinal absorption
of an orally administered medicine using *Bombyx mori* larvae. Drug Discoveries and
Therapeutics 12:7-15 (2018). DOI: 10.5582/ddt.2018.01004

International Silkworm Genome Consortium. The genome of a lepidopteran model
insect, the silkworm Bombyx mori. 38(12):1036-45 (2008). DOI:
10.1016/j.ibmb.2008.11.004.

Ito, K. et al. Deletion of a gene encoding an amino acid transporter in the midgut
membrane causes resistance to a *Bombyx* parvo-like virus. Proceedings of the
National Academy of Sciences of the United States of America 105 7523-7527
(2008). DOI: 10.1073/pnas.0711841105

James, T. et al. Integrative genomics viewer. Nature Biotechnology 29:24–26 (2011).
DOI: 10.1038/nbt.1754

Kawamoto, M. et al. High-quality genome assembly of the silkworm, Bombyx mori.
Insect Biochemistry and Molecular Biology 107:53-62 (2019). DOI:
10.1016/j.ibmb.2019.02.002

Kikuchi, A. et al. Identification of functional enolase genes of the silkworm *Bombyx mori*
from public databases with a combination of dry and wet bench processes. BMC
Genomics 18: 83 (2017). DOI: 10.1186/s12864-016-3455-y

Kinsella, R.J. et al. Ensembl BioMarts: a hub for data retrieval across taxonomic space.
Database (Oxford) 2011:bar030 (2011). DOI: 10.1093/database/bar030

Kobayashi, Y. et al. Comparative analysis of seven types of superoxide dismutases for

their ability to respond to oxidative stress in *Bombyx mori*. Scientific Reports 9, 2170

(2019). DOI: 10.1038/s41598-018-38384-8

Kondo, Y. et al. Toll ligand Spätzle3 controls melanization in the stripe pattern

formation in caterpillars. Proceedings of the National Academy of Sciences of the

United States of America 114 8336-8341 (2017). DOI: 10.1073/pnas.1707896114

Mita, K. et al.   The genome sequence of silkworm, *Bombyx mori.* DNA Research

29;11(1):27-35 (2004). DOI:10.1093/dnares/11.1.27

Pertea, M. et al. Transcript-level expression analysis of RNA-seq experiments with

HISAT, StringTie and Ballgown. Nature Protocols 11:1650-1667 (2016).

DOI:10.1038/nprot.2016.095

Sakurai, T. et al. A single sex pheromone receptor determines chemical response

specificity of sexual behavior in the silkmoth *Bombyx mori*. PLoS Genetics

7:e1002115 (2011). DOI 10.1371/journal.pgen.1002115

Suetsugu, Y.   et al. Large scale full-Length cDNA sequencing reveals a unique

genomic landscape in a lepidopteran model insect, *Bombyx mori*. G3 3(9):1481-

1492 (2013). DOI:10.1534/g3.113.006239

Takiya, S. et al. Regulation of Silk Genes by Hox and Homeodomain Proteins in the

Terminal Differentiated Silk Gland of the Silkworm *Bombyx mori*. Journal of

Developmental Biology 4(2):19 (2016). DOI:10.3390/jdb4020019

Tabunoki, H. et al. Can the silkworm (Bombyx mori) be used as a human disease

model? Drug Discoveries and Therapeutics 10:3-8 (2016).

DOI:10.5582/ddt.2016.01011

Takasu, Y. et al. The silk sericin component with low crystallinity. Sanshi-Konchu

Biotec 75:133–139 (2006) DOI: 10.11416/konchubiotec.75.133

Takasu, Y. et al. Identification and characterization of a novel sericin gene expressed

in the anterior middle silk gland of the silkworm *Bombyx mori*. Insect Biochemistry

and Molecular Biology 37:1234–1240 (2007) DOI: 10.1016/j.ibmb.2007.07.009

Tamura, T. et al. Germline transformation of the silkworm *Bombyx mori* L. using a

piggyBac transposon-derived vector. Nature Biotechnology 18:81–84 (2000)

DOI:10.1038/71978

Tan, A. et al. Precocious metamorphosis in transgenic silkworms overexpressing

juvenile hormone esterase. Proceedings of the National Academy of Sciences of the

United States of America. 102:11751-11756. (2005) DOI:

10.1073/pnas.0500954102

Tatematsu, K. et al. Construction of a binary transgenic gene expression system for
recombinant protein production in the middle silk gland of the silkworm *Bombyx*
*mori*. Transgenic Research, 19:473-487 (2010). DOI:10.1038/71978

Xia, Q. et al. A draft sequence for the genome of the domesticated silkworm (*Bombyx*
*mori*). Science 10;306(5703):1937-40 (2004). DOI:10.1126/science.1102210

Xia, Q. et al. Microarray-based gene expression profiles in multiple tissues of the
domesticated silkworm, *Bombyx mori*. Genome Biology 8(8):R162 (2007).
DOI:10.1186/gb-2007-8-8-r162

Yamatomo, K. et al. A BAC-based integrated linkage map of the silkworm Bombyx
mori. Genome Biology. 9:R21 (2008). DOI:10.1186/gb-2008-9-1-r21

Yukuhiro, K. et al. Insect silks and cocoons: structural and molecular aspects.
Extracellular composite matrices in Arthropods (eds. E. Cohen & B. Moussian),
pp.515-555. Springer, Cham. (2016) DOI: 10.1007/978-3-319-40740-1_18

Zhou, C. et al. Fine organization of *Bombyx mori* fibroin heavy chain gene. Nucleic
Acids Research 28:2413–2419 (2000). DOI: 10.1093/nar/28.12.2413

Zhou, Y. et al. Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. Nature Communications 10(1):1523 (2019).
DOI:10.1038/s41467-019-09234-6

# 435   Funding

This work was supported by the National Bioscience Database Center of the Japan
Science and Technology Agency (JST) to HB.

This work was supported by the Cabinet Office, Government of Japan, Cross-
ministerial Strategic Innovation Promotion Program (SIP), "Technologies for Smart Bio-
industry and Agriculture" (funding agency: Bio-oriented Technology Research
Advancement Institution, NARO) to KY, TT, JS, and HS.

This work was supported by a grant from the Ministry of Agriculture, Forestry
and Fisheries of Japan (Research Project for Sericultural
Revolution) to KY, TT, and HS.

# 445   Acknowledgments

# 449   Author contributions

Conceived and designed the experiments: K.Y., T.T., and H.S.

Performed the experiments: T.T.

Contributed reagents/materials/analysis tools: H.S.

Analyzed the data: K.Y., J.S., A.J., and H.B.

Contributed to the writing of the manuscript under draft version: K.Y., T.T., and H.B.

All authors discussed the data and helped with manuscript preparation. K.Y.

supervised the project.

All authors read and approved the final manuscript.

## 458  Competing interests

459  The authors declare no conflicts of interest.

## 460  Figures

Fig. 1 Workflow of the data analysis performed in the present study. To obtain
reference transcriptome sequences, Fastq data of 10 tissues from $5^{th}$ instar larvae
were mapped to the new reference genome (Kawamoto et al., 2019). Kallisto software
was used to estimate the expression abundance of each transcript in these tissues
plus other *B. mori* samples of which RNA-seq data were obtained from a public
database (Accession numbers are listed in Table 2B). We performed a Blast search
against human and *Drosophila* genome data to perform functional annotations of the
reference transcriptome. Insect, human, database image, and sequencer drawings
(http://togotv.dbcls.jp/ja/pics.html) are licensed at
(http://creativecommons.org/licenses/by/4.0/deed.en).

Fig. 2 Basal characteristics of the reference transcriptome. (A) Comparison of gene

model data (Kawamoto et al., 2019) and the reference transcriptome data of the

present study. The number of loci and transcripts are shown. These numbers were

calculated from gff files of the two data sets. (B) Classification of 51,926 transcripts.

Each transcript was classified into three categories, and the numbers of the three

categories are shown in a pie chart. Definitions of the three categories were described

in the main text.

Fig. 3 Hierarchical clustering of expression data in 45 samples. Hierarchical clustering

was performed using transcriptome expression data (tpm values). Abbreviations of the

samples are shown and described in the main text. The numbers added to the

abbreviations mean biological replicates.

Fig. 4 Longest isoforms of *sericin-1* and *sericin-4* in the reference transcriptome.

(A) A schematic drawing showing the exon positions of isoform MTSRG.2477.1

(longest isoform of *sericin-1*) and a gene model of *sericin-1* (Kawamoto et al., 2019).

Squares indicate exons with exon numbers in the gff file (Supporting data 1). Exon 6 of

MSTRG.2477.1 corresponds to exon 5' and exons 7-10 correspond to exons 6-9 in

KWMTBOMO06216 (each group of exons are connected with dashed lines).

(B) Exon positions of isoform MTSRG.2477.1 (the longest isoform of *sericin-4*) and the

gene model of *sericin-4* (KWMTBOMO06325, KWMTBOMO06325, and

KWMTBOMO06326) in the new reference genome is shown by IGV. The scale above

the transcript indicates the location of chromosome 11.

Fig. 5 Strongly expressed transcripts within the silk gland. The numbers in the Venn
diagrams indicate the number of transcripts of which values of tpm were more than 30
in the corresponding silk gland parts.

Fig. 6 Results of the enrichment analysis by Metascape. Using annotation data against
the human gene set of the reference transcripts expressed in one or several parts of
the silk gland (Fig. 5), an enrichment analysis was performed (numbers in brackets
after the silk gland parts indicate the numbers of transcripts). -log10 (P) represents -
log10 (P-value). For example, -log10 (P)=5 represents P-value= $10^{-5}$ (A) Transcripts
showing tpm > 30 in MGM_M (99),MGM_P (71), and MGM_M and MGM_P (345).

(B) Transcripts showing tpm > 30 in MSG_A (180).

(C) Transcripts showing tpm > 30 in ASG (351).

(D) Transcripts showing tpm > 30 in PSG (100).

Table

Table 1

*Sericin, fibroin*, and *fibrohexamerin* genes and isoform IDs

| Gene name | Gene model ID in Kawamoto et al., 2019 | Isoform IDs in Supporting data 1 (GenBank IDs) |
|---|---|---|
| *sericin-1* | KWMTBOMO06216 | MSTRG.2477.1 - MSTRG.2477.16, KWMTBOMO06216.mrna1 (ICPK01006046 -ICPK01006062) |
| *sericin-2* | KWMTBOMO06334 | MSTRG.2627.1, MSTRG.2627.2, KWMTBOMO06334.mrna1 (ICPK01006484 -ICPK01006486) |
| *sericin-3* | KWMTBOMO06311 | MSTRG.2595.1 - MSTRG.2595.9, KWMTBOMO06311.mrna1 (ICPK01006421 -ICPK01006430) |
| *sericin-4* | KWMTBOMO06324-06326 | MSTRG.2610.1 (ICPK01006453) |
| fibroin heavy chain (*h-fib*) | KWMTBOMO15365 | MSTRG.14927.1 - MSTRG.14927.23, KWMTBOMO15365.mrna1 (ICPK01035046 -ICPK01035068) |

| fibroin light chain (*l-fib*) | KWMTBOMO08464 | MSTRG.5511.1, KWMTBOMO08464.mrna1 (ICPK01013031 -ICPK01013032) |
|---|---|---|
| *fibrohexamerin* | KWMTBOMO01001 | |

520    Table 2

521    A. Samples for RNA-seq and run accession IDs

| Sample | SRA Run ID | Sex |
|---|---|---|
| Anterior silk gland (ASG) | DRR186474, DRR186475, DRR186476 | Male |
| Anterior part of the middle silk gland (MSG_A) | DRR186477, DRR186478, DRR186479 | Male |
| Middle part of the middle silk gland (MSG_M) | DRR186480, DRR186481, DRR186482 | Male |
| Posterior part of the middle silk gland (MSG_P) | DRR186483, DRR186484, DRR186485 | Male |
| Posterior silk gland (PSG) | DRR186486, DRR186487, DRR186488 | Male |
| Fat body (FB) | DRR186489, DRR186490, DRR186491 | Male |
| Midgut (MG) | DRR186492, DRR186493, DRR186494 | Male |
| Malpighian tubules (MT) | DRR186495, DRR186496, DRR186497 | Male |
| Testis (TT) | DRR186498, DRR186499, DRR186500 | Male |

| | | |
|---|---|---|
| Ovary (OV) | DRR186501, DRR186502, DRR186503 | Female |

523 B. RNA-seq data from SRA

| Sample | SRA Run ID | Reference |
|---|---|---|
| Testis | DRR068893, DRR068894, DRR068895 | Kikuchi et al. 2017 |
| Fat body | DRR095105, DRR095106, DRR095107 | Kobayashi et al. 2019 |
| Midgut | DRR095108, DRR095109, DRR095110 | Ichino et al. 2018 |
| Malpighian tubules | DRR095111, DRR095112, DRR095113 | Kobayashi et al. 2019 |
| Silk gland | DRR095114, DRR095115, DRR095116 | Kobayashi et al. 2019 |

# 525 Supporting data

All supporting data are available in The Life Science Database Archive

(https://dbarchive.biosciencedbc.jp/index-e.html).

Supporting data 1

Metadata of reference transcriptome data

URL:https://togodb.biosciencedbc.jp/db/kaiko_trascnript_data

DOI:10.18908/lsdba.nbdc02443-001.V001

Supporting data 2

Predicted amino acid sequences of the reference transcriptome

DOI:10.18908/lsdba.nbdc02443-004.V001

Supporting data 3

Annotations of each transcript (blast against human and *Drosophila* gene sets)

URL:https://togodb.biosciencedbc.jp/db/kaiko_annotation_human_drosophila_data

DOI:10.18908/lsdba.nbdc02443-003.V001

Supporting data 4

Expression data of each transcript in multiple tissues

URL:https://togodb.biosciencedbc.jp/db/kaiko_transcript_tpm_data

DOI:10.18908/lsdba.nbdc02443-002.V001

# Supplemental Figure

Supplemental Fig. 1

Predicted amino acid sequences of the longest *sericin1* isoforms (MSTRG.2477.1.p1

).  Glycine, asparagine and threonine residues are colored in red. The region of exon7

is underlined.

DOI:10.6084/m9.figshare.998056