Reference transcriptome data in silkworm

Bombyx mori

Kakeru Yokoi[1,2,*,**], Takuya Tsubota[3,*], Akiya Jouraku[1], Hideki Sezutsu[3], Hidemasa Bono[4,5]

1 Insect Genome Research and Engineering Unit, Division of Applied Genetics, Institute of Agrobiological Sciences (NIAS), National Agriculture and Food Research Organization (NARO), 1-2 Owashi, Tsukuba, Ibaraki 305-8634, Japan

2 Research Center for Agricultural Information Technology (RCAIT), National Agriculture and Food Research Organization (NARO), Kintetsu Kasumigaseki Building Kasumigaseki 3-5-1 Chiyoda-ku, Tokyo 100-0013, Japan

3 Transgenic Silkworm Research Unit, Division of Biotechnology, Institute of Agrobiological Sciences (NIAS), National Agriculture and Food Research Organization (NARO), 1-2 Owashi, Tsukuba, Ibaraki 305-8634, Japan

4 Database Center for Life Science (DBCLS), Joint Support-Center for Data Science Research, Research Organization of Information and Systems, 1111 Yata, Mishima, Shizuoka 411-8540, Japan

5 Program of Biomedical Science, Graduate School of Integrated Sciences for Life, Hiroshima University, 3-10-23 Kagamiyama, Higashi-Hiroshima city, Hiroshima, 739-0046, Japan

*These authors equally contributed to this work.

**Corresponding author: Kakeru Yokoi

E-mail addresses

Kakeru Yokoi: yokoi123@affrc.go.jp

Takuya Tsubota: tsubota@affrc.go.jp

Akiya Jouraku: joraku@affrc.go.jp

Hideki Sezutsu: hsezutsu@affrc.go.jp

Hidemasa Bono: bonohu@hiroshima-u.ac.jp

32   Abstract

Background:

The silkworm *Bombyx mori* is a lepidopteran model insect with biological and industrial

importance. Its high-quality genome sequence has recently become available and the

utilization of this information in combination with extensive transcriptomic analyses is

expected to provide an elaborate gene model. It will also be possible to clarify gene

expression in detail using this approach.

Results:

We herein performed RNA-seq analysis of ten major tissues/subparts of silkworm

larvae. Sequences were mapped onto the reference genome assembly and reference

transcriptome data was successfully constructed. The reference data provided a nearly

complete sequence for *sericin-1*, a major silk gene with a complex structure. We also

markedly improved the gene model for other genes. Transcriptomic expression was

investigated in each tissue and a number of transcripts were identified that were

exclusively expressed in tissues such as the testis. Transcripts strongly expressed in

the midgut formed tight genomic clusters, suggesting that they originated from tandem

gene duplication. Transcriptional factor genes expressed in specific tissues or the silk

gland subparts were also identified.

Conclusions:

We successfully constructed reference transcriptome data in the silkworm and found

that a number of transcripts showed unique expression profiles. These results will

facilitate basic studies on the silkworm and accelerate its applications, which will

contribute to further advances in lepidopteran and entomological research and the

practical use of these insects.

Keywords: silkworm, *Bombyx mori*, transcriptome analysis, RNA-seq, gene model,

transcriptional factor

Background

The silkworm *Bombyx mori* is a lepidopteran insect that has been utilized in

studies of physiology, genetics, molecular biology, and pathology. Functional analyses

of genes related to hormone synthesis/degradation, pheromone reception, larval

marking formation, and virus resistance have been performed using this silkworm [1-5],

and the findings obtained have contributed to the promotion of insect science. The

silkworm has the ability to produce large amounts of silk proteins, which is one of the

most prominent characteristics of this species. Silk proteins are mainly composed of

the fibrous protein Fibroin and aqueous protein Sericin, which are produced in the

larval tissue silk gland (SG) [6]. A transgenic technique has been applied to the

silkworm [7], which has enabled the production of a large amount of recombinant

proteins through the introduction of transgenes for overexpression in the SG [8]. The

silkworm can be utilized as a significant bioreactor by this approach.

Based on its biological and industrial importance, the whole genome sequence

of the silkworm was reported in 2004 by two research groups [9, 10]. This was the first

lepidopteran genomic analysis and has served as a fundamental basis for genomic

studies on Lepidoptera. This silkworm genome data was updated in 2008 [11] and

related data have since become available, including microarray-based gene expression

profiles, a BAC-based linkage map, and full-length cDNA data [12-14]. These data

have strongly promoted studies on *B. mori* and other lepidopteran insects in the past

few decades.

A new and high-quality reference genome assembly of the silkworm p50T

(*daizo*) strain using PacBio long-read and Illumina short-read sequencers was reported

in 2019 [15]. The new genome assembly consists of 696 scaffolds with N50 of 16.8 Mb

and only 30 gaps. Predicted new gene model based on this novel genome assembly,

using cDNA, protein and RNA-seq data as hints, was constructed and was more

precise that the previous model made via the old genome assembly [15]. The next step

in the establishment of genome-related data is transcriptome data, which contains

reference transcriptome sequence data as well as gene expression profiles in major

tissues. These data will significantly contribute to advances in research on the silkworm

and other lepidopterans.

In the present study, we constructed a reference transcriptome dataset using

RNA-seq data obtained from ten major tissues/subparts of silkworm larvae (Fig. 1).

RNA-seq data were mapped on the new genome assembly and reference

transcriptome sequence data were successfully constructed. We also performed

functional annotation of the reference transcriptome using human and *Drosophila*

protein datasets in addition to NCBI-nr data. Established reference transcriptome

sequence data provided a nearly complete structure for *sericin-1* (*ser1*), a major silk

gene with a complex sequence. The expression of the transcriptome was investigated

in each tissue, and the expression of a number of transcripts was found to be confined

to tissues such as testis (TT). Among them, genes with transcripts that were strongly

expressed in the midgut (MG) formed tight genomic clusters, suggesting that they

originated via gene duplication. The transcripts of transcriptional factor (TF) genes

expressed in specific tissues or SG subparts were also detected, and we speculate

that these genes play key roles in the major biological process of these

tissues/territories. The present results will accelerate molecular biological studies on

the silkworm and other related species, and this is an essential milestone to promote

entomological research as well as the practical use of insects.

Results

Reference transcriptome data

We performed a transcriptomic analysis of major silkworm larval tissues,

namely, the SG, fat body (FB), MG, Malpighian tubules (MT), TT, and ovary (OV), to

acquire more expanded RNA-seq data (Fig. 2; Table 1). Gene expression was clearly

differentiated among subregions in the SG [16, 17], and, thus, we subdivided the SG

into five subparts and investigated gene expression in each region (the anterior silk

gland (ASG), anterior part of the middle silk gland (MSG_A), middle part of the middle

silk gland (MSG_M), posterior part of the middle silk gland (MSG_P) and posterior silk

gland (PSG)) (Fig. 2; Table 1). Totally ten tissues/subparts were dissected from the fifth

instar third day larvae of p50T strain with three biological replicates (Table 1). Thirty

sets of RNAs were used for RNA-seq. We mapped RNA-seq data on the reference

genome assembly using information from the previously established gene model [15]

(hereafter referred to as "gene model data (GMD)") and constructed reference

transcriptome data (RTD). RTD comprises 51,926 transcripts in 24,236 loci (Fig. 3A),

the numbers of which are higher than those of GMD (24,236 vs 16,845 loci and 51,926

vs 16,880 transcripts; see Fig. 3A and 3B). Therefore, RTD is an extension of GMD. To

perform functional annotations, coding sequence (CDS) regions and amino acid

sequence data were constructed by using RTD (Additional file 1), and found that

39,619 transcripts, derived from 16,632 loci, had at least one CDS in RTD. The

predicted amino acid sequences were used for gene functional annotations by a

homology search against human and *Drosophila* gene sets. This analysis revealed that

26,698 genes showed homology to human genes and 29,177 to fruit fly genes

(Additional file 2). We also performed a blastp analysis using the NCBI nr database

and found that 43,358 amino acids had homological proteins in this database

(Additional file 3).

Comparison between constructed reference transcriptome data and previous gene

model data

RTD represents a marked improvement over GMD. Several misassembled genes

are present in GMD, as represented by KWMTBOMO00087-88, 00196-197, or 00222-

223 genes [15]. These genes are split into two structures even though full-length cDNA

data define them as single genes [14, 15]. We investigated their structures in our

model and found that all were accurately predicted (MSTRG.494.1, MSTRG.649.1-2,

and MSTRG.704.1-3) (Additional file 4: Fig. S1). The elucidated gene structure was

attributed to the extensive RNA-seq analysis in the present study; the previous study

lacked gene expression data from tissues such as TT, OV, MT, and PSG, and these

genes were all strongly expressed in these tissues (Additional file 5: Table S1).

RTD also provided a number of novel genes/isoforms. Comparisons of RTD with

GMD revealed that among the 51,926 transcripts identified, 7,704 belonged to the new

loci group, whereas 27,342 were categorized as new isoforms (Fig. 3B, see Methods).

Among the 7,704 new loci group transcripts, a number of genes were also present in

the previously established full-length cDNA-based gene model [14]. However, 2,324

transcripts did not hit this gene set and, thus, these genes to which these transcripts

belong were perceived to be novel genes. An expression analysis revealed that many

of these genes were commonly expressed in all of the tissues investigated herein,

whereas other genes were exclusively expressed in specific tissues, such as TT

(Additional file 6: Fig. S2). The functional annotation analysis revealed that newly

identified genes included a trypsin inhibitor (MSTRG.14562.2), carboxypeptidase

(MSTRG.16874.1-3), and pyruvate kinase (MSTRG.18651.2). Therefore, our RTD

represents a significant improvement over GMD (Additional file 3).

<u>Elaborated structures of silk genes</u>

Silk production is one of the most prominent characteristics of the silkworm. Silk

genes are strongly expressed in the silk-producing tissue SG and our extensive RNA-

seq data are expected to show highly elaborated models for these genes. *ser1* is one

of the major silk genes, is strongly expressed in the MSG, and encodes a >400-kDa

serine-rich protein [6, 18]. *ser1* is composed of 9 exons, among which exon 6 has a

long repetitive sequence with a length of ~6,500 bp [6 ,19]. The full-length sequence of

this exon has yet to be elucidated due to its complexity. We herein demonstrated that

our model MSTRG.2477.1 provided an almost complete sequence for this exon (6,234

bp, Additional file 7: Fig. S3). Exon 6 encodes serine, glycine, threonine, asparagine,

and aspartic acid-rich residues (Additional file 8: Fig. S4), which is consistent with

previous findings showing that Ser1 comprises large numbers of these residues

(Additional file 9: Table S2) [20]. The detailed structural analysis revealed that the long

repetitive motif identified here comprised 53 repeats of a 38-amino acid unit (Additional

file 10: Fig. S5). Each unit had serine-rich residues and a slight difference was

observed in sequences among units (Additional file 10: Fig. S5) [6]. The 38-amino acid-

based repeat unit was also observed in exon 8 of *ser1* (Additional file 8: Fig. S4;

Additional file 10: Fig. S5) or in the sericins of saturniid species [6, 21], and, thus, the

repeat unit of this length is expected to have a structural function in a number of sericin

proteins.

We also observed significant improvements for other *sericin* genes. Sericin-3

(Ser3) is another major silk protein that has a relatively soft texture and possesses

serine-rich residues [22, 23]. In GMD, a 73-bp deletion was detected in exon 3, and

due to this structural error, a frame shift was present in the predicted amino acid

sequence (KWMTBOMO06311; Additional file 11: Fig. S6). In contrast, our RTD

(MSTRG.2595.1) successfully provided an accurate gene structure (Additional file 11:

Fig. S6). Sericin-4 (Ser4) is another sericin protein that is composed of 34 exons [24].

This gene is split into three distinct structures in GMD (KWMTBOMO06324,

KWMTBOMO06325, and KWMTBOMO06326), whereas RTD provided an exact model

(MSTRG.2610.1) (Additional file 12: Fig. S7). Collectively, these results suggest that

our RTD provided highly defined structures, even for complex silk genes.

<u>Estimating the abundance of the reference transcriptome in multiple tissues</u>

Our extensive transcriptomic analysis provided fundamental insights into gene

expression profiles in multiple silkworm tissues. The expression abundance of each

transcript was calculated as transcripts per million (tpm) (Additional file 13) and overall

gene expression was compared among tissues through two independent methods, one

for hierarchical clustering (HC) and another for a principal component analysis (PCA).

To avoid the effects of low expression transcripts, we performed these analyses using

transcripts with tpm values > 30 in at least one sample. HC using all transcript tpm data

was also performed for comparisons. These analyses revealed that the biological

replicate samples collected herein were highly reproducible because they were derived

from the same tissues forming tight clusters (Fig. 4; Additional file 14: Fig. S8). In the

HC analysis, a single cluster was formed for the MSG_M and MSG_P in each sample

(Fig. 4; Additional file 14: Fig. S8B), and we speculate that this was due to highly

conserved gene expression between these SG territories. A correlation analysis of

gene expression using SG transcriptome data supported this hypothesis (Additional file

15: Table S3). A previous study performed RNA-seq analysis of multiple silkworm

larval tissues in another silkworm strain, o751 [25-27], and these RNA-seq data were

added to our analysis (Fig. 4; Table 1; Additional file 14: Fig. S8; these samples are

referred to as BN_MG, BN_FB, BN_MT, BN_SG, and BN_TT). We found that the

samples collected from the same tissues clearly formed clusters (Fig. 4). In the o751

strain, the SG was collected as a whole tissue. Although all the SG subparts (MSG_A,

MSG_M, MSG_P, ASG and PSG) were closely located in the HC analysis of

transcripts with > 30 (tpm) in at least one sample, BN_SG formed a single cluster

distant from those of the SG subparts (Fig. 4). Therefore, our transcriptomic data is a

robust platform for analyzing and comparing gene expression in multiple tissues.

<u>Transcript abundance in each tissue and in the silk gland</u>

Using the data described above, we investigated gene expression in detail in

each tissue. We focused on genes with a tpm value > 30, which approximately

accounted for the top 5% of the most strongly expressed genes (data not shown), and

regarded such transcripts as being expressed in each tissue. We found that 711 genes

were expressed in all tissues (Fig. 5), suggesting ubiquitous functions. We also

detected genes expressed in specific tissues (Fig. 5, Additional file 16: Table S4).

Among them, genes solely expressed in TT were the most abundant (1882), followed

by those expressed in OV (799), MT (499), and MG (440) (Fig. 5, Additional file 16:

Table S4). We also identified genes expressed in more than two tissues, such as TT

and OV (397, Fig. 5). A functional enrichment analysis (FEA) was performed on genes

with tissue-restricted expression, and functional clusters enriched in each tissue were

very diversified; for example, genes exclusively expressed in the TT were strongly

enriched for "cilium organization", "Huntington's disease", and "cilium or flagellum-

dependent cell motility" (Fig. 6A) whereas "Metabolism of RNA", "regulation of mRNA

metabolic process", and "ribonucleoprotein complex biogenesis" were enriched in the

OV (Fig. 6B). Comparisons of gene expression levels among tissues revealed that the

expression levels of strongly expressed genes were very high in the ASG (Fig. 7).

These genes comprise fungal protease inhibitors, cuticular protein genes and others

(Additional file 17: Table S5). In contrast, the levels of strongly expressed genes were

lower in the MSG-M/MSG-P (Fig. 7; Additional file 17: Table S5). We also investigated

the expression profiles of genes strongly expressed in each tissue and found tissue-

restricted expression for these genes in the MG, FB, and MSG_A and ubiquitous

expression for those in the other tissues examined (Additional file 17: Table S5;

Additional file 18: Fig. S9). The genomic positions of tissue-enriched genes were

examined, and genes strongly expressed in the MG formed tight genomic clusters (Fig.

8; Additional file 17: Table S5). We also found clusters for strongly expressed genes in

other tissues (Additional file 17: Table S5; Additional file 19: Fig. S10).

We then investigated gene expression in the SG in more detail. Previous

studies revealed that a number of genes showed territory-specific expression in the SG

[16, 17]. However, overall gene expression in each territory remains unclear. We

herein demonstrated that >1000 genes were commonly expressed in all SG subparts

and also that a number of genes were expressed in specific territories (Fig. 9). They

included 351 ASG-restricted, 180 MSG_A-restricted, 99 MSG_M-restricted, 71

MSG_P-restricted, and 100 PSG-restricted genes, respectively (Fig. 9, Additional file

20: Table S6). Furthermore, we identified genes that were commonly expressed in

more than two territories (Fig. 9). They included genes expressed in the MSG_M and

MSG_P (Fig. 9), and combined with the results of HC and the correlation analysis (Fig.

4; Additional file 15: Table S3), we speculate that gene expression is highly conserved

between the MSG_M and MSG_P. This result was supported by the presence of a

smaller number of genes expressed solely in the MSG_M or MSG_P (Fig. 9). We also

found that the genes that were exclusively expressed in the ASG were more abundant

than in other territories, and fewer genes were commonly expressed in the ASG and

other subparts (Fig. 9). We speculate that this reflects the functional diversification of

the ASG because the numbers of genes exclusively expressed in ASG (351) and those

expressed in the MSG and PSG (395) were comparable, suggesting similar diversified

functions (Fig. 9). This may also be the case for the MSG_A based on the presence of

similar characteristics (Fig. 9). FEA revealed that the functional clusters enriched in

each SG subpart were largely diversified (Additional file 21: Fig. S11).

Expression analysis of transcriptional factor genes in the silkworm

A transcriptomic analysis is a powerful tool for identifying genes with low levels

of expression. TF genes are considered to show low expression levels even though

they have many important functions in developmental, physiological, and other major

biological processes. Therefore, an expression analysis of TF genes will contribute to a

more detailed understanding of the silkworm biology. Silkworm TF genes have recently

been catalogued [28] and we investigated their expression levels in various silkworm

tissues using this information. According to the low level of expression of TF genes, we

herein perceived TF genes with a tpm value > 5 as those expressed in each tissue.

This analysis revealed that a number of TF genes were exclusively expressed in the

OV and/or TT (Fig. 10A; Additional file 22: Table S7). These genes included

KWMTBOMO02002 (*traffic jam*), KWMTBOMO002212 (*mirror*), KWMTBOMO01693

(*vismay*), and KWMTBOMO06584 (*Sox100B*), all of which play significant roles in

gonad morphogenesis, oogenesis, spermatogenesis, and TT differentiation in

Drosophila melanogaster (Additional file 22: Table S7) [29-32]. We also found that

KWMTBOMO09369/KWMTBOMO10218, the silkworm counterparts of human GATA4,

were expressed in the OV, similar to that in humans (Additional file 22: Table S7) [33].

We speculate that these TF genes have conserved functions in a wide variety of

organisms, possibly in the development, differentiation, or homeostasis of reproductive

tissues.

We also investigated TF gene expression in the SG. Previous studies revealed

the central roles of the TF genes *forkhead* (*fkh*), *Antennapedia* (*Antp*), and *Arrowhead*

(*Awh*) in the regulation of silk gene expression [34-38]. Other TF genes that function in

the regulation of silk gene expression may also exist because silk genes are strongly

expressed and a number of silk genes are expressed in specific territories in the SG

(Fig. 9). In addition to previously identified TF genes, the expression of novel TF

genes, such as those belonging to the High Mobility Group (HMG), Zinc Finger (ZF),

Paired box (PAX), or forkhead family, was confined to the SG (Fig. 10B, Additional file

23: Table S8). The tpm values of these genes are shown in Additional file 23: Table

S8, and we identified TF genes expressed solely in the ASG or MSG_A. Genes

expressed in more than two subparts were also detected (Additional file 23: Table S8).

We confirmed their expression in an RT-PCR analysis and experimental results were

consistent with tpm values (Additional file 24: Fig. S12). The TF genes identified herein

include those with indispensable roles in the development of *D. melanogaster*; for

example, *Dichaete* (*D*), *gooseberry-neuro* (*gsb-n*), and *sloppy paired 2* (*slp2*) are

essential for embryonic segmentation [39-41], *Enhancer of split mβ-HLH* (*E*(*spl*)*mβ-*

HLH) for neurogenesis [42], and *twin of eyegone* (*toe*) for eye development [43]. We

consider that it is very interesting if these genes have significant roles in the regulation

of the silk gene expression, a function that is independent from the developmental

regulation, in the silkworm.

Discussion

In the present study, we performed RNA-seq analysis of multiple larval tissues

from the silkworm *B. mori*. We established RTD using a recently reported high-quality

reference genome assembly [15] and RNA-seq data newly obtained herein. RTD

showed marked improvements over GMD, most notably the establishment of a nearly

complete structure for *ser1* (Additional file 7: Fig. S3; Additional file 8: Fig. S4); its full

sequence has never been entirely elucidated due to its complexity. Our results indicate

that an extensive RNA-seq analysis in combination with high-quality reference genome

data provides highly refined gene structures, even for complex genes. The cost of

performing a deep sequencing analysis has recently decreased and, thus, it has

become affordable for every researcher to conduct not only a short-read RNA-seq

analysis, but also long-read genome sequencing using their own species. The present

results are a significant proof-of-concept that highly refined gene structures may be

established using the combination of these data, even for non-model organisms.

Furthermore, more elaborate gene structures may be constructed using the RNA-seq

data derived from other tissues and/or stages.

We herein found that a number of genes showed tissue-restricted expression in

the silkworm (Fig. 5). Among them, genes exclusively expressed in the TT were the

most abundant (Fig. 5). A previous study identified a number of TT-specific genes in

the silkworm [12], which is consistent with the present results. The presence of a

number of TT-specific genes was also demonstrated in the jewel wasp, *Nasonia*

vitripennis [44], indicating that this is a common feature in insects. Recent studies on

Drosophila revealed that newly emerging genes were strongly biased for expression in

the male reproductive system [45]. Therefore, the TT-specific genes identified in the

present study may have similar traits. This issue may be confirmed by investigating the

evolutionary ages of these genes, and, if this is the case, addressing the question of

why the TT is a tissue that is permissive for new gene birth, a phenomenon observed

not only in insects, but also in vertebrates [46], which will become possible using the

silkworm. Another important result obtained from our cross-tissue gene expression

analysis is that genes strongly expressed in the MG showed strong tissue-restricted

expression and also formed tight genomic clusters (Fig. 8; Additional file 17: Table S5;

Additional file 18: Fig. S9). Comparisons of the sequences of these genes revealed

that genes in each cluster encoded homological proteins; chr.7 genes encoded

trypsins, chr.15 juvenile hormone-binding proteins (JHBPs), chr.16 fatty acid-binding

proteins, chr.19 actin cytoskeleton-regulatory complex proteins, and chr.20 multiprotein

bridging factor 2 (MBF2) (Additional file 17: Table S5). Among these genes, strong

expression in the MG has already been demonstrated for *jhbp*s [47], and Trypsin

proteins were present in digestive juices [48]. Homological genes that cluster in the

genome are generally considered to have originated via tandem gene duplication, and

to the best of our knowledge, few studies have investigated clustered genes that are

expressed in the MG. In one case study on *Drosophila*, neutral lipase genes expressed

in the MG clustered in the genome and were presumably under positive selection to

retain different substrate specificities towards new lipid components of the diet [49].

Based on these findings, the genes identified in the present study may also have

advantages in the silkworm MG, such as enhancing activity for digestion and/or

xenobiotic detoxification. The present study provides valuable insights into gene

evolution as well as neofunctionalization in insects, which may be validated in more

detail in future studies. In addition, the results obtained herein will facilitate the practical

application of the silkworm; targeted gene integration into the clusters identified in the

present study will enable strong gene expression in the MG, which will contribute to the

establishment of strains with valuable properties, such as increased antiviral or

antibiotic activities. The latest genome editing technologies should promote the

establishment of useful silkworm strains.

Our detailed gene expression analysis provides fundamental information on the

traits of the SG, particularly each subpart. The SG is a tissue that arises from a single

embryonic segment and has a long tubular structure [50]. Gene expression in each SG

subpart is largely diversified, as demonstrated herein (Fig. 9; Additional file 20: Table

S6) and in previous studies [16, 17]. The most important result of the present study is

that overall gene expression in the ASG was diversified the most within the SG, as

demonstrated by the presence of a number of genes exclusively expressed in the ASG

(351, Fig. 9), a number of genes expressed both in the MSG and PSG (395, Fig. 9),

and the location of the ASG at the outermost site in the SG cluster in the HC analysis

(Fig. 4). We speculate that these results indicate the functional diversification of the

ASG, which is consistent with previous findings showing that the ASG functions in silk

fiber processing and the MSG/PSG in silk protein production [51]. FEA revealed that

functional clusters enriched in the ASG were "carbohydrate metabolic process" and

"Transport of small molecules" (Additional file 21: Fig. S11A), while those enriched in

the MSG/PSG were "ribonucleoprotein complex biogenesis" and "Translation"

(Additional file 21: Fig. S11F), further supporting this concept. In this context, another

result showing that gene expression in the MSG_A also showed diversification is of

interest. We demonstrated that genes specifically expressed in the MSG_A were

abundant (180, Fig. 9), a number of genes were commonly expressed in the

MSG_M/MSG_P/PSG (335, Fig. 9), and the MSG_A was located at the outermost site

in the MSG/PSG cluster in the HC analysis (Fig. 4). MSG_A is a part of the MSG and

functions in the production of the sericin proteins Ser2 and Ser3 [23, 52], similar to the

function of the MSG_M/MSG_P in the production of Ser1 [20]. Nevertheless, our

results indicate that gene expression in the MSG_A was more diverse than that in the

MSG_M/MSG_P compared with that of the PSG; the PSG is the territory that produces

fibroin and not sericin [53], whereas gene expression appeared to be more conserved

between the PSG and MSG_M/MSG_P than between the MSG_A and

MSG_M/MSG_P (Figs. 4 and 9). These results may be attributed to the presence of a

number of genes that are strongly and specifically expressed in the MSG_A, including

ecdysone oxidase, fatty acid hydroperoxide dehydratase, and other genes (Fig. 9;

Additional file 20: Table S6). We found that genes strongly expressed in the MSG_A

were enriched for the functional clusters of "Metabolism of vitamins and cofactors" and

"organic hydroxy compound transport" (Additional file 21: Fig. S11B), and speculated

that these clusters define the biological functions that are unique to this territory.

Therefore, our extensive RNA-seq analysis provides fundamental insights into the

functions of the SG as well as its evolution, which has not yet been elucidated in detail.

In the Lepidoptera, the morphology of the SG is largely diversified among species [54]

and the Saturniidae, a family that is phylogenetically close to the Bombycidae, have

MSGs with one territory and no morphological separation [6]. Therefore, the

differentiation of MSG gene expression observed herein may be specific to *B. mori*

and/or other closely related species. Further studies are needed to clarify whether

differences in gene expression among species are a driving force that generates

diversity in cocoon properties, including shape, size, and physical activity. The TF

genes identified in the present study may be one of the key factors responsible for the

differences in gene expression within the SG or among species.

In the present study, we established RTD for the silkworm and performed a

detailed examination of gene expression in silkworm tissues. The results obtained will

contribute to our understanding of silkworm biology and further promote the industrial

application of the silkworm and other insects.

416 Conclusions

417 We performed RNA-seq analysis of the major larval tissues of the silkworm and

418 established RTD. Using these data, we successfully improved the gene structure and

419 clarified gene expression in detail in each tissue. The present results are a

420 fundamental basis for the further promotion of silkworm research and will contribute to

421 the practical application of insects.

423 Methods

<u>Silkworm rearing, RNA extraction, and sequencing</u>

The silkworm p50T strain was reared on an artificial diet (Nihon Nosan Kogyo,

Yokohama, Japan) at 25°C under a 12-hour light/dark photoperiod. The SG, FB, MG,

MT, TT, and OV were dissected on the third day of the fifth instar larvae. The SG was

further subdivided into the ASG, MSG_A, MSG_M, MSG_P, and PSG. Each

tissue/subpart was dissected from one individual, and three biological replicates were

obtained and separately analyzed (see Table 1). Tissues were homogenized using

ISOGEN (NIPPON GENE, Tokyo, Japan) and the SV Total RNA Isolation System

(Promega, Madison, WI) was used for RNA extraction. The total RNA samples

extracted were sequenced by Illumina NovaSeq6000 (Macrogen Japan Corp., Kyoto,

Japan).

<u>Construction of RTD and estimation of the expression of each transcript</u>

The raw RNA-seq data of 30 samples were trimmed by Trimmomatic version

0.36 [55]. The trimmed RNA-seq data of each tissue were mapped to the new

reference genome with a new gene model [15] by HISAT2 version 2.1.0 [56]. Mapped

data were each assembled to transcriptome data by StringTie version 1.3.3 [57]. The

30 transcriptome datasets were merged into one transcriptome dataset, referred to as

"a reference transcriptome" by StringTie. GffCompare version 0.10.6 was used [58] for

comparisons with the reference transcriptome and previously reported gene sets [15].

The transcripts detected at the newly identified loci were categorized into the "New loci"

group, those newly detected at the previously identified loci into the "New isoform"

group, and other genes and transcripts into "Identified in gene models" (see Fig. 3B).

To estimate the expression of the reference transcriptome in 30 samples, the

raw fastq data of each sample and reference transcript data were used with Kallisto

version 0.44.0 [59]. The raw RNA-seq data of multiple tissues in *B. mori* strain o751

from the Sequence Read Archive (SRA) and reference transcript data were used in

comparisons of transcriptome data: the accession numbers of raw RNA-seq data are

DRA005094, DRA005878, and DRA005094 [25-27].

We used TIBCO Spotfire Desktop (version 7.6.0) software with the "Better

World" program license (TIBCO, Inc., Palo Alto, CA; [60]) for the classification of

differentially expressed samples in silkworm tissues in HC using Ward's method.

Morpheus was also used [61] for HC. R (version 3.6.0) was used in the PCA analysis

[62]. Regarding Venn diagram construction and the scatter plot analysis, R (version

4.0.2) was used [63]. The relationships among gene expression profiles in the SG

territories were evaluated using Spearman's rank correlation.

Annotation for the reference transcriptome and FEA

Transcoder (version 5.5.0) was used to identify coding regions within transcript

sequences and convert transcript sequences to amino acid sequences

[64]. Transcriptome sequence sets were compared at the predicted amino acid

sequence level by the successive execution of the blastp program in the NCBI BLAST

software package (version 2.9.0+) with default parameters and an e-value cut-off of 1e-

10 [65]. Regarding the reference database sets to be blasted, human and fruit fly (*D.*

melanogaster) protein datasets in the Ensembl database (v.97) were used because the

sequences of these organisms were functionally well-annotated and amenable to

computational methods, such as a pathway analysis [66]. The names of the top-hit

genes in the human and fruit fly datasets were annotated to *B. mori* transcripts utilizing

Ensembl Biomart [67] and Spotfire Desktop software under TIBCO Spotfire's "Better

World" program license (TIBCO Software, Inc., Palo Alto, CA, USA) [60]. Functional

enrichment analyses were performed using the metascape portal site [68]

with annotation results against the human gene set.

<u>Comparison of gene structures among different models</u>

Gene structures among RTD, GMD, and cDNA-based data were compared in the

silkbase [69] or KAIKObase [70]. Amino acid sequences were aligned using CLC

Genomics Workbench 20.0.04 (QIAGEN, Aarhus, Denmark).

<u>RT-PCR</u>

cDNA was synthesized using Superscript IV (Thermo Fisher Scientific Inc., Waltham,

MA, USA) according to the manufacturer's instructions. Five hundred nanograms of

total RNAs extracted from the ASG, MSG_A, MSG_M, MSG_P, and PSG were used

for the cDNA synthesis. KOD FX neo polymerase (Toyobo, Osaka, Japan) was used

for RT-PCR. PCR conditions were as follows: 95°C for 1 min followed by 22 cycles (for

rp49) or 30 cycles (for TF genes) of 95°C for 30 sec, 58°C for 30 sec, followed by 68°C

for 1 min, and additional 68 °C for 1 min after the cyclic phase. Primer sequences are

listed in Additional file 25: Table S9.

491   Declarations

Availability of data and materials

The RNA-seq reads supporting the conclusions of the present study are available in

the SRA with accession number DRA008737 (The accession number of RNA-seq data

of each sample is shown in Table 1).

Reference transcriptome data is available at the Transcriptome Shotgun Assembly

(TSA) database under accession IDs ICPK01000001-ICPK01051926, and the Gff file

of the transcriptome was available via DOI: 10.18908/lsdba.nbdc02443-001.V001.

The estimated abundance of transcripts is available from the Gene Expression Archive

(GEA) in DDBJ under accession ID E-GEAD-315 or from The Life Science Database

Archive (DOI: 10.18908/lsdba.nbdc02443-002.V001).

Additional data are available in The Life Science Database Archive (the title in the

Archive is "KAIKO - Metadata of reference transcriptome data"

(DOI:10.18908/lsdba.nbdc02443-000.V001)) and figshare (DOI:

10.6084/m9.figshare.c.5333894).

Additional file 1

Predicted amino acid sequences of reference transcriptome.

DOI:10.18908/lsdba.nbdc02443-004

Additional file 2

Functional annotations of the reference transcriptome (blast against human and

Drosophila gene sets). DOI:10.18908/lsdba.nbdc02443-003

Additional file 3

Functional annotations of the reference transcriptome (blast against the NCBI nr

database). DOI:10.6084/m9.figshare.14192741.

Additional file 4: Fig. S1

Comparison of gene structures among gene model data, cDNA-based data, and

reference transcriptomic data. (A) Locus around KWMTBOMO00087, (B)

KWMTBOMO00196, and (C) KWMTBOMO00222. GMD; gene model data, CBD;

cDNA-based data, RTD; reference transcriptomic data.

DOI:10.6084/m9.figshare.14205785

Additional file 5: Table S1

Tpm values in MSTRG.494.1, MSTRG.649.1-2 and MSTRG.704.1-3. DOI:

10.6084/m9.figshare.14206034

Additional file 6: Fig. S2

Expression of new loci transcripts that did not hit the cDNA-based gene model.

Numbers indicate transcripts with a tpm value >0.01 in each tissue. SG, silk gland

(using average tpm values in the five SG subparts); FB, fat body; MG, mid gut; MT,

Malpighian tubules; OV, ovary; TT, testis. DOI: 10.6084/m9.figshare.14217335

Additional file 7: Fig. S3

Structure of the *sericin-1* gene. MSTRG.2477.1 has a long exon 6.GMD; gene model

data, CBD; cDNA-based data, RTD; reference transcriptomic data. DOI:

10.6084/m9.figshare.14206166

Additional file 8: Fig. S4

The presumptive full-length amino acid sequence of sericin-1 deduced by reference

transcriptomic data (MSTRG.2477.1). The orange characters show the amino acids

encoded by exon 6 and blue characters by exon 8. DOI:

10.6084/m9.figshare.14206736

Additional file 9: Table S2

Comparisons of the sericin-1 amino acid composition elucidated by an amino acid

analysis and gene model. Mole% is shown. DOI: 10.6084/m9.figshare.14206748

<u>Additional file 10: Fig. S5</u>

Sequence of the 38-amino acid-based repeat unit encoded by exon 6 and exon 8 in

sericin-1. Exon 6 comprises 53 repeats and exon 8 13 repeats. DOI:

10.6084/m9.figshare.14212673

<u>Additional file 11: Fig. S6</u>

Comparison of the sericin-3 sequence among different gene models. The amino acid

sequences identified in the previous study (NM_001114644; [23]), derived from cDNA-

based data (BMgn014348), reference transcriptomic data (MSTRG.2595.1), and gene

model data (KWMTBOMO06311), are shown. Amino acids that differ in

KWMTBOMO06311 are shown as white letters. Note that a frame shift occurs in the

gene model data. DOI: 10.6084/m9.figshare.14216927

<u>Additional file 12: Fig. S7</u>

Comparison of the *sericin-4* gene structure. The gene structures elucidated by the

previous study (*sericin-4*; [24]) and modeled by gene model data (GMD) and reference

transcriptomic data (RTD) are shown. DOI: 10.6084/m9.figshare.14216951

<u>Additional file 13</u>

Expression data of each transcript in multiple tissues. DOI: 10.18908/lsdba.nbdc02443-

002.V001

<u>Additional file 14: Fig. S8</u>

(A) Principal Component Analysis (PCA) results with expression profiles in 45 samples

of transcripts showing a tpm value > 30 in at least one sample. Abbreviations and the

numbers of samples are the same as in Fig. 4. The X axis and Y axis are principal

components 1 (PC1) and PC2, respectively. (B) Hierarchical clustering of expression

data in 45 samples using all transcript tpm values. DOI:

10.6084/m9.figshare.14216993.

Additional file 15: Table S3

Spearman's rank correlation coefficient among silk gland territories. DOI:

10.6084/m9.figshare.14217071

Additional file 16: Table S4

List of genes expressed in specific tissues. DOI: 10.6084/m9.figshare.14217206

Additional file 17: Table S5

List of genes strongly expressed in each tissue or in silk gland subparts. The top 50

strongly expressed genes are shown. DOI: 10.6084/m9.figshare.14217218

Additional file 18: Fig. S9

A scatter plot of gene expression in each tissue. Each spot shows the tpm value. The

top ten ranked strongly expressed genes in (A) ASG, (B) MSG_A, (C) MSG_M, (D)

MSG_P, (E) PSG, (F) FB, (G) MG, (H) MT, (I) OV and (J) TT are marked in red. DOI:

10.6084/m9.figshare.14217227

Additional file 19: Fig. S10

Genomic position of genes strongly expressed in each tissue. In (A), genes strongly

expressed in the ASG only are shown with a red bar, in subparts other than the ASG

with a green bar, and those commonly expressed in the ASG and other tissues with a

yellow bar. This same applies to MSG_A (B), MSG_M (C), MSG_P (D), PSG (E), FB

(F), MT (G), OV (T), and TT (I). (A') shows the chr. 11 of the ASG and (F') shows chr.

20 of the FB. Regarding the MG, see Fig. 8. The top 50 strongly expressed genes are

shown. The black bar indicates the chromosome, and the number at the left side of

each chromosome indicates the chromosomal number. DOI:

10.6084/m9.figshare.14217233

Additional file 20: Table S6

List of territory-specific genes in the silk gland. DOI: 0.6084/m9.figshare.14217242

<u>Additional file 21: Fig. S11</u>

Results of the enrichment analysis for territory-specific transcripts in the silk gland. The

number of transcripts used for the analysis is shown in the bracket. -log10 (P)

represents -log10 (P-value). For example, -log10 (P)=5 represents P-value= 10^{-5}. DOI:

10.6084/m9.figshare.14217257

<u>Additional file 22: Table S7</u>

List of tissue-specific TF genes. DOI: 10.6084/m9.figshare.14217272

<u>Additional file 23: Table S8</u>

List of territory-specific TF genes in the silk gland. DOI: 10.6084/m9.figshare.14217296

<u>Additional file 24: Fig. S12</u>

RT-PCR of TF genes showing territory-specific expression in the SG. DOI:

10.6084/m9.figshare.14217299

<u>Additional file 25: Table S9</u>

Primer sequences used for RT-PCR. DOI: 10.6084/m9.figshare.14217308

Competing interests

The authors declare no conflicts of interest.

Funding

This work was supported by the National Bioscience Database Center of the Japan

Science and Technology Agency (JST) to HB.

This work was supported by the Cabinet Office, Government of Japan, Cross-

ministerial Strategic Innovation Promotion Program (SIP), "Technologies for Smart Bio-

industry and Agriculture" (funding agency: Bio-oriented Technology Research

Advancement Institution, NARO) to KY, TT, and HS.

This work was also supported by a grant from the Ministry of Agriculture, Forestry and

Fisheries of Japan (Research Project for Sericultural Revolution) to KY, TT, and HS.

Author contributions

Conceived and designed experiments: KY, TT, and HS.

Performed experiments: TT.

Contributed reagents/materials/analysis tools: HS.

Analyzed data: KY, TT, AJ, and HB.

Contributed to the writing of the manuscript under the draft version: KY, TT, and HB.

All authors discussed the data and helped with manuscript preparation. KY supervised

the project.

All authors read and approved the final manuscript.

Acknowledgments

The computing resource was partly provided by the super computer system at the

National Institute of Genetics (NIG), Research Organization of Information and

Systems (ROIS), Japan. The authors thank Mrs. Kaoru Nakamura and Toshihiko

Misawa for rearing the silkworms, Ms. Satoko Kawamoto for technical assistance, and

Dr. Jianqiang Sun for data analysis advice.

Legends

Fig. 1. Workflow of the data analysis performed in the present study. To obtain

reference transcriptome sequences, Fastq data of 10 tissues/subparts from 5th instar

larvae were mapped to the new reference genome [15]. Kallisto software was used to

estimate the expression abundance of each transcript in these tissues. Regarding

RNA-seq data obtained from the public database (o751 strain), see Table 1. We

performed a Blast search against human and *Drosophila* genome data to perform

functional annotations of the reference transcriptome. Insect, human, database

images, and sequencer drawings [71] are licensed at Creative Commons [72].

Fig. 2. (A) Final (5th) instar larva of the silkworm p50T (*daizo*) strain. Scale bar = 5 mm.

(B, C) Male (B) or (C) female individuals dissected on the third day of fifth instar larvae.

Scale bar = 5 mm. MG, mid gut; TT, testis; MT, Malpighian tubules; SG, silk gland; FB,

fat body; OV, ovary. (D, D') Image (D) and schematic (D') of the silk gland. ASG,

anterior silk gland; MSG_A, anterior part of the middle silk gland; MSG_M, middle part

of the middle silk gland; MSG_P, posterior part of the middle silk gland; PSG, posterior

silk gland. Scale bar = 2.5 mm.

Fig. 3. Basal characteristics of the reference transcriptome. (A) Comparison of gene

model data [15] and the reference transcriptome data of the present study. The

numbers of loci and transcripts are shown. These numbers were calculated from gff

files of the two data sets. (B) Classification of 51,926 transcripts. Each transcript was

classified into three categories, and the numbers of the three categories are shown in a

pie chart. Definitions of the three categories are described in the main text.

Fig. 4. Hierarchical clustering of expression data in 45 samples of transcripts showing a

tpm value > 30 in at least one sample. Abbreviations that start with "BN" indicate

samples collected in the previous study [25-27]. The numbers added to the

abbreviations mean biological replicates.

Fig. 5. Venn diagram showing transcripts expressed in each tissue. The number of

transcripts with a tpm value >30 is shown. SG, silk gland; FB, fat body; MG, mid gut;

MT, Malpighian tubules; OV, ovary; TT, testis.

Fig. 6. Results of the enrichment analysis by Metascape in testis (TT) (A) and ovary

(OV) (B). An enrichment analysis was performed using annotation data against the

human gene set of the reference transcripts expressed in specific tissues. -log10 (P)

represents -log10 (P-value). For example, -log10 (P)=5 represents P-value= 10^{-5}.

Fig. 7. A scatter plot of gene expression in each tissue. Each spot shows the tpm

value.

Fig. 8. Genomic position of genes strongly expressed in the MG only (red bar), in other

than the MG (green bar), and commonly in the MG and other tissues (yellow bar). The

top 50 strongly expressed genes are shown. The black bar indicates the chromosome.

The number at the left side of each chromosome indicates the chromosomal number.

(A) Genomic positions of all chromosomes. (B) Genomic positions of chromosomes 7,

15, 16, 19 and 20, in which the tight clusters of strongly expressed genes in the MG

are present.

Fig. 9. Venn diagram showing the number of transcripts with a tpm value >30 in each

silk gland part.

Fig. 10. Venn diagrams showing the number of TF genes with a tpm value >5 in each

tissue (A) and in silk gland subparts (B).

References

1.   Tan A, Tanaka H, Tamura T, Shiotsuki T. Precocious metamorphosis in transgenic

silkworms overexpressing juvenile hormone esterase. Proc Natl Acad Sci U S A.

2005; doi:10.1073/pnas.0500954102.

2.   Ito K, Kidokoro K, Sezutsu H, Nohata J, Yamamoto K, Kobayashi I, et al. Deletion

of a gene encoding an amino acid transporter in the midgut membrane causes

resistance to a *Bombyx* parvo-like virus. Proc Natl Acad Sci U S A. 2008;

doi:10.1073/pnas.0711841105.

3.   Sakurai T, Mitsuno H, Haupt SS, Uchino K, Yokohari F, Nishioka T, et al. A single

sex pheromone receptor determines chemical response specificity of sexual

behavior in the silkmoth *Bombyx mori*. PLoS Genet. 2011;

doi:10.1371/journal.pgen.1002115.

4.   Daimon T, Uchibori M, Nakao H, Sezutsu H, Shinoda T. Knockout silkworms reveal

a dispensable role for juvenile hormones in holometabolous life cycle. Proc Natl

Acad Sci U S A. 2015; doi:10.1073/pnas.1506645112.

5.   Kondo Y, Yoda S, Mizoguchi T, Ando T, Yamaguchi J, Yamamoto K, et al. Toll

ligand Spätzle3 controls melanization in the stripe pattern formation in caterpillars.

Proc Natl Acad Sci U S A. 2017; doi:10.1073/pnas.1707896114.

6.   Yukuhiro K, Sezutsu H, Tsubota T, Takasu Y, Kameda T, Yonemura Y. Insect silks

and cocoons: structural and molecular aspects. In: Cohen E, Moussian B, editors.

Extracellular composite matrices in Arthropods. Cham: Springer; 2016;

doi:10.1007/978-3-319-40740-1_18.

7.   Tamura T, Thibert C, Royer C, Kanda T, Abraham E, Kabma B, et al. Germline

transformation of the silkworm *Bombyx mori* L. using a *piggyBac* transposon-

derived vector. Nat Biotechnol. 2000; doi:10.1038/71978.

8. Tatematsu K, Kobayashi I, Uchino K, Suzutsu H, Iizuka T, Yonemura N, et al.

Construction of a binary transgenic gene expression system for recombinant

protein production in the middle silk gland of the silkworm *Bombyx mori*. Transgenic

Res. 2010; doi:10.1007/s11248-009-9328-2.

9. Mita K, Kasahara M, Sasaki S, Nagayasu Y, Yamama T, Kanamori H, et al. The

genome sequence of silkworm, *Bombyx mori*. DNA Res. 2004;

doi:10.1093/dnares/11.1.27.

10. Xia Q, Zhou Z, Lu C, Cheng D, Dai F, Li B, et al. A draft sequence for the genome

of the domesticated silkworm (*Bombyx mori*). Science. 2004;

doi:10.1126/science.1102210.

11. The International Silkworm Genome Consortium. The genome of a lepidopteran

model insect, the silkworm *Bombyx mori*. Insect Biochem Mol Biol. 2008;

doi:10.1016/j.ibmb.2008.11.004.

12. Xia Q, Cheng D, Duan J, Wang G, Cheng T, Zha X, et al. Microarray-based gene

expression profiles in multiple tissues of the domesticated silkworm, *Bombyx mori*.

Genome Biol. 2007; doi:10.1186/gb-2007-8-8-r162.

13. Yamamoto K, Nohata J, Kadono-Okuda K, Narukawa J, Sasanuma M, Sasanuma

S, et al. A BAC-based integrated linkage map of the silkworm *Bombyx mori*.

Genome Biol. 2008; doi:10.1186/gb-2008-9-1-r21.

14. Suetsugu Y, Futahashi R, Kanamori H, Kadono-Okuda K, Sasanuma S, Narukawa

J, et al. Large scale full-Length cDNA sequencing reveals a unique genomic

landscape in a lepidopteran model insect, *Bombyx mori*. G3. 2013;

doi:10.1534/g3.113.006239.

15. Kawamoto M, Jouraku A, Toyoda A, Yokoi K, Minakuchi Y, Katsuma S, et al. High-

quality genome assembly of the silkworm, *Bombyx mori*. Insect Biochem Mol Biol.

2019; doi:10.1016/j.ibmb.2019.02.002.

16. Chang H, Cheng T, Wu Y, Hu W, Long R, Liu C, et al. Transcriptomic analysis of

the anterior silk gland in the domestic silkworm (*Bombyx mori*) – insight into the

mechanism of silk formation and spinning. PLoS One. 2015;

doi:10.1371/journal.pone.0139424.

17. Shi R, Ma S, He T, Peng J, Zhang T, Chen X, et al. Deep insight into the

transcriptome of the single silk gland of *Bombyx mori*. Int J Mol Sciences. 2019;

doi:10.3390/ijms20102491.

18. Ishikawa E, Suzuki Y. Tissue- and stage-specific expression of sericin genes in the

middle silk gland of *Bombyx mori*. Dev Growth Differ. 1985; doi:10.1111/j.1440-

169X.1985.00073.x.

19. Garel A, Deleage G, Prudhomme J-C. Structure and organization of the *Bombyx

mori* Sericin 1 gene and of the Sericins 1 deduced from the sequence of the Ser 1B

cDNA. Insect Biochem Mol Biol. 1997; doi:10.1016/s0965-1748(97)00022-2.

20. Takasu Y, Yamada H, Tsubouchi K. Isolation of three main sericin components

from the cocoon of the silkworm, *Bombyx mori*. Biosci Biotechnol Biochem. 2002;

doi:10.1271/bbb.66.2715.

21. Zurovec M, Yonemura N, Kludkiewicz B, Sehnal F, Kodrik D, Viera LC, et al.

Sericin composition in the silk of *Antheraea yamamai*. Biomacromolecules. 2016;

doi:10.1021/acs.biomac.6b00189.

22. Takasu Y, Yamada H, Tsubouchi K. The silk sericin component with low

crystallinity. Sanshi-Konchu Biotec. 2006; doi:10.11416/konchubiotec.75.133.

23. Takasu Y, Yamada H, Tamura T, Sezutsu H, Mita K, Tsubouchi K. Identification

and characterization of a novel sericin gene expressed in the anterior middle silk

gland of the silkworm *Bombyx mori*. Insect Biochem Mol Biol. 2007;

doi:10.1016/j.ibmb.2007.07.009.

24. Dong Z, Guo K, Zhang X, Zhang T, Zhang Y, Ma S. Identification of *Bombyx mori*

sericin 4 protein as a new biological adhesive. Int J Biol Macromol. 2019;

doi:10.1016/j.ijbiomac.2019.03.166.

25. Kikuchi A, Nakazato T, Ito K, Nojima Y, Yokoyama T, Iwabuchi K, et al.

Identification of functional enolase genes of the silkworm *Bombyx mori* from public

databases with a combination of dry and wet bench processes. BMC Genomics.

2017; doi:10.1186/s12864-016-3455-y.

26. Ichino F, Bono H, Nakazato T, Toyoda A, Fujiyama A, Iwabuchi K, et al.

Construction of a simple evaluation system for the intestinal absorption of an orally

administered medicine using *Bombyx mori* larvae. Drug Discov Ther. 2018;

doi:10.5582/ddt.2018.01004.

27. Kobayashi Y, Nojima Y, Sakamoto T, Iwabuchi K, Nakazato T, Bono H, et al.

Comparative analysis of seven types of superoxide dismutases for their ability to

respond to oxidative stress in *Bombyx mori*. Sci Rep. 2019; doi:10.1038/s41598-

018-38384-8.

28. Zhu Z, Guan Z, Liu G, Wang Y, Zhang Z. SGID: a comprehensive and interactive

database of the silkworm. Database. 2019; doi:10.1093/database/baz134.

29. Zhao D, Woolner S, Bownes M. The Mirror transcription factor links signaling

pathways in *Drosophila* oogenesis. Dev Genes Evol. 2000;

doi:10.1007/s004270000081.

30. Li MA, Alls JD, Avancini RM, Koo K, Godt D. The large Maf factor Traffic Jam

controls gonad morphogenesis in *Drosophila*. Nat Cell Biol. 2003;

doi:10.1038/ncb1058.

31. Wang Z, Mann RS. Requirement for two nearly identical TGIF-related homeobox

genes in *Drosophila* spermatogenesis. Development. 2003; doi:10.1242/dev.00510.

32. Nanda S, DeFalco TJ, Loh SHY, Phochanukul N, Camara N, Doren MV, et al.

Sox100B, a *Drosophila* group E Sox-domain gene, is required for somatic testis

differentiation. Sex Dev. 2009; doi: 10.1159/000200079.

33. Vaskivuo TE, Anttonen M, Herva R, Billig H, Dorland M, Velde ERT, et al. Survival

of human ovarian follicles from fetal to adult life: apoptosis, apoptosis-related

proteins, and transcription factor GATA-4. J Clin Endocrinol Metab. 2001;

doi:10.1210/jcem.86.7.7679.

34. Mach V, Takiya S, Ohno K, Handa H, Imai T, Suzuki Y. Silk gland factor-1 involved

in the regulation of *Bombyx* sericin-1 gene contains fork head motif. J Biol Chem.

1995; doi:10.1074/jbc.270.16.9340.

35. Ohno K, Sawada J, Takiya S, Kimoto M, Matsumoto A, Tsubota T, et al. Silk gland

factor-2, involved in fibroin gene transcription, consists of LIM homeodomain, LIM-

interacting, and single-stranded DNA-binding proteins. J Biol Chem. 2013;

doi:10.1074/jbc.M113.514471.

36. Kimoto M, Tsubota T, Uchino K, Sezutsu H, Takiya S. Hox transcription factor Antp

regulates *sericin-1* gene expression in the terminal differentiated silk gland of

Bombyx mori. Dev Biol. 2014; doi:10.1016/j.ydbio.2013.12.002.

37. Kimoto M, Tsubota T, Uchino K, Sezutsu H, Takiya S. LIM-homeodomain

transcription factor Awh is a key component activating all three fibroin genes, *fibH*,

fibL and *fhx*, in the silk gland of the silkworm, *Bombyx mori*. Insect Biochem. Mol.

Biol. 2015; doi:10.1016/j.ibmb.2014.11.003.

38. Tsubota T, Tomita T, Uchino K, Kimoto M, Takiya S, Kajiwara H, et al. A *Hox* gene,

Antennapedia, regulates expression of multiple major silk protein genes in the

silkworm *Bombyx mori*. J Biol Chem. 2016; doi:10.1074/jbc.M115.699819.

39. Baumgartner S, Bopp D, Burri M, Noll M. Structure of two genes at the *gooseberry*

locus related to the *paired* gene and their spatial expression during *Drosophila*

embryogenesis. Genes Dev. 1987; doi:10.1101/gad.1.10.1247.

40. Grossniklaus U, Pearson RK, Gehring WJ. The *Drosophila sloppy paired* locus

encodes two proteins involved in segmentation that show homology to mammalian

transcription factors. Genes Dev. 1992; doi:10.1101/gad.6.6.1030.

41. Russell SRH, Sanchez-Soriano N, Wright CR, Ashburner M. The *Dichaete* gene of

Drosophila melanogaster encodes a SOX-domain protein required for embryonic

segmentation. Development. 1996;122:3669-76.

42. Couturier L, Mazouni K, Corson F, Schweisguth F. Regulation of Notch output

dynamics via specific E(spl)-HLH factors during bristle patterning in *Drosophila*. Nat

Commun. 2019; doi:10.1038/s41467-019-11477-2.

43. Yao J-G, Weasner BM, Wang L-H, Jang C-C, Weasner B, Tang C-Y, et al.

Differential requirements for the Pax6(5a) genes *eyegone* and *twin of eyegone*

during eye development in Drosophila. Dev Biol. 2008;

doi:10.1016/j.ydbio.2007.12.037.

44. Ferree PM, Fang C, Mastrodimos M, Hay BA, Amrhein H, Akbari OS. Identification

of genes uniquely expressed in the germ-line tissues of the jewel wasp *Nasonia*

vitripennis. G3(Bethesda). 2015; doi:10.1534/g3.115.021386.

45. Kondo S, Vedanayagam J, Mohammed J, Eizadshnass S, Kan L, Pang N, et al.

New genes often acquire male-specific functions but rarely become essential in

Drosophila. Genes Dev. 2017; doi:10.1101/gad.303131.117.

46. Kaessmann H. Origins, evolution, and phenotypic impact of new genes. Genome

Res. 2010; doi:10.1101/gr.101386.109.

47. Li W, Cheng T, Hu W, Peng Z, Liu C, Xia Q. Genome-wide identification and

analysis of JHBP-domain family members in the silkworm *Bombyx mori*. Mol Genet

Genomics. 2016; doi:10.1007/s00438-016-1245-5.

48. Hu X, Zhu M, Wang S, Zhu L, Xue R, Cao G, et al. Proteomics analysis of digestive

juice from silkworm during *Bombyx mori* nucleopolyhedrovirus infection.

Proteomics. 2015, doi:10.1002/pmics.201400475.

49. Horne I, Haritos, VS. Multiple tandem gene duplications is a neural lipase gene

cluster in *Drosophila*. Gene 2008, doi:10.1016/j.gene.2008.01.002.

50. Julien E, Coulon-Bublex M, Garel A, Royer C, Chavancy G, Prudhomme J-C, et al.

Silk gland development and regulation of silk protein genes. In: Gilbert LI, editor.

Comprehensive molecular insect science. Amsterdam: Elsevier; 2005;369-384.

51. Akai H. The silk gland. In: Akai H, editor. Ultrastructural morphology of insects.

Tokyo: University of Tokyo press; 1976;57-114.

52. Takasu Y, Hata T, Uchino K, Zhang Q. Identification of Ser2 proteins as major

sericin components in the non-cocoon silk of *Bombyx mori*. Insect Biochem Mol

Biol. 2010. doi:10.1016/j.ibmb.2010.02.010.

53. Bello B, Horard B, Couble P. The selective expression of silk-protein-encoding

genes in *Bombyx mori* silk gland. Bull Inst Pasteur. 1994;92;81-100.

54. Akai H. Porous cocoon filaments – Their characteristic and formation. Int J wild

silkmoth & silk. 2005;10:57-74.

55. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics. 2014. doi:10.1093/bioinformatics/btu170.

56. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment

and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol. 2019.

doi:10.1038/s41587-019-0201-4.

57. Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. Transcript-level expression

analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. Nat Protoc.

2016. doi:10.1038/nprot.2016.095.

58. gffcompare [https://ccb.jhu.edu/software/stringtie/gffcompare.shtml].

59. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq

quantification. Nat Biotechnol. 2016. doi:10.1038/nbt.3519.

60. TIBCO Spotfire [http://spotfire.tibco.com/better-world-donation-program/].

61. Morpheus [https://software.broadinstitute.org/morpheus].

62. R Core Team (2019). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

63. R Core Team (2020). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

64. TransDecoder [https://transdecoder.github.io/].

65. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped

BLAST and PSI-BLAST: a new generation of protein database search programs.

Nucleic Acids Res. 1997. doi:10.1093/nar/25.17.3389.

66. Tabunoki H, Ono H, Ode H, Ishikawa K, Kawana N, Banno Y, et al. Identification of

key uric acid synthesis pathway in a unique mutant silkworm *Bombyx mori* model of

Parkinson's disease. 2013. doi:10.1371/journal.pone.0069130.

67. Kinsella RJ, Kähäri A, Haider S, Zamora J, Proctor G, Spudich G, et al. Ensembl

BioMarts: a hub for data retrieval across taxonomic space. Database (Oxford).

2011. doi:10.1093/database/bar030.

68. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al.

Metascape provides a biologist-oriented resource for the analysis of systems-level

datasets. Nat Commun. 2019. doi:10.1038/s41467-019-09234-6.

69. SilkBase [http://silkbase.ab.a.u-tokyo.ac.jp/cgi-bin/index.cgi].

70. KAIKObase [https://kaikobase.dna.affrc.go.jp/].

71. TogoTV [https://togotv.dbcls.jp/pics.html].

72. Creative Commons [http://creativecommons.org/licenses/by/4.0/deed.en].

Tables

Table 1. Samples for RNA-seq and run accession IDs.

Sample	SRA Run ID	Strain	Sex	Reference
ASG-1,2,3	DRR186474,DRR186475,DRR186476	p50T	Male	This study
MSG_A-1,2,3	DRR186477,DRR186478,DRR186479	p50T	Male	This study
MSG_M-1,2,3	DRR186480,DRR186481,DRR186482	p50T	Male	This study
MSG_P-1,2,3	DRR186483,DRR186484,DRR186485	p50T	Male	This study
PSG-1,2,3	DRR186486,DRR186487,DRR186488	p50T	Male	This study
FB-1,2,3	DRR186489,DRR186490,DRR186491	p50T	Male	This study
MG-1,2,3	DRR186492,DRR186493,DRR186494	p50T	Male	This study
MT-1,2,3	DRR186495,DRR186496,DRR186497	p50T	Male	This study
TT-1,2,3	DRR186498,DRR186499,DRR186500	p50T	Male	This study
OV-1,2,3	DRR186501,DRR186502,DRR186503	p50T	Female	This study
BN_TT-1,2,3	DRR068893,DRR068894,DRR068895	o751	Male	[25]
BN_FB-1,2,3	DRR095105,DRR095106,DRR095107	o751	Male	[27]
BN_MG-1,2,3	DRR095108,DRR095109,DRR095110	o751	Male	[26]
BN_MT-1,2,3	DRR095111,DRR095112,DRR095113	o751	Male	[27]
BN_SG-1,2,3	DRR095114,DRR095115,DRR095116	o751	Male	[27]

Figures

Fig. 1

P50T strain
5th instar day 3 larva

Male

Female → Ovary

Testis
Midgut
Fat body
Malpighian tuble
Silk gland (divided into multiple parts)

Sequencing

RNA-seq data

Bombyx mori RNA-seq data from SRA

Reference genome data (Kawamoto et al., 2019)

Mapping (Hisat2) and assembling (Stringtie)

Estimation of expression (Kallisto)

Further analyses

Expression and functional annotation data

Functional prediction (Blast)

Reference transcriptome data

Human and *Drosophila* protein data set from Ensembl DB

Fig. 2

Fig. 3

A

60,000
50,000
40,000
30,000
20,000
10,000
0

Loci   Transcripts

■ Gene model data (Kawamoto et al., 2019)
■ Reference transcriptome data

B

Reference transcriptome
51,926

New loci
7,704

Identified
in gene
models
16,880

New isoform
27,342

Fig. 4

Fig. 5

Fig. 6

(A) TT

(B) OV

Fig. 7

Fig. 8

(A)

(B)

Fig. 9

Fig. 10

(A)

(B)