Frost and drought: effects of extreme weather events on stem carbon

dynamics in a Mediterranean beech forest

Ettore D'Andrea[1*], Negar Rezaie[1*], Peter Prislan[2], Jozica Gričar[2], Jan Muhr[3,4], Alessio Collalti[5,6],

Giorgio Matteucci[1]

[1]CNR-ISAFOM, via Patacca 2, 80056 Ercolano (NA), Italy

[2]Department of Yield and Silviculture, Slovenian Forestry Institute, SI-1000 Ljubljana, Slovenia

[3]University of Göttingen, Bioclimatology, Büsgenweg 2, 37077 Göttingen, Germany

[4]Max-Planck-Institute for Biogeochemistry, Department of Biogeochemical Processes, Hans-Knoell-

Str. 10. 07745 Jena, Germany

[5]CNR-ISAFOM, Via Cavour 4/6, 87036, Rende (CS), Italy

[6] Department of Innovation in Biological, Agro-food and Forest Systems, University of Tuscia,

01100 Viterbo, Italy

*__Corresponding Authors__:

ettore.dandrea@isafom.cnr.it, +393389119720

rezaie.negar@gmail.com, +393349410230

Summary

The effects of short-term extreme events on tree functioning and physiology are still rather elusive. European beech is one of the most sensitive species to late frost and water shortage. We investigated the intra-annual C dynamics in stems under such conditions.

Wood formation and stem CO_2 efflux were monitored in a Mediterranean beech forest for three years (2015–2017), including a late frost (2016) and a summer drought (2017).

The late frost reduced radial growth and, consequently, the amount of carbon fixed in the stem biomass by 80%. Stem carbon efflux in 2016 was reduced by 25%, which can be attributed to the reduction of effluxes due to growth respiration. Counter to our expectations, we found no effects of the 2017 summer drought on radial growth and stem carbon efflux.

The studied extreme weather events had various effects on tree growth. Even though late spring frost had a devastating impact on beech radial growth in the current year, trees fully recovered in the following growing season, indicating high resilience of beech to this stressful event.

Keywords: drought, extreme weather events, *Fagus sylvatica* L. (beech), growth, late frost, resilience, stem carbon efflux, wood formation.

Introduction

Even small changes in the mean or variation of a climate variable cause disproportionally large changes in the frequency of extreme weather events, recognized as major drivers of current and future ecosystem dynamics (Frank *et al.* 2015). In the near future, the Mediterranean region is predicted to be the most vulnerable of the European regions to global change (Schröter *et al.* 2005). Changes in temperature and precipitation regimes may increase drought risk (Schröter *et al.* 2005), which can negatively affect physiological performance (Rezaie *et al.* 2018), as well as the growth and competitive strength (Peuke, Schraml, Hartung

& Rennenberg 2002) of common beech, one of the most important and widespread

broadleaved trees in Europe.

Increasing spring temperatures can trigger earlier leaf unfolding (Gordo & Sanz 2010;

Allevato *et al.* 2019), which in turn results in a higher risk that young leaves are exposed to

late spring frost (Augspurger 2013), especially at higher elevations (Vitasse, Schneider,

Rixen, Christen & Rebetez 2018). Temperatures below −4°C can kill the developing new

shoots and leaves, thus reducing the photosynthetic area and ultimately the trees' growth. In

the case of late frost, depending on the intensity of damage, the formation of new leaves

requires a high amount of reserves (Dittmar, Fricke & Elling 2006; D'Andrea *et al.* 2019).

Tree stems play an important role in the carbon balance of forest ecosystems (Yang, He,

Aubrey, Zhuang & Teskey 2016). Part of the carbon (C) fixed by photosynthesis is allocated

to the stem, and some is respired by stems and emitted into the atmosphere. Radial growth –

an often used proxy for the overall allocation of C to the stem (Bascietto, Cherubini &

Scarascia-Mugnozza 2004; Cuny *et al.* 2015; Chan, Berninger, Kolari, Nikinmaa & Hölttä

2018) – is largely related to the process of wood formation, which can be divided into five

(main) developmental phases: *i)* cambial cell division; *ii)* cell enlargement; *iii)* secondary wall

deposition and, *iv)* cell wall thickening (lignification); while *v)* in the case of vessels and

fibres, also genetically-programmed cell death or *apoptosis* (Prislan, Čufar, De Luis & Gričar

2018). The whole process is sensitive to many factors, such as leaf phenology (Michelot,

Simard, Rathgeber, Dufrêne & Damesin 2012), temperature (Begum, Nakaba, Oribe, Kubo &

Funada 2007), drought (Linares, Camarero & Carreira 2009), tree-size and social status

(Rathgeber, Rossi & Bontemps 2011) and tree vigour (Gričar, Krže & Čufar 2009).

As also reported by Damesin (2002), stem respiration may represent up to 1/3 of the overall

above ground respiration and 1.4% of the annual carbon assimilation. However, field

measurements of actual stem respiration (RS) are difficult if not impossible (Teskey, Saveyn,

Steppe & McGuire 2008), and the most commonly measured proxy, stem CO_2 efflux (ES), is

likely to underestimate local respiration (Trumbore, Angert, Kunert, Muhr & Chambers 2013).

Previous studies have reported a strong correlation between RS and ES, with ES ranging

between 82–94% and 86–91% of RS in *Populus deltoides* W.Bartram ex Marshall (Saveyn,

Steppe, Mc Guire, Lemeur & Teskey 2008) and *Dacrydium cupressinum* Lamb stems

(Bowman *et al.* 2005), respectively. Another study using O_2 uptake, as an alternative proxy

for actual respiration, and comparing it to traditionally measured ES, showed that ES on

average underestimated RS by about 41% (CO_2 was not emitted locally at the point of

measurement) (Hilman *et al.* 2019). ES and RS are different because part of the CO_2 produced

by respiration is not released directly through the bark into the atmosphere, but is dissolved in

xylem sap and is carried upward by the transpiration stream (Bloemen *et al.* 2014). In

addition, ES is affected by CO_2 deriving from root respiration, which is also carried upward

into the stem (Bloemen, McGuire, Aubrey, Teskey & Steppe 2013). Moreover, part of respired

CO_2 can be fixed in xylem storage pools (Ubierna *et al.* 2009). A recent global estimate

showed that the stem CO_2 efflux (ES) alone, from boreal to tropical forests combined, was 6.7

($\pm$ 1.1) Pg C yr^{-1}, accounting for 11% and 20% of global forest ecosystem gross primary

production (GPP) and net primary production (NPP), respectively (Yang *et al.* 2016).

ES is influenced by many factors, such as air temperature (Yang et al., 2016), growth rate

(Damesin, Ceschia, Le Goff, Ottorini & Dufrêne 2002), distribution and turnover of living

cells (Collalti *et al.* 2019), nitrogen concentration (Ceschia, Damesin, Lebaube, Pontailler &

Dufrêne 2002) and tree social class (Guidolotti, Rey, D'Andrea, Matteucci & De Angelis

2013). Both RS and ES are also affected by growth respiration (R_G), which provides the

energy for synthesizing new tissues; and by maintenance respiration (R_M), which maintains

existing living cells (Ceschia *et al.* 2002). Separating ES into these components allows further

investigation of stem carbon budgeting and tissue costs (Chan *et al.* 2018).

Despite the crucial role of extreme events and increasing attention on their prospective

increasing role in future climate scenarios, information on the effect of short-term extreme

events on tree functioning is still fairly elusive (Carrer, Brunetti & Castagneri 2016; Gazol *et*

al. 2019). In this respect, not much is known about the interaction between wood formation

(xylogenesis) and ES. At the seasonal timescale, the capacity of micro-coring technique to

identify phenological phases of wood formation allows to attribute metabolic costs to each

one of them (Meir, Mencuccini & Coughlin 2019).

A deeper investigation of this link is crucial, especially in the context of climate change,

associated with increased frequency of extreme weather events (e.g., drought and late frost),

which may greatly modify the contribution of these processes to the C cycle.

In this context, we monitored xylogenesis, together with ES and overall growth, in a mature

Mediterranean beech forest (*Fagus sylvatica* L.) from 2015 to 2017 − a period characterized

by a spring late frost (2016) and a summer drought (2017) − with the objective of unravelling

the intra-annual C dynamics in stems under different climatic conditions and in response to

extreme weather events. On the hypothesis that extreme weather events would alter the stem

C dynamics at both tree and stand scales, we investigated tree scale physiological processes;

specifically, of growth (xylogenesis) and respiration (proxied by stem CO_2 efflux), and then

upscaled to stand-scale growth and C emission related to this process. We hypothesized that:

1) cambial activity and radial growth may have ceased soon after leaf death due to the 2016

spring late frost; 2) second leaf re-sprouting starts at the expense of stem growth; 3) the 2017

summer drought would have negatively impacted stem biomass production and effluxes; and

that, 4) climatic variability and extreme weather events are important factors in C dynamics

on tree and stand scales.

Material and methods

Study site and description of weather events

The measurements were carried out between 2015 and 2017 on a long-term monitored beech stand *(Fagus sylvatica* L.) located at Selva Piana (41°50'58" N, 13°35'17" E, 1,560 m elevation), close to Collelongo (Abruzzi Region, Italy) in the Central Apennines. The site, established in 1991 and since 2006 part of the long-term ecological research (LTER) network, is located in a 3000 ha forest included in the wider forest area of the external belt of Abruzzi National Park. In 2017, the stand density was 725 trees ha^{-1}, the basal area was 45.77 m^2 ha^{-1} with a mean diameter at breast height (DBH) of 28.5 cm, and a mean tree height of 23 m. Mean tree age in 2013 was estimated to be about 110 years. Site topography is gently sloping and the soil is humic alisol with a variable depth (40–100 cm), developed on calcareous bedrock (Chiti *et al.* 2010). The climate is of Mediterranean mountain type, during the period 1989 – 2014 the mean annual temperature was 7.2°C, and the mean annual precipitation was 1178 mm, of which ~10% falls in summer (Guidolotti *et al.* 2013).

In the night between the 25[th] and 26[th] of April 2016 (Day of Year, DOY 115), a spring late frost occurred in Central and Southern Italy, causing leaf damage in many beech stands (Bascietto, Bajocco, Mazzenga & Matteucci 2018; Greco *et al.* 2018; Nolè, Rita, Ferrara & Borghetti 2018; Allevato *et al.* 2019). At the Selva Piana site, the air temperature reached −6°C at canopy level, destroying the whole-stand canopy and leaving the trees without leaves for almost two months (Bascietto *et al.* 2018; D'Andrea *et al.* 2019).

In the summer of 2017, a widespread positive temperature anomaly affected Central Italy and the Balkans, with a duration ranging from 20 to 35 days (Rita *et al.* 2019). In the same year, annual precipitation was 950 mm, with only 54 mm of precipitation throughout the entire summer; while only 1 mm of rain and a mean air temperature (23.9 °C) ~ 2°C warmer than

the long-term average (1989–2014) characterized August 2017, leading to a Standardized

Precipitation Evapotranspiration Index (SPEI) > 1.5 during the vegetative season (Fig. 1).

Tree selection, wood formation dynamics, xylem phenology and C fixation

Sampling was performed on five trees that were selected for their similarity, with a site tree

ring chronology following the methodology described in Rezaie *et al*. (2018) and as done in

other studies on wood formation and stem CO_2 efflux (e.g. Ceschia *et al.*, 2002; Damesin *et*

al., 2002; Gruber *et al.*, 2009; Delpierre *et al.*, 2019). Microcore collection and ES

measurements were carried out from April 2015, before leaf unfolding, until November 2017,

when the trees had completely lost their leaves.

Microcores (2 mm diameter and 15 mm long) were extracted from each tree at 1.1-1.7 m

above ground using a Trephor tool (Rossi, Menardi, Fontanella & Anfodillo 2005).

Microcores were collected 15, 12, and 14 times in 2015, 2016, and 2017, respectively. To

avoid wound effects, cores were sampled at a distance of at least 5 cm from each other. The

microcores, containing bark, cambium, newly developing xylem and 1-2 older xylem rings,

were immediately stored in formaldehyde-ethanol-acetic acid solution (FEA) in the field.

Cross-sections of the microcores were prepared following the standard methodology (Prislan,

Gričar, De Luis, Smith & Cufar 2013) and were photographed in high definition under a

Leica DM 4000 microscope (Leica Microsystems, Wetzlar, Germany) using transmission and

polarized light. Histometrical analyses were performed on images taken with a Leica DFC

280 digital camera using the LAS (Leica Application Suite) image analysis system (Leica

Microsystems, Germany). On each photographed cross-section, the number of cambium cells

was counted and the widths of the developing xylem were measured along three radial

directions.

The dynamics of xylem formation were analysed by fitting the Gompertz function to xylem

increments (Prislan *et al.* 2018; Rathgeber, Santenoise & Cuny 2018), corrected for the

previous tree ring width (Camarero, Guerrero-Campo & Gutierrez 1998; Oladi, Pourtahmasi,

Eckstein & Bräuning 2011), as follows:

$$y = \alpha \exp\left[-e^{(\beta - kt)}\right] \qquad (1)$$

where y is the cumulative ring width (μm) at time t (day of the year), α is the final asymptotic

size representing the annual potential growth, β is the x-axis placement parameter and k is the

rate of change parameter.

For each tree and monitoring year, the following phenological xylem formation phases were

recorded: *i*) cambium *reactivation*, *ii*) beginning of cell *enlargement* (bE); *iii*) beginning of

cell wall *thickening* (bW); *iv*) beginning of cell *maturation* (bM); *v*) *cessation* of cell

enlargement (cE); and *vi*) cessation of cell wall thickening and *lignification* (cW). The date of

cambium reactivation was assessed as the average between dates when an increase of

cambium cells was observed (i.e., from 3-4 to 6-7 cells in a radial row) (Čufar, Prislan, De

Luis & Gričar 2008; Deslauriers, Rossi, Anfodillo & Saracino 2008). Phases of xylem growth

and ring formation were computed using logistic regression, spanning from the 50%

probability that phenophases have started or ended (Rathgeber *et al.* 2018). Based on

phenological phases, the durations of key wood formation phases were calculated: *i*) the

overall duration of the enlargement period (dE = cE – bE); *ii*) the duration of the wall-

thickening period (dW = cW – bW); and *iii*) the total duration of wood formation (i.e., the

duration of xylogenesis) (dX = cW – bE). Data were analysed using the CAVIAR (v2.10-0)

package (Rathgeber *et al.* 2018) built for R statistical software (R Development Core Team

2018).

Starting from the detailed time-resolved data from tree microcores, the annual C fixed in the

stem (SG_t) was estimated for each of the sampled tree, as follows:

$$SG_t = \frac{0.46 \times (BS_t - BS_{t-1})}{\Delta t} \qquad (2)$$

where SG_t is the amount of C fixed in the stem per year expressed in Mg C yr^{-1}, 0.46 is the

carbon content of the woody tissues (Scarascia-Mugnozza, Bauer, Persson, Matteucci &

Masci 2000), BS_t and BS_{t-1} are the stem biomass in Mg of Dry Matter (DW) of at the

beginning and at the end of each sampling year, Δt is the time variation (one year).

The site-specific allometric equation for beech used for BS was that proposed by Masci

(2002):

$$BS = \frac{283.734 \times DBH^{2.134}}{10^6} \qquad (3)$$

208   Where BS is in Mg DW, and DBH is the diameter (in cm) at 1.30 m ($R^2 = 0.96$, p-value $<$

209   0.01).

210   Upscaling to stand scale was performed according to the following equation:

$$SG = SG_t \times n \times \frac{d_{avg}}{d_{st}} \qquad (4)$$

212   with SG expressed in Mg C ha^{-1} yr^{-1}, n being the number of trees per hectare, d_{avg} the average

213   diameter (cm) and d_{st} the diameter of the sampled tree (cm).

Stem CO_2 efflux (ES)

Two PVC collars (10 cm diameter and 5 cm high, one facing north and one south) were fixed

on each tree with flexible plastic ties and sealed leak tight with Terostat (Henkel KgaA,

Germany). When present, bark mosses and lichens were removed. Stem CO_2 efflux was

measured with a portable IRGA (EGM 4, PP-System, Hitchin, UK), equipped with a closed-

dynamic chamber (SRC-1, PP-System, Hitchin, UK), which was tightened. Each

measurement consisted of a 120-second loop, in which the CO_2 concentration inside the

chamber was measured every 5 seconds. During measurements, the CO_2 concentration

typically increased by 10 to 50 μmol mol^{-1}. ES measurements were performed 16, 12, and 12

times in 2015, 2016, and 2017, respectively.

Stem CO_2 efflux (ES) was calculated as:

$$ES = K_{CO_2} \div V_{mol} \times \frac{V_{cuv}}{A} \qquad (5)$$

where ES is the stem CO_2 efflux per unit surface area (μmol m^{-2} s^{-1}), K_{CO2} (μmol mol^{-1} s^{-1}) is

the slope of the regression between CO_2 concentration and time during measurements, V_{mol},

the molar volume, is the volume occupied by one mole of CO_2 (m^3 mol^{-1}), at the air pressure

(measured by built-in sensor of the EGM-4) and air temperature (T_{air} in °C) at the

measurement time, A is the exposed lateral surface area of the stem (m^2), and V_{cuv} is the sum

of SRC-1 and collar volumes (m^3).

An exponential function was used to assess the relationship between ES and T_{air}

$$ES = a \times e^{T_{air} \times b} \qquad (6)$$

and ES overall temperature sensitivity for a 10 °C increase (Q_{10}) was calculated according to

Gruber *et al.* (2009) as:

$$Q_{10} = 10^{K_T \times 10} \qquad (7)$$

where K_T is the regression slope taken from linear regression of log10 of ES versus T_{air}.

From the wood formation and xylem phenology analysis described above, we identified wood

(*w*) and non-wood (*nw*) formation periods for each tree, making it possible to divide the

measured ES into two groups, ES_w and ES_{nw}. According to Eq. 5, we calculated for each

group the specific CO_2 efflux at a base air temperature of 15°C (ES_{15w} and ES_{15nw}) and the

specific Q_{10} (Q_{10w} and Q_{10nw}).

During the non-wood formation period, ES_{nw} ($\mu mol\ m^{-2}\ s^{-1}$) was constituted only by the

effluxes derived by maintenance respiration (ESb, $\mu mol\ m^{-2}\ s^{-1}$), which was calculated as:

$$ES_{nw} \equiv ESb = ES_{15nw} \times Q_{10nw}^{\frac{(T_{air}-15)}{10}} \quad (8)$$

255   During the wood formation period, ES_w ($\mu mol\ m^{-2}\ s^{-1}$), which is affected by both

256   maintenance and growth respiration, was calculated as:

$$ES_w = ES_{15w} \times Q_{10w}^{\frac{(T_{air}-15)}{10}} \quad (9)$$

We assumed that ESb and its relationship with air temperature was also valid during the wood

formation period, although this approach does not account for the acclimatisation of

maintenance respiration to temperature during warmer periods (Collalti *et al*., 2018 and

references therein). However, there are contrasting hypotheses on the magnitude of

acclimatisation (Carey *et al*., 1997; Stockfors *&* Linder, 1998). Under the above-mentioned

assumptions, we calculated the stem CO_2 efflux due to growth respiration, ESg ($\mu mol\ m^{-2}$

s^{-1}), as:

$$ESg = ES_w - ESb \quad (10)$$

The daily C effluxes of the whole stem were obtained by integrating, over the entire stem

area, the effluxes through equation 7, 8, 9, using half-hourly T_{air} values measured at the site.

The stem area was calculated as follow:

$$LA = 0.464 \times DBH - 2.083 \qquad (11)$$

where LA is the stem lateral area (m^2) ($R^2 = 0.828$, p-value < 0.01, for more details on the

equation see additional material Methods S1). Using DBH, we considered the measurement at

1.30 m to be representative of the whole stem, even though contrasting effects of height on

stem CO_2 effluxes have been reported (see Damesin *et al.*, 2002; Katayama *et al.*, 2019).

Annual values of each of the C fluxes of the five sampled trees (TES, TESb, TESg, see Table

1 for definitions) were obtained by summing up the daily values.

On the assumption that the selected trees were representative of the stand, annual values of

each of the fluxes at stand scale (AES, AESb, AESg, Mg C ha^{-1} yr^{-1}, see Table 1 for

definitions) were calculated:

$$AES_x = 10^{-6} \times TES_x \times n \times \frac{d_{avg}}{d_{st}} \quad (12)$$

where TES_x was the effluxes at tree level (TES, TESb, TESg, g C yr^{-1}), n is the number of

trees per hectare, d_{avg} is the average diameter (cm) and d_{st} is the diameter of the sampled tree

(cm).

Meteorological and phenological data

For the period 1989-2014, FLUXNET2015 release half-hourly air temperature and

precipitation were used (Reyer *et al.* 2019). For the period of the study (2015-2017),

measured data were gap filled using downloaded data by the ERA5 database of the European Centre for Medium-Range Weather Forecasts (ECMWF) (https://www.ecmwf.int/en/forecasts/datasets/archive-datasets/reanalysis-datasets/era5), according to FLUXNET 2015 release equations.

The Standardized Precipitation Evapotranspiration Index (SPEI), considered the most appropriate index for the Mediterranean climate (Vicente-Serrano *et al.* 2013), was used to assess the magnitude of the drought in 2017. This index is based on the difference between precipitation and potential evapotranspiration (PET), computed according to Hargreaves' equation. The 3-month SPEI was calculated for the site for the period 1989-2017, using the SPEI package in R.

Leaf phenology was monitored using the MODIS Leaf Area Index product (LAI, MOD15A2H, https://modis.gsfc.nasa.gov/) with 8-day temporal resolution and 500- meter spatial resolution. The date of onset of photosynthetic activity (green up) and the date at which plant green leaf area peaked at its annual maximum (maturity) were assessed through the rate of change in the curvature of the fitted logistic models (Zhang *et al.* 2003).

Statistical data analysis

Descriptive parameters of growth and xylem phenology were tested using one-way repeated measures analysis of variance, with years as factor, followed by post-hoc (Holm-Sidak method). An exponential equation was used to evaluate the relationship between ES and T_{air}. Differences among ES parameters (Q_{10} and ES_{15}) were tested using two-way repeated measures Anova (two factor repetition), using year and period (non-wood formation, wood formation) as predictive factors. Multiple comparisons were performed by the Holm-Sidak method. Linear regressions were used to assess the relationship between tree ring widths (TRW) and mean annual ES. We tested data normality and constant variance using the

Shapiro-Wilk test and the Spearman rank correlation between the absolute values of the

residuals and the observed value of the dependent variable.

Results

Wood formation dynamics

The date of onset of photosynthetic activity, based on leaf area index (LAI) dynamics,

differed among the study years, occurring at DOY 118, 95 and 127 in 2015, 2016 and 2017,

respectively. In all three years, cambium reactivation occurred after leaf unfolding at DOY

123 ± 4, 118 ± 8, 138 ± 6 in 2015, 2016 and 2017, respectively (Fig. 2). In 2016, cambium

cell production also continued after the late frost event, but at considerably lower rates.

Different intra-annual growth patterns were observed during the three study years, especially

in the year of the late frost (2016, Fig. 3, Table 1). In 2016, the maximum growth rate (rx) (F

= 8.469, p-value = 0.014) was lower and was reached 3 weeks earlier (tx) (F = 22.667, p-

value < 0.001) than in the other two years. The different intra-annual growth patterns also

resulted in significantly narrower tree rings in 2016 (230.12 ± 1.54 μm) (F = 13.272, p-value

< 0.01) than in 2015 (1312.17 ± 196.15 μm) and 2017 (1234.80 ± 269.32 μm).

Differences were also observed for the beginning, cessation, and duration of wood formation

phases (Fig. S2). The beginning of the enlargement phase occurred earliest in 2016 and latest

in 2017 (F = 34.789, p-value < 0.001). In contrast, the cessation of this phase was observed

latest in 2015 (F = 17.155, p-value < 0.01). Consequently, the duration of the enlargement

phase was longer in 2015 (110 ± 22 days) than in 2016 (82 ± 4 days) and 2017 (78 ± 4 days)

(F = 8.025, p-value = 0.01).

The beginning of the wall thickening phase did not differ among the years (F = 4.188, p-value

= 0.06). The cessation of this phase occurred latest in 2015 (F = 69.167, p-value < 0.001). The

duration of the wall thickening phase was thus shorter in 2016 (57 ± 5 days) than in 2015 (99

± 6 days) and 2017 (75 ± 4 days) (F = 26.561, *p*-value < 0.001). We also observed a delay in

the beginning of cell maturation in 2016 (at DOY 200 ± 4) with respect to 2015 (at DOY 178

± 4) and 2017 (at DOY 176 ± 2) (F = 11.650, *p*-value < 0.01). The overall duration of wood

formation was longer in 2015 (128 ± 5 days) than in 2016 (98 ± 8 days) and 2017 (97 ± 8

days) (F = 12.561, *p*-value < 0.001).

Stem CO_2 efflux (ES)

During the monitoring period (April 2015 – November 2017), the measured ES ranged

between 0.16 ± 0.03 µmol CO_2 m^{-2} s^{-1} (December 2015) and 3.01 ± 0.40 µmol CO_2 m^{-2} s^{-1}

(August 2017) (Fig. 5). Mean ES measured in 2016 (0.68 ± 0.19 µmol CO_2 m^{-2} s^{-1}) was

lower (F = 24.476, *p*-value < 0.01) than in 2015 (1.11 ± 0.40 µmol CO_2 m^{-2} s^{-1}) and 2017

(1.29 ± 0.30 µmol CO_2 m^{-2} s^{-1}).

In each year, ES was strongly related to air temperature through the standard exponential

function (Fig. 6). The relation was confirmed at different aggregation levels of measurements

(whole year, wood formation and non-wood formation periods; see also Table S1 in

Supporting Information).

Average Q_{10} was 2.71 ± 0.15, 2.11 ± 0.18 and 2.68 ± 0.15, in 2015, 2016 and 2017,

respectively. The Q_{10} parameter was not strongly affected by the sampling year (*p*-value =

0.059), although the values in 2016 were 22% lower than in the other two years. Wood

formation affected the Q_{10} parameter (F = 31.563, *p*-value < 0.01) with Q_{10w} and Q_{10nw}

calculated to be 3.06 ± 0.15 and 1.93 ± 0.14 (t = 5.571, *p*-value < 0.01), respectively. This

difference was confirmed for all of the sampled years.

ES_{15} was also affected by the different conditions of the monitoring years (F = 7.094, *p*-value

= 0.01) with mean values in 2016 (0.63 ± 0.07 µmol CO_2 m^{-2} s^{-1}) lower than in 2015 (0.93 ±

0.12 µmol CO_2 m^{-2} s^{-1}) and 2017 (0.82 ± 0.12 µmol CO_2 m^{-2} s^{-1}). Similar to Q_{10}, the wood

formation period also affected ES_{15}, with ES_{15w} (0.84 ± 0.22 µmol CO_2 m^{-2} s^{-1}) higher than

ES_{15nw} (0.73 ± 0.02 µmol CO_2 m^{-2} s^{-1}, F = 7.094, p-value = 0.01). Furthermore, during the

wood formation period, ES_{15w} was higher in 2015 (1.03 ± 0.07 µmol CO_2 m^{-2} s^{-1}) and 2017

(0.90 ± 0.07 µmol CO_2 m^{-2} s^{-1}) than in 2016 (0.60 ± 0.09 µmol CO_2 m^{-2} s^{-1}). No differences

among years were found for ES during the non-wood formation periods.

Annual ES for individual trees, per unit of lateral surface area, ranged between 112 g C m^{-2}

yr^{-1} in 2016 (tree 4) and 349 g C m^{-2} yr^{-1} in 2017 (tree 2). Average total ES for all sampled

trees was lower in 2016 (182 ± 25 g C m^{-2} yr^{-1}, F = 12.007, p-value < 0.01) than in 2015 (258

± 27 g C m^{-2} yr^{-1}) and 2017 (233 ± 35 g C m^{-2} yr^{-1}).

The estimated contribution of maintenance respiration to ES for individual trees ranged

between 112 g C m^{-2} yr^{-1} in 2016 (tree number 4 showed no growth) and 284 g C m^{-2} yr^{-1} in

2017 (tree number 2), and was lower, on average, in 2016 (169 ± 21 g C m^{-2} yr^{-1}) than in

2015 (211 ± 18 g C m^{-2} yr^{-1}) (q = 5.104, p-value = 0.017).

Likewise, the estimated contribution of wood formation to ES for individual trees varied

between 0 in 2016 (tree number 4) and 70 g C m^{-2} yr^{-1} in 2015 (tree number 2) and was

significantly lower (F = 8.144, p-value = 0.012) on average in 2016 (14 ± 5 g C m^{-2} y^{-1}) than

in 2015 (48 ± 9 g C m^{-2} yr^{-1}) and 2017 (39 ± 8 g C m^{-2} yr^{-1}). In relative terms, the

contribution of wood formation to ES was estimated to be 18 ± 2%, 9 ± 3% and 16 ± 3% in

2015, 2016 and 2017, respectively, with the remaining CO_2 efflux originating from

maintenance respiration.

Radial growth and stem C effluxes

During the study period, annual average measured ES and tree ring widths were closely

related (Fig. 5). Seasonal patterns of ES were similar in the three years, but with different

amplitudes (Fig. 8). Moreover, ESb, the stem C effluxes affected by maintenance respiration,

showed a similar pattern among the three years. We observed a time-lag between the time of

maximum growth rate (tx) and maximum ES values of 23 ± 2 days, 31 ± 2 days and 29 ± 1

days in 2015, 2016 and 2017, respectively. Differences between years were not significant (F

= 3.317, p-value = 0.07).

From tree to stand level

Annual stand-level stem C emissions (AES) were lower in 2016 (1.58 ± 0.22 Mg C ha^{-1} yr^{-1})

than in 2015 (2.24 ± 0.24 Mg C ha^{-1} yr^{-1}) and 2017 (2.02 ± 0.27 Mg C ha^{-1} yr^{-1}) (Table 3).

Annual stem C effluxes due to maintenance respiration (AESb) in 2015 (1.47 ± 0.19 Mg C

ha^{-1} yr^{-1}) were lower than in 2016 (1.82 ± 0.16 Mg C ha^{-1} yr^{-1}); AES due to growth

respiration (AESg) was lower in 2016 (0.11 ± 0.04 Mg C ha^{-1} yr^{-1}) than in 2015 (0.42 ± 0.08

Mg C ha^{-1} yr^{-1}) and 2017 (0.34 ± 0.06 Mg C ha^{-1} yr^{-1}). The contribution of AESg to the

annual stem effluxes was 19%, 7% and 16% in 2015, 2016 and 2017, respectively.

The amount of carbon fixed in the stem biomass (SG) was lower in 2016 (0.48 ± 0.13 Mg C

ha^{-1} yr^{-1}) than in 2015 (2.78 ± 0.41 Mg C ha^{-1} y^{-1}) and 2017 (2.50± 0.49 Mg C ha^{-1} yr^{-1}). At

the studied beech forest, the mean C construction cost of wood, defined as the slope of the

relationship between AESg and SG at tree level (R^2 = 0.849, p-value < 0.01, see Supporting

Information Fig.S3), was 0.2 g C emitted per g C fixed. On an annual scale, this parameter

was 0.15 ± 0.01 for 2015, 0.24 ± 0.05 for 2016 and 0.14 ± 0.02 for 2017.

Discussion

Cambial activity and radial growth are not entirely linked to leaf phenology

To the best of our knowledge, this is the first time that the effects of a spring late frost with a

subsequently close summer drought on the wood formation dynamic of beech have been

described.

In all three years, cambium reactivation and wood formation occurred within 1-3 weeks after

leaf development, confirming the tight dependence of radial growth on leaf phenology and

photosynthesis in diffuse-porous species (Čufar *et al.* 2008; Michelot *et al.* 2012).

Nevertheless, in diffuse-porous trees, stem conductivity to water occurs in several outermost growth rings and is not limited to the youngest formed xylem, as found in ring-porous species (Schume, Grabner & Eckmüllner 2004). Hence, in beech allocation to the current year wood, it is not as decisive as in ring-porous species, and newly formed photosynthates at the beginning of the season are preferably used for other crucial processes, such as foliage and fine root growth. At the same site, Matteucci (1998) analysed in parallel net ecosystem exchange (NEE) and carbon allocation to foliage and stem radial growth, finding that the latter started approximately 15 days after photosynthesis exceeded respiration (i.e., NEE was negative). Until then, net absorbed carbon was allocated mostly to foliage growth. This can be related to the C allocation hierarchy, which identifies newly developing leaves as the main C sink at the beginning of the growing season (Campioli *et al.*, 2013; Collalti *et al.*, 2016; Merganičová *et al.*, 2019) Interestingly, and even counter to our expectations, in 2016, the cambium remained active at low rates even after complete canopy defoliation, probably fuelled by old C reserves (D'Andrea *et al.* 2019).

After the second re-sprouting, cambium cell production decreased and became non-productive, although the environmental conditions were potentially still favourable for radial growth, being related to day-length (Rossi *et al.* 2006; Camarero, Olano & Parras 2010). Stem radial growth of beech in Selva Piana was greatly affected by the extreme late spring frost in 2016 because of the premature cessation of cambial cell production and the lower growth rate during the active period, which resulted in 82% narrower annual xylem increments compared to 2015 and 2017. This can be related to a somewhat hypothesised, genetically controlled, form of hierarchy in C allocation (composed by old C reserve and recently fixed photosynthates), which identifies newly developing leaves as the main C sink rather than radial growth. In beech, previously reported growth reduction – as a consequence of late frost – has ranged from 48 to 83% in beech, with the maximum occurring at the northern fringe of

the Alps (Dittmar *et al.* 2006). Radial growth rates had fully recovered in 2017, with no

visible long-term effects of the late spring frost event in 2016, showing the high resilience of

beech growth to late frost (Dittmar *et al.* 2006; Principe, Struwe, Wilmking & Kreyling

2017). However, as recently shown in D'Andrea *et al.* (2019), beech trees during 2016 were

able to compensate the lost C reserve, completely refilling the pool to the same level as before

the frost event. Hence, there was no need to prioritize reserve recharge over stem biomass

production the subsequent year.

The 2017 summer drought became severe only in August (SPEI > 1.5), when the trees had

already completed most of their radial growth, as already seen for other tree species growing

in the Mediterranean adjusting the end xylem growth before potential stressful conditions may

occur (e.g. Lempereur *et al.*, 2015; Forner *et al.*, 2018). Instead, the importance of spring

climatic conditions on beech growth has been reported for the Apennines and the Eastern

Alps (Piovesan, Biondi, Di Filippo, Alessandrini & Maugeri 2008; Di Filippo, Biondi,

Maugeri, Schirone & Piovesan 2012).

Effluxes from stem are not entirely synchronised to radial growth

Mean annual values of Q_{10} ranged between 2.11 (2016) and 2.71 (2015) and were similar to

the values estimated at the same site for co-dominant (2.59) and dominant (2.34) trees

(Guidolotti *et al.* 2013). Q_{10nw} and Q_{10w} estimated in this study are closely comparable with

the dataset of various coniferous and broadleaf tree species reported in Damesin et al. (2002).

Similar intra-annual variability of Q_{10} has been observed in many studies on different species

with higher Q_{10} during the growing period (Paembonan, Hagihara & Hozumi 1992; Carey,

Callaway & DeLucia 1997; Stockfors & Linder 1998; Gruber *et al.* 2009). However, other

studies found (or assume) stable Q_{10} throughout the year (Ceschia *et al.* 2002; Damesin *et al.*

2002).

Our results suggests that wood formation affects Q_{10}, indeed Q_{10w} was higher than Q_{10nw},

despite higher temperatures during the wood formation period, exactly contrary to a case of

sole control of temperature on the parameters (Atkin & Tjoelker 2003).

As reported in other studies, ES15, stem CO_2 efflux at an air temperature of 15°C, was

sensitive to wood formation processes, showing an increase during the growing period

(Ceschia *et al.* 2002; Damesin *et al.* 2002).

Maximum xylem production and maximum ES were not synchronized, while a constant

delay, of about a month, was observed, as similarly found in a young beech forest in which

peak ES occurred *c.* 27 days after the maximum stem growth rate (Ceschia *et al.* 2002).

Furthermore, our results confirmed that peak ES occurred when xylem cells were still in the

phase of wall thickening and lignification, as previously hypothesized (Ceschia *et al.* 2002).

Moreover, when maximum ES was observed, it is very likely that trees were already refilling

the stem C reserves pool (Scartazza, Moscatello, Matteucci, Battistelli & Brugnoli 2013).

Only spring frost affects negatively stem C fluxes

The amount of C fixed by stem biomass formation in 2015 and 2017 ranged from 2.58 to 2.70

Mg C ha^{-1} yr^{-1}, a bit higher than the values reported for a beech forest in Germany, ranging

from 1.69 to 2.41 Mg C ha^{-1} yr^{-1} (Mund *et al.* 2010). In 2016, however, we measured only

0.48 ± 0.13 Mg C ha^{-1} yr^{-1}, i.e., only about 20% of the fixation during the two reference

years, emphasizing how exceptionally negative this year was.

Annual stem CO_2 efflux (AES) is known to be highly variable in temperate forests (Yang *et*

al. 2016). Our data ranges from 1.58 to 2.24 Mg C ha^{-1} yr^{-1} and thus is similar to the 1.65 to

2.25 Mg C ha^{-1} yr^{-1} reported for a younger beech forest (Damesin *et al.* 2002).

An earlier estimate of AES at the study site was 0.63 Mg C ha^{-1} yr^{-1} (Guidolotti *et al.* 2013)

for 2007, year characterized by an extreme summer drought (SPEI > 2). The stem C efflux of

the drought year presented in this study (2017, 1.58 Mg C ha^{-1} yr^{-1}) was about 150% higher

than in 2007, which could be due to an increase in stem biomass (*c*. 15% lower in 2007 than

in 2017, see also Collalti *et al*., 2019) and to different measurement tools. The contribution of

AESg to annual stem effluxes, ranging from 7% to 19%, was lower than that measured in a

young beech forest (Ceschia et al., 2002), evidencing the importance of forest developmental

stage in determining wood formation and growth respiration. The construction cost we found

of 0.23 g C fixed per g C emitted is consistent with the values reported for beech, 0.2 – 0.38 g

C fixed per g C emitted (Ceschia *et al.* 2002; Damesin *et al.* 2002).

While the late frost event in 2016 reduced both wood growth and stem CO_2 efflux with

respect to those measured in the other two years, the percentage of growth reduction (80%)

was much larger than the reduction of ES (25%). Hence, it seems that in mature beech the

contribution of growth respiration to total stem CO_2 fluxes is lower than that of maintenance

respiration, as reported in Collalti *et al.*(2019). In the year of late frost, the strong reduction of

fixed growth C and the contemporary lower reduction of stem CO_2 efflux strongly affected

the overall stem carbon balance. In contrast, the summer drought did not have any effect on

stem growth, and thus neither on C efflux due to growth respiration.

In conclusion, this study further highlights the sensitivity of beech to leaf damage due to late

spring frost. Since leaf development is forecast to start earlier for beeches due to global

warming (Augspurger 2013), the likelihood that spring frost may damage leaves will increase.

We demonstrated that stem growth was significantly reduced due to the prolonged absence of

photosynthesizing leaves after frost, since beech trees tapped their pool of old C reserves.

However, the loss in growth was not completely compensated for after re-growth of leaves,

but rather the cambium activity ceased shortly thereafter. Consequently, the trees fixed less C

in the stem biomass, showing also a reduction of the stem carbon efflux related to growth

respiration.

Moreover, the summer drought occurred too late to affect wood formation and stem CO_2

effluxes. However, more investigations are needed to evaluate its effects on other

physiological processes. This study also underlines the crucial role of spring weather

conditions on the growth and physiology of beech trees.

A better understanding of fine scale C dynamic will help in evaluating a medium- to long-

term response to climate change under an increasing frequency of extreme events.

Acknowledgements

The activities of Negar Rezaie at the wood anatomy laboratory of Slovenian Forestry Institute

were supported by an Excellence Research Award of the National Research Council of Italy,

Department of Biology, Agriculture, and Food Secures (Prot. 71951, 06/11/2017).

Collelongo-Selva Piana is one of the sites of the Italian Long Term Ecological Research

network (LTER-Italy), part of the International LTER network (ILTER). Research at the site

in the years of this study was funded by the eLTER H2020 project (grant agreement no.

654359). Authors are grateful to Martin for English language editing.

Author contribution

533　　E.D'A., N.R., G.M. contributed to the design of the research. Fieldwork was carried out by

534　　E.D'A., N.R.; laboratory analysis for wood formation N.R., J.M., P.P.; data analysis was done

535　　by E.D'A.; data interpretation by E.D'A., N.R., P.P., J.M., A.C., G.M., J.K.. The manuscript

536　　was written by E.D'A and N.R. with major contributions by P.P., J.M., A.C., G.M., J.K..

References

Allevato E., Saulino L., Cesarano G., Chirico G.B., D'Urso G., Falanga Bolognesi S., …

Bonanomi G. (2019) Canopy damage by spring frost in European beech along the

Apennines: effect of latitude, altitude and aspect. *Remote Sensing of Environment* **225**,

431–440.

Atkin O.K. & Tjoelker M.G. (2003) Thermal acclimation and the dynamic response of plant

respiration to temperature. *Trends in Plant Science* **8**, 343–351.

Augspurger C.K. (2013) Reconstructing patterns of temperature , phenology , and frost

damage over 124 years□: Spring damage risk is increasing. *Ecology* **94**, 41–50.

Bascietto M., Bajocco S., Mazzenga F. & Matteucci G. (2018) Assessing spring frost ff ects

on beech forests in Central Apennines from remotely-sensed data. *Agricultural and*

Forest Meteorology **248**, 240–250.

Bascietto M., Cherubini P. & Scarascia-Mugnozza G. (2004) Tree rings from a European

beech forest chronosequence are useful for detecting growth trends and carbon

sequestration. *Canadian Journal of Forest Research* **34**, 481–492.

Begum S., Nakaba S., Oribe Y., Kubo T. & Funada R. (2007) Induction of Cambial

Reactivation by Localized Heating in a Deciduous Hardwood Hybrid Poplar (Populus

sieboldii 3 P . grandidentata). *Annals of botany*, 439–447.

Bloemen J., Agneessens L., Van Meulebroek L., Aubrey D.P., Mcguire M.A., Teskey R.O. &

Steppe K. (2014) Stem girdling affects the quantity of CO2 transported in xylem as well

as CO2 efflux from soil. *New Phytologist* **201**, 897–907.

Bloemen J., McGuire M.A., Aubrey D.P., Teskey R.O. & Steppe K. (2013) Transport of root-

respired CO 2 via the transpiration stream affects aboveground carbon assimilation and

CO 2 efflux in trees. *New Phytologist* **197**, 555–565.

Bowman W.P., Barbour M.M., Turnbull M.H., Tissue D.T., Whitehead D. & Griffin K.L.

(2005) Sap flow rates and sapwood density are critical factors in within- and between-

tree variation in CO2 efflux from stems of mature Dacrydium cupressinum trees. *New*

Phytologist **167**, 815–828.

Camarero J.J., Guerrero-Campo J. & Gutierrez E. (1998) Tree-Ring Growth and Structure of

Pinus uncinata and Pinus sylvestris in the Central Spanish Pyrenees. *Arctic and Alpine*

Research **30**, 1.

Camarero J.J., Olano J.M. & Parras A. (2010) Plastic bimodal xylogenesis in conifers from

continental Mediterranean climates. *New Phytologist* **185**, 471–480.

Campioli M., Verbeeck H., Van den Bossche J., Wu J., Ibrom A., D'Andrea E., … Granier A.

(2013) Can decision rules simulate carbon allocation for years with contrasting and

extreme weather conditions? A case study for three temperate beech forests. *Ecological*

Modelling **263**.

Carey E. V., Callaway R.M. & DeLucia E.H. (1997) Stem respiration of ponderosa pines

grown in contrasting climates: implications for global climate change. *Oecologia* **111**,

19–25.

Carrer M., Brunetti M. & Castagneri D. (2016) The Imprint of Extreme Climate Events in

Century-Long Time Series of Wood Anatomical Traits in High-Elevation Conifers.

Frontiers in Plant Science **7**, 1–12.

Ceschia É., Damesin C., Lebaube S., Pontailler J.Y. & Dufrêne É. (2002) Spatial and seasonal

variations in stem respiration of beech trees (Fagus sylvatica). *Annals of Forest Science*

59, 801–812.

Chan T., Berninger F., Kolari P., Nikinmaa E. & Hölttä T. (2018) Linking stem growth

respiration to the seasonal course of stem growth and GPP of Scots pine. *Tree*

Physiology **38**, 1356–1370.

Chiti T., Papale D., Smith P., Dalmonech D., Matteucci G., Yeluripati J., … Valentini R.

(2010) Predicting changes in soil organic carbon in mediterranean and alpine forests

during the Kyoto Protocol commitment periods using the CENTURY model. *Soil Use*

and Management **26**, 475–484.

Collalti A., Marconi S., Ibrom A., Trotta C., Anav A., D'Andrea E., … Santini M. (2016)

Validation of 3D-CMCC Forest Ecosystem Model (v . 5 . 1) against eddy covariance

592   data for 10 European forest sites. *Geoscientific Model Development* **9**, 479–504.

593   Collalti A., Tjoelker M.G., Hoch G., Mäkelä A., Guidolotti G., Heskel M., … Prentice I.C.

594   (2019) Plant respiration: Controlled by photosynthesis or biomass? *Global Change*

595   *Biology* **0**.

596   Collalti A., Trotta C., Keenan T.F., Ibrom A., Bond-lamberty B., Grote R., … Reyer C.P.O.

597   (2018) Thinning Can Reduce Losses in Carbon Use Efficiency and Carbon Stocks in

598   Managed Forests Under Warmer Climate. *Journal of Advances in Modeling Earth*

599   *Systems*, 2427–2452.

600   Čufar K., Prislan P., De Luis M. & Gričar J. (2008) Tree-ring variation, wood formation and

601   phenology of beech (Fagus sylvatica) from a representative site in Slovenia, SE Central

602   Europe. *Trees - Structure and Function* **22**, 749–758.

603   Cuny H.E., Rathgeber C.B.K., Frank D., Fonti P., Mäkinen H., Prislan P., … Fournier M.

604   (2015) Woody biomass production lags stem-girth increase by over one month in

605   coniferous forests. *Nature Plants* **1**, 15160.

606   D'Andrea E., Rezaie N., Battistelli A., Kuhlmann I., Matteucci G., Moscatello S., … Muhr J.

607   (2019) Winter's bite : beech trees survive complete defoliation due to spring late-frost

608   damage by mobilizing old C reserves. **224**, 625–631.

609   Damesin C., Ceschia E., Le Goff N., Ottorini J.-M. & Dufrêne E. (2002) Stem and branch

610   respiration of beech: from tree measurements to estimations at the stand level. *New*

611   *Phytologist* **153**, 159–172.

612   Delpierre N., Lireux S., Hartig F., Camarero J.J., Cheaib A., Čufar K., … Rathgeber C.B.K.

613   (2019) Chilling and forcing temperatures interact to predict the onset of wood formation

614   in Northern Hemisphere conifers. *Global Change Biology* **25**, 1089–1105.

615   Deslauriers A., Rossi S., Anfodillo T. & Saracino A. (2008) Cambial phenology, wood

616   formation and temperature thresholds in two contrasting years at high altitude in

southern Italy. *Tree physiology* **28**, 863–871.

Dittmar C., Fricke W. & Elling W. (2006) Impact of late frost events on radial growth of common beech (Fagus sylvatica L .) in Southern Germany. *European Journal of Forest Research* **125**, 249–259.

Di Filippo A., Biondi F., Maugeri M., Schirone B. & Piovesan G. (2012) Bioclimate and growth history affect beech lifespan in the Italian Alps and Apennines. *Global Change Biology* **18**, 960–972.

Forner A., Valladares F., Bonal D., Granier A. & Grossiord C. (2018) Extreme droughts affecting Mediterranean tree species' growth and water-use efficiency: the importance of timing. *Tree physiology*.

Frank D., Reichstein M., Bahn M., Thonicke K., Frank D., Mahecha M.D., … Zscheischler J. (2015) Effects of climate extremes on the terrestrial carbon cycle: Concepts, processes and potential future impacts. *Global Change Biology* **21**, 2861–2880.

Gazol A., Camarero J.J., Colangelo M., de Luis M., Martínez del Castillo E. & Serra-Maluquer X. (2019) Summer drought and spring frost, but not their interaction, constrain European beech and Silver fir growth in their southern distribution limits. *Agricultural and Forest Meteorology* **278**, 107695.

Gordo O. & Sanz J.J. (2010) Impact of climate change on plant phenology in Mediterranean ecosystems. *Global Change Biology* **16**, 1082–1106.

Greco S., Infusino M., De Donato C., Coluzzi R., Imbrenda V., Lanfredi M., … Scalercio S. (2018) Late spring frost in mediterranean beech forests: Extended crown dieback and short-term effects on moth communities. *Forests* **9**, 1–18.

Gričar J., Krže L. & Čufar K. (2009) Number of cells in xylem, phloem and dormant cambium in silver fir (ABIES ALBA), in trees of different vitality. *IAWA Journal,* **30**, 121–133.

Gruber A., Wieser G. & Oberhuber W. (2009) Intra-annual dynamics of stem CO2 efflux in

relation to cambial activity and xylem development in Pinus cembra. *Tree Physiology*

29, 641–649.

Guidolotti G., Rey A., D'Andrea E., Matteucci G. & De Angelis P. (2013) Effect of

environmental variables and stand structure on ecosystem respiration components in a

Mediterranean beech forest. *Tree Physiology* **33**, 960–972.

Hilman B., Muhr J., Trumbore S.E., Kunert N., Carbone M.S., Yuval P., … Angert A. (2019)

Comparison of CO2 and O2 fluxes demonstrate retention of respired CO2 in tree stems

from a range of tree species. *Biogeosciences* **16**, 177–191.

Katayama A., Kume T., Ichihashi R. & Nakagawa M. (2019) Vertical variation in wood CO2

efflux is not uniformly related to height: measurement across various species and sizes of

Bornean tropical rainforest trees. *Tree Physiology* **39**, 1000–1008.

Lempereur M., Martin-stpaul N.K., Damesin C., Joffre R., Ourcival J., Rocheteau A. &

Rambal S. (2015) Growth duration is a better predictor of stem increment than carbon

supply in a Mediterranean oak forest□: implications for assessing forest productivity

under climate change. 579–590.

Linares J.C., Camarero J.J. & Carreira J.A. (2009) Plastic responses of Abies pinsapo

xylogenesis to drought and competition. *Tree Physiology* **29**, 1525–1536.

Matteucci G. (1998) *Bilancio del carbonio in una faggeta dell'Italia Centro-Meridionale:*

determinanti ecofisiologici, integrazione a livello di copertura e simulazione

dell'impatto dei cambiamenti ambientali. (ed Università degli studi di Padova), Padova.

Meir P., Mencuccini M. & Coughlin S.I. (2019) Respiration in wood: integrating across

tissues, functions and scales. *New Phytologist*, 1824–1827.

Merganičová K., Merganič J., Lehtonen A., Vacchiano G., Sever M.Z.O., Augustynczik

A.L.D., … Reyer C.P.O. (2019) Forest carbon allocation modelling under climate

change. *Tree Physiology* **39**, 1937–1960.

Michelot A., Simard S., Rathgeber C., Dufrêne E. & Damesin C. (2012) Comparing the intra-annual wood formation of three European species (Fagus sylvatica, Quercus petraea and Pinus sylvestris) as related to leaf phenology and non-structural carbohydrate dynamics. *Tree Physiology* **32**, 1033–1045.

Mund M., Kutsch W.L., Wirth C., Kahl T., Knohl a, Skomarkova M. V & Schulze E.-D. (2010) The influence of climate and fructification on the inter-annual variability of stem growth and net primary productivity in an old-growth, mixed beech forest. *Tree physiology* **30**, 689–704.

Nolè A., Rita A., Ferrara A.M.S. & Borghetti M. (2018) Effects of a large-scale late spring frost on a beech (Fagus sylvatica L.) dominated Mediterranean mountain forest derived from the spatio-temporal variations of NDVI. *Annals of Forest Science* **75**, 83.

Oladi R., Pourtahmasi K., Eckstein D. & Bräuning A. (2011) Seasonal dynamics of wood formation in Oriental beech (Fagus orientalis Lipsky) along an altitudinal gradient in the Hyrcanian forest, Iran. *Trees - Structure and Function* **25**, 425–433.

Paembonan S.A., Hagihara A. & Hozumi K. (1992) Long-Term respiration in relation to growth and maintenance processes of the aboveground parts of a hinoki forest tree. *Tree Physiology* **10**, 21–31.

Peuke A.D., Schraml C., Hartung W. & Rennenberg H. (2002) Identification of drought-sensitive beech ecotypes by physiological parameters. *New Phytologist* **154**, 373–387.

Piovesan G., Biondi F., Di Filippo A., Alessandrini A. & Maugeri M. (2008) Drought-driven growth reduction in old beech (Fagus sylvatica L.) forests of the central Apennines, Italy. *Global Change Biology* **14**, 1265–1281.

Principe A., Struwe T., Wilmking M. & Kreyling J. (2017) Low resistance but high resilience in growth of a major deciduous forest tree (Fagus sylvatica L .) in response to late spring

frost in southern Germany. *Trees - Structure and Function*, 743–751.

Prislan P., Čufar K., De Luis M. & Gričar J. (2018) Precipitation is not limiting for xylem formation dynamics and vessel development in European beech from two temperate forest sites. *Tree Physiology* **38**, 186–197.

Prislan P., Gričar J., De Luis M., Smith K.T. & Cufar K. (2013) Phenological variation in xylem and phloem formation in Fagus sylvatica from two contrasting sites. *Agricultural and Forest Meteorology*, 142–151.

R Development Core Team (2018) R: A Language and Environment for Statistical Computing.

Rathgeber C.B.K., Rossi S. & Bontemps J.-D. (2011) Cambial activity related to tree size in a mature silver-fir plantation. *Annals of Botany* **108**, 429–438.

Rathgeber C.B.K., Santenoise P. & Cuny H.E. (2018) CAVIAR: An R package for checking, displaying and processing wood-formation-monitoring data. *Tree Physiology* **38**, 1246–1260.

Reyer C.P., Silveyra Gonzalez R., Dolos K., Hartig F., Hauf Y., Noack M., … Frieler K. (2019) The PROFOUND database for evaluating vegetation models and simulating climate impacts on forests. *Earth System Science Data Discussion*, 1–47.

Rezaie N., D'Andrea E., Bräuning A., Matteucci G., Bombi P. & Lauteri M. (2018) Do atmospheric CO2 concentration increase, climate and forest management affect iWUE of common beech? Evidences from carbon isotope analyses in tree rings. *Tree Physiology* **1975**, 1110–1126.

Rita A., Camarero J.J., Nolè A., Borghetti M., Brunetti M., Pergola N., … Ripullone F. (2019) The impact of drought spells on forests depends on site conditions : The case of 2017 summer heat wave in southern Europe. 1–13.

Rossi S., Deslauriers A., Anfodillo T., Morin H., Saracino A., Motta R. & Borghetti M.

(2006) Conifers in cold environments synchronize maximum growth rate of tree-ring

formation with day length. *New Phytologist* **170**, 301–310.

Rossi S., Menardi R., Fontanella F. & Anfodillo T. (2005) Campionatore Trephor: un nuovo

strumento per l'analisi della xilogenesi in specie legnose. *Dendronatura* **1**, 60–67.

Saveyn A., Steppe K., Mc Guire M.A., Lemeur R. & Teskey Æ.R.O. (2008) Stem respiration

and carbon dioxide efflux of young Populus deltoides trees in relation to temperature and

xylem carbon dioxide concentration. *Oecologia* **154**, 637–649.

Scarascia-Mugnozza G., Bauer G.A., Persson H., Matteucci G. & Masci A. (2000) Tree

Biomass, Growth and Nutrient Pools. In *Carbon and Nitrogen Cycling in European*

Forest Ecosystems, Ecological. (ed E.-D. Schulze), pp. 49–62. Springer Verlag.

Scartazza A., Moscatello S., Matteucci G., Battistelli A. & Brugnoli E. (2013) Seasonal and

inter-annual dynamics of growth, non-structural carbohydrates and C stable isotopes in a

Mediterranean beech forest. *Tree physiology* **33**, 730–42.

Schröter D., Cramer W., Leemans R., Prentice I.C., Araújo M.B., Arnell N.W., … Zierl B.

(2005) Ecosystem service supply and vulnerability to global change in Europe. *Science*

(New York, N.Y.) **310**, 1333–7.

Schume H., Grabner M. & Eckmüllner O. (2004) The influence of an altered groundwater

regime on vessel properties of hybrid poplar. *Trees* **18**, 184–194.

Stockfors J. & Linder S. (1998) Effect of nitrogen on the seasonal course of growth and

maintenance respiration in stems of Norway spruce trees. *Tree Physiology* **18**, 155–166.

Teskey R.O., Saveyn A., Steppe K. & McGuire M.A. (2008) Origin, fate and significance of

CO_2 in tree stems. *New Phytologist* **177**, 17–32.

Trumbore S.E., Angert A., Kunert N., Muhr J. & Chambers J.Q. (2013) What's the flux?

Unraveling how CO2 fluxes from trees reflect underlying physiological processes. *New*

Phytologist **197**, 353–355.

Ubierna N., Kumar A.S., Cernusak L.A., Pangle R.E., Gag P.J. & Marshall J.D. (2009)

Storage and transpiration have negligible effects on $\delta 13C$ of stem CO2 efflux in large

conifer trees. *Tree Physiology* **29**, 1563–1574.

Vicente-Serrano S.M., Gouveia C., Camarero J.J., Beguería S., Trigo R., López-Moreno J.I.,

… Sanchez-Lorenzo A. (2013) Response of vegetation to drought time-scales across

global land biomes. *Proceedings of the National Academy of Sciences of the United*

States of America **110**, 52–57.

Vitasse Y., Schneider L., Rixen C., Christen D. & Rebetez M. (2018) Increase in the risk of

exposure of forest and fruit trees to spring frosts at higher elevations in Switzerland over

the last four decades. *Agricultural and Forest Meteorology* **248**, 60–69.

Yang J., He Y., Aubrey D.P., Zhuang Q. & Teskey R.O. (2016) Global patterns and

predictors of stem CO 2 efflux in forest ecosystems. *Global Change Biology* **22**, 1433–

1444.

Zhang X., Friedl M. a., Schaaf C.B., Strahler A.H., Hodges J.C.F., Gao F., … Huete A.

(2003) Monitoring vegetation phenology using MODIS. *Remote Sensing of Environment*

84, 471–475.

Figure

Fig. 1: 3-month Standardized Precipitation Evapotranspiration Index (SPEI) at the

experimental site of Selva Piana- Collelongo.

Fig. 2: Number of cambium cells (N. cells vs. leaf area index (LAI, $m^2 m^{-2}$)). Grey points and

dashed lines are the MODIS-LAI values and modelled intra-annual dynamic of Selva Piana

beech forest, respectively. Green and blue vertical lines represent the green-up and maturity

phases of leaf phenology, respectively. The red vertical line represents the late frost of 25[th] April

2016. Black triangles are the average number of cambial cells of five beech trees. Bars are the

standard error.

Fig. 3: Intra-annual dynamics of xylem formation (µm) in 2015 (red dots and solid line), 2016 (green dots and solid line) and 2017 (blue dots and solid line). Gompertz functions were fitted to the total xylem increment comprised of enlarging, wall thickening and mature cells. Each point is the mean of 5 sampled *Fagus sylvatica* trees and bars are standard errors. *** *p*-value < 0.001

Fig. 4: Critical dates and duration of wood formation phases. Different letters (a, b and c) represent significant differences among the beginning of the enlargement phase (grey dots), the

beginning of the wall thickening phase (black dots), cessation of the enlargement phase (grey

triangle) and cessation of the wall thickening phase (black triangle) (*p*-value < 0.05). Grey and

black rectangles represent the duration of the enlargement and wall thickening phases,

respectively. The sum of their length is the duration of wood formation. Each shape is the mean

of 5 sampled trees per year and the bars error represent standard error.

Fig. 5: Top panel: T_{air} (°C) at the measuring time in 2015 (red), 2016 (green) and 2017 (blue).

Bottom panel: Measured stem CO_2 effluxes (μmol CO_2 m^{-2} s^{-1}) in 2015 (red), 2016 (green) and

2017 (blue). Each point is the mean of 5 *Fagus sylvatica* trees. Bars are standard errors.

Fig. 6: Relationship between ES (µmol CO$_2$ m^{-2} s^{-1}) and air temperature (T$_{air}$, °C). Annual, considering the whole measurements for each year, each point is the mean of five sampled trees. Each point represents the mean of those trees, during non-wood and wood formation phases, at a given sampling date. Bars are the standard error. *** *p*-value < 0.001, ** *p*-value < 0.01, * *p*-value < 0.05.

Fig. 7: Relationships between ring widths (mm) and the mean annual ES (μmol CO$_2$ m^{-2} s^{-1})

measured in 2015 (green), 2016 (blue), and 2017 (red). Each point represents a sampled *Fagus*

***sylvatica* tree per year. Bars are the standard error. *** *p*-value < 0.001.**

Fig. 8: Daily increment and stem Carbon effluxes of *Fagus sylvatica*. The black line is the daily increment during 2015, 2016 and 2017, respectively. The red line represents ES, the daily C effluxes. The lower panel, ESb/ES shows the contribution of maintenance respiration to daily stem C efflux.

 Tables

 Table 1: List of terms used in the text

Terms	Definition	Spatial Scale
ES	stem CO_2 efflux per surface area (μmol m^{-2} s^{-1})	Local
ES$_{15w}$	specific CO_2 efflux at an air temperature of 15°C during the wood formation period (μmol m^{-2} s^{-1})	Local
ES$_{15nw}$	specific CO_2 efflux at an air temperature of 15°C during the non-wood formation period (μmol m^{-2} s^{-1})	Local
Q$_{10w}$	ES temperature sensitivity for a 10 °C increase during the wood formation period	Local
Q$_{10nw}$	ES temperature sensitivity for a 10 °C increase during the non-wood formation period	Local
ESw	stem CO_2 efflux per surface area (μmol m^{-2} s^{-1}) during wood formation	Local
ESb	stem CO_2 efflux per surface area (μmol m^{-2} s^{-1}) due to maintenance respiration	Local
ESg	stem CO_2 efflux per surface area (μmol m^{-2}	Local

	s^{-1}) due to growth respiration	
TES	annual stem C efflux (g C yr^{-1})	Tree
TESb	annual stem C efflux due to maintenance respiration (g C yr^{-1})	Tree
TESg	annual stem C efflux due to growth respiration (g C yr^{-1})	Tree
AES	annual stem C efflux (Mg C ha^{-1} yr^{-1})	Stand
AESb	annual stem C efflux due to maintenance respiration (Mg C ha^{-1} yr^{-1})	Stand
AESg	annual stem C efflux due to growth respiration (Mg C ha^{-1} yr^{-1})	Stand
SG	annual C fixed in stem biomass (Mg C ha^{-1} yr^{-1})	Stand

Table 2: Parameters describing the intra annual radial growth derived from the Gompertz

function for the total xylem increment comprised of enlarging, wall thickening and mature cells.

α is the upper asymptote, representing the final ring width at the end of the growing season; tx is

the DOY at which the daily increment is maximum (Gompertz curve inflection point); rx is the

maximum daily increment (μm day^{-1}). Different letters represent significant differences among

the monitored years.

Total xylem increment				
Parameter	Year	Mean (± S.E.)	F	*p*-value
α	2015	1312.17 (± 196.15) a		
α	2016	230.12 (± 31.54) b	13.722	< 0.01
α	2017	1234.80 (± 269.32) a		
rx	2015	25.77 (± 2.95) a		
rx	2016	6.09 (± 2.00) b	8.469	0.014
rx	2017	22.71 (± 3.51) a		
tx	2015	174 (± 2.19) a		
tx	2016	157 (± 5.65) b	22.667	< 0.001
tx	2017	174.80 (± 1.24) a		

829 **Table 3: Annual C stem fluxes. AES is the annual stem C efflux assessed using specific**

830 **parameters for wood formation (Q_{10w} and ES_{15w}) and non-wood formation phases (Q_{10nw} and**

831 **ES_{15nw}); AESb is the annual stem C efflux due to maintenance respiration; AESg is the annual**

832 **stem C efflux due to growth respiration; SG is the annual amount of C fixed in the stem**

833 **biomass; different letters represent significant differences p-value < 0.05**

Year	Flux type	Mean ±(S.E.) Mg C ha^{-1} y^{-1}	F	p-value
2015	AES	2.24±(0.24) a		
2016	AES	1.58±(0.22) b	11.800	0.004
2017	AES	2.02±(0.27) a		
2015	AESb	1.82±(0.16) a		
2016	AESb	1.47±(0.19) b	8.145	0.012
2017	AESb	1.73±(0.18) a/b		
2015	AESg	0.42±(0.08) a		
2016	AESg	0.11±(0.04) b	7.573	0.014
2017	AESg	0.34±(0.06) a		
2015	SG	2.78±(0.41) a		
2016	SG	0.48±(0.13) b	16.534	0.001
2017	SG	2.50±(0.49) a		