Proximity proteomics in a marine diatom reveals a putative cell surface-to-chloroplast iron trafficking pathway

Jernej Turnšek[1,2,3,4,5,6,#], John K. Brunson[6,7], Maria del Pilar Martinez Viedma[8,§], Thomas J. Deerinck[9], Aleš Horák[10,11], Miroslav Oborník[10,11], Vincent A. Bielinski[12], Andrew E. Allen[4,6*]

[1]Biological and Biomedical Sciences, The Graduate School of Arts and Sciences, Harvard University, Cambridge, MA 02138, USA; [2]Department of Systems Biology, Harvard Medical School, Boston, MA 02115, USA; [3]Wyss Institute for Biologically Inspired Engineering, Harvard University, Boston, MA 02115, USA; [4]Integrative Oceanography Division, Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA 92093, USA; [5]Center for Research in Biological Systems, University of California San Diego, La Jolla, CA 92093, USA; [6]Microbial and Environmental Genomics, J. Craig Venter Institute, La Jolla, CA 92037, USA; [7]Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA 92093, USA; [8]Informatics, J. Craig Venter Institute, La Jolla, CA 92037, USA; [9]National Center for Microscopy and Imaging Research, University of California San Diego, La Jolla, CA 92093, USA; [10]Biology Centre CAS, Institute of Parasitology, Branišovská 31, 370 05 České Budějovice, Czech Republic; [11]University of South Bohemia, Faculty of Science, Branišovská 31, 370 05 České Budějovice, Czech Republic; [12]Synthetic Biology and Bioenergy, J. Craig Venter Institute, La Jolla, CA 92037, USA; #Present affiliations: Department of Plant & Microbial Biology, University of California, Berkeley, Berkeley, CA 94720, USA; Howard Hughes Medical Institue, 4000 Jones Bridge Rd, Chevy Chase, MD 20815; §Present affiliation: PHARMA MAR, S.A., Avda. de los Reyes, 1, Pol. Ind. La Mina, 28770 Colmenar Viejo, Madrid, Spain

*For correspondence: aallen@ucsd.edu & aallen@jcvi.org (A.E.A.)

Abstract

Iron is a biochemically critical metal cofactor in enzymes involved in photosynthesis, cellular respiration, nitrate assimilation, nitrogen fixation, and reactive oxygen species defense. Marine microeukaryotes have evolved a phytotransferrin-based iron uptake system to cope with iron scarcity, a major factor limiting primary productivity in the global ocean. Diatom phytotransferrin is endocytosed, however proteins downstream of this environmentally ubiquitous iron receptor are unknown. We applied engineered ascorbate peroxidase APEX2-based subcellular proteomics to catalog proximal proteins of phytotransferrin in the model marine diatom *Phaeodactylum tricornutum*. Proteins encoded by poorly characterized iron-sensitive genes were identified including three that are expressed from a chromosomal gene cluster. Two of them showed unambiguous colocalization with phytotransferrin adjacent to the chloroplast. Further phylogenetic, domain, and biochemical analyses suggest their involvement in intracellular iron processing. Proximity proteomics holds enormous potential to glean new insights into iron acquisition pathways and beyond in these evolutionarily, ecologically, and biotechnologically important microalgae.

Keywords

ocean, iron cycle, diatoms, *Phaeodactylum tricornutum*, phytotransferrin, endocytosis, metal trafficking, chloroplast, proximity-dependent proteomic mapping, APEX2

Glossary

46	AA/AAs	amino acid residue/amino acid residues
47	APEX	engineered ascorbate peroxidase (APX)
48	CREG	cellular repressor of E1A-stimulated genes
49	DAB	3,3'-diaminobenzidine
50	EYFP	enhanced yellow fluorescent protein
51	Fe'	dissolved labile iron (all unchelated iron species)
52	FMN	flavin mononucleotide
53	HNLC	high-nutrient, low-chlorophyll
54	ISIP	iron starvation induced protein
55	MS	mass spectrometry
56	PBS	phosphate-buffered saline
57	pTF	phytotransferrin
58	RT	room temperature
59	TEM	transmission electron microscopy
60	TM	transmembrane
61	TMT	tandem mass tag
62	UTR	untranslated region
63	V-ATPase	vacuolar-type H^+-ATPase
64	WT	wild type

Introduction

Iron (Fe) likely played an important role in the origin of life (*Bonfio et al., 2017*; *Jin et al., 2018*) and is fundamental to almost all extant metabolisms (*Aguirre et al., 2012*) serving as a cofactor in enzymes involved in DNA replication, photosynthesis, cellular respiration, nitrate assimilation, nitrogen fixation, and reactive oxygen species defense (*Crichton, 2016*). Early anoxic oceans were rich in readily bioavailable ferrous, Fe^{2+}, iron, but with the rise of oxygenic photosynthesis and the Great Oxygenation Event (GOE) in the Paleoproterozoic ~2.3 billion years ago followed by the Neoproterozoic Oxygenation Event (NOE) ~1.5 billion years later, most of the ferrous iron oxidized into insoluble ferric, Fe^{3+}, minerals which are not bioavailable (*Camacho et al., 2017*; *Knoll et al., 2017*; *Och and Shields-Zhou, 2012*). Conceivably, these large global shifts in ocean chemistry could have had a major role in driving the evolution of novel iron uptake mechanisms.

Iron availability limits primary productivity in high nutrient, low chlorophyll (HNLC) regions which cover ~25% of the modern ocean habitat as demonstrated by numerous large-scale Fe fertilization experiments and natural Fe upwelling events invariably resulting in diatom-dominated phytoplankton blooms (*Ardyna et al., 2019*; *de Baar et al., 2005*; *Martin et al., 1994*). Although dissolved Fe in marine environments is primarily found complexed to organic ligands (*Hutchins and Boyd, 2016*; *Tagliabue et al., 2018*) such as bacterially produced siderophores and hemes (*Boiteau et al., 2016*; *Hogle et al., 2014*), low (pM) amounts of unchelated labile Fe, Fe', are crucial for eukaryotic phytoplankton, particularly diatoms (*Morel et al., 2008*), widespread single-celled microalgae critical to oceanic primary production (*Benoiston et al., 2017*). Diatoms have convergently evolved phytotransferrin (pTF), which serves as the basis of a high-affinity ferric, Fe^{3+}, iron binding and acquisition pathway (*McQuaid et al., 2018*; *Morrissey et al., 2015*). Phytotransferrin sequences have a broad taxonomic distribution and are abundant in marine transcriptomic datasets (*Bertrand et al., 2015*; *Marchetti et al., 2012*).

Phytotransferrins are estimated to have emerged concurrently with the NOE exemplifying the link between large environmental changes and molecular innovation (*McQuaid et al., 2018*). They are phylogenetically related to Fe-assimilation (FEA) domain-containing proteins found in freshwater and marine single-celled algae, and represent functional analogues of iron delivery proteins transferrins (*Behnke and LaRoche, 2020*; *Cheng et al., 2004*; *McQuaid et al., 2018*; *Scheiber et al., 2019*). *pTF* (Ensembl ID: Phatr3_J54465, UniProt ID: B7FYL2) from the model marine diatom *Phaeodactylum tricornutum* (*Bowler et al., 2008*) was first identified as an iron

deficiency-sensitive gene and named *ISIP2a* (iron starvation induced protein 2a) (*Allen et al.,*

2008), although its expression is relatively high in iron-replete conditions as well (*Smith et al.,*

2016). pTF localizes to the cell surface and intracellular puncta, presumably endosomal vesicles

(*McQuaid et al., 2018*), which is further supported by the presence of an endocytosis motif in the

C-terminus of the protein (*Lommer et al., 2012*). In diatom pTF, dissolved labile ferric iron is

coordinated synergistically with carbonate anion (CO_3^{2-}), a common feature of all transferrins

(*Cheng et al., 2004*; *McQuaid et al., 2018*; *Zak et al.*, 2002). Thus, decline in seawater carbonate

concentrations due to ocean acidification caused by elevated atmospheric CO_2 may negatively

impact this prevalent iron uptake system in diatoms and other marine phytoplankton (*McQuaid*

et al., 2018).

While accessing and binding dilute dissolved labile ferric iron on the cell surface is important, its

subsequent internalization and delivery to target sites within complex cellular milieu are also

critical. Fe^{3+} is highly insoluble and conversion between Fe^{3+} and Fe^{2+} can lead to toxic reactive

oxygen species causing damage to proteins, lipids, and nucleic acids (*Cheng et al., 2004*).

Precise control of iron internalization and intracellular trafficking in either of its redox states is thus

crucial for maintenance of cellular homeostasis (*Philpott and Jadhav, 2019*; *Wang and*

Pantopoulos, 2011). Human ferric iron-laden transferrin (Tf) bound to transferrin receptor is

internalized via clathrin-mediated endocytosis. Endosome acidification leads to carbonate

protonation and Tf conformational change resulting in iron release. Fe^{3+} is then reduced by six-

transmembrane epithelial antigen of prostate 3 (STEAP3), exported to the cytoplasm through

divalent metal transporter 1 (DMT1; also NRAMP2) or Zrt- and Irt-like protein (ZIP) 14, and

offloaded to iron chaperones for distribution to cellular iron sinks (*Bogdan et al., 2016*; *Eckenroth*

et al., 2011; *Ohgami et al., 2005*; *Philpott and Jadhav, 2019*; *Wang and Pantopoulos, 2011*).

In contrast, proteins that mediate intracellular allocation of internalized Fe^{3+} and conduct specific

biochemical transformations downstream of diatom pTF are unknown. Elucidation of these

pathways will advance our understanding of diatom adaptation to survival in iron-deficient ocean

waters and provide insight into how climate change will impact diatom physiology and fitness.

To investigate the proximal proteomic neighborhood of *P. tricornutum* pTF, we employed APEX2,

an engineered soybean ascorbate peroxidase that functions as a dual probe for electron

microscopy and proximity proteomics (*Hung et al., 2016*; *Lam et al., 2015*; *Martell et al., 2017*).

When fused with a protein of interest (bait protein), the ~27 kDa APEX2 enzyme permits spatially

resolved proteomic mapping by oxidizing biotin-phenol to short lived (<1 ms) phenoxyl radicals

which can covalently react with electron-rich amino acid residues—primarily tyrosine—on the surface of nearby endogenous proteins. Tagged proteins can then be isolated by purification with streptavidin beads and analyzed using mass spectrometry (MS). While the "biotinylation radius" in APEX2 experiments is estimated to be in the 10–20 nm range, it should rather be seen as a "probability gradient" where the likelihood of tagging decreases with distance away from the APEX2-tagged protein of interest (*Gingras et al., 2019*; *Hung et al., 2016*; *Lam et al., 2015*). Proteomic hits can therefore include vicinal proteins which interact with the APEX2-tagged bait protein indirectly in addition to direct interactors of varying strength (*Lundberg and Borner, 2019*). Tha ability to capture weakly and transiently interacting proteins in a biological process of interest is a major advantage APEX2 offers over traditional pull-down approaches (*Gingras et al., 2019*). Swaping biotin-phenol for diaminobendizine (DAB)—another APEX2 substrate— enables one to determine fusion protein localization via high-resolution electron microscopy making APEX2 a powerful bifunctional probe (*Martell et al., 2017*; *Martell et al., 2012*).

APEX2 and related enzyme-based chemical biology tools have over the past decade been applied to investigate cellular compartments and processes in diverse model systems including mammalian cell culture, yeast, protists, and plant protoplasts (*Gingras et al., 2019*; *Trinkle, 2019*). APEX2 was recently used to map proteomes associated with post-Golgi vesicle trafficking and endocytosis (*Del Olmo et al., 2019*; *Otsuka et al., 2019*) as well as receptor-mediated signaling (*Paek et al., 2017*), successes that are all directly pertinent to the goals of our study.

Here, we provide further evidence for *P. tricornutum* phytotransferrin pTF endocytosis, demonstrate correct subcellular localization and activity of its APEX2 fusion, and identify nearly 40 likely proximal proteins through a proximity-dependent proteomic mapping experiment conducted in quintuplicate. To the best of our knowledge, this study represents the first application of APEX2 in a marine microbial model system (*Figure 1*). We then provide initial characterization, using dual fluorophore protein tagging fluorescence microscopy, phylogenetic, domain, and biochemical analyses, for three proteins expressed from a chromosomal gene cluster (pTF.CREG1, pTF.CatCh1, pTF.ap1) believed to act in association with endocytosed pTF downstream of ferric iron binding at the cell surface. Bacterially expressed recombinant pTF.CREG1 was found to be catalytically dead indicating it may have a regulatory, rather than enzymatic, role in pTF endocytosis, similar to functions proposed for human CREG1. Meanwhile, bioinformatic interrogation of pTF.CatCh1 and its localization pattern suggest it might be a

166   chloroplast-associated iron-binding protein. Finally, we introduce a model for an intracellular ferric

167   iron allocation pathway in *P. tricornutum* based on our proximity proteomics hits.

 Results

P. tricornutum **phytotransferrin (pTF) is localized to intracellular vesicles.**

To confirm pTF association with intracellular vesicles as suggested by *McQuaid et al., 2018*, a
pTF-mCherry encoding episome was conjugated into *ΔpTF P. tricornutum* cells (*Diner et al.,*
2016; *Karas et al., 2015*; *McQuaid et al., 2018*). After labeling the fluorescent transconjugant
cell line with 100 μM of the membrane dye MDY-64 for 10 min at room temperature, colocalized
mCherry and MDY-64 signals were observed on the cell surface and within intracellular vesicles
(*Figure 2A*).

APEX2 fused with *P. tricornutum* phytotransferrin (pTF) is enzymatically active *in vivo*.

To generate pTF-APEX2 expressing diatom cell lines, an episome encoding pTF with C-terminal
APEX2 was assembled and conjugated into WT *P. tricornutum* cells (*Figure 2B*). Considering the
predicted pTF domains (*Figure 2—figure supplement 1A*), APEX2 was likely facing the cytosol
at the cell surface and once internalized into vesicles. Western blot with pTF-specific antibodies
(*McQuaid et al., 2018*) confirmed the ~84.7 kDa fusion protein expression in four out of five tested
transconjugant cell lines. The protein was present in the insoluble cell lysate fraction further
indicative of its membrane localization (*Figure 2C*). Amplex UltraRed, a highly sensitive APEX2
substrate (*Hung et al., 2016*), was used to assay live pTF-APEX2 expressing cells. A resorufin
(reaction product of Amplex UltraRed and APEX2) signal up to 4- and 40-fold above WT
background was observed in experiments performed at room temperature (data not shown) and
on ice (*Figure 2D*; *Figure 2—source data 1*), respectively, indicating active APEX2 with
incorporated heme. Resorufin was also directly visualized by confocal microscopy and a strong
cytosolic signal not tightly localized to the expected site of origin, similar to previous reports
(*Martell et al., 2012*), was observed (*Figure 2—figure supplement 1B*), perhaps indirectly
supporting our predicted APEX2 orientation. APEX2 was active only in the presence of both
Amplex UltraRed and hydrogen peroxide (*Figure 2—figure supplement 1C*) implying that
endogenous H_2O_2 levels are not sufficient to drive APEX2-catalyzed reactions, and that the
overall cell surface and intracellular milieu in *P. tricornutum* is permissive to the APEX2 catalytic
cycle.

pTF-APEX2 is localized to the cell membrane and intracellular vesicles.

To confirm pTF-APEX2 is localized to the cell surface and intracellular vesicles similar to the pTF-mCherry fusion protein, pTF-APEX2 expressing cells (cell line s2, *Figure 2C–D*) were treated with 25 mM 3,3'-diaminobenzidine (DAB) in the presence of 3 mM H_2O_2 for 15 min on ice (*Figure 3A*). This reaction leads to DAB polymerization and local precipitation around APEX2 that can be stained with osmium tetroxide and visualized with an electron microscope. Cells were embedded in a 3% agar matrix to prevent losses during numerous washing steps (*Figure 3—figure supplement 1A*). Tightly localized signal was observed on the cell membrane and in intracellular vesicles indicating that pTF-APEX2 fusion trafficked to the correct subcellular sites (*Figure 3B*). Additionally, we observed mitochondrial signal in WT and pTF-APEX2 cell lines subjected to the DAB reaction (*Figure 3B*; *Figure 3—figure supplement 1B*). Analysis of the *P. tricornutum* proteome revealed eight peroxidases with Arg38, His42, His163, and Asp208; catalytic amino acid residues that are conserved across all identified ascorbate peroxidases (APX) including soybean APX (*Raven et al., 2003*) and its derivative APEX2 (*Figure 3—figure supplement 1C*). Three of them contain proline in place of alanine at position 134 relative to APEX2, a substrate-binding loop mutation rendering APEX2 highly active (*Lam et al., 2015*), and another one is predicted to localize to mitochondria (*Figure 3—figure supplement 1D–E*). It is plausible that one or more of these endogenous APEX2-like peroxidases are responsible for the observed mitochondrial signal. APEX2 was inactivated in iron-deplete conditions (40 nM total Fe) and could be reactivated by supplementing fixed cells with 10 μM hemin chloride for 3 hrs (data not shown), but this substantially increased background in the Amplex UltraRed assay. Therefore, subsequent proximity labeling experiments were carried out on cells growing in iron-replete conditions.

Identification of the proximal phytotransferrin proteome with biotin-phenol labeling.

To identify proteins proximal to pTF, quintuplicate cultures from one WT and one pTF-APEX2 expressing cell line (cell line s2, *Figure 2C–D*) were grown to mid- to late-exponential phase and supplemented with 2.5 mM biotin-phenol and 1 mM hydrogen peroxide, following the APEX2 proximity labeling protocol developed for yeast (*Hwang and Espenshade, 2016*; *Hwang et al., 2016*). These labeling reaction steps were performed at 4°C and the incubation with hydrogen peroxide was extended to 20 min to mirror Amplex UltraRed assay and DAB reaction conditions (*Figure 4A*). Increasing biotin-phenol concentration from 0.5 mM, usually used in mammalian cells, to 2.5 mM, and exposing cells to osmotic stress with 1.2 M sorbitol—both of which are critical for efficient labeling in yeast (*Hwang and Espenshade, 2016*)—was necessary to detect enrichment of biotinylated proteins in experimental samples (*Figure 4B*). This result confirmed our hypothesis that the lack of heavy silicification in *P. tricornutum* (*Francius et al., 2009*; *Tesson et al., 2009*), making its cell wall composition and cell membrane permeability likely similar to that of yeast, would permit labeling. Streptavidin pull-downs were then performed with clarified cell lysates followed by tandem mass tag (TMT)-based quantitative proteomics. WT and pTF-APEX2 proteomic replicates, with the exception of one WT and one pTF-APEX2 sample, formed two distinct clusters (*Figure 4—figure supplement 1*; *Figure 4—figure supplement 1—source data 1*) indicating minimal technical variability. Thirty-eight statistically significant proteins (P value ≤0.05) with APEX2/WT ratios of at least 1.5 were identified (*Figure 4B*; *Figure 4—source data 1*). These ratios were obtained from average total peptide counts across quintuplicates (*Figure 4—source data 1*). Endogenous biotinylated proteins were also detected and had APEX2/WT ratios close to one, thus acting as an intrinsic pull-down control (*Supplemental file 1—Table S1*; *Figure 4—source data 1*). Some background enrichment in WT cells was likely due to endogenous APEX2-like peroxidases and one would therefore expect mitochondrial proteins with APEX2/WT ratios close to one to be present in our MS dataset. Indeed, at least three were detected: mitochondrial chaperonin CPN60 (Phatr3_J24820, UniProt ID: B7FQ72, APEX2/WT = 0.99), mitochondrial import receptor subunit TOM70 (Phatr3_J47492, UniProt ID: B7G3J4, APEX2/WT = 0.91), and acetyl-CoA dehydrogenase (Phatr3_J11014, UniProt ID: B7FTR6, APEX2/WT = 1.23) (*Figure 4—source data 1*). Fourteen proteins with an APEX2/WT ratio of at least two were detected. Of these 14 proteins, nine are known to be transcriptionally sensitive to iron availability; this is also the case for an additional five proteins with APEX2/WT ratio of at least 1.5 (*Figure 5—figure supplement 1—source data 1*; *Smith et al., 2016*). We note that pTF was present, but not enriched, in our pTF-APEX2 proteomics samples (*Figure 4—source data 1*).

One possible explanation for this result is that pTF may not have surface exposed amino acid residues that are permissive to biotinylation by APEX2-generated phenoxyl radicals and so its presence in all of our pull-down samples could be due to unspecific binding to streptavidin beads.

Some of the most apparent biologically interesting hits are summarized in *Figure 4C*. Considering iron is critically involved in all of the major photosynthetic complexes—photosystem II, photosystem I, and cytochrome *b6f* (*Rochaix, 2011*)—we asked if some of the proteins are perhaps predicted to be targeted to the chloroplast. Indeed, five are, albeit with low confidence, by ASAFind (*Gruber et al., 2015*), including two (Phatr3_J51183 and Phatr3_J54986) that are part of a gene cluster on chromosome 20 and two (Phatr3_J41423 and Phatr3_J55031) that are known to associate with the *P. tricornutum* chloroplast (*Allen et al., 2012*; *Kazamia et al., 2019*; *Kazamia et al., 2018*; *Figure 4C*). We elaborate on the potential role for these proteins in intracellular iron trafficking in the Discussion.

Proteins encoded by an iron-sensitive gene cluster on chromosome 20 colocalize with pTF.

We focused on three proteins with APEX2/WT ratio of at least two—Phatr3 IDs: J51183 (hereafter pTF.CREG1), J52498 (hereafter pTF.CatCh1), and J54986 (hereafter pTF.ap1)—that are expressed from a previously identified iron- and silicon-sensitive gene cluster on chromosome 20 (*Allen et al., 2008*; *Sapriel et al., 2009*; *Figure 5A*). One additional protein, ISIP2b (Phatr3_J54987, UniProt ID: B7G9B1), which is also transcriptionally sensitive to iron and silicon and that we did not detect in our proximity proteomics experiment, is also encoded by this uncharacterized locus. We note that *pTF* is not co-located with these genes, but instead lies on chromosome 7 (genomic location 1,000,053–1,001,833, forward strand). All three genes exhibit a transcriptional profile similar to *pTF* (*Figure 5—figure supplement 1*), two proteins (pTF.CREG1 and pTF.ap1) are predicted to go to the chloroplast (with low confidence as determined by ASAFind), and two (pTF.CatCh1 and pTF.ap1) contain a C-terminal transmembrane domain (*Figure 4C*; *Supplemental file 1—Table S2*). To test whether they colocalize with pTF, co-expression episomes were assembled and conjugated into WT *P. tricornutum* cells which resulted in diatom cell lines expressing pTF-mCherry and MS hit-EYFP fusion proteins (*Figure 5—figure supplement 2A–B*). Imaging conditions were optimized with mCherry and Venus (yellow fluorescent protein with spectral properties very similar to those of EYFP) expressing *P. tricornutum* cell lines for minimal cross-channel bleed-through (*Figure 5—figure supplement 2C*). pTF.CREG1 and pTF.CatCh1 colocalized with pTF in the chloroplast vicinity and on the chloroplast margin, respectively (*Figure 5B*). Both proteins consistently exhibited these distinct colocalization patterns as demonstrated by additional three mCherry- and EYFP-positive cells from corresponding co-expression transconjugant cell lines (*Figure 5—figure supplement 3A–B*). Colocalization of pTF.ap1 with pTF close to the chloroplast was also evident, although somewhat less precise (*Figure 5—figure supplement 4*).

pTF.CREG1 has homologs across the tree of life and may possess a non-enzymatic role similar to human CREG1.

To shed light on possible functions of the uncharacterized proteins colocalizing with pTF and to examine their occurrence in other diatoms and marine phytoplankton beyond *Phaeodactylum tricornutum*, phylogenetic analysis was performed. pTF.CREG1 homologs were identified across the tree of life (*Figure 6*; *Figure 6—source data 1*). Further investigation of the protein sequence alignment underlying the phylogenetic tree revealed 12 amino acid residues that are at least 90% conserved across all homologs (*Figure 6—figure supplement 1*). Diatom homologs can be seen in a crown group with proteins from other complex plastid-containing algae such as cryptophytes, haptophytes, pelagophytes, chlorarachniophytes, and dinoflagellates (*Figure 6*; *Archibald, 2009*; *Oborník and Füssy, 2018*). pTF.CREG1 and its iron-insensitive paralog Phatr3_J10972—which phylogenetically cluster together—have ~50% amino acid sequence similarity and belong to the flavin mononucleotide (FMN)-binding split barrel fold protein superfamily (SSF50475) according to their Ensembl Protists profiles.

We analysed pTF.CREG1 amino acid sequence (UniProt ID: B7G9B3, 234 AAs, ~26.4 kDa) and compared it to the human CREG1 ortholog (UniProt ID: O75629, 220 AAs, ~24 kDa). While human CREG1 is evolutionarily and structurally closely related to flavin mononucleotide (FMN)-coordinating dimeric oxidases and oxidoreductases such as pyridoxine 5'-phosphate (PNP) oxidases, it is unable to bind FMN and catalyze NADPH oxidation (*Ghobrial et al., 2018*; *Sacher et al., 2005*). pTF.CREG1 has a predicted N-terminal signal peptide ending in SGA that may be recognized by an endoplasmic reticulum-localized signal peptidase (*Benham, 2012*; *Tuteja, 2005*), and is likely modified by both N- and O-glycosylation, in line with human CREG1 (*Figure 6—figure supplement 2A*; *Sacher et al., 2005*). These features suggest that pTF.CREG1 enters and travels along the secretory pathway (*Bard and Chia, 2016*; *Benham, 2012*). Furthermore, the amino acid sequence alignment between pTF.CREG1 and human CREG1 revealed a similar tetrapeptide motif DP(Q/E)S which is located on the loop occluding the FMN-binding pocket at the human CREG1 dimer interface (*Figure 6—figure supplement 1*; *Figure 6—figure supplement 2A*; *Sacher et al, 2005*). This short amino acid sequence—while thought to contribute to the lack of catalytic activity in human CREG1—was found to be critical for mediating cell growth inhibition in human teratocarcinoma cell line NTERA-2 (*Sacher et al., 2005*). Structural comparison between a pTF.CREG1 homology model created using Phyre2 web portal (*Kelley et al., 2015*) and the human CREG1 monomer (*Sacher et al., 2005*; PDB ID: 1XHN) supports our hypothesis that pTF.CREG1 also contains a loop preventing FMN binding (*Figure*

6—figure supplement 2B). Two arginine amino acid residues involved in binding the terminal

FMN phosphate in *S. cerevisiae* pyridoxine 5′-phosphate oxidase (UniProt ID: P38075, PDB ID:

1CI0) are mutated to aspartic acid and methionine amino acid residues in human CREG1 (*Sacher*

et al., 2005), and one of these two key arginine residues is absent from pTF.CREG1 (*Figure 6—*

figure supplement 3A). Finally, we note that a putative homodimerization region in pTF.CREG1

homology model contains many hydrophobic amino acid residues and overall follows the

compositional character of the human CREG1 dimerization interface suggesting pTF.CREG1 is

acting as a dimer *in vivo* (*Figure 6—figure supplement 3B*).

Given the apparent structural similarity of the *P. tricornutum* pTF.CREG1 to the catalytically

inactive human ortholog, we opted to test its activity, or lack thereof, to delineate it from the

evolutionarily related FMN-binding enzymes. Various truncations of His6-tagged pTF.CREG1

were expressed in *Escherichia coli* and purified using cobalt, Co^{2+}, immobilized metal affinity

chromatography (IMAC) to test solubility. A construct encoding pTF.CREG1 without the 31 N-

terminal amino acid residues and leading to a soluble protein was selected for larger-scale

purification (*Figure 6—figure supplement 4A*). The Co^{2+}-bound protein fraction (*Figure 6—*

figure supplement 4B) exhibited no flavin reductase activity as measured by the oxidation of

NADPH in the presence of potential flavin substrates flavin mononucleotide (FMN) and riboflavin

(*Figure 6—figure supplement 2C*; *Figure 6—figure supplement 2—source data 1*).

In conclusion, our homology modeling analyses and enzymatic assays indicate that pTF.CREG1

does not bind FMN and does not reduce flavins in an NADPH-dependent manner, respectively,

both in agreement with our current knowledge about human CREG1. As such, pTF.CREG1 is an

enticing candidate for further study as a key participant in the endocytic pTF receptor-mediated

iron uptake pathway in diatoms. We put forth a hypothesis for its function in the Discussion.

pTF.CatCh1 is a heterokont-specific protein with conserved CX(X)C motifs.

In contrast to pTF.CREG1, pTF.CatCh1 homologs were identified only in diatoms and other single-celled heterokonts (*Figure 7A*; *Figure 7—source data 1*). pTF.CatCh1 is a ~37.4 kDa (349 AAs) large protein with a predicted signal peptide (AAs 1–26) ending in ASA, two N-glycosylation NX(S/T) sequons (i.e., asparagines in the identified NST and NLS motifs may be glycosylated), and 13 predicted O-glycosylation sites (*Rao and Wollenweber, 2010*; data not shown). This protein is paralogous to the iron starvation induced protein 2b (ISIP2b/Phatr3_J54987) whose gene lies adjacently to *pTF.CatCh1* in the same chromosome 20 gene cluster (*Figure 5A*). They share three CXC and one CXXC motifs typical of metal-binding, redox-active, and iron–sulfur cluster-containing proteins (*Figure 7B*; *Blaby-Haas et al., 2014*; *Fomenko and Gladyshev, 2003*; *Poole, 2014*; *Przybyla-Toscano et al., 2018*). One of the CXC motifs and the CXXC motif are conserved across all but two homologs in our phylogenetic analysis (*Figure 7B*; *Figure 7—figure supplement 1A*), and the latter is predicted to be located in one of the two disordered C-terminal regions in pTF.CatCh1 (AAs 196–292, AAs 321–349; *Figure 7—figure supplement 1B*). With additional 12 cysteines conserved between pTF.CatCh1 and ISIP2b, pTF.CatCh1 contains a total of 20 cysteine amino acid residues (all between AAs 34 and 226). Six of these 20 are 100% conserved across all homologs (*Figure 7B*). The predicted transmembrane domain (AAs 296–315) is flanked by a polyserine ($[Ser]_6$) and a short arginine-rich stretch ($RKL[R]_3$). Flexible polyserine linkers are found in modular, multidomain proteins (*Uversky, 2015a*), and positively charged amino acid residue tracts in the vicinity of transmembrane regions were shown to be orientation determinants in outer chloroplast membrane proteins (*May and Soll, 1998*). These observations are synthesized in a domain organization schematic shown in *Figure 7—figure supplement 1C*. We elaborate on possible roles of the conserved CX(X)C motifs, as well as other identified motifs and domains in pTF.CatCh1 in the Discussion.

pTF.ap1 homologs are present in marine microeukaryotes including diatoms.

Phylogenetic characterization was performed with pTF.ap1 as well and numerous homologs with at least five highly conserved motifs were identified in diatoms and other marine microeukaryotes, including chlorophytes, cryptophytes, and haptophytes (*Figure 7—figure supplement 2A–B*; *Figure 7—figure supplement 2—source data 1*). These pTF.ap1 homologs contain five 100% conserved cysteine amino acid residues, two of which are shown in *Figure 7—figure supplement 2B*, as well as 100% conserved tyrosine, histidine, and aspartic acid residues,

389 indicating this protein may also be redox-active and/or have a metal ion processing role

390 (*Dokmanić et al., 2007*).

Discussion

In this study, we used the iron receptor protein phytotransferrin (pTF) to implement APEX2-mediated proximity-dependent proteomic mapping in the model marine diatom *Phaeodactylum tricornutum*. The resulting data provide several advances regarding the function of pTF and its putative protein interaction partners downstream of iron binding at the cell surface.

pTF (ISIP2a; iron starvation induced protein 2a) was identified in *P. tricornutum* as a marker for iron (Fe) limitation (*Allen et al., 2008*). It was subsequently shown to be involved in Fe acquisition (*Morrissey et al., 2015*) and to be widespread in marine and freshwater microeukaryotic phytoplankton communities (*Bertrand et al., 2015*; *Carradec et al., 2018*; *McQuaid et al., 2018*). In a breakthrough study, *McQuaid et al., 2018*, demonstrated that ISIP2a is a type of transferrin, phytotransferrin (pTF), that is crucial for acquisition of dissolved labile ferric, Fe^{3+}, iron, a critical micronutrient for cellular biochemistry. *McQuaid et al., 2018*, showed that pTF is essential for high-affinity iron uptake in *P. tricornutum*, and that carbonate, CO_3^{2-}, and Fe^{3+} interact synergistically to control the iron uptake rate. The finding that carbonate anions are required for the activity of a key diatom iron transport system is noteworthy and suggests that ocean acidification might inhibit microbial iron uptake. The occurrence of transferrin in marine diatoms and other single-celled marine and freshwater phytoplankton raises significant questions concerning its evolutionary origin and cellular role.

McQuaid et al., 2018, observed pTF in intracellular puncta, showed that the clathrin-mediated endocytosis inhibitor Pitstop 2 reduces iron (Fe) uptake rates, visualized vesicles after adding iron to an iron-limited WT *P. tricornutum* culture stained with the membrane dye FM 1-43, but our MDY-64 labeling data provide the first direct evidence that diatom pTF is indeed associated with intracellular membranous compartments. Further work is needed to elucidate pTF dynamic upon cellular internalization—especially after iron addition to iron-deplete cells which mimics oceanic iron repletion events—and its trafficking in and potential exit from the endomembrane system in relation to cell cycle phases and light-dark cycles (*Mayle et al., 2012*; *Naslavsky and Kaplan, 2018*; *Smith et al., 2016*).

While APEX2 was expected to be active in *P. tricornutum* given previous heterologous expression of functional horseradish peroxidase (HRP) in the model marine diatom *Thalassiosira pseudonana* (*Sheppard et al., 2012*), it was unclear whether endogenous peroxidases would

contribute to the background signal in various APEX2 assays. In our Amplex UltraRed assay
experiments, an order of magnitude higher signal-to-noise ratio (i.e., resurofin signal in pTF-
APEX2 expressing versus WT *P. tricornutum* cells) was detected when live cells were reacted on
ice as opposed to on room temperature, suggesting endogenous *P. tricornutum* peroxidases, but
not exogenous APEX2, are largely inhibited on ice. This is notable as it indicates that performing
APEX2 assays on ice may represent a generalizable experimental strategy for mesophilic
microeukaryotic phytoplankton model systems.

Identifying Amplex UltraRed assay conditions which minimized WT background was promising,
but not a guarantee that some endogenous enzymes would not be able to process other APEX2
substrates, such as those crucial for electron microscopy (3,3'-diaminobenzidine/DAB) and
proximity proteomics (biotin-phenol). Indeed, mitochondrial background signal was detected in
our electron microscopy experiments both in WT and pTF-APEX2 expressing cells, and our *P.*
tricornutum proteome analysis revealed candidate endogenous APEX2-like peroxidases that
could explain this result. Nevertheless, C-terminal tagging of pTF with APEX2 resulted in the
expected cell surface and what appeared to be vesicular fusion protein localization away from
mitochondria indicating no spatial overlap between endogenous APEX2-like enzymes and pTF-
APEX2. Inconsistent size and shape of vesicle-like structures observed in our electron
micrographs were conceivably due to a combination of arbitrary cell orientation inside agar blocks
(where diatom cells were embedded) and the exact z position of ultramicrotome cut sites. On the
whole, APEX2-based imaging is complementary to super-resolution microscopy in diatoms and
represents a basis for electron tomography applications (*Gröger et al., 2016*; *Sengupta et al.,*
2019). Some off-target biotinylation in our proximity labeling experiments was likely due to
background mitochondrial peroxidase activity as indicated by the presence of mitochondrial
proteins in our mass spectrometry data. Their APEX2/WT ratios close to one give us more
confidence that those with elevated ratios above 1.5 were enriched due to APEX2 activity. Many
of the latter have functional annotations for proteins we might expect to play a role in endosomal
iron assimilation, and many have previously been shown to have iron-sensitive transcripts, both
of which further support this notion. Future efforts to express pTF tagged with N-terminal APEX2,
which should lead to intravesicular proximity labeling according to our pTF orientation prediction,
optimizing conditions for APEX2 reactivation in iron-deplete conditions with hemin chloride, and
more cell wall-permeant APEX2 probes (*Tian et al., 2020*) will allow for identification of additional
candidate proteins. Considering iron metabolism genes in *P. tricornutum*, including *pTF*, are
highly expressed at night, proteomic maps with improved spatial and temporal resolution may

emerge from proximity-dependent proteomic mapping experiments conducted with synchronized cell lines sampled at different timepoints throughout diel cycle (*Huysman et al., 2014*; *Lundberg and Borner, 2019*; *Smith et al., 2016*). The successful implementation of TurboID—an engineered biotin ligase for proximity proteomics (*Branon et al., 2018*)—in model plants *Arabidopsis thaliana* (*Kim et al., 2019*; *Mair et al., 2019*) and *Nicotiana benthamiana* (*Zhang et al., 2019*) suggests its adoption in existing and emerging single-celled marine phytoplankton model systems, including diatoms, is likely (*Falciatore et al., 2020*; *Faktorová et al., 2020*). TurboID is an iron-independent protein and could thus be particuarly valuable to further advance iron metabolism studies in *P. tricornutum*. These proposed improvements and alternative experimental approaches should cross-validate the proteins identified in this study as well as illuminate new ones.

Functionally uncharacterized gene cluster on chromosome 20 we zoomed in on is transcriptionally upregulated in iron-deplete (*Allen et al., 2008*; *Smith et al., 2016*) and silicic acid-replete (*Sapriel et al., 2009*) conditions corroborating a known link between iron and silicon metabolism in diatoms (*Brzezinski et al., 2015*; *Durkin et al., 2012*; *Hutchins et al., 1998*; *Leynaert et al., 2004*). Physically clustered functionally related genes are a hallmark of prokaryotic biology and operon-like genomic elements have been described in fungi and plants (*Osbourn and Field, 2009*; *Nützmann et al., 2018*), however identifying and characterizing them in marine diatom genomes is in its relative infancy. For example, genes encoding light harvesting fucoxanthin-chlorophyll-binding proteins (FCPs) are co-localized on chromosome 2 in *P. tricornutum* (*Bhaya and Grossman, 1993*), and a compact ~7 kbp gene cluster encoding enzymes for domoic acid neurotoxin production was characterized in a cosmopolitan diatom *Pseudo-nitzschia multiseries* (*Brunson et al., 2018*). Given that diatom genomes have likely been extensively shaped by horizontal gene transfer (*Bowler et al., 2008*; *Diner et al., 2017*), it is interesting to note that some yeast species acquired iron acquisition-enabling operon-like genomic elements via horizontal operon transfer (HOT) from an ancient bacterium (*Kominek et al., 2019*). Our phylogenetic analyses suggest all three proteins—pTF.CREG1, pTF.CatCh1, pTF.ap1—encoded by this locus are present in diatoms and other (marine) microeukaryotes, while pTF.CREG1 homologs are found across the tree of life. They, alongside the rest of our proteomic data, paint an emerging view of intracellular ferric, Fe^{3+}, iron processing events downstream of its phytotransferrin-mediated binding at the cell surface.

In humans, endosome acidification causes structural rearrangements in the transferrin receptor protein and protonation of synergistically coordinated carbonate anion leading to ferric iron release (*Cheng et al., 2004*). This event is concurrent with or followed by endosomal reductase STEAP3-catalyzed ferric iron reduction and ferrous, Fe^{2+}, iron export through divalent metal transporter DMT1/NRAMP2 or Zrt- and Irt-like protein (ZIP) 14 (*Bogdan et al., 2016*). Two putative proton, H^+, pumps were present in our mass spectrometry data: Phatr3_J23497—predicted as a P-type ATPase—and Phatr3_J27923—a vacuolar-type H^+-ATPase subunit A. Multisubunit V-ATPases regulate pH homeostasis in virtually all eukaryotes and are known to function as endosome acidifying protein complexes (*Finbow and Harrison, 1997*; *Maxson and Grinstein, 2014*), which makes Phatr3_J27923 a candidate protein involved in acidification of phytotransferrin-rich endosomes in *P. tricornutum*. Notably, V-ATPase transcripts were overrepresented in diatoms following iron enrichment, possibly to account for an increase in the number of endosomes associated with pTF-driven iron acquisition (*Marchetti et al., 2012*), and V-ATPase was shown to be involved in diatom silica deposition vesicle (SDV) acidification (*Hildebrand et al., 2018*; *Yee et al., 2019*). Endosome acidification, perhaps via V-ATPase, may facilitate phytotransferrin-bound ferric, Fe^{3+}, iron release, possibly for processing by an iron reducing enzyme similar to the human metalloreductase STEAP3.

Human cellular repressor of E1A-stimulated genes 1 (CREG1) is a glycoprotein involved in embryonic development, growth, differentiation, and senescence as part of the endosomal-lysosomal system (*Ghobrial et al., 2018*; *Schähs et al., 2008*). Contrary to evolutionarily related flavin mononucleotide (FMN)-binding split barrel fold oxidases and oxidoreductases such as pyridoxine 5′-phosphate (PNP) oxidase, an enzyme involved in vitamin B_6 metabolism (*Musayev et al., 2003*), CREG1 forms a homodimer unable to bind FMN *in vitro* suggesting it does not act as an enzyme *in vivo* (*Sacher et al., 2005*). The lack of FMN in human CREG1 is due to a short loop protruding into the FMN-binding pocket at the dimer interface, and aspartic acid and methionine amino acid residues replacing the corresponding two yeast PNP oxidase (PDB ID: 1CI0) arginines responsible for binding the terminal FMN phosphate (*Sacher et al., 2005*).

CREG1-like protein pTF.CREG1 was proposed to be central to sustained growth of iron-limited *P. tricornutum* cells (*Allen et al., 2008*) or to serve as a cell surface iron receptor (*Lommer et al., 2012*). CREG transcripts were more common in diatom transcriptomes from the Southern Ocean as opposed to non-Southern Ocean regions indicating these proteins are important in coping with iron limitation in this major high nutrient, low chlorophyll (HNLC) zone (*Moreno et al., 2018*). This

may also explain why more diatom homologs are not present in our phylogenetic tree as transcriptomes from the Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP) database used in our phylogenetic analysis were not obtained from iron-limited cultures (*Caron et al., 2016*; *Keeling et al., 2014*). Our protein feature analysis revealed a signal peptide and multiple putative glycosylation sites which suggests that pTF.CREG1 enters the secretory pathway (*Bard and Chia, 2016*). While the nature of pTF.CREG1 recruitment to and colocalization with pTF-rich endosomes in the vicinity of chloroplast periphery—as suggested by our fluorescence microscopy results—is unclear, it is possible that pTF.CREG1-containing vesicles intersect with pTF-rich endosomes *en route* through the secretory pathway. Vesicular trafficking is highly dynamic and bidirectional (*Naslavsky and Kaplan, 2018*; *Progida and Bakke, 2016*), and early endosomal traffic consists of sequential vesicle fusion events (*Brenner, 2012*), so this is at least in principle plausible. Punctate pTF.CREG1 colocalization with pTF suggests that this protein acts in an endosomal-lysosomal system, and this localization pattern is consistent with the observation that proteins with low confidence ASAFind chloroplast prediction (such as pTF.CREG1) can be associated with "blob-like structures" (BLBs) adjacent to, but not inside, the chloroplast (*Gruber et al., 2015*; *Kilian and Kroth, 2005*). Given that our transgenic colocalization *P. tricornutum* cell lines were neither synchronized nor grown in iron-deplete conditions, it is plausible the protein also acts closer to the cell surface. Likely, but not confirmed, APEX2 orientation away from the cell surface and endosome interior in our proximity labeling experiments, and the reported inability of APEX2-generated phenoxyl radicals to penetrate endomembranes (*Rhee et al., 2013*), would suggest that pTF.CREG1—at some point during its recruitment to pTF-rich endosomes—has to be exposed to the cytosol to promote efficient biotinylation. Alternatively, it is impossible to rule out the possibility that lipid composition of *P. tricornutum* endomembranes is permissive to phenoxyl radical diffusion (*Zulu et al., 2018*).

Domain annotation in pTF.CREG1 (UniProt ID: B7G9B3) encouraged us to perform amino acid sequence and structural alignments with human CREG1 (UniProt ID: O75629, PDB ID: 1XHN). These analyses revealed that pTF.CREG1 lacks at least one arginine amino acid residue required for terminal FMN phosphate binding and likely contains an FMN binding pocket-occluding loop, two key features delineating human CREG1 from the yeast PNP oxidase (*Sacher et al., 2005*). Our hypothesis that pTF.CREG1 is unable to coordinate FMN was initially supported by our protein purifications which were not yellow as expected from a typical flavoprotein (*Theorell, 1935*), and was subsequently strengthened by their catalytic inactivity upon supplementation with biologically relevant flavin cofactors. The demonstrated lack of enzymatic activity in pTF.CREG1

and localization data from *in vivo* fluorescent reporter experiments represent an important extension of the human CREG1 literature to the single-celled eukaryotic phytoplankton field.

Secreted glycosylated human CREG1 was shown to delay G1/S cell cycle phase transition via direct interaction with mannose 6-phosphate/insulin-like growth factor II receptor (M6P/IGF2R), a receptor internalized via clathrin-mediated endocytosis (*Di Bacco and Gill, 2003*; *Ghosh et al., 2003*; *Han et al., 2009*). This cellular growth inhibition was demonstrated to, at least partially, depend on a tetrapeptide motif within an FMN binding pocket-occluding loop (*Sacher et al., 2005*). In conjunction with our amino acid sequence alignments and structural prediction which point to the conservation of these two features in pTF.CREG1, these studies lend additional support to our data suggesting colocalization and potential direct physical association between phytotransferrin and pTF.CREG1 is non-enzymatic in nature. Internalization of the human transferrin-transferrin receptor complex is not a constitutive, but rather a regulated kinase activity-dependent process (*Cao et al., 2016*). We currently hypothesize that pTF.CREG1 may serve as a "checkpoint" to keep phytotransferrin endocytosis at a low level in iron-replete conditions, but when iron is scarce instead positively regulate this pathway. Indeed, transcriptomics experiments show that a second transcriptional peak for *pTF.CREG1* occurs in iron-deplete medium approx. 8 hrs prior to the *pTF* peak, a transcriptional dynamic which would be consistent with our hypothesis (*Figure 5—figure supplement 1*; *Smith et al., 2016*). Given that glycosylation in human CREG1 appears to be important for interaction with its cognate receptor (*Han et al., 2011*; *Sacher et al., 2005*; *Schähs et al., 2008*), posttranslational modifications, perhaps glycosylation(s), of pTF.CREG1 may further drive this regulatory activity. The presence of Phatr3_J10972, an iron-insensitive pTF.CREG1 paralog, in the *P. tricornutum* genome suggests that CREG-like proteins are implicated in diverse endocytosis pathways in single-celled eukaryotic phytoplankton. However, the association between iron-sensitive pTF.CREG1 and phytotransferrin in *P. tricornutum* implies a specific pTF.CREG1 function in non-reductive iron uptake. It will be very interesting to define the exact pTF.CREG1 role in further cellular and mechanistic studies and compare it to that of its homologs in multicellular organisms. Conceivably, an ancient protein like pTF.CREG1 could have remained at the core of linking cell cycle progression, cellular growth, and nutrient homeostasis across the tree of life.

In contrast to pTF.CREG1, no functional annotation could be assigned to and no structural homology models were generated for pTF.CatCh1 in the Pfam database (*El-Gebali et al., 2019*) and by Phyre2, respectively. This protein has a predicted N-terminal signal peptide and numerous

N- and O-glycosylation sites suggesting it enters and trafficks along the secretory pathway. The

outermost complex plastid membrane in diatoms corresponds to the chloroplast endoplasmic

reticulum (cER) and is continuous with the nuclear envelope (*Gibbs et al., 1981*; *Prihoda et al.,*

2012). Nuclear-encoded *P. tricornutum* proteins destined for the periplastidial compartment

(vastly reduced endosymbiont cytoplasm between the inner two and the outer two complex plastid

membranes) are first cotranslationally imported into the cER lumen (*Grosche et al., 2014*;

Peschke et al., 2013). Our microscopy results show that pTF.CatCh1 is localized on the

chloroplast margin, perhaps anchored to the outer chloroplast (cER) membrane with its C-terminal

transmembrane domain, thus raising a possibility that pTF.CatCh1 gets to its final destination via

cotranslational membrane insertion.

The cysteine-rich domain contains one CXC and one CXXC motifs which are nearly invariably

present in pTF.CatCh1 homologs. Two cysteines spaced by two non-cysteine amino acids (the

CXXC motif) are commonly found in metal-binding proteins, including those that bind iron

(*Fomenko et al., 2008*; *Gladyshev et al., 2004*). This motif in pTF.CatCh1 is preceded by an

aspartic acid residue resulting in DCXXC which is known to be particularly permissive to binding

divalent metal ions (*Figure 7—figure supplement 1A*; *Banci et al., 2002*; *Banci et al., 2006*).

Histidine, tyrosine, methionine, aspartic acid, and glutamic acid residues can also be involved in

iron cation coordination, but none of these are conserved across pTF.CatCh1 homologs (*Figure*

7—source data 1; *Dokmanić et al., 2007*). Given the nearly completely conserved C(X)XC

motifs and additional six 100% conserved cysteine residues, it is conceivable they are critical to

pTF.CatCh1 activity.

Based on the fluorescence microscopy data, the conserved cysteine amino acid residues and

cysteine-containing motifs, and given that disordered protein domains—two are predicted to flank

the pTF.CatCh1 transmembrane domain—are mediators of protein-protein interactions and play

an important role in the formation of protein complexes (*Uversky, 2015b*; *Uversky, 2016*), we

are considering two hypotheses for *in vivo* pTF.CatCh1 function.

First, pTF.CatCh1 may serve as a metallochaperone that binds ferrous, Fe^{2+}, iron downstream of

the endosomal ferric, Fe^{3+}, iron reduction step, and directs it to the chloroplast interior.

Metallochaperones are intracellular proteins responsible for protecting and guiding metal ions on

their way to correct metalloenzyme sinks (*O'Halloran and Culotta, 2000*). They engage in

protein-protein interactions and help prevent metal-induced toxicity caused by free radical-

generating Fenton reactions which are especially common with iron and copper (*Philpott and Jadhav, 2019*; *Rosenzweig, 2002*; *Valko et al., 2005*).

Second, pTF.CatCh1 may recruit pTF to the chloroplast periphery via one of its two C-terminal disordered regions flanking the predicted transmembrane domain, and thus serve as a docking site for iron-laden pTF. It could achieve this via the "fly-casting mechanism" which involves binding-induced protein folding (*Shoemaker et al., 2000*). The observed colocalization of pTF.CatCh1 with pTF may indicate that Fe^{3+} can be offloaded directly from pTF to pTF.CatCh1 which would imply an additional, non-endosomal, intracellular Fe^{3+} reduction step. Notably, a chloroplast-associated Fe^{3+} reduction pathway mediated by Fe^{3+} chelate reductase FRO7 is present in *Arabidopsis thaliana* (*Jeong et al., 2008*). Abberant cellular localization of heterologously expressed fluorescent protein fusions is a widely observed and acknowledged phenomenon (*Jensen, 2012*), but we note that pTF localization on the chloroplast margin insdistinguishable from that in this study was reported by *McQuaid et al., 2018* suggesting it is biologically relevant. Similarly to *pTF.CREG1*, *pTF.CatCh1* exhibits a second transcriptional peak in iron-deplete medium approx. 16 hrs prior to the *pTF* peak (*Figure 5—figure supplement 1*). It would make sense to synthesize a protein critical for iron-laden pTF recruitment to the chloroplast margin prior to significant increase in basal pTF endocytosis.

All in all, our data suggest pTF.CatCh1 is a chloroplast margin-localized, pTF-associated, *P. tricornutum* protein with a possible role in chloroplast iron homeostasis. The two hypotheses are put forth here to help frame its further genetic, cellular, biochemical, and structural dissection.

Chloroplasts are the most iron-rich system in plant cells (*López-Millán et al., 2016*). Iron—in particular in the form of iron–sulfur (Fe–S) clusters—serves as an essential cofactor for the photosynthetic electron transfer chain, chlorophyll biosynthesis, and chloroplast protein import (*Balk and Schaedler, 2014*; *Marchand et al., 2018*; *Soll and Schleiff, 2004*). A complex vesicle-based system possibly in perpetual dynamic exchange with cytosolic vesicles is present in chloroplasts (*Hertle et al., 2020*; *Lindquist and Aronsson, 2018*), an endoplasmic reticulum to Golgi to chloroplast protein trafficking pathway exists (*Radhamony and Theg, 2006*; *Villarejo et al., 2005*), and additional hypotheses for such specialized vesicle-mediated plastid targeting of proteins are established (*Baslam et al., 2016*). Therefore, a direct link between endocytic internalization of phytotransferrin-bound iron and its offloading to or adjacent to chloroplasts as suggested by our results seems plausible. "Kiss and run" mechanism that allows erythroid cell

mitochondria to access iron via direct interaction with transferrin-rich endosomes further
underscores this idea (*Hamdi et al., 2016*). Analysis of our mass spectrometry (MS) hits that were
not co-expressed with pTF-mCherry reveals three additional proteins—ISIP1 (Phatr3_J55031),
FBP1 (Phatr3_J46929), and FBAC5 (Phatr3_J41423)—that support the idea of a direct cell
surface-to-chloroplast iron trafficking pathway in *Phaeodactylum tricornutum*.

ISIP1 (iron starvation induced protein 1; Phatr3_J55031) gene expression in *P. tricornutum* and
other marine microeukaryotes is induced by iron limitation (*Allen et al., 2008*; *Kazamia et al.,*
2018; *Marchetti et al., 2012*). *P. tricornutum* ISIP1—the second most enriched protein in our MS
dataset—has a role in siderophore-bound iron uptake and colocalizes with phytotransferrin (pTF)
adjacently to the chloroplast in an "iron-processing compartment" (*Kazamia et al., 2019*; *Kazamia*
et al., 2018). FBP1 (Phatr3_J46929)—iron-sensitive siderophore ferrichrome-binding protein with
punctate intracellular localization and central to the carbonate-independent iron acquisition
pathway in *P. tricornutum* (*Allen et al., 2008*; *Coale et al., 2019*)—was another highly enriched
protein in our dataset indicating the internalization pathways for inorganic (i.e., phytotransferrin-
bound) and organic (i.e., siderophore-bound) iron in *P. tricornutum* may be physically coupled
intracellularly and directed to iron-accumulating vesicles close to the chloroplast. Metal
homeostasis by such discrete membranous compartments is widespread in nature (*Blaby-Haas*
and Merchant, 2014). Copper- and zinc-storing compartments—cuprosomes and zincosomes,
respectively—are present in the model chlorophyte microalga *Chlamydomonas reinhardtii* (*Aron*
et al., 2015; *Hong-Hermesdorf et al., 2014*; *Merchant, 2019*), and lipid-bound iron-accumulating
ferrosome compartments exist in diverse bacteria and archaea (*Grant and Komeili, 2020*).

Finally, the most enriched protein in our MS dataset was FBAC5 (Phatr3_J41423)—iron-sensitive
iron-independent class I fructose-biophosphate aldolase—which was shown to be localized to the
P. tricornutum pyrenoid—proteinaceous RuBisCO-containing subchloroplastic organelle—and
proposed to link Calvin-Benson-Bassham cycle activity with the CO_2 concentrating mechanism
(*Allen et al., 2012*; *Matsuda et al., 2017*). FBAC5 is one of the most reliable markers for diatom
iron stress as demonstrated by laboratory culture (*Cohen et al., 2018*; *Lommer et al., 2012*),
microcosm incubation (*Cohen et al., 2017*; *Cohen et al., 2017*), and field studies (*Bertrand et*
al., 2015), however its exact role has remained elusive. Fructose-biophosphate aldolases (FBAs)
are glycolytic enzymes known for their ability to moonlight (*Boukouris et al., 2016*), a term for
proteins with multiple physiologically relevant functions encoded by one polypeptide chain
(*Jeffery, 2018*). Moonlighting FBAs have roles in actin cytoskeleton dynamic, cellular growth

arrest, and apoptosis (in human cancer cell lines; *Gizak et al., 2019*), transcriptional regulation

(in yeast; *Cieśla et al., 2014*), and endosome acidification via direct interaction with vacuolar-type

ATPase (in a mouse cell line; *Merkulova et al., 2011*). These reported non-catalytic FBA activities

suggest that FBAC5 acting outside of the *P. tricornutum* chloroplast, perhaps in the endosomal-

lysosomal system as our data would imply, is possible.

Our initial characterization of pTF.CREG1 and pTF.CatCh1, together with the discussion on the

four other mass spectrometry hits, enable proposition of a model connecting cell surface ferric

iron binding, internalization, and intracellular trafficking in *Phaeodactylum tricornutum* with many

outstanding questions ripe for future investigation (*Figure 8*). Importantly, it remains to be

determined which protein is responsible for intracellular ferric, Fe^{3+}, iron reduction in *P.*

tricornutum, activity that would be consistent with the non-reductive cell surface ferric, Fe^{3+}, iron

uptake proposed for phytotransferrin (pTF) (*McQuaid et al., 2018*; *Morrissey et al., 2015*).

Perhaps notably, neither of the five proteins with clear ferric reductase annotation (FRE1–5,

Phatr3 IDs: J54486, J46928, J54940, J54409, J54982) nor additional five hypothetical ferric

reductases (*Coale et al., 2019*; *Smith et al., 2016*) were present in our mass spectrometry data.

Further proximity-dependent proteomic mapping experiments will be essential for determining the

full set of proteins in the non-reductive iron acquisition pathway in *P. tricornutum* as enzymatic

Fe^{3+} reduction may happen later in the pathway and/or in a separate subcellular compartment.

In conclusion, the identified proteins and the associated model we introduce in this work represent

an important advance for the marine phytoplankton field, given that diatoms are central to the

study of cellular iron homeostasis and its link to fluctuating iron levels in the ocean. The presented

insights are vital as our understanding of intracellular iron trafficking is relatively incomplete even

in significantly more established model systems such as mammalian cells, and our knowledge of

chloroplast iron uptake is in its infancy (*Blaby-Haas and Merchant, 2012*; *Philpott and Jadhav,*

2019). We anticipate further molecular dissection of our proteomics dataset will not only provide

details on the transformation dynamic of acquired ferric iron and its cellular sinks but will thereby

also illuminate the extent to which phytotransferrin pathway mirrors the metazoan transferrin

cycle.

Implementing the use of APEX2 in a model marine diatom is itself a significant and timely advance

as the field is expanding and moving from broad -omics-based surveys to detailed molecular

studies (*Falciatore et al., 2020*; *Faktorová et al., 2020*). APEX2 and related molecular tools hold

great promise to dissect additional key subcellular compartments in diatoms such as pyrenoids

and silica deposition vesicles (*Figure 8—figure supplement 1*; *Barrett et al., 2021*; *Hildebrand*

et al., 2018; *Matsuda et al. 2017*; *Wang and Jonikas, 2020*), two exciting future avenues for

these single-celled heterokonts uniquely positioned at the nexus of evolutionary cell biology and

blue biotechnology.

<u>**Vector cloning**</u>

pJT_NR_pTF-AP2: Gibson Assembly (*Gibson et al., 2009*) was performed with three DNA

fragments: (1) linearized pPtPBR1 episome backbone (Addgene, Cambridge, MA, plasmid

pPtPBR1, Catalog #80388) opened ~280 bp downstream of the tetracycline resistance gene, (2)

amplicon with nitrate reductase gene (*NR*) promoter, 5' *NR* untranslated region (UTR), and *pTF*,

and (3) gBlocks Gene Fragment (Integrated DNA Technologies (IDT), Coralville, IA) with linker

sequence, codon optimized *APEX2* (using IDT Codon Optimization Tool and selecting

"*Thalassiosira pseudonana*" from the "Organism" drop-down menu), 3' *NR* UTR, and *NR*

terminator.

pJT_native_pTF-mCherry: Gibson Assembly was performed with three DNA amplicons: (1)

pPtPBR1 episome backbone split into two fragments and (2) expression cassette including *pTF*,

mCherry, and *pTF* promoter and terminator.

pJT_pTF-mCherry_MS hit-EYFP: pJT_native_pTF-mCherry was split into two fragments via

PCR stitching (keeping ampicillin resistance gene split). Genes corresponding to MS hits were

amplified using cDNA from iron-starved WT *P. tricornutum* cells and assembled with *EYFP* and

Phatr3_J23658 (flavodoxin-encoding gene) promoter and terminator through two PCR stitching

rounds into a single fragment. Gibson Assembly was then performed to combine all three final

amplicons. Detailed assembly scheme is presented in *Figure 5—figure supplement 2A*.

pJT_Δ31_pTF.CREG1-His6: Cloning of gene fragments into the *E. coli* protein expression vector

PtpBAD-CTHF was performed as described previously (*Brunson et al., 2018*). Briefly, PtpBAD-

CTHF was linearized by digestion with XhoI (New England Biolabs (NEB), Ipswich, MA, Catalog

#R0146S) and the resulting DNA was column purified. *Δ18_pTF.CREG1* gene was obtained by

PCR from *P. tricornutum* gDNA with PrimeSTAR GXL DNA Polymerase (Takara Bio, Kusatsu,

Japan, Catalog #R050B) and primer set JT31/JT32 to incorporate the appropriate Gibson

Assembly overhangs and remove the predicted N-terminal signal peptide (amino acid residues

1–18). Insertion of truncated *pTF.CREG1* into linearized PtpBAD-CTHF was performed using

Gibson Assembly Master Mix (NEB, Catalog #E2611S), 1 uL of the Gibson Assembly reaction

mixture was transformed via heat shock into chemically competent NEB 5-alpha cells (NEB,

Catalog #C2988J), and cells were incubated on lysogeny broth with 10 μg/mL tetracycline (LB-

Tet10) 1% agar plates overnight at 37°C. Transformants were screened by colony PCR using the

primer set JT37/JT38 and Sapphire polymerase (Takara Bio), and positive clones were selected

for outgrowth. Isolated plasmids were sequence-validated by Sanger sequencing (Eurofins,

Luxembourg, Luxembourg). A sequence-validated clone was designated as PtpBAD-

Δ18_pTF.CREG1-CTHF and transformed into chemically competent BL21 *E. coli* cells (NEB,

Catalog #C2530H) which were spread on LB-Tet10 1% agar plates. The resulting transformants

were used for subsequent Δ18_pTF.CREG1 expression experiments. Following unsuccessful

expression testing of the Δ18 N-terminal truncation construct, an additional set of genes encoding

pTF.CREG1 with N-terminal truncations (Δ26, Δ31, Δ39, Δ43) was generated by PCR using

PrimeStar GXL DNA Polymerase (Takara Bio), primer sets JT33/JT32, JT34/JT32, JT35/JT32,

JT36/JT32, and PtpBAD-Δ18_pTF.CREG1-CTHF as a template. Assembly, colony PCR

screening, plasmid isolation, sequencing, and transformation into chemically competent BL21 *E.*

coli cells of all additional expression vectors, including PtpBAD-Δ31_pTF.CREG1-CTHF

(=pJT_Δ31_pTF.CREG1-His6), was performed as above. An additional construct encoding

pTF.CREG1 with an N-terminal His6 and Δ39 N-terminal truncation was built (PtpBAD-NTH-

Δ39_pTF.CREG1) using the vector NTH-PtpBAD (constructed similarly to PtpBAD-CTHF; see

Brunson et al., 2018 and *Savitsky et al., 2010*), but was not pursued beyond initial expression

testing.

pTF (*Phatr3_J54465*) in all episomes was in its native form (3 exons, 2 introns). Molecular cloning

primers are listed in *Supplemental file 1—Table S3*. Vector details, further amplicon information,

and fusion protein sequences are catalogued in *Supplemental file 1—Table S4*.

Related resource: Turnšek J. and Gholami P. 2017. Guidelines for highly efficient construction

of diatom episomes using Gibson Assembly. *protocols.io*. DOI:

dx.doi.org/10.17504/protocols.io.jy7cpzn

Diatom culturing, conjugation, and genotyping

Culturing: Sequenced *Phaeodactylum tricornutum* strain CCMP632 (synonymous to CCMP2561

and CCAP 1055/1; National Center for Marine Algae and Microbiota (NCMA), East Boothbay,

ME) was used throughout the study and grown at 18°C, 300 μmol quanta m^{-2} s^{-1}, and a 10 h:14

h dark:light cycle in biotin-free L1 medium prepared by mixing 1 L Aquil salts, 2 mL nitrate and

phosphate (NP) nutrient stock, 1 mL trace metal stock, and 1 mL thiamine hydrochloride and cyanocobalamin (TC) stock unless otherwise noted. Preparation of Aquil salts: 0.5 L anhydrous salts (0.5 L Milli-Q, 24.5 g NaCl, 4.09 g Na_2SO_4, 0.7 g KCl, 0.2 g $NaHCO_3$, 0.1 g KBr, 900 μL 33.3 mg/mL H_3BO_3 stock, 300 μL 10 mg/mL NaF stock) and 0.5 L hydrous salts (0.5 L Milli-Q, 11.1 g $MgCl_2$ x $6H_2O$, 1.54 g $CaCl_2$ x $2H_2O$, 100 μL 170 mg/mL $SrCl_2$ x $6H_2O$ stock) were prepared separately, combined, filter sterilized (0.2 μm), and stored at room temperature. Preparation of NP nutrient stock: 37.5 g $NaNO_3$ and 2.5 g NaH_2PO_4 were dissolved in 100 mL Milli-Q, filter sterilized (0.2 μm), and stored at 4°C. Preparation of trace metal stock (for 1 L 1,000x stock): 3.15 g $FeCl_3$ x $6H_2O$, 4.36 g Na_2EDTA x $2H_2O$, 0.25 mL 9.8 g/L $CuSO_4$ x $5H_2O$, 3.0 mL 6.3 g/L Na_2MoO_4 x $2H_2O$, 1.0 mL 22 g/L $ZnSO_4$ x $7H_2O$, 1.0 mL 10 g/L $CoCl_2$ x $6H_2O$, 1.0 mL 180 g/L $MnCl_2$ x $4H_2O$, 1.0 mL 1.3 g/L H_2SeO_3, 1.0 mL 2.7 g/L $NiSO_4$ x $6H_2O$, 1.0 mL 1.84 g/L Na_3VO_4, 1.0 mL 1.94 g/L K_2CrO_4, and Milli-Q up to 1 L were combined, filter sterilized (0.2 μm), and kept at 4°C. Preparation of TC stock: 20 mg thiamine hydrochloride and 0.1 mL 1 g/L cyanocobalamin stock were mixed in 100 mL Milli-Q. The resulting solution was stored at 4°C. *ΔpTF P. tricornutum* cell line (maintained in the Allen Lab; *McQuaid et al., 2018*) was additionally supplemented with 200 μg/mL nourseothricin (GoldBio, Saint Louis, MO, Catalog #N-500-1). All transconjugant *P. tricornutum* cell lines were supplemented with 50 or 100 μg/mL phleomycin (InvivoGen, San Diego, CA, Catalog #ant-ph-10p).

Conjugation: (1) Bacterial donor preparation: Chemically competent pTA-Mob-containing TransforMax™ EPI300™ *E. coli* cells (*Strand et al., 2014*; Lucigen, Middleton, WI, Catalog #EC300110) were transformed via heat shock with sequence-verified pPtPBR1 episomes. Transformants were selected on gentamycin-, carbenicillin-, and tetracycline-containing LB 1% agar plates. 3 mL overnight LB cultures supplemented with antibiotics were inoculated from glycerol stocks. (2) *P. tricornutum* preparation: ~2 x 10^8 *P. tricornutum* cells (in 200 μL) in mid- to late-exponential phase were spread on pre-dried (i.e., plates with lid half open and kept in the laminar flow hood for at least 90 min) ½ L1 1% agar plates with 5% LB and left growing for 1 or 2 days. Plates were additionally supplemented with 200 μg/mL nourseothricin for conjugation of *ΔpTF P. tricornutum* cells. (3) Conjugation: Overnight donor bacterial cultures were diluted 1:50 in 25 mL LB supplemented with antibiotics, grown at 37°C until OD_{600} 0.8–1, spun down, resuspended in 150 uL Super Optimal broth with Catabolite repression (SOC) medium, and spread as evenly as possible on top of a *P. tricornutum* lawn. Plates with donor-*P. tricornutum* co-culture were first left in dark and 30°C for 90 min, then for 1 or 2 days at standard growth conditions. (4) Selection: Co-culture lawn was scraped off of plates with 1 mL fresh L1 medium,

transferred to a microcentrifuge tube, and 200 µL spread on pre-dried (see above) ½ L1 1% agar

plates with 50 or 100 µg/mL phleomycin. Plates were additionally supplemented with 200 µg/mL

nourseothricin for conjugation of *ΔpTF P. tricornutum* cells. Porous adhesive tape was used to

seal the plates and transconjugants emerged after ~10 days of incubation under standard growth

conditions. Please see *Karas et al., 2015* and *Diner et al., 2016* for further description of diatom

conjugation.

Genotyping: Candidate transconjugant colonies were inoculated in 300 µL L1 medium

supplemented with 50 or 100 µg/mL phleomycin (and 200 µg/mL nourseothricin in case of *ΔpTF*

P. tricornutum cells) and typically grown for ~1 week. 0.5 µL liquid culture was then genotyped

using either Phire Plant Direct PCR Master Mix (Thermo Fischer Scientific, Waltham, MA, Catalog

#F160L) or Phire Plant Direct PCR Kit (Thermo Fischer Scientific, Catalog #F130WH). 200 µL of

each genotype positive cell line was passaged in 30 mL L1 medium supplemented with 50 or 100

µg/mL phleomycin (and 200 µg/mL nourseothricin in case of *ΔpTF P. tricornutum* cells).

<u>Related resource:</u> Turnšek J. 2017. Simple & rapid genotyping of marine microeukaryotes.

protocols.io. DOI: dx.doi.org/10.17504/protocols.io.jcdcis6

RNA extraction and cDNA synthesis

Diatom culture: 100 mL WT *P. tricornutum* culture was grown in iron-deplete conditions (L1

medium with 7.5 nM total iron) for two weeks. Cells were then centrifuged, supernatants

discarded, pellets flash frozen in liquid nitrogen, and stored at -80°C.

RNA extraction: Direct-zol™ RNA Miniprep Plus RNA extraction kit was used (Zymo Research,

Irvine, CA, Catalog #R2070). Briefly, cell pellets were resuspended in 800 µL Trizol, equal amount

of 100% ethanol, and centrifuged in spin columns. Columns were then washed with 400 µL RNA

Wash Buffer followed by on-column DNA digestion with 5 µL DNase I in 75 µL DNA Digestion

Buffer for 15 min at RT, washing twice with 400 µL Direct-zol RNA PreWash, and once with 700

µL RNA Wash Buffer. RNA was eluted with 50 µL DNase/RNase-Free Water, RNA integrity

number (RIN) evaluated with 2200 TapeStation (Agilent Technologies, Santa Clara, CA;

measured RIN was 7.0), concentration estimated with Qubit 2.0 Fluorometer (Thermo Fischer

Scientific; measured concentration was 27.6 ng/µL), and samples stored at -80°C.

cDNA synthesis: cDNA synthesis kit from Thermo Fischer Scientific was used (SuperScript™ III First-Strand Synthesis System, Catalog #18080-051). Briefly, 1 µL total RNA was combined with 0.5 µL Oligo(dT)$_{20}$ Primer, 0.5 µL 10 mM dNTP Mix, and 3 µL nuclease-free water followed by 5 min and 1 min incubation at 65°C and on ice, respectively. After 5 µL cDNA Synthesis Mix (1 µL 10x Reverse Transcription buffer, 2 µL 25 mM MgCl$_2$, 1 µL 0.1 M DTT, 0.5 µL RNase OUT, 0.5 µL SuperScript III Reverse Transcriptase) was added, samples were incubated for 50 min and 5 min at 50°C and 85°C, respectively, then chilled on ice. Finally, 1 µL RNase H was added and sample kept for 20 min at 37°C. cDNA was stored at -20°C until use.

MDY-64 labeling and imaging

10 mL of a pTF-mCherry expressing *ΔpTF P. tricornutum* cell line grown in iron-deplete conditions (L1 medium with 7.5 nM total iron) were spun down, supernatant discarded, and pellet resuspended in 50 µL phosphate-buffered saline (PBS) (pH 7.4). 0.5 µL 10 mM MDY-64 stock (in DMSO; Thermo Fischer Scientific, Catalog #Y7536) was added, cells incubated for 10 min at RT, pelleted, and resuspended in 50 µL fresh PBS (pH 7.4). Cell suspension was prepared for imaging as follows: 5 µL was placed between a 1.5 mm microscope slide and a cover slip (this setup applies to all imaging experiments in the study). Imaging conditions: Leica TCS SP5 confocal microscope (Leica Microsystems, Wetzlar, Germany), argon laser strength set to 30%, 458 nm laser line at 50% maximum strength, emission window set to 477–517 nm (for visualizing MDY-64; MDY-64 excitation and emission maxima are 451 nm and 497 nm, respectively), 514 nm laser line at 50% maximum strength, emission window set to 620–640 nm (for visualizing mCherry; mCherry excitation and emission maxima are 587 nm and 610 nm, respectively).

Protein expression analyses

pTF-APEX2 detection: Cell pellets from 8 mL mid- to late-exponential phase WT or transconjugant *P. tricornutum* cultures were resuspended in 150 µL cell lysis buffer (50 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 1 mM PMSF, pH 8.5) and sonicated for 5 min (30 sec on, 1 min off) with Bioruptor UCD-200TM (Diagenode, Liège, Belgium). The resulting cell lysates were centrifuged, total protein content in supernatants measured with Bradford Assay Kit (Thermo Fischer Scientific, Catalog #23236), and insoluble fractions resuspended in 150 µL cell lysis buffer. 1 µg soluble protein samples and 0.5 µL resuspended insoluble protein fractions were resolved on a NuPage 4-12% Bis-Tris 1.5 mm gel (Thermo Fischer Scientific, Catalog

#NP0335BOX), wet transferred to polyvinylidene difluoride (PVDF) membranes (Thermo Fischer Scientific, Catalog #LC2005), and visualized with WesternBreeze™ Chemiluminescent Kit, anti-rabbit (Thermo Fischer Scientific, Catalog #WB7106).

pTF-mCherry detection: Cell pellets from 400 µL mid- to late-exponential phase WT or transconjugant *P. tricornutum* cultures were resuspended in 50 µL cell lysis buffer (50 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 1 mM PMSF, pH 8.5) and sonicated 15 min (30 sec on, 1 min off) with Bioruptor UCD-200TM. 1 µL of each whole cell lysate was resolved on a NuPage 4-12% Bis-Tris 1.5 mm gel, wet transferred to PVDF membranes, and visualized with WesternBreeze™ Chemiluminescent Kit, anti-rabbit.

pTF antibody details: Amino acid residues 32–223 served as the immunogen (N-terminal pTF region just downstream of the signal peptide). The antibody was produced in a rabbit. 1:10,000 dilution of 1.14 mg/mL antibody stock was used in this study for background-free results.

Protein ladder: MagicMark™ XP Western Protein Standard (Thermo Fischer Scientific, Catalog #LC5602).

Related resource: Turnšek J. 2017. HA tag enables highly efficient detection of heterologous proteins in *Phaeodactylum tricornutum* (*Pt*) exconjugants. *protocols.io*. DOI: dx.doi.org/10.17504/protocols.io.j7ncrme

Amplex UltraRed assay and resorufin imaging

Amplex UltraRed assay: 5 mL of triplicate WT or pTF-APEX2 expressing *P. tricornutum* cultures in mid- to late-exponential phase were incubated on ice for 5 min, spun down, supernatant discarded, pellet resuspended in 500 µL ice-cold PBS (pH 7.4), and transferred to microcentrifuge tubes. Cells were spun down again, resuspended in 200 µL ice-cold reaction buffer (50 µM Amplex UltraRed (AUR; Thermo Fischer Scientific, Catalog #A36006), 2 mM H_2O_2, in PBS (pH 7.4)), and incubated on ice for 15 min unless otherwise noted. 50 µL supernatant was mixed with 50 µL PBS (pH 7.4) and resorufin fluorescence measured in a black microtiter plate with black bottom using Flexstation 3 microtiter plate reader (Molecular Devices, San Jose, CA; excitation: 544 nm, emission: 590 nm; resorufin excitation and emission maxima are 568 nm and 581 nm, respectively). Horseradish peroxidase (HRP) was always included as a positive assay control.

Fluorescence was normalized to OD_{750} of experimental *P. tricornutum* cultures. Amplex UltraRed

was prepared as a 10 mM stock in DMSO and stored in 20 µL aliquots at -20°C. 3% (w/w) H_2O_2

stock (Sigma-Aldrich, Saint Louis, MO, Catalog #323381-25ML) was stored in 100 µL aliquots at

-20°C.

Resorufin imaging: WT and pTF-APEX2 expressing *P. tricornutum* cells after performing the

Amplex UltraRed assay were imaged with Leica TCS SP5 confocal microscope using the

following parameters: argon laser strength at 30%, 514 nm laser line at 50% maximum strength,

resorufin emission window: 575–605 nm, autofluorescence emission window: 700–750 nm.

Transmission electron microscopy (TEM)

Part 1: Labeling

5 mL of triplicate WT or pTF-APEX2 expressing *P. tricornutum* cultures in mid- to late-exponential

phase were spun down (4,000 rpm, 4°C, 10 min) and fixed in 5 mL ice-cold 2% (w/v)

paraformaldehyde (PFA) and 2% (v/v) glutaraldehyde in 0.15 M sodium cacodylate buffer (pH

7.4) for 30 min on ice. Cells were rinsed in 5 mL 0.15 M sodium cacodylate buffer (pH 7.4) five

times for 3 min on ice, then once again in 5 mL 0.15 M sodium cacodylate buffer (pH 7.4) with 10

mM glycine for 3 min on ice. Cells were then treated with 25 mM 3,3'-diaminobenzidine (DAB;

Sigma-Aldrich, Catalog #D8001) as follows: 5.36 mg DAB was dissolved in 1 mL 0.1 N HCl and

sonicated for 45 min. 5 mL of 0.3 M sodium cacodylate buffer (pH 7.4) was added to dissolved

DAB, final volume adjusted to 10 ml with ddH_2O, solution filtered through a 0.22 µm syringe filter,

and 3 µL 30% (w/w) H_2O_2 added for 3 mM final concentration. Cells were incubated in 5 mL of

this reaction buffer for 15 min on ice, rinsed five times for 3 min in 5 mL 0.15 M sodium cacodylate

buffer (pH 7.4), post-fixed in 2 mL 1% osmium tetroxide (OsO_4) (Electron Microscopy Sciences,

Hatfield, PA, Catalog #19150) in 0.15 M sodium cacodylate buffer (pH 7.4) for 30 min on ice, and

rinsed in 5 mL ice-cold ddH_2O five times for 3 min. Cells were then resuspended in 300 µL melted

3% agar, poured onto a glass slide sitting on ice, left to solidify, and cut into small ~3 x 3 x 1 mm

pieces which were transferred into scintillation vials with 10 mL ice-cold ddH_2O. All shorter spin

down steps were done at 4,000 rpm and 4°C for 1.5 min. All buffers were used ice-cold.

Part 2: TEM sample preparation

Agar blocks with embedded *P. tricornutum* cells were fixed in 10 mL 2% (v/v) glutaraldehyde in

ddH_2O to crosslink agar, rinsed five times for 2 min in 5 mL ice-cold ddH_2O, and left incubating in

mL 2% uranyl acetate (Electron Microscopy Sciences, Catalog #22400) overnight at 4°C. Next morning, agar blocks were first dehydrated in the following ethanol series: 20%, 50%, 70%, 90%, 100% (on ice, 10 mL), 100%, 100% (at RT, 10 mL), then infiltrated with 10 mL 50% epoxy resin for ~1 h. To prepare 20 mL 100% resin, Durcupan ACM mixture (Electron Microscopy Sciences, Catalog #14040) components were combined as follows: 11.4 g A (epoxy resin), 10 g B (964 hardener), 0.3 g C (964 accelerator), and 0.1 g D (dibutyl phthalate) in this exact order (for 50% resin 1 part 100% resin was combined with 1 part 100% ethanol). Agar blocks were transferred into 10 mL 100% resin for ~4 h, then into fresh 10 mL 100% resin overnight, again into fresh 10 mL 100% resin for 4 h the following morning before finally being poured into aluminum boats and left to polymerize in 60°C oven for at least 48 h (over the weekend). Polymerized resins were detached from aluminum boats, agar blocks dense with cells cut out and glued to "dummy" blocks by incubation in a 60°C oven for at least 15 min. ~500 nm thick sections were cut with an ultramicrotome (Leica Ultracut), stained with 1% toluidine blue, and observed under light microscope to make sure embedded cells were exposed. ~100 nm thick sections were then cut, placed on TEM grids, labeled, and saved until imaging. Imaging was performed with JEOL JEM-1200 (Japan Electron Optics Laboratory, Akishima, Tokyo, Japan) transmission electron microscope at 80 keV. Backscatter scanning electron microscopy was performed with Zeiss Merlin (Oberkochen, Germany) scanning electron microscope (SEM) at 2 keV by placing ~80 nm thick sections on a silicon wafer and imaged with inverted contrast which gives a TEM-like image.

Proximity-dependent proteomic mapping

Part 1: Labeling

25 mL of quintuplicate WT and pTF-APEX2 expressing *P. tricornutum* cultures in mid- to late-exponential phase—cell density of all cultures just prior to harvest was ~2 x 10^7 cells/mL which corresponds to OD_{750} ~0.4—were cooled on ice for 10 min and pelleted (4,000 rpm, 4°C, 10 min). Supernatants were discarded, pellets resuspended in 0.5 mL ice-cold PBS (pH 7.4), transferred to microcentrifuge tubes, and spun down (4,000 rpm, 4°C, 10 min). Cells were then resuspended in 0.5 mL ice-cold 1.2 M D-sorbitol in PBS (pH 7.4), supplemented with 2.5 mM biotin-phenol (Berry & Associates, Dexter, MI, Catalog #BT 1015), incubated on a tube rotator at 4°C for 90 min, supplemented with 1 mM H_2O_2, and incubated on a tube rotator at 4°C for another 20 min. Labeling reaction was quenched by washing cells twice (4,000 rpm, 4°C, 5 min) with 0.5 mL ice-cold quenching solution (10 mM sodium ascorbate (VWR International, Radnor, PA, Catalog #95035-692), 5 mM Trolox (Sigma-Aldrich, Catalog #238813-5G), 10 mM sodium azide (VWR

International, Catalog #AA14314-22) in PBS (pH 7.4)). 50 μL of quenched cell suspension was saved for a streptavidin blot. The remaining 450 μL was spun down (4,000 rpm, 4°C, 10 min) and lysed in 250 μL cell lysis buffer (50 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 1 mM PMSF, pH 8.5) by sonication for 15 min (30 sec on, 1 min off). Cell lysates were spun down at 4,000 rpm and 4°C for 45 min and protein concentration in supernatants measured using Bradford Assay Kit. For streptavidin blot, 25 μL of saved quenched cells were first sonicated for 15 min (30 sec on, 1 min off). 2.5 μL of whole cell lysates and 1 ng biotinylated BSA positive control (Thermo Fischer Scientific, Catalog #29130) were then resolved on a NuPage 4-12% Bis-Tris 1.5 mm gel and wet transferred to a PVDF membrane. Membrane was washed twice with 15 mL PBST (PBS (pH 7.4) with 0.1% (v/v) Tween-20) for 5 min, left blocking overnight at RT and gentle shaking in PBST supplemented with 5% BSA (Sigma-Aldrich, Catalog #A9647-100G), and washed once the next morning with 15 mL PBST for 5 min. It was then incubated for 1 h at RT and gentle shaking in 15 mL PBST supplemented with 5% BSA and 1:15,000 HRP-conjugated streptavidin (Thermo Fischer Scientific, Catalog #S911). Finally, membrane was washed three times with 15 mL PBST supplemented with 5% BSA for 5 min and once with 15 mL PBST for 10 min after which it was visualized with SuperSignal™ West Dura Extended Duration reagent (Thermo Fischer Scientific, Catalog #34075) using C-DiGit® Blot Scanner (Li-COR, Lincoln, NE).

Protein ladder: WesternSure® Pre-stained Chemiluminescent Protein Ladder (HRP-conjugated protein ladder) (Li-COR, Catalog #926-98000).

Part 2: Pull-down and mass spectrometry

Pull-down: 50 μL streptavidin beads (Thermo Fischer Scientific, Catalog #88816) transferred to microcentrifuge tubes were pelleted in a magnetic rack (tubes were left standing for ~5 min to pellet fully). Supernatants were discarded and beads were washed twice with 1 mL ice-cold cell lysis buffer (50 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 1 mM PMSF, pH 8.5). 360 μg proteins in 500 μL total volume (x μL supernatant from cell lysis with 360 μg proteins and 500-x μL cell lysis buffer) were incubated on a tube rotator overnight at 4°C. Streptavidin beads were washed to remove nonspecific binders the next morning as follows: 2x 1 mL cell lysis buffer, 1x 1 mL 1 M KCl, 1x 1 mL 0.1 M Na_2CO_3 (pH 11.5), 1x 1 mL 2 M urea (pH 8.0) in 10 mM Tris-HCl, 2x 1 mL cell lysis buffer, 2x 1 mL PBS (pH 7.4). PBS after the final washing step was removed before storing beads at -80°C. Notes: All streptavidin beads collection steps were 5 min long. All the

1040 washing solutions were ice-cold. Microcentrifuge tubes were either very briefly vortexed (~2 sec)

1041 or tapped by hand between each washing step to promote bead resuspension.

1042

On-bead digestion and TMT labeling: Samples were prepared as previously described

(*Kalocsay, 2019*). Liquid reagents used were HPLC quality grade. Washed beads were

resuspended in 50 µL of 200 mM EPPS (4-(2-Hydroxyethyl)-1-piperazinepropanesulfonic acid)

buffer (pH 8.5) and 2% (v/v) acetonitrile with 1 µL of 2 mg/mL lysil endoproteinase Lys-C stock

solution (FUJIFILM Wako Pure Chemical Corporation, Richmond, VA, Catalog #125-05061),

briefly vortexed, and incubated at 37°C for 3 hours. 50 µL of trypsin stock (Promega, Madison,

WI, Catalog #V5111) diluted 1:100 (v/v) in 200 mM EPPS (pH 8.5) was then added. After mixing,

digests were incubated at 37°C overnight and beads were magnetically removed. Peptides were

then directly labeled as follows: acetonitrile was added to 30% (v/v) concentration and peptides

were labeled with TMT 10-plex reagent (Thermo Fisher Scientific, Catalog #90406) for 1 hour.

Labeling reactions were quenched with hydroxylamine at a final concentration of 0.3% (v/v) for

15 min and 1% of labeled peptides was analyzed for label incorporation efficiency by mass

spectrometry. After quenching, peptide solutions were first acidified with formic acid,

trifluoroacetic acid (TFA) was then added to a concentration of 0.1% (v/v), and peptides were

desalted by acidic C_{18} solid phase extraction (StageTip). Labeled peptides were finally

resuspended in 1% (v/v) formic acid and 3% (v/v) acetonitrile.

1059

Mass spectrometry: Data were collected with a MultiNotch MS3 TMT method (*McAlister et al.,*

2014) using an Orbitrap Lumos mass spectrometer coupled to a Proxeon EASY-nLC 1200 Liquid

Chromatography (LC) system (both Thermo Fisher Scientific). The used capillary column was

packed with C_{18} resin (35 cm length, 100 µm inner diameter, 2.6 µm Accucore matrix (Thermo

Fisher Scientific)). Peptides were separated for 3 or 4 hours over acidic acetonitrile gradients by

LC prior to mass spectrometry (MS) analysis. Data from two 4 hour runs and one 3 hour run were

recorded and combined. After an initial MS[1] scan (Orbitrap analysis; resolution 120,000; mass

range 400−1,400 Th), MS[2] analysis used collision-induced dissociation (CID, CE = 35) with a

maximum ion injection time of 150–300 ms and an isolation window of 0.5 m/z. In order to obtain

quantitative information, MS[3] precursors were fragmented by high-energy collision-induced

dissociation (HCD) and analyzed in the Orbitrap at a resolution of 50,000 at 200 Th. Further details

on LC and MS parameters and settings were described recently (*Paulo et al., 2016*). The mass

spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the

PRIDE partner repository (*Perez-Riverol et al., 2019*) with the dataset identifier PXD018022.

pTF-mCherry and MS hit-EYFP co-expression and imaging

Sequence-validated episomes with pTF-mCherry and MS hit-EYFP expression cassettes were conjugated into WT *P. tricornutum* cells, resulting transconjugants genotyped, and screened for fluorescence. 5 µL of fluorescent transconjugant cell lines in mid- to late-exponential phase were imaged with settings that minimized cross-channel bleed-through: argon laser strength at 30%, 514 nm laser line at 50% maximum strength, mCherry emission window: 620–640 nm, EYFP emission window: 520–540 nm, autofluorescence emission window: 700–750 nm.

pTF.CREG1 protein expression conditions

Small scale expression testing for the Δ31_pTF.CREG1-His6 (and all other pTF.CREG1 truncations not described here) was performed as follows: an overnight culture of BL21 *E. coli* cells carrying the PtpBAD-Δ31_pTF.CREG1-CTHF (=pJT_Δ31_pTF.CREG1-His6) expression vector was used to inoculate 50 mL of Terrific Broth (TB) supplemented with tetracycline (10 ug/mL). Cultures were grown in a shaking incubator (37°C, 200 rpm) until OD_{600} of 0.4–0.6 was reached. Incubator temperature was then lowered to 18°C and flasks were allowed to adjust to this temperature for about 30 min before arabinose was added to a final concentration of 0.5% (w/v). Growth at 18°C was continued overnight (12–18 hrs) after which 10 mL of cultures were harvested by centrifugation at 6,000 g for 10 min. Large-scale Δ31_pTF.CREG1-His6 expression was performed as above, but at 2 L total volume. Following arabinose induction and overnight growth of the 2 L culture, 500 mL was harvested for further processing.

pTF.CREG1 purification conditions

All purifications were performed using cobalt, Co^{2+}, TALON Metal Affinity Resin (Takara Bio, Catalog #635502) which binds His6-tagged proteins with high affinity. Pellets from small scale expression testing (10 mL culture) were resuspended in 800 uL of lysis buffer (50 mM NaH_2PO_4 (pH 7.5), 500 mM NaCl, 0.1% (v/v) Triton X-100, 10 mM imidazole, 1 mg/mL lysozyme, and 10 µM β-mercaptoethanol) and subjected to microtip sonication on ice until lysis was complete. Lysates were then clarified by centrifugation (10 min, 15,000 g, 4°C) and supernatants (700 uL) were set aside. For each purification, 25 uL of TALON resin was equilibrated by washing three times with 10 volumes of lysis buffer and pelleting by centrifugation after each wash (30 sec, 3,000 g, RT). After third wash, 25 uL of lysis buffer was added to the resin to make a 50 uL slurry.

Supernatants from previously clarified cell lysates were then added to the TALON resin slurry and

incubated for 1 hr at RT with end-over-end mixing. Following the 1 hr incubation, the resin was

pelleted by centrifugation (30 sec, 3,000 g, RT) and washed three times with 10 volumes of wash

buffer (50 mM NaH_2PO_4 (pH 7.5), 500 mM NaCl, 30 mM imidazole). Proteins were eluted with 50

uL of elution buffer (50 mM NaH_2PO_4 (pH 7.5), 500 mM NaCl, 250 mM imidazole, 10% glycerol)

and were either subjected to immediate SDS-PAGE electrophoresis or stored at -80°C.

1114

Pellets from large-scale expression (500 mL harvested bacterial culture) were processed with a

few modifications. For lysis, cell pellets were resuspended in 20 mL of lysis buffer. For purification,

1 mL of TALON resin (i.e., 2 mL of equilibrated slurry) was used in combination with approx. 19

mL of clarified cell lysate which was allowed to incubate overnight at 8°C with end-over-end

mixing. Final protein elution was performed using 5 mL of elution buffer. Purified protein was

subjected to SDS-PAGE electrophoresis and concentrated using an Amicon Ultra-15 10 kDa

cutoff concentrator (Millipore Sigma, Burlington, MA, Catalog #UFC901024). Following the

concentration step, buffer exchange into 20 mM HEPES (pH 8.0), 10% glycerol, 300 mM KCl was

performed using a PD-10 desalting column (GE Healthcare, Chicago, IL, Catalog #17-0851-01).

Protein was then concentrated again and total protein content was determined by the Bradford

method using the Protein Assay Dye Reagent (Bio-Rad, Hercules, CA, Catalog #5000006).

1126

1127 **Flavin reduction assays**

1128

To test for flavin reductase activity, we set up the following enzyme assay conditions in 100 uL

final volume: 50 mM Tris-HCl (pH 7.5), 100 mM KCl, 10% glycerol, 30 µM flavin mononucleotide

(FMN), 30 µM riboflavin, and 250 µM NADPH (*Coves and Fontecave, 1993*). Assays were

started by the addition of 50 µg of pTF.CREG1, BSA, or Milli-Q water for no-enzyme controls.

Oxidation of NADPH was measured by the decrease in absorbance at 320 nm on a Flexstation 3

microtiter plate reader and the signal for each reaction was tracked over a period of 60 min. All

assays were conducted in triplicate.

1136

1137 **Bioinformatic and data analyses**

1138

1139 **Mass spectrometry data analysis:** Peptide-spectrum matches used a SEQUEST (v. 28, rev.

1140 12) algorithm (*Eng et al., 1994*). Data were searched against a size-sorted forward and reverse

1141 database of the *Phaeodactylum tricornutum* proteome (strain CCAP 1055/1, UniProt reference

proteome UP000000759) with added common contaminant proteins and the pTF-APEX2 fusion protein sequence. Spectra were first converted to mzXML and searches were then performed using a mass tolerance of 50 ppm for precursors and a fragment ion tolerance of 0.9 Da. For the searches, maximally two missed cleavages per peptide were allowed. We searched dynamically for oxidized methionine residues (+15.9949 Da) and applied a target decoy database strategy. A false discovery rate (FDR) of 1% was set for peptide-spectrum matches following filtering by linear discriminant analysis (LDA) (*Beausoleil et al., 2006*; *Huttlin et al., 2010*). The FDR for final collapsed proteins was 1%. Quantitative peptide information was derived from MS^3 scans. Quant tables were generated with the following filter criteria: MS^2 isolation specificity of >70% for each peptide and a sum of TMT signal-to-noise (s/n) of >200 over all channels per peptide. Quant tables were exported to Excel and further processed therein. Details of the TMT intensity quantification method and additional applied search parameters were described previously (*Paulo et al., 2016*). Scaled proteomic data were subjected to two-way hierarchical clustering (Ward's method) using JMP software package (SAS Institute). Volcano plot with log_2-transformed average APEX2/WT ratios and associated P values was made in R using ggplot2 data visualization package (*R Core Team, 2013*). Gene IDs corresponding to protein hits were inferred using Ensembl Protists Phatr3 *P. tricornutum* genomic database.

Protein feature identification: Protein lengths, molecular weights, and isoelectric points (pI) were determined with ProtParam (*Gasteiger et al., 2005*). Signal peptides and transmembrane regions were identified with SignalP 4.1 (*Petersen et al., 2011*) and TMHMM Server v. 2.0 (*Krogh et al., 2001*), respectively. Protein localizations were predicted with a combination of tools: TargetP 1.1 (*Emanuelsson et al., 2000*), SignalP 4.1, and ASAFind (*Gruber et al., 2015*) version 1.1.7. All putative chloroplastic localizations in the study mean that a protein was predicted to be chloroplastic by ASAFind (state-of-the-art plastidial protein localization prediction tool for diatoms); chloroplast localization prediction confidences are noted. Low confidence prediction by ASAFind means that a protein satisfies the following filtering criteria: (1) it contains a signal peptide as detected by SignalP 4.1, (2) +1 position of ASAFind predicted cleavage site is phenylalanine (F), tryptophan (W), tyrosine (Y), or leucine (L) (making the protein "potentially plastid targeted"; it was tyrosine in pTF.CREG1), and (3) one or both of the following is false: the ASAFind predicted cleavage site coincides with the SignalP 4.1 prediction and the transit peptide score is higher than 2 (the latter is false for pTF.CREG1: transit peptide score was ~0.81). Peroxidase class prediction was done in RedoxiBase (*Savelli et al., 2019*). N-glycosylation and

1175 GalNAc-type O-glycosylation predictions were performed with NetNGlyc 1.0 (*Gupta et al., 2004*)

1176 and NetOGlyc 4.0 (*Steentoft et al., 2013*), respectively.

1177

Amino acid sequence alignments: APEX2-like *P. tricornutum* peroxidases were aligned with

Clustal Omega (*Sievers et al., 2011*) using the default settings and CLC Sequence Viewer 7.7

(QUIAGEN) using the following parameters: Gap open cost: 10.0, Gap extension cost: 1.0, End

gap cost: As any other, Alignment: Very accurate (slow). Putative substrate-binding loops in these

peroxidases were evaluated using the Clustal Omega alignment. Conserved motifs in amino acid

sequence alignments underlying the phylogenetic trees were displayed in CLC Sequence Viewer

7.7.

1185

Phylogenetic analyses: Homologs of respective *P. tricornutum* proteins were retrieved from the

non-redundant National Center for Biotechnology Information (NCBI) and the Marine Microbial

Eukaryote Transcriptome Sequencing Project (MMETSP) databases using the BLASTP algorithm

(e-value threshold set to e^{-15}) (*Altschul et al., 1990*). The BLAST search retrieved only a handful

of homologs for pTF.ap1 and no homologs for pTF.CatCh1 even among closely related diatoms.

Therefore, more sensitive HMMER (*Eddy, 1998*) searches were employed to extend the datasets,

which were afterward aligned using the Localpair algorithm as implemented in MAFFT (*Katoh et*

al., 2002). Ambiguously aligned regions, regions composed mostly of gaps, and short fragments

were manually removed in SeaView 4 (*Gouy et al., 2010*). For each alignment, the maximum

likelihood analysis was carried out in IQ-TREE (*Nguyen et al., 2015*) under the best-fitting

substitution matrix as inferred by the built-in model finder. Branching support was estimated using

"thorough" non-parametric bootstrap analysis from 500 replicates in IQ-TREE.

1198

1199 **Identification of disordered protein regions:** pTF.CatCh1 amino acid sequence was analyzed

1200 for the presence of disordered protein regions with PONDR (Predictor of Naturally Disordered

1201 Regions) VSL2 predictor (*Peng et al., 2006*) and IUPred2a long disorder prediction type

1202 (*Mészáros et al., 2018*).

1203

Homology modeling of pTF.CREG1: Full-length pTF.CREG1 amino acid sequence was used

as an input for Phyre2, an online protein structure prediction server (*Kelley et al., 2015*). Normal

modeling mode was selected. Human CREG1 (PDB ID: 1XHN) was identified as the best

modeling template (human CREG1 is ~37% identical to pTF.CREG1 as inferred from their Clustal

Omega alignment). pTF.CREG1 was modeled with >90% confidence across 182 modeled amino

acid residues (Pro43–Lys224) indicating the predicted structure is representative of the actual

structure. pTF.CREG1 homology model was aligned with human CREG1 monomer in UCSF

Chimera version 1.11.1 using MatchMaker tool with default settings (*Pettersen et al., 2004*)

1212

1213 **Other data analyses:** Amplex UltraRed assay and transcriptomic data were plotted in R using

1214 ggplot2 data visualization package (*R Core Team, 2013*). Flavin reduction assay data were

1215 plotted in Excel.

Acknowledgements

We would like to thank Andrea Thor and Mason Mackey for help with electron microscopy sample preparation and imaging; Marian Kalocsay for assistance with mass spectrometry experiments and data analyses; Pardis Gholami and Hong Zheng for laboratory assistance; Jeffrey D. Martell, Jeffrey B. McQuaid, Tyler H. Coale, and Sarah R. Smith for fruitful discussions. This project was supported by the Gordon and Betty Moore Foundation grants GBMF4958 (JT), GBMF3828 (AEA), and GBMF5006 (AEA), the National Science Foundation grants NSF-OCE-1756884 (AEA) and NSF-MCB-1818390 (AEA), the United States Department of Energy Genomics Science program grant DE-SC0018344 (AEA), the Czech Science Foundation grant 18-13458S (MO), and ERDF/ESF Centre for Research of Pathogenicity and Virulence of Parasites grant No.CZ.02.1.01/0.0/0.0/16_019/0000759 (MO). JKB was supported by the National Institutes of Health (NIH) Ruth L. Kirschstein Predoctoral Individual National Research Service Award F31 1F31ES030613-01. National Center for Microscopy and Imaging Research (NCMIR) is supported by the National Institute for General Medical Sciences (NIGMS) of the National Institutes of Health (NIH).

Additional information

Funding

Funder	Grant reference number	Author
Gordon and Betty Moore Foundation (GBMF), Marine Microbial Initiative (MMI)	GBMF4958	Jernej Turnšek*
National Institutes of Health (NIH) Ruth L. Kirschstein Predoctoral Individual National Research Service Award F31	1F31ES030613-01	John K. Brunson
The Czech Science Foundation	18-13458S	Miroslav Oborník
ERDF/ESF Centre for Research of Pathogenicity and Virulence of Parasites	No.CZ.02.1.01/0.0/0.0/ 16_019/0000759	Miroslav Oborník
Gordon and Betty Moore Foundation (GBMF), Marine Microbial Initiative (MMI)	GBMF3828	Andrew E. Allen
Gordon and Betty Moore Foundation (GBMF), Marine Microbial Initiative (MMI)	GBMF5006	Andrew E. Allen
National Science Foundation, Biological Oceanography	NSF-OCE-1756884	Andrew E. Allen
National Science Foundation, Molecular and Cellular Biology	NSF-MCB-1818390	Andrew E. Allen
Department of Energy (DOE), Office of Biological and Environmental Research (BER)	DE-SC0018344	Andrew E. Allen

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

*Jernej Turnšek originated and proposed the idea to implement the use of APEX2 in model marine diatoms and wrote the proposal jointly with James J. Russell, Pamela A. Silver, and Julie A. Theriot.

Author ORCIDs

1241 Jernej Turnšek   https://orcid.org/0000-0002-9056-3565

1242 Maria del Pilar Martinez Viedma   https://orcid.org/0000-0002-8885-5008

1243 Aleš Horák   https://orcid.org/0000-0002-3364-7383

1244 Vincent A. Bielinski   https://orcid.org/0000-0002-8107-8752

1245 Andrew E. Allen   https://orcid.org/0000-0001-5911-6081

Additional files

Supporting files

Supplemental file 1.

Table S1: Predicted endogenous biotinylated proteins are present at similar levels in WT and pTF-APEX2 proteomic samples. **Table S2:** Features of the three proteins co-expressed from a gene cluster on chromosome 20. **Table S3:** Molecular cloning primers. **Table S4:** Constructed *Phaeodactylum tricornutum* episomes and *Escherichia coli* vectors with expression cassette details and fusion protein amino acid sequences.

Key Resources Table.

The quantitative mass spectrometry proteomics data used to generate *Figure 4—source data 1* and *Figure 4—figure supplement 1—source data 1* have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository (*Perez-Riverol et al., 2019*) with the dataset identifier PXD018022. Data can be accessed via a reviewer account using Username: reviewer15941@ebi.ac.uk and Password: ZCd0fJ8p.

The amino acid sequence alignment LINSI files *Figure 6—source data 1*, *Figure 7—source data 1*, and *Figure 7—figure supplement 2—source data 1* can be read and interrogated with AlignmentViewer available at https://alignmentviewer.org/.

Data availability

All data generated or analyzed during this study are included in the manuscript and supporting files.

Competing interests

The authors declare that no competing interests exist.

References

Aguirre JD, Clark HM, McIlvin M, Vazquez C, Palmere SL, Grab DJ, Seshu J, Hart PJ, Saito M, Culotta VC. 2013. A Manganese-rich Environment Supports Superoxide Dismutase Activity in a Lyme Disease Pathogen, *Borrelia burgdorferi*. *J Biol Chem* **288**:8468–8478. DOI: https://doi.org/10.1074/jbc.M112.433540, PMID: 23376276

Allen AE, Laroche J, Maheswari U, Lommer M, Schauer N, Lopez PJ, Finazzi G, Fernie AR, Bowler C. 2008. Whole-cell response of the pennate diatom *Phaeodactylum tricornutum* to iron starvation. *Proc Natl Acad Sci U S A* **105**:10438–10443. DOI: https://doi.org/10.1073/pnas.0711370105, PMID: 18653757

Allen AE, Moustafa A, Montsant A, Eckert A, Kroth PG, Bowler C. 2012. Evolution and Functional Diversification of Fructose Bisphosphate Aldolase Genes in Photosynthetic Marine Diatoms. *Mol Biol Evol* **29**:367–379. DOI: https://doi.org/10.1093/molbev/msr223, PMID: 21903677

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* **215**:403–410. DOI: https://doi.org/10.1016/S0022-2836(05)80360-2, PMID: 2231712

Archibald JM. 2009. The puzzle of plastid evolution. *Curr Biol* **19**:R81–8. DOI: https://doi.org/10.1016/j.cub.2008.11.067, PMID: 19174147

Aron AT, Ramos-Torres KM, Cotruvo JA Jr, Chang CJ. 2015. Recognition- and Reactivity-Based Fluorescent Probes for Studying Transition Metal Signaling in Living Systems. *Acc Chem Res* **48**:2434–2442. DOI: https://doi.org/10.1021/acs.accounts.5b00221, PMID: 26215055

Ardyna M, Lacour L, Sergi S, d'Ovidio F, Sallée J-B, Rembauville M, Blain S, Tagliabue A, Schlitzer R, Jeandel C, Arrigo KR, Claustre H. 2019. Hydrothermal vents trigger massive phytoplankton blooms in the Southern Ocean. *Nat Commun* **10**:2451. DOI: https://doi.org/10.1038/s41467-019-09973-6, PMID: 31165724

Balk J, Schaedler TA. 2014. Iron Cofactor Assembly in Plants. *Annu Rev Plant Biol* **65**:125–153. DOI: https://doi.org/10.1146/annurev-arplant-050213-035759, PMID: 24498975

Banci L, Bertini I, Ciofi-Baffoni S, Finney LA, Outten CE, O'Halloran TV. 2002. A New Zinc-protein Coordination Site in Intracellular Metal Trafficking: Solution Structure of the Apo and Zn(II) forms of ZntA(46-118). *J Mol Biol* **323**:883–897. DOI: https://doi.org/10.1016/s0022-2836(02)01007-0, PMID: 12417201

Banci L, Bertini I, Ciofi-Baffoni S, Su X-C, Miras R, Bal N, Mintz E, Catty P, Shokes JE, Scott RA. 2006. Structural basis for metal binding specificity: the N-terminal cadmium binding domain of the P1-type ATPase CadA. *J Mol Biol* **356**:638–650. DOI: https://doi.org/10.1016/j.jmb.2005.11.055, PMID: 16388822

Bard F, Chia J. 2016. Cracking the Glycome Encoder: Signaling, Trafficking, and Glycosylation. *Trends Cell Biol* **26**:379–388. DOI: https://doi.org/10.1016/j.tcb.2015.12.004, PMID: 26832820

Barrett J, Girr P, Mackinder LCM. 2021. Pyrenoids: CO_2-fixing phase separated liquid
organelles. *Biochim Biophys Acta Mol Cell Res* **1868**:118949. DOI:
https://doi.org/10.1016/j.bbamcr.2021.118949, PMID: 33421532

Baslam M, Oikawa K, Kitajima-Koga A, Kaneko K, Mitsui T. 2016. Golgi-to-plastid trafficking of
proteins through secretory pathway: Insights into vesicle-mediated import toward the plastids.
Plant Signal Behav **11**:e1221558. DOI: https://doi.org/10.1080/15592324.2016.1221558, PMID:
27700755

Beausoleil SA, Villén J, Gerber SA, Rush J, Gygi SP. 2006. A probability-based approach for
high-throughput protein phosphorylation analysis and site localization. *Nat Biotechnol* **24**:1285–
1292. DOI: https://doi.org/10.1038/nbt1240, PMID: 16964243

Behnke J, LaRoche J. 2020. Iron uptake proteins in algae and the role of Iron Starvation-
Induced Proteins (ISIPs). *Eur J Phycol* **55**:339–360. DOI:
https://doi.org/10.1080/09670262.2020.1744039

Benham AM. 2012. Protein Secretion and the Endoplasmic Reticulum. *Cold Spring Harb*
Perspect Biol **4**:a012872. DOI: https://doi.org/10.1101/cshperspect.a012872, PMID: 22700933

Benoiston A-S, Ibarbalz FM, Bittner L, Guidi L, Jahn O, Dutkiewicz S, Bowler C. 2017. The
evolution of diatoms and their biogeochemical functions. *Philos Trans R Soc Lond B Biol Sci*
372:20160397. DOI: https://doi.org/10.1098/rstb.2016.0397, PMID: 28717023

Bertrand EM, McCrow JP, Moustafa A, Zheng H, McQuaid JB, Delmont TO, Post AF, Sipler
RE, Spackeen JL, Xu K, Bronk DA, Hutchins DA, Allen AE. 2015. Phytoplankton–bacterial
interactions mediate micronutrient colimitation at the coastal Antarctic sea ice edge. *Proc Natl*
Acad Sci U S A **112**:9938–9943. DOI: https://doi.org/10.1073/pnas.1501615112, PMID:
26221022

Bhaya D, Grossman AR. 1993. Characterization of gene clusters encoding the fucoxanthin
chlorophyll proteins of the diatom *Phaeodactylum tricornutum*. *Nucleic Acids Res* **21**:4458–
4466. DOI: https://doi.org/10.1093/nar/21.19.4458, PMID: 8233779

Blaby-Haas CE, Merchant SS. 2012. The ins and outs of algal metal transport. *Biochim*
Biophys Acta **1823**:1531–1552. DOI: https://doi.org/10.1016/j.bbamcr.2012.04.010, PMID:
22569643

Blaby-Haas CE, Merchant SS. 2014. Lysosome-related Organelles as Mediators of Metal
Homeostasis. *J Biol Chem* **289**:28129–28136. DOI: https://doi.org/10.1074/jbc.R114.592618,
PMID: 25160625

Blaby-Haas CE, Padilla-Benavides T, Stübe R, Argüello JM, Merchant SS. 2014. Evolution of a
plant-specific copper chaperone family for chloroplast copper homeostasis. *Proc Natl Acad Sci*
U S A **111**:E5480–E5487. DOI: https://doi.org/10.1073/pnas.1421545111, PMID: 25468978

Bogdan AR, Miyazawa M, Hashimoto K, Tsuji Y. 2016. Regulators of Iron Homeostasis: New
Players in Metabolism, Cell Death, and Disease. *Trends Biochem Sci* **41**:274–286. DOI:
https://doi.org/10.1016/j.tibs.2015.11.012, PMID: 26725301
1374

1375 **Boiteau RM**, Mende DR, Hawco NJ, McIlvin MR, Fitzsimmons JN, Saito MA, Sedwick PN,
1376 DeLong EF, Repeta DJ. 2016. Siderophore-based microbial adaptations to iron scarcity across
1377 the eastern Pacific Ocean. *Proc Natl Acad Sci U S A* **113**:14237–14242. DOI:
1378 https://doi.org/10.1073/pnas.1608594113, PMID: 27911777
1379
1380 **Bonfio C**, Valer L, Scintilla S, Shah S, Evans DJ, Jin L, Szostak JW, Sasselov DD, Sutherland
1381 JD, Mansy SS. 2017. UV-light-driven prebiotic synthesis of iron-sulfur clusters. *Nat Chem*
1382 **9**:1229–1234. DOI: https://doi.org/10.1038/nchem.2817, PMID: 29168482
1383
1384 **Boukouris AE**, Zervopoulos SD, Michelakis ED. 2016. Metabolic Enzymes Moonlighting in the
1385 Nucleus: Metabolic Regulation of Gene Transcription. *Trends Biochem Sci* **41**:712–730. DOI:
1386 https://doi.org/10.1016/j.tibs.2016.05.013, PMID: 27345518
1387
1388 **Bowler C**, (75 authors), Grigoriev IV. 2008. The *Phaeodactylum* genome reveals the
1389 evolutionary history of diatom genomes. *Nature* **456**:239–244. DOI:
1390 https://doi.org/10.1038/nature07410, PMID: 18923393
1391
1392 **Branon TC**, Bosch JA, Sanchez AD, Udeshi ND, Svinkina T, Carr SA, Feldman JL, Perrimon N,
1393 Ting AY. 2018. Efficient proximity labeling in living cells and organisms with TurboID. *Nat*
1394 *Biotechnol* **36**:880–887. DOI: https://doi.org/10.1038/nbt.4201, PMID: 30125270
1395
1396 **Brenner MP**. 2012. Endocytic Traffic: Vesicle Fusion Cascade in the Early Endosomes. *Curr*
1397 *Biol*. **22**:R597–598. DOI: https://doi.org/10.1016/j.cub.2012.06.009, PMID: 22877778
1398
1399 **Brunson JK**, McKinnie SMK, Chekan JR, McCrow JP, Miles ZD, Bertrand EM, Bielinski VA,
1400 Luhavaya H, Oborník M, Smith GJ, Hutchins DA, Allen AE, Moore BS. 2018. Biosynthesis of the
1401 neurotoxin domoic acid in a bloom-forming diatom. *Science* **361**:1356–1358. DOI:
1402 https://doi.org/10.1126/science.aau0382, PMID: 30262498
1403
1404 **Brzezinski MA**, Krause JW, Bundy RM, Barbeau KA, Franks P, Goericke R, Landry MR, Stukel
1405 MR. 2015. Enhanced silica ballasting from iron stress sustains carbon export in a frontal zone
1406 within the California Current. *J Geophys Res C: Oceans* **120**:4654–4669. DOI:
1407 https://doi.org/10.1002/2015JC010829
1408
1409 **Camacho A**, Walter XA, Picazo A, Zopfi J. 2017. Photoferrotrophy: Remains of an Ancient
1410 Photosynthesis in Modern Environments. *Front Microbiol* **8**:323. DOI:
1411 https://doi.org/10.3389/fmicb.2017.00323, PMID: 28377745
1412
1413 **Cao H**, Schroeder B, Chen J, Schott MB, McNiven MA. 2016. The Endocytic Fate of the
1414 Transferrin Receptor Is Regulated by c-Abl Kinase. *J Biol Chem* **291**:16424–16437. DOI:
1415 https://doi.org/10.1074/jbc.M116.724997, PMID: 27226592
1416
1417 **Caron DA**, Alexander H, Allen AE, Archibald JM, Armbrust EV, Bachy C, Bell CJ, Bharti A,
1418 Dyhrman ST, Guida SM, Heidelberg KB, Kaye JZ, Metzner J, Smith SR, Worden AZ. 2017.
1419 Probing the evolution, ecology and physiology of marine protists using transcriptomics. *Nat Rev*
1420 *Microbiol* **15**:6–20. DOI: https://doi.org/10.1038/nrmicro.2016.160, PMID: 27867198
1421
1422 **Carradec Q**, (37 authors), Wincker P. 2018. A global ocean atlas of eukaryotic genes. *Nat*
1423 *Commun* **9**:373. DOI: https://doi.org/10.1038/s41467-017-02342-1, PMID: 29371626
1424

Cheng Y, Zak O, Aisen P, Harrison SC, Walz T. 2004. Structure of the Human Transferrin Receptor-Transferrin Complex. *Cell* **116**:565–576. DOI: https://doi.org/10.1016/s0092-8674(04)00130-8, PMID: 14980223

Cieśla M, Mierzejewska J, Adamczyk M, Farrants A-KÖ, Boguta M. 2014. Fructose bisphosphate aldolase is involved in the control of RNA polymerase III-directed transcription. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* **1843**:1103–1110. DOI: https://10.1016/j.bbamcr.2014.02.007, PMID: 24576411

Coale TH, Mooseburner M, Horák A., Oborník M., Barbeau KA, Allen AE. 2019. Reduction-dependent siderophore assimilation in a model pennate diatom. *Proc Natl Acad Sci U S A*. **116**:23609–23617. DOI: https://doi.org/10.1073/pnas.1907234116, PMID: 31685631

Cohen NR, A. Ellis K, Burns WG, Lampe RH, Schuback N, Johnson Z, Sañudo-Wilhelmy S, Marchetti A. 2017. Iron and vitamin interactions in marine diatom isolates and natural assemblages of the Northeast Pacific Ocean: Iron and vitamin interactions in diatoms. *Limnol Oceanogr* **62**:2076–2096. DOI: https://doi.org/10.1002/lno.10552

Cohen NR, Ellis KA, Lampe RH, McNair H, Twining BS, Maldonado MT, Brzezinski MA, Kuzminov FI, Thamatrakoln K, Till CP, Bruland KW, Sunda WG, Bargu S, Marchetti A. 2017. Diatom Transcriptional and Physiological Responses to Changes in Iron Bioavailability across Ocean Provinces. *Frontiers in Marine Science* **4**:360. DOI: https://doi.org/10.3389/fmars.2017.00360

Cohen NR, Gong W, Moran DM, Mcilvin MR, Saito MA, Marchetti A. 2018. Transcriptomic and proteomic responses of the oceanic diatom *Pseudo-nitzschia granii* to iron limitation. *Environ Microbiol* **20**:3109–3126. DOI: https://doi.org/10.1111/1462-2920.14386, PMID: 30117243

Coves J, Fontecave M. 1993. Reduction and mobilization of iron by a NAD(P)H:flavin oxidoreductase from *Escherichia coli*. *Eur J Biochem* **211**:635–641. DOI: https://doi.org/10.1111/j.1432-1033.1993.tb17591.x, PMID: 8436123

Crichton R. 2016. Iron Metabolism: From Molecular Mechanisms to Clinical Consequences, Fourth Edition. John Wiley & Sons, Inc., Hoboken, NJ, USA.1–556. DOI: https://doi.org/10.1002/9781118925645

de Baar HJW, (32 authors), Wong CS. 2005. Synthesis of iron fertilization experiments: From the Iron Age in the Age of Enlightenment. *J Geophys Res* **110**:C09S16. DOI: https://doi.org/10.1029/2004JC002601

Del Olmo T, Lauzier A, Normandin C, Larcher R, Lecours M, Jean D, Lessard L, Steinberg F, Boisvert F-M, Jean S. 2019. APEX2-mediated RAB proximity labeling identifies a role for RAB21 in clathrin-independent cargo sorting. *EMBO Rep* **20**:e47192. DOI: https://doi.org/10.15252/embr.201847192, PMID: 30610016

Di Bacco A, Gill G. 2003. The secreted glycoprotein CREG inhibits cell growth dependent on the mannose-6-phosphate/insulin-like growth factor II receptor. *Oncogene* **22**:5436–5445. DOI: https:// doi.org/10.1038/sj.onc.1206670, PMID: 12934103

1475 **Diner RE**, Bielinski VA, Dupont CL, Allen AE, Weyman PD. 2016. Refinement of the Diatom
1476 Episome Maintenance Sequence and Improvement of Conjugation-Based DNA Delivery
1477 Methods. *Front Bioeng Biotechnol* **4**:65. DOI: https:// doi.org/10.3389/fbioe.2016.00065, PMID:
1478 27551676

1479

1480 **Diner RE**, Noddings CM, Lian NC, Kang AK, McQuaid JB, Jablanovic J, Espinoza JL, Nguyen
1481 NA, Anzelmatti MA Jr, Jansson J, Bielinski VA, Karas BJ, Dupont CL, Allen AE, Weyman PD.
1482 2017. Diatom centromeres suggest a mechanism for nuclear DNA acquisition. *Proc Natl Acad*
1483 *Sci U S A* **114**:E6015–E6024. DOI: https:// doi.org/10.1073/pnas.1700764114, PMID: 28673987

1484

1485 **Dokmanić I**, Sikić M, Tomić S. 2008. Metals in proteins: correlation between the metal-ion type,
1486 coordination number and the amino-acid residues involved in the coordination. *Acta Crystallogr*
1487 *D Biol Crystallogr* **64**:257–263. DOI: https:// doi.org/10.1107/S090744490706595X, PMID:
1488 18323620

1489

1490 **Durkin CA**, Marchetti A, Bender SJ, Truong T, Morales R, Mock T, Armbrust EV. 2012.
1491 Frustule-related gene transcription and the influence of diatom community composition on silica
1492 precipitation in an iron-limited environment. *Limnol Oceanogr* **57**:1619–1633. DOI:
1493 https://doi.org/10.4319/lo.2012.57.6.1619

1494

1495 **Eckenroth BE**, Steere AN, Chasteen ND, Everse SJ, Mason AB. 2011. How the binding of
1496 human transferrin primes the transferrin receptor potentiating iron release at endosomal pH.
1497 *Proc Natl Acad Sci U S A* **108**:13089–13094. DOI: https://doi.org/10.1073/pnas.1105786108,
1498 PMID: 21788477

1499

1500 **Eddy SR**. 1998. Profile hidden Markov models. *Bioinformatics* **14**:755–763. DOI:
1501 https://doi.org/10.1093/bioinformatics/14.9.755, PMID: 9918945

1502

1503 **El-Gebali S**, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M, Richardson LJ,
1504 Salazar GA, Smart A, Sonnhammer ELL, Hirsh L, Paladin L, Piovesan D, Tosatto SCE, Finn
1505 RD. 2019. The Pfam protein families database in 2019. *Nucleic Acids Res* **47**:D427–D432. DOI:
1506 https://doi.org/10.1093/nar/gky995, PMID: 30357350

1507

1508 **Emanuelsson O**, Nielsen H, Brunak S, von Heijne G. 2000. Predicting Subcellular Localization
1509 of Proteins Based on their N-terminal Amino Acid Sequence. *J Mol Biol* **300**:1005–1016. DOI:
1510 https://doi.org/10.1006/jmbi.2000.3903, PMID: 10891285

1511

1512 **Eng JK**, McCormack AL, Yates JR. 1994. An approach to correlate tandem mass spectral data
1513 of peptides with amino acid sequences in a protein database. *J Am Soc Mass Spectrom* **5**:976–
1514 989. DOI: https://doi.org/10.1016/1044-0305(94)80016-2, PMID: 24226387

1515

1516 **Faktorová D**, (104 authors), Lukeš, J. 2020. Genetic tool development in marine protists:
1517 Emerging model organisms for experimental cell biology. *Nat Methods* **17**:481–494. DOI:
1518 https://doi.org/10.1038/s41592-020-0796-x, PMID: 32251396

1519

1520 **Falciatore A**, Jaubert M, Bouly J-P, Bailleul B, Mock T. 2020. Diatom Molecular Research
1521 Comes of Age: Model Species for Studying Phytoplankton Biology and Diversity. *Plant Cell*
1522 **32**:547–572. DOI: https://doi.org/10.1105/tpc.19.00158, PMID: 31852772

1523

1524 **Finbow ME**, Harrison MA. 1997. The vacuolar H^+-ATPase: a universal proton pump of
1525 eukaryotes. *Biochem J* **324**:697–712. DOI: https://doi.org/10.1042/bj3240697, PMID: 9210392

Fomenko DE, Gladyshev VN. 2003. Identity and Functions of CxxC-Derived Motifs. *Biochemistry* **42**:11214–11225. DOI: https://doi.org/10.1021/bi034459s, PMID: 14503871

Fomenko DE, Marino SM, Gladyshev VN. 2008. Functional Diversity of Cysteine Residues in Proteins and Unique Features of Catalytic Redox-Active Cysteines in Thiol Oxidoreductases. *Mol Cells* **26**:228–235. PMID: 18648218

Francius G, Tesson B, Dague E, Martin-Jézéquel V, Dufrêne YF. 2008. Nanostructure and nanomechanics of live *Phaeodactylum tricornutum* morphotypes. *Environ Microbiol* **10**:1344–1356. DOI: https://doi.org/10.1111/j.1462-2920.2007.01551.x, PMID: 18248452

Gasteiger E, Hoogland C, Gattiker A, Duvaud S, Wilkins MR, Appel RD, Bairoch A. 2005. Protein Identification and Analysis Tools on the ExPASy Server. *(*In:*)* The Proteomics Protocols Handbook, (Editor:) John M. Walker. Humana Press, Inc., Totowa, NJ, USA. 571–607. DOI: https://doi.org/10.1385/1-59259-890-0:571

Ghobrial G, Araujo L, Jinwala F, Li S, Lee LY. 2018. The Structure and Biological Function of CREG. *Front Cell Dev Biol* **6**:136. DOI: https://doi.org/10.3389/fcell.2018.00136, PMID: 30416997

Ghosh P, Dahms NM, Kornfeld S. 2003. Mannose 6-phosphate receptors: new twists in the tale. *Nat Rev Mol Cell Biol* **4**:202–212. DOI: https://doi.org/10.1038/nrm1050, PMID: 12612639

Gibbs SP. 1981. The Chloroplast Endoplasmic Reticulum: Structure, Function, and Evolutionary Significance (In:) International Review of Cytology, (Editors:) Bourne GH, Danielli JF, Jeon KW. Academic Press, Inc., New York, NY, USA. 49–99. DOI: https://doi.org/10.1016/S0074-7696(08)61194-8

Gibson DG, Young L, Chuang R-Y, Craig Venter J, Hutchison CA, Smith HO. 2009. Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* **6**:343–345. DOI: https://doi.org/10.1038/nmeth.1318, PMID: 19363495

Gingras A-C, Abe KT, Raught B. 2019. Getting to know the neighborhood: using proximity-dependent biotinylation to characterize protein complexes and map organelles. *Curr Opin Chem Biol* **48**:44–54. DOI: https://10.1016/j.cbpa.2018.10.017, PMID: 30458335

Gizak A, Wiśniewski J, Heron P, Mamczur P, Sygusch J, Rakus D. 2019. Targeting a moonlighting function of aldolase induces apoptosis in cancer cells. *Cell Death Dis* **10**:712. DOI: https://doi.org/10.1038/s41419-019-1968-4, PMID: 31558701

Gladyshev VN, Kryukov GV, Fomenko DE, Hatfield DL. 2004. Identification of Trace Element-Containing Proteins in Genomic Databases. *Annu Rev Nutr* **24**:579–596. DOI: https://doi.org/10.1146/annurev.nutr.24.012003.132241, PMID: 15189132

Gouy M, Guindon S, Gascuel O. 2010. SeaView Version 4: A Multiplatform Graphical User Interface for Sequence Alignment and Phylogenetic Tree Building. *Mol Biol Evol* **27**:221–224. DOI: https://doi.org/10.1093/molbev/msp259, PMID: 19854763

Grant CR, Komeili A. 2020. Ferrosomes are iron storage organelles formed by broadly conserved gene clusters in bacteria and archaea. *bioRxiv*. DOI: https://doi.org/10.1101/2020.01.10.902569

Gröger P, Poulsen N, Klemm J, Kröger N, Schlierf M. 2016. Establishing super-resolution imaging for proteins in diatom biosilica. *Sci Rep* **6**:36824. DOI: https://doi.org/10.1038/srep36824, PMID: 27827427

Grosche C, Hempel F, Bolte K, Zauner S, Maier UG. 2014. The periplastidal compartment: a naturally minimized eukaryotic cytoplasm. *Curr Opin Microbiol* **22**:88–93. DOI: https://doi.org/10.1016/j.mib.2014.09.017, PMID: 25460801

Gruber A, Rocap G, Kroth PG, Armbrust EV, Mock T. 2015. Plastid proteome prediction for diatoms and other algae with secondary plastids of the red lineage. *Plant J* **81**:519–528. DOI: https://doi.org/10.1111/tpj.12734, PMID: 25438865

Gupta R, Jung E., Brunak S. 2004. Prediction of N-glycosylation sites in human proteins. In preparation. URL: https://services.healthtech.dtu.dk/service.php?NetNGlyc-1.0

Hamdi A, Roshan TM, Kahawita TM, Mason AB, Sheftel AD, Ponka P. 2016. Erythroid cell mitochondria receive endosomal iron by a "kiss-and-run" mechanism. *Biochim Biophys Acta* **1863**:2859–2867. DOI: https://doi.org/10.1016/j.bbamcr.2016.09.008, PMID: 27627839

Han Y, Cui J, Tao J, Guo L, Guo P, Sun M, Kang J, Zhang X, Yan C, Li S. 2009. CREG inhibits migration of human vascular smooth muscle cells by mediating IGF-II endocytosis. *Exp Cell Res* **315**:3301–3311. DOI: https://doi.org/10.1016/j.yexcr.2009.09.013, PMID: 19769965

Han Y, Luan B, Sun M, Guo L, Guo P, Tao J, Deng J, Wu G, Liu S, Yan C, Li S. 2011. Glycosylation-independent binding to extracellular domains 11-13 of mannose-6-phosphate/insulin-like growth factor-2 receptor mediates the effects of soluble CREG on the phenotypic modulation of vascular smooth muscle cells. *J Mol Cell Cardiol* **50**:723–730. DOI: https://doi.org/10.1016/j.yjmcc.2010.12.013, PMID: 21195083

Hertle AP, García-Cerdán JG, Armbruster U, Shih R, Lee JJ, Wong W, Niyogi KK. 2020. A Sec14 domain protein is required for photoautotrophic growth and chloroplast vesicle formation in Arabidopsis thaliana. *Proc Natl Acad Sci U S A* **117**:9101–9111. DOI: https://doi.org/10.1073/pnas.1916946117, PMID: 32245810

Hildebrand M, Lerch SJL, Shrestha RP. 2018. Understanding Diatom Cell Wall Silicification—Moving Forward. *Frontiers in Marine Science* **5**:125. DOI: https://doi.org/10.3389/fmars.2018.00125

Hogle SL, Barbeau KA, Gledhill M. 2014. Heme in the marine environment: from cells to the iron cycle. *Metallomics* **6**:1107–1120. DOI: https://doi.org/10.1039/c4mt00031e, PMID: 24811388

Hong-Hermesdorf A, Miethke M, Gallaher SD, Kropat J, Dodani SC, Chan J, Barupala D, Domaille DW, Shirasaki DI, Loo JA, Weber PK, Pett-Ridge J, Stemmler TL, Chang CJ, Merchant SS. 2014. Subcellular metal imaging identifies dynamic sites of Cu accumulation in *Chlamydomonas*. *Nat Chem Biol* **10**:1034–1042. DOI: https://doi.org/10.1038/nchembio.1662, PMID: 25344811

Hung V, Udeshi ND, Lam SS, Loh KH, Cox KJ, Pedram K, Carr SA, Ting AY. 2016. Spatially resolved proteomic mapping in living cells with the engineered peroxidase APEX2. *Nat Protoc* **11**:456–475. DOI: https://doi.org/10.1038/nprot.2016.018, PMID: 26866790

Hutchins DA, Boyd PW. 2016. Marine phytoplankton and the changing ocean iron cycle. *Nat Clim Chang* **6**:1072–1079. DOI: https://doi.org/10.1038/nclimate3147

Hutchins DA, Bruland KW. 1998. Iron-limited diatom growth and Si:N uptake ratios in a coastal upwelling regime. *Nature* **393**:561–564. DOI: https://doi.org/10.1038/31203

Huttlin EL, Jedrychowski MP, Elias JE, Goswami T, Rad R, Beausoleil SA, Villén J, Haas W, Sowa ME, Gygi SP. 2010. A Tissue-Specific Atlas of Mouse Protein Phosphorylation and Expression. *Cell* **143**:1174–1189. DOI: https://doi.org/10.1016/j.cell.2010.12.001, PMID: 21183079

Huysman MJJ, Vyverman W, De Veylder L. 2014. Molecular regulation of the diatom cell cycle. *J Exp Bot* **65**:2573–2584. DOI: https://doi.org/10.1093/jxb/ert387, PMID: 24277280

Hwang J, Espenshade PJ. 2016. Proximity-dependent biotin labelling in yeast using the engineered ascorbate peroxidase APEX2. *Biochem J* **473**:2463–2469. DOI: https://doi.org/10.1042/BCJ20160106, PMID: 27274088

Hwang J, Ribbens D, Raychaudhuri S, Cairns L, Gu H, Frost A, Urban S, Espenshade PJ. 2016. A Golgi rhomboid protease Rbd2 recruits Cdc48 to cleave yeast SREBP. *EMBO J* **35**:2332–2349. DOI: https://doi.org/10.15252/embj.201693923, PMID: 27655872

Jeffery CJ. 2018. Protein moonlighting: what is it, and why is it important? *Philos Trans R Soc Lond B Biol Sci* **373**:20160523. DOI: https://doi.org/10.1098/rstb.2016.0523, PMID: 29203708

Jensen EC. 2012. Use of fluorescent probes: their effect on cell biology and limitations. *Anat Rec* **295**:2031–2036. DOI: https://doi.org/10.1002/ar.22602, PMID: 23060362

Jeong J, Cohu C, Kerkeb L, Pilon M, Connolly EL, Guerinot ML. 2008. Chloroplast Fe(III) chelate reductase activity is essential for seedling viability under iron limiting conditions. *Proc Natl Acad Sci U S A* **105**:10619–10624. DOI: https://doi.org/10.1073/pnas.0708367105, PMID: 18647837

Jin L, Engelhart AE, Zhang W, Adamala K, Szostak JW. 2018. Catalysis of Template-Directed Nonenzymatic RNA Copying by Iron(II). *J Am Chem Soc* **140**:15016–15021. DOI: https://doi.org/10.1021/jacs.8b09617, PMID: 30335371

Kalocsay M. 2019. APEX Peroxidase-Catalyzed Proximity Labeling and Multiplexed Quantitative Proteomics. (In:) Proximity Labeling, (Editors:) Murat Sunbul and Andreas Jäschke. Human Press, Inc., Totawa, NJ, USA. 41–55. DOI: https://doi.org/10.1007/978-1-4939-9537-0_4, PMID: 31124087

Karas BJ, Diner RE, Lefebvre SC, McQuaid J, Phillips APR, Noddings CM, Brunson JK, Valas RE, Deerinck TJ, Jablanovic J, Gillard JTF, Beeri K, Ellisman MH, Glass JI, Hutchison CA 3rd, Smith HO, Venter JC, Allen AE, Dupont CL, Weyman PD. 2015. Designer diatom episomes delivered by bacterial conjugation. *Nat Commun* **6**:6925. DOI: https://doi.org/10.1038/ncomms7925, PMID: 25897682

Katoh K, Misawa K, Kuma K-I, Miyata T. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30**:3059–3066. DOI: https://doi.org10.1093/nar/gkf436/, PMID: 12136088

Kazamia E, Sutak R, Paz-Yepes J, Dorrell RG, Vieira FRJ, Mach J, Morrissey J, Leon S, Lam F, Pelletier E, Camadro J-M, Bowler C, Lesuisse E. 2018. Endocytosis-mediated siderophore uptake as a strategy for Fe acquisition in diatoms. *Sci Adv* **4**:eaar4536. DOI: https://doi.org/10.1126/sciadv.aar4536, PMID: 29774236

Kazamia E, Mach J, McQuaid J, Olsinova M, Machan R, Coale T, Lesuisse E, Allen, A, Sutak R, Bowler, C. 2019. Monitoring the localization of three iron starvation induced proteins (ISIPs) in a single transformant of *Phaeodactylum tricornutum* under a range of iron supplementation regimes. The molecular life of diatoms Programme and Abstract Book: 95–96. URL: http://meetings.embo.org/files/abstractbooks/Program_%20Abstract%20book_MLD_5_for_print.pdf

Keeling PJ, (79 authors), Worden AZ. 2014. The Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP): Illuminating the Functional Diversity of Eukaryotic Life in the Oceans Through Transcriptome Sequencing. *PLoS Biol* **12**:e1001889. DOI: https://doi.org/10.1371/journal.pbio.1001889, PMID: 24959919

Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Phyre2 web portal for protein modeling, prediction and analysis. *Nat Protoc* **10**:845–858. DOI: https://doi.org/10.1038/nprot.2015.053, PMID: 25950237

Kilian O, Kroth PG. 2005. Identification and characterization of a new conserved motif within the presequence of proteins targeted into complex diatom plastids. *Plant J* **41**:175–183. DOI: https://doi.org/10.1111/j.1365-313X.2004.02294.x, PMID: 15634195

Kim T-W, Park CH, Hsu C-C, Zhu J-Y, Hsiao Y, Branon T, Xu S-L, Ting AY, Wang Z-Y. 2019. Application of TurboID-mediated proximity labeling for mapping a GSK3 kinase signaling network in *Arabidopsis*. *bioRxiv*. DOI: https://doi.org/10.1101/636324

Knoll AH, Bergmann KD, Strauss JV. 2016. Life: the first two billion years. *Philos Trans R Soc Lond B Biol Sci* **371**:20150493. DOI: https://doi.org/10.1098/rstb.2015.0493, PMID: 27672146

Kominek J, Doering DT, Opulente DA, Shen X-X, Zhou X, DeVirgilio J, Hulfachor AB, Groenewald M, Mcgee MA, Karlen SD, Kurtzman CP, Rokas A, Hittinger CT. 2019. Eukaryotic Acquisition of a Bacterial Operon. *Cell* **176**:1356–1366.e10. DOI: https://doi.org/10.1016/j.cell.2019.01.034, PMID: 30799038

Krogh A, Larsson B, von Heijne G, Sonnhammer EL. 2001. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* **305**:567–580. DOI: https://doi.org/10.1006/jmbi.2000.4315, PMID: 11152613

Kustka AB, Allen AE, Morel FMM. 2007. SEQUENCE ANALYSIS AND TRANSCRIPTIONAL REGULATION OF IRON ACQUISITION GENES IN TWO MARINE DIATOMS. *J Phycol* **43**:715–729. DOI: https://doi.org/10.1111/j.1529-8817.2007.00359.x

Lam SS, Martell JD, Kamer KJ, Deerinck TJ, Ellisman MH, Mootha VK, Ting AY. 2015. Directed evolution of APEX2 for electron microscopy and proximity labeling. *Nat Methods* **12**:51–54. DOI: https://doi.org/10.1038/nmeth.3179, PMID: 25419960

Lampe RH, Mann EL, Cohen NR, Till CP, Thamatrakoln K, Brzezinski MA, Bruland KW, Twining BS, Marchetti A. 2018. Different iron storage strategies among bloom-forming diatoms. *Proc Natl Acad Sci U S A* **115**:E12275–E12284. DOI: https://doi.org/10.1073/pnas.1805243115, PMID: 30538208

Leynaert A, Bucciarelli E, Claquin P, Dugdale RC, Martin-Jézéquel V, Pondaven P, Ragueneau O. 2004. Effect of iron deficiency on diatom cell size and silicic acid uptake kinetics. *Limnol Oceanogr* **49**:1134–1143. DOI: https://doi.org/10.4319/lo.2004.49.4.1134

Lindquist E, Aronsson H. 2018. Chloroplast vesicle transport. *Photosynth Res* **138**:361–371. DOI: https://doi.org/10.1007/s11120-018-0566-0, PMID: 30117121

López-Millán AF, Duy D, Philippar K. 2016. Chloroplast Iron Transport Proteins - Function and Impact on Plant Physiology. *Front Plant Sci* **7**:178. DOI: https://doi.org/10.3389/fpls.2016.00178, PMID: 27014281

Lommer M, Specht M, Roy A-S, Kraemer L, Andreson R, Gutowska MA, Wolf J, Bergner SV, Schilhabel MB, Klostermeier UC, Beiko RG, Rosenstiel P, Hippler M, LaRoche J. 2012. Genome and low-iron response of an oceanic diatom adapted to chronic iron limitation. *Genome Biol* **13**:R66. DOI: https://doi.org/10.1186/gb-2012-13-7-r66, PMID: 22835381

Lundberg E, Borner GHH. 2019. Spatial proteomics: a powerful discovery tool for cell biology. *Nat Rev Mol Cell Biol* **20**:285–302. DOI: https://doi.org/10.1038/s41580-018-0094-y, PMID: 30659282

Mair A, Xu S-L, Branon TC, Ting AY, Bergmann DC. 2019. Proximity labeling of protein complexes and cell type-specific organellar proteomes in Arabidopsis enabled by TurboID. *eLife* **8**:e47864. DOI: https://doi.org/10.7554/eLife.47864, PMID: 31535972

Marchand J, Heydarizadeh P, Schoefs B, Spetea C. 2018. Ion and metabolite transport in the chloroplast of algae: lessons from land plants. *Cell Mol Life Sci* **75**:2153–2176. DOI: https://doi.org/10.1007/s00018-018-2793-0, PMID: 29541792

Marchetti A, Schruth DM, Durkin CA, Parker MS, Kodner RB, Berthiaume CT, Morales R, Allen AE, Armbrust EV. 2012. Comparative metatranscriptomics identifies molecular bases for the physiological responses of phytoplankton to varying iron availability. *Proc Natl Acad Sci U S A* **109**:E317–E325. DOI: https://doi.org/10.1073/pnas.1118408109, PMID: 22308424

Martell JD, Deerinck TJ, Lam SS, Ellisman MH, Ting AY. 2017. Electron microscopy using the genetically encoded APEX2 tag in cultured mammalian cells. *Nat Protoc* **12**:1792–1816. DOI: https://doi.org/ 10.1038/nprot.2017.065, PMID: 28796234

Martell JD, Deerinck TJ, Sancak Y, Poulos TL, Mootha VK, Sosinsky GE, Ellisman MH, Ting AY. 2012. Engineered ascorbate peroxidase as a genetically encoded reporter for electron microscopy. *Nat Biotechnol* **30**:1143–1148. DOI: https://doi.org/10.1038/nbt.2375, PMID: 23086203

Martin JH, (42 authors), Tindale NW. 1994. Testing the iron hypothesis in ecosystems of the equatorial Pacific Ocean. *Nature* **371**:123–129. DOI: https://doi.org/10.1038/371123a0

Matsuda Y, Hopkinson BM, Nakajima K, Dupont CL, Tsuji Y. 2017. Mechanisms of carbon dioxide acquisition and CO_2 sensing in marine diatoms: a gateway to carbon metabolism. *Philos Trans R Soc Lond B Biol Sci*. B **372**:20160403 DOI: https://doi.org/10.1098/rstb.2016.0403, PMID: 28717013

May T, Soll J. 1998. Positive Charges Determine the Topology and Functionality of the Transmembrane Domain in the Chloroplastic Outer Envelope Protein Toc34. *J Cell Biol* **141**:895–904. DOI: https://doi.org/10.1083/jcb.141.4.895, PMID: 9585409

Maxson ME, Grinstein S. 2014. The vacuolar-type H^+-ATPase at a glance – more than a proton pump. *J Cell Sci* **127**:4987–4993. DOI: https://doi.org/10.1242/jcs.158550, PMID: 25453113

Mayle KM, Le AM, Kamei DT. 2012. The Intracellular Trafficking Pathway of Transferrin. *Biochim Biophys Acta* **1820**:264–281. DOI: https://doi.org/10.1016/j.bbagen.2011.09.009, PMID: 21968002

McAlister GC, Nusinow DP, Jedrychowski MP, Wühr M, Huttlin EL, Erickson BK, Rad R, Haas W, Gygi SP. 2014. MultiNotch MS3 Enables Accurate, Sensitive, and Multiplexed Detection of Differential Expression Across Cancer Cell Line Proteomes. *Anal Chem* **86**:7150–7158. DOI: https://doi.org/10.1021/ac502040v, PMID: 24927332

McQuaid JB, Kustka AB, Oborník M, Horák A, McCrow JP, Karas BJ, Zheng H, Kindeberg T, Andersson AJ, Barbeau KA, Allen AE. 2018. Carbonate-sensitive phytotransferrin controls high-affinity iron uptake in diatoms. *Nature* **555**:534–537. DOI: https://doi.org/10.1038/nature25982, PMID: 29539640

Merchant S. 2019. Nutritional Copper Signaling and Homeostasis. Grantome research proposal Abstract. URL: http://grantome.com/grant/NIH/R35-GM127114-01

Merkulova M, Hurtado-Lorenzo A, Hosokawa H, Zhuang Z, Brown D, Ausiello DA, Marshansky V. 2011. Aldolase directly interacts with ARNO and modulates cell morphology and acidic vesicle distribution. *Am J Physiol Cell Physiol* **300**:C1442–C1455. DOI: https://doi.org/10.1152/ajpcell.00076.2010, PMID: 21307348

Mészáros B, Erdos G, Dosztányi Z. 2018. IUPred2A: context-dependent prediction of protein disorder as a function of redox state and protein binding. *Nucleic Acids Res* **46**:W329–W337. DOI: https://doi.org/10.1093/nar/gky384, PMID: 29860432

Morel FMM, Kustka AB, Shaked Y. 2008. The role of unchelated Fe in the iron nutrition of phytoplankton. *Limnol Oceanogr* **53**:400–404. DOI: doi.org/10.4319/lo.2008.53.1.0400

Moreno CM, Lin Y, Davies S, Monbureau E, Cassar N, Marchetti A. 2018. Examination of gene repertoires and physiological responses to iron and light limitation in Southern Ocean diatoms. *Polar Biol* **41**:679–696. DOI: https://doi.org/10.1007/s00300-017-2228-7

Morrissey J, Sutak R, Paz-Yepes J, Tanaka A, Moustafa A, Veluchamy A, Thomas Y, Botebol H, Bouget F-Y, McQuaid JB, Tirichine L, Allen AE, Lesuisse E, Bowler C. 2015. A Novel Protein, Ubiquitous in Marine Phytoplankton, Concentrates Iron at the Cell Surface and Facilitates Uptake. *Curr Biol* **25**:364–371. DOI: https://doi.org/10.1016/j.cub.2014.12.004, PMID: 25557662

Musayev FN, Di Salvo ML, Ko T-P, Schirch V, Safo MK. 2003. Structure and properties of recombinant human pyridoxine 5'-phosphate oxidase. *Protein Sci* **12**:1455–1463. DOI: https://doi.org/10.1110/ps.0356203, PMID: 12824491

Naslavsky N, Caplan S. 2018. The enigmatic endosome – sorting the ins and outs of endocytic trafficking. *J Cell Sci* **131**:jcs216499. DOI: https://doi.org/10.1242/jcs.216499, PMID: 29980602

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* **32**:268–274. DOI: https://doi.org/10.1093/molbev/msu300, PMID: 25371430

Nützmann H-W, Scazzocchio C, Osbourn A. 2018. Metabolic Gene Clusters in Eukaryotes. *Annu Rev Genet* **52**:159–183. DOI: https://doi.org/10.1146/annurev-genet-120417-031237, PMID: 30183405

Oborník M, Füssy Z. 2018. Complex Endosymbioses I: From Primary to Complex Plastids, Multiple Independent Events. *(*In:) Plastids. Methods in Molecular Biology, (Editor:) Eric Maréchal. Humana Press, Inc., Totowa, NJ, USA. **1829**:17–35. DOI: https://doi.org/10.1007/978-1-4939-8654-5_2

Och LM, Shields-Zhou GA. 2012. The Neoproterozoic oxygenation event: Environmental perturbations and biogeochemical cycling. *Earth-Sci Rev* **110**:26–57. DOI: https://doi.org/10.1016/j.earscirev.2011.09.004

Ohgami RS, Campagna DR, Greer EL, Antiochos B, McDonald A, Chen J, Sharp JJ, Fujiwara Y, Barker JE, Fleming MD. 2005. Identification of a ferrireductase required for efficient transferrin-dependent iron uptake in erythroid cells. *Nat Genet* **37**:1264–1269. DOI: https://doi.org/10.1038/ng1658, PMID: 16227996

O'Halloran TV, Culotta VC. 2000. Metallochaperones, an Intracellular Shuttle Service for Metal Ions. *J Biol Chem* **275**:25057–25060. DOI: https://doi.org/10.1074/jbc.R000006200, PMID: 10816601

Osbourn AE, Field B. 2009. Operons. *Cell Mol Life Sci* **66**:3755–3775. DOI: https://doi.org/10.1007/s00018-009-0114-3, PMID: 19662496

Otsuka Y, Satoh T, Nakayama N, Inaba R, Yamashita H, Satoh AK. 2019. Parcas is the predominant Rab11GEF for rhodopsin transport in *Drosophila* photoreceptors. *J Cell Sci*. **132**:1–13. DOI: https://doi.org/10.1242/jcs.231431, PMID: 31296556

Paek J, Kalocsay M, Staus DP, Wingler L, Pascolutti R, Paulo JA, Gygi SP, Kruse AC. 2017. Multidimensional Tracking of GPCR Signaling via Peroxidase-Catalyzed Proximity Labeling. *Cell* **169**:338–349. DOI: https://doi.org/10.1016/j.cell.2017.03.028, PMID: 28388415

Paulo JA, O'Connell JD, Everley RA, O'Brien J, Gygi MA, Gygi SP. 2016. Quantitative mass spectrometry-based multiplexing compares the abundance of 5000 *S. cerevisiae* proteins across 10 carbon sources. *J Proteomics* **148**:85–93. DOI: https://doi.org/10.1016/j.jprot.2016.07.005, PMID: 27432472

Peschke M, Moog D, Klingl A, Maier UG, Hempel F. 2013. Evidence for glycoprotein transport into complex plastids. *Proc Natl Acad Sci U S A* **110**:10860–10865. DOI: https://doi.org/10.1073/pnas.1301945110, PMID: 23754425

Peng K, Radivojac P, Vucetic S, Dunker AK, Obradovic Z. 2006. Length-dependent prediction of protein intrinsic disorder. *BMC Bioinformatics* **7**:208. DOI: https://doi.org/10.1186/1471-2105-7-208, PMID: 16618368

Perez-Riverol Y, (21 authors), Vizcaíno JA. 2019. The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic Acids Res* **47**:D442–D450. DOI: https://doi.org/10.1093/nar/gky1106, PMID: 30395289

Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods* **8**:785–786. DOI: https://doi.org/10.1038/nmeth.1701, PMID: 21959131

Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. 2004. UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* **25**:1605–1612. DOI: https://doi.org/10.1002/jcc.20084, PMID: 15264254

Philpott CC, Jadhav S. 2019. The ins and outs of iron: Escorting iron through the mammalian cytosol. *Free Radic Biol Med* **133**:112–117. DOI: https://doi.org/10.1016/j.freeradbiomed.2018.10.411, PMID: 30321701

Poole LB. 2015. The basics of thiols and cysteines in redox biology and chemistry. *Free Radic Biol Med* **80**:148–157. DOI: https://doi.org/10.1016/j.freeradbiomed.2014.11.013, PMID: 25433365

Prihoda J, Tanaka A, de Paula WBM, Allen JF, Tirichine L, Bowler C. 2012. Chloroplast-mitochondria cross-talk in diatoms. *J Exp Bot* **63**:1543–1557. DOI: https://doi.org/10.1093/jxb/err441, PMID: 22268145

Progida C, Bakke O. 2016. Bidirectional traffic between the Golgi and the endosomes - machineries and regulation. *J Cell Sci* **129**:3971–3982. DOI: https://doi.org/10.1242/jcs.185702, PMID: 27802132

Przybyla-Toscano J, Roland M, Gaymard F, Couturier J, Rouhier N. 2018. Roles and maturation of iron-sulfur proteins in plastids. *J Biol Inorg Chem* **23**:545–566. DOI: https://doi.org/10.1007/s00775-018-1532-1, PMID: 29349662

R Core Team. 2013. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: http://www.R-project.org/

Radhamony RN, Theg SM. 2006. Evidence for an ER to Golgi to chloroplast protein transport pathway. *Trends Cell Biol* **16**:385–387. DOI: https://doi.org/10.1016/j.tcb.2006.06.003, PMID: 16815014

Rao RSP, Wollenweber B. 2010. Do N-glycoproteins have preference for specific sequons? *Bioinformation* **5**:208–212. DOI: https://doi.org/10.6026/97320630005208, PMID: 21364799

Raven EL. 2003. Understanding functional diversity and substrate specificity in haem peroxidases: what can we learn from ascorbate peroxidase? *Nat Prod Rep* **20**:367–381. DOI: https://doi.org/10.1039/b210426c, PMID: 12964833

Ree R, Varland S, Arnesen T. 2018. Spotlight on protein N-terminal acetylation. *Exp Mol Med* **50**:90. DOI: https://doi.org/10.1038/s12276-018-0116-z, PMID: 30054468

Rhee H-W, Zou P, Udeshi ND, Martell JD, Mootha VK, Carr SA, Ting AY. 2013. Proteomic Mapping of Mitochondria in Living Cells via Spatially Restricted Enzymatic Tagging. *Science* **339**:1328–1331. DOI: https://doi.org/10.1126/science.1230593, PMID: 23371551

Rochaix J-D. 2011. Regulation of photosynthetic electron transport. *Biochim Biophys Acta* **1807**:375–383. DOI: https://doi.org/10.1016/j.bbabio.2010.11.010, PMID: 21118674

Rosenzweig AC. 2002. Metallochaperones: Bind and Deliver Minireview. *Chem Biol* **9**:673–677. DOI: https://doi.org/10.1016/S1074-5521(02)00156-4, PMID: 12079778

Sacher M, Di Bacco A, Lunin VV, Ye Z, Wagner J, Gill G, Cygler M. 2005. The crystal structure of CREG, a secreted glycoprotein involved in cellular growth and differentiation. *Proc Natl Acad Sci U S A* **102**:18326–18331. DOI: https://doi.org/10.1073/pnas.0505071102, PMID: 16344469

Schähs P, Weidinger P, Probst OC, Svoboda B, Stadlmann J, Beug H, Waerner T, Mach L. 2008. Cellular repressor of E1A-stimulated genes is a *bona fide* lysosomal protein which undergoes proteolytic maturation during its biosynthesis. *Exp Cell Res* **314**:3036–3047. DOI: https://doi.org/10.1016/j.yexcr.2008.06.015, PMID: 18621046

Sapriel G, Quinet M, Heijde M, Jourdren L, Tanty V, Luo G, Le Crom S, Lopez PJ. 2009. Genome-Wide Transcriptome Analyses of Silicon Metabolism in *Phaeodactylum tricornutum* Reveal the Multilevel Regulation of Silicic Acid Transporters. *PLoS ONE* **4**:e7458. DOI: https://doi.org/10.1371/journal.pone.0007458, PMID: 19829693

Savelli B, Li Q, Webber M, Jemmat AM, Robitaille A, Zamocky M, Mathé C, Dunand C. 2019. RedoxiBase: A database for ROS homeostasis regulated proteins. *Redox Biol* **26**:101247. DOI: https://doi.org/10.1016/j.redox.2019.101247, PMID: 31228650

Savitsky P, Bray J, Cooper CDO, Marsden BD, Mahajan P, Burgess-Brown NA, Gileadi O. 2010. High-throughput production of human proteins for crystallization: the SGC experience. *J Struct Biol* **172**:3–13. DOI: https://doi.org/10.1016/j.jsb.2010.06.008, PMID: 20541610

Scheiber IF, Pilátová J, Malych R, Kotabova E, Krijt M, Vyoral D, Mach J, Léger T, Camadro J-M, Prášil O, Lesuisse E, Sutak R. 2019. Copper and iron metabolism in *Ostreococcus tauri* - the role of phytotransferrin, plastocyanin and a chloroplast copper-transporting ATPase. *Metallomics* **11**:1657–1666. DOI: https://doi.org/10.1039/c9mt00078j, PMID: 31380866

Shoemaker BA, Portman JJ, Wolynes PG. 2000. Speeding molecular recognition by using the folding funnel: the fly-casting mechanism. *Proc Natl Acad Sci U S A* **97**:8868–8873. DOI: https://doi.org/10.1073/pnas.160259697, PMID: 10908673

Sengupta R, Poderycki MJ, Mattoo S. 2019. CryoAPEX – an electron tomography tool for subcellular localization of membrane proteins. *J Cell Sci* **132**:jcs222315. DOI: https://doi.org/10.1242/jcs.222315, PMID: 30886003

Sheppard VC, Scheffel A, Poulsen N, Kröger N. 2012. Live Diatom Silica Immobilization of Multimeric and Redox-Active Enzymes. *Appl Environ Microbiol* **78**:211–218. DOI: https://doi.org/10.1128/AEM.06698-11, PMID: 22057862

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, Mcwilliam H, Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* **7**:539. DOI: https://doi.org/10.1038/msb.2011.75, PMID: 21988835

Smith SR, Gillard JTF, Kustka AB, McCrow JP, Badger JH, Zheng H, New AM, Dupont CL, Obata T, Fernie AR, Allen AE. 2016. Transcriptional Orchestration of the Global Cellular Response of a Model Pennate Diatom to Diel Light Cycling under Iron Limitation. *PLoS Genet* **12**:e1006490. DOI: https://doi.org/10.1371/journal.pgen.1006490, PMID: 27973599

Soll J, Schleiff E. 2004. Protein import into chloroplasts. *Nat Rev Mol Cell Biol* **5**:198–208. DOI: https://doi.org/10.1038/nrm1333, PMID: 14991000

Steentoft C, Vakhrushev SY, Joshi HJ, Kong Y, Vester-Christensen MB, Schjoldager KT-BG, Lavrsen K, Dabelsteen S, Pedersen NB, Marcos-Silva L, Gupta R, Bennett EP, Mandel U, Brunak S, Wandall HH, Levery SB, Clausen H. 2013. Precision mapping of the human O-GalNAc glycoproteome through SimpleCell technology. *EMBO J* **32**:1478–1488. DOI: https://doi.org/10.1038/emboj.2013.79, PMID: 23584533

Strand TA, Lale R, Degnes KF, Lando M, Valla S. 2014. A New and Improved Host-Independent Plasmid System for RK2-based Conjugal Transfer. *PLoS One* **9**:e90372. DOI: https://doi.org/10.1371/journal.pone.0090372, PMID: 24595202

Tagliabue A, Bowie AR, Boyd PW, Buck KN, Johnson KS, Saito MA. 2017. The integral role of iron in ocean biogeochemistry. *Nature* **543**:51–59. DOI: https://doi.org/10.1038/nature21058, PMID: 28252066

Tesson B, Genet MJ, Fernandez V, Degand S, Rouxhet PG, Martin-Jézéquel V. 2009. Surface Chemical Composition of Diatoms. *Chembiochem* **10**:2011–2024. DOI: https://doi.org/10.1002/cbic.200800811, PMID: 19623594

Theorell H. 1935. Preparation in pure state of the effect group of yellow enzymes. *Biochemische Zeitschrift* **275**:344–346.

Li Y, Tian C, Liu K, Zhou Y, Yang J, Zou P. 2020. A Clickable APEX Probe for Proximity-Dependent Proteomic Profiling in Yeast. *Cell Chem Biol* **27**:858-865.e8. DOI: https://doi.org/10.1016/j.chembiol.2020.05.006; PMID: 32470320

Trinkle-Mulcahy L. 2019. Recent advances in proximity-based labeling methods for interactome mapping. *F1000Res* **8**:135. DOI: https://doi.org/10.12688/f1000research.16903.1, PMID: 30774936

Tuteja R. 2005. Type I signal peptidase: An overview. *Arch Biochem Biophys* **441**:107–111. DOI: https://doi.org/10.1016/j.abb.2005.07.013, PMID: 16126156

Uversky VN. 2015a. The intrinsic disorder alphabet. III. Dual personality of serine. *Intrinsically*
Disord Proteins **3**:e1027032. DOI: https://doi.org/10.1080/21690707.2015.1027032, PMID:
28232888

Uversky VN. 2015b. The multifaceted roles of intrinsic disorder in protein complexes. *FEBS Lett*
589:2498–2506. DOI: https://doi.org/10.1016/j.febslet.2015.06.004, PMID: 26073257

Uversky VN. 2016. Dancing Protein Clouds: The Strange Biology and Chaotic Physics of
Intrinsically Disordered Proteins. *J Biol Chem* **291**:6681–6688. DOI:
https://doi.org/10.1074/jbc.R115.685859, PMID: 26851286

Valko M, Morris H, Cronin MTD. 2005. Metals, Toxicity and Oxidative Stress. *Curr Med Chem*
12:1161–1208. DOI: https://doi.org/10.2174/0929867053764635, PMID: 15892631

Villarejo A, Burén S, Larsson S, Déjardin A, Monné M, Rudhe C, Karlsson J, Jansson S,
Lerouge P, Rolland N, von Heijne G, Grebe M, Bako L, Samuelsson G. 2005. Evidence for a
protein transported through the secretory pathway en route to the higher plant chloroplast. *Nat*
Cell Biol **7**:1224–1231. DOI: https://doi.org/10.1038/ncb1330, PMID: 16284624

Wang L, Jonikas MC. 2020. The pyrenoid. *Curr Biol* **30**:R456–R458. DOI:
https://doi.org/10.1016/j.cub.2020.02.051, PMID: 32428480

Wang J, Pantopoulos K. 2011. Regulation of cellular iron metabolism. *Biochem J* **434**:365–381.
DOI: https://doi.org/10.1042/BJ20101825, PMID: 21348856

Yee DP, Hildebrand M, Tresguerres M. 2019. Dynamic subcellular translocation of V-type H^+-
ATPase is essential for biomineralization of the diatom silica cell wall. *New Phytol* **225**:2411–
2422. DOI: https://doi.org/10.1111/nph.16329, PMID: 31746463.

Zak O, Ikuta K, Aisen P. 2002. The Synergistic Anion-Binding Sites of Human Transferrin:
Chemical and Physiological Effects of Site-Directed Mutagenesis. *Biochemistry* **41**:7416–7423.
DOI: https://doi.org/10.1021/bi0160258, PMID: 12044175

Zhang Y, Song G, Lal NK, Nagalakshmi U, Li Y, Zheng W, Huang P, Branon TC, Ting AY,
Walley JW, Dinesh-Kumar SP. 2019. TurboID-based proximity labeling reveals that UBR7 is a
regulator of N NLR immune receptor-mediated immunity. *Nat Commun* **10**:3252. DOI:
https://doi.org/10.1038/s41467-019-11202-z, PMID: 31324801

Zulu NN, Zienkiewicz K, Vollheyde K, Feussner I. 2018. Current trends to comprehend lipid
metabolism in diatoms. *Prog Lipid Res* **70**:1–16. DOI:
https://doi.org/10.1016/j.plipres.2018.03.001, PMID: 29524459

Figure 1. Functional APEX2-enabled proteogenomics with pTF (phytotransferrin/ISIP2a/Phatr3_J54465) in the model marine diatom *Phaeodactylum tricornutum*.

(A) pTF-APEX2 encoding episome is introduced into *P. tricornutum* cells using bacterial conjugation.

(B) Resulting transconjugants are genotyped and evaluated for fusion protein expression. APEX2 activity and fusion protein localization are then confirmed with an enzymatic assay and electron microscopy, respectively.

(C) In a proximity-dependent proteomic mapping (proximity proteomics) experiment, pTF-APEX2 expressing cell line is supplemented with biotin-phenol and hydrogen peroxide, reaction quenched, and cells lysed. Cell lysate is then subjected to streptavidin pull-down, proteins analyzed with mass spectrometry (MS), peptides mapped to a *P. tricornutum* proteome database, and corresponding genes identified.

(D) Interesting MS hits are further evaluated experimentally (e.g., for colocalization with the bait protein (i.e., pTF) and/or for predicted biochemical activity) as well as bioinformatically. Created with BioRender.com.

Figure 2. pTF is localized to intracellular vesicles and APEX2 is enzymatically active in live *Phaeodactylum tricornutum* cells.

(A) pTF-mCherry colocalizes with the membrane dye MDY-64 on the cell surface and within intracellular vesicles. The fluorescent fusion protein was expressed under the *pTF* promoter and terminator in a *ΔpTF P. tricornutum* genetic background. Cells were stained with 100 µM MDY-64 for 10 min at room temperature. Scale bar is 10 µm. Fusion protein schematic created with BioRender.com.

(B) Top: *pTF-APEX2* flanked by nitrate-inducible *NR* (nitrate reductase) promoter, terminator, and untranslated regions (UTRs). Linker sequence ("*link*") between *pTF* and *APEX2* encoded KGSGSTSGSG. Bottom: Schematic of the pTF-APEX2 expressing episome. Tetracycline (*Tet^R*) and ampicillin (*Amp^R*) resistance genes for *E. coli* selection, bacterial replication origin, yeast centromere (*CEN6-ARSH4-HIS3*) for episome maintenance, phleomycin/zeocin (*ShBle*) resistance gene for *P. tricornutum* selection, origin of conjugative transfer (*oriT*). Plasmid map created with BioRender.com.

(C) Anti-pTF western blot confirming pTF-APEX2 fusion protein (84.73 kDa) expression in four transconjugant *P. tricornutum* cell lines (lanes s1–s4; left: soluble cell lysate fraction, right: insoluble cell lysate fraction). Bacterial conjugation was performed with WT *P. tricornutum* cells which is why native pTF (57.05 kDa) bands were present in our samples. Teal circle: APEX2, white circle: linker, gray oval: pTF.

(D) Left: pTF-APEX2 expressing, but not WT, cells convert APEX2 substrate Amplex UltraRed (50 µM) into a colored product resorufin in the presence of 2 mM H_2O_2. Right: >40-fold higher resorufin signal was observed in supernatants from pTF-APEX2 expressing cells than those from WT cells. Triplicate cultures from one WT and one pTF-APEX2 expressing cell line (cell line s2) were used in this experiment. Standard deviation is shown. *P. tricornutum* cartoons created with BioRender.com.

Figure 2—source data 1. Raw Amplex UltraRed assay data underpinning the chart in *Figure 2D*.

Figure 2—source data 2. R script to reproduce the chart in *Figure 2D*.

Figure 2—source data 3. CSV file to be used in conjuction with **Figure 2—source data 2**.

Figure 3. pTF-APEX2 is localized to the cell membrane and intracellular vesicles.

(A) Electron microscopy protocol summary. Briefly: WT and pTF-APEX2 expressing cells (grown in triplicate) were fixed, treated with diaminobenzidine (DAB) and hydrogen peroxide, post-fixed with osmium tetroxide, embedded in agar, negatively stained with uranyl acetate, dehydrated, embedded in resin, and visualized. Created with BioRender.com.

(B) Expected cell surface and intracellular pTF-APEX2 localization. Teal and brown arrows point to APEX2-induced signal and cell wall, respectively. Zoomed in (top left image) is cell periphery where it can clearly be seen that cell membrane, not cell wall, is occupied by the fusion protein. pTF-APEX2 containing vesicles were observed (bottom two left images; acquired with backscatter scanning electron microscopy). Zoomed in (middle left image) are cell membrane and vesicles with pTF-APEX2. Mitochondrial signal in both WT and pTF-APEX2 expressing cells is likely due to endogenous (mitochondrial) APEX2-like peroxidases. Scale bar is 1 µm. *P. tricornutum* cartoons created with BioRender.com.

Figure 4. Proximity-dependent proteomic mapping with APEX2 reveals candidate proteins involved in phytotransferrin (pTF) endocytosis in *Phaeodactylum tricornutum.*

(A) Summary of the proximity-dependent proteomic mapping experiment. Briefly: WT and pTF-APEX2 expressing cells (grown in quintuplicate) were chilled on ice, pelleted, treated with 1.2 M sorbitol, supplemented with 2.5 mM biotin-phenol and 1 mM H_2O_2. The labeling reaction was quenched, cells were lysed, and evaluated for biotin enrichment. Cleared cell lysates were then subjected to streptavidin pull-down followed by quantitative mass spectrometry using tandem mass tags (TMT). Created with BioRender.com.

(B) Left: Enrichment of biotinylated proteins over WT background was observed with streptavidin blot; ~66.5 kDa biotinylated BSA control; equal loading. The most prominent band in all samples is likely the endogenous biotin-containing propionyl-CoA carboxylase (Phatr3_J51245, ~72.5 kDa). Right: Volcano plot of quantitative MS data highlighting proteins with APEX2/WT ratio of at least 1.5 and P value ≤0.05 (shaded area with teal data points). Proteins encoded by a known iron-sensitive gene cluster on chromosome 20 are highlighted (arrowed Ensembl IDs). *P. tricornutum* cartoons created with BioRender.com.

(C) 14/38 (10 shown here) MS hits are proteins with iron-sensitive transcripts. Proteins chosen for further characterization are bolded. Chloroplast localization and the associated prediction confidence were determined with SignalP 4.1 and ASAFind version 1.1.7. *Experimentally shown to localize to the pyrenoid—a RuBisCO-containing proteinaceus organelle—in the interior of the *P. tricornutum* chloroplast (*Allen et al., 2012*). **Experimentally shown to be localized adjacently to the chloroplast (*Kazamia et al., 2019*; *Kazamia et al., 2018*).

Figure 4—source data 1. Pre-processed quantitative mass spectrometry proteomics data underpinning the Volcano plot and the table in *Figure 4B* and *Figure 4C*, respectively. The table in *Figure 4C* was additionally informed by **Figure 5—figure supplement 1—source data 1**.

Figure 4—source data 2. R script to reproduce the chart in *Figure 4B*.

Figure 4—source data 3. CSV file to be used in conjuction with **Figure 4—source data 2**.

Figure 5. Proteins encoded by a known iron- and silicon-sensitive locus in *Phaeodactylum tricornutum* colocalize with pTF in the chloroplast vicinity.

(A) Genes corresponding to three statistically significant proteomic hits are clustered on chromosome 20. All three proteins were co-expressed with mCherry-tagged pTF as fusions with a yellow fluorescent protein. Corresponding Ensembl and UniProt IDs are noted. Numbers indicate base pairs. Created with BioRender.com.

(B) Two proteins—pTF.CREG1 and pTF.CatCh1—show clear, yet distinct, colocalization with pTF and the chloroplast periphery. pTF.CREG1-EYFP was consistently punctate whereas pTF.CatCh1-EYFP lined the chloroplast margin. pTF-mCherry punctum was almost exclusively positioned next to this "chloroplast lining" pattern. *P. tricornutum* is pleiomorphic which explains why different cell morphologies were observed in pTF.CREG1-EYFP (fusiform morphotype) and pTF.CatCh1-EYFP (triradiate morphotype) cell lines. Scale bar is 10 μm. Fusion protein schematics created with BioRender.com.

Figure 6. pTF.CREG1 homologs are present across the tree of life.
pTF.CREG1 homolog search was performed with the National Center for Biotechnology Information (NCBI)
and the Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP) databases (*Caron et*
al., 2016; *Keeling et al., 2014*). Homologs from algae with complex plastids (including diatoms) form a
crown group away from animal proteins. pTF.CREG1 clusters with its paralog Phatr3_J10972 (black
rectangle). Red dots indicate predicted gene duplication events. Scale bar: 0.5 substitutions per position.
2168
Figure 6—source data 1. Amino acid sequence alignment underpinning the phylogenetic tree in *Figure*
6.

Figure 7. pTF.CatCh1 is a cysteine-rich heterokont-specific protein.

(A) pTF.CatCh1 homolog search was performed with NCBI and marine microbial eukaryote (MMETSP)
databases. Numerous, almost exclusively diatom, proteins were identified. Interestingly, pTF.CatCh1 is
paralogous to ISIP2b (black arrows). Scale bar: 0.1 substitutions per position.

(B) pTF.CatCh1 and ISIP2b share three CXC and one CXXC motifs (boxed) commonly found in redox-
active proteins and proteins involved in cellular metal homeostasis. Black boxes indicate the two CX(X)C
motifs conserved across the majority of pTF.CatCh1 homologs. Blue triangles: additional conserved
cysteine amino acid residues between pTF.CatCh1 and ISIP2b for a total of 20. Asterisks: cysteine amino
acid residues that are 100% conserved across all homologs.

2180

2181 **Figure 7—source data 1.** Amino acid sequence alignment underpinning the phylogenetic tree in *Figure*
2182 *7A*.

Figure 8. Emerging view of intracellular inorganic iron allocation pathway in *Phaeodactylum tricornutum.* We propose a model for clathrin-mediated internalization of phytotransferrin (pTF)-bound ferric, Fe^{3+}, iron, and its trafficking to the chloroplast periphery. In this model, pTF releases Fe^{3+} after undergoing a structural change upon endosome acidification driven by a vacuolar-type H^+-ATPase. pTF.CREG1 may regulate this process at the endosome periphery and perhaps also in earlier endocytosis stages closer to the cell surface. It is unclear how intravesicular reduction of ferric iron is achieved and how the resulting ferrous iron gets exported. Given that no DMT1/NRAMP2 homologs are present in *P. tricornutum*, Zrt- and Irt-like proteins (ZIPs)—known to export Fe^{2+} from intracellular compartments—may mediate the latter (*Blaby-Haas and Merchant, 2012*; *Kustka et al., 2007*; *Lampe et al., 2018*). Ferrous iron could be offloaded to pTF.CatCh1 on the outer chloroplast membrane and ultimately transported further into the chloroplast interior for cellular assimilation and use (e.g., iron–sulfur cluster biosynthesis for incorporation into photosystems I and II). pTF.CatCh1 may alternatively (or additionally) be responsible for pTF recruitment to the chloroplast margin. The enrichment of FBP1 and ISIP1—both previously shown to localize to intracellular vesicles in *P. tricornutum* (*Coale et al., 2019*; *Kazamia et al., 2019*)—in our proximity proteomics experiment suggests the whole pathway intersects with that for internalizing organic, siderophore-bound, iron. Gray panel: pytotransferrin (pTF). Blue panels: proteins colocalized with pTF. Green panels: additional proteins enriched in our APEX2 experiment. One phytotransferrin-Fe^{3+} complex is shown for clarity. Orange vesicle interior indicates pH drop due to ATPase activity. Detailed chloroplast membrane layers are omitted (diatom chloroplasts contain four membranes). Created with BioRender.com.

Figure 2—figure supplement 1. *Phaeodactylum tricornutum* phytotransferrin (pTF) features, resorufin imaging in pTF-APEX2 expressing cells, and resorufin signal co-dependence on Amplex UltraRed and hydrogen peroxide.

(A) pTF is a 537 amino acid residues (AAs) long cell membrane-associated iron receptor protein consisting of a signal peptide (AAs 1–18; predicted with SignalP 4.1), an iron-binding domain (AAs 19–493), a transmembrane domain (AAs 494–516; TMHMM Server v. 2.0 prediction output is shown to the right), and a tail domain (AAs 517–537). The latter is predicted to contain an endocytosis motif (*Lommer et al., 2012*). Bolded and light blue are conserved putative iron- (Asp48, Tyr71, Tyr214, His294) and carbonate-coordinating (Arg183) amino acid residues, respectively.

(B) Direct visualization of resorufin (yellow) in pTF-APEX2 expressing cells. Both WT and pTF-APEX2 expressing cells were incubated in reaction buffer (50 µM Amplex UltraRed, 2 mM H_2O_2, PBS (pH 7.4)) for 6 min at room temperature prior to imaging. Red is chloroplast autofluorescence.

(C) Supernatants after incubating cells from four pTF-APEX2 expressing cell lines (s1–s4) in reaction buffer (50 µM Amplex UltraRed, 2 mM H_2O_2, PBS (pH 7.4)) for 6 min at room temperature. Resorufin signal was observed only when both APEX2 substrates (Amplex UltraRed and hydrogen peroxide) were present.

Figure 3—figure supplement 1. Analysis of transcriptionally active *Phaeodactylum tricornutum* peroxidases reveals putative APEX2-like and mitochondrial enzymes.

(A) Detail from the electron microscopy sample preparation protocol: agar blocks with embedded diaminobenzidine (DAB)-treated pTF-APEX2 expressing diatom cells. *P. tricornutum* cartoon and "zoom in" objects created with BioRender.com.

(B) Mitochondrial signal as shown here was observed in electron microscopy experiments with both WT and pTF-APEX2 expressing cells subjected to DAB and hydrogen peroxide. "Zoom in" objects created with BioRender.com.

(C) Amino acid sequence alignment of putative APEX2-like *P. tricornutum* peroxidases. Amino acid residues crucial for ascorbate peroxidase (APX) and likely crucial for APEX2 activity (Arg38, His42, His163, and Asp208; numbering is relative to APEX2; depicted with black triangles and shaded) are conserved in eight *P. tricornutum* peroxidases. Numbers above amino acid sequences indicate alignment position.

(D) Amino acid sequence alignment of putative substrate binding loops in the identified APEX2-like peroxidases. Black and blue arrows point to enzymes with proline at position 134 (shaded; numbering is relative to APEX2)—an amino acid residue change rendering APEX2 much more active than its APEX predecessor (*Lam et al., 2015*)—and predicted mitochondrial localization, respectively.

(E) Some additional properties of the identified APEX2-like peroxidases. They all belong to class I (intracellular) haem peroxidases; one may be membrane-associated. [1]RedoxiBase database. [2]APEX2 and horseradish peroxidase (HRP) have proline at this position. [3]TargetP 1.1. [4]TMHMM Server v. 2.0.

2236 **Figure 4—figure supplement 1.** WT and pTF-APEX2 proteomic replicates form distinct clusters.
2237 Hierarchical clustering (Ward's method as implemented in the JMP software package) of scaled quantitative
2238 mass spectrometry proteomics data reveals WT- and pTF-APEX2-specific clusters. The heat map as a
2239 whole suggests minimal technical variability in our proteomic experiment even though samples WT5 and
2240 pTF-APEX2.4 deviate slightly from other corresponding samples.
2241
Figure 4—figure supplement 1—source data 1. Scaled quantitative mass spectrometry proteomics data.

Figure 5—figure supplement 1. *pTF.CREG1*, *pTF.CatCh1*, and *pTF.ap1* closely resemble *pTF*
transcriptional profile.
Transcriptional profiles of *pTF* and the three genes from the identified iron- and silicon-sensitive gene
cluster on chromosome 20 across different growth conditions (L (low): 20 pM Fe', M (medium): 40 pM Fe',
H (high): 400 pM Fe') and sampling timepoints (10AM, 2PM, 6PM, 10PM, 2AM, 6AM) are shown. Fe': sum
of all iron species not complexed to EDTA. Red dashed lines correspond to genes' transcriptional peaks
(late night). Average RPKM: Reads Per Kilobase of transcript, per Million mapped reads averaged across
across two (L and M) or three (H) replicates.
2251
2252 **Figure 5—figure supplement 1—source data 1.** Transcriptomic data and functional annotations for
2253 proteins with APEX2/WT ratios ≥1.5 and P values ≤0.05 underpinning *Figure 5—figure supplement 1* and
2254 *Figure 4C*.
2255
2256 **Figure 5—figure supplement 1—source data 2.** R script to reproduce the chart in *Figure 5—figure*
2257 *supplement 1*.
2258
2259 **Figure 5—figure supplement 2—source data 3.** CSV file to be used in conjuction with **Figure 5—figure**
2260 **supplement 1—source data 2**.

Figure 5—figure supplement 2. Co-expression episome assembly overview, assessment of co-expression cell lines, and confocal microscopy bleed-through controls.

(A) Summary of the pTF-mCherry MS hit-EYFP co-expression episome assembly strategy. Created with BioRender.com.

(B) Colony PCR and western blot analyses indicate the presence of full-length fusion proteins in our co-expression cell lines. Left: Primers binding to *Phatr3_J23658* (flavodoxin-encoding gene) promoter and terminator flanking MS hit-EYFP coding regions were used in PCR reactions which resulted in the expected ~3 kbp amplicons. Positive controls: pure episomes used for conjugation. Negative controls: WT *P. tricornutum* cells and water which yielded no amplicons (not shown). Right: Anti-pTF western blot revealed the expected pTF-mCherry (83.75 kDa) bands alongside the bands corresponding to native pTF (57.05 kDa) and what appears to be a pTF-mCherry truncation (the additional band at ~70 kDa). Protein schematics created with BioRender.com.

(C) The observed lack of significant cross-channel bleed-through supports colocalization imaging data and conclusions. Imaging conditions with minimal bleed-through were determined with WT *P. tricornutum* cells (top row), *P. tricornutum* cells expressing mCherry alone (middle row), and *P. tricornutum* cells expressing a Venus-tagged protein (bottom row). Scale bar is 10 μm.

Figure 5—figure supplement 3. Additional representative examples of pTF.CREG1 and pTF.CatCh1
colocalization with pTF.
(A) Three additional cells from pTF-mCherry and pTF.CREG1-EYFP expressing cell line described in the
main text exhibit similar punctate colocalization of both fusion proteins at the chloroplast periphery.
(B) Three additional cells from pTF-mCherry and pTF.CatCh1-EYFP expressing cell line described in the
main text exhibit similar "chloroplast lining" colocalization pattern of both fusion proteins at the chloroplast
periphery. pTF-mCherry punctum was typically positioned adjacently to this signal. Scale bar is 10 μm.

 Figure 5—figure supplement 4. pTF.ap1 colocalizes with pTF.
 pTF.ap1-EYFP was co-expressed with pTF-mCherry and colocalization of the two fusion proteins—
 although not as obvious as in the case of pTF.CREG1 and pTF.CatCh1—was observed. These cells are
 representative of a pooled non-clonal *P. tricornutum* culture. Scale bar is 10 µm. Fusion protein schematics
 created with BioRender.com.

Figure 6—figure supplement 1. Conserved amino acid residues across pTF.CREG1 homologs. Protein sequence alignment (*Figure 6—source data 1*) leading to the phylogenetic tree shown in *Figure 6* was examined for amino acid residues that are at least 90% conserved across all homologs. 12 such amino acid residues (blue triangles) in nine examined regions were identified. Additional few amino acid residues surrounding each conserved site are shown. Putative loop tetrapeptide motifs (see the main text for details) shown to the far right directly precede the conserved proline and cysteine amino acid residues in region 4. The DPES tetrapeptide motif in pTF.CREG1 is also found in *Pteridomonas danica* (single-celled heterotrophic heterokont), a *Pterosperma* species (single-celled green alga), *Gossypium barbadense*, *Durio zibethinus*, and *Theobroma cacao* (all three are plants). Black arrows point to pTF.CREG1 and its paralog Phatr3_J10972. Human CREG1 and pTF.CREG1 sequences are shaded. Numbers above amino acid sequences indicate alignment position. *17 AAs between glycine (G) and arginine (R) in the conserved region 1 were omitted.

Figure 6—figure supplement 2. Bioinformatic, structural, and enzymatic insights suggest pTF.CREG1 has a non-catalytic function similar to human CREG1.

(A) pTF.CREG1 (UniProt ID: B7G9B3) features and its amino acid sequence alignment with human CREG1 (UniProt ID: O75629). pTF.CREG1 has a signal peptide that ends in SGA (blue underline; predicted with SignalP 4.1). This motif is closely related to Type I signal peptidase recognition site AXA indicating that pTF.CREG1 may enter the secretory pathway (*Tuteja, 2005*). Blue triangle points to a putative peptidase cut site. The core signal peptide sequence is enriched in hydrophobic amino acid residues, a common signal peptide feature (black underline). N-terminal methionine in Met-Phe (gray underline) may be acetylated by an N-terminal acetyltransferase (NAT), perhaps NatC or NatE which use Met-Phe sequence as their substrate (*Ree et al., 2018*). Tyrosine immediately succeeding SGA (green triangle) was used by ASAFind version 1.1.7 to classify pTF.CREG1 as a low confidence chloroplast-directed protein. Ser32 and Ser133 (red triangles) are predicted to undergo mucin-type O-glycosylation. Yellow triangle indicates an asparagine amino acid residue (Asn132) that is predicted to be N-glycosylated in human CREG1 (*Sacher et al., 2005*). These putative glycosylation sites further support the notion that the protein trafficks along the secretory pathway. Amino acid residues 141–144 in human CREG1 (DPQS)—which are located on the FMN binding pocket-occluding loop (*Sacher et al., 2005*)—are largely conserved (DPES) in pTF.CREG1 (shaded black rectangle). Finally, magenta triangle points to the arginine mutation site, Asn79, which may— like the corresponding Asp87 in human CREG1—prevent coordination of the terminal FMN phosphate (*Figure 6—figure supplement 3A*; *Sacher et al., 2005*).

(B) Structural comparison between human CREG1 (PDB ID: 1XHN) and pTF.CREG1 homology model (created with Phyre2 web portal) suggests pTF.CREG1 is unable to coordinate FMN at the possible dimer interface (*Figure 6—figure supplement 3B*). Human CREG1 does not bind FMN *in vitro* (*Sacher et al., 2005*) due to the FMN binding pocket-occluding loop (gray rectangle). Pro146 and Glu147 in pTF.CREG1 are predicted to protrude into this pocket similarly to their human CREG1 counterparts Pro142 and Gln143. Gray: human CREG1 chain A (i.e., monomer 1), colored by secondary structure: human CREG1 chain B (i.e., monomer 2), blue: pTF.CREG1 homology model. N- and C-termini in protein structures are depicted. Created with UCSF Chimera version 1.11.1 and BioRender.com.

(C) pTF.CREG1 does not reduce physiologically relevant flavins. Left: Comparison of NADPH oxidation (measured by decrease in absorbance at 320 nm) by recombinant pTF.CREG1 (long dashed black line) supplemented with flavins (flavin mononucleotide (FMN) and riboflavin) versus a no enzyme, water control (solid gray line). Right: The same assay, but here comparing a negative control protein without known NADPH oxidation activity (bovine serum albumin, BSA; short dashed black line) versus a no enzyme, water control (solid gray line). Experiments were started by the addition of protein or water blank and carried out in triplicate for each treatment. Background 320 nm absorbance of water blanks was subtracted from each replicate. Standard deviation is shown.

Figure 6—figure supplement 2—source data 1. Flavin reduction enzymatic assay data underpinning charts in *Figure 6—figure supplement 2C*.

Figure 6—figure supplement 3. pTF.CREG1 is unlikely able to coordinate the terminal FMN phosphate and may act as a dimer *in vivo*.

(A) Two mutated arginine amino acid residues in human CREG1—one (Asp87) can be inferred from *Sacher et al., 2005*—are believed to prevent the protein from coordinating the terminal FMN phosphate. Asn79 in pTF.CREG1 is a corresponding amino acid residue to the human Asp87 (inset). The steric similarity between their side chains sugests that pTF.CREG1 would also be unable to coordindate the terminal FMN phosphate even in the absence of the FMN binding pocket-occluding loop. Gray: human CREG1 chain A (i.e., monomer 1 from PDB ID: 1XHN), colored by secondary structure: human CREG1 chain B (i.e., monomer 2 from PDB ID: 1XHN), blue: pTF.CREG1 homology model. Created with UCSF Chimera version 1.11.1 and BioRender.com.

(B) Putative homodimerization region in the pTF.CREG1 homology model exhibits features of the human CREG1 dimer one. Human CREG1 chain A (i.e., monomer 1 from PDB ID: 1XHN) from the visualization presented in *Figure 6—figure supplement 2B* was hidden which left us with a chimeric human-diatom dimer. Colored by secondary structure: human CREG1 chain B (i.e., monomer 2 from PDB ID: 1XHN), blue: pTF.CREG1 homology model. Gray rectangles indicate the FMN binding pocket-occluding loop (see the main text). Black rectangle depicts a modeled β-sheet region in pTF.CREG1 that is absent from human CREG1. Colored in magenta, red, orange, and green is a possible homodimerization interface in pTF.CREG1. Amino acid residue composition (written in N-terminal→C-terminal direction) corresponding to each color is as follows: magenta: **P**ES**PV**C**A**R**LT**L**SG**, red: K**VSF**TL**T**E**ASL**, orange: **WGVL**T**TI**, green: **P**D**G**Q**PFG**N**VY**S**F**. Bold: hydrophobic, underlined: hydrophobic and/or following *Sacher et al., 2005*. Created with UCSF Chimera version 1.11.1 and BioRender.com

Figure 6—figure supplement 4. pTF.CREG1 expression and purification.
Briefly, genes encoding various pTF.CREG1 truncations were placed under arabinose-inducible promoter,
expressed in BL21 *E. coli* cells, and purified using cobalt, Co^{2+}, resin-based gravity flow columns. CTHF:
C-terminal His6 and FLAG tag. NTH: N-terminal His6 tag. L: PageRuler Plus Prestained Protein Ladder.
(A) Final elutions of various pTF.CREG1 truncations from small scale (50 mL) protein expression
experiments. Δ31_pTF.CREG1-His6 (bolded and black rectangle) was selected for scale-up.
(B) The final Δ31_pTF.CREG1-His6 elution from a large-scale (2 L) protein expression experiment. NuPage
4-12% Bis-Tris gels were used to resolve protein samples.

Figure 7—figure supplement 1. Conservation of CX(X)C motifs in pTF.CatCh1 homologs and
pTF.CatCh1 features.
(A) CXXC and the adjacent CXC motifs are present in all but two identified pTF.CatCh1 homologs. Six
additional cysteines are 100% conserved (*Figure 7—source data 1*). pTF.CatCh1 and ISIP2b (paralogs)
are marked with black arrows. Amino acid residues immediately preceding the CXXC motif are also shown
as aspartic acid residue (D) at this position is known to promote divalent metal ion binding (*Banci et al.,*
2002; *Banci et al., 2006*). Numbers above amino acid sequences indicate alignment position.
(B) pTF.CatCh1 contains two disordered C-terminal regions flanking the transmembrane domain. CXXC is
predicted to be in one of them by PONDR VSL2 (top). No cysteines are predicted to be in this disordered
region by IUPred2a (bottom). Approximate locations of the conserved CXC and CXXC motifs are depicted
with blue and orange triangles, respectively.
(C) pTF.CatCh1 features assuming disorder prediction with PONDR VSL2. Signal peptide (AAs 1–26; gray)
is followed by a cysteine-rich domain (20 cysteines between amino acid residue 34 and 226; pink), and two
disordered regions (green). The transmembrane (TM) domain (AAs 296–315; blue) nested between the
predicted disordered regions is flanked by a polyserine ($[S]_6$) and an arginine-rich ($RKL[R]_3$) amino acid
residue stretch. N- and C-terminus are shown. Created with BioRender.com.

Figure 7—figure supplement 2. pTF.ap1 phylogeny and conserved motifs.

(A) pTF.ap1 (black arrow) homologs are present in diatoms and other marine microeukaryotes. Homolog
search was performed with NCBI and marine microbial eukaryote (MMETSP) databases. Some notable
species include common polar diatoms *Fragilariopsis kergulensis* and *Fragilariopsis cylindrus*, toxin-
producing microeukaryotes (diatoms from the genus *Pseudo-nitzschia*, haptophyte *Prymnesium parvum*,
and pelagophyte *Aureococcus anophagefferens*), cosmopolitan diatom *Ditylum brightwelli*, haptophyte
Emiliana huxleyi, ultraplanktonic species *Pelagomonas calceolate*, colonial species *Prasinoderma coloniale*
and *Phaeocystis antarctica*. Scale bar: 0.1 substitutions per position.

(B) pTF.ap1 (black arrow) contains highly conserved motifs (five shown here). There are three 100%
conserved cysteine amino acid residues across pTF.ap1 homologs (*Figure 7—figure supplement 2—*
source data 1) in addition to the two here (blue triangles and gray shade). Numbers above amino acid
sequences indicate alignment position.

2396

Figure 7—figure supplement 2—source data 1. Amino acid sequence alignment underpinning the
phylogenetic tree in *Figure 7—figure supplement 2A*.

Figure 8—figure supplement 1. Outstanding biological problems in established model marine diatoms
poised to be advanced with proximity-dependent proteomic mapping approaches.
Two diatom compartments of immediate interest are pyrenoid—proteinaceous RuBisCO-containing
subchloroplastic organelle—and silica deposition vesicle (SDV)—diatom cell wall biosynthesis site. *P.*
tricornutum and *T. pseudonana*—genetically tractable model marine diatoms—are appropriate for
immediate dissection of pyrenoid and SDV proteomic composition, respectively. Created with
BioRender.com.

(A) Generate pTF-APEX2 *P. tricornutum* cell line

pTF-APEX2 episome

pTA-Mob mobilization plasmid

E. coli

conjugation

P. tricornutum

(B) Evaluate pTF-APEX2 expression, activity, and localization

genotyped pTF-APEX2 transconjugant

Western blot

Enzyme assay

Electron microscopy

(C) Proximity proteomics with pTF-APEX2 cell line

biotin-phenol H₂O₂

APEX2 pTF

pull-down

Proximity-dependent protein biotinylation

Protein ID | Gene ID

ATGC

CAGT

GCAG

Mass spectrometry and proteogenomic analysis

(D) Characterization of mass spectrometry hits

MS hit-EYFP

pTF-mCherry

Colocalization

A → B

Biochemical activity

protein alignment

Phylogeny

A

pTF-APEX2   WT

fix cell pellets in paraformaldehyde
and glutaraldehyde
↓
treat with diaminobenzidine
and H_2O_2 for 15 min
↓
post-fix in osmium
tetroxide
↓
embed in 3% agar
↓
cut and put agar
pieces into vials

pTF-APEX2   WT

fix agar pieces in
glutaraldehyde
↓
incubate in uranyl
acetate overnight
↓
dehydrate in ethanol, embed in
resin, cut ~100 nm thick slices

B

pTF-APEX2   WT

A

Part 1: Labeling

5x pTF-APEX2   5x WT

⬇

resuspend cell pellets in sorbitol

⬇

+ biotin-phenol | 90 min | 4°C
+ H_2O_2 | 20 min | 4°C

biotin-phenol ⬤
H_2O_2

⬇

quench, lyse, and perform
streptavidin blot

Part 2: Pull-down and MS

incubate cell lysates with magnetic
streptavidin beads at 4°C overnight

⬇

wash beads with cell lysis buffer,
KCl, Na_2CO_3, and urea

⬇

store beads at -80°C

⬇

on-bead trypsin digestion and tandem
mass tag (TMT) attachment

⬇

B

C

UniProt ID	APEX2/WT ratio	Ensembl ID	Iron-sensitive transcript	Chloroplast localization/ confidence	Annotation
B7GE67	6.75	Phatr3_J41423	yes	yes/low*	fructose-biphosphate aldolase FBAC5
B7GA90	4.20	Phatr3_J55031	yes	yes/low**	iron starvation induced protein (ISIP) 1
B5Y488	3.46	Phatr3_J46929	yes	no	siderophore-bound iron assimilation protein
B7G9B3	**3.15**	**Phatr3_J51183**	**yes**	**yes/low**	**CREG1-like protein**
B5Y3Y4	3.13	Phatr3_J54656	yes	no	heat shock protein (HSP)
B7G9B2	**2.43**	**Phatr3_J52498**	**yes**	**no**	**cell surface protein**
B7G9B0	**2.36**	**Phatr3_J54986**	**yes**	**yes/low**	**cell surface protein**
B7G195	2.32	Phatr3_J46448	yes	no	heat shock protein (HSP)
B7G9L9	1.82	Phatr3_J30139	yes	no	small GTPase
B7GBX2	1.71	Phatr3_J23497	yes	no	ATPase
B7FNT2	1.62	Phatr3_J42574	no	yes/low	α-carbonic anhydrase
B7GDH4	1.59	Phatr3_J41172	no	no	calreticulin/mobilferrin
B7G5Y2	1.57	Phatr3_J5651	no	no	14-3-3-like protein
B7FR50	1.52	Phatr3_J43251	no	no	small GTPase
B7G162	1.51	Phatr3_J27923	no	no	vacuolar-type H+-ATPase subunit A

A

Chr20

415,875

pTF.ap1 →	*ISIP2b* ←	*pTF.CatCh1* →	*pTF.CREG1* ←
1182	1168	1155	705

537   403   410

421,434

	Ensembl ID	UniProt ID
	Phatr3_J54986	B7G9B0
	Phatr3_J54987	B7G9B1
	Phatr3_J52498	B7G9B2
	Phatr3_J51183	B7G9B3

B

pTF.CREG1

pTF-mCherry	pTF.CREG1-EYFP	Autofluorescence
Merge	Merge + BF	Brightfield (BF)

pTF.CatCh1

pTF-mCherry	pTF.CatCh1-EYFP	Autofluorescence
Merge	Merge + BF	Brightfield (BF)

A

Signal peptide

MFSFKSFVFAALLSSCEASVRARKLSNELIAGFEPRTVVTDHNAIDL**D**QAKIEDLLATPSTDTFSSARDV**Y**EQGSHS
KSFAVLTLDPALTVAVPEGTVISGKNEDGSTVSGTALKSYSVGDTTIEIQYSTGNTQATYVDCQVGGNPDPNTSGCF
AANGTVTISGSSGSLPYSYDPLVNNDNG**R**TIQGFSTAAETRFRPSGGGELFPDFQKFFD**Y**YGTLTYADEWVQSAFAR
RSTSFSKGDADFSKYGDDGILQAVKKGTAYISIAMYVIRELEDALDDCDRGCDTADCNDDAV**H**ALDEAVAFWTGSLE
GKDGSGSGVLMYGLADTRCINFKTCGRNGDEASGKSKVNFEIFRHFDTMQRELTNKRCNSARATKEKIVPWMIVPLI
QGVLRYAYIIENDGFTEKAEAEGATFAAAVLPYVASCNAEDAKTIYDNMRVGLGGGASYSTVFDAFVRQSSCMGISC
SDIGGYWNEAQGAYEEGAGPCGGSGSGDDGANVGLAVGLSIGGLLVLVLLGVVLRRRRSASGSTVEFKDTGSTPI*

Transmembrane   **Tail**
domain   **domain**

B

pTF-APEX2

WT

C

Amplex UltraRed   +   +   −   +   +   +
H₂O₂   +   −   +   +   +   +

s3   **s1**   **s2**   **s4**

A

pTF-APEX2 expressing *P. tricornutum* cells embedded in agar

B

C

0%   100%

D

→ Phatr3_J18572   KRPLGASE
  HRP   PAPFF---
→ Phatr3_Jdraft1645 ASPDN---
→ Phatr3_J46616   PNPGL---
  Phatr3_J54731   PDAEP--G
  Phatr3_J47395   PDAEP--G
  Phatr3_J13244   PNADV--G
→ Phatr3_Jdraft1023 PDADK--G
  Phatr3_EG02288   PDATQ--G
  APEX2   PDPTK--G

E

Ensembl ID	Peroxidase class[1]	Position 134 relative to APEX2[2]	Predicted localization[3]	Number of transmembrane domains[4]
Phatr3_J18572	I	**Pro**	cytosol	/
Phatr3_Jdraft1645	I	**Pro**	secretory pathway	1
Phatr3_J46616	I	**Pro**	secretory pathway	/
Phatr3_J54731	I	Ala	cytosol	/
Phatr3_J47395	I	Ala	secretory pathway	/
Phatr3_J13244	I	Ala	secretory pathway	/
Phatr3_Jdraft1023	I	Ala	**mitochondria**	/
Phatr3_EG02288	I	Ala	cytosol	/

Growth conditions and sampling timepoints

A

B

C

A

| pTF-mCherry | pTF.CREG1-EYFP | Autofluorescence | Merge | Merge + BF | Brightfield (BF) |

B

| pTF-mCherry | pTF.CatCh1-EYFP | Autofluorescence | Merge | Merge + BF | Brightfield (BF) |

A

B

C

A

B

2x 90°

unique to
pTF.CREG1

A

B

C

Phaeodactylum tricornutum
(pennate model diatom)

fusiform

triradiate

oval

Compartment

chloroplast

→ **pyrenoid** →

Broader themes

carbon dioxide fixation
molecular self-assembly

Thalassiosira pseudonana
(centric model diatom)

Compartment

→ **silica deposition vesicle
(SDV)** →

Broader themes

biomineral morphogenesis
nanotechnology