Running title: Bacterial community profiling highlights complex diversity and novel

organisms in wildlife ticks.

Authors: Siobhon L. Egan[1], Siew-May Loh[1], Peter B. Banks[2], Amber Gillett[3], Liisa

Ahlstrom[4], Una M. Ryan[1], Peter J. Irwin[1] and Charlotte L. Oskam[1,*]

[1] Vector and Waterborne Pathogens Research Group, College of Science, Health,

Engineering and Education, Murdoch University, Perth, Western Australia, Australia

[2] School of Life and Environmental Sciences, The University of Sydney, Sydney, New South

Wales, Australia

[3] Australia Zoo Wildlife Hospital, Beerwah, Queensland, Australia

[4] Bayer Australia Ltd, Animal Health, Pymble, New South Wales, Australia

[*] Corresponding author

E-mail addresses: siobhon.egan@murdoch.edu.au (S. L. Egan); kim_lsm@yahoo.com (S-M.

Loh); peter.banks@sydney.edu.au (P. B. Banks); drambergillett@hotmail.com (A. Gillett);

liisa.ahlstrom@bayer.com (L. Ahlstrom); una.ryan@murdoch.edu.au (U. M. Ryan);

p.irwin@murdoch.edu.au (P. J. Irwin); c.oskam@murdoch.edu.au (C. L. Oskam).

Abstract

Ticks (Acari: Ixodida) transmit a greater variety of pathogens than any other blood-feeding

group of arthropods. While numerous microbes have been identified inhabiting Australian

Ixodidae, some of which are related to globally important tick-borne pathogens, little is

known about the bacterial communities within ticks collected from Australian wildlife. In this

study, 1,019 ticks were identified on 221 hosts spanning 27 wildlife species. Next-generation

sequencing was used to amplify the V1-2 hypervariable region of the bacterial 16S rRNA

gene from 238 ticks; *Amblyomma triguttatum* (n=6), *Bothriocroton auruginans* (n=11),

Bothriocroton concolor (n=20), *Haemaphysalis bancrofti* (n=10), *Haemaphysalis bremneri*

(n=4), *Haemaphysalis humerosa* (n=13)*, *Haemaphysalis longicornis* (n=4), *Ixodes antechini*

(n=29), *Ixodes australiensis* (n=26), *Ixodes fecialis* (n=13), *Ixodes holocyclus* (n=37), *Ixodes*

*myrmecobii (*n=1), *Ixodes ornithorhynchi* (n=10), *Ixodes tasmani* (n=51) and *Ixodes*

trichosuri (n=3). After bioinformatic analyses, over 14 million assigned bacterial sequences

revealed the presence of recently described bacteria '*Candidatus* Borrelia tachyglossi',

'*Candidatus* Neoehrlichia australis', '*Candidatus* Neoehrlichia arcana' and '*Candidatus*

Ehrlichia ornithorhynchi'. Furthermore, three novel Anaplasmataceae species were identified

in the present study including; a *Neoehrlichia* sp. in *I. australiensis* and *I. fecialis* collected

from quenda (*Isoodon fusciventer*) (Western Australia), an *Anaplasma* sp. from one *B.*

concolor from echidna (*Tachyglossus aculeatus*) (New South Wales), and an *Ehrlichia* sp.

from a single *I. fecialis* parasitising a quenda (WA). This study highlights the diversity of

bacterial genera harboured within wildlife ticks, which may prove to be of medical and/or

veterinary importance in the future.

Keywords: Microbiome; ticks; Ixodida; wildlife; marsupials; Anaplasmataceae

1. Introduction

Current estimates suggest that approximately 17% of all infectious diseases of humans are

vector-borne (Rinker et al., 2016) and global trends show that vector-borne diseases (VBDs)

are rising at a rapid rate (Jones et al., 2008; Morens and Fauci, 2012). The complex interplay

between pathogen, vector, host(s) and the environment make VBDs particularly challenging

to understand. In addition, factors such as climate change (Ostfeld and Brunner, 2015), land

use (Ferrell and Brinkerhoff, 2018), feral animal populations (Merrill et al., 2018) and the

microclimate within a landscape (Dobson et al., 2011) can further influence the prevalence

and distribution of VBDs.

Ticks (Acari: Ixodida) comprise a group of haematophagous (blood feeding) arthropods with

over 900 species described globally (Guglielmone et al., 2014; Mans et al., 2019). Ticks are

known to transmit various pathogens; however, they also harbour a range of endosymbiont

and commensal species (Špitalská et al., 2018). The epidemiology of recognised tick-borne

diseases (TBDs) in the northern hemisphere demonstrates that wildlife serve as sentinels and

can be used to monitor the presence and distribution of tick-borne pathogens (TBPs).

Importantly, research has shown that some wildlife species act as dilution hosts for certain

TBPs, whereas others may act as amplification hosts (LoGiudice et al., 2003).

The complexity of TBDs means that studies are increasingly shifting away from isolated

species-specific studies toward ecosystem-based, collaborative research (Estrada-Peña et al.,

2013; Pfäffle et al., 2013). Metabarcoding provides an informative molecular tool to

characterise the bacterial diversity in ticks. Worldwide, next-generation sequencing (NGS)-

based analyses have been applied to a range of tick species that are important from medical

and veterinary perspectives, including *Amblyomma americanum* (Ponnusamy et al., 2014),

Ixodes ricinus (Bonnet et al., 2014), and *Rhipicephalus microplus* (Andreotti et al., 2011).

Metabarcoding studies of the bacterial microbiome of Australian ticks have been reported

only recently, with the first bacterial profiling by NGS conducted on the human-biting tick

Ixodes holocyclus (Gofton et al., 2015a). These authors identified a highly abundant

endosymbiont 'Candidatus Midichloria mitochondrii' (CMm) and after blocking the

amplification of this organism, a greater bacterial diversity was revealed, including a number

of novel microbes (Gofton et al., 2015a,b). Critically, in contrast to many parts of the world

where multiple TBPs have been elucidated within well-studied tick-host-environment

ecologies, there is a relative dearth of such information available for Australia. With this in

mind, the aims of our study were to survey the bacterial communities present in ticks

collected from Australian wildlife and to investigate their genetic relatedness to 'taxa of

interest', i.e. tick-associated pathogenic and endosymbiotic organisms (Parola and Raoult,

2001; Mediannikov and Fenollar, 2014; Sumrandee et al., 2016). 'Taxa of interest' in the

present study were defined as genera within alphaproteobacteria, gammaproteobacteria and

spirochaetes known to be transmitted by ticks in other parts of the world, specifically;

Anaplasma, *Bartonella*, *Borrelia*, *Coxiella*, *Ehrlichia*, *Francisella*, *Midichloria*,

Neoehrlichia, *Rickettsia* and *Rickettsiella*. (Ahantarig et al., 2013; Vayssier-Taussat et al.,

2015; Bonnet et al., 2017; de la Fuente et al., 2017).

2. Materials and methods

2.1 Sample collection and identification

1,019 ticks were sourced opportunistically from wildlife in Australia by veterinarians,

veterinary nurses, wildlife carers, researchers and via submissions from members of the

public, and preserved in 70% ethanol before being shipped to Murdoch University, Western

Australia (WA), for identification. Ticks were identified morphologically to life stage and

species using keys and species' descriptions (Roberts, 1970; Barker and Walker, 2014).

Details of sample collection are available in Supplementary File S1.

2.2 DNA Extraction

A sub-sample of ticks (*n*=238) was chosen for DNA extraction and bacterial profiling. Ticks

were selected to represent as many tick species, hosts, geographical regions and life stages as

possible within the study. Prior to DNA extraction, individual ticks were surface-sterilised in

10% hypochlorite solution, rinsed in 70% ethanol and DNA-free PBS, and then air-dried.

Total genomic DNA (gDNA) was extracted using the Qiagen DNeasy Blood and Tissue kit

(Qiagen, Germany) following the manufacturer's recommendations with the following

modifications; ticks were placed in a 2 mL safe lock Eppendorf tube with a 5 mm steel bead,

frozen in liquid nitrogen for 1 minute and homogenised by shaking at 40 Hz in a Tissue Lyser

LT (Qiagen, Germany). The volume of elution buffer AE was adjusted to 200 μL for

engorged females, 100 μL for unengorged adults and 40 μL for nymphs. A double elution

was carried out to increase gDNA yield for unengorged adults and nymphs. Sterile and DNA-

free equipment and tubes were used for each step. Extraction reagents blank (EXB) controls

were performed alongside tick extractions to assess background bacterial communities.

2.3 16S amplification and library preparation for NGS sequencing

Amplicons targeting a 250-320 bp product of the V1-2 hypervariable region of the 16S rRNA

(16S) gene were generated using the primer pair 27F-Y (Gofton et al., 2015b) and 338R

(Turner et al., 1999). Previous research identified that paralysis tick, *I. holocyclus*, has a

highly abundant bacterial species ('*Ca*. M. mitochondrii'), which masks the diversity of

bacteria in 16S metabarcoding studies (see Gofton et al. 2015a). Therefore, a blocking primer

developed by Gofton et al. (2015a) was used to inhibit the amplification of '*Ca*. M.

mitochondrii' in *I. holocyclus* ticks. Depending on tick life stage and level of engorgement, 3-

10 μM of the CMm blocking primer was added to the PCR reaction. Amplicon PCRs were

conducted in 25 μL reactions each containing 1 X Buffer (KAPA Biosystems, USA), 1.5 mM

MgCl$_2$, 0.4 μM of each forward and reverse primer with MiSeq adapters, 0.4 mg/mL BSA

(Fisher Biotech, Australia), 0.25 mM dNTPs (Fisher Biotech, Australia), 0.5 U KAPA Taq

(KAPA Biosystems, USA), and 2.0 μL of undiluted genomic DNA. Samples underwent the

following thermal cycling conditions; initial denaturation at 95°C for 5 mins, followed by 35

cycles of denaturation at 95°C for 30 s, annealing at 62°C (CMm blocking primer present) or

55°C (CMm blocking primer absent) for 30 s, and extension at 72°C for 45 s with a final

extension at 72°C for 5 mins. Libraries were then prepared using the Nextera XT DNA

library preparation kit in 25 μL reactions following manufacturer's recommendations.

Libraries were purified using Agencourt® AMPure® XP PCR purification beads (Beckman

Coulter Life Sciences, USA) and pooled in equimolar amounts. Final libraries were then

quantified using the Qubit® 2.0 Fluorometer (Thermo Fisher, Australia). Libraries were

sequenced on an Illumina MiSeq™ using v2 chemistry (2x250 paired end).

Extraction reagents blanks (n=12) and PCR no-template controls (NTC; n=7) were included

in all stages of the workflow. All pre- and post-PCR procedures were performed in physically

separate dedicated laboratories and sterile protocols were maintained through library

preparation in order to minimise amplicon contamination.

2.4 Bioinformatics and statistical analysis

Raw fastq files were downloaded from the Illumina BaseSpace Sequence Hub for analysis in

a custom pipeline using USEARCH (Edgar, 2010). Raw paired-end sequences were merged

in USEARCH v10, a minimum of 50 nucleotide (nt) overlap and maximum number of

mismatches increased to 15 nt due to long overlap of paired-end sequences. Only sequences

with perfect primer sequences were retained, primer sequences and distal bases were trimmed

using USEARCH v8.0. Sequences were quality filtered in USEARCH v10, allowing a <1%

expected error rate and singletons were discarded (Edgar and Flyvbjerg, 2015). Sequences

were then clustered into operational taxonomic units (OTUs) of 97% similarity using the

UPARSE algorithm (Edgar, 2013) in USEARCH v10. Taxonomy was assigned in QIIME2

v2018.4 using the q2-feature-classifier (Bokulich et al., 2018) with reference to a trained

Greengenes database (DeSantis et al., 2006) (release May 2013) using the primer pair 27F-Y/

338R. Taxonomic assignments were confirmed using NCBI MegaBLAST (Morgulis et al.,

2008) on a random subsample of OTUs and in the case of tick-associated microbes, GenBank

accession numbers and percentage identity of top hits were recorded for 'taxa of interest'.

The profiles from EXB controls and NTCs were first assessed to ensure quality of sampling

and absence of tick-associated bacteria. The following criteria were used to assess inclusion

of sequences: all OTUs that appeared exclusively in controls were removed; OTUs that had a

higher relative sequence abundance in controls compared to tick samples were removed; and

OTUs were removed that appeared in over half of controls (i.e. at least eight) that had a

taxonomic identity associated with environmental bacteria (e.g. members of the phyla

Acidobacteria and Cyanobacteria). In addition, potential cross-contamination during library

preparation or 'cross-talk' at the sequencing level was assessed by inspecting the presence of

expected tick-associated bacteria (e.g. members of the obligate intracellular bacterial families

Anaplasmataceae and Midichloriaceae) in controls. The profiles of EXB controls and NTCs

were then removed bioinformatically from associated samples to eliminate background

bacteria.

Data analysis and visualisation was carried out in RStudio (RStudio Team, 2015) using

packages metacoder (Poisot et al., 2017), phyloseq (McMurdie and Holmes, 2013) and vegan

(Oksanen et al., 2019). Alpha diversity of samples was measured using observed OTUs,

Chao1 index, Shannon index and Simpson index. Removal of samples with a low sequencing

depth (<1000 assigned OTUs after data filtering), did not significantly alter alpha diversity

measurements among tick species (data not shown). Rarefaction curves for samples were

calculated to assess sequencing depth based on observed number of OTUs. Principal

coordinate analysis (PCoA) on weighted unifrac dissimilarity measurements was used to

assess the differences in microbial composition between tick species. Investigation into 'taxa

of interest' warranted a more rigorous assessment of sequence number than was required for

diversity measures in order to avoid any potential cross-contamination and machine cross-

talk (Dong et al., 2017; Wang et al., 2017), as such only samples with >100 sequences were

considered positive. Taxonomic assignment to 'taxa of interest' was assessed based on NCBI

MegaBLAST top hit of named organism. Where precent identity was ≤97%, the terminology

of nearest named Genus-like was employed. Due to sample collection biases in the present

study, prevalence data of these microbes were not considered statistically relevant.

Nucleotide sequences from 'taxa of interest' were aligned by MUSCLE (Edgar, 2004) using

default parameters and aligned sequences were then imported into MEGA7 (Kumar et al.,

2016) with the most appropriate nucleotide substitution model chosen based on the lowest

Bayesian Information Criterion (BIC) score. Evolutionary histories were inferred using the

Neighbour-Joining method based on the Tamura 3-parameter model (Tamura, 1992).

Bootstrap analysis was conducted using 10,000 replicates to assess the reliability of inferred

tree topologies.

3. Results

3.1 Tick-host associations

Ticks were collected from 221 wildlife hosts including 24 native and three introduced species

(Fig. 1, Supplementary File S1). Hosts recorded in the present study included; agile

antechinus (*Antechinus agilis*), brown antechinus (*Antechinus stuartii*), spotted-tail quoll

(*Dasyurus maculatus*), water rat (*Hydromys chrysogaster*), quenda (*Isoodon fusciventer*),

northern brown bandicoot (*Isoodon macrourus*), western grey kangaroo (*Macropus*

fuliginosus), eastern grey kangaroo (*Macropus giganteus*), red-necked wallaby

(*Notamacropus rufogriseus*), red kangaroo (*Osphranter rufus*), platypus (*Ornithorhynchus*

anatinus), eastern barred bandicoot (*Perameles gunnii*), long-nosed bandicoot (*Perameles*

nasuta), sugar glider (*Petaurus breviceps*), eastern ring-tailed possum (*Pseudocheirus*

peregrinus), black fruit-bat (*Pteropus alecto*), fruit-bat sp. (*Pteropus* sp.), bush rat (*Rattus*

fuscipes), black rat (*Rattus rattus*), Tasmanian devil (*Sarcophilus harrisii*), wild pig (*Sus*

scrofa), short-beaked echidna (*Tachyglossus aculeatus*), rufous-bellied pademelon

(*Thylogale billardierii*), short-eared brush-tailed possum (*Trichosurus caninus*), common

brush-tailed possum (*Trichosurus vulpecula*), wombat (*Vombatus ursinus*), red fox (*Vulpes*

vulpes), and swamp wallaby (*Wallabia bicolor*). Female ticks were the dominant life stage

recorded (n=547) followed by nymphs (n=319) and males (n=153). Ticks were received from

animals in the Northern Territory (NT) (n=16), Queensland (QLD) (n=249), New South

Wales (NSW) (n=316), Tasmania (TAS) (n=136), Victoria (VIC) (n=44) and Western

Australia (WA) (n=258). Together, *Bothriocroton concolor* (n=123), *Ixodes australiensis*

(n=210), *I. holocyclus* (n=173) and *Ixodes tasmani* (n=184) accounted for over two-thirds of

all ticks submitted.

3.2 16S rRNA bacterial profiling

A sub-sample of 238 ticks and 19 controls underwent 16S NGS profiling. A total of 23.9

million raw paired-end sequences were generated on the Illumina MiSeq. 17.9 million

sequences were retained after merging, and subsequent quality filtering yielded 14.8 million

sequences for clustering and taxonomic assignment. A total of 4,864 OTUs (average length

299 bases) were retained after background profiles were removed. After removal of

background sequences, a total of 14,328,059 bacterial sequences were assigned to tick

samples. Despite a high number of OTUs, only 1,535 OTUs had greater than 100 total

sequences from tick samples. Tick samples had an average of 60,201 assigned sequences (see

Supplementary File S2). *Amblyomma triguttatum* had the highest median alpha diversity as

measured by the observed number of OTUs and the chao1 index, whereas *Ixodes antechini*

had the highest median alpha diversity as measured by the Shannon and Simpson indexes

(Fig. 2). Rarefaction analysis of the sequence depth shows that the observed number of OTUs

plateaued at 50,000 sequences (Fig. 3). After the removal of sequences from controls, eight

phyla were retained. An ordination plot of OTUs (Fig. 4) show that bacteria belonging to the

Proteobacteria phylum were the most abundant and diverse taxa classified, followed by

Firmicutes and Actinobacteria. Bacterial families identified in tick species, represented as

relative number of sequences in Fig. 5, show 26 dominant taxa. While bacterial composition

varied between tick species, sequences from the members of Coxiellaceae, Francisellaceae,

and Rickettsiaceae families (Phylum: Proteobacteria) were the most abundant. Although

sample sizes varied among tick species, beta diversity analysis incorporating abundance

(sequence number) and taxonomic relatedness, showed evidence of different bacterial

communities among tick species (Supplementary File S3).

3.3 Presence of 'taxa of interest'

In total, 37 OTUs were identified as 'taxa of interest' in the present study. Their taxonomic

identity and abundance (number of sequences) in each sample is available in Supplementary

File S4 (see Supplementary File S5 for fasta file of sequences). Phylogenetic analysis of the

'taxa of interest' is represented in Fig. 6, showing tick species and Australian states and

territory, where each OTU was identified.

A proposed novel *Anaplasma* sp. (OTU230, MK814412, 96.3% identity) was identified in a

single *B. concolor* (1/20) collected from (ex) echidna from NSW. A second *Anaplasma*

bovis-like OTU (OTU312, JN862824, 97.3% identity) was identified in *Haemaphysalis*

bancrofti (3/10) ex red-necked wallaby and long-nosed bandicoot and *Haemaphysalis*

humerosa (n = 1/13) tick ex long-nosed bandicoot, all from NSW.

OTU5509 was assigned to the genus *Bartonella* and had a top BLAST hit of *Bartonella*

bacilliformis (LN624026, 92.1% identity). OTU5509 was identified in two samples,

however, was present in an extremely low number of sequences (2) in each case. Both

samples were *I. fecialis* collected from two different quenda in WA. The low number of

sequences means that in the case of statistical analysis, this OTU would have been filtered

out. While there was no sufficient match to this sequence, it is noted that in many instances

native Australian *Bartonella* species are lacking sequence information for this region of the

16S gene (V1-2).

Two OTUs (OTU14, CP025785, 100% identity; and OTU4629, CP025785, 100% identity)

were identified as *'Ca.* B. tachyglossi'. Sequences were identified in *Bothriocroton concolor*

(8/20) ex echidnas from QLD & NSW and *H. humerosa* (1/13) ex northern brown bandicoot

from QLD.

A *Coxiella*-like organism (OTU12, CP032542, 95.5% identity) was identified exclusively in

B. concolor (19/20) ex echidna from NSW and QLD. This was the second most abundant

OTU identified in *B. concolor*, accounting for ~20% of the overall sequences.

A *Coxiella* sp. (OTU15, KC170757, 100% identity) was identified in *Haemaphysalis*

longicornis (3/4) ex red fox from NSW and, OTU represented ~61.4% of the overall assigned

bacterial sequences.

A novel *Ehrlichia* sp. (OTU33, AY309970, 96.3% identity) was identified in *I. fecialis* (1/13)

ex quenda from WA. Other *Ehrlichia* OTUs (OTU16 and 1632) represented sequences from

'Ca. Ehrlichia ornithorhynchi' in *Ixodes ornithorhynchi* (6/10) ex platypus QLD and these

sequences accounted for the majority of sequences (~50.2%) from *I. ornithorhynchi*.

A *Francisella* endosymbiont (OTU5, AF001077, 98.0% identity) was identified in *H.*

humerosa (11/13) ticks ex northern brown bandicoots from NT and QLD, and

Haemaphysalis bremneri (2/4) ex possum (species unknown) in QLD. OTU5 was the most

abundance sequence in *H. bremneri* and *H. humerosa*, representing 27.3% and 52.4% of the

assigned sequences, respectively. A second *Francisella* endosymbiont (OTU10, AB001522,

99% identity) was identified in *H. bancrofti* (8/11) ex red-necked wallabies from QLD and

NSW, and ex a long-nosed bandicoot in NSW; and in a single *H. humerosa* (1/13) ex red-

necked wallaby from NSW. OTU10 was the most abundance sequences in *H. bancrofti*

representing 27.0% of the assigned sequences. OTU40 (AF001077, 97.6% identity) was also

identified as a *Francisella*-like endosymbiont from *A. triguttatum* (6/6) ex red kangaroo from

WA. It was highly abundant in these ticks, accounting for 76.9% of the overall sequences.

Midichloria (OTU6, FM992372, 100% identity) was identified in *I. holocyclus* (20/36) ex

long-nosed bandicoots from NSW and QLD, *I. fecialis* (1/13) ex red-necked wallaby NSW,

H. bancrofti (1/11) ex long-nosed bandicoot NSW, and *H. humerosa* (1/13) ex long-nosed

bandicoot NSW. *Midichloria* (OTU7, FM992373, 100% identity) was identified in *I.*

holocyclus (15/36) ex long-nosed bandicoots from NSW and QLD, *I. fecialis* (1/13) ex red-

necked wallaby NSW and *H. bancrofti* (1/11) ex long-nosed bandicoot NSW. Overall a total

of 15 ticks had both OTU6 and OTU 7, which included *I. holocyclus* (13/36) ex long-nosed

bandicoots from NSW and QLD, *I. fecialis* (1/13) ex red-necked wallaby NSW and *H.*

bancrofti (1/11) ex long-nosed bandicoot NSW. While the CMm blocking primer was

incorporated into the PCRs conducted on *I. holocyclus, Midichloria* sequence were still

observed in 2/9 females, 3/6 males and 16/21 nymphs. *Midichloria* was the most abundant

sequence from *I. holocyclus* accounting for 36.1% of assigned sequences.

Three *Neoehrlichia* OTUs were identified. '*Ca*. Neoehrlichia arcana' (OTU25, KT203914,

99.3% identity) was identified in eight ticks, *I. tasmani* (4/51), *I. holocyclus* (2/36)*, H.*

bancrofti (1/11) and *H. humerosa* (1/13), ex long-nosed bandicoots NSW and QLD. '*Ca*.

Neoehrlichia australis' (OTU29, KT803957, 100% identity) was identified in 10 ticks, *I.*

holocyclus (8/36) and *I. tasmani* (2/51)*,* ex long-nosed bandicoots NSW and QLD. Three

ticks (*I. tasmani* and *I. holocyclu*s) were co-infected with both '*Ca*. N arcana' and '*Ca*. N.

australis'. A novel *Neoehrlichia* species (OTU31, MG670107, 97.3% identity) was identified

in *I. fecialis* (2/13) and *I. australiensis* (2/26) ex quenda from WA, and *I. antechini* (2/29) ex

brown antechinus from NSW.

Due to the conserved nature of the *Rickettsia* genus at the 16S gene, reported previously

(Roux and Raoult, 1995; Stothard and Fuerst, 1995), resolution to species level was not

possible using the sequences obtained in the present study. *Rickettsia* sp. (OTU2, KF318168,

99.7% identity) had a widespread distribution among samples and was detected in 61 ticks

(B. concolor, H. bancrofti, H. humerosa, I. antechini, I. australiensis, I. holocyclus and *I.*

tasmani) from three states and one territory (NSW, NT, QLD, WA). *Rickettsia* was the most

abundant sequence from *B. concolor* and *I. antechini* ticks accounting for 63.3% and 12.0%

of the assigned sequences, respectively

A *Rickettsiella*-like sp. (OTU1, LC388767, 92.7% identity) was identified in *I. australiensis*

(22/26) and was the most abundant sequence accounting for 76.8% of assigned sequences. A

Rickettsiella-like sp. (OTU3, EU430251, 95.1% identity) was identified in *I. tasmani* (11/51)

ex Tasmanian devils and eastern barred bandicoot from TAS. A *Rickettsiella*-like sp. (OTU4,

U97547, 98.4% identity) was identified in, *I. tasmani* (36/51) ex bandicoots, possum and a

sugar glider from NSW, TAS and QLD; *I. australiensis* (1/26) ex western grey kangaroo

from WA; and *I. holocyclus* (1/36) ex long nosed bandicoot NSW. OTU4 accounted for

30.2% of the overall sequences assigned for *I. tasmani.*

4. Discussion

Currently there is a scarcity of detailed information about the life cycles of Australian ticks,

however it is generally assumed that, as with other hard tick species around the world, the

majority will exhibit a three-host life cycle (Barker and Walker, 2014). Despite these

limitations, the opportunistic sampling strategy used in this study provides an economical

way to survey a wide range of tick and wildlife fauna across a range of geographical areas

across Australia. The paucity of larvae from this data set is also a limitation of this type of

sampling. Tick larval stages, which may indicate the presence of transovarially-transmitted

organisms (Kwan et al., 2017), are difficult to see with the naked eye, and in situations where

collection of ticks is not the main objective, they are easily overlooked (Lydecker et al.,

2019). Furthermore, the opportunistic nature of the sampling precluded assessment of the

infestation rates of the various hosts. In future, if TBPs of humans and the wildlife

themselves are identified, further extensive surveys of tick-pathogen-wildlife ecologies will

require more targeted and comprehensive approaches in order to gather sufficient relevant

epidemiological data.

As with previous studies (Swei and Kwan, 2016; Zolnik et al., 2016) alpha diversity

measures were highly variable both within and between tick species. This may be driven by a

variety of factors such as; starting material (i.e. size/life stage of tick), extraction methods,

library preparation, normalisation of DNA input concentrations for sequencing, batch

sequencing effects and bioinformatics (Greay et al. 2018). Within the present study,

sequencing depth was likely most impacted by sample input type and normalisation of DNA

concentrations, as all samples went through the same library preparation and bioinformatic

analysis. Overall, bacterial diversity of ticks started to plateau by 50,000 sequences however,

it is noted that in some cases this plateau was not achieved and deeper sequencing would be

required in order to confidently characterise the full suite of bacterial taxa present. This was

most evident in *A. triguttatum*, *B. auruginans*, *I. trichosuri* and *I. ornithorhynchi* and

therefore we recommend a minimum of 100,000 sequences per sample for future bacterial

16S amplicon studies in these species to be confident that the complete bacterial community

has been sampled. Studies that investigate the shift of the core microbiome (i.e. most

abundant bacteria taxa) relative to a given parameter may not require the same depth of

sequencing, as seen in previous studies investigating the effect of temperature (Thapa et al.,

2019) and life stage (Andreotti et al., 2011).

After bioinformatic analysis, including stringent quality filtering, a large number of OTUs

remained with relatively low number of reads. This has been noted in previous studies of the

tick microbiome (Budachetri et al., 2016; Zolnik et al., 2016) and in the broader field of high-

throughput microbiome studies (Pollock et al., 2018). Depending on bioinformatic analysis

and quality filtering, the number of OTUs and therefore diversity can vary greatly among

studies (Greay et al., 2018). In addition, current practice in microbiome studies is to

normalise count data, however models show that this can oversimplify the data (McMurdie

and Holmes, 2014).

Ordination analysis demonstrated that tick species was the strongest predictor of bacterial

composition. Beta-diversity analysis showed that the common marsupial tick, *I. tasmani*,

exhibited a variable bacterial composition. This diversity may be explained by the wide

geographic distribution of this tick in Australia, its own high genetic diversity (Burnard and

Shao, 2019), and its ability to parasitise many marsupial species. In the present study,

bacterial profiling of *I. tasmani* included ticks from three states (NSW, QLD, TAS) and seven

host species. With this in mind, we suggest that future studies should combine careful

taxonomic status identity with data on microbial communities of *I. tasmani*. Although sample

sizes were limited in 'host-specialist' ticks *B. auruginans* (wombats)*, H. bremneri* (possums)*,

I. antechini (antechinus) and *I. ornithorhynchi* (platypus), these specimens showed less

diversity of bacterial communities between samples.

An important caveat on the relative diversity of bacterial communities in the present study

was the use of a blocking primer inhibiting the amplification of '*Ca*. M. mitochondrii'. The

inclusion of this blocking primer it vital to explore the full bacterial community of *I.*

holocyclus (see Gofton et al., 2015), however it does impact the analysis and interpretation of

the data. Due to the inhibition of this bacteria, there is an inherent bias in the bacterial

community, and comparisons in alpha and beta-diversity measures must account for this. In

the case of highly abundant organisms, the use of a blocking primer assay will likely not

completely inhibit amplification, and thus a reduced level of the organisms may be still

detected, as was the case in the present study. In this instance, the use of diversity measures

that rely on presence/absence data will not be affected by this granted there is still some level

of detection (i.e. number of species observed for alpha diversity and Jaccard index for beta-

diversity). The use of more advanced, and usually preferred, diversity indexes may be

impacted by the manipulation of bacterial composition, and care must be taken when

comparing results with other studies (Greay et al., 2018). Despite these limitations, the use of

blocking primers has shown to be vital in the context of investigating known tick-borne

pathogens and novel related taxa from tick samples. In the present study we have

characterised the bacterial communities of ten tick species for the first time. This information

provides the first step of focusing future research and where this established technique should

be applied.

Members of the Proteobacteria represented the most diverse and abundant (as determine by

number of sequences) in Australian hard ticks analysed in the present study. This finding is

consistent with previous microbiome studies from tick species, such as *Ixodes scapularis*

(Sperling et al., 2017; Thapa et al., 2019), *Ixodes persulcatus* (Zhang et al., 2014; Kurilshikov

et al., 2015) and *Amblyomma americanum* (Fryxell and DeBruyn, 2016). Additionally the

identification of novel bacterial taxa in native Australian ticks species is consistent with

recent findings (Gofton et al., 2015b; Panetta et al., 2017).

Currently in Australia, the ecology, epidemiology, and incidence of human TBDs remains

largely a matter of conjecture, having received little scientific study compared with many

other parts of the world (Graves and Stenos, 2017). Despite significant national interest,

including a federal government senate inquiry (Radcliffe et al., 2016), the prevailing

scientific opinion concludes that Lyme borreliosis (caused by *B. burgdorferi* sensu lato), for

example, is absent from Australia (Chalada et al., 2016; Irwin et al., 2017). In the unique

Australian environment, long isolated in geological terms, it is likely that unidentified tick-

host life cycles have evolved, given the endemic wildlife (including tick fauna) present

(Long, 2017; Beati and Klompen, 2019). It is possible that these cycles may contribute to

zoonotic illness when humans encroach these sylvatic ecologies and become exposed to

native ticks. Indeed, it is well documented that in endemic areas in Europe and North

America, the causative agent of Lyme borreliosis is readily identified in wildlife (such as

white-tailed deer, *Odocoileus virginianus*, and white-footed mice, *Peromyscus leucopus*) as

well as their natural tick species (Bosler et al., 1984). The present study therefore is one of

the first to explore the concept that Australian wildlife ticks should be a promising source to

identify exotic and novel TBPs, and we focussed our search towards taxa known to be

associated with TBDs overseas ('taxa of interest'), namely *Anaplasma*, *Bartonella*, *Borrelia*,

Coxiella, *Ehrlichia*, *Francisella*, *Midichloria*, *Neoehrlichia*, *Rickettsia* and *Rickettsiella*.

A new genetic variant of *Anaplasma bovis* was recently described from questing *A.*

triguttatum ticks in WA and the nearby Barrow Island (Gofton et al., 2017). In the present

study we report a range expansion of a genetically similar *A. bovis* along the east coast of

Australia from two widespread *Haemaphysalis* species. As suggested by Gofton et al.,

(2017), this finding further supports the hypothesis that it is likely that endemic *A. bovis*

genotypes exist in sylvatic cycles within native Australian ticks and wildlife fauna. Further

research on the phylogenetic position of these *A. bovis* sequences is needed to understand

their likely evolutionary history and relatedness to *A. bovis* genotype Y11 identified in

Western Australia. While three other species of *Anaplasma* (*A. marginale*, *A. centrale* and *A.*

platys) have been introduced to Australia (Rogers and Shiels, 1979; Callow, 1984; Angus,

1996), the present study did not identify any of these species in ticks from wildlife. This is

likely due to the absence of the cattle tick (*R.* (*B.*) *australis*) and the brown dog tick (*R.*

sanguineus) specimens from wildlife hosts in the present study. A species of *Anaplasma*

identified exclusively in *B. concolor* from echidnas sheds more light on the diverse range of

microbes that have been described from this specialist tick (Loh, 2018).

Although the presence of a *Bartonella* sp. was identified by only four sequences, its absence

in control samples means that this likely represents a true finding in female *I. fecialis*, which

were parasitising different individuals of the same host (quenda) in south-west WA. Recent

studies have shown that Australian marsupials and native rodents harbour a range of distinct

Bartonella species; '*Ca*. Bartonella antechini' has been identified in ticks (*I. antechini*) and

fleas (*Acanthopsylla jordani*) from mardo (*Antechinus flavipes*) in south-west WA

(Kaewmongkol et al., 2011c), '*Ca*. Bartonella woyliei' in woylie ticks (*I. australiensis*) and

fleas (*Pygiopsylla hilli*) in south-west WA, and '*Ca*. Bartonella bandicootii' in fleas

(*Pygiopsylla tunneyi*) from western barred bandicoots (*Perameles bougainville*) on Bernier

and Dorre Island (Kaewmongkol et al., 2011a). Reports of *Bartonella* spp. occurring outside

south-west WA include *Bartonella australis* ex eastern grey kangaroos (*Macropus giganteus)*

(Fournier et al., 2007). Molecular detection of *Bartonella* DNA from Australian ticks has also

been reported in ticks (*I. tasmani*) parasitising koalas (*Phascolarctos cinereus*) from Philip

Island, VIC (Vilcins et al., 2009b). Additional studies on *Bartonella* in Australia wildlife

include those by Gundi et al. (2009), Kaewmongkol et al. (2011a) and Dybing et al. (2016).

Due to the limited size and region of the 16S amplicon generated in the present study, there

were insufficient relevant reference sequences available from other Australian *Bartonella*

species for comparison. A study by Kaewmongkol et al. (2011b) into flea-derived *Bartonella*

from native and introduced Australian species suggests co-evolution of marsupial hosts, their

fleas and the *Bartonella* species.

'Ca. Borrelia tachyglossi' was identified in 8/20 B. concolor ticks from echidnas in QLD and

NSW, an anticipated finding given previous research into this organism (Loh et al., 2016).

Importantly however, the present study provides the first evidence of 'Ca. B. tachyglossi'

sequences from a female H. humerosa tick parasitising a northern brown bandicoot from

QLD. In addition, no Bothriocroton ticks were present in the extraction batch of the positive

H. humerosa sample, and there was no evidence of Borrelia sequences from any controls.

Despite the wide geographical range of H. humerosa ticks, the restricted finding of 'Ca. B.

tachyglossi' from QLD supports Loh et al. (2016), suggesting a restricted geographical

distribution along the east coast of Australia.

In tick microbiome studies overseas, Coxiella spp. are commonly identified (Khoo et al.,

2016; Machado-Ferreira et al., 2016). Additionally, recent studies on Australian ticks have

also identified Coxiella spp. in A. triguttatum (Cooper et al., 2013; Gofton et al., 2015b), B.

auruginans (Vilcins et al., 2009c), I. holocyclus (Cooper et al., 2013) and R. sanguineus

(Oskam et al., 2017). Interestingly the present study did not identify the widespread presence

of Coxiella spp. with only two unique OTUs identified from B. concolor and H. longicornis,

both of which appear to be host specific to those tick species, and the causative agent of Q-

Fever (C. burnetii) was not identified from ticks in the present study.

Ehrlichia sequences ('Ca. E. ornithorhynchi') from platypus ticks, I. ornithorhynchi, were

recently described (Gofton et al., 2018). In the present study, 'Ca. E. ornithorhynchi' was

exclusively observed within I. ornithorhynchi, suggesting this genetically distinct Ehrlichia

species has a unique and host-specific relationship with the platypus. During this study, a

potentially novel species of Ehrlichia was identified in an I. fecialis female tick from a

quenda in WA. BLAST results show it is a close relative to an Ehrlichia sp. detected in

Haemaphysalis ticks from Japan (Inokuma et al., 2004). No sequences from the recently

described '*Ca*. E. occidentalis' (Gofton et al., 2017) were identified in the present study,

however it is noted *A. triguttatum* were only represented by six samples.

Francisella-like endosymbionts have been widely reported in ticks overseas, such as

Dermacentor spp. (Scoles, 2005), *Dermacentor occidentalis* (Gurfield et al., 2017),

Haemaphysalis longicornis (Wang et al., 2018) and *Hyalomma rufipes* (Szigeti et al., 2014),

and *Francisella* sequences have previously been identified in *A. fimbriatum* ticks from

reptiles in the Northern Territory of Australia (Vilcins et al., 2009a). In the present study, a

Francisella-like endosymbiont was identified in 100% of *A. triguttatum* ticks, and in a high

proportion. The high prevalence and relative abundance of the *Francisella*-like organisms in

A. triguttatum may be of medical relevance with respect to this common human-biting tick,

particularly given the recent report of *Francisella* bacteraemia in WA (Aravena-Román et al.,

2015). Importantly, as noted previously for *I. holocyclus* ticks, organisms in lower abundance

may be masked by abundant endosymbionts, unless samples are sequenced deeply and/or a

blocking primer is used (Gofton et al., 2015a). Therefore, future bacterial profiling studies of

A. triguttatum should take these factors into consideration.

While '*Ca*. M. mitochondrii' has been documented in *I. holocyclus,* the present study

provides the first report of '*Ca*. M. mitochondrii' within additional species of native

Australian ticks (*I. fecialis, H. bancrofti* and *H. humerosa*). Despite this expansion among

tick species, identification of '*Ca*. M. mitochondrii' remains confined to NSW and QLD as

previously reported (Gofton et al., 2015a). '*Ca*. M. mitochondrii' has been detected overseas

in laboratory models (Cafiso et al., 2019), wildlife (Serra et al., 2018), and humans

(Mariconti et al., 2012). Despite the use of a CMm blocking primer by Gofton et al. (2015a)

in the library preparation of *I. holocyclus*, the bacteria was still identified in the majority of

ticks (21/36); however its abundance was greatly reduced to an average of 26.1% of reads per

sample in comparison to previous research demonstrating a relative abundance of 98.2%

(Gofton et al., 2015a).

A recent addition to the Anaplasmataceae family, '*Ca*. Neoehrlichia mikurensis' was first

isolated from wild rats and their ticks (*Ixodes ovatus*) in Japan (Kawahara et al., 2004). In

Europe, the hedgehog (*Erinaceus* spp.), a common peri-urban dweller, has been shown to

play an important role in the life cycle of '*Ca*. N. mikurensis' (Földvári et al., 2014; Jahfari et

al., 2017). A high proportion of bandicoot ticks were infected with *Neoehrlichia* spp. in this

study, suggesting that these marsupials may play an important role in the life cycle of this

bacterium on the Australian continent. Bandicoots also frequently inhabit gardens and have

relatively close contact with humans (Carthey and Banks, 2012). Closer study of these

marsupials may provide vital information about the life cycle of these microbes and the

potential risk for human infection.

Rickettsia spp. are among the only currently recognised TBPs affecting people in Australia,

however, as previously outlined, the conserved nature of the 16S gene in *Rickettsia* precludes

rigorous species delimitation from the relatively short sequences generated in this dataset. In

addition to the known human pathogens (*R. australis* and *R. honei*) (Graves and Stenos,

2017), the presence of *Rickettsia* has been demonstrated from a variety of Australian tick

species. *Rickettsia gravesii* has been described from *A. triguttatum* (Li et al., 2010; Abdad et

al., 2017) and *Rickettsia* sequences have been previously identified in *I. tasmani* ex

Tasmanian devils (Vilcins et al., 2009c) and *Amblyomma fimbriatum* from reptiles (Vilcins et

al., 2009a).

The diversity and significance of the *Rickettsiella* genus remains largely unknown globally.

Phylogenetic analysis of sequence data places the genus within the Coxiellaceae family

(Fournier and Raoult, 2005; Leclerque and Kleespies, 2012). While the presence of these

organisms has been well documented in tick species around the world and in some cases, they

have been identified as pathogenic to arthropods (Kurtti et al., 2002; Leclerque et al., 2012),

disease causation within the vertebrate hosts remains unknown at the present time. A study

by Vilcins et al. (2009c) identified *Rickettsiella* in *I. tasmani* ex koalas from Phillip Island,

just off the coast of south eastern Australia. A genetically similar sequence was identified in

the present study (OTU3) from *I. tasmani* exclusively in Tasmania.

While this is the first study to characterise the bacterial communities within native Australian

wildlife ticks, it is apparent that much research is still required in order to better understand

the tick-associated microbial life cycles and their ecologies, let alone elucidating their

potential for transmission and pathogenicity in vertebrate hosts, including humans.

Furthermore, whilst cost-effective, ticks were collected opportunistically resulting in an

inherent geographical bias, confining sampling largely to urban areas along the east coast of

the Australian continent. Nevertheless, this area is also where most humans receive tick bites,

although reliable data about this is also lacking. In addition, the present study suggests that

'taxa of interest' are largely restricted to a combination of geographical location and tick

species. For example, '*Ca*. Midichloria' was identified in *H. humerosa* and *I. fecialis* from

NSW only and was absent in samples from NT for *H. humerosa* and WA and TAS for *I.*

fecialis.

The analysis of ticks removed from wildlife hosts comes with the inherent complication of a

blood meal. The host blood meal has been recently shown to influence both the tick

microbiome composition and the presence of pathogens (Landesman et al., 2019; Swei and

Kwan, 2017). Further targeted studies are needed to assess the source of microbes with

respect to a tick or host origin (Irwin et al., 2018). In addition, limitations in the current

sequencing technologies mean that high-throughput methods favour short amplicons which

may not be able to accurately discriminate bacterial species, as is the case with *Rickettsia* in

particular (Gihring et al., 2012). Furthermore, important caveats to consider in metabarcoding

microbial diversity analysis include PCR efficiency (including primers used and length of

target amplicon), variations in 16S gene copy number (Ahn et al., 2012), sequencing depth,

machine cross-talk, bioinformatic analysis and taxonomic assignment. While there have been

some recent attempts to correct these biases (Kembel et al., 2012; Rosselli et al., 2016), their

application to uncharacterised metagenomic samples remains limited. While 16S

metabarcoding continues to be vital in bacterial biodiversity discovery (Carpi et al., 2011;

Tessler et al., 2017), future molecular studies incorporating metagenomics and

metatranscriptomics, to detect actively expressed genes (Cabezas-Cruz et al., 2018; Greay et

al., 2018), will be useful in characterising the full suite of micro-organisms present in

Australian ticks. A multi-disciplinary approach incorporating cell culture, in vitro tick studies

and morphological techniques (e.g. fluorescence microscopy) will also be needed to assess

the potential for transmissibility and pathogenicity of these novel tick-associated organisms.

5. Conclusions

With over 75% of emerging human infectious pathogens originating from wildlife (King,

2014), surveillance methods that target these species are important for investigations of

emerging and exotic infectious disease. Research into the interactions of wildlife hosts, ticks

and pathogens in Europe and North America continues to highlight the complexity of these

dynamic systems (Ostfeld et al., 2018; Tomassone et al., 2018). Results from the present

study build on recent research into Australian tick-associated microbes, further highlighting

the diversity of organisms present that appear related, yet distinct from their overseas

counterparts. With the evolutionary history of Australia's unique tick species (Beati and

Klompen, 2019) and wildlife fauna, it is likely that reports of their tick-associated microbes

(potential pathogens) will continue to reveal taxonomic differences from those described in

the northern hemisphere. As such, future research on emerging tick-borne zoonoses should

include methods that are able to detect novel micro-organisms in humans and reservoirs and

include a variety of sample types such as blood, tissue (e.g. skin, spleen etc.) and tick from

both the host and environment (questing). As anthropogenic changes to the environment

continue to grow in Australia, a greater emphasis on wildlife disease surveillance is critical to

ensure the early detection of potential infectious diseases affecting humans, livestock,

companion animals and wildlife (Woods et al., 2019). Advances in pathogen detection and

characterisation are greatly enhanced by collaboration; the authors advocate for continued

multidisciplinary efforts between health professionals, researchers, land managers and local

communities.

Supporting Information

Data availability

Associated metadata output from bioinformatic analysis and subsequent data visualisation is

available on FigShare repository https://doi.org/10.6084/m9.figshare.c.4608803.v1. Next-

generation sequencing data can be accessed from NCBI Sequence Read Archive under

BioProject: PRJNA559059 (BioSample accession numbers: SAMN12512474 –

SAMN12512711).

Acknowledgements

The authors would like to thank collaborating research teams, veterinary clinics, wildlife

centres and members of the public for contributing samples to this study. Supplementary File

S1 contains complete list of collectors and institutions. This study was part-funded by the

Australian Research Council (LP160100200) and Bayer HealthCare (Germany) and Bayer

Australia. S.E. received funding from a Murdoch University honours scholarship. S-M.L. was

support by a Murdoch International Postgraduate scholarship. The authors also acknowledge

the Pawsey Supercomputing Centre and the Western Australia State Agriculture

Biotechnology Centre for use of its facilities. We thank Dr. Alexander Gofton and Ms.

Telleasha Greay for advice on next-generation sequencing methods. The funders had no role

in the experimental design, data collection and analyses.

Compliance with ethical standards

This study was conducted under the compliance of the Australian Code for the Responsibility

Conduct of Research (2007) and Australian Code for the Care and Use of Animals for

Scientific Purposes, 2013. Tick collection was carried out opportunistically with the approval

from the Murdoch University Animal Ethics Committee.

References

Abdad, M.Y., Abou Abdallah, R., Karkouri, K., Beye, M., Stenos, J., Owen, H., Unsworth, N., Robertson, I., Blacksell, S.D., Nguyen, T.T., Nappez, C., Raoult, D., Fenwick, S., Fournier, P.E., 2017. *Rickettsia gravesii* sp. nov.: a novel spotted fever group rickettsia in Western Australian *Amblyomma triguttatum triguttatum*. Int. J. Syst. Evol. Microbiol. 67, 3156–3161.

Ahantarig, A., Trinachartvanit, W., Baimai, V., Grubhoffer, L., 2013. Hard ticks and their bacterial endosymbionts (or would be pathogens). Folia Microbiol. (Praha) 58, 419–28.

Ahn, J.H., Kim, B.Y., Song, J., Weon, H.Y., 2012. Effects of PCR cycle number and DNA

polymerase type on the 16S rRNA gene pyrosequencing analysis of bacterial communities. J. Microbiol. 50, 1071–1074.

Andreotti, R., Perez deLeon, A.A., Dowd, S.E., Guerrero, F.D., Bendele, K.G., Scoles, G.A., 2011. Assessment of bacterial diversity in the cattle tick *Rhipicephalus* (*Boophilus*) *microplus* through tag-encoded pyrosequencing. BMC Microbiol. 11, 6.

Angus, B.M., 1996. The history of the cattle tick *Boophilus microplus* in Australia and achievement in its control. Int. J. Parasitol. 26, 1341–1355.

Aravena-Román, M., Merritt, A., Inglis, T.J., 2015. First case of *Francisella* bacteraemia in Western Australia. New Microbes New Infect. 8, 75–77.

Barker, S.C., Walker, A.R., 2014. Ticks of Australia. The species that infest domestic animals and humans. Zootaxa 3816, 1-144

Beati, L., Klompen, H., 2019. Phylogeography of Ticks (Acari: Ixodida). Annu. Rev. Entomol. 64, 379–397.

Bokulich, N.A., Kaehler, B.D., Rideout, B.A., Dillon, M., Bolyen, E., Knight, R., Huttley, G.A., Caporaso, J.G., 2018. Optimizing taxonomic classification of marker gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. Microbiome 6, 90.

Bonnet, S., Michelet, L., Moutailler, S., Cheval, J., Hebert, C., Vayssier-Taussat, M., Eloit, M., 2014. Identification of parasitic communities within European ticks using next-generation sequencing. PLoS Negl. Trop. Dis. 8, e2753.

Bonnet, S.I., Binetruy, F., Hernández-Jarguín, A.M., Duron, O., 2017. The tick microbiome: why non-pathogenic microorganisms matter in tick biology and pathogen transmission. Front. Cell. Infec. Microbiol. 7, 236.

Bosler, E.M., Ormiston, B.G., Coleman, J.L., Hanrahan, J.P., Benach, J.L., 1984. Prevalence of the Lyme Disease spirochete in populations of white-tailed deer and white-footed Mice. Yale J. Biol. Med. 57, 651–659.

Budachetri, K., Gaillard, D., Williams, J., Mukherjee, N., Karim, S., 2016. A snapshot of the microbiome of *Amblyomma tuberculatum* ticks infesting the gopher tortoise, an endangered species. Ticks Tick Borne Dis. 7, 1225-1229.

Burnard, D., Shao, R., 2019. Mitochondrial genome analysis reveals intraspecific variation within Australian hard tick species. Ticks Tick Borne Dis. 10, 677–681.

Cabezas-Cruz, A., Vayssier-Taussat, M., Greub G., 2018. Tick-borne pathogen detection: what's new? Microbes Infect 20, 441-444.

Cafiso, A., Sassera, D., Romeo, C., Serra, V., Hervet, C., Bandi, C., Plantard, O., Bazzocchi, C., 2019. *Midichloria mitochondrii*, endosymbiont of *Ixodes ricinus*: evidence for the transmission to the vertebrate host during the tick blood meal. Ticks Tick Borne Dis. 10, 5–12.

Callow, L.L., 1984. Animal Health in Australia. Volume 5: Protozoal and Rickettsial Diseases. Australian Government Publishing Services, Canberra, Australia.

Carthey, A.J.R., Banks, P.J., 2011. When does an alien become a native species? A vulnerable native mammal recognizes and responds to its long-term alien predator. PLoS One 7, e31804.

Carpi, G., Cagnacci, F., Wittekindt, N.E., Zhao, F., Qi, J., Tomsho, L.P., Drautz, D.I., Rizzoli, A., Schuster, S.C., 2011. Metagenomic profile of the bacterial communities associated with *Ixodes ricinus* ticks. PLoS One 6, e25604.

Chalada, M.J., Stenos, J., Bradbury, R.S., 2016. Is there a Lyme-like disease in Australia? Summary of findings to date. One Health 2, 42–54.

Chalada, M.J., Stenos, J., Vincent, G., Barker, D., Bradbury, R.S., 2018. A molecular survey of tick-borne pathogens from ticks collected in central Queensland, Australia. Vector Borne Zoonotic Dis 18, 151–163.

Cooper, A., Stephens, J., Ketheesan, N., Govan, B., 2013. Detection of *Coxiella burnetii* DNA in Wildlife and Ticks in Northern Queensland, Australia. Vector Borne Zoonotic Dis. 13, 12–16.

de la Fuente, J., Antunes, S., Bonnet, S., Cabezas-Cruz, A., Domingos, A.G., Estrada-Peña, A., Johnson, N., Kocan, K.M., Mansfield, K.L., Nijhof, A.M., Papa, A., Rudenko, N., Villar, M., Alberdi, P., Torina, A., Ayllon, N., Vancova, M., Golovchenko, M., Grubhoffer, L., Caracappa, S., Fooks, A.R., Gortazar, C., Rego, R.O.M., 2017. Tick-pathogen interactions and vector competence: Identification of molecular drivers for tick-borne diseases. Front. Cell. Infect. Microbiol. 7, 114.

DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber, T., Dalevi, D., Hu, P., Andersen, G.L., 2006. Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. Appl. Environ. Microbiol. 72, 5069–5072.

Dobson, A.D.M., Finnie, T.J.R., Randolph, S.E., 2011. A modified matrix model to describe the seasonal population ecology of the European tick *Ixodes ricinus*. J. Appl. Ecol. 48, 1017–1028.

Dong, X., Kleiner, M., Sharp, C.E., Thorson, E., Li, C., Liu, D., Strous, M., 2017. Fast and simple analysis of MiSeq amplicon sequencing data with MetaAmp. Front. Microbiol. 8, 1461.

Dybing, N.A., Jacobson, C., Irwin, P., Algar, D., Adams, P.J., 2016. *Bartonella* species identified in rodent and feline hosts from island and mainland Western Australia. Vector Borne Zoonotic Dis. 16, 238–44.

Edgar, R.C., 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. Nat. Methods 10, 996–968.

Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics 26, 2640–2461.

Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res. 32, 1792–1797.

Edgar, R.C., Flyvbjerg, H., 2015. Error filtering, pair assembly and error correction for next-generation sequencing reads. Bioinformatics 31, 3476–82.

Estrada-Peña, A., Gray, J., Kahl, O., Lane, R.S., Nijhof, A.M., 2013. Research on the ecology of ticks and tick-borne pathogens - methodological principles and caveats. Front. Cell. Infect. Microbiol. 3, 29.

Ferrell, A., Brinkerhoff, R., 2018. Using landscape analysis to test hypotheses about drivers of tick abundance and infection prevalence with *Borrelia burgdorferi*. Int. J. Environ. Res. Public Health 15, 737.

Földvári, G., Jahfari, S., Rigó, K., Jablonszky, M., Szekeres, S., Majoros, G., Tóth, M., Molnár, V., Coipan, E.C., Sprong, H., 2014. *Candidatus* Neoehrlichia mikurensis and *Anaplasma phagocytophilum* in Urban Hedgehogs. Emerg. Infect. Dis. 20, 496–498.

Fournier, P., Taylor, C., Rolain, J.M., Barrassi, L., Smith, G., Raoult, D., 2007. *Bartonella australis* sp. nov. from kangaroos, Australia. Emerg. Infect. Dis. 13, 1961–1963.

Fournier, P.-E., Raoult, D., 2005. Genus II. Rickettsiella Philip 1956, in: Garrity, G.M., Brenner, D.J., Krieg, N.R., Staley, J.T. (Eds.), Bergey's Manual of Systematic Bacteriology, Part B. Springer, New York, pp. 241–247.

Fryxell, R.T.T., DeBruyn, J.M., 2016. The microbiome of *Ehrlichia*-infected and uninfected lone star ticks (*Amblyomma americanum*). PLoS One 11, e0146651.

Gihring, T.M., Green, S.J., Schadt, C.W., 2012. Massively parallel rRNA gene sequencing exacerbates the potential for biased community diversity comparisons due to variable library sizes. Environ. Microbiol. 14, 285–290.

Gofton, A.W., Doggett, S.L., Ratchford, A., Oskam, C.L., Paparini, A., Ryan, U.M., Irwin,

P.J., 2015b. Bacterial profiling reveals novel "Ca. Neoehrlichia", *Ehrlichia*, and *Anaplasma* species in Australian human-biting ticks. PLoS One 10, e0145449.

Gofton, A.W., Loh, S.M., Barbosa, A.D., Paparini, A., Gillett, A., Macgregor, J., Oskam, C.L., Ryan, U.M., Irwin, P.J., 2018. A novel *Ehrlichia* species in blood and *Ixodes ornithorhynchi* ticks from platypuses (*Ornithorhynchus anatinus*) in Queensland and Tasmania, Australia. Ticks Tick Borne Dis. 9, 435–442.

Gofton, A.W., Oskam, C.L., Lo, N., Beninati, T., Wei, H., McCarl, V., Murray, D.C., Paparini, A., Greay, T.L., Holmes, A.J., Bunce, M., Ryan, U.M., Irwin, P.J., 2015a. Inhibition of the endosymbiont "*Candidatus* Midichloria mitochondrii" during 16S rRNA gene profiling reveals potential pathogens in Ixodes ticks from Australia. Parasit. Vectors 8, 345.

Gofton, A.W., Waudby, H.P., Petit, S., Greay, T.L., Ryan, U.M., Irwin, P.J., 2017. Detection and phylogenetic characterisation of novel *Anaplasma* and *Ehrlichia* species in *Amblyomma triguttatum* subsp. from four allopatric populations in Australia. Ticks Tick Borne Dis. 8, 749–756.

Graves, S.R., Stenos, J., 2017. Tick-borne infectious diseases in Australia. Med. J. Aust. 206, 320–324.

Greay, T.L, Gofton, A.W., Paparini, A., Ryan, U.M., Oskam, C.L., Irwin, P.J., 2018. Recent insights into the tick microbiome gained through next-generation sequencing. Parasit Vectors 11, 12.

Guglielmone, A.A., Robbins, R.G., Apanaskevich, D.A., Petney, T.N., Estrada-Peña, A., Horak, I.G., 2014. The Hard Ticks of the World: Acari: Ixodida: Ixodidae. Springer, Dordrecht.

Gundi, V.A., Taylor, C., Raoult, D., La Scola, B., 2009. *Bartonella rattaustraliani* sp. nov., *Bartonella queenslandensis* sp. nov. and *Bartonella coopersplainsensis* sp. nov., identified in Australian rats. Int. J. Syst. Evol. Microbiol. 59, 2956–2961.

Gurfield, N., Grewal, S., Cua, L.S., Torres, P.J., Kelley, S.T., 2017. Endosymbiont interference and microbial diversity of the Pacific coast tick, *Dermacentor occidentalis*, in San Diego County, California. PeerJ 5, e3202.

Inokuma, H., Beppu, T., Okuda, M., Shimada, Y., Sakata, Y., 2004. Detection of Ehrlichial DNA in *Haemaphysalis* ticks recovered from dogs in Japan that is slosely related to a novel *Ehrlichia* sp. found in cattle yicks from Tibet, Thailand, and Africa. J. Clin. Microbiol. 42, 1353–1355.

Irwin, P.J., Egan, S., Greay, T., Oskam, C.L., 2018. Bacterial tick-associated infections in Australia: current studies and future direction. Microbiol. Aust. 39, 200–202.

Irwin, P.J., Robertson, I.D., Westman, M.E., Perkins, M. Straubinger, R.K., 2017. Searching for Lyme borreliosis in Australia: results of a canine sentinel study. Parasit. Vectors 10, 114.

Jahfari, S., Ruyts, S.C., Frazer-Mendelewska, E., Jaarsma, R., Verheyen, K., Sprong, H., 2017. Melting pot of tick-borne zoonoses: the European hedgehog contributes to the maintenance of various tick-borne diseases in natural cycles urban and suburban areas. Parasit. Vectors 10, 134.

Jones, K.E., Patel, N.G., Levy, M.A., Storeygard, A., Balk, D., Gittleman, J.L., Daszak, P., 2008. Global trends in emerging infectious diseases. Nature 451, 990–993.

Kaewmongkol, G., Kaewmongkol, S., Burmej, H., Bennett, M.D., Fleming, P.A., Adams, P.J., Wayne, A.F., Ryan, U., Irwin, P.J., Fenwick, S.G., 2011a. Diversity of *Bartonella* species detected in arthropod vectors from animals in Australia. Comp. Immunol. Microbiol. Infect. Dis. 34, 411–417.

Kaewmongkol, G., Kaewmongkol, S., McInnes, L.M., Burmej, H., Bennett, M.D., Adams, P.J., Ryan, U., Irwin, P.J., Fenwick, S.G., 2011b. Genetic characterization of flea-

derived *Bartonella* species from native animals in Australia suggests host-parasite co-evolution. Infect. Genet. Evol. 11, 1868–1872.

Kaewmongkol, G., Kaewmongkol, S., Owen, H., Fleming, P.A., Adams, P.J., Ryan, U., Irwin, P.J., Fenwick, S.G., 2011c. Candidatus Bartonella antechini: a novel *Bartonella* species detected in fleas and ticks from the yellow-footed antechinus (*Antechinus flavipes*), an Australian marsupial. Vet. Microbiol. 149, 517–521.

Kawahara, M., Rikihisa, Y., Isogai, E., Takahashi, M., Misumi, H., Suto, C., Shibata, S., Zhang, C., Tsuji, M., 2004. Ultrastructure and phylogenetic analysis of "*Candidatus* Neoehrlichia mikurensis" in the family Anaplasmataceae, isolated from wild rats and found in *Ixodes ovatus* ticks. Int. J. Syst. Evol. Microbiol. 54, 1837–1843.

Kembel, S.W., Wu, M., Eisen, J.A., Green, J.L., 2012. Incorporating 16S gene copy number information improves estimates of microbial diversity and abundance. PLoS Comput. Biol. 8, e1002743.

Khoo, J.J., Lim, F.S., Chen, F., Phoon, W.H., Khor, C.S., Pike, B.L., Chang, L.Y., AbuBakar, S., 2016. *Coxiella* detection in ticks from wildlife and livestock in Malaysia. Vector Borne Zoonotic Dis. 16, 744–751.

King, L.J., 2014. Combating the triple threat: the need for a one health approach, in: Atlas, R.M., Maloy, S. (Eds.), One Health: People, Animals and the Environment. American Society for Microbiology, Washington, pp. 3–16.

Kumar, S., Stecher, G., Tamura, K., 2016. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. Mol. Biol. Evol. 33, 1870–1874.

Kurilshikov, A., Livanova, N.N., Fomenko, N.V., Tupikin, A.E., Rar, V.A., Kabilov, M.R., Livanov, S.G., Tikunova, N.V., 2015. Comparative metagenomic profiling of symbiotic bacterial communities associated with *Ixodes persulcatus*, *Ixodes pavlovskyi* and *Dermacentor reticulatus* ticks. PLoS One 10, e0131413.

Kurtti, T.J., Palmer, A.T., Oliver, J.H., 2002. *Rickettsiella*-like bacteria in *Ixodes woodi* (Acari: Ixodidae). J. Med. Entomol. 39, 534–540.

Kwan, J.Y., Griggs, R., Chicana, B., Miller, C., Swei, A., 2017. Vertical vs. horizontal transmission of the microbiome in a key disease vector, *Ixodes pacificus*. Mol. Ecol. 26, 6578–6589.

Landesman, W.J., Mulder, K., Allan, B.F., Bashor, L.A., Keesing, F., LoGiudice, K., Ostfeld, R.S., 2019. Potential effects of blood meal host on bacterial community composition in *Ixodes scapularis* nymphs. Ticks Tick Borne Dis. 10, 523-527.

Leclerque, A., Kleespies, R.G., 2012. A *Rickettsiella* bacterium from the hard tick, *Ixodes woodi*: molecular taxonomy combining multilocus sequence typing (MLST) with significance testing. PLoS One 7, e38062.

Leclerque, A., Kleespies, R.G., Schuster, C., Richards, N.K., Marshall, S.D.G., Jackson, T.A., 2012. Multilocus sequence analysis (MLSA) of '*Rickettsiella costelytrae*' and '*Rickettsiella pyronotae*', intracellular bacterial entomopathogens from New Zealand. J. Appl. Microbiol. 113(5), 1228–1237.

Li, A.Y., Adams, P.J., Abdad, M.Y., Fenwick, S.G., 2010. High prevalence of *Rickettsia gravesii* sp. nov. in *Amblyomma triguttatum* collected from feral pigs. Vet. Microbiol. 146, 59–62.

LoGiudice, K., Ostfeld, R.S., Schmidt, K.A., Keesing, F., 2003. The ecology of infectious disease: effects of host diversity and community composition on Lyme disease risk. Proc. Natl. Acad. Sci. U.S.A. 100, 567–571.

Loh, S.M., 2018. Identification and characterisation of microorganisms in Australian wildlife tick (PhD thesis). Murdoch University, Perth, Western Australia.

Loh, S.M., Gofton, A.W., Lo, N., Gillett, A., Ryan, U.M., Irwin, P.J., Oskam, C.L., 2016. Novel *Borrelia* species detected in echidna ticks, *Bothriocroton concolor*, in

Australia. Parasites & Vectors 9, 339.

Long, J.A., 2017 Why Australasian vertebrate animals are so unique – a palaeontological perspective. Gen. Comp. Endocr. 244, 2-10.

Lydecker, H.W., Banks, P.B., Hochuli, D.F., 2019. Counting ticks (Acari: Ixodida) on hosts is complex: a review and comparison of methods. J. Med. Entomol. 56, 1527-1533.

Machado-Ferreira, E., Vizzoni, V.F., Balsemao-Pires, E., Moerbeck, L., Gazeta, G.S., Piesman, J., Voloch, C.M., Soares, C.A., 2016. *Coxiella* symbionts are widespread into hard ticks. Parasitol. Res. 115(12), 4691–4699.

Mans B.J., Featherston, J., Kvas, M., Pillay, K., de Klerk D.G., Pienaar, R., de Castro, M. H., Schwan, T. G., Lopez, J. E., Teel, P., Pérez de León, A. A., Sonenshine, D. E., Egekwu, N. I., Bakkes, D. K., Kanduma, E. G., Nyangiwe, N., Bouattour, A., Latif, A. A., 2019. Argasid and ixodid systematics: Implications for soft tick evolution and systematics, with a new argasid species list. Ticks Tick Borne Dis. 10, 219-240.

Mariconti, M., Epis, S., Gaibani, P., Dalla Valle, C., Sassera, D., Tomao, P., Fabbi, M., Castelli, F., Marone, P., Sambri, V., Bazzocchi, C., Bandi, C., 2012. Humans parasitized by the hard tick *Ixodes ricinus* are seropositive to *Midichloria mitochondrii*: is *Midichloria* a novel pathogen, or just a marker of tick bite? Pathog. Glob. Health 106, 391–396.

McMurdie, P.J., Holmes, S., 2013. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. PLoS One 8(4), e61217.

McMurdie, P.J., Holmes, S., 2014. Waste not, want not: why rarefying microbiome data is inadmissible. PLoS Comput. Biol. 10, e1003531.

Mediannikov, O., Fenollar, F., 2014. Looking in ticks for human bacterial pathogens. Microb. Pathog. 77, 142–8.

Merrill, M.M., Boughton, R.K., Lord, C.C., Sayler, K.A., Wight, B., Anderson, W.M., Wisely, S.M., 2018. Wild pigs as sentinels for hard ticks: A case study from south-central Florida. Int. J. Parasitol. Parasites Wildl. 7, 161–170.

Morens, D.M., Fauci, A.S., 2012. Emerging infectious diseases in 2012: 20 years after the institute of medicine report. MBio 3, e00494-12.

Morgulis, A., Coulouris, G., Raytselis, Y., Madden, T.L., Agarwala, R., Schaffer, A.A., 2008. Database indexing for production MegaBLAST searches. Bioinformatics 25, 1757-1764.

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2019. Vegan: community ecology package. R package version 2.5-5

Oskam, C.L., Gofton, A.W., Greay, T.L., Yang, R., Doggett, S., Ryan, U.M., Irwin, P.J., 2017. Molecular investigation into the presence of a *Coxiella* sp. in *Rhipicephalus sanguineus* ticks in Australia. Vet. Microbiol. 201, 141–145.

Ostfeld, R.S., Brunner, J.L., 2015. Climate change and *Ixodes* tick-borne diseases of humans. Philos. Trans. R. Soc. Lond. B. Biol. Sci. 370, 20140051.

Ostfeld, R.S., Levi, T., Keesing, F., Oggenfuss, K., Canham, C.D., 2018. Tick-borne disease risk in a forest food web. Ecology 99, 1562–1573.

Panetta, J.L., Šíma, R., Calvani, N.E.D., Hajdušek, O., Chandra, S., Panuccio, J., Šlapeta, J., 2017. Reptile-associated *Borrelia* species in the goanna tick (*Bothriocroton undatum*) from Sydney, Australia. Parasit. Vectors 10, 616.

Parola, P., Raoult, D., 2001. Ticks and tickborne bacterial diseases in humans: an emerging infectious threat. Clin. Infect. Dis. 32, 897–928.

Pfäffle, M., Littwin, N., Muders, S.V., Petney, T.N., 2013. The ecology of tick-borne diseases. Int. J. Parasitol. 43, 1059–1077.

Poisot, T., Foster, Z.S.L., Sharpton, T.J., Grünwald, N.J., 2017. Metacoder: An R package for

visualization and manipulation of community taxonomic diversity data. PLOS Comput. Biol. 13, e1005404.

Pollock, J., Glendinning, L., Wisedchanwet, T., Watson, M., 2018. The madness of microbiome: attempting to find consensus "best pratice" for 16S microbiome studies. Appl. Environ. Microbiol. 19, pii: e02627-17

Ponnusamy, L., Gonzalez, A., Treuren, W.V., Welss, S., Parobek, C.M., Jullano, J.J., Knight, R., Roe, R.M., Apperson, C.S., Meshnick, S.R., 2014. Diversity of *Rickettsiales* in the microbiome of the lone star tick, *Amblyomma americanum*. Appl. Environ. Microbiol. 80, 354–359.

Radcliffe, J., Rusjakovski, N., Forkert, J., Jones, M., Stewart, C., 2016. Growing evidence of an emerging tick-borne disease that causes a Lyme-like illness for many Australian patients. Community Affairs References Committee, Canberra, Australia.

Rinker, D.C., Pitts, R.J., Zwiebel, L.J., 2016. Disease vectors in the era of next generation sequencing. Genome Biol. 17, 95.

Roberts, F.S.H., 1970. Australian Ticks. CSIRO, Melbourne.

Rogers, R.J., Shiels, I.A., 1979. Epidemiology and control of anaplasmosis in Australia. J. S. Afr. Vet. Assoc. 50, 363–366.

Rosselli, R., Romoli, O., Vitulo, N., Vezzi, A., Campanaro, S., de Pascale, F., Schiavon, R., Tiarca, M., Poletto, F., Concheri, G., Valle, G., Squartini, A., 2016. Direct 16S rRNA-seq from bacterial communities: a PCR-independent approach to simultaneously assess microbial diversity and functional activity potential of each taxon. Sci. Rep. 6, 32165.

Roux, V., Raoult, D., 1995. Phylogenetic analysis of the genus *Rickettsia* by 16S rDNA sequencing. Res. Microbiol. 146, 385–396.

RStudio Team, 2015. RStudio: Integrated Development for R. RStudio Inc, Boston, MA.

Scoles, G.A., 2004. Phylogenetic analysis of the *Francisella*-like endosymbionts of *Dermacentor* ticks. J. Med. Entomol. 41, 277–286.

Serra, V., Cafiso, A., Formenti, N., Verheyden, H., Plantard, O., Bazzocchi, C., Sassera, D., 2018. Molecular and serological evidence of the presence of *Midichloria mitochondrii* in roe deer (*Capreolus capreolus*) in France. J. Wildl. Dis. 54, 597–600.

Sperling, J.L., Silva-Brandão, K.L., Brandão, M.M., Lloyd, V.K., Dang, S., Davis, C.S., Sperling, F.A.H., Magor, K.E., 2017. Comparison of bacterial 16S rRNA variable regions for microbiome surveys of ticks. Ticks Tick Borne Dis. 8, 453–461.

Špitalská, E., Sparagano, O., Stanko, M., Schwarzová, K., Špitalský, Z., Škultéty, Ľ., Havlíková, S.F., 2018. Diversity of *Coxiella*-like and *Francisella*-like endosymbionts, and *Rickettsia* spp., *Coxiella burnetii* as pathogens in the tick populations of Slovakia, Central Europe. Ticks Tick Borne Dis. 9, 1207–1211.

Stothard, D.R., Fuerst, P.A., 1995. Evolutionary analysis of the spotted fever and thyphus groups of *Rickettsia* using 16S rRNA gene sequences. Syst. Appl. Microbiol. 18, 52–61.

Sumrandee, C., Baimai, V., Trinachartvanit, W., Ahantarig, A., 2016. Molecular detection of *Rickettsia*, *Anaplasma*, *Coxiella* and *Francisella* bacteria in ticks collected from Artiodactyla in Thailand. Ticks Tick Borne Dis. 7, 678-689.

Swei, A., Kwan, J.Y., 2017. Tick microbiome and pathogen acquisition altered by host blood meal. ISME J. 11, 813–816.

Szigeti, A., Kreizinger, Z., Hornok, S., Abichu, G., Gyuranecz, M., 2014. Detection of *Francisella*-like endosymbiont in *Hyalomma rufipes* from Ethiopia. Ticks Tick Borne Dis. 5, 818–820.

Tamura, K., 1992. Estimation of the number of nucleotide substitutions when there are strong transition-transversion and G+C-content biases. Mol. Biol. Evol. 9, 678–687.

Tessler, M., Neumann, J.S., Afshinnekoo, E., Pineda, M., Hersch, R., Velho, L.F.M., Segovia, B.T., Lansac-Toha, F.A., Lemke, M., DeSalle, R., Mason, C.E., Brugler, M.R., 2017. Large-scale differences in microbial biodiversity discovery between 16S amplicon and shotgun sequencing. Sci. Rep. 7, 6589.

Thapa, S., Zhang, Y., Allen, M.S., 2019. Bacterial microbiomes of *Ixodes scapularis* ticks collected from Massachusetts and Texas, USA. BMC Microbiol. 19, 138.

Tomassone, L., Berriatua, E., De Sousa, R., Duscher, G.G., Mihalca, A.D., Silaghi, C., Sprong, H., Zintl, A., 2018. Neglected vector-borne zoonoses in Europe: Into the wild. Vet. Parasitol. 251, 17–26.

Turner, S., Pryer, K.M., Miao, V.P., Palmer, J.D., 1999. Investigating deep phylogenetic relationships among cyanbacteria and plastids by small subunit rRNA sequence analysis. J Eukaryot Microbiol 46, 327–338.

Vayssier-Taussat, M., Kazimirova, M., Hubalek, Z., Hornok, S., Farkas, R., Cosson, J.F., Bonnet, S., Vourch, G., Gasqui, P., Mihalca, A.D., Plantard, O., Silaghi, C., Cutler, S., Rizzoli, A., 2015. Emerging horizons for tick-borne pathogens: from the 'one pathogen-one disease' vision to the pathobiome paradigm. Future Microbiol. 10, 2033–2043.

Vilcins, I.-M., Fournier, P., Old, J.M., Deane, E., 2009a. Evidence for the presence of *Francisella* and spotted fever group *Rickettsia* DNA in the tick *Amblyomma fimbriatum* (Acari: Ixodidae), Northern Territory Australia. J. Med. Entomol. 46, 926–33.

Vilcins, I.M., Kosoy, M., Old, J.M., Deane, E.M., 2009b. *Bartonella*-like DNA detected in *Ixodes tasmani* ticks (Acari: Ixodida) infesting Koalas (*Phascolarctos cinereus*) in Victoria, Australia. Vector Borne Zoonotic Dis. 9, 499–503.

Vilcins, I.M., Old, J.M., Deane, E., 2009c. Molecular detection of *Rickettsia*, *Coxiella* and *Rickettsiella* DNA in three native Australian tick species. Exp. Appl. Acarol. 49, 229–242.

Wang, B., Wan, L., Wang, A., Li, L.M., 2017. An adaptive decorrelation method removes Illumina DNA base-calling errors caused by crosstalk between adjacent clusters. Sci. Rep. 7, 41348.

Wang, Y., Mao, L., Sun, Y., Wang, Z., Zhang, Jiayong, Zhang, Jibo, Peng, Y., Xia, L., 2018. A Novel *Francisella*-Like endosymbiont in *Haemaphysalis longicornis* and *Hyalomma asiaticum*, China. Vector Borne Zoonotic Dis. 18, 669–676.

Woods, R., Reiss, A., Cox-Witton, K., Grillo, T., Peters, A., 2019. The importance of wildlife disease monitoring as part of global surveillance for zoonotic diseases: the role of Australia. Trop. Med. Infect. Dis. 4, 29.

Zhang, X., Yang, Z., Lu, B., Ma, X., Zhang, C., Xu, H., 2014. The composition and transmission of microbiome in hard tick, *Ixodes persulcatus*, during blood meal. Ticks Tick Borne Dis. 5, 864–870.

Zolnik, C.P., Prill, R., Falco, R.C., Daniels, T.J., Kolokotronis, S.O., 2016. Microbiome changes through ontogent of a tick pathogen vector. Mol. Ecol. 25, 4963-4977.

Legends to Figures

Fig. 1. Chord diagram showing tick-wildlife associations recorded in the present study. The

wildlife hosts recorded are represented on the lower half of the plot, and tick species across

the top. Thickness of bar relative to number of ticks. Host records that could not be assigned

to species level are not represented in this plot. A complete list of records is available in

Supplementary File S1. Introduced wildlife species are denotes with an asterisk. Silhouette

wildlife images sourced from phylopic.org.

Fig. 2. Alpha diversity of bacterial communities in ticks parasitising Australian wildlife

measured as; observed number of operational taxonomic units (OTUs), Chao1 index,

Shannon index and Simpson index. Tick species abbreviated to; *A. triguttatum* (*A. tri*), *B.*

aurugians (*B. aur*), *B. concolor* (*B. con*), *H. bancrofti* (*H. ban*), *H. bremneri* (*H. bremneri*),

H. humerosa (*H. hum*), *H. longicornis* (*H. lon*), *I. antechini* (*I. ant*), *I. australiensis* (*I. aus*), *I.*

fecialis (*I. fec*), *I. holocyclus* (*I. hol*), *I. myrmecobii* (*I. myr*), *I. ornithorhynchi* (*I. orn*), *I.*

tasmani (*I. tas*), and *I. trichosuri* (*I. tri*).

Fig 3. Rarefaction plot of 16S rRNA bacterial sequences clustered into 97% operational

taxonomic units (OTUs) from ticks parasitising Australian wildlife.

Fig 4. Non-metric multidimensional scaling (NMDS) plot of operational taxonomic units

(OTUs) based on Bray-Curtis dissimilarity matrix. Taxa filtered to only display OTUs that

were present at least twice in >10% of tick samples.

Fig 5. Stacked barplot of bacterial composition (shown at family level) from Australian ticks

represented as percentage of assigned sequences. Taxa at the family level that represented

<10% of the relative sequences within each tick species were grouped as "Low abundant".

Sequences not able to be accurately assigned to family taxa are displayed as "Unclassified".

Fig 6. Neighbour-joining phylogenetic tree displaying taxa of interest prevalence and

distribution among tick samples. Geographic data relating to tick collection is represented by

state and territory; New South Wales (NSW), Northern Territory (NT), Queensland (QLD),

Tasmania (TAS) and Western Australia (WA). Evolutionary histories were inferred based on

the Tamura 3-parameter model with bootstrap analysis (10,000 replicates) (bootstrap values

>60 are displayed). Tick samples were considered positive for taxa of interest if >100

sequences present. Information on number of sequences, taxonomic identity (as inferred from

NCBI MegaBLAST analysis with nucleotide database), and sequences can be found in

supplementary information (Supplementary File S4 & S5).

Supplementary Files

Supplementary File S1. Sample metadata of tick-wildlife records identified from the present

study.

Supplementary File S2. Sequence information from 16S rRNA bacterial profiling of ticks

on the Illumina MiSeq platform after bioinformatic analysis; (a) number sequences assigned

to 97% operational taxonomic units (OTUs) and (b) number of sequences per sample.

Supplementary File S3. Principal coordinate analysis (PCoA) plot of the weighted UniFrac

distance matrix. Plots show beta diversity of tick bacterial communities, displayed by tick

species and life stage. Operational taxonomic units (OTUs) that represented <100 sequences

in a tick sample were removed.

Supplementary File S4. Number of sequences relating to the taxa of interest in tick samples.

The top hit from NCBI BLAST results showing GenBank accession number, percent identity

and E-value are given, and final evaluation of taxonomic assignment.

Supplementary File S5. Fasta sequences for operational taxonomic units (OTUs) used in the

taxa of interest investigation.