Running title: Bacterial community profiling highlights complex diversity and novel

organisms in wildlife ticks.

Authors: Siobhon L. Egan[1], Siew-May Loh[1], Peter B. Banks[2], Amber Gillett[3], Liisa A.

Ahlstrom[4], Una M. Ryan[1], Peter J. Irwin[1] and Charlotte L. Oskam[1,*]

[1] Vector and Waterborne Pathogens Research Group, College of Science, Health,

Engineering and Education, Murdoch University, Perth, Western Australia, Australia

[2] School of Life and Environmental Sciences, The University of Sydney, Sydney, New South

Wales, Australia

[3] Australia Zoo Wildlife Hospital, Beerwah, Queensland, Australia

[4] Bayer Australia Ltd, Animal Health, Pymble, New South Wales, Australia

[*] Corresponding author

S.L.E siobhon.egan@murdoch.edu.au; https://orcid.org/0000-0003-4395-4069

S-M.L. kim_lsm@yahoo.com

P.B.B. peter.banks@sydney.edu.au; https://orcid.org/0000-0002-4340-6495

A.G. drambergillett@hotmail.com

L.A.A liisa.ahlstrom@bayer.com

U.M.R. una.ryan@murdoch.edu.au; https://orcid.org/0000-0003-2710-9324

P.J.I. p.irwin@murdoch.edu.au; https://orcid.org/0000-0002-0006-8262

C.L.O. c.oskam@murdoch.edu.au; https://orcid.org/0000-0001-8886-2120

Abstract

Ticks (Acari: Ixodida) transmit a greater variety of pathogens than any other blood-feeding

arthropod. While numerous microbes have been identified inhabiting Australian Ixodidae,

some of which are related to globally important tick-borne pathogens, little is known about

the bacterial communities within ticks collected from Australian wildlife. In this study, 1,019

ticks were identified on 221 hosts spanning 27 wildlife species. Next-generation sequencing

was used to amplify the V1-2 hypervariable region of the bacterial 16S rRNA gene from 238

ticks; *Amblyomma triguttatum* (n=6), *Bothriocroton auruginans* (n=11), *Bothriocroton*

concolor (n=20), *Haemaphysalis bancrofti* (n=10), *Haemaphysalis bremneri* (n=4),

Haemaphysalis humerosa (n=13)*, Haemaphysalis longicornis* (n=4), *Ixodes antechini* (n=29),

Ixodes australiensis (n=26), *Ixodes fecialis* (n=13), *Ixodes holocyclus* (n=37), *Ixodes*

*myrmecobii (*n*=1), Ixodes ornithorhynchi* (n=10), *Ixodes tasmani* (n=51) and *Ixodes*

trichosuri (n=3). After bioinformatic analyses, over 14 million assigned bacterial sequences

revealed the presence of newly described bacteria '*Ca*. Borrelia tachyglossi', '*Ca.*

Neoehrlichia australis', '*Ca.* Neoehrlichia arcana' and '*Ca*. Ehrlichia ornithorhynchi'.

Furthermore, three novel Anaplasmataceae species were identified including; a *Neoehrlichia*

sp. in *I. australiensis* and *I. fecialis* ex quenda (*Isoodon fusciventer*) (Western Australia), an

Anaplasma sp. from one *B. concolor* ex echidna (*Tachyglossus aculeatus*) (New South

Wales), and an *Ehrlichia* sp. from a single *I. fecialis* parasitising a quenda (WA). This study

highlights the diversity of bacterial genera harboured within wildlife ticks, which may prove

to be of medical and/or veterinary importance in the future.

Keywords: Microbiome; ticks; Ixodida; wildlife; marsupials; Anaplasmataceae

1. Introduction

Current estimates suggest that approximately 17% of all infectious diseases of humans are

vector-borne (Rinker et al., 2016) and global trends show that vector-borne diseases (VBDs)

are rising at a rapid rate (Jones et al., 2008; Morens and Fauci, 2012). The complex interplay

between pathogen, vector, host(s) and the environment make VBDs particularly challenging

to understand. In addition, factors such as climate change (Ostfeld and Brunner, 2015), land

use (Ferrell and Brinkerhoff, 2018), feral animal populations (Merrill et al., 2018) and the

microclimate within a landscape (Dobson et al., 2011) can further influence the prevalence

and distribution of VBDs.

Ticks (Acari: Ixodida) comprise a group of haematophagous (blood feeding) arthropods with

almost 900 species described globally (Guglielmone et al., 2010). Ticks are known to

transmit various pathogens, however, they also harbour a range of endosymbiont and

commensal species (Špitalská et al., 2018). The epidemiology of recognised tick-borne

diseases (TBDs) in the northern hemisphere demonstrates that wildlife serve as sentinels and

can be used to monitor the presence and distribution of tick-borne pathogens (TBPs).

Importantly, research has shown that some wildlife species act as dilution hosts for certain

TBPs, whereas others may act as amplification hosts (LoGiudice et al., 2003).

The complexity of TBDs means that studies are increasingly shifting away from isolated

species-specific studies toward ecosystem-based, collaborative research (Estrada-Pena et al.,

2013; Pfaffle et al., 2013). Metabarcoding provides an informative molecular platform to

characterise the bacterial diversity in ticks. Worldwide, next-generation sequencing (NGS)-

based analyses have been applied to a range of tick species that are important from medical

and veterinary perspectives, including *Amblyomma americanum* (Ponnusamy et al., 2014),

Ixodes ricinus (Bonnet et al., 2014), and *Rhipicephalus microplus* (Andreotti et al., 2011).

Metabarcoding studies of the bacterial microbiome of Australian ticks have been reported

only recently, with the first bacterial profiling by NGS conducted on human-biting *Ixodes*

holocyclus (Gofton et al., 2015a). These authors identified a highly abundant endosymbiont

'Candidatus Midichloria mitochondrii' (CMm) and after blocking the amplification of this

organism, a greater bacterial diversity was revealed, including a number of novel microbes

(Gofton et al., 2015a,b). Critically, in contrast to many parts of the world where multiple

TBPs have been elucidated within well-studied tick-host-environment ecologies, there is a

relative dearth of such information available for Australia. With this in mind, the aims of our

study were to survey the bacterial communities present in ticks collected from Australian

wildlife and to investigate their genetic relatedness to 'taxa of interest', i.e. tick-associated

pathogenic and endosymbiotic organisms (Parola and Raoult, 2001; Mediannikov and

Fenollar, 2014; Sumrandee et al., 2016). Taxa of interest in the present study were defined as

genera within alphaproteobacteria, gammaproteobacteria and spirochaetes known to be

transmitted by ticks in other parts of the world, specifically; *Anaplasma*, *Bartonella*, *Borrelia*,

Coxiella, *Ehrlichia*, *Francisella*, *Midichloria*, *Neoehrlichia*, *Rickettsia* and *Rickettsiella*.

(Ahantarig et al., 2013; Vayssier-Taussat et al., 2015; Bonnet et al., 2017; de la Fuente et al.,

2017).

2. Materials and methods

2.1 Sample collection and identification

1,019 ticks were sourced opportunistically from wildlife in Australia by veterinarians,

veterinary nurses, wildlife carers, researchers and via submissions from members of the

public, and preserved in 70% ethanol before being shipped to Murdoch University, Western

Australia, for identification. Ticks were identified morphologically to instar and species using

keys and species' descriptions (Roberts, 1970; Barker and Walker, 2014). Details of sample

collection are available in Supplementary File S1.

2.2 DNA Extraction

A sub-sample of ticks (*n*=238) was chosen for DNA extraction and bacterial profiling based

on geography, instar, tick and host species. Prior to DNA extraction, individual ticks were

surface-sterilised in 10% hypochlorite solution, rinsed in 70 % ethanol and DNA-free PBS,

and then air-dried. Total genomic DNA (gDNA) was extracted using the Qiagen DNeasy

Blood and Tissue kit (Qiagen, Germany) following the manufacturer's recommendations

with the following modifications; ticks were placed in a 2 mL safe lock Eppendorf tube with

a 5 mm steel bead, frozen in liquid nitrogen for 1 minute and homogenised by shaking at 40

Hz in a Tissue Lyser LT (Qiagen, Germany). The volume of elution buffer AE was adjusted

to 200 μL for engorged females, 100 μL for unengorged adults and 40 μL for nymphs. A

double elution was carried out to increase gDNA yield for unengorged adults and nymphs.

Sterile and DNA-free equipment and tubes were used for each step. Extraction reagents blank

(EXB) controls were performed alongside tick extractions to assess background bacterial

communities.

2.3 16S amplification and library preparation for NGS sequencing

Amplicons targeting a 250-320 bp product of the V1-2 hypervariable region of the 16S rRNA

(16S) gene were generated using the primer pair 27F-Y (Gofton et al., 2015b) and 338R

(Turner et al., 1999). To inhibit the amplification of abundant '*Ca*. Midichloria mitochondrii'

in *I. holocyclus* ticks, 3-10 μM of the CMm blocking primer was added to the PCR reaction

(Gofton et al., 2015a). Amplicon PCRs were carried in 25 μL reactions each containing 1 X

Buffer (KAPA Biosystems, USA), 1.5 mM MgCl$_2$, 0.4 μM of each forward and reverse

primer with MiSeq adapters, 0.4 mg/mL BSA (Fisher Biotech, Australia), 0.25 mM dNTPs

(Fisher Biotech, Australia), 0.5 U KAPA Taq (KAPA Biosystems, USA), and 2.0 μL of

undiluted genomic DNA. Samples underwent the following thermal cycling conditions;

initial denaturation at 95°C for 5 mins, followed by 35 cycles of denaturation at 95°C for 30

s, annealing at 62°C (CMm blocking primer present) or 55°C (CMm blocking primer absent)

for 30 s, and extension at 72°C for 45 s with a final extension at 72°C for 5 mins. Libraries

were then prepared using the Nextera XT DNA library preparation kit in 25 μL reactions

following manufacturer's recommendations. Libraries were purified using Agencourt®

AMPure® XP PCR purification beads (Beckman Coulter Life Sciences, USA) and pooled in

equimolar amounts. Final libraries were then quantified using the Qubit® 2.0 Fluorometer

(Thermo Fisher, Australia). Libraries were sequenced on an Illumina MiSeq™ using v2

chemistry (2x250 paired end).

Extraction reagent blank (n=12) and PCR no-template controls (NTC; n=7) were included in

all stages of the workflow. All pre- and post-PCR procedures were performed in physically-

separate dedicated laboratories, and sterile protocols were maintained through library

preparation in order to minimise amplicon contamination.

2.4 Bioinformatics and statistical analysis

Raw fastq files were downloaded from the Illumina BaseSpace Sequence Hub for analysis in

a custom pipeline using USEARCH (Edgar, 2010). Raw paired-end sequences were merged

in USEARCH v10, a minimum of 50 nucleotide (nt) overlap and maximum number of

mismatches increased to 15 nt due to long overlap of paired-end sequences. Only sequences

with perfect primer sequences were retained, primer sequences and distal bases were trimmed

using USEARCH v8.0. Sequences were quality filtered in USEARCH v10, allowing a <1%

expected error rate and singletons were discarded (Edgar and Flyvbjerg, 2015). Sequences

were then clustered into operational taxonomic units (OTUs) of 97% similarity using the

UPARSE algorithm (Edgar, 2013) in USEARCH v10. Taxonomy was assigned in QIIME2

v2018.4 using the q2-feature-classifier (Bokulich et al., 2018) with reference to a trained

Greengenes database (DeSantis et al., 2006) (release May 2013) using the primer pair 27F-Y/

338R. Taxonomic assignments were confirmed using NCBI blast (Altschul et al., 1990) on a

random subsample of OTUs and in the case of tick-associated microbes, GenBank accession

numbers and percentage similarity of top hits were recorded for 'taxa of interest' (see

Supplementary File S2). The profile from EXB controls and NTCs was first assessed to

ensure quality of sampling and absence of tick-associated bacteria. The profiles of EXB and

NTC were then removed bioinformatically from associated samples to eliminate potential

contamination and background bacteria.

Data analysis and visualisation was carried out in RStudio (RStudio Team, 2015) using

packages metacoder (Poisot et al., 2017), phyloseq (McMurdie and Holmes, 2013) and vegan

(Oksanen et al., 2019). Alpha diversity of samples was measured using observed OTUs,

Chao1 index, Shannon index and Simpson index. Removal of samples with a low sequencing

depth (<1000 assigned OTUs after data filtering), did not significantly alter alpha diversity

measurements among tick species (data not shown). Rarefaction curves for samples were

calculated to assess sequencing depth based on observed number of OTUs. Principal

coordinate analysis (PCoA) on weighted unifrac dissimilarity measurements was used to

assess the differences in microbial composition between tick species. Investigation into 'taxa

of interest' warranted a more rigorous assessment of sequence number than was required for

diversity measures in order to avoid any potential cross-contamination and machine cross-

talk (Dong et al., 2017; Wang et al., 2017), as such only samples with >100 sequences were

considered positive. Due to sample collection biases in the present study, prevalence data of

these microbes were not considered statistically relevant. Nucleotide sequences from 'taxa of

interest' (Supplementary File S3) were aligned by MUSCLE (Edgar, 2004) using default

parameters and aligned sequences were then imported into MEGA7 (Kumar et al., 2016) with

the most appropriate nucleotide substitution model chosen based on the lowest Bayesian

Information Criterion (BIC) score. Evolutionary histories were inferred using the Neighbour-

Joining method based on the Tamura 3-parameter model (Tamura, 1992). Bootstrap analysis

was conducted using 10,000 replicates to assess the reliability of inferred tree topologies.

3. Results

3.1 Tick-host associations

Ticks were collected from 221 wildlife hosts including 25 native and three introduced species

(Fig 1, Supplementary File S1). Hosts recorded in the present study were; agile antechinus

(*Antechinus agilis*), brown antechinus (*Antechinus stuartii*), spotted-tail quoll (*Dasyurus*

maculatus), water rat (*Hydromys chrysogaster*), quenda (*Isoodon fusciventer*), northern

brown bandicoot (*Isoodon macrourus*), western grey kangaroo (*Macropus fuliginosus*),

eastern grey kangaroo (*Macropus giganteus*), red-necked wallaby (*Notamacropus*

rufogriseus), red kangaroo (*Osphranter rufus*), platypus (*Ornithorhynchus anatinus*), eastern

barred bandicoot (*Perameles gunnii*), long-nosed bandicoot (*Perameles nasuta*), sugar glider

(*Petaurus breviceps*), eastern ring-tailed possum (*Pseudocheirus peregrinus*), black fruit-bat

(*Pteropus alecto*), fruit-bat sp. (*Pteropus* sp.), bush rat (*Rattus fuscipes*), black rat (*Rattus*

rattus), Tasmanian devil (*Sarcophilus harrisii*), wild pig (*Sus scrofa*), short-beaked echidna

(*Tachyglossus aculeatus*), rufous-bellied pademelon (*Thylogale billardierii*), short-eared

brush-tailed possum (*Trichosurus caninus*), common brush-tailed possum (*Trichosurus*

vulpecula), wombat (*Vombatus ursinus*), red fox (*Vulpes vulpes*), and swamp wallaby

(*Wallabia bicolor*). Female ticks were the dominant instar recorded (*n=547*) followed by

nymphs (*n=319*) and males (*n=153*). Ticks were received from animals in the Northern

Territory (NT) (*n=16*), Queensland (QLD) (*n=249*), New South Wales (NSW) (*n=316*),

Tasmania (TAS) (*n=136*), Victoria (VIC) (*n=44*) and Western Australia (WA) (*n=258*).

Together, *Bothriocroton concolor* (*n=123*), *Ixodes australiensis* (*n=210*), *I. holocyclus*

(*n=173*) and *Ixodes tasmani* (*n=184*) accounted for over two-thirds of all ticks submitted.

3.2 16S rRNA bacterial profiling

A sub-sample of 238 ticks and 19 controls underwent 16S NGS profiling. A total of 23.9

million raw paired-end sequences were generated on the Illumina MiSeq. 17.9 million

sequences were retained after merging, and subsequent quality filtering yielded 14.8 million

sequences for clustering and taxonomic assignment. A total of 4,864 OTUs (average length

299 bases) were retained after background profiles were removed. After removal of

background sequences, a total of 14,328,059 bacterial sequences were assigned to tick

samples. Despite a high number of OTUs, only 1,535 OTUs had greater than 100 total

sequences from tick samples. Tick samples had an average of 60,201 assigned sequences (Fig

2). *Amblyomma triguttatum* had the highest alpha diversity as measured by the observed

number of OTUs and the chao1 index, whereas *Ixodes antechini* had the highest alpha

diversity as measured by the Shannon and Simpson indexes (Fig 3). Analysis of sequence

depth shows that in most cases observed number of OTUs, began to plateau around 50,000

sequences (Fig 4). An ordination plot of taxa (Fig 5) show that bacteria belonging to the

Proteobacteria phylum were the most abundant taxa classified. Principal coordinate analysis

(PCoA) of weighted Uni-Frac distances (Fig 6) demonstrate that microbiome composition

was largely grouped by tick species, with the most notable exception being *I. tasmani*.

Overall eight phyla were recorded, however after removal of low abundant and background

bacterial communities (identified from controls) six phyla were retained consisting of 91

families (Fig 7). Bacterial families identified in tick species are represented as relative

number of sequences in Fig 8.

3.3 Presence of 'taxa of interest'

Phylogenetic analysis of the 'taxa of interest' is represented in Fig 9, showing tick species

and Australian state where each OTU was identified.

A proposed novel *Anaplasma* sp. (OTU230, MK814412, 96.3% similarity) was identified in a

single *B. concolor* (1/20) ex echidna from NSW. A second *Anaplasma bovis*-like OTU

(OTU312, JN862824, 97.3% similarity) was identified in *Haemaphysalis bancrofti* (3/10) ex

red-necked wallaby and long-nosed bandicoot and *Haemaphysalis humerosa* (n = 1/13) tick

ex long-nosed bandicoot, all from NSW.

OTU5509 was assigned to the genus *Bartonella* and had a top BLAST hit of *Bartonella*

bacilliformis (LN624026, 92.1% similarity). OTU5509 was identified in two samples,

however was present in an extremely low number of sequences (2) in each case. Both

samples were *Ixodes fecialis* collected from two different quenda in WA. The low number of

sequences means that in the case of statistical analysis it would have been filtered out. While

there was no sufficient match to this sequence, it is noted that in many instances native

Australian *Bartonella* species are lacking sequence information for this region of the 16S

gene (v1-2).

Two OTUs (OTU14, CP025785, 100% similarity; and OTU4629, CP025785, 100%

similarity) were identified as *'Ca.* B. tachyglossi*'*. Sequences were identified in *B. concolor*

(8/20) ex echidnas from QLD & NSW and *H. humerosa* (1/13) ex northern brown bandicoot

from QLD.

A *Coxiella*-like organism (OTU12, CP032542, 95.5% similarity) was identified exclusively

in *B. concolor* (19/20) ex echidna from NSW and QLD. This was the second most abundant

OTU identified in *B. concolor*, accounting for ~20% of the overall sequences.

A *Coxiella* sp. (OTU15, KC170757, 100% similarity) was identified in *Haemaphysalis*

longicornis (3/4) ex red fox from NSW, this OTU represented ~61.4% of the overall assigned

bacterial sequences.

A novel *Ehrlichia* sp. (OTU33, AY309970, 96.3% similarity) was identified in *I. fecialis*

(1/13) ex quenda from WA. Other *Ehrlichia* OTUs (OTU16 and 1632) represented sequences

from '*Ca.* Ehrlichia ornithorhynchi' in *Ixodes ornithorhynchi* (6/10) ex platypus QLD and

these sequences accounted for the majority of sequences (~50.2%) from *I. ornithorhynchi*.

A *Francisella* endosymbiont (OTU5, AF001077, 98.0% similarity) was identified in *H.*

humerosa (11/13) ticks ex northern brown bandicoots from NT and QLD, and

Haemaphysalis bremneri (2/4) ex possum (species unknown) in QLD. OTU5 was the most

abundance sequence in *H. bremneri* and *H. humerosa*, representing 27.3% and 52.4% of the

assigned sequences, respectively. A second *Francisella* endosymbiont (OTU10, AB001522,

99% similarity) was identified in *H. bancrofti* (8/11) ex red-necked wallabies from QLD and

NSW, and ex a long-nosed bandicoot in NSW; and in a single *H. humerosa* (1/13) ex red-

necked wallaby from NSW. OTU10 was the most abundance sequences in *H. bancrofti*

representing 27.0% of the assigned sequences. OTU40 (AF001077, 97.6% similarity) was

also identified as a *Francisella*-like endosymbiont from *A. triguttatum* (6/6) ex red kangaroo

from WA. It was highly abundant in the ticks, accounting for 76.9% of the overall sequences.

Midichloria (OTU6, FM992372, 100% similarity) was identified in *I. holocyclus* (20/36) ex

long-nosed bandicoots from NSW and QLD, *I. fecialis* (1/13) ex red-necked wallaby NSW,

H. bancrofti (1/11) ex long-nosed bandicoot NSW, and *H. humerosa* (1/13) ex long-nosed

bandicoot NSW. *Midichloria* (OTU7, FM992373, 100% similarity) was identified in *I.*

holocyclus (15/36) ex long-nosed bandicoots from NSW and QLD, *I. fecialis* (1/13) ex red-

necked wallaby NSW and *H. bancrofti* (1/11) ex long-nosed bandicoot NSW. Overall a total

of 15 ticks had both OTU6 and OTU 7, which included *I. holocyclus* (13/36) ex long-nosed

bandicoots from NSW and QLD, *I. fecialis* (1/13) ex red-necked wallaby NSW and *H.*

bancrofti (1/11) ex long-nosed bandicoot NSW. *Midichloria* was the most abundant sequence

from *I. holocyclus* accounting for 36.1% of assigned sequences.

Three *Neoehrlichia* OTUs were identified. '*Ca*. Neoehrlichia arcana' (OTU25, KT203914,

99.3% similarity) was identified in eight ticks, *I. tasmani* (4/51), *I. holocyclus* (2/36), *H.*

bancrofti (1/11) and *H. humerosa* (1/13), ex long-nosed bandicoots NSW and QLD. '*Ca*.

Neoehrlichia australis' (OTU29, KT803957, 100% similarity) was identified in 10 ticks, *I.*

holocyclus (8/36) and *I. tasmani* (2/51), ex long-nosed bandicoots NSW and QLD. Three

ticks (*I. tasmani* and *I. holocyclu*s) were co-infected with both '*Ca*. N arcana' and '*Ca*. N.

australis'. A novel *Neoehrlichia* species (OTU31, MG670107, 97.3% similarity) was

identified in *I. fecialis* (2/13) and *I. australiensis* (2/26) ex quenda from WA, and *I. antechini*

(2/29) ex brown antechinus from NSW.

Due to the conserved nature of the *Rickettsia* genus at the 16S gene, reported previously

(Roux and Raoult, 1995; Stothard and Fuerst, 1995), resolution to species level was not

possible using the sequences obtained in the present study. *Rickettsia* sp. (OTU2, KF318168,

99.7% similarity) had a widespread distribution among samples and was detected in 61 ticks

(B. concolor, H. bancrofti, H. humerosa, I. antechini, I. australiensis, I. holocyclus and *I.*

tasmani) from four states (NSW, NT, QLD, WA). *Rickettsia* was the most abundance

sequence from *B. concolor* and *I. antechini* ticks accounting for 63.3% and 12.0% of the

assigned sequences, respectively

A *Rickettsiella*-like sp. (OTU1, LC388767, 92.7% similarity) was identified in *I.*

australiensis (22/26) and was the most abundant sequence accounting for 76.8% of assigned

sequences. A *Rickettsiella*-like sp. (OTU3, EU430251, 95.1% similarity) was identified in *I.*

tasmani (11/51) ex Tasmanian devils and eastern barred bandicoot from TAS. A

Rickettsiella-like sp. (OTU4, U97547, 98.4% similarity) was identified in, *I. tasmani* (36/51)

ex bandicoots, possum and a sugar glider from NSW, TAS and QLD; *I. australiensis* (1/26)

ex western grey kangaroo from WA; and *I. holocyclus* (1/36) ex long nosed bandicoot NSW.

OTU4 accounted for 30.2% of the overall sequences assigned for *I. tasmani.*

4. Discussion

Currently there is a scarcity of detailed information about the life cycles of Australian ticks,

however it is generally assumed that, as with other hard tick species around the world, the

majority will exhibit a three-host life cycle (Barker and Walker, 2014). Despite these

limitations, the opportunistic sampling strategy used in this study provides an economical

way to survey a wide range of tick and wildlife fauna across a range of geographical areas

across Australia. The paucity of larvae from this data set is also a limitation of this type of

sampling. Tick larval stages, which may indicate the presence of transovarially-transmitted

organisms (Kwan et al., 2017), are difficult to see with the naked eye, and in situations where

collection of ticks is not the main objective, they are easily overlooked (Lydecker et al.,

2019). Furthermore, the opportunistic nature of the sampling precluded assessment of the

infestation rates of the various hosts. In future, if TBPs of humans and the wildlife

themselves are identified, further extensive surveys of tick-pathogen-wildlife ecologies will

require more targeted and comprehensive approaches in order to gather sufficient relevant

epidemiological data.

Overall bacterial diversity of ticks started to plateau at about 50,000 sequences, however, it is

noted that in some cases this plateau was not achieved and deeper sequencing would be

required in order to confidently characterise the full suite of bacterial species present. This

was most evident in *A. triguttatum*, *B. auruginans*, *I. trichosuri* and *I. ornithorhynchi* and

therefore we recommend a minimum of 100,000 sequences per sample for future bacterial

16S amplicon studies in these species. Ordination analysis demonstrated that tick species was

the strongest predictor of bacterial composition. Beta-diversity analysis showed that the

common marsupial tick, *I. tasmani*, exhibited a variable bacterial composition. This diversity

may be explained by the wide geographic distribution of this tick in Australia, its own high

genetic diversity (Burnard and Shao, 2019), and its ability to parasitise many marsupial

species. In the present study, bacterial profiling of *I. tasmani* included ticks from three states

(NSW, QLD, TAS) and seven host species. With this in mind, we suggest that future studies

should combine careful taxonomic status identity with data on microbial communities of *I.*

tasmani. Although sample sizes were limited in 'host-specialist' ticks *B. auruginans*

(wombats)*, H. bremneri* (possums)*, I. antechini* (antechinus) and *I. ornithorhynchi*

(platypus), these specimens showed less diversity of bacterial communities between samples.

The composition and relative proportion of bacterial taxa identified in ticks during the present

study are in line with previous research overseas (Zhang et al., 2014; Kurilshikov et al., 2015;

Sperling et al., 2017) and in Australia (Gofton et al., 2015b; Panetta et al., 2017). Molecular

tools continue to develop at a rapid rate with an increasing shift towards whole-genome or

shotgun sequencing (sequencing the full range of nucleic acids present in a metagenomic

sample). However, comparisons between whole-genome and amplicon sequencing continue

to indicate that the detection of low abundant bacteria is greatly enhanced by using targeted

16S amplicon bacterial profiling methods as used in the present study (Carpi et al., 2011;

Tessler et al., 2017).

Currently in Australia, the ecology, epidemiology, and incidence of human tick-borne

diseases remains largely a matter of conjecture, having received little scientific study

compared with many other parts of the world (Graves and Stenos, 2017). Despite significant

national interest, including a federal government senate inquiry (Radcliffe et al., 2016), the

prevailing scientific opinion concludes that Lyme borreliosis (caused by *B. burgdorferi* sensu

lato), for example, is absent from Australia (Chalada et al., 2016; Irwin et al., 2017). In the

unique Australian environment, long isolated in geological terms, it is likely that unidentified

tick-host life cycles have evolved that may contribute to zoonotic illness when humans

encroach these sylvatic ecologies and become exposed to native ticks. Indeed, it is well

documented that in endemic areas in Europe and North America, the causative agent of Lyme

borreliosis is readily identified in wildlife reservoirs (such as white-tailed deer, *Ococoileus*

virginianus, and white-footed mice, *Peromyscus leucopus*) as well as their natural tick

species (Bosler et al., 1984). The present study therefore is one of the first to explore the

concept that Australian wildlife ticks should be a promising source to identify exotic and

novel TBPs, and we focussed our search towards taxa known to be associated with TBDs

overseas ('taxa of interest'), namely *Anaplasma*, *Bartonella*, *Borrelia*, *Coxiella*, *Ehrlichia*,

Francisella, *Midichloria*, *Neoehrlichia*, *Rickettsia* and *Rickettsiella*.

A new genetic variant of *Anaplasma bovis* was recently described from questing *A.*

triguttatum ticks in WA and the nearby Barrow Island (Gofton et al., 2017). In the present

study we report a range expansion of a genetically similar *A. bovis* along the east coast of

Australia from two widespread *Haemaphysalis* tick species. As suggested by Gofton et al.,

(2017), this finding further supports the hypothesis that it is likely that endemic *A. bovis*

genotypes exist in sylvatic cycles within native Australian ticks and wildlife fauna. Further

research on the phylogenetic position of these *A. bovis* sequences is needed to understand

their likely evolutionary history and relatedness to *A. bovis* genotype Y11 identified on the

west coast of Australia. While three other species of *Anaplasma* (*A. marginale*, *A. centrale*

and *A. platys*) have been introduced to Australia (Rogers and Shiels, 1979; Callow, 1984;

Angus, 1996) the present study did not identify any of these species in ticks from wildlife. A

species of *Anaplasma* identified exclusively in *B. concolor* from echidnas sheds more light on

the diverse range of microbes that have been described from this specialist tick (Loh, 2018).

Although the presence of a *Bartonella* sp. was identified by only four sequences, its absence

in control samples means that this likely represents a true finding in female *I. fecialis*, which

were parasitising different individuals of the same host (quenda) in south-west WA. Recent

studies have shown that Australian marsupials and native rodents harbour a range of distinct

Bartonella species; '*Ca*. Bartonella antechini' has been identified in ticks (*I. antechini*) and

fleas (*Acanthopsylla jordani*) from mardo (*Antechinus flavipes*) in south-west WA

(Kaewmongkol et al., 2011c), '*Ca*. Bartonella woyliei' in woylie ticks (*I. australiensis*) and

fleas (*Pygiopsylla hilli*) in south-west WA, and '*Ca*. Bartonella bandicootii' in fleas

(*Pygiopsylla tunneyi*) from western barred bandicoots (*Perameles bougainville*) on Bernier

and Dorre Island (Kaewmongkol et al., 2011a). Reports of *Bartonella* spp. occurring outside

south-west WA include *Bartonella australis* ex eastern grey kangaroos (*Macropus giganteus)*

(Fournier et al., 2007). Molecular detection of *Bartonella* DNA from Australian ticks has also

been reported in ticks (*I. tasmani*) parasitising koalas (*Phascolarctos cinereus*) from Philip

Island, VIC (Vilcins et al., 2009b). Additional studies on *Bartonella* in Australia wildlife

include those by Gundi et al. (2009), Kaewmongkol et al. (2011a) and Dybing et al. (2016).

Due to the limited size and region of the 16S amplicon generated in the present study, there

were insufficient relevant reference sequences available from other Australian *Bartonella*

species for comparison. A study by Kaewmongkol et al. (2011b) into flea-derived *Bartonella*

from native and introduced Australian species suggests co-evolution of marsupial hosts, their

fleas and the *Bartonella* species.

'*Ca*. Borrelia tachyglossi' was identified in 8/20 *B. concolor* ticks from echidnas in QLD and

NSW, an anticipated finding given previous research into this organism (Loh et al., 2016).

Importantly however, the present study provides the first evidence of '*Ca*. B. tachyglossi'

sequences from a female *H. humerosa* tick parasitising a northern brown bandicoot from

QLD. In addition, no *Bothriocroton* ticks were present in the extraction batch of the positive

H. humerosa sample, and there was no evidence of *Borrelia* sequences from any controls.

Despite the wide geographical range of *H. humerosa* ticks, the restricted finding of '*Ca*. B.

tachyglossi' from QLD supports Loh et al. (2016), suggesting a restricted geographical

distribution along the east coast of Australia.

In tick microbiome studies overseas, *Coxiella* spp. are commonly identified (Khoo et al.,

2016; Machado-Ferreira et al., 2016). Additionally, recent studies on Australian ticks have

also identified *Coxiella* spp. in *A. triguttatum* (Cooper et al., 2013; Gofton et al., 2015b), *B.*

auruginans (Vilcins et al., 2009c), *I. holocyclus* (Cooper et al., 2013) and *R. sanguineus*

(Oskam et al., 2017). Interestingly the present study did not identify the widespread presence

of *Coxiella* spp. with only two unique OTUs identified from *B. concolor* and *H. longicornis*,

both of which appear to be host specific to those tick species, and the causative agent of Q-

Fever (*C. burnetii*) was not identified from ticks in the present study.

Ehrlichia sequences ('*Ca*. E. ornithorhynchi') from platypus ticks, *I. ornithorhynchi*, were

recently described (Gofton et al., 2018). In the present study, '*Ca*. E. ornithorhynchi' was

exclusively observed within *I. ornithorhynchi,* suggesting this genetically distinct *Ehrlichia*

species has a unique and host-specific relationship with the platypus. During this study, a

potentially novel species of *Ehrlichia* was identified in an *I. fecialis* female tick from a

quenda in WA. BLAST results show it is a close relative to an *Ehrlichia* sp. detected in

Haemaphysalis ticks from Japan (Inokuma et al., 2004). No sequences from the recently

described '*Ca*. E. occidentalis' (Gofton et al., 2017) were identified in the present study,

however it is noted *A. triguttatum* were only represented by six samples.

Francisella-like endosymbionts have been widely reported in ticks overseas, such as

Dermacentor spp. (Scoles, 2005), *Dermacentor occidentalis* (Gurfield et al., 2017),

Haemaphysalis longicornis (Wang et al., 2018) and *Hyalomma rufipes* (Szigeti et al., 2014),

and *Francisella* sequences have previously been identified in *A. fimbriatum* ticks from

reptiles in the Northern Territory of Australia (Vilcins et al., 2009a). In the present study, a

Francisella-like endosymbiont was identified in 100% of *A. triguttatum* ticks, and in a high

proportion. The high prevalence and relative abundance of the *Francisella*-like organisms in

A. triguttatum may be of medical relevance with respect to this common human-biting tick,

particularly given the recent report of *Francisella* bacteraemia in WA (Aravena-Román et al.,

2015). Importantly, as noted previously for *I. holocyclus* ticks, organisms in lower abundance

may be masked by plentiful endosymbionts, unless samples are sequenced deeply and/or a

blocking primer is used (Gofton et al., 2015a). Therefore, future bacterial profiling studies of

A. triguttatum should take these factors into consideration.

While '*Ca*. M. mitochondrii' has been documented in *I. holocyclus,* the present study

provides the first report of '*Ca*. M. mitochondrii' within additional species of native

Australian ticks (*I. fecialis, H. bancrofti* and *H. humerosa*). Despite this expansion among

tick species, identification of '*Ca*. M. mitochondrii' remains confined to NSW and QLD as

previously reported (Gofton et al., 2015a). '*Ca*. M. mitochondrii' has been detected overseas

in laboratory models (Cafiso et al., 2019), wildlife (Serra et al., 2018), and humans

(Mariconti et al., 2012).

A recent addition to the Anaplasmataceae family, '*Ca*. Neoehrlichia mikurensis' was first

isolated from wild rats and their ticks (*Ixodes ovatus*) in Japan (Kawahara et al., 2004). In

Europe, the hedgehog (*Erinaceus* spp.), a common peri-urban dweller, has been shown to

play an important role in life cycle of '*Ca*. N. mikurensis' (Földvári et al., 2014; Jahfari et al.,

2017). A high proportion of bandicoot ticks were infected with *Neoehrlichia* spp. in this

study, suggesting that these marsupials may play an important role in the life cycle of this

bacterium on the Australian continent. Bandicoots also frequently inhabit gardens and have

relatively close contact with humans (Carthey and Banks, 2012) closer study (including blood

samples and tissue biopsies) of these marsupials may provide vital information about the life

cycle of these microbes and the potential risk for human infection.

Rickettsia spp. are among the only currently recognised TBPs affecting people in Australia,

however, as previously outlined, the conserved nature of the 16S gene in *Rickettsia* precludes

rigorous species delimitation from the relatively short sequences generated in this dataset. In

addition to the known human pathogens (*R. australis* and *R. honei*) (Graves and Stenos,

2017), the presence of *Rickettsia* has been demonstrated from a variety of Australian tick

species. *Rickettsia gravesii* has been described from *A. triguttatum* (Li et al., 2010; Abdad et

al., 2017) and *Rickettsia* sequences have been previously identified in *I. tasmani* ex

Tasmanian devils (Vilcins et al., 2009c) and *Amblyomma fimbriatum* from reptiles (Vilcins et

al., 2009a).

The diversity and significance of the *Rickettsiella* genus remains largely unknown globally.

Phylogenetic analysis of sequence data places the genus within the Coxiellaceae family

(Fournier and Raoult, 2005; Leclerque and Kleespies, 2012). While the presence of these

organisms has been well documented in tick species around the world and in some cases they

have been identified as pathogenic to arthropods (Kurtti et al., 2002; Leclerque et al., 2012),

disease causation within the vertebrate hosts remains unknown at the present time. A study

by Vilcins et al. (2009c) identified *Rickettsiella* in *I. tasmani* ex koalas from Phillip Island,

just off the coast of south eastern Australia. A genetically similar sequence was identified in

the present study (OTU3) from *I. tasmani* exclusively in Tasmania.

While this is the first study to characterise the bacterial communities within native Australian

wildlife ticks, it is apparent that much research is still required in order to better understand

the tick-associated microbial life cycles and their ecologies, let alone elucidating their

potential for transmission and pathogenicity in vertebrate hosts, including humans.

Furthermore, whilst cost-effective, ticks were collected opportunistically resulting in an

inherent geographical bias, confining sampling largely to urban areas along the east coast of

the Australian continent. Nevertheless, this area is also where most humans receive tick bites,

although reliable data about this is also lacking.

The analysis of ticks removed from wildlife hosts comes with the inherent complication of a

blood meal. The host blood meal has recently been shown to affect both the tick microbiome

composition and the presence of pathogens (Landesman et al., 2019; Swei and Kwan, 2017).

Further targeted studies are needed to assess the source of microbes with respect to a tick or

host origin (Irwin et al., 2018). In addition, limitations in the current sequencing technologies

mean that high-throughput methods favour short amplicons which may not be able to

accurately discriminate bacterial species, as is the case with *Rickettsia* in particular (Gihring

et al., 2012). Furthermore, important caveats to consider in metabarcoding microbial diversity

analysis include PCR efficiency (including primers used and length of target amplicon),

variations in 16S gene copy number (Ahn et al., 2012), sequencing depth, machine cross-talk,

bioinformatic analysis and taxonomic assignment. While there have been some recent

attempts to correct these biases (Kembel et al., 2012; Rosselli et al., 2016), their application

to uncharacterised metagenomic samples remains limited.

5. Conclusions

With over 75% of emerging human infectious pathogens originating from wildlife (King,

2014), surveillance methods that target these species are important for investigations of

emerging and exotic infectious disease. Research into the interactions of wildlife hosts, ticks

and pathogens in Europe and North America continues to highlight the complexity of these

dynamic systems (Ostfeld et al., 2018; Tomassone et al., 2018). Results from the present

study build on recent research into Australian tick-associated microbes, further highlighting

the diversity of organisms present that appear related, yet distinct from their overseas

counterparts. With the evolutionary history of Australia's unique tick species (Beati and

Klompen, 2019) and wildlife fauna, it is likely that reports of their tick-associated microbes

(potential pathogens) will continue to reveal taxonomic differences from those described in

the northern hemisphere. As such, future research on emerging zoonoses should include

methods that are able to detect novel micro-organisms in humans and reservoirs. As

anthropogenic changes on the environment continue to grow in Australia, a greater emphasis

on wildlife disease surveillance is critical to ensure the early detection of potential infectious

diseases affecting humans, livestock, companion animals and wildlife (Woods et al., 2019).

Advances in pathogen detection and characterisation are greatly enhanced by collaboration;

the authors advocate for continued multidisciplinary efforts between health professionals,

researchers, land managers and local communities.

Supporting Information

Data availability

Associated metadata output from bioinformatic analysis and subsequent data visualisation is available on FigShare repository https://doi.org/10.6084/m9.figshare.c.4608803.v1. Next-generation sequencing data can be accessed from NCBI Sequence Read Archive under BioProject: PRJNA559059 (BioSample accession numbers: SAMN12512474 – SAMN12512711).

Acknowledgements

The authors would like to thank collaborating research teams, veterinary clinics, wildlife centres and members of the public for contributing samples to this study. Supplementary information S2 contains complete list of collectors and institutions. This study was part-funded by the Australian Research Council (LP160100200) and Bayer HealthCare (Germany) and Bayer Australia. S.E. received funding from a Murdoch University honours scholarship. S-M.L. was support by a Murdoch International Postgraduate scholarship. The authors also acknowledge the Pawsey Supercomputing Centre and the Western Australia State Agriculture Biotechnology Centre for use of its facilities. We thank Dr. Alexander Gofton and Ms. Telleasha Greay for advice on next-generation sequencing methods. The funders had no role in the experimental design, data collection and analyses.

Compliance with ethical standards

This study was conducted under the compliance of the Australian Code for the Responsibility Conduct of Research (2007) and Australian Code for the Care and Use of Animals for Scientific Purposes, 2013. Tick collection was carried out opportunistically with the approval

from the Murdoch University Animal Ethics Committee.

Figure Captions

Fig 1. Chord diagram showing tick-wildlife associations recorded in the present study. The

wildlife hosts recorded are represented on the lower half of the plot, and tick species across

the top. Host records that could not be assigned to species level are not represented in this

plot. A complete list of records are available in Supplementary File S1. Silhouette wildlife

images sourced from phylopic.org.

Fig 2. Sequence information from 16S rRNA bacterial profiling of ticks on the Illumina

MiSeq platform after bioinformatic analysis; (a) number sequences assigned to 97%

operational taxonomic units (OTUs) and (b) number of sequences per tick.

Fig 3. Alpha diversity of bacterial communities in ticks parasitising Australian wildlife

measured as; observed number of operational taxonomic units (OTUs), Chao1 index,

Shannon index and Simpson index. Tick species abbreviated to; *A. triguttatum* (*A. tri*), *B.*

aurugians (*B. aur*), *B. concolor* (*B. con*), *H. bancrofti* (*H. ban*), *H. bremneri* (*H. bremneri*),

H. humerosa (*H. hum*), *H. longicornis* (*H. lon*), *I. antechini* (*I. ant*), *I. australiensis* (*I. aus*), *I.*

fecialis (*I. fec*), *I. holocyclus* (*I. hol*), *I. myrmecobii* (*I. myr*), *I. ornithorhynchi* (*I. orn*), *I.*

tasmani (*I. tas*), and *I. trichosuri* (*I. tri*).

Fig 4. Rarefaction plot of 16S rRNA bacterial sequences clustered into 97% operational

taxonomic units (OTUs) from ticks parasitising Australian wildlife.

Fig 5. Non-metric multidimensional scaling (NMDS) plot of operational taxonomic units

(OTUs) based on Bray-Curtis dissimilarity matrix. Taxa filtered to only display OTUs that

were present at least twice in >10% of tick samples.

Fig 6. Principal coordinate analysis (PCoA) plot of the weighted UniFrac distance matrix.

Plots shows beta-diversity of tick bacterial communities, displayed by tick species and instar.

Operational taxonomic units (OTUs) that represented <100 sequences in a tick sample were

disregarded.

Fig 7. Heat tree of overall microbiome composition of ticks parasitising Australian wildlife.

Taxa is shown to family level with colour scale representing number of operational

taxonomic units (OTUs).

Fig 8. Stacked barplot of bacterial composition (shown at family level) from Australian ticks

represented as percentage of assigned sequences. Taxa at the family level that represented

<10% of the relative sequences within each tick species were grouped as "Low abundant".

Sequences not able to be accurately assigned to family taxa are displayed as "Unclassified".

Fig 9. Neighbour-joining phylogenetic tree displaying taxa of interest prevalence and

distribution among tick samples. Geographic data relating to tick collection is represented by

state; New South Wales (NSW), Northern Territory (NT), Queensland (QLD), Tasmania

(TAS) and Western Australia (WA). Evolutionary histories were inferred based on the

Tamura 3-parameter model with bootstrap analysis (10,000 replicated). Tick samples were

considered positive for taxa of interest if >100 sequences present. Information on number of

sequences, taxonomic identity (as inferred from NCBI BLAST analysis with nucleotide

database), and sequences can be sound in supplementary information (Supplementary File S2

& S3).

Supplementary Files

Supplementary File S1. Sample metadata of tick-wildlife records identified from the present

study.

Supplementary File S2. Number of sequences relating to the taxa of interest in tick samples.

The top hit from NCBI BLAST results showing Genbank accession number, percent identity

and E-value are given, and final evaluation of taxonomic assignment.

Supplementary File S3. Fasta sequences for operational taxonomic units (OTUs) used in the

taxa of interest investigation.

614   References

Abdad, M.Y., Abou Abdallah, R., Karkouri, K., Beye, M., Stenos, J., Owen, H., Unsworth, N., Robertson, I., Blacksell, S.D., Nguyen, T.T., Nappez, C., Raoult, D., Fenwick, S., Fournier, P.E., 2017. *Rickettsia gravesii* sp. nov.: a novel spotted fever group rickettsia in Western Australian *Amblyomma triguttatum triguttatum*. Int. J. Syst. Evol. Microbiol. 67(9), 3156–3161.

Ahantarig, A., Trinachartvanit, W., Baimai, V., Grubhoffer, L., 2013. Hard ticks and their bacterial endosymbionts (or would be pathogens). Folia Microbiol. (Praha) 58(5), 419–28.

Ahn, J.H., Kim, B.Y., Song, J., Weon, H.Y., 2012. Effects of PCR cycle number and DNA polymerase type on the 16S rRNA gene pyrosequencing analysis of bacterial communities. J. Microbiol. 50(6), 1071–1074.

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search tool. J Mol Biol 515(3), 403-410.

Andreotti, R., Perez deLeon, A.A., Dowd, S.E., Guerrero, F.D., Bendele, K.G., Scoles, G.A., 2011. Assessment of bacterial diversity in the cattle tick *Rhipicephalus* (*Boophilus*) *microplus* through tag-encoded pyrosequencing. BMC Microbiol. 11(1), 6.

Angus, B.M., 1996. The history of the cattle tick *Boophilus microplus* in Australia and achievement in its control. Int. J. Parasitol. 26(12), 1341–1355.

Aravena-Román, M., Merritt, A., Inglis, T.J., 2015. First case of *Francisella* bacteraemia in Western Australia. New Microbes New Infect. 8, 75–77.

Barker, S.C., Walker, A.R., 2014. Ticks of Australia. The species that infest domestic animals and humans. Zootaxa 3816, 1-144

Beati, L., Klompen, H., 2019. Phylogeography of Ticks (Acari: Ixodida). Annu. Rev. Entomol. 64, 379–397.

Bokulich, N.A., Kaehler, B.D., Rideout, B.A., Dillon, M., Bolyen, E., Knight, R., Huttley, G.A., Caporaso, J.G., 2018. Optimizing taxonomic classification of marker gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. Microbiome 6, 90.

Bonnet, S., Michelet, L., Moutailler, S., Cheval, J., Hebert, C., Vayssier-Taussat, M., Eloit, M., 2014. Identification of parasitic communities within European ticks using next-generation sequencing. PLoS Negl. Trop. Dis. 8(3), e2753.

Bonnet, S.I., Binetruy, F., Hernández-Jarguín, A.M., Duron, O., 2017. The tick microbiome: why non-pathogenic microorganisms matter in tick biology and pathogen transmission. Front. Cell. Infec. Microbiol. 7, 236.

Bosler, E.M., Ormiston, B.G., Coleman, J.L., Hanrahan, J.P., Benach, J.L., 1984. Prevalence of the Lyme Disease spirochete in populations of white-tailed deer and white-footed Mice. Yale J. Biol. Med. 57(4), 651–659.

Burnard, D., Shao, R., 2019. Mitochondrial genome analysis reveals intraspecific variation within Australian hard tick species. Ticks Tick Borne Dis. 10(3), 677–681.

Cafiso, A., Sassera, D., Romeo, C., Serra, V., Hervet, C., Bandi, C., Plantard, O., Bazzocchi, C., 2019. *Midichloria mitochondrii*, endosymbiont of *Ixodes ricinus*: evidence for the transmission to the vertebrate host during the tick blood meal. Ticks Tick Borne Dis. 10(1), 5–12.

Callow, L.L., 1984. Animal Health in Australia. Volume 5: Protozoal and Rickettsial Diseases. Australian Government Publishing Services, Canberra, Australia.

Carthey, A.J.R., Banks, P.J., 2011. When does an alien become a native species? A vulnerable native mammal recognizes and responds to its long-term alien predator. PLoS One

7(2), e31804.

Carpi, G., Cagnacci, F., Wittekindt, N.E., Zhao, F., Qi, J., Tomsho, L.P., Drautz, D.I., Rizzoli, A., Schuster, S.C., 2011. Metagenomic profile of the bacterial communities associated with *Ixodes ricinus* ticks. PLoS One 6(10), e25604.

Chalada, M.J., Stenos, J., Bradbury, R.S., 2016. Is there a Lyme-like disease in Australia? Summary of findings to date. One Health 2, 42–54.

Chalada, M.J., Stenos, J., Vincent, G., Barker, D., Bradbury, R.S., 2018. A molecular survey of tick-borne pathogens from ticks collected in central Queensland, Australia. Vector Borne Zoonotic Dis 18, 151–163.

Cooper, A., Stephens, J., Ketheesan, N., Govan, B., 2013. Detection of Coxiella burnetii DNA in Wildlife and Ticks in Northern Queensland, Australia. Vector Borne Zoonotic Dis. 13(3), 12–16.

de la Fuente, J., Antunes, S., Bonnet, S., Cabezas-Cruz, A., Domingos, A.G., Estrada-Pena, A., Johnson, N., Kocan, K.M., Mansfield, K.L., Nijhof, A.M., Papa, A., Rudenko, N., Villar, M., Alberdi, P., Torina, A., Ayllon, N., Vancova, M., Golovchenko, M., Grubhoffer, L., Caracappa, S., Fooks, A.R., Gortazar, C., Rego, R.O.M., 2017. Tick-pathogen interactions and vector competence: Identification of molecular drivers for tick-borne diseases. Front. Cell. Infect. Microbiol. 7, 114.

DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber, T., Dalevi, D., Hu, P., Andersen, G.L., 2006. Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. Appl. Environ. Microbiol. 72, 5069–5072.

Dobson, A.D.M., Finnie, T.J.R., Randolph, S.E., 2011. A modified matrix model to describe the seasonal population ecology of the European tick *Ixodes ricinus*. J. Appl. Ecol. 48(4), 1017–1028.

Dong, X., Kleiner, M., Sharp, C.E., Thorson, E., Li, C., Liu, D., Strous, M., 2017. Fast and simple analysis of MiSeq amplicon sequencing data with MetaAmp. Front. Microbiol. 8, 1461.

Dybing, N.A., Jacobson, C., Irwin, P., Algar, D., Adams, P.J., 2016. *Bartonella* species identified in rodent and feline hosts from island and mainland Western Australia. Vector Borne Zoonotic Dis. 16(4), 238–44.

Edgar, R.C., 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. Nat. Methods 10(1), 996–968.

Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics 26(19), 2640–2461.

Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res. 32(5), 1792–1797.

Edgar, R.C., Flyvbjerg, H., 2015. Error filtering, pair assembly and error correction for next-generation sequencing reads. Bioinformatics 31(21), 3476–82.

Estrada-Pena, A., Gray, J., Kahl, O., Lane, R.S., Nijhof, A.M., 2013. Research on the ecology of ticks and tick-borne pathogens - methodological principles and caveats. Front. Cell. Infect. Microbiol. 3, 29.

Ferrell, A., Brinkerhoff, R., 2018. Using landscape analysis to test hypotheses about drivers of tick abundance and infection prevalence with *Borrelia burgdorferi*. Int. J. Environ. Res. Public Health 15(4), 737.

Földvári, G., Jahfari, S., Rigó, K., Jablonszky, M., Szekeres, S., Majoros, G., Tóth, M., Molnár, V., Coipan, E.C., Sprong, H., 2014. *Candidatus* Neoehrlichia mikurensis and

Anaplasma phagocytophilum in Urban Hedgehogs. Emerg. Infect. Dis. 20(3), 496–498.

Fournier, P., Taylor, C., Rolain, J.M., Barrassi, L., Smith, G., Raoult, D., 2007. *Bartonella australis* sp. nov. from kangaroos, Australia. Emerg. Infect. Dis. 13(12), 1961–1963.

Fournier, P.-E., Raoult, D., 2005. Genus II. Rickettsiella Philip 1956, in: Garrity, G.M., Brenner, D.J., Krieg, N.R., Staley, J.T. (Eds.), Bergey's Manual of Systematic Bacteriology, Part B. Springer, New York, pp. 241–247.

Gihring, T.M., Green, S.J., Schadt, C.W., 2012. Massively parallel rRNA gene sequencing exacerbates the potential for biased community diversity comparisons due to variable library sizes. Environ. Microbiol. 14(2), 285–290.

Gofton, A.W., Doggett, S.L., Ratchford, A., Oskam, C.L., Paparini, A., Ryan, U.M., Irwin, P.J., 2015b. Bacterial profiling reveals novel "Ca. Neoehrlichia", *Ehrlichia*, and *Anaplasma* species in Australian human-biting ticks. PLoS One 10(12), e0145449.

Gofton, A.W., Loh, S.M., Barbosa, A.D., Paparini, A., Gillett, A., Macgregor, J., Oskam, C.L., Ryan, U.M., Irwin, P.J., 2018. A novel *Ehrlichia* species in blood and *Ixodes ornithorhynchi* ticks from platypuses (*Ornithorhynchus anatinus*) in Queensland and Tasmania, Australia. Ticks Tick Borne Dis. 9(2), 435–442.

Gofton, A.W., Oskam, C.L., Lo, N., Beninati, T., Wei, H., McCarl, V., Murray, D.C., Paparini, A., Greay, T.L., Holmes, A.J., Bunce, M., Ryan, U.M., Irwin, P.J., 2015a. Inhibition of the endosymbiont "*Candidatus* Midichloria mitochondrii" during 16S rRNA gene profiling reveals potential pathogens in Ixodes ticks from Australia. Parasit. Vectors 8, 345.

Gofton, A.W., Waudby, H.P., Petit, S., Greay, T.L., Ryan, U.M., Irwin, P.J., 2017. Detection and phylogenetic characterisation of novel *Anaplasma* and *Ehrlichia* species in *Amblyomma triguttatum* subsp. from four allopatric populations in Australia. Ticks Tick Borne Dis. 8(5), 749–756.

Graves, S.R., Stenos, J., 2017. Tick-borne infectious diseases in Australia. Med. J. Aust. 206(7), 320–324.

Guglielmone, A.A., Robbins, R.G., Apanaskevich, D.A., Petney, T.N., Estrada-Pena, A., Horak, I.G., Shao, R., Barker, S.C., 2010. The Argasidae, Ixodidae and Nuttalliellidae (Acari: Ixodida) of the world: a list of valid species names. Zootaxa 2528(1-4), 27.54.

Gundi, V.A., Taylor, C., Raoult, D., La Scola, B., 2009. *Bartonella rattaustraliani* sp. nov., *Bartonella queenslandensis* sp. nov. and *Bartonella coopersplainsensis* sp. nov., identified in Australian rats. Int. J. Syst. Evol. Microbiol. 59(Pt 12), 2956–2961.

Gurfield, N., Grewal, S., Cua, L.S., Torres, P.J., Kelley, S.T., 2017. Endosymbiont interference and microbial diversity of the Pacific coast tick, *Dermacentor occidentalis*, in San Diego County, California. PeerJ 5, e3202.

Inokuma, H., Beppu, T., Okuda, M., Shimada, Y., Sakata, Y., 2004. Detection of Ehrlichial DNA in *Haemaphysalis* ticks recovered from dogs in Japan that is slosely related to a novel *Ehrlichia* sp. found in cattle yicks from Tibet, Thailand, and Africa. J. Clin. Microbiol. 42(3), 1353–1355.

Irwin, P.J., Egan, S., Greay, T., Oskam, C.L., 2018. Bacterial tick-associated infections in Australia: current studies and future direction. Microbiol. Aust. 39(4), 200–202.

Irwin, P.J., Robertson, I.D., Westman, M.E., Perkins, M. Straubinger, R.K., 2017. Searching for Lyme borreliosis in Australia: results of a canine sentinel study. Parasit. Vectors 10(1), 114.

Jahfari, S., Ruyts, S.C., Frazer-Mendelewska, E., Jaarsma, R., Verheyen, K., Sprong, H., 2017. Melting pot of tick-borne zoonoses: the European hedgehog contributes to the

maintenance of various tick-borne diseases in natural cycles urban and suburban areas. Parasit. Vectors 10(1), 134.

Jones, K.E., Patel, N.G., Levy, M.A., Storeygard, A., Balk, D., Gittleman, J.L., Daszak, P., 2008. Global trends in emerging infectious diseases. Nature 451(7181), 990–993.

Kaewmongkol, G., Kaewmongkol, S., Burmej, H., Bennett, M.D., Fleming, P.A., Adams, P.J., Wayne, A.F., Ryan, U., Irwin, P.J., Fenwick, S.G., 2011a. Diversity of *Bartonella* species detected in arthropod vectors from animals in Australia. Comp. Immunol. Microbiol. Infect. Dis. 34(5), 411–417.

Kaewmongkol, G., Kaewmongkol, S., McInnes, L.M., Burmej, H., Bennett, M.D., Adams, P.J., Ryan, U., Irwin, P.J., Fenwick, S.G., 2011b. Genetic characterization of flea-derived *Bartonella* species from native animals in Australia suggests host-parasite co-evolution. Infect. Genet. Evol. 11(8), 1868–1872.

Kaewmongkol, G., Kaewmongkol, S., Owen, H., Fleming, P.A., Adams, P.J., Ryan, U., Irwin, P.J., Fenwick, S.G., 2011c. Candidatus Bartonella antechini: a novel *Bartonella* species detected in fleas and ticks from the yellow-footed antechinus (*Antechinus flavipes*), an Australian marsupial. Vet. Microbiol. 149(3-4), 517–521.

Kawahara, M., Rikihisa, Y., Isogai, E., Takahashi, M., Misumi, H., Suto, C., Shibata, S., Zhang, C., Tsuji, M., 2004. Ultrastructure and phylogenetic analysis of "*Candidatus* Neoehrlichia mikurensis" in the family Anaplasmataceae, isolated from wild rats and found in *Ixodes ovatus* ticks. Int. J. Syst. Evol. Microbiol. 54(Pt 5), 1837–1843.

Kembel, S.W., Wu, M., Eisen, J.A., Green, J.L., 2012. Incorporating 16S gene copy number information improves estimates of microbial diversity and abundance. PLoS Comput. Biol. 8(10), e1002743.

Khoo, J.J., Lim, F.S., Chen, F., Phoon, W.H., Khor, C.S., Pike, B.L., Chang, L.Y., AbuBakar, S., 2016. *Coxiella* detection in ticks from wildlife and livestock in Malaysia. Vector Borne Zoonotic Dis. 16(12), 744–751.

King, L.J., 2014. Combating the triple threat: the need for a one health approach, in: Atlas, R.M., Maloy, S. (Eds.), One Health: People, Animals and the Environment. American Society for Microbiology, Washington, pp. 3–16.

Kumar, S., Stecher, G., Tamura, K., 2016. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. Mol. Biol. Evol. 33(7), 1870–1874.

Kurilshikov, A., Livanova, N.N., Fomenko, N.V., Tupikin, A.E., Rar, V.A., Kabilov, M.R., Livanov, S.G., Tikunova, N.V., 2015. Comparative metagenomic profiling of symbiotic bacterial communities associated with I*xodes persulcatus*, *Ixodes pavlovskyi* and *Dermacentor reticulatus* ticks. PLoS One 10(7), e0131413.

Kurtti, T.J., Palmer, A.T., Oliver, J.H., 2002. *Rickettsiella*-like bacteria in *Ixodes woodi* (Acari: Ixodidae). J. Med. Entomol. 39(3), 534–540.

Kwan, J.Y., Griggs, R., Chicana, B., Miller, C., Swei, A., 2017. Vertical vs. horizontal transmission of the microbiome in a key disease vector, *Ixodes pacificus*. Mol. Ecol. 26(23), 6578–6589.

Landesman, W.J., Mulder, K., Allan, B.F., Bashor, L.A., Keesing, F., LoGiudice, K., Ostfeld, R.S., 2019. Potential effects of blood meal host on bacterial community composition in *Ixodes scapularis* nymphs. Ticks Tick Borne Dis. 10(3), 523-527.

Leclerque, A., Kleespies, R.G., 2012. A *Rickettsiella* bacterium from the hard tick, *Ixodes woodi*: molecular taxonomy combining multilocus sequence typing (MLST) with significance testing. PLoS One 7(5), e38062.

Leclerque, A., Kleespies, R.G., Schuster, C., Richards, N.K., Marshall, S.D.G., Jackson, T.A.,

2012. Multilocus sequence analysis (MLSA) of '*Rickettsiella costelytrae*' and '*Rickettsiella pyronotae*', intracellular bacterial entomopathogens from New Zealand. J. Appl. Microbiol. 113(5), 1228–1237.

Li, A.Y., Adams, P.J., Abdad, M.Y., Fenwick, S.G., 2010. High prevalence of *Rickettsia gravesii* sp. nov. in *Amblyomma triguttatum* collected from feral pigs. Vet. Microbiol. 146(1-2), 59–62.

LoGiudice, K., Ostfeld, R.S., Schmidt, K.A., Keesing, F., 2003. The ecology of infectious disease: effects of host diversity and community composition on Lyme disease risk. Proc. Natl. Acad. Sci. U.S.A. 100(2), 567–571.

Loh, S.M., 2018. Identification and characterisation of microorganisms in Australian wildlife tick (PhD thesis). Murdoch University, Perth, Western Australia.

Loh, S.M., Gofton, A.W., Lo, N., Gillett, A., Ryan, U.M., Irwin, P.J., Oskam, C.L., 2016. Novel *Borrelia* species detected in echidna ticks, *Bothriocroton concolor*, in Australia. Parasites & Vectors 9, 339.

Lydecker, H.W., Banks, P.B., Hochuli, D.F., 2019. Counting ticks (Acari: Ixodida) on hosts is complex: a review and comparison of methods. J. Med. Entomol. tjz094.

Machado-Ferreira, E., Vizzoni, V.F., Balsemao-Pires, E., Moerbeck, L., Gazeta, G.S., Piesman, J., Voloch, C.M., Soares, C.A., 2016. *Coxiella* symbionts are widespread into hard ticks. Parasitol. Res. 115(12), 4691–4699.

Mariconti, M., Epis, S., Gaibani, P., Dalla Valle, C., Sassera, D., Tomao, P., Fabbi, M., Castelli, F., Marone, P., Sambri, V., Bazzocchi, C., Bandi, C., 2012. Humans parasitized by the hard tick *Ixodes ricinus* are seropositive to *Midichloria mitochondrii*: is *Midichloria* a novel pathogen, or just a marker of tick bite? Pathog. Glob. Health 106(7), 391–396.

McMurdie, P.J., Holmes, S., 2013. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. PLoS One 8(4), e61217.

Mediannikov, O., Fenollar, F., 2014. Looking in ticks for human bacterial pathogens. Microb. Pathog. 77, 142–8.

Merrill, M.M., Boughton, R.K., Lord, C.C., Sayler, K.A., Wight, B., Anderson, W.M., Wisely, S.M., 2018. Wild pigs as sentinels for hard ticks: A case study from south-central Florida. Int. J. Parasitol. Parasites Wildl. 7(2), 161–170.

Morens, D.M., Fauci, A.S., 2012. Emerging infectious diseases in 2012: 20 years after the institute of medicine report. MBio 3(6), e00494-12.

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2019. Vegan: community ecology package. R package version 2.5-5

Oskam, C.L., Gofton, A.W., Greay, T.L., Yang, R., Doggett, S., Ryan, U.M., Irwin, P.J., 2017. Molecular investigation into the presence of a *Coxiella* sp. in *Rhipicephalus sanguineus* ticks in Australia. Vet. Microbiol. 201, 141–145.

Ostfeld, R.S., Brunner, J.L., 2015. Climate change and *Ixodes* tick-borne diseases of humans. Philos. Trans. R. Soc. Lond. B. Biol. Sci. 370(1665), 20140051.

Ostfeld, R.S., Levi, T., Keesing, F., Oggenfuss, K., Canham, C.D., 2018. Tick-borne disease risk in a forest food web. Ecology 99,(7) 1562–1573.

Panetta, J.L., Šíma, R., Calvani, N.E.D., Hajdušek, O., Chandra, S., Panuccio, J., Šlapeta, J., 2017. Reptile-associated *Borrelia* species in the goanna tick (*Bothriocroton undatum*) from Sydney, Australia. Parasit. Vectors 10(1), 616.

Parola, P., Raoult, D., 2001. Ticks and tickborne bacterial diseases in humans: an emerging infectious threat. Clin. Infect. Dis. 32(6), 897–928.

Pfäffle, M., Littwin, N., Muders, S.V., Petney, T.N., 2013. The ecology of tick-borne diseases. Int. J. Parasitol. 43(12-13), 1059–1077.

Poisot, T., Foster, Z.S.L., Sharpton, T.J., Grünwald, N.J., 2017. Metacoder: An R package for visualization and manipulation of community taxonomic diversity data. PLOS Comput. Biol. 13(2), e1005404.

Ponnusamy, L., Gonzalez, A., Treuren, W.V., Welss, S., Parobek, C.M., Jullano, J.J., Knight, R., Roe, R.M., Apperson, C.S., Meshnick, S.R., 2014. Diversity of *Rickettsiales* in the microbiome of the lone star tick, *Amblyomma americanum*. Appl. Environ. Microbiol. 80(1), 354–359.

Radcliffe, J., Rusjakovski, N., Forkert, J., Jones, M., Stewart, C., 2016. Growing evidence of an emerging tick-borne disease that causes a Lyme-like illness for many Australian patients. Community Affairs References Committee, Canberra, Australia.

Rinker, D.C., Pitts, R.J., Zwiebel, L.J., 2016. Disease vectors in the era of next generation sequencing. Genome Biol. 17(1), 95.

Roberts, F.S.H., 1970. Australian Ticks. CSIRO, Melbourne.

Rogers, R.J., Shiels, I.A., 1979. Epidemiology and control of anaplasmosis in Australia. J. S. Afr. Vet. Assoc. 50(4), 363–366.

Rosselli, R., Romoli, O., Vitulo, N., Vezzi, A., Campanaro, S., de Pascale, F., Schiavon, R., Tiarca, M., Poletto, F., Concheri, G., Valle, G., Squartini, A., 2016. Direct 16S rRNA-seq from bacterial communities: a PCR-independent approach to simultaneously assess microbial diversity and functional activity potential of each taxon. Sci. Rep. 6, 32165.

Roux, V., Raoult, D., 1995. Phylogenetic analysis of the genus *Rickettsia* by 16S rDNA sequencing. Res. Microbiol. 146(5), 385–396.

RStudio Team, 2015. RStudio: Integrated Development for R. RStudio Inc, Boston, MA.

Scoles, G.A., 2004. Phylogenetic analysis of the *Francisella*-like endosymbionts of *Dermacentor* ticks. J. Med. Entomol. 41(3), 277–286.

Serra, V., Cafiso, A., Formenti, N., Verheyden, H., Plantard, O., Bazzocchi, C., Sassera, D., 2018. Molecular and serological evidence of the presence of *Midichloria mitochondrii* in roe deer (*Capreolus capreolus*) in France. J. Wildl. Dis. 54(3), 597–600.

Sperling, J.L., Silva-Brandão, K.L., Brandão, M.M., Lloyd, V.K., Dang, S., Davis, C.S., Sperling, F.A.H., Magor, K.E., 2017. Comparison of bacterial 16S rRNA variable regions for microbiome surveys of ticks. Ticks Tick Borne Dis. 8(4), 453–461.

Špitalská, E., Sparagano, O., Stanko, M., Schwarzová, K., Špitalský, Z., Škultéty, Ľ., Havlíková, S.F., 2018. Diversity of *Coxiella*-like and *Francisella*-like endosymbionts, and *Rickettsia* spp., *Coxiella burnetii* as pathogens in the tick populations of Slovakia, Central Europe. Ticks Tick Borne Dis. 9(5), 1207–1211.

Stothard, D.R., Fuerst, P.A., 1995. Evolutionary analysis of the spotted fever and thyphus groups of *Rickettsia* using 16S rRNA gene sequences. Syst. Appl. Microbiol. 18(1), 52–61.

Sumrandee, C., Baimai, V., Trinachartvanit, W., Ahantarig, A., 2016. Molecular detection of *Rickettsia*, *Anaplasma*, *Coxiella* and *Francisella* bacteria in ticks collected from Artiodactyla in Thailand. Ticks Tick Borne Dis. 7(5), 678-689.

Swei, A., Kwan, J.Y., 2017. Tick microbiome and pathogen acquisition altered by host blood meal. ISME J. 11(3), 813–816.

Szigeti, A., Kreizinger, Z., Hornok, S., Abichu, G., Gyuranecz, M., 2014. Detection of *Francisella*-like endosymbiont in *Hyalomma rufipes* from Ethiopia. Ticks Tick Borne

Dis. 5(6), 818–820.

Tamura, K., 1992. Estimation of the number of nucleotide substitutions when there are strong transition-transversion and G+C-content biases. Mol. Biol. Evol. 9(4), 678–687.

Tessler, M., Neumann, J.S., Afshinnekoo, E., Pineda, M., Hersch, R., Velho, L.F.M., Segovia, B.T., Lansac-Toha, F.A., Lemke, M., DeSalle, R., Mason, C.E., Brugler, M.R., 2017. Large-scale differences in microbial biodiversity discovery between 16S amplicon and shotgun sequencing. Sci. Rep. 7(1), 6589.

Tomassone, L., Berriatua, E., De Sousa, R., Duscher, G.G., Mihalca, A.D., Silaghi, C., Sprong, H., Zintl, A., 2018. Neglected vector-borne zoonoses in Europe: Into the wild. Vet. Parasitol. 251, 17–26.

Turner, S., Pryer, K.M., Miao, V.P., Palmer, J.D., 1999. Investigating deep phylogenetic relationships among cyanbacteria and plastids by small subunit rRNA sequence analysis. J Eukaryot Microbiol 46(4), 327–338.

Vayssier-Taussat, M., Kazimirova, M., Hubalek, Z., Hornok, S., Farkas, R., Cosson, J.F., Bonnet, S., Vourch, G., Gasqui, P., Mihalca, A.D., Plantard, O., Silaghi, C., Cutler, S., Rizzoli, A., 2015. Emerging horizons for tick-borne pathogens: from the 'one pathogen-one disease' vision to the pathobiome paradigm. Future Microbiol. 10(12), 2033–2043.

Vilcins, I.-M., Fournier, P., Old, J.M., Deane, E., 2009a. Evidence for the presence of *Francisella* and spotted fever group *Rickettsia* DNA in the tick *Amblyomma fimbriatum* (Acari: Ixodidae), Northern Territory Australia. J. Med. Entomol. 46(4), 926–33.

Vilcins, I.M., Kosoy, M., Old, J.M., Deane, E.M., 2009b. *Bartonella*-like DNA detected in *Ixodes tasmani* ticks (Acari: Ixodida) infesting Koalas (*Phascolarctos cinereus*) in Victoria, Australia. Vector Borne Zoonotic Dis. 9(5), 499–503.

Vilcins, I.M., Old, J.M., Deane, E., 2009c. Molecular detection of *Rickettsia*, *Coxiella* and *Rickettsiella* DNA in three native Australian tick species. Exp. Appl. Acarol. 49(3), 229–242.

Wang, B., Wan, L., Wang, A., Li, L.M., 2017. An adaptive decorrelation method removes Illumina DNA base-calling errors caused by crosstalk between adjacent clusters. Sci. Rep. 7, 41348.

Wang, Y., Mao, L., Sun, Y., Wang, Z., Zhang, Jiayong, Zhang, Jibo, Peng, Y., Xia, L., 2018. A Novel *Francisella*-Like endosymbiont in *Haemaphysalis longicornis* and *Hyalomma asiaticum*, China. Vector Borne Zoonotic Dis. 18(12), 669–676.

Woods, R., Reiss, A., Cox-Witton, K., Grillo, T., Peters, A., 2019. The importance of wildlife disease monitoring as part of global surveillance for zoonotic diseases: the role of Australia. Trop. Med. Infect. Dis. 4(1), 29.

Zhang, X., Yang, Z., Lu, B., Ma, X., Zhang, C., Xu, H., 2014. The composition and transmission of microbiome in hard tick, I*xodes persulcatus*, during blood meal. Ticks Tick Borne Dis. 5(6), 864–870.