Systematic Identification of Novel Cancer Genes through Analysis of Deep shRNA

Perturbation Screens

Hesam Montazeri[1,2†], Mairene Coto-Llerena[2†], Gaia Bianco[2†], Ehsan Zangene[1], Stephanie

Taha-Mehlitz[3], Viola Paradiso[2], Sumana Srivatsa[4,5], Antoine de Weck[6], Guglielmo Roma[6],

Manuela Lanzafame[2], Martin Bolli[7], Niko Beerenwinkel[4,5], Markus von Flüe[7], Luigi M.

Terracciano[2], Salvatore Piscuoglio[2,3,7#] and Charlotte K. Y. Ng[2,8#]

[1]Department of Bioinformatics, Institute of Biochemistry and Biophysics, University of

Tehran, Tehran, Iran.

[2]Institute of Pathology, University Hospital Basel, Basel, Switzerland;

[3]Visceral surgery research laboratory, Clarunis, Department of Biomedicine, University of

Basel, Basel, Switzerland;

[4]Department of Biosystems Science and Engineering, ETH Zurich, Basel, Switzerland;

[5]SIB Swiss Institute of Bioinformatics, Basel, Switzerland;

[6]Novartis Institutes for BioMedical Research, Novartis Pharma AG, Basel, Switzerland;

[7]Clarunis, Department of Visceral Surgery, University Centre for Gastrointestinal and Liver

Diseases, St. Clara Hospital and University Hospital Basel, Switzerland

[8]Department for BioMedical Research, University of Bern, Bern, Switzerland.

†equal author contribution

#corresponding authors

Hesam Montazeri: hesam.montazeri@ut.ac.ir

Mairene Coto-Llerena: mairene.cotollerena@usb.ch

Gaia Bianco: Gaia.Bianco@usb.ch

Ehsan Zangene: e.zangene@ut.ac.ir

Stephanie Taha-Mehlitz: stephanie.taha@clarunis.ch

Viola Paradiso: viola.paradiso@usb.ch

Sumana Srivatsa: sumana.srivatsa@bsse.ethz.ch

Antoine de Weck: antoine.deweck@novartis.com

Guglielmo Roma: guglielmo.roma@novartis.com

Manuela Lanzafame: manuela.lanzafame@gmail.com

Martin Bolli: martin.bolli@claraspital.ch

Niko Beerenwinkel: niko.beerenwinkel@bsse.ethz.ch

Markus von Flüe: markus.vonfluee@clarunis.ch

Luigi M. Terracciano: Luigi.Terracciano@usb.ch

Salvatore Piscuoglio: s.piscuoglio@unibas.ch

Charlotte K. Y. Ng: charlotte.ng@dbmr.unibe.ch

Abstract

Background: Systematic perturbation screens provide comprehensive resources for the

elucidation of cancer driver genes, including rarely mutated genes that are missed by

approaches focused on frequently mutated genes and driver genes for which the basis for

oncogenicity is non-genetic. The perturbation of many genes in relatively few cell lines in

such functional screens necessitates the development of specialized computational tools

with sufficient statistical power.

Results: Here we developed APSiC (*A*nalysis of *P*erturbation *S*creens for *i*dentifying novel

*C*ancer genes) that can identify genetic and non-genetic drivers even with a limited number

of samples. Applying APSiC to the large-scale deep shRNA screen Project DRIVE, APSiC

identified well-known, pan-cancer genetic drivers, novel putative genetic drivers known to be

dysregulated in specific cancer types and the context dependency of mRNA-splicing

between cancer types. Additionally, APSiC discovered a median of 28 and 35 putative non-

genetic oncogenes and tumor suppressor genes, respectively, for individual cancer types,

including genes involved in genome stability maintenance and cell cycle. We functionally

demonstrated that *LRRC4B,* a putative non-genetic tumor suppressor gene that has not

previously been associated with carcinogenesis, suppresses proliferation by delaying cell

cycle and modulates apoptosis in breast cancer.

Conclusion: We demonstrate APSiC is a robust statistical framework for discovery of novel

cancer genes through analysis of large-scale perturbation screens. The analysis of DRIVE

using APSiC is provided as a web portal and represents a valuable resource for the

discovery of novel cancer genes.

Keywords: APSiC, perturbation screen, shRNA screen, cancer genes, *LRRC4B*, oncogene,

tumor suppressor.

Background

Advances in large-scale functional screening technologies have enabled the discovery of

gene requirements across diverse cancer entities [1,2]. Systematic perturbation screens

using short hairpin RNA (shRNA) or CRISPR are increasingly used to investigate how

genetic alterations or expression modulation of individual genes lead to phenotypic changes,

revealing novel factors in carcinogenesis [3–5]. In parallel, large-scale sequencing efforts of

10,000+ cancers have provided a comprehensive molecular portrait of human cancers and

their molecular pathogenesis [6]. Among the major findings is the unbiased discovery of

genes mutated at rates significantly higher than the expected background level [7], revealing

the global landscape of genetic 'driver genes' [8]. The discovery of these 'driver genes' forms

the critical foundations of cancer diagnostics, therapeutics, clinical trial design and selection

of rational combination therapies. Despite the large cohort size, the functional consequences

of mutations in rarely mutated genes, such as *YAP/TAZ* [9], are only revealed by functional

studies. Moreover, a systematic survey of non-genetic driver genes (i.e. driver genes for

which the basis for oncogenicity is non-genetic) is lacking.

In a recent large-scale perturbation screen, the project DRIVE (deep RNAi interrogation of

viability effects in cancer), 7,837 genes in 398 cancer cell lines were targeted with a median

of 20 shRNAs per gene across a variety of malignancies to generate a comprehensive atlas

of cancer dependencies [4]. The findings described in the study provided an overarching

view of the nature and types of cancer dependencies, but has barely scratched the surface

of the potential of the data generated. This huge resource of deep, robust and well-curated

functional data, in conjunction of molecular profiling from the Cancer Cell Line Encyclopedia

(CCLE) [10], remain largely untapped resources to be thoroughly mined and interrogated.

Analysis of shRNA perturbation screens is challenging due to the off-target effects of

shRNAs as well as low number of cell lines screened [4,11]. Project DRIVE used deep

coverage libraries to alleviate the off-target issues [4]. Additionally, several computational

tools have been developed to elicit the on-target effect from a pool of siRNAs that have both

on- and off-target effects [11–15]. RSA provides an absolute gene score based on the rank

distribution of phenotypes produced by all shRNA reagents of a given gene [14]. ATARiS

estimates a consensus shRNA profile for each gene and provides a relative gene score [13].

DEMETER is a regularized linear model for computing the effects due to knockdown of a

target gene by taking into account the effects due to the seed sequence [11], while

DEMETER2 is a complex hierarchical model structure that provides absolute gene

dependency score as opposed to ATARiS and DEMETER that provide relative dependency

score [15].

Cancer dependencies can be divided into two main groups of non-self dependencies (or

synthetic lethality) and self-dependencies. Synthetic lethality refers to the cell dependency

on the concomitant loss of two or more genes. In the context of an shRNA or CRISPR

screen, synthetic lethality refers to the loss of cell viability upon knockdown/knockout of a

gene conditional on the loss-of-function state of another gene. Various computational

approaches have been used to discover synthetic lethal pairs from perturbation studies [16–

18]. MiSL is a computational pipeline for the identification of synthetic lethal pairs using

Boolean implications applied to the The Cancer Genome Atlas (TCGA) datasets [19,20]. A

recent computational tool, called SLIdR, uses Irwin-Hall tests and causal inference for the

discovery of pan-cancer as well as cancer-specific synthetic lethal pairs using the DRIVE

data [21]. On the other hand, self-dependency refers to cell dependency on a gene with

specific molecular features such as mutation, copy number, expression, and DNA

methylation. Based on the molecular features, various statistical tests such as Fisher's Exact

and Wilcoxon tests, and differential expression analysis have been used to discover

association between molecular features and gene dependencies. Most existing methods for

finding self-dependencies are limited to the pan-cancer setting where a sufficient number of

cell lines is available [4]. These tests lack statistical power for a limited number of

observations (i.e. 5-10 observations) particularly after multiple testing corrections for a large

number of tests. While the project DRIVE is one of the largest of its kind, the number of cell

lines for some cancer types is quite small. A tool that can identify self-dependencies with

small sample size is needed to make the best use of perturbation screens to reveal the

cancer type-specific vulnerabilities.

Here, we introduce APSiC (*A*nalysis of *P*erturbation *S*creens for *i*dentifying novel *C*ancer

genes), a novel tool for the systematic and robust interrogation of large-scale perturbation

screens to discover gene (self-)dependencies for individual cancers even with limited

number of samples. Incorporating mutation and copy number status of the samples, APSiC

identifies potential genetic and non-genetic cancer genes. We consider three classes of

genetic drivers, namely mutation oncogenes, amplification oncogenes, and mutation tumor

suppressor genes as well as two classes of non-genetic drivers namely non-genetic

oncogenes and tumor suppressor genes. Of particular importance, non-genetic drivers

identified by APSiC have been largely less studied in cancer research. We applied APSiC to

26 cancers and identified both known and novel candidate genetic and non-genetic drivers.

As a proof of concept, we functionally validated *LRRC4B* as a putative tumor suppressor

gene in breast cancer. We provided the statistical analysis of DRIVE by APSiC as a web

portal (https://apsic.shinyapps.io/APSiC/) for the scientific community to explore and

functionally characterize genes that may be involved in carcinogenic processes and may

pave the way for the discovery of novel cancer-related biomarkers and drug targets.

Results and Discussion

The APSiC Algorithm

APSiC uses rank-based statistics to discover self-dependencies in perturbation screens (see

Methods for a technical description of the algorithm). Given the raw cell viability readout of a

perturbation screen, APSiC computes a rank profile for each gene by first ranking all genes

by their viabilities upon knockdown in a given sample to the range of [0, 1] then aggregating

the normalized ranks for a given gene across all samples (**Fig. 1a**). Thus ranks close to zero

represent reduced viability while the ranks close to one indicate cell growth upon

knockdown. Incorporating mutation and copy number status of the samples, APSiC identifies

potential genetic and non-genetic cancer genes by assessing deviation of the distribution of

normalized ranks from what is expected by chance using the Bates and Irwin-Hall tests. The

Irwin-Hall distribution has been successfully used in the identification of synthetic lethal gene

pairs [21] and prioritization of cancer genes based on multi-omics data [22]. The use of the

rank-based statistics with the Bates and Irwin-Hall distributions provides enhanced statistical

power when the number of cell lines is limited.

We consider three classes of genetic drivers (mutation oncogenes, amplification oncogenes,

and mutation tumor suppressor genes) and two classes of non-genetic drivers (non-genetic

oncogenes and tumor suppressor genes). We define mutation and amplification oncogenes

as genes for which reduced cell viabilities are preferentially observed in samples with

missense mutations and copy number amplifications, respectively, while mutation tumor

suppressor genes are those for which increased viabilities are preferentially observed in

samples with deleterious mutations. To identify such genetic drivers, we test, for a given

gene, whether ranks of the samples with and without the specific class of genetic alteration

are significantly different using a one-sided Bates test (**Fig. 1b, Additional files 1-3**). For

mutation and amplification oncogenes, we compute the lower-tailed *P* values (i.e. the ranks

preferentially suggest reduced viability upon gene knockdown), while for mutation tumor

suppressor genes, we compute the upper-tailed *P* values (i.e. the ranks preferentially

suggest increased cell viability upon knockdown). For the non-genetic drivers, we test

whether gene knockdown in samples without genetic alteration in the gene has any impact

on cell viability by computing lower and upper-tailed Irwin-Hall test *P* values for oncogenes

and tumor suppressor genes, respectively (**Fig. 1c, Additional files 4-5**). Optionally, we

further test whether the expression of candidate non-genetic oncogenes or tumor suppressor

genes is respectively enhanced or repressed in human tumors compared to the

corresponding normal tissue type.

APSiC identifies well-known and novel genetic cancer drivers

We applied APSiC to the DRIVE perturbation screens and the genetic data from the CCLE

[10] to identify genetic driver genes (**Additional files 1-3**). The dataset consists of 383 cell

lines across 26 cancer types, with a median of 11 (range 5-40) cell lines per cancer type

(**Fig. 2a**). In a pan-cancer analysis, APSiC reassuringly identified the well-known mutation

oncogenes *BRAF, CTNNB1, KRAS, NRAS, PIK3CA* and *TP53* (**Figs. 2b-c**, **Additional file**

6). Additionally, *DDX27, DCAF8L2* and *RBM39* were detected as mutation oncogenes

(**Additional files 6 and 7**). The top amplification oncogene were *KRAS, BRAF, CDK4,*

YAP1, IL6 and *HAS2* (**Figs. 2b, d and Additional file 6**), while the only mutation tumor

suppressor was *ARID1A* (**Figs. 2b, e and Additional file 6**). However, the identification of

mutation tumor suppressor genes in a knockdown screen is likely to have limited utility given

that mutation tumor suppressor genes are frequently associated with loss of the wild-type

allele.

One of the main strengths of APSiC is the identification of dependencies in small sample

sets. We therefore applied APSiC to the DRIVE data to identify genetic driver genes for

individual cancer types. Across the 26 cancer types, we found 20, 18 and 14 mutation

oncogenes, amplification oncogenes and mutation tumor suppressor genes, respectively

(**Fig. 2f** and **Additional files 1-3**), in 15, 14 and 11 cancer types. We found no correlation

between the number of cell lines and the number of driver genes identified (p=0.14, 0.21 and

0.97 for mutation oncogenes, amplification oncogenes and mutation tumor suppressor

genes, respectively, Spearman correlation tests). *KRAS*, *BRAF* and *TP53* were identified as

a mutation oncogene in 4, 2 and 2 cancer types, respectively, while the other 17 genes were

identified as mutation oncogenes in a single cancer type each. Other well described

mutation oncogenes include *NFE2L2* in esophagus carcinoma and *CTNNB1* in non-small

cell lung cancer. The remaining putative mutation oncogenes have not been reported as

frequently mutated in human cancers. These putative novel mutation oncogenes include

RREB1 (Ras Responsive Element Binding Protein 1) and *SBK1* (SH3 Domain Binding

Kinase 1), both of which have previously been found to be dysregulated in cancer [23,24].

No amplification oncogene or mutation tumor suppressor was identified in more than one

cancer type. We identified *MCL1* and *RHOB* as the top amplification oncogenes in the

squamous subtype of non-small cell lung cancer and glioma, respectively (**Fig. 2f** and

Additional files 2-3). *HOXA5*, a gene frequently overexpressed in hepatocellular carcinoma

[25], was identified as an amplification oncogene. Of the mutation tumor suppressor genes,

BRCA1 and *SETD2* were found in breast and kidney cancers, respectively. *BAZ2B*, a gene

involved in chromatin remodeling, is a putative mutation tumor suppressor in esophageal

cancer. Of note, 15% of *BAZ2B* somatic mutations in the TCGA pan-cancer cohort are

truncating mutations [8], suggestive of a tumor suppressor role for *BAZ2B*. Taken together,

APSiC identified both well known and putative genetic driver genes previously linked to

carcinogenesis.

Survey of non-genetic driver genes reveals cancer type specificity

By identifying strong irregular viability patterns in the rank profiles of cell lines wild-type for a

given gene using APSiC, we evaluated the non-genetic dependencies across cancer types

in the DRIVE data (**Additional files 4-5**). Consensus clustering of the most variable genes in

terms of APSiC *P* values revealed that such non-genetic dependencies segregate by organ

systems or cell-of-origin into four clusters (**Fig. 3a**). In particular, non-epithelial cancers

including leukemias/lymphomas, sarcomas, gliomas and neuroblastomas form a cluster

distinct from epithelial cancers including those of the lungs, the breasts and gastrointestinal

tract. This is consistent with the observation that multi-omics cancer classification is primarily

driven by cell-of-origin and anatomic regions [26]. Furthermore, the top-level segregation of

the cancer types was largely driven by the context-dependency of mRNA-splicing genes. We

observed that mRNA-splicing genes such as *PRPF6* (Pre-MRNA Processing Factor 6) and

SART3 (Spliceosome Associated Factor 3, U4/U6 Recycling Protein) were tumor

suppressive in the cluster enriched for non-epithelial cancers while they were oncogenic in

the epithelial cancer cluster (**Fig. 3b**). The context-dependency highlights the divergent role

of mRNA-splicing in carcinogenesis between cancer types. Our results also underscore the

necessity for an algorithm powerful enough to analyze perturbation screens for small

numbers of samples in a cancer type-specific manner.

Based on the DRIVE screen alone, we identified a median of 28 non-genetic oncogenes

(range 6-557) and 35 non-genetic tumor suppressor genes (range 1-471) per cancer type.

However, we reasoned that the many non-genetic onco- and tumor suppressor genes would

also be over- and under-expressed, respectively, in the corresponding cancer types. For the

12 cancer types for which gene expression data for the cancer and corresponding non-

cancer counterparts were available from the TCGA (**Additional file 8**), we further restricted

the putative non-genetic onco- and tumor suppressor genes to those that were over- and

under-expressed, respectively, relative to their non-cancer counterparts. After this filtering

step, there were a median of 13 non-genetic oncogenes (range 2-117) and 3 non-genetic

tumor suppressor genes (range 0-42, **Fig. 4**) per cancer type. We identified several well-

known oncogenes, including *CDK1* (a master regulator of cell cycle) and *SMC1A* (a

component of the cohesin complex involved in cell cycle checkpoint and genome

stability)[27], and some that have been shown to have oncogenic properties in some cancer

types, such as *MKI67IP* (or *NFIK*) [28]. We also identified *TEAD3*, a lesser described

member of the TEAD family involved in hippo signalling, as oncogenic in liver cancer [29].

Among the top candidate tumor suppressors were *FOXP2* in endometrial cancer and

XRCC5 in kidney carcinoma. *FOXP2* knockdown has been shown to promote tumor initiation

and metastasis in breast cancer [30] while *XRCC5*, encoding the protein Ku80, is a key DNA

damage repair protein. However, we also identified many genes that have not been

associated with carcinogenesis.

LRRC4B **is a putative tumor suppressor gene in breast cancer**

As a proof-of-concept to validate APSiC, we selected *LRRC4B,* the top putative non-genetic

tumor suppressor gene in breast cancer, for functional validation. A literature search of

LRRC4B in cancer suggests that its role and function in carcinogenesis are unknown.

Meanwhile, one of its paralogs *LRRC4* has been shown to have a putative tumor suppressor

role in glioma by modulating the extracellular signal-regulated kinase/protein kinase

B/nuclear factor-κB pathway [31]. In the DRIVE RNAi screen, nearly all breast cancer cell

lines displayed significantly increased cell viability upon *LRRC4B* knockdown and breast

cancers in TCGA showed lower *LRRC4B* expression compared to normal breast tissue

(**Additional file 9a**). We selected the breast cancer cell lines MDA-MB231, BT-549 and

MCF-7 with high, moderate and low endogenous *LRRC4B* expression to investigate whether

LRRC4B knock-down would result in other classical cancer phenotypes such as increased

migration and colony formation (**Additional file 9b**). We silenced *LRRC4B* in MDA-MB231

and BT-549 using siRNA, reducing LRRC4B protein expression by 40% and 60%,

respectively, 72 hours post-transfection (**Figs. 5a, e**). In both models, *LRRC4B*

downregulation significantly increased the proliferation and migration rates (**Figs. 5b-c, f-g**).

By contrast, *LRRC4B* overexpression significantly reduced proliferation and migration in

MCF-7 (**Figs. 5i-k**).

LRRC4 has been shown to suppress cell proliferation by delaying cell cycle in late G_1 phase

[32],[33]. To test whether *LRRC4B* may play the same role in breast, we analyzed cells with

LRRC4B overexpression or downregulation stained with DAPI by flow cytometry (FACS).

LRRC4B knockdown in MDA-MB231 and BT-549 promoted cell transition into S phase

(**Figs. 5d, h**), while *LRRC4B* overexpression in MCF-7 significantly retained cells in G_1

phase (**Fig. 5l**), suggesting a similar mechanism to *LRRC4*.

A common mechanism of oncogenicity is resistance to apoptosis [34]. To test whether

modulation of apoptosis is a mechanism of action of *LRRC4B* as an oncosuppressor, we

induced apoptosis with doxorubicin and measured it using Annexin V and propidium iodide

co-staining followed by FACS analysis (**Fig. 6a**). Forty eight hours after treatment, *LRRC4B*-

overexpressing MCF-7 cells showed 10% more apoptotic and 10% fewer live cells,

suggesting that *LRRC4B* overexpression could sensitize cells to doxorubicin-induced

apoptosis (**Fig. 6b**). By contrast, *LRRC4B*-downregulating MDA-MB231 and BT-549 cells

showed increased resistance to doxorubicin and had 25% and 10% fewer apoptotic and

25% and 10% more live cells, respectively (**Fig. 6b**). Our results provide compelling

evidence that APSiC identified *LRRC4B* as a novel oncosuppressor gene in breast cancer.

Conclusions

APSiC is a novel tool to enable systematic and robust discovery of gene dependencies from

small number of samples and can conceivably be applied to other types of perturbation

screens, such as CRISPR screens. Our analysis of the DRIVE perturbation screens using

APSiC is a valuable resource for the discovery of drug targets, cancer-related biomarkers

and novel cancer genes, in particular of non-genetic driver genes for which a systematic

analysis has been lacking. Our results would complement the associated multi-omics

profiling [35,36] and drug screens [37] to understand the vulnerabilities of cancer.

Methods

The detailed explanation of the APSiC algorithm

In this section, we give a more technical description of APSiC and also introduce a new

waterfall plot, called rank profile, for visualization of gene dependencies. First we briefly

describe some necessary definitions and background material from ordered statistics. We

consider the knockdown experiments of p genes across N cell lines. Let v_{ij} be viability of cell

line $i \in \{1,...,N\}$ upon knocking down gene $j \in \{1,..., p\}$ and m_{ij} be a binary variable indicating

whether a specific genetic alteration (i.e. mutation or copy number alteration) is present in

gene j of cell line i. In this study, we only consider deleterious (e.g. nonsense, frameshift,

splice site and mutations affecting start or stop codons) and missense mutations. Waterfall

plots are often used to show viabilities of knockdown experiments for a single gene across

different cell lines and are aimed to illustrate different gene dependencies. As an example,

waterfall plot for gene *TP53* is shown in **Fig. 1a** (left). Each vertical bar corresponds to a cell

line and is colored by the pre-existing mutation types present in *TP53*. **Fig. 1a** indicates cell

lines with the presence of deleterious or missense mutations in *TP53* tend to have lower

viabilities upon knockdown of this gene. While waterfall plot is a useful visualization tool for

demonstrating gene dependencies, it lacks sufficient interpretability in certain cases,

particularly when the number of cell lines is limited. In this paper, we introduce a new

waterfall plot, named rank viability profile or simply rank profile, to address this issue.

To make viability scores comparable across cell lines, we compute normalized rank values

per cell lines denoted as   , representing the rank of viability for gene *j* among all

knockdown experiments in cell line *i*. For mathematical convenience and without loss of

generality, we normalized ranks to the range of [0, 1]. When the number of knockdown

genes is high, normalized ranks have many distinct levels in the interval [0, 1] and we

assume normalized ranks are continuous. Let   denote random variables

associated to ranks of a gene *A* in *N* cell lines. We drop subscript A and denote ranks as

for the simplicity of notation. By placing ranks,   , in ascending orders and

renaming them, we obtain   where   is called *i*th ordered statistic. It is easy

to see that   and   . The probability density function of

ordered statistic   in general is given as

$$f(y_i) = N f(r) \binom{N-1}{i-1} F(r)^{i-1} (1-F(r))^{N-i}$$

where *f(r)* and *F(r)* denote probability density and cumulative distribution functions,

respectively. If there is no dependency between knocking down of a gene and the viability of

the cell, we can assume   for   , hence we have   .

Using this result, we can construct a no-change viability band at statistical significance *α*

using the quantiles of   at the   and   for   . Now we define a new

waterfall plot, called rank viability profile or simply rank profile, as a waterfall plot using

normalized ranks, realizations of   for a gene, overlaid with no-change viability band (**Fig.**

1a).

The APSiC algorithm identifies potential cancer genes by assessing deviation of respective

rank profiles from what is expected by chance. The algorithm can identify both genetic and

non-genetic drivers (**Fig. 1b-c**). We consider three categories for genetic drivers.

●   Mutation oncogene: defined as genes for which reduced viabilities are observed

349   preferentially in samples with missense mutation.

●   Amplification oncogene: defined as genes for which reduced viabilities are observed

preferentially in samples with copy number amplification.

●   Mutation tumor suppressor: defined as genes for which increased viabilities are

observed preferentially in samples with deleterious mutation.

We consider two categories for non-genetic drivers, namely

●   Non-genetic oncogene: defined as genes for which reduced viabilities are observed

in samples without a genetic alteration in the respective gene.

●   Non-genetic tumor suppressor: defined as genes for which increased viabilities are

observed preferentially in samples without a genetic alteration in the respective gene.

For genetic drivers, the APSiC algorithm considers rank profiles of mutated and wild-type

samples with respect to an input gene g (**Fig. 1b**). Then, it performs a one-sided statistical

test to determine whether rank scores of the two groups of samples are significantly different

in the direction of interest, according to the genetic feature of interest. Suppose

$R_1^{wt}, R_2^{wt}, \ldots, R_m^{wt}$ and $R_1^{mu}, R_2^{mu}, \ldots, R_n^{mu}$ are random variables denoting rank scores upon

knockdown of gene g for m wild-type and n mutated samples, respectively. Let $\underline{R^{wt}}$ and $\underline{R^{mu}}$

denote the average of ranks for the wild-type and mutated samples, respectively. We define

$S = \underline{R^{mu}} - \underline{R^{wt}}$ as the test statistic and s_{obs} as the observed test statistic. We assume the

null hypothesis is that the knockdown of gene g does not have any impact on the viability of

samples and therefore there is no difference in average of ranks for two groups, i.e. $S = 0$.

The general formula for the distribution of any weighted sum of uniform random variables is

given in Kamgar-Parsi, 1994[38]. We simplify the general formula thereby and obtain the null

distribution of the test statistic S as

$$P(S \leq s) = \frac{(-1)^m}{n^n m^m (n+m)!} \sum_{k=0}^{n} \sum_{p=0}^{m} (-1)^{p+k} \frac{n}{k} \frac{m}{p} (s + \frac{k}{n} - \frac{p}{m})^{m+n} \Theta(s + \frac{k}{n} - \frac{p}{m})$$

374 where

$$\theta(t) = \begin{cases} 0 & t < 0 \\ \dfrac{1}{2} & t = 0 \\ 1 & t > 0 \end{cases}$$

For mutation and amplification oncogenes, we compute lower-tailed P values,

while for mutation tumor suppressors, we compute upper-tailed P values,   . Due

to numerical issues, it is impractical to use the exact null distribution formula for large values

of m and n ($m+n$ >20). In this case, we compute an approximation for the null distribution of

S as follows. Under the null hypothesis, ___ and ___ follow $Bates$ distributions $Bates(m)$

and $Bates(n),$ respectively. The Bates distribution is a distribution that represents the mean

of a number of independent uniform random variables on the unit interval. For large values

of m and $n,$ ___ and ___ are approximately distributed by   – — and   – — . Hence,

under the null hypothesis the test statistic S is approximately distributed as   — — – .

To identify non-genetic drivers, we only consider the wild-type (i.e. without non-synonymous

mutations and without copy number amplification (GISTIC copy number state 2) or deep

deletions (GISTIC copy number state -2)) samples with respect to an input gene   (**Fig. 1c**).

The null hypothesis is that the knockdown of a gene g does not have any impact on the

viability of the samples. We define the test statistic as   and   as the

observed test statistic. Under the null hypothesis, T follows an Irwin-Hall distribution

392   , which represents the summation of   independent uniform random variables on

the unit interval. For large values of m, S is approximately distributed as   To

identify non-genetic oncogenes, we require significant lower-tailed P values,   for

wild-type cell lines with respect to the input gene. Additionally, for the respective tissue type,

the overall expression at the RNA level of a putative oncogene in tumor samples is required

to be significantly higher than the one in normal tissue samples using the t test. On the

contrary, for identifying non-genetic tumor suppressors, we require significant upper-tailed P

values, $P(T \geq t_{obs})$, for wild-type cell lines with respect to the input gene as well as lower

RNA expression of tumor samples in comparison to normal tissue samples using the *t* test.

Downloading and preprocessing of DRIVE and TCGA data

We considered the viability profiles of 383 cell lines in the project DRIVE [4] for which their

genetic profiles were available at the Cancer Cell Line Encyclopedia (**Fig. 2a**) [10]. We

computed aggregated gene-level viability scores for each experiment by the RSA and

ATARiS algorithms [13,14]. The RSA and ATARiS scores are available for 7726 and 6557

genes, respectively. We used the same method as defined in the project DRIVE to remove

essential genes, defined as genes with an RSA value of ≤ 3 in more than half of cell lines

[4]. After the removal of 184 essential genes by this method, for pan-cancer analysis, the

ATARiS scores are available for 6373 genes in 383 cell lines in 39 cancer types over 17

primary tissues. For the genetic drivers, we considered only genes for which there are at

least 2 samples harbored a genetic alteration of the corresponding class in individual cancer

types. For the non-genetic drivers, we only considered genes for which there are at least 2

samples wild-type for the gene in individual cancer types. For pan-cancer analysis, the

above threshold was at least 4 samples for both genetic and non-genetic drivers. For the

identification of mutation tumor suppressor genes, mutations annotated as *In_Frame_Ins*,

In_Frame_Del, *Frame_Shift_Ins*, *Frame_Shift_Del*, *Nonsense_Mutation*, *Splice_Site*,

Start_Codon_Del, *Stop_Codon_Del*, *Stop_Codon_Ins*, *Start_Codon_Ins* were considered

deleterious. For the analysis of individual cancer types, we considered the 26 cancer types

for which more than four cell lines are available in the DRIVE data.

TCGA gene expression data was obtained for 12 cancers for which data are available for

tumor and normal tissues (**Additional file 8**). The data were downloaded using the

TCGAbiolinks package in R [39]. The normalized expression level of genes in CPM (counts

per million) were used for the identification of non-genetic drivers.

Multiple testing

428 To address the multiple comparisons problem, we chose a significance level such that the

429 expected number of false positives due to multiple testing for each cancer and feature is

430 equal to one. To this end, we chose a significance level of $1/n$, or 0.05 if $1/n > 0.05$, where n

431 is the number of genes tested for identification of drivers. Using this approach, we were able

432 to keep many interesting hits while keeping the number of false positive cases low.

Clustering and pathway analysis

435 Clustering was performed using ConsensusClusterPlus Bioconductor package in R [40]

436 using 1-Spearman correlation as the distance metric and the Ward hierarchical clustering

437 algorithm. The number of clusters was determined based on the relative change in area

438 under the consensus cumulative distribution function over the number of evaluated clusters.

439 Pathway analysis was performed using g:Profiler [41].

Cell lines

Breast cancer derived cell lines (MCF-7, BT-549 and MDA-MB231) were maintained in a 5%

CO_2-humidified atmosphere at 37°C and cultured in DMEM supplemented with 10% FBS,

1% Pen/Strep (Bio-Concept) and 1% MEM-NEAA (MEM non-essential amino acids,

ThermoFisher Scientific). All cell lines were confirmed negative for mycoplasma infection

using the PCR-based Universal Mycoplasma Detection kit (American Type Culture

Collection, Manassas, VA) as previously described [42].

Transient gene knockdown by siRNAs

Transient gene knockdown was conducted using ON-TARGET plus siRNA transfection. ON-

TARGET plus SMARTpool siRNAs against human *LRRC4B* (Dharmacon, CO; #L-023786-

01-0005), ON-TARGET plus SMARTpool non-targeting control and DharmaFECT

transfection reagent (Dharmacon, CO; #T-2001-03) were all purchased from GE

Dharmacon. Transfection was performed according to the manufacturer's protocol. Briefly,

log-phase breast cancer cells were seeded at approximately 60% confluence. Because

antibiotics affects the knockdown efficiency of ON-TARGET plus siRNAs, growth medium

was removed as much as possible and replaced by antibiotic-free complete medium. siRNAs

were added to a final concentration of 25 nM. Cells were incubated at 37°C in 5% CO_2 for

24-48-72 hours for 48-72 hours for protein analysis. To avoid cytotoxicity, transfection

medium was replaced with complete medium after 8 hours.

Protein extraction and western blot

Proteins were extracted using Co-IP buffer (100 mmol/L NaCl, 50 mmol/L Tris pH 7.5, 1

mmol/L EDTA, 0.1% Triton X-100) supplemented with 1x protease inhibitors (cOmplete

Mini, EDTA-free Protease Inhibitor Cocktail, Roche, CO, #4693159001) and 1x phosphatase

inhibitors (PhosSTOP #4906837001, Merck). Cell lysates were then treated with 10x

reducing agent (NuPAGE Sample Reducing Agent, Invitrogen, #NP0009), 4x loading buffer

(NuPAGE LDS Sample Buffer, Invitrogen, #NP0007), boiled and loaded into neutral pH, pre-

cast, discontinuous SDS-PAGE mini-gel system (NuPAGE 10% Bis-Tris Protein Gels,

ThermoFisher). The proteins were then transferred to nitrocellulose membranes using

Trans-Blot Turbo Transfer System (Bio-Rad). The membranes were blocked for 1 hr with

Sure Block (Lubio Science) and then probed with primary antibodies overnight at 4°C. Next

day, the membranes were incubated for 1 hr at RT with fluorescent secondary goat anti-

mouse (IRDye 680) or anti-rabbit (IRDye 800) antibodies (both from LI-COR Biosciences).

Blots were scanned using the Odyssey Infrared Imaging System (LI-COR Biosciences) and

band intensity was quantified using ImageJ software. The ratio of proteins of interest/loading

control in treated samples were normalized to their counterparts in control cells. Antibodies

against LRRC4B (PA5-23529, Thermofisher) and B-actin (A5441, Sigma) were used at

dilution 1:1000 and 1:5000, respectively.

Proliferation assay

Cell proliferation was assayed using the xCELLigence system (RTCA, ACEA Biosciences,

San Diego, CA, USA) as previously described.[43] Cells were first seeded and transfected in

6 well plates and 24 h after transfection 5×10^3 cells were resuspended in 100 μl of medium

and plated in each well of an E-plate 16. Background impedance of the xCELLigence system

was measured for 12 s using 50 μl of room temperature cell culture media in each well of E-

plate 16. The final volume in each well was then 150 μl. The impedance signals were

recorded every 15 minutes until 96/120 h and expressed as cell index values, calculated

automatically and normalized by the RTCA Software Package v1.2. The values were defined

as mean ± standard deviation. Mann-Whitney test was used for statistical analysis with

GraphPad software.

Migration assay

Migration assays were performed using the CIM-plate of the xCELLigence Real-Time Cell

Analysis (RTCA, ACEA Biosciences, San Diego, CA, USA) system. Cells were first

transfected in 6-well plates and 24 h after transfection, they were harvested and seeded in

the CIM-plate. Every well of the bottom chamber was filled with 160 μl of the corresponding

medium at 10% FBS concentration. After placing the upper chamber on top of the lower

chamber, 50 μl of serum free medium was added on each CIM well for the background

measurement. After 3x PBS washing, 3×10^4 cells re-suspended in 100 μl of the

corresponding medium at 1% FBS concentration were seeded in each well of the upper

chamber. The measurements were taken every 15 minutes until 24 h after seeding and

expressed as cell index values. Mann-Whitney U test was used for statistical analysis with

GraphPad software.

Cell cycle analysis

Seventy-two hour after transfection, cells were collected, stained with DAPI and analyzed by

flow cytometry using the BD FACS Canto II cytometer (BD Biosciences, USA). Briefly cells

were harvested and washed 2X in PBS to get rid of serum proteins at 1200 rpm for 5

minutes. Pellets (up to 3x10^6 cells) were resuspended in 1.2 ml PBS (Ca and Mg free). For

crosslinking proteins 3.0 ml of 95% ice cold EtOH was added dropwise while vortexing. Cells

were fixed in this final 70% Et-OH solution for at least 30 minutes or over night. The Et-

OH/cell suspension was then diluted with 12 ml of PBS (for a total volume of 15 ml) and

centrifuge at 2000-2200 rpm for 10 min. Cells were then washed once more with 15 ml PBS

and then resuspended in 0.5-2.0 ml of DAPI stain solution (0.1% TritonX 100 and 10 ug/ml).

After 30 min of incubation on ice cells were analyzed by flow cytometry, measuring the

fluorescence emission at 461 nm. Data were analysed using the FlowJo software version

10.5.3 (https://www.flowjo.com).

Apoptosis analysis by flow cytometry

BT-549 and MDA-MB231 cells were transfected with siRNA (control or against *LRRC4B*)

and MCF-7 cells were transfected with *LRRC4B* overexpressing plasmid or control plasmid.

Eight hours after transfection medium was changed and doxorubicin added according to the

respective IC50 for each cell line[44][45]. Cells were collected 60 hours post siRNA

transfection or *LRRC4B* overexpression and 48 hours post treatment with doxorubicin

respectively, stained with annexin V (Annexin V-FITC conjugate; Invitrogen, CO; #V13242)

and propidium iodide (PI; Invitrogen, CO; #V13242), and analyzed by flow cytometry using

the BD FACS Canto II cytometer (BD Biosciences, USA). Briefly, cells were harvested after

incubation period and washed twice by centrifugation (1,200 g, 5 min) in cold phosphate-

buffered saline (DPBS; Gibco, CO; #14040133). After washing, cells were resuspended in

0.15 ml AnnV binding buffer 1X (ABB 5X, Invitrogen, CO; #V13242; 50 mM HEPES, 700 mM

NaCl, and 12.5 mM CaCl2 at pH 7.4) containing fluorochrome-conjugated AnnV and PI (PI

to a final concentration of 1 ug/ml) and incubated in darkness at room temperature for 15

min. As soon as possible cells were analyzed by flow cytometry, measuring the fluorescence

emission at 530 nm and >575 nm. Data were analysed using the FlowJo software version

10.5.3 (https://www.flowjo.com).

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The dataset supporting the conclusions of this article is included as **Additional Files 1-5**. A web portal using the Shiny framework in R has been developed to visualize rank profiles of the DRIVE shRNA screen and corresponding gene expression data from TCGA at https://apsic.shinyapps.io/APSiC/. The code for the APSiC algorithm is available at https://github.com/hesmon/APSiC/. The raw shRNA data has already published as a part of the project DRIVE (https://data.mendeley.com/datasets/y3ds55n88r/4) and copy number and mutation profiles of the cell lines are available at the Cancer Cell Line Encyclopedia portal (https://portals.broadinstitute.org/ccle/home). The gene expression data from the TCGA are available at the TCGA Genomics Data Commons data portal (https://portal.gdc.cancer.gov/).

Competing interests

A.dW. and G.R. are employees of Novartis Pharma AG. The other authors declare that they have no competing interests.

Funding

V.P. was supported by the Swiss Centre for Applied Human Toxicology. L.M.T. and S.P. were supported by the Swiss Cancer League (KLS-3639-02-2015 and KFS-3995-08-2016, respectively). S.P. was supported by Swiss National Science Foundation (PZ00P3_168165). N.B. was supported by European Research Council Synergy Grant 609883. The funders had

no role in study design, data collection and analysis, decision to publish, or preparation of

the manuscript.

Authors' contributions

H.M., S.P. and C.K.Y.N. conceived the study. S.P. and C.K.Y.N. supervised the study. H.M.

and C.K.Y.N. developed the methodology. H.M. performed the bioinformatic analyses. E.Z.

contributed to data presentation and visualization. H.M. and E.Z. implemented the Shiny web

application. M.C.-L., G.B., S.T.-M., V.P. and M.L. performed *in vitro* experiments. S.S.,

A.dW., G.R., M.B., N.B., M.vF. and L.M.T. critically discussed the results. H.M., M.C.-L.,

G.B., S.P. and C.K.Y.N. interpreted the results and wrote the manuscript. All authors agreed

to the final version of the manuscript.

Acknowledgments

References

1. Cheung HW, Cowley GS, Weir BA, Boehm JS, Rusin S, Scott JA, et al. Systematic
investigation of genetic vulnerabilities across cancer cell lines reveals lineage-specific
dependencies in ovarian cancer. Proc Natl Acad Sci U S A. 2011;108:12372–7.

2. Marcotte R, Brown KR, Suarez F, Sayad A, Karamboulas K, Krzyzanowski PM, et al.
Essential gene profiles in breast, pancreatic, and ovarian cancer cells. Cancer Discov.
2012;2:172–89.

3. Meyers RM, Bryan JG, McFarland JM, Weir BA, Sizemore AE, Xu H, et al. Computational
correction of copy number effect improves specificity of CRISPR-Cas9 essentiality screens
in cancer cells. Nat Genet. 2017;49:1779–84.

4. McDonald ER 3rd, de Weck A, Schlabach MR, Billy E, Mavrakis KJ, Hoffman GR, et al.
Project DRIVE: A Compendium of Cancer Dependencies and Synthetic Lethal Relationships
Uncovered by Large-Scale, Deep RNAi Screening. Cell. 2017;170:577–92.e10.

5. Marcotte R, Sayad A, Brown KR, Sanchez-Garcia F, Reimand J, Haider M, et al.
Functional Genomic Landscape of Human Breast Cancer Drivers, Vulnerabilities, and

Resistance. Cell. 2016;164:293–309.

6. Ding L, Bailey MH, Porta-Pardo E, Thorsson V, Colaprico A, Bertrand D, et al. Perspective
on Oncogenic Processes at the End of the Beginning of Cancer Genomics. Cell.
2018;173:305–20.e10.

7. Lawrence MS, Stojanov P, Mermel CH, Robinson JT, Garraway LA, Golub TR, et al.
Discovery and saturation analysis of cancer genes across 21 tumour types. Nature.
2014;505:495–501.

8. Bailey MH, Tokheim C, Porta-Pardo E, Sengupta S, Bertrand D, Weerasinghe A, et al.
Comprehensive Characterization of Cancer Driver Genes and Mutations. Cell.
2018;173:371–85.e18.

9. Wang Y, Xu X, Maglic D, Dill MT, Mojumdar K, Ng PK-S, et al. Comprehensive Molecular
Characterization of the Hippo Signaling Pathway in Cancer. Cell Rep. 2018;25:1304–17.e5.

10. Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The
Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity.
Nature. 2012;483:603–7.

11. Tsherniak A, Vazquez F, Montgomery PG, Weir BA, Kryukov G, Cowley GS, et al.
Defining a Cancer Dependency Map. Cell. 2017;170:564–76.e16.

12. Schmich F, Szczurek E, Kreibich S, Dilling S, Andritschke D, Casanova A, et al. gespeR:
a statistical model for deconvoluting off-target-confounded RNA interference screens.
Genome Biol. 2015;16:220.

13. Shao DD, Tsherniak A, Gopal S, Weir BA, Tamayo P, Stransky N, et al. ATARiS:
computational quantification of gene suppression phenotypes from multisample RNAi
screens. Genome Res. 2013;23:665–78.

14. König R, Chiang C-Y, Tu BP, Yan SF, DeJesus PD, Romero A, et al. A probability-based
approach for the analysis of large-scale RNAi screens. Nat Methods. 2007;4:847–9.

15. McFarland JM, Ho ZV, Kugener G, Dempster JM, Montgomery PG, Bryan JG, et al.
Improved estimation of cancer dependencies from large-scale RNAi screens using model-
based normalization and data integration. Nat Commun. 2018;9:4610.

16. Szczurek E, Misra N, Vingron M. Synthetic sickness or lethality points at candidate
combination therapy targets in glioblastoma [Internet]. International Journal of Cancer. 2013.
p. 2123–32. Available from: http://dx.doi.org/10.1002/ijc.28235

17. Jerby-Arnon L, Pfetzer N, Waldman YY, McGarry L, James D, Shanks E, et al.
Predicting cancer-specific vulnerability via data-driven detection of synthetic lethality. Cell.
2014;158:1199–209.

18. Sinha S, Thomas D, Chan S, Gao Y, Brunen D, Torabi D, et al. Systematic discovery of
mutation-specific synthetic lethals by mining pan-cancer human primary tumor data. Nat
Commun. 2017;8:15580.

19. Sahoo D, Dill DL, Gentles AJ, Tibshirani R, Plevritis SK. Boolean implication networks
derived from large scale, whole genome microarray datasets. Genome Biol. 2008;9:R157.

20. Sinha S, Tsang EK, Zeng H, Meister M, Dill DL. Mining TCGA data using Boolean
implications. PLoS One. 2014;9:e102119.

21. Srivatsa S, Montazeri H, Bianco G, Coto-Llerena M, Ng CKY, Piscuoglio S, et al.
Discovery of synthetic lethal interactions from large-scale pan-cancer perturbation screens
[Internet]. Available from: http://dx.doi.org/10.1101/810374

22. Dimitrakopoulos C, Hindupur SK, Häfliger L, Behr J, Montazeri H, Hall MN, et al.
Network-based integration of multi-omics data for prioritizing cancer genes. Bioinformatics.
2018;34:2441–8.

23. Wang P, Guo J, Wang F, Shi T, Ma D. Human SBK1 is dysregulated in multiple cancers
and promotes survival of ovary cancer SK-OV-3 cells. Mol Biol Rep. 2011;38:3551–9.

24. Kent OA, Fox-Talbot K, Halushka MK. RREB1 repressed miR-143/145 modulates KRAS
signaling through downregulation of multiple targets. Oncogene. 2013;32:2576–85.

25. Cillo C, Schiavo G, Cantile M, Bihl MP, Sorrentino P, Carafa V, et al. The HOX gene
network in hepatocellular carcinoma. Int J Cancer. 2011;129:2577–87.

26. Hoadley KA, Yau C, Hinoue T, Wolf DM, Lazar AJ, Drill E, et al. Cell-of-Origin Patterns
Dominate the Molecular Classification of 10,000 Tumors from 33 Types of Cancer. Cell.
2018;173:291–304.e6.

27. Yi F, Wang Z, Liu J, Zhang Y, Wang Z, Xu H, et al. Structural Maintenance of
Chromosomes protein 1: Role in Genome Stability and Tumorigenesis. Int J Biol Sci.
2017;13:1092–9.

28. Lin T-C, Su C-Y, Wu P-Y, Lai T-C, Pan W-A, Jan Y-H, et al. The nucleolar protein NIFK
promotes cancer progression via CK1α/β-catenin in metastasis and Ki-67-dependent cell
proliferation. Elife [Internet]. 2016;5. Available from: http://dx.doi.org/10.7554/eLife.11288

29. Perra A, Kowalik MA, Ghiso E, Ledda-Columbano GM, Di Tommaso L, Angioni MM, et
al. YAP activation is an early event and a potential therapeutic target in liver cancer
development [Internet]. Journal of Hepatology. 2014. p. 1088–96. Available from:
http://dx.doi.org/10.1016/j.jhep.2014.06.033

30. Cuiffo BG, Campagne A, Bell GW, Lembo A, Orso F, Lien EC, et al. MSC-regulated
microRNAs converge on the transcription factor FOXP2 and promote breast cancer
metastasis. Cell Stem Cell. 2014;15:762–74.

31. Wu M, Huang C, Gan K, Huang H, Chen Q, Ouyang J, et al. LRRC4, a putative tumor
suppressor gene, requires a functional leucine-rich repeat cassette domain to inhibit
proliferation of glioma cells in vitro by modulating the extracellular signal-regulated
kinase/protein kinase B/nuclear factor-kappaB pathway. Mol Biol Cell. 2006;17:3534–42.

32. Zhang Q-H, Wang L-L, Cao L, Peng C, Li X-L, Tang K, et al. Study of a novel brain
relatively specific gene LRRC4 involved in glioma tumorigenesis suppression using the Tet-
on system. Acta Biochim Biophys Sin . 2005;37:532–40.

33. Wang R, Wang Z, Yang J, Liu X, Wang L, Guo X, et al. LRRC4 inhibits the proliferation
of human glioma cells by modulating the expression of STMN1 and microtubule
polymerization. J Cell Biochem. 2011;112:3621–9.

34. Hanahan D, Weinberg RA. Hallmarks of Cancer: The Next Generation [Internet]. Cell.
2011. p. 646–74. Available from: http://dx.doi.org/10.1016/j.cell.2011.02.013

35. Ghandi M, Huang FW, Jané-Valbuena J, Kryukov GV, Lo CC, McDonald ER 3rd, et al.
Next-generation characterization of the Cancer Cell Line Encyclopedia. Nature.

2019;569:503–8.

36. Li H, Ning S, Ghandi M, Kryukov GV, Gopal S, Deik A, et al. The landscape of cancer
cell line metabolism. Nat Med. 2019;25:850–60.

37. Iorio F, Knijnenburg TA, Vis DJ, Bignell GR, Menden MP, Schubert M, et al. A
Landscape of Pharmacogenomic Interactions in Cancer. Cell. 2016;166:740–54.

38. Kamgar-Parsi B, Kamgar-Parsi B, Brosh M. Distribution and moments of the weighted
sum of uniforms random variables, with applications in reducing monte carlo simulations
[Internet]. Journal of Statistical Computation and Simulation. 1995. p. 399–414. Available
from: http://dx.doi.org/10.1080/00949659508811688

39. Colaprico A, Silva TC, Olsen C, Garofano L, Cava C, Garolini D, et al. TCGAbiolinks: an
R/Bioconductor package for integrative analysis of TCGA data [Internet]. Nucleic Acids
Research. 2016. p. e71–e71. Available from: http://dx.doi.org/10.1093/nar/gkv1507

40. Wilkerson MD, Hayes DN. ConsensusClusterPlus: a class discovery tool with confidence
assessments and item tracking. Bioinformatics. 2010;26:1572–3.

41. Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, et al. g:Profiler: a web
server for functional enrichment analysis and conversions of gene lists (2019 update).
Nucleic Acids Res. 2019;47:W191–8.

42. Ng CKY, Martelotto LG, Gauthier A, Wen H-C, Piscuoglio S, Lim RS, et al. Intra-tumor
genetic heterogeneity and alternative driver genetic alterations in breast cancers with
heterogeneous HER2 gene amplification. Genome Biol. 2015;16:107.

43. Andreozzi M, Quintavalle C, Benz D, Quagliata L, Matter M, Calabrese D, et al. HMGA1
Expression in Human Hepatocellular Carcinoma Correlates with Poor Prognosis and
Promotes Tumor Growth and Migration in in vitro Models. Neoplasia. 2016;18:724–31.

44. Pilco-Ferreto N, Calaf GM. Influence of doxorubicin on apoptosis and oxidative stress in
breast cancer cell lines. Int J Oncol. 2016;49:753–62.

45. Inao T, Iida Y, Moritani T, Okimoto T, Tanino R, Kotani H, et al. Bcl-2 inhibition sensitizes
triple-negative human breast cancer cells to doxorubicin. Oncotarget. 2018;9:25545–56.

Figure legends

Fig. 1. Overview of the APSiC algorithm. a, Illustration of the transformation from the raw cell

viability scores for a given gene (left, using *TP53* as an example) to the rank profile (right). Each bar

of the waterfall plots represents one sample and is colored by the mutational status of the given gene

in the sample. The red ellipse in the rank profile (right) represents a no-change (random) viability

band. **b-c,** Schematic representation of the APSiC algorithm for identifying **(b)** genetic and **(c)** non-

genetic drivers (See Materials and Methods for details).

Fig. 2. Analysis of the genetic drivers in the DRIVE perturbation screen. a, The number of cell

lines available for 26 cancer subtypes in the DRIVE perturbation screen. **b,** Kernel density estimation

of the P values (on a -$\log_{10}$ scale) for genetic drivers using the APSiC algorithm in a pan-cancer

analysis. Candidate mutation oncogenes, amplification oncogenes, and mutation tumor suppressors

identified by the APSiC are shown, with genes reaching significance level after multiple testing

corrections highlighted in red. **c,** Rank profile for a mutation oncogene (*BRAF*) colored by mutation

status. **d,** Rank profile of an amplification oncogene (*KRAS*) colored by copy number status. **e,** Rank

profile of a mutation tumor suppressor (*ARID1A*) colored by mutation status. **f,** Dot plots of the P

values (on a -$\log_{10}$ scale) for genetic drivers using the APSiC algorithm in a cancer type-specific

analysis, for (top) mutation oncogenes, (middle) amplification oncogenes and (bottom) mutation tumor

suppressors. Genes reaching significance level after multiple testing corrections are highlighted in

red. Cancer types are sorted by the number of cell lines.

Fig. 3: Pan-cancer analysis of the non-genetic drivers in the DRIVE perturbation screen. a,

Heatmap illustrates consensus clustering of the 500 most variable P values for non-genetic drivers

using the APSiC algorithm for the 26 cancer types in the DRIVE perturbation screen. Consensus

clustering identified 4 clusters across cancer types and 5 clusters across the 500 genes. One of the

clusters was enriched for genes involved in mRNA splicing and processing. **b,** Rank profiles for

PRPF6 in (top) non-Hodgkin B-cell lymphoma and (bottom) upper aerodigestive tract carcinoma and

for *SART3* in (top) PNET neuroblastoma and (bottom) breast carcinoma.

Fig. 4: Cancer type-specific analysis of the non-genetic drivers in the DRIVE perturbation

screen. Non-genetic driver genes identified in the 12 cancer types with corresponding gene

expression data from the TCGA. APSiC P values are shown (on a -$\log_{10}$ scale) for driver genes

significant after multiple testing corrections and over- or underexpressed in human cancers for

oncogenes (in red) and tumor suppressor genes (in blue), respectively. The top 20 genes for each

cancer type are labelled.

Fig. 5: *LRRC4B* has tumor suppressor-like properties in *in-vitro* models of breast cancer. a, e,

i, Western blotting showing LRRC4B protein level in (**a**) MDA-MB231, (**e**) BT-549 and (**i**) MCF-7 cell

lines 48 and 72 hours post transfection. Actin was used as a loading control and for normalization. **b,**

f, j, Proliferation kinetics of (**b**) MDA-MB231, (**f**) BT-549 and (**j**) MCF-7 cells upon (**b, f**)

downregulation or (**j**) upregulation of *LRRC4B* compared with the control. **c, g, k,** Migration potential

of (**c**) MDA-MB231, (**g**) BT-549 and (**k**) MCF-7 cells upon (**c, g**) downregulation or (**k**) upregulation of

LRRC4B compared with the control. **d, h, i,** Cell cycle analysis of (**d**) MDA-MB231, (**h**) BT-549 and (**i**)

MCF-7 cells upon (**d, h**) downregulation or (**i**) upregulation of *LRRC4B* compared with the control.

Error bars represent standard deviation obtained from three independent experiments. For all

experiments, statistical significance was assessed by multiple t-tests (* $P < 0.05$, ** $P < 0.01$, *** $P <$

0.001).

Fig. 6: *LRRC4B* **expression modulates induction of apoptosis by Doxorubicin. a,** Dot plot

illustrating the flow cytometry gating strategy used to assess cell viability and apoptosis using Annexin

V and propidium iodide staining of MDA-MB231, BT-549 and MCF-7 cells upon downregulation/

upregulation of *LRRC4B* compared with the control cells, with and without Doxorubicin. **b,**

Quantification of the mean (+/- SD) percentage of apoptotic cells (AnnV+) and live cells (AnnV-/PI-)

across the different groups. Error bars represent standard deviation obtained from three independent

experiments. For all experiments, statistical significance was assessed by multiple t-tests (* $P < 0.05$,

** $P < 0.01$, *** $P < 0.001$).

Additional files

Additional file 1: APSiC analysis of mutation oncogenes in DRIVE

Additional file 2: APSiC analysis of amplification oncogenes in DRIVE

Additional file 3: APSiC analysis of mutation tumor suppressors in DRIVE

Additional file 4: APSiC analysis of non-genetic oncogenes in DRIVE

Additional file 5: APSiC analysis of non-genetic tumor suppressors in DRIVE

Additional file 6: Significant genetic drivers identified by APSiC in the pan-cancer analysis

of DRIVE

Additional file 7: Rank profiles of *DDX27*, *DCAF8L2* and *RBM39* in DRIVE

Additional file 8: Pathological annotation in the DRIVE project and the corresponding TCGA

projects used for gene expression analysis.

Additional file 8: **a,** (left) Rank profile of the *LRRC4B* gene in breast cancer cell lines. Each

bar in the waterfall plots represents one cell line and is colored by the mutation status. (right)

LRRC4B transcript expression in breast cancers and normal tissues. The plot was generated

using gene expression data obtained from the TCGA dataset. **b,** Screening of LRRC4B

protein expression in a panel of breast cancer cell lines by western blot. Ratio of LRRC4B

expression relative to actin were calculated for each cell line. Ratio are shown above the

western blot plot. **c,** Dot plot showing the flow cytometry gating strategy used to assess cell

cycle status on breast cancer cell lines in both overexpressing and downregulating *LRRC4B*

cells. (*** *P* < 0.001).

Fig. 1

Fig. 2

Fig. 3

a

Thyroid_Carcinoma
Bladder_Carcinoma
CNS_Glioma
CNS_Glioma_HighGrade
Lung_Mesothelioma
Kidney_Carcinoma
Liver_HCC
Skin_Melanoma
Soft_Tissue_Sarcoma_Rhabdoid
PNET_Neuroblastoma
Leukemia_AML
Leukemia_ALL
Lymphoma_NH_B_cell
Lung_NSCLC_Others
Lung_NSCLC_Large_Cell
Pancreas_Carcinoma
Lung_NSCLC_Squamous
Upper_Aerodigestive_Tract_Carcinoma
Oesophagus_Carcinoma
Lung_NSCLC_Adeno
Ovary_Carcinoma
Breast_Carcinoma
Gastric_Carcinoma
Colorectal_Carcinoma
Endometrium_Carcinoma
Lung_SCLC

mRNA splicing

Tumor suppressive ◼️ Oncogenic

b

PRPF6

Non-Hodgkin B-cell lymphoma

Ranked Gene Score

WT

Upper aerodigestive tract carcinoma

Ranked Gene Score

WT

SART3

PNET neuroblastoma

Ranked Gene Score

WT

Breast carcinoma

Ranked Gene Score

WT

Fig. 4

Fig. 5

Fig. 6