Synergistic effect: a common theme in mixed-species litter decomposition

Jun Liu[1, 2] https://orcid.org/0000-0001-9311-1874, Xiaoyu Liu[2], Qingni Song[1, 3], Fenggang Luan[1], Hui Wang[4], Yalin Hu[2*], Qingpei Yang[1, 3*]

[1]*Jiangxi Provincial Key Laboratory for Bamboo Germplasm Resources and Utilization, Jiangxi Agricultural University, Nanchang 330045, China*

[2]*Stable Isotope Center, College of Forestry, Fujian Agriculture and Forestry University, Fuzhou 350002, China*

[3]*2011 Collaborative Innovation Center of Jiangxi Typical Trees Cultivation and Utilization, Jiangxi Agricultural University, Nanchang 330045, China*

[4]*College of Resources and Environment, Shandong Agricultural University, Taian 271018, China*

*Author for correspondence:

Yalin Hu,

Tel: +86 17720797616

E-mails: huyl@iae.ac.cn

Qingpei Yang

Tel: +86 13479161643

E-mails: qingpeiyang@126.com

Total word count (excluding summary, references and legends):	3913	No. of figures:	4 (Fig. 1, 2, 3, and 4 in colors)
Summary:	161	No. of Tables:	1
Introduction:	671	No of Supporting Information files:	6 (Fig. S1, S2, S3 Note S1, Table S1, S2)
Materials and Methods:	855		
Results:	599		
Discussion:	1571		
Acknowledgements:	56		

Abstract

• Litter decomposition plays a key role in ecosystem nutrients cycling, yet, to date

science is lacking a comprehensive understanding of the non-additive effect in

mixing litter decomposition.

• In order to fill that gap, we compiled 69 individual studies for the purpose of

performing two sub-meta-analyses on the non-additive effect.

• Our results show that a significantly synergistic effect occurs at global scale with

the average increase by 2-4% in litter mixture decomposition; In particular,

low-quality litter in mixture shows a significantly synergistic effect, while no

significant change is observed with high-quality species. Additionally, the

synergistic effect turns into the antagonistic effect when soil fauna is absent or

litter decomposition enters into humus-near stage. In contrast to temperate and

tropical areas, studies in frigid area also show a significantly antagonistic effect.

• Our meta-analysis provides a systematic evaluation of the non-additive effect in

decomposition mixed litters, which is critical for understanding and improving

the carbon forecasts and nutrient dynamics.

Keywords

Non-additive effect, litter mixture, litter quality, synergistic effect, litter

decomposition, meta-analysis

Introduction

Litter decomposition is a central component of biogeochemical cycles in ecosystems.

The rate of decomposition controls nutrients returns and energy flow, which regulates

atmospheric carbon emissions, soil organic matter composition, and supply of based

mineral nutrients to flora (Schneider *et al*. 2012). Thus, litter decomposition

influences ecosystem primary productivity (Bradford *et al.*, 2016). Over past few

decades, many studies about litter decomposition have focused on single litter decay

(Gartner & Cardon, 2004), which resulted in a profound exploration of the connection

between decomposition rate and its impact factors. However, in many ecosystems,

litter is generally a mix of multiple species. Previous studies suggest that those

various litter species interact with each other during their decomposition (Ball et al.

2008; Gessner *et al.,* 2010), implying that the decomposition rate of litter mixture is

different from that of single litter. To deeply discern how the litter mixture impacting

the decomposition is essential to understand the carbon and nutrients cycles.

When different litter species mix together, their decomposition rates generally do

not equal to the arithmetic mean value, i.e., the expected decay rate, between single

litter species (Fig. S1) (Gartner & Cardon, 2004; Steinwandter *et al.,* 2019). Instead,

one of two potential results usually develop: synergistic effect and antagonistic effect

(Collectively referred to as non-additive effect) (Fig. S1). In general, the chemical

characteristics of litter, decomposer activity, as well as other environmental factors

often constrain decomposition rate (Berg. B, 2014). In litter mixture, the underlying

mechanisms of synergistic effect are mainly from three aspects: the nutrients (such as

nitrogen) transfer from high-quality litter species to low-quality litter, the

complementary effects of soil fauna and decomposers, and the improvement of

microclimatic conditions (Schimel & Hattenschwiler, 2007; Tiunov, 2009; Madritch

and Cardinale 2007). Conversely, the antagonistic effect is often induced by the

enhancement of microbial nutrient immobilization for litter with poor nutrients or the

inhibitory secondary compounds released from low-quality litter species

(Hättenschwiler et al., 2005; Montané 2013).

Although experiments concerning the mixture effects on litter decomposition rates

have been conducted in numerous individual studies, the conflicting results have

hampered any possibility to draw general conclusions (Li *et al.,* 2016). Most of

conflicts could mainly result from the significant differences of influencing factors

related to litter decompositions in experimental design (Barbe et al. 2018, Leroy et al.

2018, Zhao et al. 2019). Thus far, those various studies on mixed decomposition were

all performed in different ecosystems across different climate regions with

decomposition durations varying from several weeks to even as long as several years.

Moreover, the various mesh sizes used in litterbag and microcosm, two methods that

the most commonly employed in litter decomposition, often make the results hard to

compare. As such, for the past several years, the scientific community has been

requesting a meta-analysis in order to determine a general global pattern (Gartner &

Cardon, 2004; Gessner *et al.* 2010).

While studies throughout the literature have summarized and analyzed the

non-additive effect in mixed litter decomposition (Gartner & Cardon, 2004; Li *et al.,*

2016), there still remains a lot of unanswered questions. In order to better understand

how the mixing influences the decomposition rate of litter, we employed a

meta-analysis that built upon the system analysis done by Gartner and Cardon (2004).

Finally, 70 individual studies were compiled to perform a global analysis (Note S1). If

the litter mixing effects occur, we also aimed to explore seven detailed questions

following:

1) How does the climate influence the effect of mixing on decomposition?

2) How does the ecosystem influence the effect of mixing on decomposition?

3) How does the mesh size of litterbag influence the effect of mixing on

decomposition?

4) How does the period of experiment influence the effect of mixing on

decomposition?

5) How do the evenness and richness of species influence the effect of litter mixing

on decomposition?

6) How does lignin content (AKA litter quality) influence the effect of mixing on

decomposition?

7) How do the leaf type (broad/needle) and habit (evergreen/deciduous) influence

the effect of mixing on decomposition?

Materials and Methods

Data compilation

Publications that reported data on litter mixture decomposition were selected from

the Web of Science resource, Google Scholar, and the China National Knowledge Infrastructure. Keyword search strings consisted of term combinations, such as (litter or debris or residue) AND (mix* or diversity or non-additive effect or additive effect) AND (decompos* or decay* or degrad*). The PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) flow diagram performed the procedure used for the selection of studies for meta-analysis (Fig S2).

Each selected publication had to satisfy the following three criteria: 1) it had to report at least one of our selected variables [expected decay rate (Rexp) vs. observed decay rate (Robs), or species-specific decay rate in mixture Rmix vs. in unmixed (Rsin)], and the decay rate should been expressed as mass loss or mass remaining (%); 2) it had to provide the means and sample sizes (n) of the variables selected for the meta-analysis or could be calculated from the chosen papers; 3) the measurements of selected variables were performed at the same temporal and spatial scale; and 4) except litter mixture treatment, the study of other experimental treatments (such as nutrient addition, warming, water controlling) was excluded. If the data were presented in figures, Getdata Graph Digitizer (version 2.24) was used to extract the numerical values. In addition, we sent Emails to corresponding authors to query the original data of k values (yr^{-1}) or relative mixture effect [(observed mass loss- expected mass loss)/ expected mass loss*100], when the selected variables used to estimate the non-additive effect were not provided in the papers.

We separated the dataset into two parts to perform two sub-meta-analyses (Fig. 1). The first sub-meta-analysis was to compare the decay rates of mixing litter between expected decay rates (Rexp) and observed decay rates (Robs). The second sub-meta-analysis was to compare the species-specific decay rates between single (Rsin) and mixture (Rmix) decomposition. As a note, the questions 1 to 5 were all answered by two sub-meta-analysis, the questions 6 and 7 were only answered by the second sub-meta-analysis. In total, 820 pair-observations in sub-meta-analysis 1 were obtained from 52 selected papers and 276 pair-observations in sub-meta-analysis 2

were obtained from 16 papers (Table S1). These studies were primarily conducted in

East Asia, Western Europe, and North America (Fig. S3).

For each study, we also noted the factors relevant to the mixing effects including

the site location, ecosystem type, species (and its nutrient contents if provided), decay

period, mesh size, response variables, and other background information (e.g. mean

annual temperatures (MAT), mean annual precipitation (MAP), etc.).

Statistical analysis

The natural log-transformed response ratio (R), defined as the effect size (Hedges *et*

al., 1999), was used as an index to measure the effect level of litter mixture on

decomposition rate:

$$\ln R = \ln\left(\overline{Xt}/\overline{Xc}\right) = \ln\left(\overline{Xt}\right) - \ln\left(\overline{Xc}\right) \qquad (1)$$

where $\overline{Xt}$ and $\overline{Xc}$ represent the Robs and Rexp, respectively; or the Rmix and Rsin,

respectively.

The variance of $\ln R$ (v) was calculated using the following formula:

$$v = \frac{S_t^2}{n_t \overline{X_t}^2} + \frac{S_c^2}{n_c \overline{X_c}^2} \qquad (2)$$

where S_t and S_c are the standard deviations (SDs) for the Robs and Rexp,

respectively, or for the Rmix and Rsin, respectively; n_t and n_c are the sample sizes for

the Robs and Rexp, respectively, or for the Rmix and Rsin, respectively. If both the

SD and standard error (SE) were lacking, we estimated the missing SD by multiplying

the average coefficient of variation (CV) from each data set by the reported mean

value (Wiebe *et al.,* 2006).

A nonparametric weighting function was used to weight each individual study; and

the mean effect size ($\overline{\ln R}$) of all observations was estimated according to the Eq. 3:

$$\overline{\ln R} = \frac{\sum_i \ln R_i}{\sum_i w_i} \qquad (3)$$

where w is the weighting factor used to calculate the inverse of the pooled variance

($1/v$); and $\ln R_i$ and w_i are the $\ln R$ and w of the ith observation, respectively.

To determine whether there was a significant difference under litter mixture

treatment (Rosenberg *et al.,* 2000), we employed a fixed-effect model to calculate 95%

confidence intervals (CIs) of the weighted effect size, using the Metawin 2.1 software.

The effect was only considered significant if the 95% CI values did not overlap with 0.

Furthermore, to clearly express the non-additive effects, the mean effect size was

converted back to the percent change, using the following equation:

$\text{The percent change} = (e^{\overline{\ln R_i}} - 1) \times 100\%$ (4)

In order to better understand the formation mechanisms of the non-additive effects,

we grouped the data according to climate zones (tropical, temperate, frigid),

ecosystem types (forest, shrubland, grassland, aquatic, peatland), mesh sizes [small

(diameter < 1 mm), medium (1mm≤diameter < 5) and large (diameter≥5mm)], and

decay periods (< 180 days, 180-360 days, 360-720 days, and >720 days) (Knorr et al.

2005). Finally, we also grouped the trees and shrubs based on different functional

types (broadleaf, needle, evergreen, and deciduous).

A continuous randomized-effect model was used to assess the potential linearity or

non-linearity between ln$R$ and climate factors (MAP and MAT)) or forcing factors

(decay period, and initial nutrients content). The total lnR' heterogeneity among the

selected studies (Q_T) was partitioned into different groups based on cumulative effect

sizes (Q_M) and the residual error (Q_E) (Rosenberg *et al.,* 2000).

Results

The mean effect size calculated across all the studies was significantly positive both

in Rexp vs. Robs and Rsin vs. Rmix, with an average increase of +2% and +4%,

respectively (Fig. 2a). When considering different climate zones, mixing litter

decomposition caused a significantly positive response in temperate areas and a

significant negative response in frigid areas in both of sub-meta-analyses (Fig. 2b).

Unlike temperate and frigid areas, in tropical area the Rexp was significantly higher

than Robs, and Rsin and Rmix differed little. The continuous randomized-effect

model suggested that MAT had a significant positive correlation with the mean effect

size in two sub-meta-analysises (Table 1). With respect to ecosystem types, the

mean effect sizes of litter mixing on decomposition rate showed a significant positive

response in Rexp vs. Robs for all five ecosystems except shrubland. (Fig. 2c).

The size level of the mesh in experimental methods resulted in different effects on

the decomposition rate (Fig. 3). When mesh size was divided into three

groups—small, middle, and large—and a marked antagonistic effect was found in

studies of small mesh size as expressed by Rexp vs. Robs. In contrast, the

decomposition rate in the studies of middle and large mesh sizes showed an obvious

synergistic response (Fig. 3a). For Rsin vs. Rmix, the small and large mesh sizes

depicted an additive effect, whereas, the middle mesh size showed a +6% increase in

decomposition rate (Fig. 3a). When the decay period was partitioned into four levels,

results showed that the decomposition rate increased under short-term duration (<180

days) and medium-term (180-360 days and 360-720 days), instead, a significant

antagonistic effect was observed under long-term duration (>720 days; -4%) in Rexp

vs. Robs (Fig. 3b).

In sub-meta-analysis 1 (Rexp vs. Robs), both of even litter mixtures and uneven

litter mixtures showed synergistic effects on decomposition rate (Fig. 3c). In

sub-meta-analysis 2 (Rsin vs. Rmix), the litter quality was divided into three levels

based on lignin content: low, medium, and high—and the results showed that the

decomposition rate of low-quality litter exhibits a greater positive change than the
middle and the high litter (Fig. 3d). When tree species were grouped into broad/needle
or evergreen/deciduous, results consistently showed synergistic effects. Besides, the
continuous randomized-effect model of meta-analyses showed a significant positive
correlation between litter initial P and the effect size of the sub-meta-analysis 2 (Rsin
vs. Rmix), yet, the remaining initial nutrients of litter did not show any correlation
(Table 1).

Table 1 Relationships between the effect size of litter mixing on decay rate and mean
air temperature (MAT), mean annual precipitation (MAP), experiment duration,
species richness, and litter initial nutrients content.

	Q_T	Q_M	Q_E	Slope	*P*-value
In sub-meta-analysis 1					
MAT	**575.37**	**28.88**	**546.49**	**0.004**	**<0.001**
MAP	543.89	27.87	516.02	<0.001	<0.001
Duration	1192.93	2.63	1190.31	<0.001	<0.001
Species richness	1047.87	2.27	1045.60	-0.008	0.13
In sub-meta-analysis 2					
MAT	**235.53**	**4.42**	**230.81**	**0.002**	**0.04**
MAP	269.57	0.39	269.18	<0.001	0.53
Duration	378.59	0.03	378.56	<0.001	0.86
Initial C	285.88	2.99	282.89	0.003	0.08
Initial N	312.32	0.52	311.80	-0.014	0.47
Initial P	**137.69**	**3.86**	**133.83**	**-0.896**	**0.05**
Initial Lignin	213.51	1.20	212.31	0.002	0.27
Initial C/N	282.12	0.14	281.98	<0.001	0.71
Initial C/P	131.48	0.18	131.29	<0.001	0.67
Initial N/P	132.68	0.19	132.49	0.001	0.66
Initial N/L	211.94	0.30	211.63	-0.141	0.58

Initial P/L	51.77	0.42	51.34	14.735	0.52

216 Statistical results were reported as total heterogeneity in effect sizes among studies (Q_T), the difference among

217 group cumulative effect sizes (Q_M), and the residual error (Q_E) from continuous randomized- effects model of

218 meta- analyses. The relationship is significant when $P < 0.05$.

Discussion

This study is primarily dedicated to calculate the general effects of litter mixture on its

decay rates. Our results showed that the litter mixture widely demonstrates a

non-additive effect, and most frequently synergistic effect, which is consistent with

two previous review studies (Gartner & Cardon 2004; Li *et al.,* 2016). Specifically,

the decay rates of litter mixture are on average 2%-4% faster when compared to that

of single litter species (Fig. 2a). This significant synergy is weak but logical. When

different litter species mixing, many processes (including the stimulating and

restraining) occur simultaneously with one counterbalancing the others more or less.

Based on our meta-analysis results, we further raised a conceptual model for the

non-additive effect of mixed litter samples of two species (Fig. 4). It should be noted

that the model reflects a general pattern, which may not be appropriated in all cases.

The variations of litter decomposers (including microorganisms and soil fauna)

play an important role in non-additive effect. Although our meta-analysis lacked

sufficient data to discuss the decomposer variations, the finding (mean effect size =

0.1604; 95% CI: 0.1432~0.1776, Table S2) suggested that microbial biomass is

significantly higher in litter mixture compared to single litter. In the decomposition of

litter mixture, on one hand, high-quality litter brings more available carbon to support

microbes growth (Hättenschwiler & Jørgensen 2010), on the other hand, microbial

activities and extracellular enzyme activities also increase simultaneously (Hu *et al*.

2006; Cong *et al*. 2015; Gogo *et al*. 2016), both of which can help the decay of

low-quality litter to generate synergistic effect.

When the data were divided into different climatic zones, an antagonistic effect of

litter mixture was observed in frigid areas in the two sub-meta-analysis (Fig.2b and 4b,

answer to Q1). Furthermore, a significant positive correlation between MAT and its

effect size on decomposition rate also supported this hypothesis (Table 1). This

phenomenon likely resulted from 1) soil fauna and microbial biomass being low in

high latitude regions compared with low latitude regions (Fierer *et al.,* 2009; Xu *et al.*

2013; Nielsen *et al.* 2014); 2) the soil organisms' activities being limited by lower

temperatures in high latitude areas. As mentioned before, soil fauna are regarded as an

important propelling force for the synergistic effect, once their quantity and activities

are substantially reduced, a synergistic effect seems unable to develop; and 3) the

fungi hyphae growth being limited by low temperature and less rain in frigid area

impeding nutrients transport from high-quality litter to the low-quality. When the data

were classified into different ecosystems, only the shrubland showed an antagonistic

effect (Fig.2c, answer to Q2), which would be because that most of the shrubland data

were acquired from frigid areas.

Soil fauna is an important group of decomposers but difficult to directly control in

the studies of litter decomposition. The size of mesh diameter is usually introduced to

distinguish different kinds of soil fauna. While a fine mesh (<1mm) is applied to

exclude most of the soil fauna, the antagonistic effect plays a leading role (Fig. 3a and

4b, answer to Q3). Interestingly, the antagonistic effect will turn into a synergistic

effect when middle (1-5 mm) or large mesh (>5 mm) is used. This emerging

conclusion was also drawn in a recent litter mixing study (Barbe *et al*. 2018).

High-quality litter with rich nutrients and energy would be palatable to soil fauna

(Zhang *et al*. 2016), which may accelerate the decay rate of litter mixture. A global

synthesis studies also suggested that the decay rate of litter fall by a third if the soil

fauna excluded (Zhang *et al*. 2015). Therefore, the existence of high-quality litter in

mixing litter decomposition can promote litter decay rate. In addition, the soil fauna

may make the litter more accessible to bacteria and fungi, which could stimulate the

microbial growth and therefore decomposition further (Smith & Bradford 2003). An

inexplicable phenomenon for the present is that a non-significant antagonistic effect

occurred when large mesh was used in sub-meta-analysis 2, which could be brought

by the error of insufficient data (35 observations). In order to determine whether these

results develop consistently, more extensive studies are required.

It is worth noting that the synergistic effect gradually weakens as a function of

decomposition time and turns into an antagonistic effect after 720 days of decay (Fig.

3b and 4a, answer to Q4). In the early stage of decomposition, the input of fresh litter

provides abundant food for the soil fauna and microorganisms. In addition, studies

suggest that nutrients transport frequently occurs in the early stage of decomposition

(Hansson *et al*., 2010; Liao *et al*., 2016). Thus, the rich food combined with

high-efficient nutrient transport may facilitate the growth and activities of soil

microorganisms, which ultimately produces a synergistic effect. As stated before, the

effects of soil fauna are important to the synergistic effect, however, their relative role

maybe be weaken and their effects maybe disappear at late stage of decomposition.

Consequently, the litter decomposition is mainly performed by the microbes that able

to degrade very recalcitrant compounds (such as lignin, tannin, etc.) (Guénon *et al*.

2017, Chapman *et al.* 2013). Besides, the remaining recalcitrant substances from

low-quality litter may resist the degradation at late stage of decay and subsequently

generate an antagonistic effect (Fig. 4).

Both of species evenness and richness influence the ecosystem functions, including

litter decomposition processes (Dangles & Malmqvist 2004; Tilman *et al*., 2014).

Although the species evenness depicts inconsistent effects on litter decomposition

(Hillebrand et al., 2008; Ward *et al*., 2010; Li *et al*., 2013), this meta-analysis

indicates that the synergistic effect of uneven litter mixtures is higher than that of

even mixtures (Fig. 3c, answer to Q5). We speculated that natural field values of

uniformity reflected in uneven mixtures more favorable for the microorganisms'

growth than in even mixtures (Swan *et al*. 2009). This opinion requires further

investigations to be confirmed. Our results also show a negative but insignificant

relationship between litter richness and its effect size on litter decomposition in

sub-meta-analysis 1 (Table 1, answer to Q5). Studies suggested that fungal diversity

increased with litter richness (Otsing *et al*. 2018), but we still do not know how fungi

change influences the decomposition of litter mixture. Although the underlying

reasons are unclear, we speculate that the following two points may contribute to this

phenomenon. First, the rate of litter decomposition might have a maximum value (Fig.

4) (Barbe *et al*. 2018), the non-additive effect would not be permitted to beyond this

maximal decomposition with litter richness. Second, the small size of data is not

enough to reveal the true rule.

With respect to the three classifications of litter species in sub-meta-analysis 2, a

noteworthy phenomenon is that a significant synergistic effect was observed in

low-quality litter species, while no significant change detected in medium and

high-quality litter species (Fig 3d and 4a, answer to Q6). Although the underlying

reasons are unclear, we assumed that the following three factors may contribute to this

result. First, the nutrients released from high-quality litter species promote low-quality

litter decay (Versini *et al*. 2016). Second, the input of high-quality litter species

promotes the growth of soil organisms (Hättenschwiler & Jørgensen 2010, Chapman

et al. 2013), thus leading to low-quality litter becoming competitive and accelerating

its decomposition. Third, the improved microenvironment through the addition of

high-quality litter, facilitates the decomposition of low-quality litter (He *et al.,* 2019).

High initial N and P content of litter indicate high decomposability. The strong

negative relationship between the mean effect size and litter initial N ($P > 0.05$) and P

($P < 0.05$) also support above result (Table 1). But we need more data to further clear

and definite this relation.

As the trees and shrubs were classified into four groups: broad, needle, evergreen,

and deciduous, the results showed that the mean effect size of needle and evergreen

groups were higher than those of broad and deciduous groups (Fig 3e, f, answer to

Q7). In general, the needle species contain more lignin than broadleaf species, and

they will act as a beneficiary of synergistic effect in litter mixing. Mean lignin content of evergreen leaves was 26% in this sub-meta-analysis database, which is higher than that of deciduous leaves (23%, Table S1).Moreover, the deciduous leaves contain more N and P than evergreen leaves based on the large scale research (Han *et al.* 2005), meaning the decomposability of deciduous leaves is greater than that of the evergreen leaf (Cornwell *et al.,* 2008; Liu *et al.,* 2016). The data indicates that the decomposition of low-quality litter species is more sensitive to the treatment of litter mixing, which is in agreement with Q6.

Conclusions

In summary, litter mixing generally increased the decay rate and tended to cause a synergistic effect compared to single litter. Moreover, when the mixing litter was separated, low quality-litter species displayed a synergistic effect, yet there was no change in high-quality litter species. A synergistic effect usually occurred at the early and late decay stage and disappeared at the humus-near stage. The soil organisms, especially soil fauna, were regarded as the important factors generating a synergistic effect. We suggest that, the synergistic effect and antagonistic effect, whose interplay give rise to the non-additive effect, occur simultaneously rather than independently. Whereas, some of our results have not been confirmed so far at the scale of a meta-analysis. Additional investigations are still needed to improve both the theory and the model in the near future.

Acknowledgements

We thank all the authors of the original studies included in our meta-analysis. The study was supported by the National Natural Science Foundation of China (41703065, U1805243, 41807028, 31460077), the Natural Science Foundation of Fujian Science and Technology Department (2018J01621), and the Education and Research Projects for Young Teachers of Fujian Provincial Education Department (JAT170187).

Author contribution

J. L, Y. H and Q. Y. designed the study. J. L conducted data checking. J. L and Q. Y. conducted the analyses. J. L and Y. H wrote the first draft of the manuscript. J. L., H. W., X. L., Q. S., F. L., Y. H., Q. Y. contributed to data collection, writing, and revision.

Supporting Information

Figure S1 The non-additive effect in mixing litter decomposition.

Figure S2 PRISMA diagram showing the process of locating studies include in this meta-analysis

Figure S3 World distribution of selected studies in this meta-analysis.

Note S1 List of all the references used in the meta-analysis.

Table S1 Summary of the references and data used in the meta-analysis of the effects of litter mixing on decomposition rate.

Table S2 Summary of the data and result used in meta-analysis of the effect of litter mixing on microbial biomass.

References

Ball, BA, Hunter MD, Kominoski JS, Swan CM, Bradford MA. 2008. Consequences of non-random species loss for decomposition dynamics: experimental evidence for additive and non-additive effects. *Journal of Ecology* **96**: 303-313.

Barbe, L, Mony C, Jung V, Santonja M, Bartish I, Prinzing A. 2018. Functionally or phylogenetically distinct neighbours turn antagonism among decomposing litter species into synergy. *Journal of Ecology* **106**: 1401-1414.

Berg B. 2014. Decomposition patterns for foliar litter – A theory for influencing factors. *Soil Biology & Biochemistry* **78**: 222-232.

Bradford MA, Berg B, Maynard DS, Wieder WR, Wood SA. 2016. Understanding the dominant controls on litter decomposition. *Journal of Ecology* **104**: 229-238.

Chapman SK, Newman GS, Hart SC, Schweitzer JA, Koch GW. 2013. Leaf Litter Mixtures Alter Microbial Community Development: Mechanisms for Non-Additive Effects in Litter Decomposition. *Plos One* **8**: e62671.

Gogo, S, Laggoun-Defarge F, Merzouki F, Mounier S, Guirimand-Dufour A, Jozja N, Huguet A, Delarue F, Defarge C. 2016. In situ and laboratory non-additive litter mixture effect on C dynamics of Sphagnum rubellum and Molinia caerulea litters. *Journal of Soils and Sediments* **16**: 13-27.

Cong, WF, Ruijven JV, Werf WVD, Deyn GBD, Mommer L, Berendse F, Hoffland E. 2015. Plant species richness leaves a legacy of enhanced root litter-induced decomposition in soil.

Soil Biology & Biochemistry **80**:341-348.

Cornwell WK, Cornelissen JHC, Amatangelo K, Dorrepaal E, Eviner VT, Godoy O, Hobbie

SE, Hoorens B, Kurokawa H, Pérez-Harguindeguy N *et al*. **2008.** Plant species traits are the

predominant control on litter decomposition rates within biomes worldwide. *Ecology Letters*

11:1065-1071.

Dangles O, Malmqvist B. 2004. Species richness–decomposition relationships depend on species

dominance. *Ecology Letters* **7**: 395-402.

Fierer N, Strickland MS, Liptzin D, Bradford MA Cleveland, CC. 2009. Global patterns in

belowground communities. *Ecology Letters* **12**: 1238-1249.

Gartner TB, Cardon ZG. 2004. Decomposition dynamics in mixed-species leaf litter. *Oikos* **104**:

230-246.

Gessner MO, Swan CM, Dang CK, McKie BG, Bardgett RD, Wall DH, Hättenschwiler. 2010.

Diversity meets decomposition. *Trends in Ecology and Evolution* **25**: 372-380.

Guénon R, Day TA, Velazco-Ayuso S, Gros R. 2017. Mixing of Aleppo pine and Holm oak litter

increases biochemical diversity and alleviates N limitations of microbial activity. *Soil Biology*

& Biochemistry **105**: 216-226.

Hansson K, Kleja DB, Kalbitz K, Larsson, H. 2010. Amounts of carbon mineralised and

leached as DOC during decomposition of Norway spruce needles and fine roots. *Soil Biology*

and Biochemistry **42**: 178-185.

Han W, Fang J, Guo D, Zhang Y. 2005. Leaf nitrogen and phosphorus stoichiometry across 753

terrestrial plant species in China. *New Phytologist* **168**: 377-385.

Hättenschwiler S, Tiunov, AV, Scheu, S. 2005. Biodiversity and litter decomposition in terrestrial

ecosystems. *Annual Review of Ecology, Evolution, and Systematics* **36**: 191-218.

Hättenschwiler, S, Jørgensen HB. 2010. Carbon quality rather than stoichiometry controls litter

decomposition in a tropical rain forest. *Journal of Ecology* **98**: 754-763.

He W, Ma Z, Pei J, Teng M, Zeng L, Yan Z, Huang Z, Zhou Z, Wang P, Luo Xin *et al*. **2019.**

Leaf Litter Decomposition in the Three Gorges Reservoir, China. *Forests* **10**: 360

Hedges LV, Gurevitch J, Curtis PS. 1999. The meta-analysis of response ratios in experimental

ecology. *Ecology* **80**: 1150-1156.

Hillebrand H, Bennett DM, Cadotte MW. 2008. Consequences of dominance: a review of

evenness effects on local and regional ecosystem processes. *Ecology* **89**: 1510-1520.

Hu, YL, Wang SL, Zeng DH. 2006. Effects of single Chinese fir and mixed leaf litters on soil

chemical, microbial properties and soil enzyme activities. *Plant and Soil* **282**: 379-386.

Knorr, M, Frey S, Curtis P. 2005. Nitrogen additions and litter decomposition: a meta-analysis.

Ecology **86**: 3252-3257.

Leroy, F, Gogo S, Buttler A, Bragazza L, Laggoun-Défarge F. 2018. Litter decomposition in

peatlands is promoted by mixed plants. *Journal of Soils and Sediments* **18**: 739-749.

Li D, Peng S, Chen, B. 2013. The effects of leaf litter evenness on decomposition depend on

which plant functional group is dominant. *Plant and Soil* **365**: 255-266.

Li YN, Zhou XM, Zhang NL, Ma KP. 2016. The research of mixed litter effects on litter

decomposition in terrestrial ecosystems. *Acta Ecologica Sinica* **36**: 4977-4987.

Liao S, Ni X, Yang W, Li H, Wang B, Fu C, Xu Z, Tan B, Wu F. 2016. Water, Rather than
Temperature, Dominantly Impacts How Soil Fauna Affect Dissolved Carbon and Nitrogen
Release from Fresh Litter during Early Litter Decomposition. *Forests* **7**: 249.

Liu C, Liu Y, Guo K, Zhao H, Qiao X, Wang S, Zhang L, Cai X. 2016. Mixing litter from
deciduous and evergreen trees enhances decomposition in a subtropical karst forest in
southwestern China. *Soil Biology & Biochemistry* **101**: 44-54.

Madritch MD, Cardinale BJ. 2007. Impacts of tree species diversity on litter decomposition in
northern temperate forests of Wisconsin, USA: a multi-site experiment along a latitudinal
gradient. *Plant and Soil* **292**: 147-159.

Montane F, Romanya J, Rovira P, Casals P. 2013. Mixtures with grass litter may hasten shrub
litter decomposition after shrub encroachment into mountain grasslands. *Plant and Soil* **368**:
459-469.

Nielsen UN, Ayres E, Wall DH, Li G, Bardgett RD, Wu T, Garey JR. 2014. Global-scale
patterns of assemblage structure of soil nematodes in relation to climate and ecosystem
properties. *Global Ecology and Biogeography* **23**: 968-978.

Otsing E, Barantal S, Anslan S, Koricheva J, Tedersoo L. 2018. Litter species richness and
composition effects on fungal richness and community structure in decomposing foliar and root
litter. *Soil Biology and Biochemistry* **125**: 328-339.

Rosenberg MS, Adams DC, Gurevitch J. 2000. *MetaWin: statistical software for meta-analysis.*
Massachusett, Sunderland: Sinauer Associates, Inc.

Schimel JP, Hattenschwiler S. 2007. Nitrogen transfer between decomposing leaves of different
N status. *Soil Biology & Biochemistry* **39**: 1428-1436.

Schneider T, Keiblinger KM, Schmid E, Sterflinger-Gleixner K, Ellersdorfer G, Roschitzki B,
Richter A, Eberl L, Zechmeister-Boltenstern S & Riedel K. 2012. Who is who in litter
decomposition? Metaproteomics reveals major microbial players and their biogeochemical
functions. *The ISME Journal* 6:1749-1762.

Smith VC, Bradford MA. 2003. Litter quality impacts on grassland litter decomposition are
differently dependent on soil fauna across time. *Applied Soil Ecology* **24**: 197-203.

Steinwandter M, Schlick-Steiner BC, Steiner FM, Seeber J. 2019. One plus one is greater than
two: mixing litter types accelerates decomposition of low-quality alpine dwarf shrub litter.
Plant and Soil **438**: 405–419

Swan CM, Gluth MA, Horne CL. 2009. Leaf litter species evenness influences nonadditive
breakdown in a headwater stream. *Ecology* **90**: 1650-1658.

Tilman D, Isbell F, Cowles JM. 2014. Biodiversity and Ecosystem Functioning. *Annual Review*
of Ecology, Evolution, and Systematics **45**: 471-493.

Tiunov AV. 2009. Particle size alters litter diversity effects on decomposition. *Soil Biology &*
Biochemistry **41**: 176-178.

Versini A, Laclau JP, Mareschal L, Plassard C, Diamesso LA, Ranger J, Zeller B. 2016.
Nitrogen dynamics within and between decomposing leaves, bark and branches in Eucalyptus
planted forests. *Soil Biology & Biochemistry* **101**: 55-64

Ward SE, Ostle NJ, McNamara NP, Bardgett RDJO. 2010. Litter evenness influences
short-term peatland decomposition processes. *Oecologia* **164**: 511-520.

Wiebe N, Vandermeer B, Platt RW, Klassen TP, Moher D, Barrowman NJ. 2006. A
systematic review identifies a lack of standardization in methods for handling missing variance
data. *Journal of clinical epidemiology* **59**: 342-353.
Xu X, Thornton, PE, Post, WM. 2013. A global analysis of soil microbial biomass carbon,
nitrogen and phosphorus in terrestrial ecosystems. *Global Ecology and Biogeography* **22**:
737-749.
Zhao W, Logtestijn RSPV, Hal JRV, Dong M, Cornelissen JHC. 2019. Non-additive effects of
leaf and twig mixtures from different tree species on experimental litter-bed flammability.
Plant and Soil **436**: 311-324.
Zhang WD, Yuan SF, Hu N, Lou YL, Wang SL. 2015. Predicting soil fauna effect on plant litter
decomposition by using boosted regression trees. *Soil Biology & Biochemistry* **82**:81-86.
Zhang W, Chao L, Yang Q, Wang Q, Fang Y, Wang S. 2016. Litter quality mediated nitrogen
effect on plant litter decomposition regardless of soil fauna presence. *Ecology* **97**: 2834-2843.

485
486
487

Figure 1 The two sub-meta-analyses in this study. (a) the sub-meta-analysis 1, comparison of mixing litter decay rate between expected rate (Rexp) and observed rate (Robs); (b) the sub-meta-analysis 2, comparison of species-specific decay rate between in single (Rsin) and in mixture (Rmix) decomposition. Rexp = w1R1+ w2R2+…+ wnRn, where wn is the weight of species n in the mixture and Rn is the decomposition rate of species n. The additive effect and non-additive effect are when "Rexp vs. Robs" or "Rsin vs. Rmix" are equal and not equal, respectively.

Figure 2 Comparison of mixing litter decay rates between expected values (Rexp) and observed values (Robs) (sub-meta-analysis 1, orange triangles), and comparison of species-specific decay rates between in single (Rsin) and in mixture (Rmix) decomposition (sub-meta-analysis 2, green circles) (a) across all studies, (b) among different climates, (c) ecosystems. If the mean effect size = 0, then means additive effect; If the mean effect size > 0, then means synergistic effect; If the mean effect size < 0, then means antagonistic effect. If the effect size 95% CI (error bars) did not cover zero, the non-additive effect was considered to be significant (*). The sample size for each variable is shown next to the point. The data on right-hand y-axis represent the mean percentage change for each variable (%).

Figure 3 Comparison of mixing litter decay rates between expected values (Rexp) and observed values (Robs) (sub-meta-analysis 1, orange triangles), and comparison of species-specific decay rates between in single (Rsin) and in mixture (Rmix) decomposition (sub-meta-analysis 2, green circles) among different experimental methods. (a) mesh size, (b) duration, (c)uniformity, (d) litter quality, (e) leaf types, and (f) leaf habit. If the mean effect size > 0, then means synergistic effect; If the mean effect size < 0, then means antagonistic effect. If the effect size 95% CI (error bars) did not cover zero, the non-additive effect was considered to be significant (*). The sample size for each variable is shown next to the point. The data on right-hand y-axis represent the mean percentage change for each variable (%).

Figure 4 Conceptual model for non-additive effect in mixed litter decomposition of two species based on our meta-analysis. The three-stages referred to (Berg 2014). (a) The non-addition effect in temperate and tropical areas; (b) The non-addition effect in frigid areas or absence of soil fauna.