Phylogenomic approaches reveal how a climatic inversion and glacial refugia

shape patterns of diversity in an African rain forest tree species

Andrew J. Helmstetter[1]*, Biowa E. N. Amoussou[1,2], Kevin Bethune[1], Narcisse G.

Kandem[3], Romain Glèlè Kakaï[4], Bonaventure Sonké[3], Thomas L. P. Couvreur[1,3,5]

[1] IRD, UMR DIADE, Univ. Montpellier, Montpellier, France

[2] Faculté des Sciences Agronomiques, Université d'Abomey-Calavi, 04 BP 1525,

Cotonou, Benin

[3] Université de Yaoundé I, Ecole Normale Supérieure, Département des Sciences

Biologiques, Laboratoire de Botanique systématique et d'Ecologie, B.P. 047,

Yaoundé, Cameroon

[4] Laboratoire de Biomathématiques et d'Estimations Forestières, Université

d'Abomey-Calavi, 03 BP 2819, Cotonou, Bénin

[5] Facultad de Ciencias Exactas y Naturales, Pontificia Universidad Católica del

Ecuador, Av. 12 de Octubre 1076 y Roca, Quito, Ecuador

* Corresponding Author

Email: andrew.j.helmstetter@gmail.com

Address:

UMR DIADE, équipe DYNADIV

Institut de recherche pour le développement (IRD)

911 avenue d'Agropolis, BP 64501

34394 Montpellier Cedex 5

France

ABSTRACT

The world's second largest expanse of tropical rain forest is in Central Africa and it

harbours enormous species diversity. Population genetic studies have consistently

revealed significant structure across central African rain forest plants, in particular a

North-South genetic discontinuity close to the equator at the level of a climatic

inversion. Here, we take a phylogeographic approach using 351 nuclear markers in

112 individuals across the distribution of the African rain forest tree species *Annickia*

affinis (Annonaceae). We show for the first time that the North-South divide is the

result of a single major colonisation event across the climatic inversion from an

ancestral population located in Gabon. We suggest that differences in ecological niche

of populations distributed either side of this inversion may have contributed to this

phylogenetic discontinuity. We find evidence for inland dispersal, predominantly in

northern areas, and variable demographic histories among genetic clusters, indicating

that populations responded differently to past climate change. We show how newly-

developed genomic tools can provide invaluable insights into our understanding of

tropical rain forest evolutionary dynamics.

Keywords: Phylogenomics, phylogeography, rain forest, sequence capture, Africa,

dispersal

Abbreviations

TRF = Tropical rain forest

CAR = central African rain forest

LGM = last glacial maximum

SFS = site frequency spectrum

1 INTRODUCTION

Tropical rain forests (TRFs) possess an incredibly diverse flora and fauna making up half of the world's biodiversity. Understanding how this diversity is generated is critical if we are to protect it (Mace et al. 2003). Central Africa hosts the world's second largest continuous extent of TRF (Linder 2001). Climatic fluctuations during the Pleistocene and associated glacial forest refugia are suggested to have influenced how genetic diversity is distributed in Central African rain forests (CAR) (Hardy et al. 2013). However, the nature (Anhuf et al. 2006; Diamond and Hamilton 1980; Maley 1996; Bonnefille 2007) and importance of forest refugia continue to be intensely debated (Cowling et al. 2008; Lezine et al. 2019). Population genetic studies within CAR plant species document differing levels of response to past climatic fluctuations (reviewed in Hardy et al. 2013). Conversely, one major phylogeographic pattern common to many CAR plant species studied is the existence of a phylogeographic barrier along a North-South axis around 0-3°N (see Fig. 1A; Hardy et al. 2013; Faye et al. 2016; Heuertz et al. 2014). There appears to be no visible geographic barrier to explain this break as continuous rain forest exists across the entire area. This North-South phylogeographic barrier corresponds, however, to the central African climatic hinge, an inversion zone between the northern and southern rainy seasons (Hardy et al; 2013). Interestingly, this barrier is rarely recovered in phylogeographic studies of

animals and thus seems to affect a greater effect on plants groups (e.g. Fuchs and

Bowie 2015; Bohoussou et al. 2015; Bell et al. 2017; Blatrix et al. 2017).

Three main hypotheses have been suggested to explain how this North-South genetic

discontinuity originated (Hardy et al. 2013). First, TRF might have disappeared

(repeatedly) along the climatic hinge during past climatic fluctuations, isolating

allopatric north/south populations, which subsequently recolonised the area. In this

case we would expect past climate models to show that suitable habitat disappeared

along the hinge during glacial periods when forests retract. Second, because seasons

are inverted across the hinge, flowering time might be displaced between northern and

southern populations preventing interbreeding and gene flow. If flowering occurs at

different times of the year either side of the inversion it may indicate this scenario

helped shape patterns in genetic structure. Third, successful colonisation across the

climatic hinge may be limited by factors such as environmental differences (e.g.

changing levels of water stress). If this was the case, we would expect ecological

niches of populations either side of the hinge to be significantly different. At present,

little is known about the relative importance of these three scenarios so we aim to

identify which, if any, have contributed to this common genetic discontinuity.

The phylogeographic approach (Avise et al. 1987) can unravel the history of

populations, and ultimately uncover which processes have shaped current patterns of

diversity. It is therefore an ideal framework to study how the climatic inversion, and

other factors, have shaped patterns of intraspecific diversity in CAR plants. High-

throughput sequencing allows the generation of phylogeographic datasets consisting

of large numbers of independently segregating nuclear loci that can account for

coalescent stochasticity (Edwards et al. 2016) where studies with small numbers of

markers fall short. Genomic data can also be used to reconstruct the spatial

evolutionary history of species by inferring phylogenetic trees among populations and

dispersal dynamics. These approaches can help us to understand relationships among

populations on either side of the climatic inversion and how often lineages traversed

this barrier.

If glacial refugia in areas proposed by Maley (1996) and Anhuf et al. (2006) have

played an important role in CAR plant dynamics we would expect to find evidence of

eastward dispersal because most putative CAR refugia are located in the Atlantic

Guineo-Congolian region. For example, within the palm species *Podococcus barteri*,

modelling past ranges and genetic data supported the hypothesis of one large coastal

refugia in western Gabon and southwestern Cameroon (Faye et al. 2016).

Furthermore, we would expect population size to increase towards the present as

populations spread out from climatically stable areas. If the climatic inversion acts as

a barrier, phylogeographic approaches should reveal limited numbers lineages that

dispersed across the region from 0-3°N. The nature of this barrier can then be

determined by a combination of phenological data and ecological niche modelling

approaches as detailed above.

To develop upon our current understanding of the phylogeographic patterns

introduced above, we present, for the first time using nuclear phylogenomic

approaches, the evolutionary dynamics of a central African tree species, *Annickia*

affinis. This species belongs to the pantropical plant family Annonaceae (Chatrou et

al. 2012) growing up to 30 metres tall and typically inhabits primary, secondary and

degraded rain forests (Versteegh and Sosef 2007). The species is widespread and

common across Lower Guinea, from southern Nigeria to the western tip of

Democratic Republic of Congo and is therefore ideal for studying CAR

phylogeography and the nature of the climatic inversion as a phylogeographic barrier.

Here, we used a newly developed baiting kit (Couvreur et al. 2019) to sequence

hundreds of nuclear markers in 112 individuals covering most of the distribution of *A.*

affinis. First, we identified the major genetic clusters within *A. affinis* and their

distribution to test if *A. affinis* shows a North-South genetic structuring along the

climatic inversion. Second, we built a phylogenetic hypothesis of relationships among

genetic clusters and conducted spatiotemporal diffusion analyses to test if lineages

have frequently crossed the climatic inversion, or if dispersal has been restricted over

time. We also test for an inland (eastwards) dispersal pattern, congruent with

expansion out of climatically stable areas. Third, we collated information on

flowering times and built ecological niche models (ENMs) of the species as a whole,

as well as populations either side of the climatic inversion, to identify the underlying

mechanism behind the common North-South genetic structure. Finally, we reconstruct

effective population size through time to infer the past demography of each identified

genetic cluster to test for recent population expansion, and if demographic histories

are congruent among clusters.

2 MATERIAL AND METHODS

2.1 Sample collection

A total of 112 individuals of *Annickia affinis* were sampled across most of the species

distribution range in Central Africa (Table S1). In addition, two individuals were

sampled from the sister species *Annickia polycarpa* as outgroups (Couvreur et al.

2019). We obtained data on flowering and fruiting times of *Annickia affinis* using

herbarium specimens of *A. affinis* retrieved from the BRAHMS database of Naturalis

NL. All specimens studied in the taxonomic revision of *Annickia* (Versteegh & Sosef,

2007) were included in BRAHMS, providing a good representation of available

specimens for the genus.

2.2 DNA extraction, gene capture and sequencing

DNA was extracted from silica gel dried leaves using the MATAB (Sigma-Aldrich,

Saint Louis, Missouri, USA) and chloroform separation methods following Couveur

et al. (2019). Nuclear markers were captured using the Annonaceae bait kit (Couvreur

et al. 2019) made of 11,583 baits 120 bp long targeting 469 exonic regions. Barcoded

Illumina libraries were constructed based on a modified protocol of Rohland and

Reich (2012). See supplementary methods for details.

2.3 Read filtering, contig assembly and multi-sequence alignment

Reads were cleaned and filtered following the protocol in Couvreur et al. (2019) and

Hybpiper (Johnson et al. 2016) was used to prepare sequence data for phylogenetic

inference. Alignments were conducted using MAFFT (Katoh and Standley 2013) and

cleaned with GBLOCKS (Castresana 2000). Putative paralogs for *A. affinis* that were

flagged by Hybpiper were verified and removed during this process. Further

information can be found in the supplementary methods.

2.4 SNP calling

We used SeCaPr (v1.1.4; Andermann et al. 2018) to call SNPs as it generates a

psuedoreference made up of the consensus sequences for each target locus (paralogs

removed) that is tailored to the given dataset, which is more efficient than the bait kit

reference made from distantly related Annonaceae species. We then mapped our

cleaned, paired reads to this psuedoreference using BWA (v0.7.12; Li and Durbin

2009). Duplicates were removed and SNPs were called using HaplotypeCaller in

GATK (v4.0; McKenna et al. 2010). We used bcftools (v1.8; Li 2011) to apply

thresholds of mapping quality (>40%) depth (>25), quality by depth (>2), minimum

depth across individuals (>10) to filter SNPs. We also removed those SNPs with a

minor allele frequency < 0.01, kept only biallelic SNPs and excluded monomorphic

sites.

182   2.5 Population genetic clustering and statistics

We examined the genetic structure of *A. affinis* using three approaches. First, we

undertook a Discriminant Analysis of Principal Components (DAPC) (Jombart et al.

2010). We used the function *find.clusters* in the R package 'adegenet' (Jombart 2008)

to identify clusters using successive K-means with 100,000 iterations per value of k

and a maximum k value of 20. We identified the most appropriate number of clusters

by examining the change in Bayesian Information Criterion (BIC) with increasing

values of k (number of clusters). We then used the function *dapc* in order to define the

diversity between the groups identified using *find.clusters*. We performed cross-

validation of our DAPC analysis to ensure our chosen number of PCs was reliable.

The data were divided into training (90%) and validation (10%) sets and DAPC was

carried out on the training set, with different numbers of PCs retained. 30 replicates

were performed to identify the optimal number of PCs – the number that minimizes

root mean squared error. We used cluster membership inferred using DAPC to define

populations for downstream analyses.

Second, we used TESS3 - an approach that takes into account geographic location

information when inferring population clusters (Caye et al. 2016). TESS3 was

implemented using the R package 'tess3r' (Caye et al. 2016). We used the projected

least squares algorithm and a maximum k of 20. We examined the cross-validation

score for each value of K to identify the appropriate number of clusters.

Third, we used fastSTRUCTURE (Raj et al., 2014) on a reduced set of unlinked SNPs

(one per locus). To maximize the number of SNPs we did not apply a depth-by-

individual filter for fastSTRUCTURE analyses. We then thinned our SNPs first by

applying more stringent filters (0% missing data) and then randomly choosing a single

SNP from each locus, where possible. We ran fastSTRUCTURE using the default

settings and the simple prior. The script 'chooseK.py' was used to identify the number

of clusters that maximized the marginal likelihood and explained the structure in the

data.

2.6 Phylogenetic inference

We filtered our dataset by choosing only those exons that had 75% of their length

reconstructed in 75% of *A. affinis* individuals. We then used the corresponding

supercontigs (i.e. targeted regions and surrounding off-target sequences) for

phylogenetic inference. We added empty sequences when individuals were missing

from locus alignments and we concatenated loci with the pxcat function from phyx

(Brown et al. 2017). We assigned a different GTR+GAMMA model to each locus to

account for differences in substitution rates. We then ran RAxML (v8.2.9)

(Stamatakis 2014) using the '-f a' option with 100 replicates. The tree was rooted

using *A. polycarpa* as outgroup. We also conducted a coalescent-based phylogenetic

analysis using ASTRAL-III (v5.5.11; Zhang et al. 2017), which uses individual gene

trees to infer a species tree.

We also constructed relationships using SNPs. We constructed a phylogenetic

network using splitstree (v4.14.6; Huson and Bryant 2006) and the full SNP dataset

using the neighbour-net algorithm. We used the coalescent-based Bayesian

phylogenetic approach SNAPP (Bryant et al. 2012) in BEAST2 (v2.5.2; Bouckaert et

al. 2019) to infer relationships among different genetic clusters. We used the same

SNP dataset as used with fastSTRUCTURE. We randomly chose 10 individuals per

DAPC genetic cluster (or all individuals if a cluster had <10) as representatives. We

ran the chain for 1 million generations, sampling every 1000 generations until

effective sample size (ESS) values, assessed with Tracer v1.7 (Rambaut et al. 2018),

were over 200 for all parameters. We performed two runs and combined these to

ensure that they reached stationarity at the same point. We then generated a consensus

tree using TREEANNOTATOR (Bouckaert et al. 2019).

2.7 Phylogeographic Diffusion in Continuous Space

We reconstructed the spatiotemporal dynamics of *A. affinis* using BEAST v1.8.4

(Drummond and Rambaut 2007) and spreaD3 v0.9.6 (Bielejec et al. 2011). As this

analysis is computationally intensive we used a subset of our dataset by selecting only

the five most informative loci based on number of phylogenetic informative sites. We

added a partition consisting of longitude and latitude coordinates as continuous trait

data. We used a HKY+G substitution model and a strict clock model for each genetic

locus and an exponential growth coalescent tree prior. We ran the analysis for 100

million generations and assessed ESS values using Tracer v1.7 (Rambaut et al. 2018).

We then used spreaD3 to visualize the output at several time points during the history

of *A. affinis*. We repeated this analysis with the next five most informative loci to

ensure similar patterns were recovered across datasets.

2.8 Demographic history

We used stairway plot (v2; Liu and Fu 2015), a model-flexible approach that uses site

frequency spectra (SFS) to infer changes in effective population size (N_e) through

time. We generated filtered VCF files representing each cluster as detailed above but

did not apply a minor allele frequency filter. We then calculated folded SFS for each

cluster. Stairway plot uses SNP counts to estimate the timing of events and changes in

N_e so the removal of SNPs with missing data may skew counts. To overcome this we

used an approach documented in Burgarella et al. (2018) where we modified each

SFS by first calculating the minor allele frequency at each SNP and then multiplying

this by the mean number of sequences (haploid samples) at each site. This results in a

new SFS that makes use of all observed site frequencies and minimizes the number of

SNPs removed. The total of samples is slightly reduced based on the amount of

missing data. The number of random breakpoints were calculated as recommended in

the manual. We used 67% of sites for training and performed 200 bootstrap replicates.

The number of observed sites was calculated as the total length of the

pseudoreference. We used an angiosperm wide mutation rate of 5.35×10^{-9} sites/year

(De la Torre et al. 2017) and a generation time of 15 years based on the generation

time of the Annonaceae species *Annona crassiflora* (Collevatti et al. 2014). In

addition, sequencing error can skew the SFS by inflating the number of singletons.

We ran analyses using the entire SFS, and then reran with singletons removed to

ensure similar histories were reached. We also ran analyses with a generation time of

50 years (Baker et al. 2014) to examine the effect this increase had on population size

change.

2.9 Modelling of current and past ranges

Current and Last Glacial Maximum (21k years ago; LGM) potential distributions

were modelled using MaxEnt (v3.3.3; Phillips et al. 2006) as implemented in

'biomod2' (v3.3-7.1;Thuiller et al.2009). Current and LGM (MIROC global

circulation model) climatic data were downloaded from WordClim ver. 1.4 (Hijmans

et al. 2005) at a resolution of 10*10 arc-minute. The LGM period represents the latest

unfavourable climate for tropical species and is therefore a good period to model the

impact of past climate change on potential range. A total of 346 presence data points

(Table S1) covering the known distribution of *A. affinis* were spatially filtered to one

point per cell to avoid overfitting due to sampling bias. Model overfitting was

constrained by using the β regularization parameter in Maxent, which limits model

complexity (Radosavljevic and Anderson 2014), and was set to 2.00 and 4.00, rather

than the default MaxEnt value of 1.00. Modelling with all 19 bioclim variables

produced unrealistic results and failed to properly model the current species range

independent of the regularization parameter (results not shown). Using just eight

bioclim variables (four precipitation and four temperature, see supplementary

methods) greatly improved the accuracy of the models to the known distribution.

Model performance was evaluated using a cross-validation procedure (Ponder et al.

2001, Muscarella et al. 2014, see supplementary methods). Model fit was assessed

using area under curve (AUC; Elith et al. 2006) and the true skill statistics (TSS,

Allouche et al. 2006). The best fitting model was then projected into the LGM.

We then constructed ecological niche models (ENMs) splitting them into two groups,

either North or South of the climatic inversion (above or below 2°N). We compared

these models using the R package 'ENMTools' (Warren et al. 2019). Models were

built using the generalized linear model (GLM) approach using the same bioclimatic

variables that were used in our MaxEnt models above. We performed niche

equivalency tests (100 replicates) using the metrics Schoener's D (Schoener 1968), I

(Warren et al. 2008) to assess whether ecological niches of individuals North and

South of the climatic inversion were more different than expected by chance. We also

performed symmetric background tests (100 replicates; Warren et al. 2008), to correct

for habitat availability. If observed values of D or I are significantly different than the

simulated distribution we can state whether the observed similarity between

populations is significantly larger or smaller than expected in the context of the

available environment. We then assessed ecological niche overlap in a phylogenetic

context using our SNAPP tree (100 replicates; Warren et al. 2008).

3 RESULTS

3.1 Sequencing

A total of 124.7 million reads were generated for 112 *A. affinis* individuals at an

average coverage depth of 77.5x across all targeted loci. Using HybPiper we

identified 366 loci where 75% of the exon length was recovered in at least 75% of

individuals. A total of 15 loci showed signs of paralogy and were removed, leading to

a final dataset of 351 supercontigs totalling 756 kb of sequence data. After cleaning

and filtering our SNP calling approach yielded 5,964 high-quality SNPs from 262

different loci. A comparison of the output of the SeCaPr and HybPipier pipeline

include amount of genetic variation and length of sequence recovered can be found in

Table S2. We found 240 loci that were reconstructed in both pipelines, representing

78% of the total number of SeCaPr loci (306) and 68% of HybPiper loci (351).

3.2 How are populations structured across the range of *A. affinis*?

After cross-validation the number of axes that was associated with the lowest

associated root mean squared error was 80 so this value was used in the DAPC.

Changes in BIC greatly decreased after $k = 4$ (Fig. S1) suggesting that four clusters

best fit our data (Fig. 1, Fig. S2). Two major clusters contained 35 and 63 individuals

that were located primarily in Western Gabon (cluster WG) and Cameroon (cluster

CA) respectively. Two smaller clusters of seven individuals each were located in

eastern Gabon (cluster EG) and Gabon / Republic of Congo (cluster GC). There is a

clear discontinuity in genetic structure across the equator, separating cluster CA from

the rest, except for a pair of individuals belonging to cluster EG.

The TESS3 analysis also found that four clusters best defined our data (Fig. 1B, S3)

with geographic discontinuities generally congruent between analyses (Fig. 1; Fig.

S4). Individual admixture proportions revealed generally limited mixed ancestry

within samples (Fig. 1B), though we detected a conspicuous amount at a single

location. The two northern most individuals belonging to cluster EG, found in at

Meyo Centre in Cameroon (labelled in Fig. 1A), had a considerable proportion of

their ancestry from cluster CA (Fig. 1B). The inverse was true of the two individuals

from cluster CA that were from the same location. The fastSTRUCTURE analysis

was similar to the aforementioned analyses, even with a reduced SNP dataset, though
the grouping of individuals belonging to DAPC clusters GC and EG was not
consistent for different values of K (Fig. S5). However, there was little evidence for
admixture when using this approach including for "Meyo Centre" individuals. These
differences are likely because of the relatively small numbers of SNPs used. Our
inferred phylogenetic network (Fig. 1C) also revealed four major clusters, and that
clusters WG & CA and EG & GC were grouped together.

3.3 How did populations of *A. affinis* disperse across central Africa?

The ASTRAL tree (Fig. 1D) was generally well-supported at deeper nodes with high
values of local posterior probability. This suggests that there are a high proportion of
gene trees supporting the topology in figure 1D, making it a reliable estimation of
evolutionary history among major clades. The tree topology reflected our clustering
inferences, lending further support to our four identified clusters and robust evidence
for phylogeographic structuring. Our SNAPP analyses reach stationarity after less
than 1 million generations and separate runs were combined as they converged at the
same point. The same pattern of divergence inferred for the four genetic clusters using
ASTRAL was found in our RAxML tree (Fig. S6) and our SNAPP tree (Fig. S7). The
phylogenetic network (Fig. 1C) also yielded relationships that supported two groups
of clusters: CA & WG and EG & WC.

We assessed the geographic locations in these clusters at a finer scale by mapping
each tip of the ASTRAL phylogenetic tree to its collection site (Fig. S8). In cluster
WG (Fig. S8C) the first of two major clades contains individuals that are found near
Mt. Cristal and in coastal rain forests in Northwest Gabon. The remaining individuals

in cluster WG formed a monophyletic group and are found to the South and East, as

far as the Southern tip of the Republic of Congo. We then examined the geographic

locations in cluster CA and identified three clades with distinct geographic

distributions (Fig. S8D) going up Cameroon's Atlantic coast. Individuals belonging to

the largest clade extend inland. Given this structure we repeated DAPC clustering

analyses using only individuals from cluster CA and revealed fine-scale genetic

structure that supported these three clades (Fig. S9).

Our diffusion analysis was based on 47.3kb of sequence data across five partitions

and converged with the ESS > 200 for all parameters after 100 million generations.

The root was inferred to be around central Gabon (Fig. 2A). We estimated a single

lineage crossed the climatic inversion, from South to North, establishing cluster CA

(Fig. 2). Late in the evolutionary history of *A. affinis* another dispersal event crossed

the barrier at Meyo Centre (see Fig. 1A). Our repetition of the diffusion analysis with

different loci matched these patterns (Fig. S10) indicating our results are reliable and

unlikely to have been biased by particular gene histories.

3.4 Do different populations share similar demographic histories?

We estimated the demographic history of the four DAPC clusters (Fig. 3; Fig S11-14

for full plots). Over the last 100 thousand years (Ka) three clusters (GC, WG and CA)

experienced similar demographic histories with population decline around 50-70 Ka

followed by an increase in N_e towards the present. We found this increase began at

slightly different times across these three clusters though all show rapid increases in

population size close to the LGM, around 20-27 Ka. Cluster CA showed evidence of a

rapid growth very close to the present, in the last 5 Ka. Cluster EG had a very

different history, exhibiting a relatively constant population size in the past with a

gradual decline in the last 10 Ka. Results were very similar when singletons were

removed indicating that sequencing errors were not affecting our analyses (results not

shown) and increasing the generation time to 50 years had little effect on the timing

and trend in population size change but did noticeably decrease N_e (Fig. S15).

3.5 Which areas have remained climatically stable over time?

A total of 113 data points were retained after filtration. The best predictors were

Precipitation of Wettest Month (Bio13) and Precipitation Seasonality (Bio15) (Table

S3). A regularization multiplier of 2 generated a better model fit than with 4, showing

a good visual match with the known distribution of the species at present (Fig. 4A).

The mean value of the AUC for the training and test data were respectively 0.77 and

0.76. The mean value of TSS was 0.454, indicating that the model is better than a

random model. During the LGM, the highest presence probabilities were all located

along the Atlantic coast in Cameroon, Equatorial Guinea and Gabon (Fig. 4B).

3.6 Does flowering time or ecological niche diverge across the climatic inversion?

We collated fruiting and flowering time data from 18 individuals (Table S4), 15 of

which had dates associated, and plotted their distribution by month (Fig. S16). This

revealed little change in flowering or fruiting time either side of the inversion.

Flowering and fruiting appears to occur all year round in the North while in the South

the majority of these events occurred from September to January (though individuals

here were also found to flower in April).

Comparisons of ENMs for individuals North and South of the climatic inversion

revealed large differences in ecological niches among these geographically separated

groups. Niche identity tests indicated that there are significant differences in niche

space of individuals North and South of the inversion ($p < 0.01$; Fig. 4C, D). Our

symmetrical background tests (Fig. 4E, F) revealed that correlation between the

available habitat between these areas is moderate-high, but niche overlap is

significantly lower than would be expected based on available habitat ($p < 0.05$).

When we examined niche overlap in a phylogenetic context, we found that overlap

was smallest between clusters CA and WG (Fig. S7A), which corresponds to

divergence across the climatic inversion.

4 DISCUSSION

4.1 Limited dispersal and divergent niches across the climatic hinge

Intraspecific diversity, based on phylogenomic nuclear sequence data, within the

widespread tree species *Annickia affinis* is highly structured with a clear North-South

divide between identified genetic clusters (Fig. 1). This is the first time this has been

observed in plants using genomic data and adds to the growing evidence of an

important phylogeographic barrier around a climatic hinge between 0-3°N in

numerous CAR distributed plants (Hardy et al. 2013; Heuertz et al. 2014; Faye et al.

2016; Ley et al. 2017; Pineiro et al. 2017). This North-South discontinuity is,

however, generally not recovered in CAR distributed animals except in rare cases

(e.g. Portik et al. 2017). This suggests that the processes taking place in relation to

this barrier affect the flora of CARs more than the fauna. Indeed, Blatrix et al. (2017)

showed that this barrier was more abrupt in the studied tree species (*Barteria*

fistulosa) than within the associated symbiotic ants. However, the reasons for this

genetic break in a seemingly continuous rain forest region remain little understood

(Hardy et al. 2013).

Here, we show that, throughout the evolutionary history of *A. affinis*, a single major

northward cross-hinge colonisation event occurred leading to the successful

establishment of the Cameroonian population (Fig. 2C). Hardy et al. (2013) posited

that environmental differences either side of the inversion may limit or prevent

successful dispersal and establishment. The rarity of lineage dispersal across the hinge

that we observed lends support to this hypothesis. It may not be environmental

constraints that limit dispersal, but instead the mechanism of dispersal itself.

However, the small red to black fleshy fruits of *Annickia affinis* (Versteegh & Sosef,

2007) are frugivore-dispersed (Poulsen et al. 2001; Holbrook and Smith 2000) and

can potentially travel long distances (> 500 m) for example by hornbills. Thus, the

genetic structure of *A. affinis* in general, and the North-South divide in particular, is

not linked to seed dispersal limitation *per se*. Our ENM comparisons give further

support to the environmental differences hypothesis - we found that ecological niches

were significantly different between individuals North and South of the inversion

(Fig. 4C-F). Furthermore, when viewed in a phylogenetic context, ecological niches

of clusters CA and WG showed much smaller values of niche overlap than values at

other nodes in our population-level tree (Fig. S7A).

We detected that one genetic cluster (cluster EG) extends across the climatic hinge

into south Cameroon (Meyo Centre site, Fig. 1A) leading to a second, more recent,

northwards migration event into the climatic hinge area (Fig. 2D). This indicates that

the barrier is not absolute, agreeing with other studies (Hardy et al. 2013; Duminil et

al. 2015; Pineiro et al. 2017). The Meyo Centre site lies within the inversion zone and

several individuals with mixed ancestry are found here, including those in the second

dispersal event (Fig. 1A, B). A similar result was found in *B. fistulosa*, with 20% of

individuals sampled near 1°N being backcrossed or F_1 individuals (Blatrix et al.

2017). Aside from these few individuals, we found a general lack of evidence for

admixture between genetic clusters on either side of the inversion (Fig. 1B; Fig. S5),

indicating that gene flow is generally rare.

The hypotheses of Hardy et al. (2013) were not mutually exclusive and given that

there is no clear barrier to dispersal of pollen or seeds, divergence in traits such as

flowering time could play a role in preventing reproduction and limiting North-South

admixture. This may in turn be linked to why successful establishment across the

inversion is rare. However, we observed no clear difference in flowering time among

individuals north and south of the climatic inversion (Fig. S16). We note that data was

available for just15 individuals, so more is needed to reliably assess whether

flowering time differs.

Interestingly, similar north-south phylogeographic breaks are also known from the

Atlantic rain forests of Brazil, due to differing climatic regimes and floral

compositions (Carnaval et al. 2014; Leite et al. 2016). This suggests that similar

processes, though not necessarily driven by exactly the same factors, might be driving

patterns of intraspecific diversity in different TRF regions.

4.2 Out-of-refugia migration in northern forests

The refuge hypothesis has received support from several population genetic studies of

CAR plants, showing concordance between putative refugia and regions of high or

unique allele/haplotype diversity (Lowe et al. 2010; Dauby et al. 2014; Heuertz et al.

2014; Faye et al. 2016). Here we find that the inferred evolutionary dynamics of *A.*

affinis support a potential role for Pleistocene forest oscillations in shaping

intraspecific genetic diversity patterns, but this role may not have been the same in

each region. we found evidence for recent demographic expansion in three clusters

(CA, GC and WG, Fig. 3), as would be expected if *A. affinis* expanded out of refugia.

These expansions were estimated to have over the last 40 Ka, around the timing of the

LGM. This timing was fairly robust to changes in generation time used (Fig. S15).

However, further work is needed to determine a more accurate mutation rate and

generation time for *A. affinis* to verify the timing of these events, and we would

therefore caution against over-interpreting the exact timing of demographic events

and instead focus on the population size trends. Sampling sizes were also small (n =7)

for clusters GC and EG meaning we are less confident in the patterns reconstructed

for these clusters. Similar patterns of recent expansion were detected in populations of

central African plants (Pineiro et al. 2017) and animals (Bell et al. 2017) as well as in

studies on neotropical flora (Vitorino et al. 2016) and fauna (Batalha-Filho et al.

2012).

In contrast to the other clusters, EG showed constant population size with a slight

decline towards the present, perhaps indicating that glacial cycles have not played an

important role in its demographic history. Similar demographic patterns were found in

populations of two central African *Erythrophleum* species (Duminil et al. 2015),

though these exhibited a more pronounced decline in the last 50 Ka. Overall, our

results indicate that demographic responses to past climate change have been different

among populations of *A. affinis* across central Africa. Similar patterns of recent

expansion were detected in populations of central African plants (Pineiro et al. 2017)

and animals (Bell et al. 2017) as well as in studies on neotropical flora (Vitorino et al.

2016) and fauna (Batalha-Filho et al. 2012).

We also find evidence to support the scenario presented by Anhuf et al. (2006) who

proposed that coastal rain forests in central Africa acted as refugia during the LGM,

though results again vary among regions. The modelled LGM distribution of *A. affinis*

indicates that suitable habitat was concentrated continuously along the coast, from

Cameroon to Gabon (Fig. 4B), like in the understory palm species *Podococcus barteri*

(Faye et al. 2016). In addition, we uncovered fine-scale genetic structure and evidence

for within-population dispersal eastwards in Cameroon (Fig. 2), demonstrating a

possible out-of-refugia pattern in this area. By contrast, we did not find an inland

pattern of migration in Gabon where dispersal was both eastwards and westwards

from a more central area. This may be a result of there having been a large area with

highly-suitable conditions (>0.8) during the LGM that extended further from the coast

in Gabon than in Cameroon (Fig. 4), meaning that populations could persist and

expand out of this area. In addition, we inferred more pronounced East-West

clustering (Fig. 1) in Gabon than in Cameroon, which has been observed in at least

four other CAR tree species (Hardy et al. 2013).

The spatial diffusion approach has been shown to recover the true ancestral locations

under a generative model that is the same as used for inference (Lemey et al. 2010).

Related categorical approaches known as "mugration" discrete trait analysis (DTA)

have received criticism and been largely replaced by structured coalescent approaches (De Maio et al. 2015). These issues may be applicable to continuous diffusion but no equivalent spatially-explicit coalescent implementation exists. As for any model, violations of its assumptions need to be considered and sampling bias is likely to be an issue for reconstructing the history of migration (Lemey et al. 2014; De Maio et al. 2015). The ways that this may affect a continuous mode are not well characterised and warrant further work. Given that our populations, in particular EG and GC, may have been underrepresented we must interpret our results with caution.Finally, the four retrieved clusters are found in allopatry or parapatry (Fig. 1A). This supports the hypothesis of incomplete mixing after post-glacial expansion and is similar to patterns found in other CAR species (Hardy et al. 2013) and within species from other TRF regions (Carnaval et al. 2009; Leite and Rogers 2013). Bringing our results together, it appears that refugia may have played a different role for populations in different areas, and that each has responded to past climate in change in its own way.

5 CONCLUSIONS

This study uncovered the evolutionary dynamics and demographic history of the CAR tree species *Annickia affinis*. Our approach is the first to use genome-wide data from hundreds of nuclear loci to infer population-level phylogeographic patterns in CAR plants. We highlighted how a climatic inversion limits lineage dispersal, drives ecological niche divergence and shapes patterns of population structure across a continuous rain forest region. We also show that the current distribution of extant populations is the result of different demographic histories and, in northern regions, migration eastwards from putative refugia in coastal rain forests. Our study is an example of how taking advantage of recently developed genomic resources, such as

the sequence capture kit used here, can help to improve our understanding of TRF

evolution, at the population level and above.

FUNDING

This study was supported by the Agence Nationale de la Recherche (grant number

ANR-15- CE02-0002-01 to TLPC).

ACKNOWLEDGEMENTS

We thank Prof. Moutsambote, Raoul Niangadouma, Théophile Ayole for help in the

field. The authors acknowledge the IRD itrop HPC (South Green Platform) at IRD

Montpellier for providing HPC resources. We are grateful to the Centre National de la

Recherche Scientifique et Technique (CENAREST), the Agence National des Parques

Nationaux (ANPN) and Prof. Bourobou Bourobou for research permits (AR0020/16;

AR0036/15 (CENAREST) and AE16014 (ANPN). Fieldwork in Cameroon was

undertaken under the "accord cadre de cooperation" between the IRD and Ministère

de la Recherche Scientifique et Technique (MINRESI). Prof. Bouka Biona of the

Institut National de Recherche en Sciences Excates et Naturelles (IRSEN) of the

Republic of Congo is thanked for research permits.

Data availability

The fastq (R1 and R2) sequences for all individuals are available in Genbank SRA

under Bioproject number PRJNA508895

(http://www.ncbi.nlm.nih.gov/bioproject/508895). Scripts for bioinformatics can be

found at https://github.com/ajhelmstetter/afrodyn

Competing interest statement

The authors of this article declare that they have no financial conflict of interest with

the content of this article. TLPC is one of the *PCIEvolBiol* recommenders.

FIGURES

Fig. 1. (A) Map of the study region showing genetic clusters inferred using

Discriminant analysis of principal components (DAPC, k=4). Individuals are colour

coded by cluster membership. Superimposed upon the map are the locations of

putative glacial refugia (adapted from Faye et al. 2016). The climatic hinge is shown

by a dashed red line. Inset is a map of the African continent showing highlighting the

study area. (B) Barplot of ancestry proportions inferred using TESS (k=4). Colours

were made to correspond to those in Fig. 1A as clustering was almost identical

between approaches. A single individual, "A_affinis_Ndjole_5", was inferred as

cluster EG (red) in DAPC but TESS suggests the majority of its ancestry is instead

from cluster WG (blue). Individuals from the "Meyo Centre" show evidence of

admixture between clusters (red and purple) across the North-South climatic

inversion. (C) Phylogenetic network among *A. affinis* individuals constructed in

splitstree using NeighbourNet algorithm based on 6787 SNPs. (D) ASTRAL tree

613   representing relationships among *Annickia affinis* samples, rooted on two *A.*

614   *polycarpa* samples. Local posterior probabilities are shown as pie charts on nodes and

615   tips are coloured based on genetic clustering results.

Fig. 2. Phylogeographic diffusion analysis split into four time slices. Images were

rendered using spreaD3 and move forward through time starting from the

(uncalibrated) time of the most recent common ancestor (A) to the present day (D).

White circles represent ancestrally estimated geographic locations for nodes in the

inferred phylogenetic tree, as well as current, real locations at tips. Polygons around

points represent uncertainty of estimated ancestral locations at 80% highest posterior

density (HPD). Putative refugia following Maley (1996) are shown in dashed blue

lines.

628   **Fig. 3.** Plots of effective population size through time for each of the four clusters

629   inferred using stairway plot. The present is located on the left side of each graph. The

630   dotted line represents the median population size and the shaded polygon represents

631   the 80% central posterior density intervals. Colours correspond to the colours used in

632   figure 1. Full plots of each species can be found in figures S11-14.

Fig. 4. Ecological niche models (ENMs) for the present (A) and projected into the

past, during the last glacial maximum (B). ENMs were constructed using MaxEnt and

bioclimatic variables. The colour scale represents habitat suitability for each cell

where green indicates more suitable cells. Red circles in (A) indicate sites where *A.*

affinis individuals were collected and used in building the model. Comparisons of

ENMs constructed from individuals North and South of the climatic inversion (see

641   panel A) are shown in panels (C-F). Panels (C) and (E) show histograms of 100

642   simulations of identity tests using two different metrics, D and I. Similarly, (E) and

643   (F) are background tests, taking into account the available environments when

644   comparing models. Values of D and I range from 0 (no niche overlap) to 1 (identical

645   niches among the two groups). Dotted lines indicate observed values.

REFERENCES

Allouche O, Tsoar A, Kadmon R. 2006 Assessing the Accuracy of Species
Distribution Models: Prevalence, Kappa and the True Skill Statistic (TSS).
Journal of Applied Ecology 43(6):1223–32. https://doi.org/10.1111/j.1365-2664.2006.01214.x.

Andermann, T, Cano Á, Zizka A, Bacon C, and Antonelli A. 2018 SECAPR—a
Bioinformatics Pipeline for the Rapid and User-Friendly Processing of
Targeted Enriched Illumina Sequences, from Raw Reads to Alignments.
PeerJ 6:e5175. https://doi.org/10.7717/peerj.5175.

Anhuf D, Ledru M-P, Behling H, Da Cruz FW, Cordeiro RC, Van der Hammen
T, Karmann I et al. 2006 Paleo-Environmental Change in Amazonian and
African Rainforest during the LGM. *Palaeogeography, Palaeoclimatology,
Palaeoecology* 239(3–4):510–27.
https://doi.org/10.1016/J.PALAEO.2006.01.017.

Avise, JC, Arnold J, Ball RM, Bermingham E, Lamb T, Neigel JE, Reeb CA,
and Saunders NC. 1987. Intraspecific Phylogeography: The Mitochondrial
DNA Bridge Between Population Genetics and Systematics. *Annual Review
of Ecology and Systematics* 18(1):489–522.
https://doi.org/10.1146/annurev.es.18.110187.002421.

Baker TR, Pennington RT, Magallon S, Gloor E, Laurance WF, Alexiades M,
Alvarez E, Araujo A, Arets, EJ, Aymard G and De Oliveira AA. 2014 Fast
demographic traits promote high diversification rates of Amazonian trees.
Ecology letters 17(5):27-536.

Batalha-Filho H, Cabanne GS, Miyaki CY. 2012 Phylogeography of an Atlantic
Forest Passerine Reveals Demographic Stability through the Last Glacial

Maximum. *Molecular Phylogenetics and Evolution* 65(3):892–902.

https://doi.org/10.1016/j.ympev.2012.08.010.

Bell, RC, MacKenzie JB, Hickerson MJ, Chavarría KL, Cunningham M,

Williams S, and Moritz C. 2011 Comparative Multi-Locus Phylogeography

Confirms Multiple Vicariance Events in Co-Distributed Rainforest Frogs.

Proceedings of the Royal Society B: Biological Sciences 279(1730):991–99.

https://doi.org/10.1098/rspb.2011.1229.

Bielejec F, Rambaut A, Suchard MA and Lemey P. 2011 SPREAD: spatial

phylogenetic reconstruction of evolutionary dynamics. *Bioinformatics*

27(20):2910-2912.

Blatrix R, Peccoud J, Born C, Piatscheck F, Benoit L, Sauve M, Djiéto-Lordon C

et al. 2017 Comparative Analysis of Spatial Genetic Structure in an Ant–

Plant Symbiosis Reveals a Tension Zone and Highlights Speciation

Processes in Tropical Africa. *Journal of Biogeography* 44(8):1856–68.

https://doi.org/10.1111/jbi.12972.

Bohoussou KH, Cornette R, Akpatou B, Colyn M, Peterhans JK, Kennis J,

Šumbera R et al. 2015 The Phylogeography of the Rodent Genus

Malacomys Suggests Multiple Afrotropical Pleistocene Lowland Forest

Refugia. *Journal of Biogeography* 42(11):2049–61.

https://doi.org/10.1111/jbi.12570.

Bonnefille R. 2007. Rainforest responses to past climate changes in tropical

Africa. In: Tropical rainforest responses to climate change (eds Bush MB,

Flenley JR), pp.117–170, Praxis Publishing, Chichester

Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M,

Gavryushkina A, Heled J, Jones G, Kühnert D, De Maio N and Matschiner

M. 2019 BEAST 2.5: An advanced software platform for Bayesian

evolutionary analysis. *PLoS computational biology* 15(4), p.e1006650.

Brown JW, Walker JF, Smith SA. 2017 Phyx: Phylogenetic Tools for Unix.

Bioinformatics 33(12):1886–88.

https://doi.org/10.1093/bioinformatics/btx063.

Bryant D, Bouckaert R, Felsenstein J, Rosenberg NA, RoyChoudhury A. 2012

Inferring species trees directly from biallelic genetic markers: bypassing

gene trees in a full coalescent analysis. *Molecular biology and evolution,*

29(8), 1917-1932.

Burgarella C, Cubry P, Kane NA, Varshney RK, Mariac C, Liu X, Shi C, Thudi

M, Couderc M, Xu X and Chitikineni A. 2018 A western Sahara centre of

domestication inferred from pearl millet genomes. *Nature ecology &*

evolution 2(9):1377-1380.

Carnaval, AC, Hickerson MJ, Haddad CFB, Rodrigues MT, and Moritz C. 2009

Stability Predicts Genetic Diversity in the Brazilian Atlantic Forest Hotspot.

Science 323 (5915):785–89. https://doi.org/10.1126/science.1166955.

Carnaval AC, Waltari E, Rodrigues MT, Rosauer D, VanDerWal J, Damasceno

R, Prates I et al. 2014 Prediction of Phylogeographic Endemism in an

Environmentally Complex Biome. *Proceedings of the Royal Society B:*

Biological Sciences 281(1792):20141461.

https://doi.org/10.1098/rspb.2014.1461.

Castresana J. 2000. Selection of Conserved Blocks from Multiple Alignments for

Their Use in Phylogenetic Analysis. *Molecular Biology and Evolution* 17

(4): 540–52. https://doi.org/10.1093/oxfordjournals.molbev.a026334.

Caye K, Deist TM, Martins H, Michel O, François O. 2016 TESS3: Fast

Inference of Spatial Population Structure and Genome Scans for Selection.

Molecular Ecology Resources 16(2):540–48. https://doi.org/10.1111/1755-

0998.12471.

Chatrou LW, Pirie MD, Erkens RHJ, Couvreur TLP, Neubig KM, Abbott JR,

Mols JB, Maas JW, Saunders RMK, Chase MW. 2012 A New Subfamilial

and Tribal Classification of the Pantropical Flowering Plant Family

Annonaceae Informed by Molecular Phylogenetics. *Botanical Journal of*

the Linnean Society 169(1):5–40. https://doi.org/10.1111/j.1095-

8339.2012.01235.x.

Collevatti RG, Telles MP, Lima JS, Gouveia FO, Soares TN. 2014 Contrasting

spatial genetic structure in Annona crassiflora populations from fragmented

and pristine savannas. *Plant systematics and evolution* 300(7):1719-1727.

Couvreur TLP, Helmstetter AJ, Koenen EJM, Bethune K, Brandão RD, Little

SA, Sauquet H, Erkens RHJ. 2019 Phylogenomics of the Major Tropical

Plant Family Annonaceae Using Targeted Enrichment of Nuclear Genes.

Frontiers in Plant Science 9 (January). Frontiers:1941.

https://doi.org/10.3389/fpls.2018.01941.

Cowling SA, Cox PM, Jones CD, Maslin MA, Peros M, Spall SA. 2008

Simulated glacial and interglacial vegetation across Africa: implications for

species phylogenies and trans-African migration of plants and animals.

Global Change Biology 14:827–840. https://doi.org/10.1111/j.1365-

2486.2007.01524.x

Dauby G, Duminil J, Heuertz M, Koffi GK, Stévart T, Hardy OJ. 2014

Congruent Phylogeographical Patterns of Eight Tree Species in Atlantic

Central Africa Provide Insights into the Past Dynamics of Forest Cover.

Molecular Ecology 23(9):2299–2312. https://doi.org/10.1111/mec.12724.

De La Torre AR, Li Z, Van De Peer Y, Ingvarsson PK. 2017 Contrasting rates of

molecular evolution and patterns of selection among gymnosperms and

flowering plants. *Molecular Biology and Evolution* 34(6):1363–1377.

De Maio N, Wu CH, O'Reilly KM, Wilson D. 2015 New routes to

phylogeography: a Bayesian structured coalescent approximation. *PLoS*

genetics 11(8). https://doi.org/10.1371/journal.pgen.1005421

Diamond AW, Hamilton AC. 1980 The Distribution of Forest Passerine Birds

and Quaternary Climatic Change in Tropical Africa. *Journal of Zoology* 191

(3): 379–402. https://doi.org/10.1111/j.1469-7998.1980.tb01465.x.

Drummond AJ, Rambaut A. 2007 BEAST: Bayesian Evolutionary Analysis by

Sampling Trees. *BMC Evolutionary Biology* 7(1):214.

https://doi.org/10.1186/1471-2148-7-214.

Duminil J, Mona S, Mardulyn P, Doumenge C, Walmacq F, Doucet JL, Hardy

OJ. 2015 Late Pleistocene Molecular Dating of Past Population

Fragmentation and Demographic Changes in African Rain Forest Tree

Species Supports the Forest Refuge Hypothesis. *Journal of Biogeography*

42(8):1443–54. https://doi.org/10.1111/jbi.12510.

Edwards SV, Potter S, Schmitt CJ, Bragg JG, Moritz C. 2016 Reticulation,

Divergence, and the Phylogeography–Phylogenetics Continuum.

Proceedings of the National Academy of Sciences 113(29):8025–32.

https://doi.org/10.1073/pnas.1601066113.

Elith J, Graham CH, Anderson RP, Dudík M, Ferrier S, Guisan A, Hijmans RJ et

al. 2006 Novel Methods Improve Prediction of Species' Distributions from

Occurrence Data. *Ecography* 29(2):129–51.

https://doi.org/10.1111/j.2006.0906-7590.04596.x.

Faye A, Deblauwe V, Mariac C, Richard D, Sonké B, Vigouroux Y, Couvreur

TLP. 2016 Phylogeography of the Genus *Podococcus* (Palmae/Arecaceae)

in Central African Rain Forests: Climate Stability Predicts Unique Genetic

Diversity. *Molecular Phylogenetics and Evolution* 105:126–38.

https://doi.org/10.1016/j.ympev.2016.08.005.

Fuchs J, Bowie, RC. 2015 Concordant genetic structure in two species of

woodpecker distributed across the primary West African biogeographic

barriers. *Molecular phylogenetics and evolution* 88:64-74.

Hardy OJ, Born C, Budde K, Daïnou K, Dauby G, Duminil J, Ewédjé EBK et al.

2013 Comparative Phylogeography of African Rain Forest Trees: A Review

of Genetic Signatures of Vegetation History in the Guineo-Congolian

Region. *Comptes Rendus - Geoscience* 345(7–8):284–96.

https://doi.org/10.1016/j.crte.2013.05.001.

Heuertz M, Duminil J, Dauby G, Savolainen V, Hardy OJ. 2014 Comparative

Phylogeography in Rainforest Trees from Lower Guinea, Africa. *PLoS*

ONE 9(1):e84307. https://doi.org/10.1371/journal.pone.0084307.

Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005 Very High

Resolution Interpolated Climate Surfaces for Global Land Areas.

International Journal of Climatology 25(15):1965–78.

https://doi.org/10.1002/joc.1276.

Holbrook KM, Smith TB. 2000 Seed dispersal and movement patterns in two

species of *Ceratogymna* hornbills in a West African tropical lowland forest.

Oecologia 125:249–257. https://doi.org/10.1007/s004420000445

Huson DH, Bryant D. 2006 Application of Phylogenetic Networks in

Evolutionary Studies. *Molecular Biology and Evolution*. 23(2):254-267.

https://doi.org/10.1093/molbev/msj030.

Johnson MG, Gardner EM, Liu Y, Medina R, Goffinet B, Shaw AJ, Zerega NJC,

Wickett NJ. 2016 HybPiper: Extracting Coding Sequence and Introns for

Phylogenetics from High-Throughput Sequencing Reads Using Target

Enrichment. *Applications in Plant Sciences* 4(7):1600016.

https://doi.org/10.3732/apps.1600016.

Jombart T. 2008 Adegenet: A R Package for the Multivariate Analysis of

Genetic Markers. *Bioinformatics* 24(11):1403–5.

https://doi.org/10.1093/bioinformatics/btn129.

Jombart T, Devillard S, Balloux F. 2010 Discriminant Analysis of Principal

Components: A New Method for the Analysis of Genetically Structured

Populations. *BMC Genetics* 11(1):94. https://doi.org/10.1186/1471-2156-

11-94.

Katoh K, Standley DM. 2013 MAFFT Multiple Sequence Alignment Software

Version 7: Improvements in Performance and Usability. *Molecular Biology*

and Evolution 30 (4): 772–80. https://doi.org/10.1093/molbev/mst010.

Leite RN, Rogers DS. 2013 Revisiting Amazonian Phylogeography: Insights into

Diversification Hypotheses and Novel Perspecitves. *Organisms Diversity &*

Evolution 13(4):639–64. https://doi.org/10.1007/s13127-013-0140-8.

Leite YLR, Costa LP, Loss AC, Rocha RG, Batalha-Filho H, Bastos AC,

Quaresma VS et al. 2016 Neotropical Forest Expansion during the Last

Glacial Period Challenges Refuge Hypothesis. *Proceedings of the National*

Academy of Sciences 113(4):1008–13.

https://doi.org/10.1073/pnas.1513062113.

Lemey P, Rambaut A, Welch JJ, Suchard MA 2010. Phylogeography takes a

relaxed random walk in continuous space and time. *Molecular biology and*

evolution 27(8):1877-1885. https://doi.org/10.1093/molbev/msq067

Lemey P, Rambaut A, Bedford T, Faria N, Bielejec F, Baele G, Russell CA et al.

2014. Unifying viral genetics and human transportation data to predict the

global transmission dynamics of human influenza H3N2. *PLoS*

*pathogens:*10(2). https://doi.org/10.1371/journal.ppat.1003932

Ley AC, Heuertz M, Hardy OJ. 2017 The Evolutionary History of Central

African Rain Forest Plants: Phylogeographical Insights from Sister Species

in the Climber Genus *Haumania* (Marantaceae). *Journal of Biogeography*

44(2):308–21. https://doi.org/10.1111/jbi.12902.

Lézine AM, Izumi K, Kageyama M, Achoundong G. 2019 A 90,000-year record

of Afromontane forest responses to climate change. *Science* 363(6423):177-

181.

Li H. 2011 A statistical framework for SNP calling, mutation discovery,

association mapping and population genetical parameter estimation from

sequencing data. *Bioinformatics* 27(21), 2987-2993.

Li H, Durbin R. 2009 Fast and Accurate Short Read Alignment with Burrows-

Wheeler Transform. *Bioinformatics* 25(14):1754–60.

https://doi.org/10.1093/bioinformatics/btp324.

Liu X, Fu YX. 2015 Exploring population size changes using SNP frequency

spectra. *Nature Genetics* 47, 555-559.

845   Linder HP. 2001 Plant Diversity and Endemism in Sub-Saharan Tropical Africa.

846   In *Journal of Biogeography*, 28:169–82. https://doi.org/10.1046/j.1365-

847   2699.2001.00527.x.

848   Lowe AJ, Harris D, Dormontt E, Dawson IK. 2010 Testing Putative African

849   Tropical Forest Refugia Using Chloroplast and Nuclear DNA

850   Phylogeography. *Tropical Plant Biology* 3(1):50–58.

851   https://doi.org/10.1007/s12042-010-9045-2.

852   Mace GM, Gittleman JL, Purvis A. 2003 Preserving the Tree of Life. *Science*

853   300 (5626):1707–9. https://doi.org/10.1126/science.1085510.

854   Maley J. 1996 The African Rain Forest - Main Characteristics of Changes in

855   Vegetation and Climate from the Upper Cretaceous to the Quaternary.

856   *Proceedings of the Royal Society of Edinburgh Section B: Biological*

857   *Sciences* 104:31–73. https://doi.org/10.1017/S0269727000006114.

858   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K et al. 2010 The

859   Genome Analysis Toolkit: A MapReduce Framework for Analyzing next-

860   Generation DNA Sequencing Data. *Genome Research* 20(9):254–60.

861   https://doi.org/10.1101/gr.107524.110.20.

862   Muscarella R, Galante PJ, Soley-Guardia M, Boria RA, Kass JM, Uriarte M,

863   Anderson RP. 2014 ENMeval: An R Package for Conducting Spatially

864   Independent Evaluations and Estimating Optimal Model Complexity for

865   Maxent Ecological Niche Models. Methods in Ecology and Evolution

866   5(11):1198–1205. https://doi.org/10.1111/2041-210x.12261.

867   Phillips SJ, Anderson RP, Schapire, RE. 2006 Maximum Entropy Modeling of

868   Species Geographic Distributions. *Ecological Modelling* 190(3–4):231–59.

869   https://doi.org/10.1016/j.ecolmodel.2005.03.026.

Piñeiro R, Dauby G, Kaymak E, Hardy OJ. 2017 Pleistocene Population

Expansions of Shade-Tolerant Trees Indicate Fragmentation of the African

Rainforest during the Ice Ages. *Proceedings of the Royal Society B:*

Biological Sciences 284(1866):20171800.

https://doi.org/10.1098/rspb.2017.1800.

Portik DM, Leaché AD, Rivera D, Barej MF, Burger M et al. 2017 Evaluating

mechanisms of diversification in a Guineo-Congolian tropical forest frog

using demographic model selection. *Molecular Ecology* 26(19):5245-5263.

Ponder WF, Carter GA, Flemons P, Chapman RR. 2001 Evaluation of Museum

Collection Data for Use in Biodiversity Assessment. *Conservation Biology*

15 (3): 648–57. https://doi.org/10.1046/j.1523-1739.2001.015003648.x.

Poulsen JR, Clark CJ, Smith TB. 2001 Seed Dispersal by a Diurnal Primate

Community in the Dja Reserve, Cameroon. *Journal of Tropical Ecology*

17:787–808.

Radosavljevic A, Anderson RP. 2014 Making better Maxent models of species

distributions: complexity, overfitting and evaluation. *Journal of*

Biogeography 41:629–643. https://doi.org/10.1111/jbi.12227

Raj A, Stephens M, Pritchard JK. 2014 FastSTRUCTURE: Variational Inference

of Population Structure in Large SNP Data Sets. *Genetics* 197(2):573–89.

https://doi.org/10.1534/genetics.114.164350.

Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. 2018 Posterior

summarisation in Bayesian phylogenetics using Tracer 1.7. Systematic

Biology. syy032. doi:10.1093/sysbio/syy032

Rohland N, Reich D. 2012. Cost-Effective, High-Throughput DNA Sequencing

Libraries for Multiplexed Target Capture. *Genome Research* 22(5):939–46.

https://doi.org/10.1101/gr.128124.111.

Schoener TW. 1968 Anolis lizards of Bimini: resource partitioning in a complex

fauna. *Ecology* 49:704- 726.

Stamatakis A. 2014 RAxML Version 8: A Tool for Phylogenetic Analysis and

Post-Analysis of Large Phylogenies. *Bioinformatics* 30(9):1312–13.

https://doi.org/10.1093/bioinformatics/btu033.

Thuiller W, Lafourcade B, Engler R, Araújo MB. 2009 BIOMOD – a platform

for ensemble forecasting of species distributions. *Ecography* 32:369–373.

https://doi.org/10.1111/j.1600-0587.2008.05742.x

Turelli M, Barton NH, Coyne JA. 2001 Theory and speciation. *Trends in*

Ecology & Evolution 16:330–343. https://doi.org/10.1016/S0169-

5347(01)02177-2

Versteegh CP, Sosef M. 2007 Revision of the African Genus *Annickia*.

Systematics and Geography of Plants 77(1):91–118.

https://doi.org/10.1600/036364411X553108.

Vitorino LC, Lima-Ribeiro MS, Terribile LC, Collevatti RG. 2016

Demographical History and Palaeodistribution Modelling Show Range Shift

towards Amazon Basin for a Neotropical Tree Species in the LGM. *BMC*

Evolutionary Biology 16(1):1–15. https://doi.org/10.1186/s12862-016-

0779-9.

Warren DL, Glor RE and Turelli M 2008. Environmental niche equivalency

versus conservatism: quantitative approaches to niche evolution. *Evolution*

62(11):2868-2883.

Warren DL, Matzke N, Cardillo M, Baumgartner J, Beaumont L, Huron N,

Simões M and Dinnage R. 2019 ENMTools (Software Package). URL:

https://github.com/danlwarren/ENMTools. doi:10.5281/zenodo.3268814

Zhang C, Sayyari E, Mirarab S. 2017 ASTRAL-III: Increased Scalability and

Impacts of Contracting Low Support Branches. In *Lecture Notes in*

Computer Science (Including Subseries Lecture Notes in Artificial

Intelligence and Lecture Notes in Bioinformatics) 10562 LNBI:53–75.

https://doi.org/10.1007/978-3-319-67979-