An ecophysiological model of plant-pest interactions: the role of nutrient and water availability

Marta Zaffaroni[1]   Nik J. Cunniffe[2]   Daniele Bevacqua[1*]

[1] INRA, UR1115 Plantes et Systèmes de culture Horticoles (PSH), Site Agroparc, 84914 Avignon, France. [2] Department of Plant Sciences, University of Cambridge, Cambridge CB2 3EA, United Kingdom. [*] Corresponding author: `daniele.bevacqua@inrae.fr`

Abstract

Empirical studies have shown that particular irrigation/fertilization

regimes can control pest populations in agroecosystems. This appears

to promise that the ecological concept of bottom-up control can be

applied to pest management. However, a conceptual framework is nec-

essary to develop a mechanistic basis for empirical evidence. Here we

couple a mechanistic plant growth model with a pest population model.

We demonstrate its utility by applying it to the peach - green aphid

system. Aphids are herbivores which feed on the plant phloem, deplete

plants' resources and (potentially) transmit viral diseases. The model

reproduces system properties observed in field studies and shows un-
der which conditions the diametrically-opposed plant vigour and plant
stress hypotheses find support. We show that the effect of fertiliza-
tion/irrigation on the pest population cannot be simply reduced as
positive or negative. In fact, the magnitude and direction of any effect
depends on the precise level of fertilization/irrigation and on the date
of observation. We show that a new synthesis of experimental data can
emerge by embedding a mechanistic plant growth model, widely stud-
ied in agronomy, in a consumer-resource modelling framework, widely
studied in ecology. The future challenge is to use this insight to inform
practical decision making by farmers and growers.

Keywords: agroecology, aphid population model, induced plant defence, plant growth model, plant stress and plant vigour hypotheses, plant-aphid interactions

Introduction

In agriculture, pest control mostly relies on the use of chemical pesticides. However, widespread application of agrochemicals carries an inherent environmental cost. There is also the significant challenge of declining efficacy due to the emergence and spread of insecticide resistance in pest populations (1). In recent decades, agroecology has developed as discipline which aims to provide alternatives to the use of chemicals in agronomy to control pest. The rationale is that ecological concepts and principles can be applied to control pest populations while reducing the use of chemicals (2). The concept of 'bottom-up' control, according to which population dynamics are driven by quantity and quality of resources, is particularly highlighted by agroecologists. There are a number of agricultural practices that can affect plant physiology and alter resources offered by plants to pests (3; 4). For example, fertilization modifies nutrient balance in plants, enhancing plant tissue nutritional status, and influences the synthesis of defence compounds (5). Similarly, irrigation controls plant vigour, phloem nutritional quality and viscosity, possibly regulating pest abundance (6; 7; 8; 9; 10).

Unfortunately, how pests might be affected by plant nutrient and irrigation status is far from obvious. Empirical evidence is ambiguous, potentially supporting diametrically-opposed hypotheses. On the one hand, the Plant Vigour Hypothesis (PVH) (11) argues that pest populations should increase most rapidly on vigorously growing plants (or organs), since these habitats provide more resources. In support of this hypothesis, there is some experimental evidence suggesting that practices such as fertilization and irrigation,

or favourable conditions for plant growth such as increased organic soil fertility, can be associated with abundant pest populations (12; 13). On the other hand, the Plant Stress Hypothesis (PSH) (14) argues that pests perform better on stressed plants that would not have resources to deploy defences and/or whose nutritional quality might be enhanced. This as been determined experimentally to be the case for some aphid species feeding on plants subjected to controlled irrigation deficit (15; 16).

In order to efficiently use the concepts of bottom-up control in agroecology, it is necessary to shed light on the mechanisms that are responsible for the observed patterns. We require a unified conceptual framework sufficiently flexible for both the PVH and PSH hypotheses to find support. Developing and validating such a framework requires integration of information from field experiments with mathematical modelling. Experimental data is clearly necessary to test the validity of theoretical hypotheses, but is often extremely costly and time consuming to obtain. Mathematical modelling, particularly mechanistic models, represent a useful tool to investigate which processes can be responsible for the observed patterns and to explore the consequences of different agricultural practices (17).

Here we present a new, explicitly agro-ecological, model synthesising elements of models as commonly used within the separate disciplines of agronomy and plant pest dynamics. Agronomic models tend to empirically parametrize the detrimental effects of pests on plant biological rates (*e.g.* photosynthetic, growth, solutes transport) . However such models invariably neglect the dynamical interaction between the plant (or some of its component parts) and the pest (see *e.g.* 18; 19; 20). That is, the impact of a pest

on the plant is modelled by varying one or more plant parameters, according to the pest disturbance level with no further interaction or feedback. On the other hand, in ecology, there is a very broad literature of models on interactions (*e.g.* predation, consumption, competition etc.) between different species or organisms. These types of models have been widely used to study temporal and spatial dynamics in plant-pest (*e.g.* 21; 22; 23) and particularly plant-pathogen systems (*e.g.* 24; 25; 26). However these types of model usually present a simplistic description of the plant (but see (27)), which in turn limits the possibility to consider the effects of agronomic practices.

Aiming to bridge the gap between the classical agronomic and the ecological modelling approaches, here we couple a plant growth model, that describes carbon and nitrogen assimilation and allocation to shoot and root compartments of a plant, with a pest population model. With regard to the plant, we use the modelling framework proposed by Thornley in the early 70s (28), and refined in the following decades (29; 30; 31), which represents a cornerstone in plant and crop modelling. With regard to the pest, we propose a novel population model which includes intraspecific competition in which pest birth and mortality rates depend on resource availability and quality. Moreover, we assume that the presence of the pest can induce the plant to produce defensive traits or compounds (32). We demonstrate the utility of our model by applying it to the peach (*Prunus persica*) - green aphid (*Myzus persicae*) system. Aphids are specialized herbivores which feed on the phloem of vascular plants. This depletes plants' resources, affecting growth and reproduction, as well as eventually impacting upon yield (33). Moreover, aphids are the most common vector of plant viral diseases and so

can often cause indirect damage far exceeding direct impacts via herbivory (34). We use likelihood-based techniques to calibrate model parameters and select model assumptions against field data obtained under different conditions of irrigation and fertilization. The resulting model has the ability to reproduce different system properties observed in field studies, as well as showing under which conditions the PVH and PSH find more support. Our model also provides insights to conceive new targeted experiments to better understand this class of system and rethink the control of plant-aphid systems.

Material and Methods

Model outline and assumptions

The model, which describes the temporal variation, during a growing season, of plant dry mass (partitioned into shoots and roots, in turn composed of structural mass, carbon and nitrogen substrates), its induced defensive level and the aphid population dwelling on the plant, is schematically represented in Figure 1. According to Thornely et al's seminal works (28; 17; 29; 30), carbon is assimilated from the atmosphere via photosynthesis and stored in shoots, as shoot carbon substrate (C_S), or transported and then stored in roots as root carbon substrate (C_R). Similarly, nitrogen is assimilated from the soil, stored in roots as root nitrogen substrate (N_R), or transported and then stored in shoots as shoot nitrogen substrate (N_S). Carbon and nitrogen substrates are utilized, in a fixed ratio, to constitute structural

shoot (S) and root (R) dry mass. With respect to the original model of Thornley, we added the assumption that the constitution of new plant mass is regulated by changes in the photo-period (35). Such an assumption permits us to model the fact that perennial plants suspend growth, in favour of reserve constitution, before entering winter dormancy (36). The assimilation of substrate (C_S or N_R) per unit of plant organ (S or R) decreases with organ mass due to shoot self-shading and root competition for nitrogen and it is inhibited by substrate concentration in the organ (30).

We coupled the plant model of carbon and nitrogen assimilation and partitioning with an aphid population model by assuming that aphids, which penetrate growing shoots of the host plant with a stylet and feed on the phloem (37), intercept a fraction of the substrates (C_S and N_S) directed towards the shoot structural mass compartment (S) to support their growth (38). We assume that aphids act in a scramble competition context (39) and therefore any aphid ingests its maximum daily amount of food when the per-capita available resource is sufficient, but that otherwise the resource is evenly shared among all the individuals. The aphid birth rate depends on the per-capita ingested food (40) and crowding can induce aphids to leave the plant (41).

We assumed that the infested plant can be induced to use carbon and nitrogen substrates to defend itself, to the detriment of growth (42; 37; 43). This can result in the production of chemical and/or in morphological and physiological changes that can reduce aphid accessibility to the resource (e.g. by phloem sealing) (44; 45) and/or decrease the rate at which ingested food is converted into progeny, e.g. by releasing toxic components in the sieve

 that can even repel or kill the aphid (32).

Model equations

148 In quantitative terms, we describe the temporal variation of the plant-aphid

149 system with the following system of ordinary differential equations.

$$\dot{C}_S = \sigma_C S[(1+\frac{S}{\nu})(1+\frac{C_S}{S\iota_C})]^{-1} - \varphi_C \kappa \frac{C_S}{S}\frac{N_S}{S}S - (\frac{C_S}{S}-\frac{C_R}{R})(SR)^q \cdot (S^q+R^q)^{-1} - \alpha\frac{C_S}{S}A \tag{1a}$$

$$\dot{N}_S = (\frac{N_R}{R}-\frac{N_S}{S})(SR)^q \cdot (S^q+R^q)^{-1} - \varphi_N \kappa \frac{C_S}{S}\frac{N_S}{S}S - \alpha\frac{N_S}{S}A \tag{1b}$$

$$\dot{S} = \begin{cases} \Phi\kappa\frac{C_S}{S}\frac{N_S}{S}S(1 - \frac{\theta A}{\Phi\kappa\frac{C_S}{S}\frac{N_S}{S}S}) & \text{if } \theta \cdot A \le \Phi\kappa\frac{C_S}{S}\frac{N_S}{S}S(1-\beta_1\frac{\frac{D^{\delta_1}}{S}}{\pi_1^{\delta_1}+\frac{D^{\delta_1}}{S}}) \\ \Phi\kappa\frac{C_S}{S}\frac{N_S}{S}S(\beta_1\frac{\frac{D^{\delta_1}}{S}}{\pi_1^{\delta_1}+\frac{D^{\delta_1}}{S}}) & \text{otherwise} \end{cases} \tag{1c}$$

$$\dot{C}_R = (\frac{C_S}{S}-\frac{C_R}{R})(SR)^q \cdot (S^q+R^q)^{-1} - \varphi_C \kappa \frac{C_R}{R}\frac{N_R}{R}R \tag{1d}$$

$$\dot{N}_R = \sigma_N R[(1+\frac{R}{\nu})(1+\frac{N_R}{R\iota_N})]^{-1} - \varphi_N \kappa \frac{C_S}{S}\frac{N_S}{S}S - (\frac{N_R}{R}-\frac{N_S}{S})(SR)^q(S^q+R^q)^{-1} \tag{1e}$$

$$\dot{R} = \Phi\kappa\frac{C_R}{R}\frac{N_R}{R}R \tag{1f}$$

$$\dot{D} = (\varepsilon_C\alpha\frac{C_S}{S}+\varepsilon_N\alpha\frac{N_S}{S})A \tag{1g}$$

$$\dot{A} = \begin{cases} (\xi\theta(1-\beta_2\frac{\frac{D^{\delta_2}}{S}}{\pi_2^{\delta_2}+\frac{D^{\delta_2}}{S}})-\mu-\omega\frac{A}{S})A & \text{if } \theta \cdot A \le \Phi\kappa\frac{C_S}{S}\frac{N_S}{S}S(1-\beta_1\frac{\frac{D^{\delta_1}}{S}}{\pi_1^{\delta_1}+\frac{D^{\delta_1}}{S}}) \\ (\xi\Phi\kappa\frac{C_S}{S}\frac{N_S}{S}S(1-\beta_1\frac{\frac{D^{\delta_1}}{S}}{\pi_1^{\delta_1}+\frac{D^{\delta_1}}{S}})\frac{1}{A}(1-\beta_2\frac{\frac{D^{\delta_2}}{S}}{\pi_2^{\delta_2}+\frac{D^{\delta_2}}{S}})-\mu-\omega\frac{A}{S})A & \text{otherwise} \end{cases} \tag{1h}$$

In our model C_S, N_S, S, C_R, N_R and R are expressed in grams (g); D is

expressed in an arbitrary defence unit (DU) and A in individuals (ind.); t rep-

resents the number of days (d) that have passed since the January 1st of the

year of the considered growing season. In equation 1a, $\sigma_C S[(1+\frac{S}{\nu})(1+\frac{C_S}{S\iota_C})]^{-1}$

is the carbon substrate assimilated in shoots, $\varphi_C\kappa\frac{C_S}{S}\frac{N_S}{S}S$ is the shoot carbon

substrate allocated to shoot growth or reserves, $(\frac{C_S}{S}-\frac{C_R}{R})(SR)^q \cdot (S^q+R^q)^{-1}$

is the shoot carbon substrate transported toward roots and $\alpha\frac{C_S}{S}A$ is the

shoot carbon substrate diverted to defences, in a unit of time. In equation 1b, $(\frac{N_R}{R} - \frac{N_S}{S})(SR)^q \cdot (S^q + R^q)^{-1}$ is the nitrogen substrate transported from roots towards shoots; $\varphi_N \kappa \frac{C_S}{S} \frac{N_S}{S} S$ is the shoot nitrogen substrate allocated to shoot growth or reserves, and $\alpha \frac{N_S}{S} A$ is the shoot nitrogen substrate diverted to defences, with each of these quantities being measured as rates per unit of time. In equation 1c, the time dependent parameter $\Phi = \frac{\lambda^\eta}{\lambda^\eta + t^\eta}$ determines the suspension of plant growth driven changes in the photo-period The term $\Phi \kappa \frac{C_S}{S} \frac{N_S}{S} S$ is the increase in structural shoot dry mass in the absence of any phloem withdrawal by the aphids, with k being the maximum rate of utilization of the substrates. The term $\frac{\theta A}{\Phi \kappa \frac{C_S}{S} \frac{N_S}{S} S}$ represents the fraction of substrates diverted from allocation to plant growth, because of ingestion by aphids, when aphid per-capita intake is limited by aphid maximum daily food intake, θ, and not by resource availability. The term $\beta_1 \frac{\frac{D}{S}^{\delta_1}}{\pi_1^{\delta_1} + \frac{D}{S}^{\delta_1}}$ indicates the fraction of phloem that is protected by plant defences and therefore inaccessible for aphids. When aphid per-capita intake is limited by the resource availability, aphids ingest all the phloem they can access and the per-capita intake is reduced. The dynamics of the variables in the root compartments (C_R, N_R, R) follow similar rules as for assimilation of substrates, transport and allocation to root growth and we assumed that they are not directly affected by the presence of aphids. In equation 1h, we assume that the aphid birth rate is proportional to the per-capita food intake (θ or $\Phi \kappa \frac{C_S}{S} \frac{N_S}{S} S(1 - \beta_1 \frac{(\frac{D}{S})^{\delta_1}}{\pi_1^{\delta_1} + (\frac{D}{S})^{\delta_1}}) \frac{1}{A}$) and that it can decrease due to a possible action of the defences. In other words, we assume that plant defences can determine an extra mortality rate, per unit of ingested food, modelled as $\beta_2 \frac{\frac{D}{S}^{\delta_2}}{\pi_2^{\delta_2} + \frac{D}{S}^{\delta_2}}$. We modelled both the fraction of the phloem that can be protected and the phloem "toxicity" as

an increasing function of the concentration of defences, $\frac{D}{S}$. The shape of this function is given by the value of parameter δ_i. Namely, if $\delta_i > 1$ it is convex for $\frac{D}{S} < \pi_i$ and concave for $\frac{D}{S} > \pi_i$, if $0 < \delta_i < 1$, it is strictly concave. The parameter ω is the strength of possible density dependent mechanisms inducing aphid migration. Details of the model variables and parameters are reported in Table 1.

Model calibration

We apply the model to a peach-green aphid system subjected to different fertilization and irrigation treatments. Details on the experimental data are reported in a previous work (8) and in the Supporting Information, SI. According to available data, we set initial conditions of the system at the first observation date (i.e. April 29$^{\text{th}}$, 119$^{\text{th}}$ day of the year 2013) (see SI). We set the value of model parameters according to information available from peer-reviewed literature or experimental data whenever possible (Table 1 and SI). On the other hand, no information was available to *a priori* derive reliable estimates for parameters σ_{N} (net N assimilation rate) and κ (maximum rate of utilization of the substrates), which depend on environmental conditions that possibly varied in the different treatments; parameter q, affecting substrates transport within the plant and depending on the plant architecture (30), and six parameters relevant to the production of defences (α) and their effect ($\pi_1, \delta_1, \beta_2, \pi_2, \delta_2$). We estimated these unknown parameters by minimizing a cost function expressed as the sum of two negative log-likelihood functions, computed with respect to observations of shoot dry mass and

aphid abundance (see SI for details). We assessed the empirical distribu-

tions of calibrated parameters by making use of the moving block bootstrap

(46). In particular, we reconstructed bootstrapped time series for each of

the observed variables and we fitted the values of the unknown parameters.

We repeated this process 1,000 times and we generated the 99% confidence

intervals CI for each parameter via the percentile method (47).

Model selection

To account for possible different mechanisms regarding aphid ecology, the

plant response to aphid infestation and its consequences, we contrasted the

'full' model reported in eq.1 with a set of nested models lacking some pro-

cesses (Fig. 2). Namely, the full model (M10) assumes that aphid crowding

promotes aphid migration, that the plant produces defences that make a frac-

tion of resources inaccessible to aphids and kill, or repel, aphid if ingested.

Three models nested in M10 assume a crowding effect on aphid migration and

the induced production of defences. Yet, they can differ regarding the effect

of defences: killing/repulsion effect (M9), reduction of phloem accessibility

(M8), or no effect (M7). There is also a simpler model neglect the produc-

tion of defences (M6). We also considered five analogous models ignoring the

effect of aphid crowding, $\omega = 0$, (M1, M2, M3, M4, M5).

We tested if the effect of irrigation and fertilization can be represented in

the models thorough a variation in some parameters κ and σ_{N}. The rationale

is that the rate of utilization of the substrates (parameter k) and the nitrogen

assimilation rate (parameter σ_N) are expected to decrease in water (48; 7)

and nutrient (49; 50) stress conditions, respectively. We then contrasted each of the ten models assuming that *i)* κ and σ_N respectively vary with irrigation and fertilization treatments; *ii)* κ varies with irrigation and σ_N does not vary with fertilization; *iii)* κ does not vary with irrigation and σ_N varies with fertilization; *iv)* neither σ_N nor κ vary with fertilization and irrigation. Therefore, we calibrated two values for nitrogen assimilation rate per unit of root (σ_N^+, σ_N^-) in cases *i* and *iii* and a unique value ($\sigma_N^\pm$) in cases *ii* and *iv*. Analogously, we calibrated two values for the allocation of substrates to plant growth (κ^+ and κ^-) in cases *i* and *ii* and a unique value ($\kappa^\pm$) in cases *iii* and *iv*.

Overall, we compared 40 different models (Fig. 2), obtained by incorporating five hypotheses on plant defences, two on density dependence of aphid migration, and four on the effect of irrigation and fertilization, to one another. We selected the best model, that is the one providing the best compromise between goodness of fit to observed data and parsimony, through a model selection procedure based upon Akaike information criterion (51). See SI for details.

Sensitivity Analysis

To assess the robustness of model outputs to uncertainty affecting model parameters, we performed a sensitivity analysis of the model to (small) perturbations of the default parameter values reported in Table 1. According to (17), we computed the sensitivity of the variable Y (where for Y we considered the maximum value of S, of A and of their ratio A/S over the growing

season), to small variations of parameter p_i as

$$\psi(Y, p_i) = \frac{\partial Y}{\partial p_i} \frac{p_i}{Y} \simeq \frac{\delta Y}{Y} \frac{p_i}{\delta p_i} \qquad (2)$$

In practice, after having changed the value of each parameter by $+5\%$, we computed the value of ψ and if $\psi(Y, p_i) > 1$ we concluded that parameter has a more-than-linear effect on the variable.

The role played by fertilization and irrigation

After having ascertained that parameters κ and σ_N are likely to vary with irrigation and fertilization treatments, respectively, we used the selected model to simulate the temporal dynamics of the system for different values of these parameters. This allowed us to perform an *in silico* experiment to explore whether or not the model was able to reproduce the observed empirical patterns that claimed support for the plant vigour or the plant stress hypotheses. The *in silico* experiment is intended to test if the aphid density is affected by the fertilization (or irrigation) treatment. We considered five levels for the fertilization treatment (i.e. σ_N equals to 0.0012, 0.0024, 0.012, 0.06 and 0.12 d^{-1}) and five levels for the irrigation treatment (i.e. κ equals to 18, 36, 182, 910 and 1820 d^{-1}) corresponding to very low - low - average - high - very high levels of fertilization (or irrigation). We varied the level of one treatment while keeping the other fixed at its average value thus obtaining nine different combinations of factorial levels. Since in real factorial experiments the number of replicates (i.e. different plant individuals) is limited, we chose to simulate ten replicates for each factors combination, which corresponds

to a realistic experiment with 90 plants being monitored. We simulated ten possible trajectories of the system variables, for the same factors combination, by running the model with ten different parameter sets drawn from the empirical distribution obtained in the calibration process.

Results

Model calibration and selection

The best model ('the model', hereafter) assumes that i) aphid migration due to crowding can be neglected; ii) aphid presence induces the plant to divert resources from growth to defence; iii) defences reduce phloem accessibility to aphids and, at higher concentrations, make the phloem sufficiently toxic to kill or repel aphids (Fig. S1 in the SI); iv) the rates of nitrogen assimilation and substrates utilization differ for different levels of fertilisation and irrigation, respectively.

The model fitted all four data sets, reproducing the main observed temporal patterns and differences between treatments (Fig.3). Shoot growth is enhanced in N^+ treatments while the water treatment considered here plays only a relatively minor role. The time course for shoot mass is linear and followed by a stop towards the end of June. This is consistent with a potential exponential course, in the first part of the season (52), which has been prevented by the presence of the aphids. On the other hand, the stop in shoot growth at the end of June is induced by changes in the day-length. Note that parameter $\phi(t) = 0.5$ for $t = \lambda = 169$, corresponding to June 18$^{\text{th}}$.

Aphid population growth is initially sigmoidal, followed by a decay towards the end of June when the plant growth is halted (Fig.3) and the concentration of defences attains the critical value of $\pi_2 = 0.03 - 0.08$ which makes ingestion from the phloem detrimental rather than beneficial to aphids. The initial phase of aphid growth is enhanced in N^+ treatments characterized by more vigorous plants.

The model gives biologically plausible parameter estimates (Table 1) and the estimated variability in parameters permits most of the variability observed in the data to be captured. The calibrated values of σ_n, k and q are consistent with previously published values (i.e. $\sigma_n = 0.02d^{-1}$, $k = 200d^{-1}$ and $q = 0.67 - 1$ in (30)). The estimated values of parameters $\delta_1 < 1$ and $\delta_2 > 1$ suggest that the fraction of phloem that is protected from aphid withdrawal quickly increases for low concentrations of defences, whereas the phloem toxicity is switched-on when the concentration of defences exceeds a threshold value (Fig. 1 in the SI). On the other hand, the model parameters relevant to the production of defences and their effect on aphids have no equivalent in the literature for a direct comparison.

Sensitivity analysis

Ranked values of the sensitivity, ψ, of shoot production, and maximum aphid abundance and density to small changes in the parameter values are reported in Table S2 in the S.I. Negative values of ψ indicate a negative correlation between a change in a parameter value and the corresponding variable of interest. As expected, increasing the parameter λ results in an increase

of shoot production, as it determines an increase of the growing season of 8.45 d, being $0.05\lambda = 8.45$, and consequently more resources to sustain a bigger aphid population, maintaining similar aphid densities. Similarly, an increase of q results in an increase of both shoot production and in the peak of aphid abundance and density, as it determines a more efficient transport of substrates C and N between roots and shoots. This translates into bigger plants able to sustain higher peaks of aphid population densities.

With the exception of q and λ, our sensitivity analysis indicates none of the model parameters has important (e.g. $\psi > 1$) consequences, indicating that the model is robust. However, our sensitivity analysis nevertheless provides some interesting insights. For instance, it shows that an increase in all those parameters positively related to the plant growth ($\sigma_c, \sigma_n, \iota_C, \iota_N, k, \nu$) determine an increase in the maximum aphid abundance and, to a lower extent, in maximum aphid density. If the aphids were more efficient in converting food into progeny (higher ξ), aphid density would increase but the overall population abundance would diminish as the resource would be overexploited. An increase of the parameter α, determining a higher rate of resources devoted to defences, would have almost no effect on the shoot production but it would decrease aphid abundance and density. Yet, the plant could take advantage of a lower aphid abundance being aphids important vectors of viral diseases (37).

The role played by fertilization and irrigation

Shoot growth follows a sigmoidal pattern and it increases with fertilization

and irrigation (Fig. 4A-B). The concentration of carbon substrates in shoots varies between 3-23% during the growing season with peaks at its beginning, when plant growth is limited by the nitrogen supply, and at its end, when plant growth halts in response to day length decreases, but carbon assimilation continues. Carbon concentration is enhanced in stressful conditions (very low to low fertilization/irrigation treatments) that limit plant growth rather than carbon assimilation (Fig. 4C-D). The concentration of nitrogen substrates varies between 0.1-1.4 % during the growing season (Fig. 4E-F). It decreases in the first weeks of growth, but, in the case of very high/high fertilization, or very low irrigation, it increases until the second week of May. In fact, for high fertilization treatments, nitrogen is not initially consumed by plant growth which is limited by carbon supply and, for low watering, nitrogen concentration increases as plant growth is impaired while N assimilation is not. Peak concentration of defences is delayed in time for higher fertilization and irrigation (Fig. 4G-H). When plant is well watered, the time of the peak aphid population density is delayed by one week. This is due to the fact that defences need more time to reach significant concentrations in bigger plants (Fig. 4I-J). The positive effect of fertilization and irrigation upon aphid abundance becomes evident in the end of May. In the first part of the season, aphid density is enhanced by a low/average value of fertilization (or irrigation) while later in the season aphid density is higher in a well fertilized (irrigated) plant (Fig. 4K-L).

The results of our virtual experiment show that one can draw very different conclusions depending on the considered fertilization/irrigation levels and the date of observations. For instance, one could infer that i) fertiliza-

tion enhances aphid population by observing aphid density in the mid-late part of the season for very low to average values of fertilization (Fig. 5C-E); *ii*) decreases it, by observing aphid density in the early-mid season for average to very high values of fertilization (Fig. 5A-C); *iii)* has no effect, by observing aphid density early and late in the season, for high to very high values of fertilization (Fig. 5A-E). Similarly, different conclusions can be drawn regarding the effect of irrigation: positive (Fig. 5F), negative (Fig. 5B) or null (Fig. 5D, from average to very high values of irrigation).

Discussion

In this work we showed that embedding a mechanistic plant growth model, widely studied in agronomy, in a consumer-resource modelling framework, widely studied in ecology, might be a promising approach for agroecology. We demonstrated the ability of such a novel approach in understanding the consequences of irrigation and fertilization treatments in a plant-aphid system.Yet, the proposed model has the ambition of being physiologically rigorous and general enough to be applied to different plant-pest systems and to incorporate the description of other agronomic practices.

The selected model and model calibration and selection

A recent review (37) suggested that infested plants can put in place phloem-sealing mechanisms to interfere with aphids' access to plant resources and produce a number of secondary metabolites (e.g. cardenolides, glucosinolates and benzoxazinoids) which, if ingested, impair aphid viability. Our study

suggests that both defensive mechanisms are at play in the peach-green aphid system. According to our calibration, impairing phloem accessibility is the most effective at low defences concentration, while 'intoxicating' aphids is the most effective at higher concentration. This is in accordance with works on the arabidopsis-*Myzus persicae* system, for which reductions of aphids fecundity, up to 100%, have been reported in response to high concentrations of some plant defensive compounds (53; 54). The model application to a real study case subjected to different irrigation × fertilization treatments indicates that parameters relevant to plant nitrogen assimilation (σ_{N}) and plant utilization of substrates (κ), originally proposed within a theoretical framework (28), can be linked to agronomic practices and then manipulated by the grower. However, in order to effectively use the proposed model to define effective agronomic recommendations, further studies on the response of the model parameters to effective practices are clearly required.

One of the main features of the peach-green aphid system is that, at the beginning of summer, aphid populations dwelling on peach trees drop. This occurs because aphids die, or abandon their primary host, or give birth to winged newborns that migrate to secondary herbaceous hosts (55). However, the underlying mechanisms triggering these processes are far from being clear. Our findings suggest that the reduction of resource availability, due to the investment in defensive traits and to photo-period driven interruption of shoot growth, along with the reduction of the phloem nutritional value, due to the accumulation of defensive compounds possibly toxic to the aphid, are the mechanisms most likely to be responsible for the observed patterns. In principle, the crash in aphid population could be due to other factors

such as the arrival of predators attracted by high aphid density (56) or the possible reduction of the phloem nutritional value due to plant ageing (5). However, if the aphid population drop were driven by density dependent mechanisms, one would probably expect to observe fluctuations in the aphid population rather than a constant decline (57) and according to (58) there is no evidence that aphidophagous predators play an effective role in regulating aphid abundance. Moreover, in previous modelling works, it has been shown that observed population trends in different aphid species could be reproduced by considering a per capita death rate positively related to the aphid cumulative population size (59; 60; 61). Such a relationship coherently emerges as a property of our model if the pest presence induces the plant to produce defences that accumulate, and not if the phloem nutritional value declines throughout the season, independently from aphid presence.

Performing experiments to find correct numerical values for parameters of biological models is virtually impossible because many parameters cannot be directly measured. For this reason, we were forced to numerically calibrate nine parameters via our likelihood-based model fitting procedure. However, biologically plausible parameter estimates and good fitting does not guarantee that parameter estimates are correct, due to possible correlations among the parameters (62) and model identifiability problems that can arise due to an imbalance between model complexity and available data (63). The proposed modelling framework would therefore enormously benefit from experimental works dedicated to the measurement, or at least a sound assessment, of some model parameters. Despite the importance of the parameter q in Thornley's models, we found no studies on its assessment.

Similarly, although it is well known that a plant can divert resources from growth to defence (64), we found no quantitative relationships relevant to the cost of making defences (parameter α in our model) in terms of growth loss, neither between the presence of defences and pest performances. Our exercise provides a preliminary assessment of these parameters that need to be confirmed or confuted by dedicated field and/or laboratory works.

The role played by fertilization and irrigation

Variations in plant growth, and in the concentration of C and N substrates in plant tissues, for different levels of fertilization and/or irrigation are well acknowledged (65; 48) and they have already been shown to be emerging properties of the original model for plant growth used in this work (28). Our pest-plant model maintains these properties (Fig. 4A-B-C-D-E-F) and allows further insights regarding the variations observed in aphid population. The aphid population response to fertilization and irrigation has been explored in a number of empirical works not providing a straightforward picture. Some authors observed no effect of fertilization in the wheat-Russian wheat aphid system (66), or negative effects of irrigation in the apple-rosy apple aphid and in the cotton-cotton aphid systems, respectively (67; 68). Other authors observed the highest aphid abundance at an average level of fertilization, and no effect of irrigation, in the chrysanthemum-cotton aphid system (69). The intrinsic rate of oat aphid population increase in three grass species was observed to be favoured by irrigation (70). On the other hand, aphid population was observed to be maximal for moderate water stress in

the cabbage-green aphid and cabbage-cabbage aphid systems (16), and in one out of three genotypes tested for the poplar-wolly poplar aphid system (71). Our model, parametrized for the peach-green aphid system, shows that all these apparently contrasting empirical evidences can emerge from the same biological principles governing plant-pest dynamics and that both plant vigour and plant stress hypotheses can find support when observing a plant-pest system evolving in time and subject to different level of changes in the environment conditions. The aphid population dynamics reproduced by our model (Fig. 5) indicate that the effect of fertilization and irrigation on the pest population cannot be simply reduced as positive or negative In fact, its sign and strength depends on the considered levels of fertilization/irrigation and on the date of observation along the growing season. The contribution of our work is to show how a new synthesis of the experimental data can emerge by using mechanistic modelling. The challenge for our future work is to show how this insight – as well as the model developed here – can be used to inform practical decision making by farmers and growers.

Aknowledgments

The field work to create the dataset used in this work was funded by the ARIMNET (ANR-12-AGR-0001) 'APMed' project (Apple and Peach in-Mediterranean orchards) and the 'RegPuc' project (Quelles stratégies d'irrigation et de fertilisation pour réguler les populations de pucerons verts en vergers de pêcher). The PhD grant of MZ is funded by the PACA region (Provence-Alpes-Côtes d'Azur) and INRA.

References

[1] N. J. Hawkins, C. Bass, A. Dixon, and P. Neve, "The evolutionary origins of pesticide resistance," *Biological Reviews*, vol. 94, no. 1, pp. 135–155, 2019.

[2] S. R. Gliessman, *Agroecology: The Ecology of Sustainable Food Systems*. Boca Raton, USA: CRC Press, third ed., 2015.

[3] P. D. Coley and J. P. Bryant, "Resource availability and plant antiherbivore defense," *Science*, vol. 230, no. 4728, pp. 895–899, 1985.

[4] M. Kytö, P. Niemelä, and S. Larsson, "Insects on Trees : Population and Individual Response to Fertilization," *Oikos*, vol. 75, no. 2, pp. 148–159, 1996.

[5] C. S. Awmack and S. R. Leather, "Host plant quality and fecundity in herbivorous insects," *Annual Review of Entomology*, vol. 47, pp. 817–44, 2002.

[6] C. Girousse, R. Bournoville, and J. L. Bonnemain, "Water deficit-induced changes in concentrations in proline and some other amino acids in the phloem sap of alfalfa," *Plant Physiology*, vol. 111, no. 1, pp. 109–113, 1996.

[7] S. Sevanto, "Phloem transport and drought," *Journal of Experimental Botany*, vol. 65, no. 7, pp. 1751–1759, 2014.

[8] A. Rousselin, M. H. Sauge, M. O. Jordan, G. Vercambre, F. Lescourret, and D. Bevacqua, "Nitrogen and water supplies affect peach tree-green

peach aphid interactions: the key role played by vegetative growth," *Agricultural and Forest Entomology*, vol. 18, no. 4, pp. 367–375, 2016.

[9] A. Rousselin, D. Bevacqua, G. Vercambre, M. Sauge, F. Lescourret, and M. Jordan, "Rosy apple aphid abundance on apple is shaped by vegetative growth and water status," *Crop Protection*, vol. 105, pp. 1–9, 2018.

[10] P. Han, N. Desneux, C. Becker, R. Larbat, J. Le Bot, S. Adamowicz, J. Zhang, and A. V. Lavoir, "Bottom-up effects of irrigation, fertilization and plant resistance on Tuta absoluta: implications for Integrated Pest Management," *Journal of Pest Science*, vol. 92, no. 4, pp. 1359–1370, 2019.

[11] P. W. Price, "The Plant Vigor Hypothesis and Herbivore Attack," *Oikos*, vol. 62, no. 2, p. 244, 1991.

[12] A. F. Huberty and R. F. Denno, "Consequences of nitrogen and phosphorus limitation for the performance of two planthoppers with divergent life-history strategies," *Oecologia*, vol. 149, no. 3, pp. 444–455, 2006.

[13] G. Tamburini, S. van Gils, M. Kos, W. van der Putten, and L. Marini, "Drought and soil fertility modify fertilization effects on aphid performance in wheat," *Basic and Applied Ecology*, vol. 30, pp. 23–31, 2018.

[14] T. C. R. White, "The abundance of invertebrate herbivores in relation to the availability of nitrogen in stressed food plants," *Oecologia*, vol. 63, no. 1, pp. 90–105, 1984.

[15] C. J. Oswald and M. J. Brewer, "Aphid-barley interactions mediated by water stress and barley resistance to Russian wheat aphid (Homoptera: Aphididae)," *Environmental Entomology*, vol. 26, no. 3, pp. 591–602, 1997.

[16] M. Tariq, D. J. Wright, J. T. Rossiter, and J. T. Staley, "Aphids in a changing world: Testing the plant stress, plant vigour and pulsed stress hypotheses," *Agricultural and Forest Entomology*, vol. 14, no. 2, pp. 177–185, 2012.

[17] J. H. M. Thornley and I. R. Johnson, *Plant and Crop Modelling.* Oxford, UK: Oxford: Claredon, 1990.

[18] P. K. Aggarwal, N. Kalra, S. Chander, and H. Pathak, "InfoCrop: A dynamic simulation model for the assessment of crop yields, losses due to pests, and environmental impact of agro-ecosystems in tropical environments. I. Model description," *Agricultural Systems*, vol. 89, no. 1, pp. 1–25, 2006.

[19] L. Willocquet, J. N. Aubertot, S. Lebard, C. Robert, C. Lannou, and S. Savary, "Simulating multiple pest damage in varying winter wheat production situations," *Field Crops Research*, vol. 107, no. 1, pp. 12–28, 2008.

[20] M. C. Dietze and J. H. Matthes, "A general ecophysiological framework for modelling the impact of pests and pathogens on forest ecosystems," *Ecology Letters*, vol. 17, no. 11, pp. 1418–1426, 2014.

[21] R. Levins and B. B. Schultz, "Effects of Density Dependence , Feedback and Environmental Sensitivity on Correlations Among Predators, Prey and Plant Resources: Models and Practical Implications," *Journal of Animal Ecology*, vol. 65, pp. 802–812, 1996.

[22] S. Bewick, R. S. Cantrell, C. Cosner, and W. F. Fagan, "How Resource Phenology Affects Consumer Population Dynamics," *The American Naturalist*, vol. 187, no. 2, pp. 151–166, 2016.

[23] D. Bevacqua, I. Grechi, M. Génard, and F. Lescourret, "The consequences of aphid infestation on fruit production become evident in a multi-year perspective: Insights from a virtual experiment," *Ecological Modelling*, vol. 338, pp. 11–16, 2016.

[24] N. J. Cunniffe, R. C. Cobb, R. K. Meentemeyer, D. M. Rizzo, and C. A. Gilligan, "Modeling when, where, and how to manage a forest epidemic, motivated by sudden oak death in California," *Proceedings of the National Academy of Sciences*, vol. 13, no. 20, pp. 5640–5645, 2016.

[25] S. R. Hyatt-Twynam, S. Parnell, R. O. J. H. Stutt, T. R. Gottwald, C. A. Gilligan, and N. J. Cunniffe, "Risk-based management of invading plant disease," *New Phytologist*, vol. 214, no. 3, pp. 1317–1329, 2017.

[26] F. F. Laranjeira, S. X. B. Suely, R. E. Murray-Watson, A. C. F. Soares, H. P. Santos-Filho, and N. J. Cunniffe, "Spatiotemporal dynamics and modelling support the case for area-wide management of citrus greasy spot in a Brazilian smallholder farming region," *Plant Pathology*, vol. 69, no. 3, pp. 467–483, 2020.

[27] J. A. Newman, D. J. Gibson, A. J. Parsons, and J. H. M. Thornley, "How predictable are aphid population responses to elevated co2?," *Journal of Animal Ecology*, vol. 72, no. 4, pp. 556–566, 2003.

[28] J. H. M. Thornley, "A Balanced Quantitative Model for Root: Shoot Ratios in Vegetative Plants," *Annals of Botany*, vol. 36, no. 2, pp. 431–441, 1972.

[29] J. H. M. Thornley, "Modelling Water in Crops and Plant Ecosystems," *Annals of Botany*, vol. 77, no. 3, pp. 261–275, 1996.

[30] J. H. M. Thornley, "Modelling Shoot : Root Relations : the Only Way Forward ?," *Annals of Botany*, vol. 81, no. 1994, pp. 165–171, 1998.

[31] J. H. M. Thornley and M. G. R. Cannel, "Modelling the Components of Plant Respiration: Representation and Realism," *Annals of Botany*, vol. 85, no. 6, pp. 55–67, 2000.

[32] T. Zust and A. A. Agrawal, "Trade-Offs Between Plant Growth and Defense Against Insect Herbivory: An Emerging Mechanistic Synthesis," *Annual Review of Plant Biology*, vol. 68, no. 1, pp. 513–534, 2017.

[33] E. L. Zvereva, V. Lanta, and M. V. Kozlov, "Effects of sap-feeding insect herbivores on growth and reproduction of woody plants: A meta-analysis of experimental studies," *Oecologia*, vol. 163, no. 4, pp. 949–960, 2010.

[34] J. Ng and K. Perry, "Transmission of plant viruses by aphid vectors," *Molecular Plant Pathology*, pp. 505–511, 2004.

[35] O. M. Heide, "Interaction of photoperiod and temperature in the control of growth and dormancy of *Prunus* species," *Scientia Horticulturae*, vol. 115, no. 3, pp. 309–314, 2008.

[36] T. O. Perry, "Dormancy of trees in winter.," *Science*, vol. 171, no. 3966, pp. 29–36, 1971.

[37] T. Zust and A. A. Agrawal, "Mechanisms and evolution of plant resistance to aphids," *Nature Plants*, vol. 2, no. 1, pp. 1–9, 2016.

[38] F. L. Goggin, "Plant-aphid interactions: molecular and ecological perspectives," *Current Opinion in Plant Biology*, vol. 10, no. 4, pp. 399–408, 2007.

[39] A. F. G. Dixon, *Aphid Ecology: an optimization approach*. New York, USA: Springer Science+Business Media New York, second ed., 1998.

[40] A. F. G. Dixon, "Structure of aphid populations," *Annual Review of Entomology*, vol. 30, no. 30, pp. 155–74, 1985.

[41] C. B. Müller, I. S. Williams, and J. Hardie, "The role of nutrition, crowding and interspecific interactions in the development of winged aphids," *Ecological Entomology*, vol. 26, no. 3, pp. 330–340, 2001.

[42] T. Will, A. C. U. Furch, and M. R. Zimmermann, "How phloem-feeding insects face the challenge of phloem-located defenses," *Frontiers in Plant Science*, vol. 4, pp. 1–12, 2013.

[43] M. Vyska, N. Cunniffe, and C. Gilligan, "Trade-off between disease resistance and crop yield: a landscape-scale mathematical modelling per-

spective," *Journal of the Royal Society - Interface*, vol. 13, no. 123, pp. 1–9, 2016.

[44] K. J. Medina-Ortega and G. P. Walker, "Does aphid salivation affect phloem sieve element occlusion in vivo?," *Journal of Experimental Botany*, vol. 64, no. 18, pp. 5525–5535, 2013.

[45] E. van Velzen and R. S. Etienne, "The importance of ecological costs for the evolution of plant defense against herbivory," *Journal of Theoretical Biology*, vol. 372, pp. 89–99, 2015.

[46] J. P. Kreiss and S. N. Lahiri, "Bootstrap Methods for Time Series," in *Handbook of statistics:Time Series Analysis: Methods and Applications*, vol. 30, pp. 3–26, Elsevier, 2012.

[47] B. Efron, "Bootstrap Methods: Another Look at the Jakknife," *The Annals of Statistics*, vol. 7, no. 1, pp. 1–26, 1979.

[48] B. Muller, F. Pantin, M. Génard, O. Turc, S. Freixes, M. Piques, and Y. Gibon, "Water deficits uncouple growth from photosynthesis, increase C content, and modify the relationships between C and growth in sink organs," *Journal of Experimental Botany*, vol. 62, no. 6, pp. 1715–1729, 2011.

[49] D. J. Connor, R. S. Loomis, and K. Cassman, *CROP ECOLOGY: Productivity and management in agricultural systems.* New York, USA: Cambridge University Press, second ed., 2011.

[50] J. H. M. Thornley and A. J. Parsons, "Allocation of new growth between

shoot, root and mycorrhiza in relation to carbon, nitrogen and phosphate

supply: Teleonomy with maximum growth rate," *Journal of Theoretical*

Biology, vol. 342, pp. 1–14, 2014.

[51] H. Akaike, "A New Look at the Statistical Model Identification," *IEEE*

Transactions on Automatic Control, vol. 19, no. 6, pp. 716–723, 1974.

[52] Y. Grossman and T. DeJong, "PEACH: A simulation model of repro-

ductive and vegetative growth in peach trees," *Tree Physiology*, vol. 14,

no. 4, pp. 329–345, 1994.

[53] I. Mewis, H. M. Appel, A. Hom, R. Raina, and J. C. Schultz, "Major sig-

naling pathways modulate *Arabidopsis glucosinolate* accumulation and

response to both phloem-feeding and chewing insects," *Plant Physiology*,

vol. 138, no. 2, pp. 1149–1162, 2005.

[54] J. H. Kim, B. W. Lee, F. C. Schroeder, and G. Jander, "Identifica-

tion of indole glucosinolate breakdown products with antifeedant effects

on *Myzus persicae* (green peach aphid)," *Plant Journal*, vol. 54, no. 6,

pp. 1015–1026, 2008.

[55] A. Dixon, "Aphid ecology," *Ann. Rev. Ecol. Syst.*, vol. 8, no. 47, pp. 329–

53, 1977.

[56] A. C. Costamagna and D. A. Landis, "Predators exert top-down control

of soybean aphid across a gradient of agricultural management systems,"

Ecological Applications, vol. 16, p. 1619–1628, 2006.

[57] W. W. Murdoch, "Population regulation in theory and practice," *Ecology*, vol. 2, no. 75, pp. 271–287, 1994.

[58] A. F. G. Dixon, *Insect Herbivore-Host Dynamics: Tree-Dwelling Aphids*. Cambridge, U.K.: Cambridge University Press, 2005.

[59] J. H. Matis, T. R. Kiffe, T. I. Matis, and D. E. Stevenson, "Nonlinear stochastic modeling of aphid population growth," *Mathematical Biosciences*, vol. 198, no. 2, pp. 148–168, 2005.

[60] J. H. Matis, T. R. Kiffe, W. van der Werf, A. C. Costamagna, T. I. Matis, and W. E. Grant, "Population dynamics models based on cumulative density dependent feedback: A link to the logistic growth curve and a test for symmetry using aphid data," *Ecological Modelling*, vol. 220, no. 15, pp. 1745–1751, 2009.

[61] J. H. Matis, T. R. Kiffe, T. I. Matis, and C. Chattopadhyay, "Generalized aphid population growth models with immigration and cumulative-size dependent dynamics," *Mathematical Biosciences*, vol. 215, no. 2, pp. 137–143, 2008.

[62] P. Li and Q. D. Vu, "Identification of parameter correlations for parameter estimation in dynamic biological models," *BMC Systems Biology*, vol. 7, no. 1, p. 91, 2013.

[63] D. J. W. De Pauw, K. Steppe, and B. De Baets, "Identifiability analysis and improvement of a tree water flow and storage model," *Mathematical Biosciences*, vol. 211, no. 2, pp. 314–332, 2008.

[64] D. A. Herms and W. J. Mattson, "The dilemma of plants: to grow or defend," *The Quarterly Review of Biology*, vol. 67, no. 3, pp. 283–335, 1992.

[65] Y. Jia and V. M. Gray, "Influence of phosphorus and nitrogen on photosynthetic parameters and growth in *Vicia faba* L.," *Photosynthetica*, vol. 42, no. 4, pp. 535–542, 2004.

[66] T. L. Archer, E. D. Bynum, A. B. Onken, and C. W. Wendt, "Influence of water and nitrogen fertilizer on biology of the Russian wheat aphid (Homoptera: Aphididae) on wheat," *Crop Protection*, vol. 14, no. 2, pp. 165–169, 1995.

[67] T. I. Matis, M. N. Parajulee, J. H. Matis, and R. B. Shrestha, "A mechanistic model based analysis of cotton aphid population dynamics data," *Agricultural and Forest Entomology*, vol. 10, no. 4, pp. 355–362, 2008.

[68] A. Rousselin, D. Bevacqua, M. H. Sauge, F. Lescourret, K. Mody, and M. Jordan, "Harnessing the aphid life cycle to reduce insecticide reliance in apple and peach orchards. A review," *Agronomy for Sustainable Development*, vol. 37, no. 5, 2017.

[69] J. A. Bethke, R. A. Redak, and U. K. Schuch, "Melon aphid performance on chrysanthemum as mediated by cultivar, and differential levels of fertilization and irrigation," *Entomologia Experimentalis et Applicata*, vol. 88, no. 1, pp. 41–47, 1998.

[70] B. K. Hale, J. S. Bale, J. Pritchard, G. J. Masters, and V. K. Brown, "Effects of host plant drought stress on the performance of the bird

cherry-oat aphid, *Rhopalosiphum padi* (L.): A mechanistic analysis," *Ecological Entomology*, vol. 28, no. 6, pp. 666–677, 2003.

[71] F. Dardeau, A. Berthier, M. Feinard-Duranceau, F. Brignolas, F. Laurans, F. Lieutier, and A. Sallé, "Tree genotype modulates the effects of water deficit on a plant-manipulating aphid," *Forest Ecology and Management*, vol. 353, pp. 118–125, 2015.

[72] L. M. Schoonhoven, J. J. A. Van Loon, and M. Dicke, *Insect-Plant Biology.* Oxford, UK: Oxford University Press Inc., second ed., 2005.

[73] M. F. Day and H. Irzykiewicz, "Feeding behaviour of the aphids *Myzus persicae* and *Brevicoryne brassicae*, studied with radiophosphorus.," *Australian journal of biological sciences*, vol. 6, no. 1, pp. 98–108, 1953.

[74] J. Saguez, R. Hainez, A. Cherqui, O. Van Wuytswinkel, H. Jeanpierre, G. Lebon, N. Noiraud, A. Beaujean, L. Jouanin, J. C. Laberche, C. Vincent, and P. Giordanengo, "Unexpected effects of chitinases on the peach-potato aphid (*Myzus persicae* Sulzer) when delivered via transgenic potato plants (*Solanum tuberosum* Linné) and in vitro," *Transgenic Research*, vol. 14, no. 1, pp. 57–67, 2005.

[75] A. C. Gange, E. Bower, and V. K. Brown, "Positive effects of an arbuscular mycorrhizal fungus on aphid life history traits," *Oecologia*, vol. 120, pp. 123–131, 1999.

 # Figures

Figure 1: Schematic representation of the plant-aphid model where the plant is constituted by shoot (S) and root (R) structural dry mass, carbon (C_i) and nitrogen (N_i) substrates in shoots (i = S) and roots (i = R). The aphid population (A) intercepts a fraction of substrates allocated to constitute shoot structural mass and the plant diverts shoot substrates (carbon and nitrogen) to produce defensive compounds (D). More details are given in the main text.

Figure 2: Schematic representation of the mechanisms considered in the different models Mi ($i \in [1, 10]$) nested in eq.1: *i)* density dependent aphid migration (ω), *ii)* plant induced defences development (α) and *iii)* effect of induced defences on phloem availability to aphids (β_1) and on phloem toxicity (β_2). When the model parameter is set to zero, the relevant mechanism is ignored. Each model can be based on different hypotheses about the variation of the nitrogen assimilation rate σ_N (equal (A, C) or different (B, D) across fertilization treatments) and the substrates utilization rate k(equal (A, B) or different (C, D) across irrigation treatments)

Figure 3: Observed (black points) and predicted (black lines) values of average shoot dry mass (top row), average aphid abundance per shoot (central row) and induced defences concentration (bottom row) under different fertilization and irrigation treatments: high fertilization and irrigation (A-E-I), high fertilization and low irrigation (B-F-J), low fertilization and high irrigation (C-G-K), low fertilization and irrigation (D-H-L). Grey shaded areas indicate the predicted 99% confidence bands.

Figure 4: Simulated effect of fertilization (left column) and irrigation (right column) on the plant-aphid system: average shoot dry mass (A, B), carbon (C, D) and nitrogen (E, F) substrate concentration in shoots, defences concentration in shoot (G, H), aphid abundance (I, J) and density (K, L). Lines colour identifies fertilization (or irrigation) level: very low (red), low (orange), average (green), high (light blue), very high (blue).

Figure 5: Simulated effect of fertilization (A, C, E) and irrigation (B, D, F) on aphids density on May 15[th], June 1[st] and June 15[th]. Boxes represent the first and third quartiles [25% and 75%] with a line inside indicating the median of ten simulated replicates of each treatment. The whiskers extend $\pm$ 1.5 $\times$ the interquartile range (75th percentile – 25th percentile) from the third and first quartiles. Values outside the whiskers are considered outliers and plotted individually using the '+' symbol.

724 Table

Table 1: Model variables and parameters. For those parameters calibrated in the present work calibrated confidence interval of 99% are reported in brackets.

Variable	Dim.	Description
S	g	Shoot structural dry mass
R	g	Root structural dry mass
C_S	g	Shoot carbon substrate
C_R	g	Root carbon substrate
N_S	g	Shoot nitrogen substrate
N_R	g	Root nitrogen substrate
D	DU	Plant induced defences
A	ind.	Aphid population

Parameter	Value	Dim.	Description	Source
C and N assimilation				
σ_C	0.1	d^{-1}	Assimilation rate of C	(1)
σ_N	6.70 - 17.08 10^{-3} [1.76-19.54] [4.56-43.04] 10^{-3} [a]	d^{-1}	Assimilation rate of N	(6)
ν	1000	g	Shoot (root) mass halving substrate assimilation due to self shading (competition)	(1)
ι_C	0.1	/	Semi-saturation C concentration	(1)
ι_N	0.01	/	Semi-saturation N concentration	(1)
C and N substrates allocation to plant growth				
φ_C	0.50	/	Unit of substrate C per unit of structural dry mass	(1)
φ_N	$2.50 \, 10^{-2}$	/	Unit of substrate N per unit of structural dry mass	(1)
κ	142 - 223 [56-601] [48-876] [b]	d^{-1}	Maximum rate of substrate utilization	(6)
η	73	/	Switch-off function of plant growth: steepness	(7)
λ	169	d	Switch-off function of plant growth: date of equal partitioning between growth and reserves	(7)
Transport				
q	0.86 [0.84-0.89]	/	Plant architecture scaling parameter	(6)
Defences development				
α	0.02 [0.01-0.03]	g d^{-1} ind^{-1}	Allocation of substrates to defences per unit of aphid	(6)
ε_C	$5 \, 10^{-2}$	DU g^{-1}	Conversion efficiency of C substrate in defences	(2)
ε_N	1	DU g^{-1}	Conversion efficiency of N substrate in defences	(2)
Aphid				
θ	$1.12 \, 10^{-3}$	g d^{-1} ind^{-1}	Maximum food intake per aphid	(3)
ξ	171	ind g^{-1}	Maximum conversion efficiency of ingested food into descendants	(4)
μ	0.04	d^{-1}	Aphid natural mortality rate	(5)
π_1	$8.52 \, 10^{-3}$ [3.70-23.12] 10^{-3}	DU g^{-1}	Swich-on function of defences protected phloem fraction: defences concentration at which defences effect is half-saturated	(6)
δ_1	0.65 [0.55-0.73]	/	Swich-on function of defences protected phloem fraction: steepness	(6)
π_2	0.05 [0.03-0.08]	DU g^{-1}	Swich-on function of defences induced mortality/repulsion rate: defences concentration at which defences effect is half-saturated	(6)
δ_2	118 [27-251]	/	Swich-on function of defences induced mortality/repulsion rate: steepness	(6)
β_2	28 [13-76]		Swich-on function of defences induced mortality/repulsion rate: asymptotic value	(6)

Sources: (1) (30); (2) (72); (3) (73); (4) (74); (5) (75); (6) Calibrated in the present work; (7) Fixed in the present work, see SI.
[a] values refer to different fertilization treatments.
[b] Values refer to different irrigation treatments