1 **A fibronectin mechanotransduction pathway translates segmentation clock oscillations into**

2 **periodic somite formation**

Patrícia Gomes de Almeida[1,2,3], Pedro Rifes[1,#], Ana Patrícia Martins-Jesus[2,3], Gonçalo G. Pinheiro[1,2],

Raquel P. Andrade[2,3,4] and Sólveig Thorsteinsdóttir[1,*]

[1] Centre for Ecology, Evolution and Environmental Change – cE3c, Departmento de Biologia

Animal, Faculdade de Ciências, Universidade de Lisboa, 1740-016 Lisboa, Portugal

[2] Centre for Biomedical Research - CBMR, Universidade do Algarve, 8005-139 Faro, Portugal

[3] Algarve Biomedical Center, Campus Gambelas, Edificio 2. Ala Norte, 8005-139 Faro, Portugal

[4] Department of Medicine and Biomedical Sciences, Universidade do Algarve, 8005-139 Faro,

Portugal

[#]Present address: The Novo Nordisk Foundation Center for Stem Cell Biology (DanStem), University

of Copenhagen, 2200 Copenhagen N, Denmark

*Correspondence to: Sólveig Thorsteinsdóttir (solveig@fc.ul.pt)

Abstract:

Somitogenesis starts with cyclic waves of expression of segmentation clock genes in the presomitic

mesoderm (PSM) and culminates with periodic budding of somites in its anterior-most region. How

cyclic clock gene expression is translated into timely morphological somite formation remains

unclear. A posterior to anterior gradient of increasing fibronectin matrix complexity surrounds the

PSM, suggesting that the mechanical properties of the matrix may play a role in regulating this

transition. Here we interfered with the chick PSM fibronectin mechanotransduction pathway by

perturbing (1) actomyosin-mediated contractility, (2) Rho-associated protein kinase function, (3)

integrin-fibronectin binding and (4) fibronectin matrix assembly. Each treatment perturbed

segmentation clock expression dynamics, *meso1* expression and resulted in defective somitic

clefts. A model is presented where the fibronectin matrix gradient participates in the PSM

wavefront of maturation by activating a mechanotransduction signaling cascade ensuring the

correct spatio-temporal conversion of cyclic segmentation clock gene expression into periodic

somite formation.

Keywords:

Fibronectin, Mechanotransduction, Segmentation Clock, Cleft Formation, Somitogenesis

Introduction:

Cells in the developing embryo are constantly receiving and integrating information, including tension-derived signals generated by the adhesion to neighboring cells and/or the surrounding extracellular matrix (ECM). Cell-surface integrins bind to ECM ligands in the extracellular space and connect the ECM to the intracellular actomyosin cytoskeleton through adaptor proteins (Campbell and Humphries, 2011; Wolfenson et al., 2013). These multimolecular complexes (i.e. focal adhesions) act as mechanotransduction centers allowing cells to perceive the chemical and physical properties of the ECM, such as its molecular composition, density and stiffness. While morphogens have been extensively studied as major chemical regulators of developmental processes (Marek and Kubícek, 1981; Slack, 1987; Tiedemann, 1976), the importance of mechanical forces in embryo development has, until recently, been largely overlooked (Chan et al., 2017; Merle and Farge, 2018). In fact, the ability of cells to sense and respond to mechanical signals has been shown to regulate critical developmental processes, including cell proliferation, migration and differentiation (Barriga et al., 2018; Cosgrove et al., 2016; Smutny et al., 2017; Wolfenson et al., 2016).

One of the most conspicuous morphogenetic events during early vertebrate embryogenesis is the formation of somites, which are the source of axial skeleton and skeletal muscle precursor cells (Christ et al., 2007). Somites are spheres of epithelioid cells and are formed periodically from the anterior portion of the mesenchymal presomitic mesoderm (PSM), bilateral to the axial structures (Bailey and Dale, 2015). Temporal control of somite formation is dependent on cyclic waves of expression of segmentation clock genes, many of which are targets of the Notch signaling pathway (Dequéant et al., 2006; Masamizu et al., 2006; Palmeirim et al., 1997). These waves periodically sweep the PSM in a posterior to anterior direction (Aulehla and Pourquié, 2010) and, as they reach the anterior PSM, oscillations first slow down and then arrest (Morimoto et al., 2005; Shih et al., 2015). The transcription factor Mesp2/Meso1 is upregulated downstream of the segmentation clock in the anterior PSM, leading to Eph/Ephrin signaling and somitic cleft

formation (Barrios et al., 2003; Nakajima et al., 2006; Saga, 2012; Watanabe et al., 2009), followed

by progressive cell rearrangements into a somite (Martins et al., 2009; Morimoto et al., 2005; Shih

et al., 2015).

The PSM of vertebrate embryos is surrounded by a fibronectin-rich ECM, and both

fibronectin and its integrin receptors are essential for somite formation (George et al., 1993;

Georges-Labouesse et al., 1996; Goh et al., 1997; Jülich et al., 2005; Koshida et al., 2005; Kragtorp

and Miller, 2007; Rifes et al., 2007; Takahashi et al., 2007; Yang et al., 1999, 1993). Moreover,

perturbing fibronectin binding to its integrin receptors, by substituting the fibronectin RGD binding

site with an RGE sequence (mouse $Fn1^{RGE/RGE}$), causes dramatic somitogenesis defects (Girós et al.,

2011; Takahashi et al., 2007). Indeed, fibronectin provides the structural support for cells to attach,

polarize, change shape and position, required for somite epithelialization (Martins et al., 2009).

We have previously shown that the fibronectin matrix assembled around the chick PSM

becomes progressively denser and more complex as the tissue matures (Rifes and

Thorsteinsdóttir, 2012), and have proposed that this posterior to anterior gradient culminates in

a fibronectin-dependent matrix assembly state that supports somite epithelialization in the

anterior PSM (Rifes et al., 2007; Rifes and Thorsteinsdóttir, 2012). More recently, adhesion to a

fibronectin substrate was noted as a regulator of *Lfng* oscillations in cultured mouse tailbud cells

(Hubaud et al., 2017; Lauschke et al., 2013) and cell adhesion to fibronectin was linked to

dampening and eventual arrest of *Lnfg* oscillations (Hubaud et al., 2017), reminiscent of what is

observed in the anterior PSM prior to somite formation. Nevertheless, how PSM cells sense and

respond to an increasing fibronectin matrix complexity gradient, and how it affects segmentation

clock oscillations *in vivo* remains unknown.

In this study, we addressed the involvement of the fibronectin ECM and its downstream

mechanotransduction pathway in the regulation of both segmentation clock dynamics and

subsequent somite formation *in vivo*, using the chick embryo as a model. We experimentally

perturbed (1) non-muscle myosin II (NM II) ATPase activity (2), Rho-associated protein kinase

(ROCK), (3) integrin-fibronectin binding through RGD and (4) extracellular fibronectin matrix

assembly, and found that all these treatments resulted in abnormal segmentation clock

oscillations, mis-positioning of *meso1* expression in the rostral PSM and perturbations in somite

morphogenesis. These results strongly suggest that tissue tension generated by the fibronectin

matrix surrounding the PSM plays an important role in converting segmentation clock oscillations

into periodic somite formation.

Results:

The intracellular mechanotransduction machinery of PSM cells regulates segmentation clock

oscillations and *meso1* activation

In the chick embryo, sequential pairs of somites bud off from the anterior PSM every 90

min, which corresponds to the period of segmentation clock oscillations (Figure 1 A). To investigate

the involvement of intracellular actomyosin contractility in this process, the expression of the

segmentation clock gene *hairy1* (Palmeirim et al., 1997), a target of Notch signaling, was analyzed

in the PSM of embryo half explants cultured in the presence of either Blebbistatin, which directly

inhibits the ATPase activity of NM II (direct inhibition) and consequently all actomyosin

contractility, or RockOut (indirect inhibition), a chemical inhibitor of ROCK I and II (ROCK I/II)

enzymes involved in activating NM II (Figure 1 B, C; Ringer et al., 2017; Straight et al., 2003; Yarrow

et al., 2005). The contralateral control sides were cultured with an equal volume of DMSO.

Explants cultured for 6 hours in each experimental condition presented significantly

altered *hairy1* expression. *hairy1* expression was either absent or in a different phase of the cycle

relative to the contralateral control in 80% of the Blebbistatin- (n=7/9) and RockOut-treated

explants (n= 8/10; Figure 1 D, a-c), suggesting that temporal control of *hairy1* oscillations requires

the generation and transduction of tensional cues mediated by NM II and ROCK I/II activity.

Figure 1. Blocking NM II or ROCK I/II activity leads to misregulation of *hairy1*, *meso1* and *dll1* expression and to defects in somite formation.

(A) Schematic representation of chick PSM maturation and somite formation over time. A new pair of somites buds off from the anterior PSM every 90 min. This is also the period of segmentation clock oscillations in the chick embryo. With each new pair of somites, the previously formed somites mature (SI becomes SII, SII becomes SIII, etc). (**B**) Schematic representation of the explant culture system. Posterior explants of HH11-14 chick embryos were bisected along the midline and cultured for 6 (or 10.5) hours. One side of the explant was cultured with either Blebbistatin (Blebb) or RockOut, while the contra-lateral half was cultured with an equal volume of DMSO. **(C)** Schematic representation of the action of Blebbistatin and RockOut. Blebbistatin inhibits NM II ATPase activity directly while RockOut inhibits ROCK I/II-mediated phosphorylation of myosin light chain, thus indirectly decreasing NM II ATPase activity. (**D**) *In situ* hybridization for *hairy1* (a, b) *meso1* (d, e) and *dll1* (**g, h**) after 6 (a, b, d, e) or 10.5 hours of culture (g, h) in Blebbistatin (Blebb; a, d, g) and in RockOut-containing (b, e, h) media. Straightened images of the respective explant pairs (right) were aligned by SIV. Rostral is on top. Percentage of Blebbistatin- or RockOut-treated explants with altered *hairy1*, *meso1* and *dll1* expression compared to the contralateral controls is shown in c, f and i, respectively. **(E)** Number of somites (black bars) or somite-like condensations (dotted bars) formed in cultured explants. Explants cultured with DMSO formed sharp somite boundaries and clearly individualized somites. Blebbistatin-treated explants only formed 1-2 somite-like condensations. In RockOut treated explants the first two somites were normal while the remaining ones were cell condensations with poorly defined boundaries. ns – not significant, *** – p<0.01. Scale bar in D**:** 500 µm. Bars - standard error of the mean.

Segmentation clock oscillations are required for the correct spatial and temporal

upregulation of *Mesp2* in the anterior PSM (Niwa et al., 2011; Saga and Takeda, 2001; Sato et al.,

2002), which regulates downstream targets required for the formation of the somitic cleft (Saga,

2012). The expression of the chick *Mesp2* homolog, *meso1*, was altered in Blebbistatin- (n=9/11)

and RockOut-treated explants (n=10/10; Figure 1 D, d-f). *meso1* expression was either absent,

located more rostrally or presented a different number of bands of expression, clearly indicating

that the normal cycles of activation and suppression of *meso1* in the rostral PSM were altered.

Importantly, *meso1* expression was also perturbed after 3 hours in culture with Blebbistatin or

RockOut (n=8/9 and 5/5, respectively; Supplementary Figure 1 A-B), corresponding to an effect

within two segmentation clock cycles. Furthermore, timely downregulation of *dll1* in the anterior-

most PSM (Palmeirim et al., 1998), which normally occurs downstream of Meso1/Mesp2 activity

(Takahashi et al., 2000; Takahashi et al., 2003), was not observed in either Blebbistatin- (80%,

n=8/10) or RockOut-treated (75%, n=6/8) explants after 10.5h of culture (Figure 1 D, g-i). Together

these data indicate that interfering with actomyosin contractility perturbs three sequential events:

the spatio-temporal expression dynamics of *hairy1*, timely *meso1* expression and the

downregulation of *dll1* expression in the anterior-most PSM.

Alterations in somite formation were observed concomitantly with the perturbations in

hairy1 and *meso1* expression. Control explants formed an average of 3.6 somites after 6 hours,

consistent with a 90 min periodicity (Figure 1 E), while contralateral RockOut-treated explants

formed 3.1 somites, of which only the first two somites were clearly individualized, while

subsequent somite-like condensations were poorly defined (Figure 1 E). After 10.5 hours, control

explants had formed an average of 7.5 somites, while RockOut-treated explants formed 7.1

somites of which the first two appeared normal, but the remaining ones were ill-defined (Figure 1

E). Importantly, explants cultured with Blebbistatin were unable to form more than 1-2 somite-

like aggregates after 6 hours, or even 10.5 hours, of culture (Figure 1 E), evidencing an absolute

requirement for NM II ATPase activity in somite formation. These effects were not due to an

increase in apoptosis (Supplementary Figure 2 A-D).

Our data reveal a previously unknown role for NM II- and ROCK I/II- mediated

mechanotransduction in the temporal regulation of the segmentation clock, *meso1* expression,

dll1 downregulation and somite formation.

<u>NM II activity is required for somite cleft formation and cell polarization</u>

Somite formation involves a mesenchymal-to-epithelial transition (MET) of anterior PSM

cells (Martins et al., 2009; Saga, 2012). To determine to what extent this process is impaired upon

NM II or ROCK I/II inhibition, we performed a detailed analysis of the morphology of S0 to SIII in

explants after a 6 hour culture period (regions e-b in Figure 1 A).

In control explants, S0 showed apically enriched N-cadherin (Figure 2 A, D) and some

zonula occludens protein 1 (ZO-1) accumulation was observed apically (Figure 2 B, D, arrowheads).

Peripheral cell alignment occurred (Figure 2 C, D) and fibronectin matrix was detected in the

nascent somitic clefts (Figure 2 E, F, arrows). In contrast, in explants cultured with Blebbistatin, N-

cadherin was homogeneous (Figure 2 G, J), ZO-1 immunostaining was absent (Figure 2 H, J) and

neither peripheral cell alignment (Figure 2 I, J) nor fibronectin matrix accumulation within the

tissue was observed (Figure 2 K, L). Furthermore, the continuous and dense fibronectin matrix

observed surrounding the rostral PSM and S0 in control explants was disrupted in Blebbistatin-

treated explants (compare Figure 2 E and K, arrowheads). Moreover, the characteristic nuclear

alignment and F-actin apical enrichment observed in control SI (Supplementary Figure 3 A-C) was

absent in Blebbistatin-treated explants and no signs of somitic boundaries could be detected

(Supplementary Figure 3 D-F). We conclude that exposure of the S-IV and S-III regions of the PSM

(regions e and d in Figure 1 A) to Blebbistatin for 6 hours completely blocks their capacity to form

somites.

186
Figure 2. NM II inhibition abolishes N-cadherin and ZO-1 polarization and impairs fibronectin
fibrillogenesis.
(A-X) Sagittal views of control explants (A-F, M-R) and their Blebbistatin-treated contralateral halves after 6
hours of culture, immunostained for N-cadherin (Ncad), ZO-1, fibronectin (FN), stained for DNA and imaged
at S0 (A-L; region e in Figure 1 A) and SIII (M-X; region b in Figure 1 A) levels. S0 of DMSO-treated explants
shows apically enriched N-cadherin (A) and ZO-1 (B, arrowheads) and peripheral nuclei are aligned (C, D),
while no signs of polarized cell-cell adhesions (G, H) or nuclear alignment (I, J) are found in the contralateral
Blebbistatin-treated explant. Apical polarization of N-cadherin and ZO-1 is maintained in SIII cells of DMSO-
treated explants (M, N, arrowheads), but no polarized N-cadherin (S) or ZO-1 (T, arrowheads) are observed
in contralateral Blebbistatin-treated halves. Nuclear alignment (U, arrowheads) and fibronectin assembly
around somites (K, W, arrowheads) are also deficient in Blebbistatin-treated explants compared to
contralateral controls (E, Q, arrowheads). Arrows in E show fibronectin assembly in the nascent somitic
cleft. Rostral to the left and dorsal on top. Dashed lines mark borders of S0 (B, H) and SIII (N, T). FN –
fibronectin. Ncad – N-cadherin. ZO-1 – *Zonula occludens* 1. Blebb – Blebbistatin. Scale bars: 50 µm.
201

We next turned our attention to SII and SIII somites after 6 hours of culture. These were

at stage S-II and S-I in the PSM, respectively (regions c and b in Figure 1 A), when the explants were

placed in culture and had thus already upregulated *meso1* (Buchberger et al., 1998). As before, in

the presence of Blebbistatin, apical enrichment of N-cadherin failed to occur (Figure 2 S, V), ZO-1

was only detected in a few small foci (Figure 2 T, V) and, although a fibronectin matrix was present,

it appeared less dense (Figure 2 W, X). An incipient nuclear alignment was sometimes observed

(Supplementary Figure 3 K, arrowheads), but cells did not polarize their F-actin into apically

enriched adhesion belts (compare Supplementary Figure 3 G-I with J-L). Epithelial tissues other

than somites (e.g. ectoderm and neural tube) did not present significant alterations after

incubation with Blebbistatin (Supplementary Figure 4). Altogether, these results point to an

indispensable role for NM II activity for the MET underlying somite formation.

RockOut-treated explants also showed perturbations in somite formation, although to a

lesser extent (Figure 1 E). When compared to control explants (Figure 3 A-D; Supplementary Figure

5 A-C), RockOut treatment resulted in incomplete somitic clefts, such that S0 shared the

somitocoel with SI and sometimes also with SII (Fig 3 E-H, arrows, Supplementary Figure 5 D-F,

arrows). In contrast to the accumulation of fibronectin in the nascent clefts in controls (Figure 3 B,

arrow), no fibronectin was observed in the incipient somitic clefts of RockOut-treated explants

(Figure 3 F, arrows). These results suggest that ROCK I/II activity in the S-IV and S-III regions of the

PSM is required for the formation of individualized somites. In contrast, when the rostral-most

PSM (stage S-I before culture, region b in Figure 1 A) was exposed to RockOut for 6 hours, it was

undistinguishable from control explants, showing apical accumulation of ZO-1 (Figure 3 I, M,

arrowhead) and N-cadherin (Supplementary Figure 5, G, J), nuclear alignment (Figure 3, K, O,

Supplementary Figure 5, H, K) and a complete, fibronectin matrix-containing cleft (Figure 3 J, N).

This indicates that ROCK I/II activity is not required for S-I to develop into a somite.

Altogether, our data indicate that the intracellular mechanotransduction machinery of

PSM cells is required for periodic somite cleft formation, and that ROCK I/II- independent NM II

activity is essential for MET.

Figure 3. ROCK I/II inhibition impairs morphological somite formation, leading to deficient ZO-1 polarization and fibronectin assembly.

(A-P) Sagittal views of explants cultured in control (DMSO) medium (A-D, I-L) and their contralateral RockOut-treated halves (E-H, M-P) at S0 (A-H; region e in Figure 1 A) and SIII (I-P; region b in Figure 1 A) levels, immunostained for ZO-1 (first column), fibronectin (second column), stained for DNA (third column) and the respective merged image (fourth column). Explants were cultured for 6 hours. S0 in control explants show normal accumulation of ZO-1 (A, arrowhead), fibronectin assembly in the nascent cleft (B, arrow) and nuclear alignment (C, arrowhead). In contrast, S0 in contralateral RockOut-treated explants fails to form a clear cleft (E-H, arrows), although ZO-1 is generally polarized (E, arrowhead) and nuclei are aligned (G, arrowhead). At SIII level, both explants show normal ZO-1 polarization (I, M, arrowheads), fibronectin assembly (J, N) and nuclear alignment (K, O). Rostral on the left and dorsal on top. FN – fibronectin. ZO-1 – *Zonula occludens* protein 1. Scale bar: 50 μm.

Integrin-fibronectin binding regulates segmentation clock oscillations and is required for somitic

cleft positioning and morphogenesis

Fibronectin signaling through α5β1 integrin plays a crucial role during somitogenesis

(Takahashi et al., 2007). To evaluate if this interaction could be regulating segmentation clock

oscillations and somite formation upstream of the intracellular mechanotransduction pathway,

embryo half explants were cultured in the presence of the RGD peptide which competes with

fibronectin for integrin binding (Figure 4 A, B). RGD-treated explants formed ill-defined somite-like

condensations, although in approximately the same number as the contralateral control (Figure 4

C). This was not due to cell death (Supplementary Figure 2 E-F). Concomitantly, RGD-treated

explants displayed alterations in *hairy1* (Figure 4 D, a-c), *meso1* (Figure 4 D, d-f) and *dll1* expression

patterns (Figure 4 D, g-i), evidencing that integrin-RGD interactions are required for proper

segmentation clock oscillations, *meso1* positioning and timely downregulation of *dll1* in the

anterior PSM.

When compared to the contralateral control, the area corresponding to S0 after 6 hours

of culture with RGD (region e in Figure 1 A) showed deficient nuclear alignment (Figure 5 B, E, H,

K, arrowheads) and N-cadherin polarization (Figure 5 A, G, arrowheads), accompanied by deficient

fibronectin assembly in the nascent cleft (Figure 5 D, J arrowheads). At the level of SII (region c in

Figure 1 A), complete somite individualization was impaired in RGD-treated explants (Figure 5 N,

Q, T, W, arrowheads) and, although N-cadherin polarization appeared normal (Figure 5 M, S,

arrowheads), cleft formation (Figure 5 Q, W, arrowheads; R, X) and fibronectin assembly between

adjacent somites was deficient (Figure 5 P, V, arrowheads; R, X).

These findings implicate cell-ECM interactions, mediated by integrin-fibronectin binding,

in temporal control of *hairy1* oscillations, correct positioning of *meso1* expression, downregulation

of *dll1* in the anterior PSM and somite morphogenesis.

Figure 4. Integrin-fibronectin binding through RGD is required for timely *hairy1* and *meso1* expression and *dll1* downregulation.

(A) Schematic representation of the explant culture system. Posterior explants of HH11-14 chick embryos were bisected along the midline and cultured for 6 to 12 hours. One side of the explant was cultured with RGD, while the contralateral half was cultured in control medium. **(B)** Schematic representation of the action of RGD. RGD competes with the RGD-binding pockets of integrins, interfering with their binding to the ECM. **(C)** Number of somites (black bars) or somite-like condensations (dotted bars) formed in culture in control

and RGD-treated explants. *p<0.05. **(D)** Expression of *hairy1* (a, b), *meso1* (d, e) and *dll1* (g, h) in RGD-treated explants and contralateral controls at representative timepoints of culture. Straightened images of respective explant pairs (right) aligned by SIV. Rostral is on top. Percentage of RGD-treated explants with altered *hairy1*, *meso1* and *dll1* expression compared to the contralateral controls is shown in c, f and i, respectively. Impairing integrin-fibronectin binding with RGD alters *hairy1* and *meso1* expression relative to contralateral controls at 6 hours of culture onwards (a-f, arrowheads). *dll1* expression was altered at 9 hours of culture onwards (g-i, arrowheads). Scale bar: 500 μm.

Figure 5. Inhibition of integrin-fibronectin binding impairs somitic cleft formation.

(A-X) Longitudinal (left) and sagittal (right) views of explants cultured in control medium (A-F, M-R) and their contralateral RGD-treated halves (G-L, S-X) at S0 (A-L; region e in Figure 1 A) and SII (M-X; region c in Figure 1 A) levels, immunostained for N-cadherin (first column), fibronectin (fourth column) and stained for DNA (second and fifth columns). Third and sixth columns show the respective merged images. Explants were cultured for 6 hours. At the S0 level, control explants show normal apical accumulation of N-cadherin (A, arrowheads) and nuclear alignment (B, E, arrowheads) in peripheral somitic cells as well as fibronectin assembly in the nascent cleft (D, F, arrowheads). All of these are deficient in their RGD-treated contralateral halves (G-L, arrowheads). At the level of SII, N-cadherin accumulation and nuclear alignment occurs normally in both control and RGD-treated explants (M, N, S, T, arrowheads), but fibronectin assembly between adjacent somites (P, V, arrowheads) and cleft formation (Q, W, arrowheads) are deficient in RGD-treated explants compared to contralateral controls. Ncad – N-cadherin; FN – fibronectin. Scale bars: 50 μm.

<u>Impaired fibronectin assembly results in altered segmentation clock dynamics, *meso1* expression</u>

<u>and defects in somite morphogenesis</u>

To interrogate the importance of the mechanical properties of the extracellular fibronectin

matrix on somite formation, we perturbed fibronectin assembly in the PSM and somites by

electroporating primitive streak-stage embryos with a construct expressing the 70kDa fibronectin

fragment, a dominant-negative inhibitor of fibronectin assembly (Figure 6 A, B; McKeown-Longo

and Mosher, 1985; Sato et al., 2017). 70kDa-electroporated embryos exhibited a disrupted

fibronectin matrix, composed of thinner fibrils when compared with control pCAGGs-

electroporated embryos (Supplementary Figure 6 A; also see Figure 7 A, B), and displayed multiple

morphological defects, including kinked neural tube and detached tissues as well as perturbations

in somite morphogenesis, which are all reminiscent of phenotypes obtained in previous studies

interfering with fibronectin-integrin binding (Supplementary Figure 6 B, C; Drake et al., 1992;

Drake and Little, 1991; George et al., 1993; Girós et al., 2011; Takahashi et al., 2007). Although the

average number of somite-like structures formed in 70kDa-electroporated embryos was similar to

the number of somites in controls (Figure 6 C) they were ill-defined, often appearing fused or

crammed (Supplementary Figure 6 B e-f, C), and thus resemble the somite-like condensations

formed in RockOut- and RGD-treated explants (Figures 3 and 5).

Next, we sought to evaluate the impact of inhibiting fibronectin matrix assembly on the

molecular machinery underlying somite formation by using unilateral electroporation of the 70kDa

fragment (Figure 6 A) to allow the direct comparison of gene expression patterns within the same

embryo. We observed a significant increase in the frequency of perturbations in the expression of

embryonic clock genes *hairy1* ($p<0.01$) and *hairy2* ($p<0.05$), as well as in *meso1* ($p<0.01$) and *dll1*

($p<0.01$), when compared with embryos electroporated with pCAGGs alone (Figure 6 D).

Figure 6. Impairing fibronectin assembly by electroporation with a 70kDa expressing vector perturbs segmentation clock oscillations, timely *meso1* expression and *dll1* downregulation.

(**A**) Schematic representation of the electroporation strategy. PSM/ectoderm progenitors of primitive-streak stage embryos were electroporated with either a pCAGGs GFP-expressing vector (pCAGGs) alone, or co-electroporated with a 70kDa-expressing vector (70kDa) and were incubated for about 26 hours. (**B**) Schematic representation of the action of 70kDa. 70kDa disrupts the assembly of fibronectin matrix by competitively binding to the N-terminal self-assembly domains of the protein, impairing fibronectin fibril formation. (**C**) Number of somites (black bars) or somite-like condensations (dotted bars) formed in pCAGGs- and 70kDa-electroporated embryos after 26 hours. (**D**) Examples of the expression of *hairy1* (a-b), *hairy2* (d-e), *meso1* (g-h) and *dll1* (j-l) in pCAGGs- (top row) and 70kDa-electroporated embryos (middle row). Electroporated side is on right (a, b, g) or left (d, e, h, j, k) Perturbing the assembly of fibronectin on one side of the PSM leads to an asymmetric pattern of *hairy1* (b, arrowheads), *hairy2* (e, arrowheads), *meso1* (h, arrowheads) and *dll1* expression (k, arrowheads) in a higher percentage of embryos than in controls. Percentage of pCAGGs- and 70kDa-electroporated embryos with asymmetric expression of *hairy1*, *hairy2*, *meso1* and *dll1* between the electroporated PSM and the contralateral non-electroporated control PSM is shown in c, f, i and l, respectively. The number of embryos with an asymmetric pattern was significantly higher in 70kDa-electroporated embryos for all four genes studied. * p<0.05, *** p<0.01. Rostral is on top. Scale bar: 200 μm.

Consistent with these data, 70kDa-electroporated embryos have deficiencies in somite

morphogenesis which, in severe cases, leads to incomplete somitic clefts (Figure 7 A, B k, w,

arrows). Peripheral cells of nascent somites of 70kDa-electroporated embryos did, however,

accumulate ZO-1 apically (Figure 7 B e, h, i, l) which was maintained as the somites matured (Figure

7 B q, u, t, x). Nevertheless, these somites were abnormal in shape and appeared smaller in

severely affected embryos (Figure 7 B, i-l, u, x). In fact, the SI of the electroporated sides of 70kDa-

treated embryos were significantly smaller in width than those of the contralateral control, while

SV was significantly shorter in length (n=151; Supplementary Figure 7). In addition to defects in

somite morphology, the ectoderm and endoderm were separated from the paraxial mesoderm in

70kDa-electroporated embryos, indicating that their fibronectin matrix was insufficient to hold

these tissues together (brackets in Figure 7 B g, k, s, w).

We conclude that unperturbed fibronectin assembly in the PSM is required for tuning clock

gene expression dynamics, positioning of *meso1* expression and timely *dll1* downregulation in the

rostral-most PSM, as well as for the complete separation and morphogenesis of somites.

Figure 7. Somite morphology of 70kDa-electroporated embryos is severely compromised.

(A) Sagittal sections of embryos electroporated with pCAGGs and 70kDa, the latter with a severe phenotype.

Arrows point to deficient somitic clefts. Rostral to the left and dorsal on top. **(B)** Sagittal views of embryos

electroporated with pCAGGs (a-d, m-p) and 70kDa (e-l, q-x) with either mild (e-h, q-t) or severe (i-l, u-x)

phenotypes, at S0 (a-l) and SII (m-x) levels, immunostained for ZO-1 (first column) and fibronectin (second

column) and stained for DNA (third column). Fourth column shows the merge of the respective channels. S0

of pCAGGs- and 70kDa-electroporated embryos all polarize ZO-1 normally (a, e, i, arrowheads) and this is

maintained at SII levels (m, q, u), but the fibronectin matrix surrounding the somites of 70kDa-treated embryos is disrupted compared to pCAGGs-electroporated embryos (b, f, j, v, arrows). Somites of 70kDa-electroporated embryos are also severely detached from both the ectoderm and endoderm compared to embryos electroporated only with pCAGGS (third column, brackets), and the fibronectin matrix connecting the endoderm to the somites is severely compromised (second column, empty arrowheads). Somites of embryos electroporated with 70kDa with more severe defects also fail to fully detach from adjacent somites (k, w, arrows). Rostral to the left and dorsal to the top. Dashed lines indicate altered somite morphology. FN – fibronectin. ZO-1 – *Zonula occludens* protein 1. Scale bars: 50 μm.

Discussion:

Fibronectin matrix and its downstream mechanotransduction pathway coordinate segmentation clock dynamics and segment boundary formation

We have identified a mechanotransduction pathway involving a fibronectin – integrin – ROCK – NM II axis, which regulates the dynamics of segmentation clock gene expression in the chick embryo. Four independent treatments interfering with different levels of this pathway (from intracellular actomyosin contractility to fibronectin matrix assembly; Figure 8 A), all lead to altered segmentation clock dynamics, misregulated *meso1* expression, impaired or deficient cleft formation and alterations in somite morphology (Figure 8 B).

The similarity of the phenotypes obtained in experiments blocking ROCK I/II and impairing normal cell-fibronectin interactions with RGD or 70kDa (Figure 8 B) suggest that the fibronectin matrix is transducing mechanical signals via ROCK and α5β1 integrin, the major fibronectin receptor in the PSM (Jülich et al., 2005; Rifes et al., 2007; Yang et al., 1999), and that these signals affect segmentation clock oscillations. Indeed, the fibronectin – α5β1 integrin – ROCK – NM II axis is believed to play key roles in cellular force generation in a variety of cell types (Bharadwaj et al., 2017; Daley et al., 2011; Schiller et al., 2013; Torr et al., 2015). Our data show that blocking different elements of this axis leads to asymmetric patterns of *hairy1* expression as well as incorrect positioning of *meso1* expression on the experimental versus the control sides of the same embryo. Moreover, they suggest that the mechanical environment is regulating Notch signaling in

the PSM. This is in agreement with a previous report, evidencing that β1 integrins are needed for

correct Notch signaling in neural progenitors (Campos et al., 2006). Combined roles of integrin

α5β1 and Notch are also required for zebrafish somitogenesis (Jülich et al., 2005) and chicken

embryos electroporated with RNAi constructs against integrin β1 show alterations in *hairy2, lfng*

and *meso1* expression in the PSM (Rallis et al., 2010). Finally, mouse embryos expressing

fibronectin where the RGD sequence was substituted with RGE (*Fn1^{RGE/RGE}* embryos) showed

asymmetric and/or dampened expression of *Lnfg* and *Hes7* in the PSM (Girós et al., 2011) and

EphA4, a direct target of Mesp2 in the anterior PSM (Nakajima et al., 2006), was diffusely

expressed or absent (Girós et al., 2011).

Exactly how the mechanical environment regulates segmentation clock oscillations is

unknown. However, it is becoming increasingly clear that mechanics play a crucial role in Notch

activation (Gordon et al., 2015; Luca et al., 2017; Meloty-Kapella et al., 2012) and it was recently

demonstrated that sustained Notch signaling in the *Drosophila* notum requires actomyosin

contractility in both signal sending and receiving cells (Hunter et al., 2019). It is thus possible that

cells in the PSM become detached from their neighbors when their mechanical environment or

intracellular contractility is impaired. This may place Notch receptors and ligands of adjacent PSM

cells further apart, leading to a reduction in the incidence or duration of Notch signaling events.

Altogether, our results strongly suggest the existence of a link between the fibronectin –

α5β1 integrin – ROCK – NM II pathway and Notch signaling in the PSM. When perturbed, this leads

to a dysregulation of segmentation clock oscillations and the mispositioning of the segmental

border.

<u>Somite cleft formation and cell epithelialization have different mechanical requirements</u>

In the rostral PSM, Mesp2/Meso1 activates the expression of EphA4, which interacts with

EphrinB2 in cells rostral to the presumptive cleft, causing cell-cell repulsion and the formation of

an incipient cleft (Nakajima et al., 2006; Watanabe et al., 2009). Then, fibronectin matrix assembly

stabilizes it (Jülich et al., 2015; Rifes and Thorsteinsdóttir, 2012) and promotes the epithelialization

of cells rostral to the cleft (Martins et al., 2009). This can be defined as the first step of

morphological somite individualization. The second step is defined as the complete

epithelialization of the remaining cells of the nascent somite and lasts until SII, when all somitic

cells have acquired a spindle-like shape and are organized into a rosette (Martins et al., 2009).

RockOut-, RGD- and 70kDa-treated embryos (Figure 8 A, B) all show perturbations in

segmentation clock gene expression, abnormal positioning of *meso1* and defects in somite

boundary development. Although somite morphology is also perturbed, the acquisition of the

spindle-shape cell morphology which occurs as S0 develops into SII does not appear to be

significantly perturbed. Thus, the first step of morphological somite formation is affected but the

second step is not. In contrast, Blebbistatin-treated explants not only have the defects listed

above, but cells that had already upregulated *meso1* before the addition of the drug and formed

an incipient cleft during culture, were completely unable to epithelialize. In fact, Blebbistatin-

treated explants formed only 1 or 2 somites, and their cells did not acquire the elongated, spindle-

shape typical of SI and SII somites. Hence, Blebbistatin affects both steps of morphological somite

formation.

RockOut targets NM II activity indirectly by inhibiting ROCK I and II, two of the kinases that

activate NM II (Newell-Litwa et al., 2015). In contrast, Blebbistatin directly targets the NM II

ATPase. Our results thus raise the possibility that the acquisition of the spindle-shaped

morphology of cells is dependent on another NM II activator. Interestingly, Ca^{++}/calmodulin

signaling can activate NM II and inhibiting calmodulin was shown to block the acquisition of this

morphology during chick somitogenesis (Chernoff and Hilfer, 1982).

<u>Fibronectin matrix complexity gradient as a player in the PSM wavefront of maturation</u>

While the waves of Notch oscillatory activity travel through the entire length of the PSM,

they are only translated into segments in the anterior-most region of the tissue. Opposing

gradients of Fgf/Wnt and Retinoic acid (RA) in the PSM are thought to define the wavefront, which

marks the region where PSM cells become competent for somite formation (Hubaud and

Pourquié, 2014). Rostral to the wavefront, segmentation clock oscillations progressively slow

down until they reach a halt and cells become part of a somite (Palmeirim et al., 1997).

Herein, we propose a model (Figure 8 C) where the posterior to anterior gradient of

fibronectin matrix complexity is interpreted by the PSM cells as an increasing gradient of

mechanical tension (Figure 8 C). We propose that the tensional state created by the fibronectin

matrix surrounding the PSM at the level of the wavefront acts as a threshold which activates a

mechanotransduction signaling cascade ensuring the correct spatio-temporal conversion of the

cyclic expression of segmentation clock genes into periodic *meso1* expression which defines the

next somitic cleft (Figure 8 C). In support of this hypothesis, a gradient of increasing paraxial

mesoderm stiffness from the tail to rostral somites has recently been identified in the chick

embryo (Marrese et al., 2019). The PSM thus integrates a combination of chemical and mechanical

signals, namely gradients of Fgf/Wnt and RA (Aulehla and Pourquié, 2010) and, simultaneously, an

increase in fibronectin matrix complexity and stiffness (Figure 8 C). The fibronectin matrix would

thus be a key contributor to the PSM wavefront of maturation in that it regulates where the next

somitic cleft is positioned while simultaneously supporting a cohort of competent cells as they bud

off from the anterior PSM as somites (Figure 8 C). Hence, we propose that the

mechanotransduction pathway downstream of fibronectin plays a major role in the translation of

cyclic waves of expression of segmentation clock genes into the periodic morphogenesis of

somites.

Materials and Methods:

Embryos

Fertile eggs were obtained from commercial sources (Sociedade Agrícola Quinta da Freiria or Pintobar Exploração Avícola, Lda, Portugal) and incubated at 37.5ºC in a humidified chamber until the desired HH stage (Hamburger and Hamilton, 1992). Somite nomenclature is according to Pourquié and Tam (2001).

Embryo explant culture and chemical treatments

Explant tissues of HH11-14 embryos were collected and cultured as previously described (Palmeirim et al., 1997; Rifes et al., 2007). Embryos were bisected along the midline and then cut transversally rostral to somites IV and Hensen's node. The two contralateral halves thus retained half of the neural tube and notochord as well as the first four somites and the PSM, with all remaining neighboring tissues intact. Explants were placed on top of a polycarbonate filter floating on M199 medium supplemented with 10% chick serum, 5% fetal calf serum and 100 U/ml of penicillin and streptomycin (Palmeirim et al., 1997). Explants were then cultured at 37ºC with 5% CO_2 from 6 to 12 hours.

InSolution™Blebbistatin (Calbiochem) and RockOut (Calbiochem) diluted in DMSO were used at a final concentration of 50 µM in culture medium. Equal volumes of DMSO (Sigma-Aldrich) were used as control for both drugs. The RGD peptide (Sigma) was diluted in culture medium and used at 0.9 mM, while control explants were cultured in medium only. RGD peptide efficiency was confirmed in a cell adhesion assay (Danen et al., 2002; Pierschbacher and Ruoslahti, 1984) before using it on explants.

Embryo electroporation and *ex ovo* culture

HH4-5 embryos were electroporated on one (randomly selected) side of the primitive streak in the presumptive PSM and/or ectoderm and cultured *ex ovo* using the Early Chick culture

method (Chapman et al., 2001). The electroporation mixture contained plasmid DNA at 0.5-1 µg/µl

mixed with 0.4% Fast Green for visualization. Embryos were submerged in an electroporation

chamber filled with Tyrode's saline and three pulses of 6-9 V, 50 ms each, at 350 ms intervals were

applied. Control embryos were electroporated with pCAGGs containing a GFP reporter (pCAGGs-

GFP; abbreviated pCAGGs). pCAGGs-70kDa qFN1 was kindly provided by Yuki Sato (Sato et al.,

2017) and was co-electroporated with the pCAGGs-GFP plasmid in experimental embryos

(treatment abbreviated 70kDa). Electroporated embryos were screened for GFP after fixation to

select embryos with an intense signal on only one side to process for whole mount morphological

analysis, *in situ* hybridization experiments and transverse sectioning. For morphological analysis in

sagittal sections, the embryo side electroporated with pCAAGGs was compared to the 70kDa-

electroporated side of other same stage embryos.

<u>Cryosectioning and immunohistochemistry</u>

Cryosectioning was performed on whole embryos and embryo explants fixed in 4%

paraformaldehyde in 0.12 M phosphate buffer containing 4% sucrose and processed for

cryoembedding as previously described. Briefly, fixed samples were embedded in 7.5% gelatin in

0.12 M phosphate buffer containing 15% sucrose, frozen on dry ice-chilled isopentane and stored

at -80ºC until sectioning. Cryostat sections (10-30 µm) were processed for immunofluorescence as

previously described (Gomes de Almeida et al., 2016). Permeabilization of sections was performed

with 0.2% Triton-X100 in phosphate buffered saline (PBS). 5% bovine serum albumen (BSA) or a

combination of 1% BSA and 10% Normal Goat Serum (NGS) in PBS were used for blocking

depending on the presence or absence of anti-fibronectin antibodies, respectively. Primary and

secondary antibodies were diluted in 1% BSA in PBS. Sections were incubated with primary

antibodies overnight at 4ºC and with secondary antibodies for 1 hour at room temperature. For

whole-mount immunodetection, explants were fixed in 4% paraformaldehyde in PBS and

processed as previously described (Martins et al., 2009; Rifes and Thorsteinsdóttir, 2012). 1%

Triton-X100 in PBS was used for permeabilization and 1% BSA in PBS was used for blocking and

antibody dilution. Antibody incubation was performed overnight at 4ºC.

The following primary antibodies were used: anti-ZO-1 (Zymed, 1:100 or Invitrogen,

1:100); anti-N-cadherin (BD Biosciences, 1:100); anti-fibronectin (Sigma, 1:400), anti-activated

caspase3 (Cell Signaling, 1:1000) and anti-GFP (Invitrogen, 1:100). For F-actin staining we used

Alexa 488-conjugated phalloidin (Invitrogen, 1:40) and for staining DNA we used ToPro3

(Invitrogen, 1:500) in conjunction with ribonuclease A (Sigma, 10 µg/ml), 4% Methyl Green (Sigma,

diluted 1:250; Prieto et al., 2015) or 4',6-diamidino-2-phenylindole (DAPI, 5µg/ml in PBS with 0.1%

Triton-X100). For detection of the primary antibodies the adequate secondary goat anti-mouse

and anti-rabbit Alexa 488-, Alexa 568- or Alexa 546-conjugated F'ab fragments from Invitrogen

were used (1:1000).

In situ hybridization

In situ hybridization using DIG-labeled RNA probes was performed as described previously

(Henrique et al., 1995) with minor alterations (Gomes de Almeida et al., 2016). RNA probes were

synthetized from linearized plasmids: *dll1* (Henrique et al., 1995), *meso1* (Buchberger et al., 1998),

hairy1 (Palmeirim et al., 1997) and *hairy2* (Jouve et al., 2000).

Statistical analysis

Paired Student's t-tests were performed to assess for differences in the number of somites

formed in Blebbistatin-, RockOut- and RGD-treated explants relative to the respective controls,

and in embryos electroporated with pCAGGs only and pCAGGs + 70kDa. Differences in the

frequency of morphological and gene expression phenotypes found in 70kDa-electroporated

embryos compared to pCAGGs-electroporated control embryos was tested through a Chi-square

test. Differences in somite size between pCAGGs- or 70kDa-electroporated sides compared to the

control (non-electroporated) sides of embryos was tested through a nested ANOVA. The side in

which the embryo was electroporated (left or right) was nested in the treatment (non-

electroporated, pCAGGs-electroporated or 70kDa-electroporated) to account for a potential

variability between the two sides. Statistical significance was set at $p<0.05$. Statistical analyses

were performed in Statistica 10, Graphpad Prism 5 and RStudio.

<u>Sample preparation and imaging</u>

Whole mount explants were gradually dehydrated in methanol and cleared in

methylsalicylate (Sigma-Aldrich) as described previously (Martins et al., 2009; Rifes and

Thorsteinsdóttir, 2012), except for phalloidin-labelled embryos and explants, where a shorter

series of ethanol dehydration series was used. Cryostat sections were mounted in Vectashield

(Vector Laboratories) or in 5mg/ml propyl gallate in glycerol/PBS (9:1) with 0.01% azide.

Immunofluorescence images were taken on a confocal Leica SPE microscope, following imaging

acquisition steps described previously (Rifes and Thorsteinsdóttir, 2012). Imaging of

electroporated embryos and explants processed for *in situ* hybridization was performed using a

Zeiss LUMAR V12 Stereoscope coupled to a Zeiss Axiocam 503 color 3MP camera. Image analysis

was performed using Fiji v. 1.49 and Amira V.5.3.3 (Visage Imaging Inc.) software. Image histogram

corrections were performed in Fiji and exported as TIFF files. When applicable, contiguous images

were stitched together into a single image using the pairwise stitching Fiji plugin (Preibisch et al.,

2009). For the analysis of *in situ* hybridization patterns along the PSM explants, the Fiji plugin

Straighten (Kocsis et al., 1991) was used.

Acknowledgements:

We thank Dr. Yuki Sato for generously sharing the pCAGGs-q70kDa construct and Inês Fragata for

help with the statistical analysis. This work was supported by Fundação para a Ciência e a

Tecnologia (FCT, Portugal) projects PTDC/SAU-OBD/103771/2008, PTDC/BEXBID/5410/2014,

UID/BIA/00329/2013, UID/BIM/04773/2019 CBMR, and FCT scholarships SFRH/BD/86980/2012

(PGA) and SFRH/BD/37423/2007 (PR). Imaging and image analysis were done in the Microscopy

Facility at the Faculty of Sciences of the University of Lisbon and the Light Microscopy Unit of

CBMR-UAlg, nodes of the Portuguese Platform for BioImage (reference PPBI-POCI-01-0145-FEDER-

022122). Finally, we are grateful to all members of our laboratories for helpful discussions.

Competing interests:

The authors declare no competing interests.

References:

Aulehla A, Pourquié O. 2010. Signaling gradients during paraxial mesoderm development. *Cold*

Spring Harb Perspect Biol **2**:1–17. doi:10.1101/cshperspect.a000869

Bailey C, Dale K. 2015. Somitogenesis in vertebrate developmentELS. Chichester, UK: John Wiley

& Sons, Ltd. pp. 1–15. doi:10.1002/9780470015902.a0003820.pub2

Barriga EH, Franze K, Charras G, Mayor R. 2018. Tissue stiffening coordinates morphogenesis by

triggering collective cell migration in vivo. *Nature* **554**:523–527. doi:10.1038/nature25742

Barrios A, Poole RJ, Durbin L, Brennan C, Holder N, Wilson SW. 2003. Eph/Ephrin signaling

regulates the mesenchymal-to-epithelial transition of the paraxial mesoderm during somite

morphogenesis. *Curr Biol* **13**:1571–82. doi:10.1016/j.cub.2003.08.030

Bharadwaj M, Strohmeyer N, Colo GP, Helenius J, Beerenwinkel N, Schiller HB, Fässler R, Müller

DJ. 2017. αV-class integrins exert dual roles on α5β1 integrins to strengthen adhesion to

fibronectin. *Nat Commun* **8**:14348. doi:10.1038/ncomms14348

Buchberger A, Seidl K, Klein C, Eberhardt H, Arnold HH. 1998. cMeso-1, a novel bHLH transcription

factor, is involved in somite formation in chicken embryos. *Dev Biol* **199**:201–15.

doi:10.1006/dbio.1998.8919

Campbell ID, Humphries MJ. 2011. Integrin structure, activation, and interactions. *Cold Spring*

Harb Perspect Biol **3**:a004994–a004994. doi:10.1101/cshperspect.a004994

Campos LS, Decker L, Taylor V, Skarnes W. 2006. Notch, epidermal growth factor receptor, and

beta1-integrin pathways are coordinated in neural stem cells. *J Biol Chem* **281**:5300–9.

doi:10.1074/jbc.M511886200

Chan CJ, Heisenberg C-P, Hiiragi T. 2017. Coordination of morphogenesis and cell-fate specification

in development. *Curr Biol* **27**:R1024–R1035. doi:10.1016/j.cub.2017.07.010

Chapman SC, Collignon J, Schoenwolf GC, Lumsden A. 2001. Improved method for chick whole-

embryo culture using a filter paper carrier. *Dev Dyn* **220**:284–9. doi:10.1002/1097-

0177(20010301)220:3<284::AID-DVDY1102>3.0.CO;2-5

Chernoff EA, Hilfer SR. 1982. Calcium dependence and contraction in somite formation. *Tissue Cell*

14:435–49. doi:10.1016/0040-8166(82)90038-6

Christ B, Huang R, Scaal M. 2007. Amniote somite derivatives. *Dev Dyn* **236**:2382–2396.

doi:10.1002/dvdy.21189

Cosgrove BD, Mui KL, Driscoll TP, Caliari SR, Mehta KD, Assoian RK, Burdick JA, Mauck RL. 2016. N-

cadherin adhesive interactions modulate matrix mechanosensing and fate commitment of

mesenchymal stem cells. *Nat Mater* **15**:1297–1306. doi:10.1038/nmat4725

Daley WP, Kohn JM, Larsen M. 2011. A focal adhesion protein-based mechanochemical checkpoint

regulates cleft progression during branching morphogenesis. *Dev Dyn* **240**:2069–2083.

doi:10.1002/dvdy.22714

Danen EHJ, Sonneveld P, Brakebusch C, Fässler R, Sonnenberg A. 2002. The fibronectin-binding

integrins α5β1 and αvβ3 differentially modulate RhoA-GTP loading, organization of cell

matrix adhesions, and fibronectin fibrillogenesis. *J Cell Biol* **159**:1071–1086.

doi:10.1083/jcb.200205014

Dequéant M-L, Glynn E, Gaudenz K, Wahl M, Chen J, Mushegian A, Pourquié O. 2006. A complex

oscillating network of signaling genes underlies the mouse segmentation clock. *Science*

314:1595–8. doi:10.1126/science.1133141

Drake CJ, Davis LA, Hungerford JE, Little CD. 1992. Perturbation of beta 1 integrin-mediated

adhesions results in altered somite cell shape and behavior. *Dev Biol* **149**:327–38.

doi:10.1016/0012-1606(92)90288-R

Drake CJ, Little CD. 1991. Integrins play an essential role in somite adhesion to the embryonic axis.

Dev Biol **143**:418–21. doi:10.1016/0012-1606(91)90092-H

George EL, Georges-Labouesse EN, Patel-King RS, Rayburn H, Hynes RO. 1993. Defects in

mesoderm, neural tube and vascular development in mouse embryos lacking fibronectin.

Development **119**:1079–1091. PMID: 8306876.

Georges-Labouesse EN, George EL, Rayburn H, Hynes RO. 1996. Mesodermal development in

mouse embryos mutant for fibronectin. *Dev Dyn* **207**:145–56. doi:10.1002/(SICI)1097-

0177(199610)207:2<145::AID-AJA3>3.0.CO;2-H

Girós A, Grgur K, Gossler A, Costell M. 2011. α5β1 integrin-mediated adhesion to fibronectin is

required for axis elongation and somitogenesis in mice. *PLoS One* **6**:e22002.

doi:10.1371/journal.pone.0022002

Goh KL, Yang JT, Hynes RO. 1997. Mesodermal defects and cranial neural crest apoptosis in alpha5

integrin-null embryos. *Development* **124**:4309–4319. PMID: 9334279.

Gomes de Almeida P, Pinheiro GG, Nunes AM, Gonçalves AB, Thorsteinsdóttir S. 2016. Fibronectin

assembly during early embryo development: A versatile communication system between

cells and tissues. *Dev Dyn* **245**:520–35. doi:10.1002/dvdy.24391

Gordon WR, Zimmerman B, He L, Miles LJ, Huang J, Tiyanont K, McArthur DG, Aster JC, Perrimon

N, Loparo JJ, Blacklow SC. 2015. Mechanical Allostery: Evidence for a Force Requirement in

the Proteolytic Activation of Notch. *Dev Cell* **33**:729–36. doi:10.1016/j.devcel.2015.05.004

Hamburger V, Hamilton HL. 1992. A series of normal stages in the development of the chick

embryo. *Dev Dyn* **195**:231–272. doi:10.1002/aja.1001950404

Henrique D, Adam J, Myat A, Chitnis A, Lewis J, Ish-Horowicz D. 1995. Expression of a Delta

homologue in prospective neurons in the chick. *Nature* **375**:787–790. doi:10.1038/375787a0

Hubaud A, Pourquié O. 2014. Signalling dynamics in vertebrate segmentation. *Nat Rev Mol Cell*

Biol **15**:709–721. doi:10.1038/nrm3891

Hubaud A, Regev I, Mahadevan L, Pourquié O. 2017. Excitable dynamics and Yap-dependent

mechanical cues drive the segmentation clock. *Cell* **171**:668–682.e11.

doi:10.1016/j.cell.2017.08.043

Hunter GL, He L, Perrimon N, Charras G, Giniger E, Baum B. 2019. A role for actomyosin contractility

in Notch signaling. *BMC Biol* **17**:12. doi:10.1186/s12915-019-0625-9

Jouve C, Palmeirim I, Henrique D, Beckers J, Gossler A, Ish-Horowicz D, Pourquié O. 2000. Notch

signalling is required for cyclic expression of the hairy-like gene HES1 in the presomitic

mesoderm. *Development* **127**:1421–1429. doi:10.1016/S0092-8674(00)80451-1

Jülich D, Cobb G, Melo AM, McMillen P, Lawton AK, Mochrie SGJ, Rhoades E, Holley SA. 2015.

Cross-scale integrin regulation organizes ECM and tissue topology. *Dev Cell* **34**:33–44.

doi:10.1016/j.devcel.2015.05.005

Jülich D, Geisler R, Holley SA, Tübingen 2000 Screen Consortium. 2005. Integrinalpha5 and

delta/notch signaling have complementary spatiotemporal requirements during zebrafish

somitogenesis. *Dev Cell* **8**:575–86. doi:10.1016/j.devcel.2005.01.016

Kocsis E, Trus BL, Steer CJ, Bisher ME, Steven AC. 1991. Image averaging of flexible fibrous

macromolecules: the clathrin triskelion has an elastic proximal segment. *J Struct Biol* **107**:6–

14. doi:10.1093/bioinformatics/btp184

Koshida S, Kishimoto Y, Ustumi H, Shimizu T, Furutani-Seiki M, Kondoh H, Takada S. 2005. Integrin

alpha5-dependent fibronectin accumulation for maintenance of somite boundaries in

zebrafish embryos. *Dev Cell* **8**:587–98. doi:10.1016/j.devcel.2005.03.006

Kragtorp KA, Miller JR. 2007. Integrin alpha5 is required for somite rotation and boundary

formation in Xenopus. *Dev Dyn* **236**:2713–20. doi:10.1002/dvdy.21280

Lauschke VM, Tsiairis CD, François P, Aulehla A. 2013. Scaling of embryonic patterning based on

phase-gradient encoding. *Nature* **493**:101–105. doi:10.1038/nature11804

Luca VC, Kim BC, Ge C, Kakuda S, Wu D, Roein-Peikar M, Haltiwanger RS, Zhu C, Ha T, Garcia KC.

2017. Notch-Jagged complex structure implicates a catch bond in tuning ligand sensitivity.

Science (80-) **355**:1320–1324. doi:10.1126/science.aaf9739

Marek M, Kubícek M. 1981. Morphogen pattern formation and development in growth. *Bull Math*

Biol **43**:259–70.

Marrese M, Antonovaite N, Nelemans BKA, Smit TH, Iannuzzi D. 2019. Micro-indentation and

optical coherence tomography for the mechanical characterization of embryos: Experimental

setup   and   measurements   on   chicken   embryos.   *Acta   Biomater*.

doi:10.1016/j.actbio.2019.07.056

Martins GG, Rifes P, Amândio R, Rodrigues G, Palmeirim I, Thorsteinsdóttir S. 2009. Dynamic 3D

cell rearrangements guided by a fibronectin matrix underlie somitogenesis. *PLoS One*

4:e7429. doi:10.1371/journal.pone.0007429

Masamizu Y, Ohtsuka T, Takashima Y, Nagahara H, Takenaka Y, Yoshikawa K, Okamura H,

Kageyama R. 2006. Real-time imaging of the somite segmentation clock: Revelation of

unstable oscillators in the individual presomitic mesoderm cells. *Proc Natl Acad Sci*

103:1313–1318. doi:10.1073/pnas.0508658103

McKeown-Longo PJ, Mosher DF. 1985. Interaction of the 70,000-mol-wt amino-terminal fragment

of fibronectin with the matrix-assembly receptor of fibroblasts. *J Cell Biol* **100**:364–74.

doi:10.1083/jcb.100.2.364

Meloty-Kapella L, Shergill B, Kuon J, Botvinick E, Weinmaster G. 2012. Notch Ligand Endocytosis

Generates Mechanical Pulling Force Dependent on Dynamin, Epsins, and Actin. *Dev Cell*

22:1299–1312. doi:10.1016/j.devcel.2012.04.005

Merle T, Farge E. 2018. Trans-scale mechanotransductive cascade of biochemical and

biomechanical patterning in embryonic development: the light side of the force. *Curr Opin*

Cell Biol **55**:111–118. doi:10.1016/j.ceb.2018.07.003

Morimoto M, Takahashi Y, Endo M, Saga Y. 2005. The Mesp2 transcription factor establishes

segmental borders by suppressing Notch activity. *Nature* **435**:354–359.

doi:10.1038/nature03591

Nakajima Y, Morimoto M, Takahashi Y, Koseki H, Saga Y. 2006. Identification of Epha4 enhancer

required for segmental expression and the regulation by Mesp2. *Development* **133**:2517–25.

doi:10.1242/dev.02422

Newell-Litwa KA, Horwitz R, Lamers ML. 2015. Non-muscle myosin II in disease: mechanisms and

therapeutic opportunities. *Dis Model Mech* **8**:1495–515. doi:10.1242/dmm.022103

Niwa Y, Shimojo H, Isomura A, Gonzalez A, Miyachi H, Kageyama R. 2011. Different types of

oscillations in Notch and Fgf signaling regulate the spatiotemporal periodicity of

somitogenesis. *Genes Dev* **25**:1115–1120. doi:10.1101/gad.2035311

Palmeirim I, Dubrulle J, Henrique D, Ish-Horowicz D, Pourquié O. 1998. Uncoupling segmentation

and somitogenesis in the chick presomitic mesoderm. *Dev Genet* **23**:77–85.

doi:10.1002/(SICI)1520-6408(1998)23:1<77::AID-DVG8>3.0.CO;2-3

Palmeirim I, Henrique D, Ish-Horowicz D, Pourquié O. 1997. Avian hairy gene expression identifies

a molecular clock linked to vertebrate segmentation and somitogenesis. *Cell* **91**:639–648.

doi:10.1016/S0092-8674(00)80451-1

Pierschbacher MD, Ruoslahti E. 1984. Cell attachment activity of fibronectin can be duplicated by

small synthetic fragments of the molecule. *Nature* **309**:30–3. PMID: 6325925.

Pourquié O, Tam PPL. 2001. A nomenclature for prospective somites and phases of cyclic gene

expression in the presomitic mesoderm. *Dev Cell* **1**:619–20. doi:10.1016/S1534-

5807(01)00082-X

Preibisch S, Saalfeld S, Tomancak P. 2009. Globally optimal stitching of tiled 3D microscopic image

acquisitions. *Bioinformatics* **25**:1463–1465. doi:10.1093/bioinformatics/btp184

Prieto D, Aparicio G, Machado M, Zolessi FR. 2015. Application of the DNA-specific stain methyl

green in the fluorescent labeling of embryos. *J Vis Exp* e52769. doi:10.3791/52769

Rallis C, Pinchin SM, Ish-Horowicz D. 2010. Cell-autonomous integrin control of Wnt and Notch

signalling during somitogenesis. *Development* **137**:3591–3601. doi:10.1242/dev.050070

Rifes P, Carvalho L, Lopes C, Andrade RP, Rodrigues G, Palmeirim I, Thorsteinsdóttir S. 2007.

Redefining the role of ectoderm in somitogenesis: a player in the formation of the fibronectin

matrix of presomitic mesoderm. *Development* **134**:3155–3165. doi:10.1242/dev.003665

Rifes P, Thorsteinsdóttir S. 2012. Extracellular matrix assembly and 3D organization during paraxial

mesoderm development in the chick embryo. *Dev Biol* **368**:370–381.

doi:10.1016/j.ydbio.2012.06.003

Ringer P, Colo G, Fässler R, Grashoff C. 2017. Sensing the mechano-chemical properties of the

extracellular matrix. *Matrix Biol* **64**:6–16. doi:10.1016/j.matbio.2017.03.004

Saga Y. 2012. The mechanism of somite formation in mice. *Curr Opin Genet Dev* **22**:331–338.

doi:10.1016/j.gde.2012.05.004

Saga Y, Takeda H. 2001. The making of the somite: molecular events in vertebrate segmentation.

Nat Rev Genet **2**:835–845. doi:10.1038/35098552

Sato Y, Nagatoshi K, Hamano A, Imamura Y, Huss D, Uchida S, Lansford R. 2017. Basal filopodia and

vascular mechanical stress organize fibronectin into pillars bridging the mesoderm-

endoderm gap. *Development* **144**:281–291. doi:10.1242/dev.141259

Sato Y, Yasuda K, Takahashi Y. 2002. Morphological boundary forms by a novel inductive event

mediated by Lunatic fringe and Notch during somitic segmentation. *Development* **129**:3633–

44. PMID: 12117813.

Schiller HB, Hermann M-R, Polleux J, Vignaud T, Zanivan S, Friedel CC, Sun Z, Raducanu A,

Gottschalk K-E, Théry M, Mann M, Fässler R. 2013. β1- and αv-class integrins cooperate to

regulate myosin II during rigidity sensing of fibronectin-based microenvironments. *Nat Cell*

Biol **15**:625–636. doi:10.1038/ncb2747

Shih NP, François P, Delaune EA, Amacher SL. 2015. Dynamics of the slowing segmentation clock

reveal alternating two-segment periodicity. *Development* **142**:1785–93.

doi:10.1242/dev.119057

Slack JMW. 1987. We have a morphogen! *Nature* **327**:553–554. doi:10.1038/327553a0

Smutny M, Ákos Z, Grigolon S, Shamipour S, Ruprecht V, Čapek D, Behrndt M, Papusheva E, Tada

M, Hof B, Vicsek T, Salbreux G, Heisenberg CP. 2017. Friction forces position the neural

anlage. *Nat Cell Biol* **19**:306–317. doi:10.1038/ncb3492

Straight AF, Cheung A, Limouze J, Chen I, Westwood NJ, Sellers JR, Mitchison TJ. 2003. Dissecting

temporal and spatial control of cytokinesis with a myosin II Inhibitor. *Science* **299**:1743–7.

doi:10.1126/science.1081412

Takahashi S, Leiss M, Moser M, Ohashi T, Kitao T, Heckmann D, Pfeifer A, Kessler H, Takagi J,

Erickson HP, Fässler R. 2007. The RGD motif in fibronectin is essential for development but

dispensable for fibril assembly. *J Cell Biol* **178**:167–178. doi:10.1083/jcb.200703021

Takahashi Y, Inoue T, Gossler A, Saga Y. 2003. Feedback loops comprising Dll1, Dll3 and Mesp2,

and differential involvement of Psen1 are essential for rostrocaudal patterning of somites.

Development **130**:4259–68. doi:10.1242/dev.00629

Takahashi Y, Koizumi K, Takagi A, Kitajima S, Inoue T, Koseki H, Saga Y. 2000. Mesp2 initiates somite

segmentation through the Notch signalling pathway. *Nat Genet* **25**:390–6.

doi:10.1038/78062

Tiedemann H. 1976. Pattern formation in early developmental stages of amphibian embryos. *J

Embryol Exp Morphol* **35**:437–44. PMID: 781172.

Torr EE, Ngam CR, Bernau K, Tomasini-Johansson B, Acton B, Sandbo N. 2015. Myofibroblasts

exhibit enhanced fibronectin assembly that is intrinsic to their contractile phenotype. *J Biol

Chem* **290**:6951–61. doi:10.1074/jbc.M114.606186

Watanabe T, Sato Y, Saito D, Tadokoro R, Takahashi Y. 2009. EphrinB2 coordinates the formation

of a morphological boundary and cell epithelialization during somite segmentation. *Proc Natl

Acad Sci* **106**:7467–7472. doi:10.1073/pnas.0902859106

Wolfenson H, Lavelin I, Geiger B. 2013. Dynamic regulation of the structure and functions of

integrin adhesions. *Dev Cell* **24**:447–58. doi:10.1016/j.devcel.2013.02.012

Wolfenson H, Meacci G, Liu S, Stachowiak MR, Iskratsch T, Ghassemi S, Roca-Cusachs P,

O'Shaughnessy B, Hone J, Sheetz MP. 2016. Tropomyosin controls sarcomere-like

contractions for rigidity sensing and suppressing growth on soft matrices. *Nat Cell Biol* **18**:33–

42. doi:10.1038/ncb3277

Yang JT, Bader BL, Kreidberg J a, Ullman-Culleré M, Trevithick JE, Hynes RO. 1999. Overlapping and

independent functions of fibronectin receptor integrins in early mesodermal development.

Dev Biol **215**:264–77. doi:10.1006/dbio.1999.9451

Yang JT, Rayburn H, Hynes RO. 1993. Embryonic mesodermal defects in alpha 5 integrin-deficient

mice. *Development* **119**:1093–105. PMID: 781172.

Yarrow JC, Totsukawa G, Charras GT, Mitchison TJ. 2005. Screening for cell migration inhibitors via

automated microscopy reveals a Rho-kinase inhibitor. *Chem Biol* **12**:385–95.

doi:10.1016/j.chembiol.2005.01.015

Supplementary Figures

809
810
811 **Supplementary Figure 1. Blocking NM II or ROCK I/II activity for 3 hours leads to altered *meso1* expression.**
812 **(A, B)** m*eso1* expression evaluated by *in situ* hybridization in explants treated for 3 hours with Blebbistatin
813 (**A**) or RockOut (**B**) and in the respective contralateral controls, shows that *meso1* expression is already
814 altered two clock cycles after adding the drugs. Straightened images of the respective explant pairs (right)
815 were aligned by SIV. Rostral is on top.
816

Supplementary Figure 2. Apoptosis is not enhanced in the experimental culture conditions used.
(A, H) Immunostaining for activated Caspase3 in transverse sections of control and contralateral explants treated with either Blebbistatin (**A, B**), RockOut (**C, D**), or RGD (**E, F**) for 6 hours, and in pCAGGs- and 70kDa-electroporated embryos (**G, H**). DNA (blue), activated Caspase3 (magenta) and GFP (green). Arrows indicate Caspase3-positive cells. Apoptosis levels are not increased by any of the treatments. Blebb – Blebbistatin; nt – neural tube; s – somite; i – intermediate mesoderm; lp – lateral plate mesoderm. Dorsal is on top. Scale bars: 50 µm.

Supplementary Figure 3. NM II inhibition impairs F-actin apical enrichment and nuclear alignment in
somites formed during culture.

(A, L) Sagittal sections of DMSO-treated explants (A-C, G-I) and contralateral Blebbistatin-treated explants
(D-F, J-L) stained for F-actin and DNA. Rostral PSM epithelialization of DMSO-treated explants occurs
normally, with F-actin apical enrichment (A) and nuclear alignment (B) in S0 and SI. At the same axial level
in the contralateral Blebbistatin-treated explant (D-F), F-actin staining is dispersed (D) and nuclei do not align
(E). SII and SIII of DMSO-treated explants are epithelial (G-I), composed of an outer cell layer with aligned
nuclei (H) and elongated cells with apically enriched F-actin (G). At the equivalent axial level in the
Blebbistatin-treated explant (J-L), the somitic segments were severely affected. There is only a slight nuclear
alignment at the prospective inter-somitic border (K, arrowheads) and F-actin aggregates into dispersed and
separate foci (J, L). Rostral to the left and dorsal on top. Blebb – Blebbistatin; Phall – phalloidin F-actin
staining. Scale bars: 50 μm.

Supplementary Figure 4. Epithelial structure of the neural tube and ectoderm are unaffected by NM II
inhibition.

(A, N) Control explants (A-D and I-K) showed strong apically located ZO-1 labeling (B, I, arrowheads) both in
the neural tube (A-D; transverse optical section, medial on top) and in the overlying surface ectoderm (I-K;
dorsal view of ectoderm). In the presence of Blebbistatin (E-H, L-N), ZO-1 labeling remained restricted to the
apical end of neural tube cells (F, arrowheads) and of surface ectoderm cells (L, arrowheads). Blebb –
Blebbistatin; Ncad – N-cadherin. ZO-1 – *Zonula occludens* protein 1. Arrowheads point to ZO-1 labeling.
Scale bars: 20 µm.

Supplementary Figure 5. ROCK I/II inhibition impairs apical polarization of N-cadherin in nascent somites.
(A, L) Longitudinal views of explants cultured for 6 hours in control (DMSO) medium (A-C, G-I) and their contralateral RockOut-treated halves (D-F, J-L) at SI (A-F) and SIII (G-L) levels, immunostained for N-cadherin (first column) and stained for DNA (second column). Third column shows the respective merge of all stainings. SI in control explants shows normal accumulation of N-cadherin (A, arrowhead) and nuclear alignment (B, arrowhead). In contrast, SI in contralateral RockOut-treated explants fail to form a clear cleft (D-F, arrows) leading to partially fused somites. N-cadherin is, however, partially polarized (D, arrowhead) and nuclei are aligned (E, arrowhead). At SIII level, both explants show normal N-cadherin polarization (G, J) and nuclear alignment (H, L). Rostral on the left and midline on top. Ncad - N-cadherin. Scale bars: 50 µm.

Supplementary Figure 6. Embryos electroporated with the 70kDa construct exhibit numerous morphological defects.

(A) Representative images of the morphology of embryos electroporated with either pCAGGs only (a) or 70kDa, the latter being a severe phenotype (e). Severe phenotypes included severely detached tissues (e, arrow) and a truncated A-P axis. Transverse sections of pCAGGs- (b-d) and 70kDa-electroporared embryos (f-h) immunostained for fibronectin (b-c, f-g), GFP (d, h) and stained for DNA (c-d, g-h). Electroporated side is on left (a-d) or right (e-h). Detachment of tissues is clearly visible in 70kDa-electroporated embryos (e), which is accompanied by a severe disruption in the fibronectin matrix (compare b-c with f-g, arrows). **(B)** Close up of embryos electroporated with 70kDa showing kinked neural tube (d, arrowheads), fused somites (e, arrowheads) and fewer somites on the electroporated side (f, arrowheads). Electroporated sides are on left. Ventral view and rostral on top. **(C)** Percentage of pCAGGS- (green bars) and 70kDa-electroporated (pink

bars) embryos with morphological defects, including detached (pCAGGs: 39/154, 70kDa: 97/144) and
severely detached tissues (pCAGGs: 6/154, 70kDa: 79/144), kinked neural tube (pCAGGs: 15/154, 70kDa:
34/144), truncated A-P axis (pCAGGs: 13/154, 70kDa: 21/144), abnormal somite morphology (pCAGGs:
18/154, 70kDa: 41/144), and fewer somites on the electroporated side compared to the control non-
electroporated side (pCAGGs: 0/154, 70kDa: 15/144). ***p<0.001. FN – fibronectin. Scale bars: (A, a, e; C,
a-c) 500 µm, (C, d-f) 200 µm, (A, b-d, f-h) 50 µm.

Supplementary Figure 7. Quantification of somite length and width in pCAGGs- and 70kDa-electroporated
embryos.
(A-D) Length of SI (A) and SV (B) and width of SI (C) and SV (D) from pCAGGS- and 70kDa-electroporated
embryos compared to the control non-electroporated side. The measurements were made on images from
whole mount embryos. Somites from pCAGGs-electroporated embryos did not show a significant difference
in either length or width between electroporated vs non-electroporated sides (n=151), but the width of SI
and length of SV of the electroporated side of 70kDa-treated embryos were significantly smaller than that
of the contralateral non-electroporated control side (n=143). Bars represent the standard error of the mean.
ns – not significant; *** p<0.001.