1

Song overlapping, ambient noise and territorial aggression in great tits

Çağlar Akçay[1,2]*; Y. Kağan Porsuk[1]; Alican Avşar[3]; Dilan Çabuk[1]; C. Can Bilgin[3]

1. Department of Psychology, Koç University, Istanbul, Turkey
2. Department of Biological Sciences, Virginia Tech, Blacksburg, VA, USA
3. Department of Biological Sciences, Middle East Technical University, Ankara, Turkey

*corresponding author: akcay.caglar@gmail.com

Çağlar Akçay

Department of Psychology

Koc University

Rumelifeneri Kampusu, Sariyer 34450

Istanbul, Turkey

Abstract

Animals often communicate with each other in noisy environments where interference from

the ambient noise and other signallers may reduce the effectiveness of signals. Signallers also

may evolve behaviours to interfere with signals of their opponents, e.g. by temporally

overlapping them with their own, such as the song overlapping behaviour that is seen in some

songbirds during aggressive interactions. Song overlapping has been proposed to be a signal

of aggressive intent although direct evidence for this hypothesis has been lacking. In the

present paper we examined the question of whether song overlapping is correlated with

aggressive behaviours in a population of great tits (*Parus major*) living in an urban-rural

gradient. We also examined whether aggressive behaviours are correlated with the ambient

noise levels. We found that overlapping was associated negatively while ambient noise levels

were associated positively with aggressive behaviours males displayed against a simulated

intruder. Ambient noise levels did not correlate with song rate, song duration or song

overlapping. These results suggest that song overlapping of male great tits is not a signal of

aggression and great tits in noisy urban habitats may display higher levels of aggression due

to either interference of noise in aggressive communication or another indirect effect of

noise.

Keywords: song overlapping, urbanization, noise, aggressive signalling, great tits

Introduction

The environment through which signals are sent and received is open to interference by either sources of noise or other signals traveling through the medium (Bee and Micheyl, 2008; Brumm, 2006; Gil and Brumm, 2014; Wiley, 2006). Signals should therefore evolve to reduce the level of interference from noise and other signals present in the environment (Endler, 1992; Endler and Basolo, 1998; Ey and Fischer, 2009). At the same time, signallers should evolve behaviours that would reduce the interference from noise or other signals, such as adjusting the timing of signalling to minimize overlapping with other signals (Brumm, 2006; Ficken et al., 1974; Gochfeld, 1978; Wasserman, 1977) or acoustic noise (Gil et al., 2014), adjusting the distance to the receivers (Halfwerk et al., 2012).

Generally speaking, signallers tend to space their signals temporally to minimize interference from other signalling individuals (Brumm, 2006; Ficken et al., 1974; Wasserman, 1977; Wilson et al., 2016). In some circumstances, however, actively overlapping the signal of another individual by starting to signal before the other signal has ended, may be a deliberate behaviour used as a signal itself. Indeed, overlapping has been proposed to be a signal of aggressive intent in songbirds (Dabelsteen et al., 1997; Mennill and Ratcliffe, 2004; Naguib and Mennill, 2010). Signals of aggressive intent (also called threat signals) are signals that are not costly to produce but carry information about the likelihood of escalation by the signaller (Caryl, 1979; Searcy et al., 2006). Under this hypothesis, overlapping and therefore interfering with the transmission of an opponent's song may carry information to the opponent that the overlapping bird is ready to escalate an aggressive interaction.

Although song overlapping has been assumed to be an aggressive signal by a large number of studies (Naguib and Mennill, 2010), a review by Searcy and Beecher (2009, 2011) questioned whether song overlapping was a signal at all. Searcy and Beecher (2009) argued that a signalling behaviour should satisfy three empirical criteria to be considered a signal of aggressive intent: 1) the behaviour should occur more often in agonistic interactions than other non-agonistic interactions (context criterion); 2) the behaviour should correlate with other aggressive behaviours or predict a subsequent escalation (predictive criterion) and 3) the behaviour should elicit a differential reaction (either stronger or weaker) in receivers (response criterion).

Most studies of song overlapping have focused on the perspective of receivers, asking
whether birds respond differentially when their songs being overlapped experimentally.
These studies generally find changes in singing behaviours such as song rate and duration
that are consistent with the singers attempting to avoid the interference due to being
overlapped. Changes in aggressive behaviours such as attack or approach due to being
overlapped however, are less consistent across studies (Naguib and Mennill, 2010; Searcy
and Beecher, 2009).

Very few studies addressed the question whether song overlapping is correlated with
aggressive behaviours or subsequent escalation, i.e. the predictive criterion (Baker et al.,
2012; Fitzsimmons et al., 2008; van Dongen, 2006; Vehrencamp et al., 2007). Only one of
these studies found potential support for overlapping being an aggressive signal: golden
whistler males (*Pachycephala pectoralis*) were closer to the speaker when they overlapped
the speaker than when they did not (van Dongen, 2006). Two other studies in black-capped
chickadees (*Poecile atricapillus*) however, found no correlation between overlapping and
concurrent aggressive behaviours or subsequent attack (Baker et al., 2012; Fitzsimmons et
al., 2008). Finally a study in banded wrens (*Thryophilus pleurostictus*) found a negative
correlation such that overlapping was correlated with lower levels of aggression
(Vehrencamp et al., 2007).

In the present study we aim to assess the predictive criterion for song overlapping by
examining the relationship between song overlapping and aggressive behaviours in great tits
(*Parus major*). Although song overlapping has long been assumed to be an aggressive signal
in great tits (Dabelsteen et al., 1996; Langemann et al., 2000; Peake et al., 2001), no previous
study has asked whether this behaviour is correlated with the overlapping individual's
aggressive behaviours in this species.

A second aim of our study is to ask whether interference due to ambient noise also
has an effect on aggressive behaviours. Although noise is a feature of natural habitats as well,
the impact of noise in animal communication has garnered special attention due to the
increasing levels of anthropogenic noise that affects social behaviour of wildlife (Gil and
Brumm, 2014; Johnson and Munshi-South, 2017; Shannon et al., 2016). Anthropogenic noise
may make signals less detectable (Kleist et al., 2016; Templeton et al., 2016) or less effective
(Halfwerk et al., 2011). In response, animals may change the frequency, amplitude or length
characteristics of their signals (Brumm and Todt, 2002; Slabbekoorn and den Boer-Visser,

2006; Wood and Yezerinac, 2006) or change their signalling or social behaviours (Halfwerk
et al., 2012; Halfwerk and Slabbekoorn, 2009).

If noise interferes with effective signalling during aggressive interactions than we
may expect signallers to display higher levels of aggression in noisier habitats. Consistent
with this, in several species of songbirds birds living in noisier urban habitats display higher
levels of aggression than birds living in rural habitats (Davies and Sewall, 2016; Evans et al.,
2010; Fokidis et al., 2011; Foltz et al., 2015; Hardman and Dalesman, 2018). Multiple recent
studies found more direct evidence for this hypothesis. In the first study male white-crowned
sparrows (*Zonotrichia leucophrys*) holding territories with greater ambient noise displayed
higher levels of aggression in response to song playback (Phillips and Derryberry, 2018). In
another study, male house wrens (*Troglodytes aedon*) responded to male song playback near
the nests with higher levels of aggression (Grabarczyk and Gill, 2019). Another recent study
in chiffchaffs (*Phylloscopus collybita*) found that birds that were exposed to airport noise also
responded more aggressively to conspecific playbacks compared to birds that were not
exposed to noise. In contrast to these studies, a study examining territorial response in
chipping sparrows (*Spizella passerina*) and spotted towhees (*Pipilo maculatus*) found that
ambient noise was negatively correlated with approach latency to speaker in both of these
species (Kleist et al., 2016). This effect may come about if ambient noise makes playbacks
less detectable or harder to locate.

Noise also has an effect on singing behaviours, most notably in change of amplitude
and frequency (Brumm and Slabbekoorn, 2005; Brumm and Zollinger, 2013), but also song
rate (Brumm and Slater, 2006) and song duration (Ríos-Chelén et al., 2013). These changes
may be adaptive in terms of ensuring the transmission of the signal to the receivers. Song
overlapping may also change as a result of ambient noise: individuals may be less likely to
overlap their opponent's songs in noisy environments if ambient noise already interferes with
the transmission of the signals. To the best of our knowledge, no previous study examined
whether song overlapping varied with ambient noise levels.

In the present study, we ask whether song overlapping and ambient noise levels are
associated with territorial aggression in a population of great tits that live in habitats that
range from urban to forest. As noted above, overlapping has been proposed to be an
aggressive signal but evidence for the predictive criterion has been lacking. Previous research
on this species showed that male great tits respond to anthropogenic noise by changing their
song and behaviours: for instance, great tits in noisy urban areas sing songs with higher

minimum frequencies than great tits in quieter rural areas, which may benefit them as low-

frequency songs are more prone to interference by anthropogenic noise (Hamao et al., 2011;

Mockford and Marshall, 2009; Slabbekoorn and den Boer-Visser, 2006; Slabbekoorn and

Peet, 2003). Urban-living great tits have also been found to be more aggressive than rural-

living great tits (Hardman and Dalesman, 2018).

We studied great tits in an urban-forest gradient in Ankara, Turkey. In our population,

located on the campus of Middle East Technical University, great tits hold territories

throughout built-up areas as well as the forested areas adjoining the populated areas.

Background noise in these areas vary significantly depending on proximity to buildings and

roads, allowing us to ask whether ambient noise is correlated with aggression. We asked

three specific questions: 1) Is song overlapping positively correlated with aggression across

males? 2) Does ambient noise on territory predict aggressive behaviours? 3) Is ambient noise

correlated with singing behaviours such as song overlapping, song rate, and song duration?

Methods

Study site and subjects

We studied great tits holding territories on the campus of Middle East Technical University,

in suburban Ankara, Turkey (39°53'32"N, 32°47'03"E). Although the campus is located in a

steppe habitat, it includes a large area that has been afforested mostly with conifers in the last

six decades. Great tits nest in human structures on campus (e.g. access ports left open in

telephone and electricity poles or cavities on the side of buildings) as well as nest boxes

provided in the forested parts and natural cavities spread throughout the campus grounds. We

located 42 territories of great tits by observation of singing posts or locating nests in the

boxes or human structures. The birds were not captured at any time and therefore were not

banded. The trials were carried out between 7 and 13 April 2019 at the start of the nesting

period in the morning hours between 0600 and 1130. We avoided testing neighbouring males

on the same day and carried out consecutive trials in locations that are outside of earshot of

each other.

Playback Stimuli

We recorded male songs from great tits on campus in Spring 2018 and 2019 using a Marantz

PMD660 or 661 recorder and a Sennheiser ME66/K6 shotgun microphone. Using the

software Syrinx (John Burt, Seattle, WA) we viewed and selected high quality recordings of

songs to create playback stimuli. After manually filtering out low-frequency noise (<~1000

Hz), we added a silent period to create a 7-second wave file. The average (±SD) playback

song duration was 2.99 (±0.42) seconds (range= 2.01-3.55 seconds). We created 24 playback

stimuli. For each subject we used a stimulus song that was recorded at least 1km away from

the subject's territory. This constraint meant we used some stimulus songs multiple times for

different males (11 stimuli were used in two trials, 2 stimuli were used in three trials, and one

stimulus was used in 4 trials- the rest of the songs were used once).

Procedure

We placed a wireless speaker (Anker SoundCore, Anker, Inc) was placed face-up at a natural

perch inside the territory of the male (either 5 m from the nest if the nest was known or at a

location that was central to the various singing posts the male was observed to be singing).

We then set up flagging at 1m, 3m and 5m from the speaker on either side of the speaker to

help with distance estimation during the trial. The observers then stepped back to about 15 m

from the speaker. The playbacks were controlled from a smartphone connected to the speaker

via Bluetooth.

The trials were recorded with the same equipment as above. We started the trials by

playing back the stimulus song until the subject responded either by singing a song or

approaching the playback. After this first response, we continued the playback for another 3

minutes and we narrated the behaviour of the subject by noting each flight, distance to the

speaker and singing behaviours. The behaviours during this 3-minute period are the main

response variables of the study.

Noise measurements

After the trial, we removed the flagging, took a GPS reading (Garmin eTrek, Garmin Inc.) of

the trial location and carried out the ambient noise measurements using the method described

by Brumm (2004). Briefly, we took two measurements in each cardinal direction at the

location of the playback using a sound level meter (VLIKE VL6708, VLIKE Inc.) with A

weighting and fast response (125 msec) settings. In a subset of territories (n=32 out of 42) we

took a second measurement of ambient noise later in the day to assess how consistent

variation in the ambient noise was. For logistical reasons we could not take second noise

measurements in 10 territories. The first and second noise measurements were highly

correlated with each other (linear regression: β=0.94, SE= 0.17, t=5.46, p= 6.4 x 10^{-6}). We

used the first noise measurement in the analyses as this measurement was available for all

territories.

 Response measures

Viewing the trial recordings in Syrinx, we extracted the following behaviours: flights,

distance to speaker with each flight, and songs. Because the trial durations slightly varied

across trials (ranging between 179 seconds to 192 seconds), we converted the counts of

flights and songs into rates by dividing it with the trial duration. We also calculated the

proportion of the trial subjects spent within 1 m of the speaker and noted the closest approach

of the distance. For each song the subject sang, we determined whether the song overlapped

with the playback song and calculated the proportion of the songs that the subject sang that

overlapped the playback song. This measure therefore corrects for variation in song rate, but

it is undefined for subjects that did not sing any songs (n=7). Because of variation in duty

cycle (duration of stimulus with song/entire duration of stimulus) between stimulus songs, we

also classified subjects based on whether the observed song overlapping (as defined above)

was higher (n=11) or lower (n=24) than the duty cycle. We classified the subjects who did

not sing any songs as a third category in this variable. Finally, for each subject we measured

duration of each of their song and calculated the average song duration for each subject.

Data Analysis

The rates of flights, closest approach distance and proportion of time spent within 1m of

speaker were all highly correlated with each other (Table 1). We therefore used a principle

component analysis (PCA) to arrive at a single measure of aggressive response. The first

component of the PCA (PCA1) explained 68% of variation and was taken as the aggressive

score. The first component was positively correlated with flight rates and time spent within

1m, and negatively correlated with closest approach distance (Table 1).

We first ran two linear mixed models (LMM) with aggression scores (PCA1) as the

response variable and stimulus song as a random factor. In the first model we entered the

ambient noise levels measured after the trial as the fixed factor while in the second model we

entered song overlapping as the fixed factor. We ran two separate models because song

overlapping (as a proportion of subject songs that overlapped) was not defined for 7 subjects

who did not sing any songs during the trial, thus reducing the sample size.

We then ran a parallel LMM to the first set of models with aggression scores again as

response variable and stimulus song as random factor. In this model we entered ambient

noise levels and the categorical overlapping variable as the predictor. This variable had three

categories high-overlapping (overlapping rates higher than the stimulus duty cycle), low-

overlapping (overlapping rates lower than the stimulus duty cycle) or no song. This model
included all subjects.

We additionally asked whether ambient noise was related to singing behaviours. To
answer that question we ran LMMs on song rate, song overlapping and average song duration
as response variables, and ambient noise levels as predictor variables, with stimulus song as a
random factor, to determine whether ambient noise had an effect on other singing behaviours.
Finally, we compared the observed level of song overlapping with the expected levels
(represented by the duty cycle), by carrying out a paired t-test. The analyses were carried out
in R using the package nlme (Pinheiro et al., 2017; R Core Team, 2012).

Results

Contrary to the hypothesis that song overlapping is a signal of aggressive intent, overlapping
(proportion of subject songs that overlapped the playback songs) was negatively related to the
aggression scores: high overlapping was indicative of lower aggression scores (Table 2,
Figure 1). Ambient noise was positively related to aggression scores: birds in noisier
territories were more aggressive in response to the playback (Table 2, Figure 2). In the model
with the categorical overlapping variable (Table 3), both overlapping and ambient noise
significantly predicted attack: birds that overlapped the playback less than expected were
more aggressive compared to high-overlapping birds (unpaired t test: t(33)=2.78, p=0.0088).
The most aggressive birds were birds who did not sing any songs who were more aggressive
than low-overlapping birds (unpaired t-test: t(29)=2.10, p=0.044; Figure 3).

Ambient noise was not related to song overlapping, song rate or average song
duration (Table 4). The observed levels of overlapping (mean ± SD: 0.33 ± 0.29) was lower
than the duty cycle of stimulus songs (0.42 ± 0.06) although not significantly so; unpaired t-
test unequal variances: t(34)=-1.77, p=0.085.

Discussion

In this study we asked whether song overlapping and ambient noise was related to territorial
aggression in male great tits in a population that spans an urban-rural gradient. We found that
song overlapping was negatively related to aggressive behaviours suggesting that high levels
of overlapping is not a signal males use to threaten opponents. We also found that ambient
noise levels were positively related to aggression scores, providing support for the hypothesis
that high ambient noise levels may select for higher aggression levels. Nonetheless, ambient

noise levels were not related to singing behaviours and parameters such as song rate, duration
or overlapping.

Is song overlapping an aggressive signal?

The present results suggest that song overlapping is not necessarily a reliable signal of
aggressive intent in great tits. Our finding of a negative correlation between aggression and
song overlapping is similar to what Vehrencamp and colleagues (2007) found in banded
wrens. In another species, the black-capped chickadees, song overlapping was not associated
with aggressive behaviours or attack (Baker et al., 2012). Taken together, these results imply
that song overlapping does not necessarily carry information regarding high likelihood of
escalation to the receivers although it remains to be seen whether the negative conclusion will
apply generally to other species. Additionally, song overlapping tended to happen slightly
lower than expected based on the duty cycle of stimulus songs but this difference was not
significant, indicating that overlapping may not be a deliberate behaviour in any case.

It is worth noting that there is some debate recently about what the correct null
expectation should be in studies of song overlapping (Masco et al., 2016; Naguib and
Mennill, 2010; Searcy and Beecher, 2009, 2011). We used stimulus duty cycle (the ratio of
the stimulus song duration over the sum of the stimulus song duration and silent period
between two stimulus songs) as the expected levels simply because it does not rely on any
additional assumptions. Choosing a different "null" model, with varying assumptions, may
yield a result indicating a significant difference between expected and observed levels of
overlapping (Masco et al., 2016). Nonetheless, whatever the null model, if song overlapping
is a deliberate behaviour that individuals use to signal their intent to escalate, individual
variation in song overlapping should correlate positively with aggressive behaviours.

If song overlapping is not an aggressive signal in great tits, what does it indicate? One
possible answer is that high levels of overlapping may simply indicate that the overlapping
bird is not engaged with the intruder, whereas low levels of overlapping may indicate a closer
engagement. In fact, birds who approached the speaker more closely and flew around the
speaker more in search of the opponent may be avoiding overlapping the opponent as an
assessment strategy (Arnott and Elwood, 2009). Great tit song has been found to contain
information regarding the singer's traits such as age and condition (Rivera-Gutierrez et al.,
2010). Thus, it may be advantageous for receivers that intend to engage the intruder at close
range to listen carefully to the song features of the opponent. This listening strategy may

explain the low rates of overlapping among aggressive individuals as well as the fact that
birds that do not sing are the most aggressive group.

It is also possible that non-overlapping the intruder may simply be due to a search
strategy to locate the opponent acoustically while the opponent is singing. Under this
hypothesis, the males are avoiding overlapping the intruder because they are trying to figure
out where the intruder is. Indeed, singing while the opponent is also singing is likely to
interfere with the processing of spatial information that can be obtained from the song of the
intruder. If that is the case, then birds that are aggressively responding to their opponents and
therefore more motivated to locate them would be less likely to overlap their opponent's
songs. This hypothesis is essentially an extension of the "readiness hypothesis" that was
proposed for explaining the occurrence of low amplitude "soft" songs that is used as a signal
of aggressive intent in several songbirds (Akçay and Beecher, 2012; Akçay et al., 2011). The
readiness hypothesis posits that for soft songs, the low amplitude is simply a by-product of
the fact that the beak and body movements required to sing loudly might interfere with visual
tracking of an opponent. Therefore, in an aggressive interaction birds that intend to escalate
might be signing with their beak closed, which leads to lower amplitudes (Akçay and
Beecher, 2012; Goller et al., 2004). Similarly, singing while the intruder is also singing (i.e.
overlapping) may interfere with localizing of the opponent acoustically and birds that intend
to escalate the interaction may therefore avoid overlapping the opponent's song. This
hypothesis can be tested with an experiment in which the simulated intruder can be either
located by the subject either easily or with difficulty.

Another possibility is that overlapping, instead of being a signal of aggressive intent,
may be correlated with more permanent traits of the signallers, such as some aspect of
genetic quality (Helfer and Osiejuk, 2015). In their recent review, Helfer and Osiejuk (2015)
presented several hypotheses to that positing that song overlapping is a signal of male quality
due to some a physical or physiological constraint or a handicapping mechanism. However,
given that song overlapping is a rather flexible behaviour that primarily depends on the
chosen timing of the singing it seems unlikely that it would be subject to a physical or
physiological constraint or handicapping mechanism. Interestingly, a study on great tits
found that great tits that were exposed to an ectoparasite as nestlings were found to be less
likely to overlap the song playback carried out in the territory after they were recruited to the
population (Bischoff et al., 2009). Song overlapping however, was not correlated with the
condition of the male as a nestling, and there was only a trend towards a positive association

between overlap and male condition as an adult. Thus, although these data suggest a potential

developmental constraint on song overlapping, the mechanism is unclear. Bischoff et al.

(2009) do not present any information on aggressive behaviours such as approach, thus it is

not clear whether the developmental handicap was associated with aggressive behaviours as

well. In summary, while it remains possible that overlapping reflects either male quality,

condition or developmental history, more data are clearly needed to test these possibilities.

Noise and aggressive behaviours

Ambient noise was positively correlated with aggressive behaviours in the present study,

consistent with the hypothesis that noise may play a significant role in determining

aggressive phenotypes observed in urban populations of birds (Davies and Sewall, 2016;

Evans et al., 2010; Hardman and Dalesman, 2018). The effect of noise on aggression may

come about via multiple mechanisms. First, noise may disrupt effective communication

(McMullen et al., 2014; Ríos-Chelén et al., 2015). When effective communication is

disrupted, aggressive behaviours may increase in prevalence (Kareklas et al., 2019; Logue et

al., 2010). Interestingly, unlike other species (Brumm and Slater, 2006; Ríos-Chelén et al.,

2013) great tits in our population did not show any evidence that they change their song rate,

average song duration or overlapping depending on ambient noise levels. Given that previous

research showed that songs in great tits and other songbirds become less effective in urban

noise (Halfwerk et al., 2011; Luther et al., 2015), this finding suggests communication may

indeed be hampered in noisier territories.

A second, non-mutually exclusive possibility is that noise may lead to higher levels of

physiological arousal or stress which may increase aggression, as has been documented in

humans (Baron and Richardson, 2004). In particular, exposure to noise may elevate baseline

levels of glucocorticoids (Creel et al., 2002; Crino et al., 2011; Davies et al., 2017; but see

Kleist et al., 2018) which in turn may lead to a higher level of aggression in individuals living

in noisier areas. Interestingly, in a recent study on urban and rural song sparrows (*Melospiza*

melodia), Davies and colleagues (2018) found that baseline levels of corticosterone (the main

avian glucocorticoid) in blood samples after a simulated territorial intrusion did not differ

between urban and rural birds, and the levels of corticosterone were positively related to

territorial aggression males displayed. For a given level of plasma corticosterone however,

urban birds displayed more aggression than rural birds, suggesting that urban birds

potentially show differences in glucocorticoid signalling from rural birds. Thus, the exact

physiological mechanisms that determines the relationship between urbanization and noise on

the one hand, and aggression on the other is yet unclear (Bonier, 2012). Further experimental
studies are needed to quantify the effect of noise on plasticity at behavioural and
physiological levels to tests hypotheses on the physiological mechanisms of aggression under
noisy conditions.

Conclusion

In summary our findings on song overlapping suggest that this signal is unlikely to be a
signal of aggressive intent, and researchers should pursue new hypotheses for this interesting
behaviour. Our findings on ambient noise meanwhile, corroborate the recent studies linking
noise and aggression and suggest new avenues to investigate the effect of anthropogenic
noise on animal social behaviour and the physiological mechanisms underlying it. Together,
the current study highlights how animals may manage interference from noise or other
signallers in their social interactions.

References

Akçay Ç, Beecher MD, 2012. Signalling while fighting: further comments on soft song. Anim Behav 83:e1-e3.

Akçay Ç, Tom ME, Holmes D, Campbell SE, Beecher MD, 2011. Sing softly and carry a big stick: signals of aggressive intent in song sparrows. Anim Behav 82:377- 382.

Arnott G, Elwood RW, 2009. Assessment of fighting ability in animal contests. Anim Behav 77:991-1004.

Baker TM, Wilson DR, Mennill DJ, 2012. Vocal signals predict attack during aggressive interactions in black-capped chickadees. Anim Behav 84:965-974.

Baron RA, Richardson DR, 2004. Human aggression: Springer Science & Business Media.

Bee MA, Micheyl C, 2008. The cocktail party problem: what is it? How can it be solved? And why should animal behaviorists study it? Journal of comparative psychology 122:235.

Bischoff LL, Tschirren B, Richner H, 2009. Long-term effects of early parasite exposure on song duration and singing strategy in great tits. Behav Ecol 20:265-270. doi: 10.1093/beheco/arp012.

Bonier F, 2012. Hormones in the city: endocrine ecology of urban birds. Horm Behav 61:763-772.

Brumm H, 2004. The impact of environmental noise on song amplitude in a territorial bird. J Anim Ecol 73:434-440. doi: doi:10.1111/j.0021-8790.2004.00814.x.

Brumm H, 2006. Signalling through acoustic windows: nightingales avoid interspecific competition by short-term adjustment of song timing. Journal of Comparative Physiology a-Neuroethology Sensory Neural and Behavioral Physiology 192:1279-1285.

Brumm H, Slabbekoorn H, 2005. Acoustic communication in noise. Advances in the Study of Behavior 35:151-209.

Brumm H, Slater P, 2006. Ambient noise, motor fatigue, and serial redundancy in chaffinch song. Behav Ecol Sociobiol 60:475-481.

Brumm H, Todt D, 2002. Noise-dependent song amplitude regulation in a territorial songbird. Anim Behav 63:891-897.

Brumm H, Zollinger SA, 2013. Avian vocal production in noise. Animal communication and noise: Springer. p. 187-227.

Caryl PG, 1979. Communication by agonistic displays: what can game theory contribute to ethology? Behaviour 68:136-169.

Creel S, Fox JE, Hardy A, Sands J, Garrott B, Peterson RO, 2002. Snowmobile activity and
glucocorticoid stress responses in wolves and elk. Conserv Biol 16:809-814.

Crino O, Van Oorschot BK, Johnson E, Malisch J, Breuner C, 2011. Proximity to a high traffic
road: glucocorticoid and life history consequences for nestling white-crowned
sparrows. Gen Comp Endocrinol 173:323-332.

Dabelsteen T, McGregor PK, Holland J, Tobias JA, Pedersen SB, 1997. The signal function of
overlapping singing in male robins. Anim Behav 53:249-256.

Dabelsteen T, McGregor PK, Shepherd M, Whittaker X, Pedersen SB, 1996. Is the Signal
Value of Overlapping Different from That of Alternating during Matched Singing in
Great Tits? J Avian Biol 27:189-194. doi: 10.2307/3677221.

Davies S, Beck ML, Sewall KB, 2018. Territorial aggression in urban and rural Song Sparrows
is correlated with corticosterone, but not testosterone. Horm Behav 98:8-15. doi:
https://doi.org/10.1016/j.yhbeh.2017.11.010.

Davies S, Haddad N, Ouyang JQ, 2017. Stressful city sounds: glucocorticoid responses to
experimental traffic noise are environmentally dependent. Biol Lett 13:20170276. doi:
doi:10.1098/rsbl.2017.0276.

Davies S, Sewall KB, 2016. Agonistic urban birds: elevated territorial aggression of urban song
sparrows is individually consistent within a breeding period. Biol Lett 12:20160315.

Endler JA, 1992. Signals, signal conditions, and the direction of evolution. The American
Naturalist 139:S125-S153.

Endler JA, Basolo AL, 1998. Sensory ecology, receiver biases and sexual selection. Trends
Ecol Evol 13:415-420.

Evans J, Boudreau K, Hyman J, 2010. Behavioural Syndromes in Urban and Rural Populations
of Song Sparrows. Ethology 116:588-595. doi: 10.1111/j.1439-0310.2010.01771.x.

Ey E, Fischer J, 2009. The "acoustic adaptation hypothesis"—a review of the evidence from
birds, anurans and mammals. Bioacoustics 19:21-48.

Ficken RW, Ficken MS, Hailman JP, 1974. Temporal pattern shifts to avoid acoustic
interference in singing birds. Science 183:762-763.

Fitzsimmons LP, Foote JR, Ratcliffe LM, Mennill DJ, 2008. Frequency matching, overlapping
and movement behaviour in diurnal countersinging interactions of black-capped
chickadees. Anim Behav 75:1913-1920.

Fokidis HB, Orchinik M, Deviche P, 2011. Context-specific territorial behavior in urban birds:
no evidence for involvement of testosterone or corticosterone. Horm Behav 59:133-
143.

Foltz SL, Ross AE, Laing BT, Rock RP, Battle KE, Moore IT, 2015. Get off my lawn:
increased aggression in urban song sparrows is related to resource availability. Behav
Ecol 26:1548-1557.

Gil D, Brumm H, 2014. Acoustic communication in the urban environment: patterns,
mechanisms, and potential consequences of avian song adjustments. In: Gil D, Brumm
H, editors. Avian Urban Ecology Oxford: Oxford University Press.

Gil D, Honarmand M, Pascual J, Pérez-Mena E, Macías Garcia C, 2014. Birds living near
airports advance their dawn chorus and reduce overlap with aircraft noise. Behav Ecol
26:435-443.

Gochfeld M, 1978. Intraspecific Social Stimulation and Temporal Displacement of Songs of
the Lesser Skylark, Alauda gulgula. Zeitschrift für Tierpsychologie 48:337-344. doi:
10.1111/j.1439-0310.1978.tb00264.x.

Goller F, Mallinckrodt MJ, Torti SD, 2004. Beak gape dynamics during song in zebra finch. J
Neurobiol 59:289-303.

Grabarczyk EE, Gill SA, 2019. Anthropogenic noise affects male house wren response to but
not detection of territorial intruders. PLOS ONE 14:e0220576. doi:
10.1371/journal.pone.0220576.

Halfwerk W, Bot S, Buikx J, van der Velde M, Komdeur J, ten Cate C, Slabbekoorn H, 2011.
Low-frequency songs lose their potency in noisy urban conditions. Proceedings of the
National Academy of Sciences 108:14549-14554.

Halfwerk W, Bot S, Slabbekoorn H, 2012. Male great tit song perch selection in response to
noise-dependent female feedback. Funct Ecol 26:1339-1347.

Halfwerk W, Slabbekoorn H, 2009. A behavioural mechanism explaining noise-dependent
frequency use in urban birdsong. Anim Behav 78:1301-1307.

Hamao S, Watanabe M, Mori Y, 2011. Urban noise and male density affect songs in the great
tit Parus major. Ethol Ecol Evol 23:111-119.

Hardman SI, Dalesman S, 2018. Repeatability and degree of territorial aggression differs
among urban and rural great tits (Parus major). Scientific Reports 8:5042.

Helfer B, Osiejuk TS, 2015. It takes all kinds in acoustic communication: a new perspective
on the song overlapping phenomenon. Ethology 121:315-326.

Johnson MT, Munshi-South J, 2017. Evolution of life in urban environments. Science
358:eaam8327.

Kareklas K, Wilson J, Kunc HP, Arnott G, 2019. Signal complexity communicates aggressive
intent during contests, but the process is disrupted by noise. Biol Lett 15:20180841.
doi: doi:10.1098/rsbl.2018.0841.

Kleist NJ, Guralnick RP, Cruz A, Francis CD, 2016. Anthropogenic noise weakens territorial
response to intruder's songs. Ecosphere 7:e01259.

Kleist NJ, Guralnick RP, Cruz A, Lowry CA, Francis CD, 2018. Chronic anthropogenic noise
disrupts glucocorticoid signaling and has multiple effects on fitness in an avian
community. Proceedings of the National Academy of Sciences 115:E648-E657.

Langemann U, Tavares JP, Peake TM, McGregor PK, 2000. Response of great tits to escalating
patterns of playback. Behaviour 137:451-471.

Logue DM, Abiola IO, Rains D, Bailey NW, Zuk M, Cade WH, 2010. Does signalling mitigate
the cost of agonistic interactions? A test in a cricket that has lost its song. Proceedings
of Royal Society of London, B doi:10.1098/rspb.2010.0421.

Luther DA, Phillips J, Derryberry EP, 2015. Not so sexy in the city: urban birds adjust songs
to noise but compromise vocal performance. Behav Ecol 27:332-340.

Masco C, Allesina S, Mennill DJ, Pruett-Jones S, 2016. The Song Overlap Null model
Generator (SONG): a new tool for distinguishing between random and non-random
song overlap. Bioacoustics 25:29-40.

McMullen H, Schmidt R, Kunc HP, 2014. Anthropogenic noise affects vocal interactions.
Behav Processes 103:125-128. doi: https://doi.org/10.1016/j.beproc.2013.12.001.

Mennill DJ, Ratcliffe LM, 2004. Overlapping and matching in the song contests of black-
capped chickadees. Anim Behav 67:441-450.

Mockford EJ, Marshall RC, 2009. Effects of urban noise on song and response behaviour in
great tits. Proceedings of the Royal Society of London B: Biological
Sciences:rspb20090586.

Naguib M, Mennill DJ, 2010. The signal value of birdsong: empirical evidence suggests song
overlapping is a signal. Anim Behav 80:e11-e15.

Peake TM, Terry AM, McGregor PK, Dabelsteen T, 2001. Male great tits eavesdrop on
simulated male-to-male vocal interactions. Proc R Soc Lond, Ser B: Biol Sci 268:1183-
1187.

Phillips JN, Derryberry EP, 2018. Urban sparrows respond to a sexually selected trait with
increased aggression in noise. Scientific Reports 8:7505. doi: 10.1038/s41598-018-
25834-6.

Pinheiro J, Bates D, DebRoy S, Sarkar D, Heisterkamp S, Van Willigen B, Maintainer R, 2017.
Package 'nlme'. R package:3-1.

R Core Team, 2012. R: A language and environment for statistical computing. Vienna, Austria:
R Foundation for Statistical Computing.

Ríos-Chelén AA, Lee GC, Patricelli GL, 2015. Anthropogenic noise is associated with changes
in acoustic but not visual signals in red-winged blackbirds. Behav Ecol Sociobiol
69:1139-1151.

Ríos-Chelén AA, Quirós-Guerrero E, Gil D, Garcia CM, 2013. Dealing with urban noise:
vermilion flycatchers sing longer songs in noisier territories. Behav Ecol Sociobiol
67:145-152.

Rivera-Gutierrez HF, Pinxten R, Eens M, 2010. Multiple signals for multiple messages: great
tit,< i> Parus major</i>, song signals age and survival. Anim Behav 80:451-459.

Searcy WA, Anderson RC, Nowicki S, 2006. Bird song as a signal of aggressive intent. Behav
Ecol Sociobiol 60:234-241.

Searcy WA, Beecher MD, 2009. Song as an aggressive signal in songbirds. Anim Behav
78:1281-1292.

Searcy WA, Beecher MD, 2011. Continued scepticism that song overlapping is a signal. Anim
Behav 81:e1-e4.

Shannon G, McKenna MF, Angeloni LM, Crooks KR, Fristrup KM, Brown E, Warner KA,
Nelson MD, White C, Briggs J, 2016. A synthesis of two decades of research
documenting the effects of noise on wildlife. Biological Reviews 91:982-1005.

Slabbekoorn H, den Boer-Visser A, 2006. Cities change the songs of birds. Curr Biol 16:2326-
2331.

Slabbekoorn H, Peet M, 2003. Ecology: Birds sing at a higher pitch in urban noise. Nature
424:267.

Templeton CN, Zollinger SA, Brumm H, 2016. Traffic noise drowns out great tit alarm calls.
Curr Biol 26:R1173-R1174.

van Dongen WF, 2006. Variation in singing behaviour reveals possible functions of song in
male golden whistlers. Behaviour 143:57-82.

Vehrencamp SL, Hall ML, Bohman ER, Depeine CD, Dalziell AH, 2007. Song matching,
overlapping, and switching in the banded wren: the sender's perspective. Behav Ecol
18:849-859.

Wasserman FE, 1977. Intraspecific acoustical interference in the white-throated sparrow
(Zonotrichia albicollis). Anim Behav 25:949-952.

Wiley RH, 2006. Signal detection and animal communication. Advances in the study of

behavior 36:217-247.

Wilson DR, Ratcliffe LM, Mennill DJ, 2016. Black-capped chickadees, Poecile atricapillus,

avoid song overlapping: Evidence for the acoustic interference hypothesis. Anim Behav

114:219-229.

Wood WE, Yezerinac SM, 2006. Song sparrow (Melospiza melodia) song varies with urban

noise. Auk 123:650-659.

 Figures.

Figure 1. Scatterplot of song overlapping (proportion of the subject's songs that started while

the playback song was playing), and aggression scores (PCA 1).

Figure 2. Scatterplot of ambient noise levels (average of 8 measurements with A settings, dB)

and aggression scores (PCA1)

Figure 3. Aggression scores of subjects a) overlapping the playback songs at more than the
duty cycle (n=11), less than the duty cycle (n=24) , singing no song (n=7).

558
559

Table 1: Pearson correlation coefficients (p-values) for the correlation between the aggressive response variables and their loadings to the first component of principle component analysis (PCA).

	Flight rate	Proportion of trial spent within 1m	Closest approach
Flight rate	_	0.60 (2×10^{-5})	-0.57 (7×10^{-5})
Proportion of trial spent within 1m	_	_	-0.42 (0.005)
Loading Coefficient	0.614	0.564	-0.551

Table 2. Linear mixed models (LMM) with stimulus song as random factor and aggression scores as response variable. The first model has song overlapping as the fixed factor (n=35 subjects, 7 subjects are excluded because they did not sing a song, and proportion of their songs that overlapped the stimulus song was therefore undefined). The second model has ambient noise as the fixed factor (n=42 subjects).

Model	Coefficient (SE)	χ^2	p-value
Song overlapping	-1.89 (0.76)	6.27	0.012
Ambient noise	0.09 (0.035)	6.94	0.0083

Table 3. Linear mixed model on aggression scores with the categorical overlap variable and ambient noise as predictor variables. Stimulus song is a random factor.

	χ^2	p-value
Overlap	16.41	0.00027
Ambient noise	6.01	0.0083

Table 4. Linear mixed models on song rate, average song duration and song overlapping as response variables and ambient noise as predictor variable. Stimulus song is a random factor in all models.

Model	χ^2	p-value
Song rate	0.063	0.80
Song duration	0.69	0.79
Song overlapping	2.18	0.13