Using the Tea Bag Index to unravel how human pharmaceuticals affect aquatic

microbial activity.

William Ross Hunter[1,2*]; Ashley Williamson[2]; Judith Maria Sarneel[2]

[1]Agri-Food and Bioscience Institute Northern Ireland, Fisheries and Aquatic

Ecosystems Branch, Belfast, BT9 5PX, United Kingdom.

[2]University of Ulster, School of Geography and Environmental Science, Coleraine,

BT52 1SA, United Kingdom

[3]Umeå Universitet, Department of Ecology and Environmental Sciences, 901 87

Umeå, Sweden

Corresponding Author. Email: Billy.Hunter@afbini.gov.uk

Abstract

The constant release of complex mixture of pharmaceuticals, including antimicrobials

and endocrine disruptors, has the potential to affect aquatic microbial metabolism

and alter biogeochemical cycling of carbon and nutrients. Here we advance the Tea

Bag Index (TBI) for decomposition by using it in a series of contaminant exposure

experiments testing how interactions between two human pharmaceuticals, the

broad spectrum antibiotic trimethoprim and the artificial estrogen 17a-Ethinylestradiol

(EE2), affects microbial activity in an aquatic system. The TBI is a citizen science

tool used to test microbial activity by measuring the differential degradation of green

and rooibos tea as proxies for labile and recalcitrant organic matter decomposition.

Exposure to trimethoprim and EE2 had significant independent negative effects upon

decomposition of labile organic matter (green tea), suggesting additive effects upon

microbial activity. Exposure to EE2 alone negatively affected the degradation of

more recalcitrant organic matter (rooibos tea). Consequently, trimethoprim and EE2

stabilized labile organic matter against microbial degradation and restricted

degradation rates. We propose that the method outlined could provide a powerful

tool for testing the impacts of multiple interacting pollutants upon microbial activity, at

a range of scales, across aquatic systems and over biogeochemically relevant time

scales.

1. Introduction

Globally, human pharmaceutical use has increased by 3 % per annum since the year 2000, leading to constant discharge of these compounds into the aquatic environment from both point (Waste Water Treatment Works and Septic Tanks) and diffuse sources (e.g. agricultural run-off) (Gros et al, 2007; Rosi-Marshall & Royer, 2012; Van Boeckel et al, 2014;). Despite being present in the environment at low concentrations, pharmaceuticals have the potential to affect aquatic ecosystem processes because they are designed to be effective at micromolar or nanomolar concentrations (Rosi-Marshall & Royer, 2012). The specific effects of pharmaceuticals in the environment are likely to be complex as a consequence of the myriad of potential interactions between these compounds, and their potential effects upon aquatic plants, animals and microorganisms (Hernando et al, 2006; Rosi-Marshall & Royer, 2012; Brodin et al, 2014). For example, the presence of antimicrobial compounds has the potential to affect the microbial processes that control aquatic carbon and nitrogen cycling (Rosi-Marshall & Royer, 2012; Brodin et al, 2014), whilst other compounds such as artificial estrogens may stimulate microbial activity (Ribeiro et al., 2010; Pieper and Rotard, 2011; McClean and Hunter, 2020). As such, we need to understand how pharmaceuticals influence microorganism-mediated ecosystem processes.

Antibiotics and artificial estrogens represent some of the most widely detected pharmaceuticals in the environment (Álvarez-Muñoz et al, 2015; Archer et al, 2017; Rosi-Marshall & Royer, 2012). Within aquatic systems, a wide range of human antibiotics such as tetracyclines, sulphonimides and the broad spectrum antibiotic trimethoprim (Metcalfe et al., 2003; Ashton et al., 2004; Thomas and Hilton, 2004, Wiegel et al., 2004) are detectable. Their presence will, therefore, directly affect the structure of aquatic microbial communities (Wieser et al 2016; Yuan et al 2017), potentially altering the microbial processes which drive carbon and nitrogen cycling (Roose-Amsleg and Laverman, 2016). Whilst bacterial populations can rapidly evolve antibiotic resistance, this comes at a physiological cost. As a consequence antibiotic resistant bacteria tend to grow more slowly and use nutrients less efficiently than their non-resistant conspecifics (Anderson and Levin, 1999, Jansen et al 2013).

EE2 and other artificial estrogens, by contrast, can provide a novel carbon source for
aquatic microorganisms, supporting increased microbial metabolism (Ribeiro et al.,
2010; Pieper and Rotard, 2011), whilst their adsorption onto microbial biofilms may,
also, provide protection against antimicrobial agents (Writer et al., 2012; Zhang et
al., 2014). Artificial estrogens (such as EE2) may mediate the effects of antimicrobial
agents upon aquatic microbial communities (McClean and Hunter, 2020).
Consequently, interactions between antibiotics and artificial estrogens are of wider
environmental relevance.

Contaminant exposure experiment provides a powerful tool for testing the sensitivity
of aquatic microbial communities to pharmaceuticals and other pollutants (Tank et al,
2006; Costello et al, 2016). A well-refined method is provided by Costello et al (2016)
to test how pharmaceuticals affect microbial biofilm growth and community structure
and ecophsysiological responses of the biofilm, such as community respiration
(Rosi-Marshall et al, 2013; McClean & Hunter, 2020). These methods, however, do
not provide information on heterotrophic processes such as microbial degradation
organic matter, which occurs over times-scales measured in weeks or months
(Vannote et al, 1980; Raymond & Bauer, 2001). The Tea Bag Index (TBI) provides a
powerful low-cost tool for investigating microbial activity in soils and aquatic systems,
based upon the traditional use of leaf-litter bags in ecology (Keuskemp et al., 2013;
Whigham et al., 2017; Mueller et al., 2018; Seelen et al., 2019. The key strength of
TBI is its ability to provide a standardized method of quantifying microbial activity by
comparing the relative degradation of a labile (green tea) and recalcitrant (rooibos
tea) organic matter source. The use of TBI within contaminant exposure experiments
could, therefore, allow the impacts of exposure to a pollutant to be quantified in
terms of microbially-mediated organic matter degradation.

We tested how interactions between the broad-spectrum antibiotic Trimethoprim and
artificial estrogen EE2 affect in-stream microbial activity, using a modified
contaminant exposure experiments. These experiments integrate the TBI
(Keuskemp et al., 2013) into the contaminant exposure experiment methodology
developed by Tank et al (2006) and Costello et al (2016). Using this method we seek
to test the hypothesis that concurrent exposure to EE2 will reduce the effects of

trimethoprim upon the microbial degradation pathways for both labile and recalcitrant

organic matter in a mixed-use catchment stream.

2. Materials and Methods

Experiments were carried out between 22nd March and 13th June 2019 in the

Ballysally Blagh (Latitude: 55°08'48.9"N Longitude: 6°40'08.0"W), a ground-water fed

second-order stream draining a mixed agricultural (consisting of 21.9 % arable; 55.9

% grassland; 13.7 % heathland; 1.9 % woodland) and urban (7.3 %) catchment of

14.2 km^2. Mean volumetric water flow in the Ballysally Blagh is 0.21 (± 0.27) m^3 s^{-1},

measured at a V-shaped weir. Water temperatures at the study site (50 cm water

depth) were recorded at 1-hour intervals throughout the experiment using a HOBO

MX2204 Bluetooth temperature logger and ranged between 5.51 ^{o}C and 15.57 ^{o}C,

with a mean temperature over the study period of 10.17 (± 2.08) ^{o}C.

Contaminant exposure experiments were conducted using forty 120 ml screw cap

sample pots, with a 35 mm hole drilled into the lid (following Costello et al, 2016).

Each pot was filled with 100 ml of % agar impregnated, with ten spiked with a 688

μmol.l^{-1} dose of trimethoprim, ten with a 688 μmol.l^{-1} dose of EE2 and ten with 688

μmol.l^{-1} dose trimethoprim and a 688 μmol.l^{-1} EE2. The remaining ten sample pots

received no pharmaceutical treatment (control). Trimethoprim and EE2 stock

solutions for were made up in 70 % ethanol, with 1 ml aliquots used to spike each

contaminant exposure experiment, whilst the agar in the control treatments was

spiked with 1 ml doses of 70 % ethanol. All pots were weighed to determine mass of

agar within. One pre-weighed unused Lipton® Green Tea Sencha pyramidal tea bag

with non-woven fabric (Product Number: EAN 87 22700 05552 5) and one pre-

weighed unused Lipton® Rooibos tea bag with non-woven fabric (Product Number:

EAN 87 22700 18843 8) were the placed into each pot screw cap secured in place.

The contaminant exposure experiments were then secured to four L-shaped metal

bars (Length = 1000 mm; Width = 35 mm; Height = 35 mm) and deployed onto the

streambed at 30 – 40 cm depth (Figure 1).

Contaminant exposure experiments were left in place for 83 days. Upon recovery,

the teabags were removed from the experiment and the pots were then weighed to

determine the agar mass loss (following Costello et al, 2016). Teabags were dried to

constant weight over 72 hours at 60 ^{o}C, excess silt and detritus adhering to each

teabag was gently removed, and their dry-weight determined. The tea was then

removed from each teabag, re-weighed and combusted at 550 ^{o}C for four hours to

determine ash free dry mass (following Keuskamp et al, 2013).

We calculated percentage mass loss of green tea and rooibos tea in each

contaminant exposure experiment as the difference between the dry weight of the

teabag at the start and end of the incubation. We corrected for mass loss during the

drying process based on the mean mass loss for five air-dry green teabags and five

air-dry rooibos teabags, dried at 60 ^{o}C over 72 hours. Following Keuskamp et al.

(2013) we then determined the organic matter stabilization factor (S) as:

[1] $S = 1 - \dfrac{a_g}{0.842}$

Where a_g is the decomposable fraction of the green tea (ash free dry weight) and

0.842 is the hydrolysable fraction of the green tea (from Keuskamp et al, 2013).

Using S we then estimated the decomposable fraction of the rooibos tea (a_r) as:

[2] $a_r = 0.552\,(1 - S)$

Where S is the stabilization factor and 0.552 is the hydrolysable fraction of the

rooibos tea (from Keuskamp et al, 2013). The organic matter Initial decomposition

rate of the labile material (k) can then be calculated as:

[3] $k = \dfrac{\left(\ln\left(\frac{a_r}{W}\right)\right)}{t}$

Where W is the mass loss of the Rooibos teabag in each experiment (in grams) and

t is the duration of the experimental incubation (in days).

Data analysis and visualization was carried out in R using the *base* and *ggplot*

packages (R Development Core Team, 2009; Wickham, 2016). We tested for

independent and combined effects of trimethoprim and EE2 upon the percentage

mass loss of the Green tea and Rooibos teabags, the stabilization factor (S) and

Initial decomposition rate (k) of the labile organic material using two-way analysis of

variance (ANOVA). Post-hoc testing of significant interactions were conducted using

Tukey's test for Honest Significant Difference. All data were visually explored to

ensure they conformed to the assumptions of normality and homoscedacity

(following Zuur et al, 2010), Initial decomposition rate (k) data were $\log_{10}$ transformed

to ensure that the residuals from the ANOVA model conformed to a normal

distribution. See supplementary table 1 for a summary of the ANOVA test results.

3. Results and Discussion

Over the course of the experiment, dissolution of the agar delivered estimated daily

doses of approximately 275 nmol $.d^{-1}$ of trimethoprim and EE2 in both the single and

combined pharmaceutical treatments. As such, the pharmaceutical doses which the

microbial community with each teabag where exposed to was broadly comparable

with the concentration of these compounds in inland waters (e.g. Álvarez-Muñoz et

al, 2015; Archer et al, 2017; Tousova et al., 2017). Rooibos Tea mass loss

decreased as a consequence of EE2 exposure, with no additional/significant effects

of Trimethoprim (Figure 2 A, Table 1a). By contrast for green tea, we observed

additive and inhibiting effects of both trimethoprim and EE2 (Figure 2 B, Table 1b).

Based on these data we can demonstrate that EE2 and trimethoprim had significant

independent effects upon both the organic matter stabilization factor (Figure 3 A,

Table 1c) initial decomposition rate (Figure 3 B, Table 1 d).

Our results demonstrate that exposure to low doses of both an antibiotic

(trimethoprim) and an endocrine disruptor (EE2) inhibits microbial degradation, with

differential effects on different phases of the decomposition process. Our results,

however, do not show that EE2 reduces the impact of trimethoprim upon aquatic

organic matter degradation. Both trimethoprim and EE2 had significant additive

positive effects upon the organic matter stabilization factor (s) and its initial

decomposition rate (k). As such, our study indicates that pollution from both

compounds may increase the residence time for organic matter in aquatic systems.

Inland waters (rivers, lakes and streams) typically receive large inputs of terrestrial

organic matter, which is then partially metabolized, temporarily buried within the

sediment or transported to the ocean (Battin et al, 2009). Our study adds further

evidence that pharmaceutical contaminants are likely to have important implications

for the rates and pathways for aquatic carbon and nutrient cycling (e.g. Jobling et al.

2003; Hernando et al. 2006; Gros et al. 2007; Rosi-Marshall and Royer 2012; Rosi-

Marshall et al. 2013; Álvarez-Muñoz et al. 2015; Ruhí et al. 2016; Archer et al. 2017;

Rosi et al. 2018; Gallagher and Reisinger 2020; McClean and Hunter, 2020).

Contaminant exposure experiments are typically used to determine the effects of a
selected contaminant on biofilm forming bacteria and algae (Tank et al., 2006;
Costello et al., 2016). By integrating the TBI (Keuskemp et al. 2013) into contaminant
exposure experiments we have developed a low-cost tool to quantify how exposure
to pollutants such as pharmaceuticals affects instream decomposition of organic
matter. Whilst our study was restricted to one stream, the method could easily be
replicated at multiple sites, within more complex experimental designs. We,
therefore, foresee potential applications of this method to allow the effects of multiple
pollutants upon organic matter degradation and residence times to be made across a
range of spatial and temporal scales in both freshwater and marine systems. As our
method is characterized by its low-cost, simplicity and use of readily available
consumer products, it also provides a tool which can be used to achieve levels
replications within experimental designs testing the interactions between a suite of
multiple pollutants and their effects upon aquatic biogeochemical processes, Finally,
we propose TBI Contaminant Exposure Experiments could provide a valuable
pedagogical tool for research-led environmental science teaching across a range of
education contexts, as well as in citizen science based initiatives to investigate the
impacts of pollutants in aquatic systems.

Acknowledgements

This work was funded through start-up funds provided to WRH by the University of
Ulster School of Geography and Environmental Science. JMS acknowledges the
Swedish Research Council (Vetenskapsrådet) for funding.

Author Information

Experiments were designed by WRH following discussions with JS, and were
conducted by WRH and AW. All authors contributed to the writing of the manuscript.

Conflict of Interest

The authors declare that we have no conflicts of interest.

Data Availability

All data are freely available Creative Commons (CC-BY) license through the Zenodo
data repository (DOI: 10.5281/zenodo.3528009).

Animal Research (Ethics)

No animals were used in this research.

Consent to Participate (Ethics)

No human subjects were used in this research.

Consent to Publish (Ethics)

This is not applicable to this study.

Plant Reproducibility

This is not applicable to this study.

Clinical Trials Registration

This is not applicable to this study.

4. References

Álvarez-Muñoz D, Rodríguez-Mozaz S, Maulvault AL, Tediosi A, Fernández-Tejedor
M, Van den Heuvel F*, et al.* (2015). Occurrence of pharmaceuticals and endocrine
disrupting compounds in macroalgaes, bivalves, and fish from coastal areas in
Europe. Environmental Research 143:56-64.

Anderson DI, Levin BR (1999). The biological cost of antibiotic resistance. Current
Opinion in Microbiology. 2:489-493.

Archer E, Petrie B, Kasprzyk-Hordern B, Wolfaardt GM (2017). The fate of
pharmaceuticals and personal care products (PPCPs), endocrine disrupting
contaminants (EDCs), metabolites and illicit drugs in a WWTW and environmental
waters. Chemosphere 174:437-446.

Battin TJ, Luyssaert S, Kaplan LA, Aufdenkampe AK, Richter A, Tranvik LJ (2009).
The boundless carbon cycle. Nature Geoscience 2:598.

Brodin T, Piovano S, Fick J, Klaminder J, Heynen M, Jonsson M (2014). Ecological
effects of pharmaceuticals in aquatic systems: impacts through behavioural alter-
ations. Philosophical Transactions of the Royal Society of London B: Biological
Sciences 369:20130580.

Costello DM, Rosi-Marshall EJ, Shaw LE, Grace M, Kelly JJ (2016). A novel method to assess effects of chemical stressors on natural biofilm structure and function. Freshwater Biology 61:2129-2140.

Gros M, Petrović M, Barcelo D (2007). Wastewater treatment plants as a pathway for aquatic contamination by pharmaceuticals in the Ebro river basin (northeast Spain). Environmental Toxicology and Chemistry 26:1553-1562.

Hernando MD, Mezcua M, Fernández-Alba AR, Barceló D (2006). Environmental risk assessment of pharmaceutical residues in wastewater effluents, surface waters and sediments. Talanta 69:334-342.

Jansen G, Barbosa C, Schulenburg H (2013). Experimental evolution as an efficient tool to dissect adaptive paths to antibiotic resistance. Drug Resistance Updates 16: 96-107

Keuskamp JA, Dingemans BJJ, Lehtinen T, Sarneel JM, Hefting MM (2013). Tea Bag Index: a novel approach to collect uniform decomposition data across ecosystems. Methods in Ecology and Evolution 4:1070-1075.

McClean P, Hunter WR (2019). 17α-ethynylestradiol (EE2) limits the impact of ibuprofen upon respiration by streambed biofilms in a sub-urban stream. Enviromental Science and Pollution Research. 27: 37149–37154.

Mueller P, Schile-Beers, LM, Mozdzer TJ, Chmura GL, Dinter T, Kuzyakov Y, de Groot AV, Esselink P, Smit C, D'Alpaos A, Ibáñez C, Lazarus M, Neumeier U, Johnson BJ, Baldwin AH, Yarwood SA, Montemayor DI, Yang Z, Wu J, Jensen K, Nolte S (2018). Global-change effects on early-stage decomposition processes in tidal wetlands: implications from a global survey using standardized litter. Biogeosciences, 15, 3189–3202, 2018

Raymond PA, Bauer JE (2001). Riverine export of aged terrestrial organic matter to the North Atlantic Ocean. Nature 409:497-500.

Roose-Amsaleg C, Laverman AM (2016). Do antibiotics have environmental side-effects? Impact of synthetic antibiotics on biogeochemical processes. Environmental Science and Pollution Research. 23:4000–4012.

Rosi-Marshall EJ, Kincaid DWL, Bechtold H, Royer TV, Rojas M, Kelly JJ (2013). Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial communities in stream biofilms. Ecological Applications 23:583-593.

Rosi-Marshall EJ, Royer TV (2012). Pharmaceutical compounds and ecosystem function: an emerging research challenge for aquatic ecologists. Ecosystems. 15:867-880.

Seelen LMS, Flaim G, Keuskamp J, Teurlincx S, Arias Font R, Tolunay D, *et al.* (2019). An affordable and reliable assessment of aquatic decomposition: Tailoring the Tea Bag Index to surface waters. Water Research 151:31-43, doi:https://doi.org/10.1016/j.watres.2018.11.081.

Tank JL, Bernot MJ, Rosi-Marshall EJ (2006). Nitrogen limitation and uptake. In:
Methods in Stream Ecology. Academic Press: San Diego, CA., pp 213-238.

Tousova Z, Oswald P, Slobodnika J, Blaha L, Muzc M, Huc M, Brack W, Krauss M,
Di Paolo C, Tarcai Z, Seilerd T-B, Hollert H, et al. (2017). European demonstration
program on the effect-based and chemical identification and monitoring of organic
pollutants in European surface waters. Science of the Total Environment. 601–602:
1849-1868.

Van Boeckel TP, Gandra S, Ashok A, Caudron Q, Grenfell BT, Levin SA*, et al.*
(2014). Global antibiotic consumption 2000 to 2010: an analysis of national
pharmaceutical sales data. Lancet Infectious Diseases 14:742-750.

Vannote RL, Minshall GW, Cummins KW, Sedell JR, Cushing CE (1980). The River
Continuum Concept. Canadian Journal of Fisheries and Aquatic Sciences 37:130-
137.

Waiser MJ, Swerhone GD, Roy J, Tumber V, Lawrence JR (2016). Effects of
erythromycin, trimethoprim and clindamycin on attached microbial communities from
an effluent dominated prairie stream. Ecotoxicology and Environmental Safety 132:
31-39. doi: 10.1016/j.ecoenv.2016.05.026.

Whigham DF, Walker CM, Maurer J, King, RS, Hauser W, Baird S, Keuskamp JA,
Neale PJ (2017). Watershed influences on the structure and function of riparian
wetlands associated with headwater streams - Kenai Peninsula, Alaska. Science of
the Total Environment. 599–600: Pages 124-134

Yuan K, Xiao S, Jiang X, Yang L, Chen B, Luan T, Lin L, Tam NFY (2017). Effects of
endocrine disrupting chemicals (EDCs) on bacterial communities in mangrove
sediments. Marine Pollution Bulletin 122: 122-128

Table 1. Two-way analysis of variance testing for significant effects of trimethoprim
and EE2 upon the percentage mass loss of the Green tea and Rooibos teabags, the
organic matter stabilization factor (S) and initial decomposition rate (k).

a) Percentage mass loss of Green tea					
	Df	SSq	MSq	F	p
Trimethoprim	1	105.89	105.889	10.486	0.003 **
EE2	1	108.43	108.428	10.738	0.002 **
Trimethoprim:EE2	1	16.26	16.259	1.610	0.213
Residuals	36	363.52	10.098		
Adjusted R^2	0.3371				
b) Percentage mass loss of Rooibos tea					
	Df	SSq	MSq	F	p
Trimethoprim	1	8.430	8.430	1.342	0.254
EE2	1	70.510	70.510	11.225	0.002 **
Trimethoprim:EE2	1	8.399	8.399	1.337	0.255
Residuals	36	226.119	6.281		
Adjusted R^2	0.219				
c) stabilization factor (S) of the labile material					
	Df	SSq	MSq	F	p
Trimethoprim	1	0.015	0.015	10.486	0.003 **
EE2	1	0.015	0.015	10.738	0.002 **
Trimethoprim:EE2	1	0.002	0.002	1.610	0.213
Residuals	36	0.051	0.001		
Adjusted R^2	0.337				
d) Initial decomposition rate (k) of the labile material					
	Df	SSq	MSq	F	p
Trimethoprim	1	1.073	1.073	5.389	0.026 *
EE2	1	2.594	2.594	13.025	<0.001 ***
Trimethoprim:EE2	1	0.112	0.112	0.564	0.458
Residuals	35	6.972	0.199		
Adjusted R^2	0.29				

310 **Figures**

311 Figure 1. Images of (A) the TBI contaminant exposure experiments, (B) how they are

312 secured prior to the L-shaped metal bars and (C) their deployment at the streambed

313 (right panel)

316   Figure 2. Effects of Trimethoprim (

) and EE2 (♀) on the degradation (% mass
317   loss) of A) Green Tea and B) Rooibos Tea, as labile and refractory organic matter
318   sources. Inserts show pooled data where significant independent effects of either
319   Trimethoprim or EE2 where detected. Significance levels: *** $p < 0.001$; ** $p < 0.01$; *
320     $p < 0.05$.

321
322

323 Figure 3. Effects of Trimethoprim (💊) and EE2 (♀) on the A) labile organic matter

324 stabilization factor and B) organic matter degradation rate, calculated following

325 Keuskamp et al (2013). Inserts show pooled data where significant independent

326 effects of either Trimethoprim or EE2 where detected. Significance levels: *** p <

327 0.001; ** p < 0.01; * p < 0.05.

329