Using the Tea Bag Index to determine how two human pharmaceuticals affect

litter decomposition by aquatic microorganisms.

William Ross Hunter[1,2*]; Ashley Williamson[2]; Judith Maria Sarneel[3]

[1]Agri-Food and Bioscience Institute Northern Ireland, Fisheries and Aquatic

Ecosystems Branch, Belfast, BT9 5PX, United Kingdom.

[2]University of Ulster, School of Geography and Environmental Science, Coleraine,

BT52 1SA, United Kingdom

[3]Umeå University, Department of Ecology and Environmental Sciences, 901 87

Umeå, Sweden

Corresponding Author. Email: Billy.Hunter@afbini.gov.uk

Abstract

This study demonstrates that independent additive effects of two human

pharmaceuticals, the antibiotic trimethoprim and the artificial estrogen 17a-

Ethinylestradiol (EE2), inhibit plant litter decomposition by aquatic microorganisms.

The constant release of pharmaceuticals, such as these, has the potential to affect

aquatic microbial metabolism and alter biogeochemical cycling of carbon and

nutrients. Here we advance the Tea Bag Index (TBI) for decomposition by using it in

a series of contaminant exposure experiments testing how interactions between

trimethoprim and EE2 affect aquatic microbial activity. The TBI is a citizen science

tool used to test microbial activity by measuring the differential degradation of green

and rooibos tea as proxies for respectively labile and recalcitrant litter

decomposition. Exposure to either trimethoprim or EE2 decreased decomposition of

green tea, suggesting additive effects upon microbial activity. Exposure to EE2 alone

decreased rooibos tea decomposition. Consequently, trimethoprim and EE2

stabilized labile organic matter against microbial degradation and restricted

decomposition. We propose that the method outlined could provide a powerful tool

for testing the impacts of multiple interacting pollutants upon microbial activity, at a

range of scales, across aquatic systems and over ecologically relevant time scales.

1. Introduction

Globally, human pharmaceutical use has increased by 3 % per annum since the year 2000, leading to constant discharge of these compounds into the aquatic environment from both point (Waste Water Treatment Works and Septic Tanks) and diffuse sources (e.g. agricultural run-off) (Gros et al, 2007; Rosi-Marshall & Royer, 2012; Van Boeckel et al, 2014;). Despite being present in the environment at low concentrations, pharmaceuticals have the potential to affect aquatic ecosystem processes because they are designed to be effective at micromolar or nanomolar concentrations (Rosi-Marshall & Royer, 2012; van Broeckel et al., 2014; Álvarez-Muñoz et al. 2015). The specific effects of pharmaceuticals in the environment are likely to be complex as a consequence of the myriad of potential interactions between these compounds, and their potential effects upon aquatic plants, animals and microorganisms (Hernando et al, 2006; Rosi-Marshall & Royer, 2012; Brodin et al, 2014). For example, the presence of antimicrobial compounds has the potential to affect the microbial processes that control aquatic carbon and nitrogen cycling (Rosi-Marshall & Royer, 2012; Brodin et al, 2014; Rosi et al., 2018), whilst other compounds such as artificial estrogens may actually stimulate microbial activity (Ribeiro et al., 2010; Pieper and Rotard, 2011; McClean and Hunter, 2020). As such, we need to understand how the interactions between pharmaceuticals influence microorganism-mediated biogeochemical processes, to safeguard ecosystem services.

Antibiotics and artificial estrogens represent some of the most widely detected pharmaceuticals in the environment (Rosi-Marshall & Royer, 2012; Álvarez-Muñoz et al, 2015; Archer et al, 2017; Gaston et al., 2019). Within aquatic systems, a wide range of human antibiotics such as tetracyclines, sulphonimides and trimethoprim are detectable in the water column, sediments and biota (Metcalfe et al., 2003; Ashton et al., 2004; Thomas and Hilton, 2004, Wiegel et al., 2004; Ebele et al., 2017; Kotke et al., 2019). Trimethoprim is of particular environmental concern, due to its widespread usage in human and veterinary healthcare, where it is mainly used to treat bladder and urinary tract infections, gastro-intestinal problems, ear infections and as prohylaxis against opportunistic infections (World Health Organisation, 2019). Approximately 50 – 60 % of any administered trimethoprim dose is excreted in a

subject's urine, combined with its widespread use this provides a definite route into

the environment via municipal wastewater and leaching of agricultural effluents. As

such, alongside its categorization as an essential medicine (World Health

Organisation, 2019), trimethoprim is recognized as a potential environmental hazard

(Straub, 2013; Tell et al., 2019).

The presence of antibiotics and their residues, both in the water column and aquatic

sediments may, directly affect the structure (Wieser et al 2016; Yuan et al 2017) and

functioning of aquatic microbial communities. In inland waters, leaf litter and other

terrigenous phytodetritus represent an important energy and nutrient source within

the food-web (e.g. Vannote et al, 1980; Raymond & Bauer, 2001; Raymond et al.,

2016). Microbial decomposition is an essential process, which supports the transfer

of litter-derived energy and nutrients to organisms at higher trophic levels.

Consequently, antimicrobials are likely to directly affect the processes that heavily

depend on microbial communities, and alter ecosystem-scale carbon and nitrogen

cycling will be affected (Roose-Amsleg and Laverman, 2016; Robson et al., 2020).

Furthermore, whilst bacterial populations rapidly evolve antibiotic resistance, this

comes at a physiological cost. Antibiotic resistant bacteria tend to grow more slowly

and use nutrients less efficiently than their non-resistant conspecifics (Anderson and

Levin, 1999, Jansen et al 2013). As such, longer-term exposure to antimicrobial

contaminations is likely to have implications for the efficiency of microbially-mediated

processes in aquatic ecosystems (e.g. Rosi et al. 2018).

By contrast, artificial estrogens, such as EE2, may provide a novel carbon source for

aquatic microorganisms, supporting increased microbial metabolism (Ribeiro et al.,

2010; Pieper and Rotard, 2011), their adsorption onto microbial biofilms can provide

protection against antimicrobial agents (Writer et al., 2012; Zhang et al., 2014).This

was recently demonstrated by McClean and Hunter (2020), who showed that

exposure of streambed biofilms to EE2 counteracted the inhibitory effects of

ibuprofen upon microbial respiration. Consequently, the interactions between

antibiotics and artificial estrogens upon aquatic microbial processes are likely to

have wider environmental relevance.

Contaminant exposure substrata (CES) experiment provides a powerful tool for

testing the sensitivity of aquatic microbial communities to pharmaceuticals and other

pollutants (Tank et al, 2006; Costello et al, 2016). A well-refined method is provided
by Costello et al (2016) to test how pharmaceuticals affect microbial biofilm growth
and community structure and ecophsysiological responses of the biofilm, such as
community respiration (Rosi-Marshall et al, 2013; Rosi et al., 2018; Gallagher and
Reisinger, 2020; McClean & Hunter, 2020 Robson et al., 2020). These methods,
however, provide insufficient information on heterotrophic processes such as
microbial litter decomposition, which occurs over times-scales measured in weeks or
months (Vannote et al, 1980; Raymond & Bauer, 2001). The Tea Bag Index (TBI)
provides a powerful low-cost tool for investigating microbial activity in soils and
aquatic systems, based upon the traditional use of leaf-litter bags in ecology
(Keuskemp et al., 2013; Whigham et al., 2017; Mueller et al., 2018; Seelen et al.,
2019. The key strength of TBI is its ability to provide a standardized method of
quantifying microbial activity by comparing the relative degradation of green tea
(easy to decompose, or labile material) and rooibos tea (resistant to decomposition,
recalcitrant, lignified material) organic matter source. The use of TBI within
contaminant exposure substrata experiments will allow quantification of the impacts
of pollutant exposure on microbially-mediated litter decomposition.

Within this study we investigate how interactions between the antibiotic trimethoprim
and the artificial estrogen EE2 affect aquatic litter decomposition, using modified
contaminant exposure experiments. We integrate the TBI (Keuskemp et al., 2013)
into the existing CES methodology developed by Tank et al (2006) and Costello et al
(2016). This will provide a simple, low-cost tool for assessing how pharmaceuticals
(and other contaminants) may affect microbial litter decomposition. Using the TBI-
CES method we test how exposure of the microbial community to both trimethoprim
and EE2 affects the decomposition of both labile and recalcitrant litter in a mixed-use
catchment stream. We hypothesize that trimethoprim will inhibit decomposition while
EE2 may enhance it. We therefore expect that EE2 may mediate against any
inhibitory effects of trimethoprim upon the microbial activity that drives litter
decomposition.
122

2. Materials and Methods

2.1. Site Description

Experiments were carried out between 22nd March and 13th June 2019 in the Ballysally Blagh (Latitude: 55°08'48.9"N Longitude: 6°40'08.0"W), a ground-water fed second-order stream draining a mixed agricultural (consisting of 21.9 % arable; 55.9 % grassland; 13.7 % heathland; 1.9 % woodland) and urban (7.3 %) catchment of 14.2 km^2. Mean volumetric water flow in the Ballysally Blagh is 0.21 (± 0.27) m^3 s^{-1}, measured at a V-shaped weir. Water temperatures at the study site (50 cm water depth) were recorded at 1-hour intervals throughout the experiment using a HOBO MX2204 Bluetooth temperature logger and ranged between 5.51 ^{o}C and 15.57 ^{o}C, with a mean temperature over the study period of 10.17 (± 2.08) ^{o}C.

2.2. Contaminant Exposure Substrata Experiments

Contaminant exposure substrata (CES) experiments were conducted using forty 120 ml screw cap sample pots, with a 35 mm hole drilled into the lid (following Costello et al, 2016). Each pot was filled with 100 ml of 2 % agar, with ten pots spiked with 688 µmol.l^{-1} Trimethoprim (Sigma-Aldich, product no: 92131), ten with 688 µmol.l^{-1} of EE2 (Sigma-Aldrich, Product No. E4876) and ten with 688 µmol.l^{-1} trimethoprim and 688 µmol.l^{-1} EE2. The remaining ten sample pots received no pharmaceutical treatment (control). Trimethoprim and EE2 stock solutions for were made up in 70 % ethanol, with 1 ml aliquots used to spike each contaminant exposure experiment, whilst the agar in the control treatments was spiked with 1 ml doses of 70 % ethanol. All pots were weighed to determine mass of agar within. One pre-weighed unused Lipton® Green Tea Sencha pyramidal tea bag with non-woven fabric (Product Number: EAN 87 22700 05552 5) and one pre-weighed unused Lipton® Rooibos tea bag with non-woven fabric (Product Number: EAN 87 22700 18843 8) were the placed into each pot screw cap secured in place. The contaminant exposure experiments were then secured to four L-shaped metal bars (Length = 1000 mm; Width = 35 mm; Height = 35 mm) and deployed onto the streambed at 30 – 40 cm depth (Figure 1).

CES experiments were left in place for 83 days. Upon recovery, the teabags were removed from the experiment and the pots were then weighed to determine the agar mass loss (following Costello et al, 2016). Teabags were dried to constant weight

over 72 hours at 60 ^{o}C, excess silt and detritus adhering to each teabag was gently

removed, and their dry-weight determined (following Keuskamp et al, 2013). The tea

was then removed from each teabag, re-weighed and combusted at 550 ^{o}C for four

hours to determine ash free dry mass.

2.3.   Data Treatment

We calculated percentage mass loss of green tea and rooibos tea in each

contaminant exposure experiment as the difference between the dry weight of the

teabag at the start and end of the incubation. We corrected for mass loss during the

drying process based on the mean mass loss for five air-dry green teabags and five

air-dry rooibos teabags, dried at 60 ^{o}C over 72 hours. Following Keuskamp et al.

(2013) we then determined the organic matter stabilization factor (S_{TBI}) as:

[1] $S_{TBI} = 1 - \dfrac{a_g}{0.842}$

Where a_g is the decomposable fraction of the green tea (ash free dry weight) and

0.842 is the hydrolysable fraction of the green tea (from Keuskamp et al, 2013).

Using S we then estimated the decomposable fraction of the rooibos tea (a_r) as:

[2] $a_r = 0.552 \, (1 - S_{TBI})$

Where S_{TBI} is the stabilization factor and 0.552 is the hydrolysable fraction of the

rooibos tea (from Keuskamp et al, 2013). The organic matter Initial decomposition

rate of the labile material (k_{TBI}) can then be calculated as:

[3] $k_{TBI} = \dfrac{\left(\ln\left(\frac{a_r}{W}\right)\right)}{t}$

Where W is the mass loss of the Rooibos teabag in each experiment (in grams) and

t is the duration of the experimental incubation (in days).

2.4.   *Data Analysis*

Data analysis and visualization was carried out in R using the *base* and *ggplot*

packages (R Development Core Team, 2009; Wickham, 2016). We tested for

independent and combined effects of trimethoprim and EE2 upon the percentage

mass loss of the Green tea and Rooibos teabags, the stabilization factor (S_{TBI}) and

Initial decomposition rate (k_{TBI}) of the labile organic material using two-way analysis

of variance (ANOVA). Post-hoc testing of significant interactions were conducted

using Tukey's test for Honest Significant Difference. All data were visually explored

to ensure they conformed to the assumptions of normality and homoscedacity

(following Zuur et al, 2010), Initial decomposition rate (k_{TBI}) data were $\log_{10}$

transformed to ensure that the residuals from the ANOVA model conformed to a

normal distribution. See supplementary table 1 for a summary of the ANOVA test

results.

3. Results and Discussion

Over the course of the experiment, dissolution of the agar was assumed to deliver

constant daily doses of approximately 275 nmol $.d^{-1}$ of trimethoprim and/or EE2 in

both the single and combined pharmaceutical treatments. As such, each teabag

would have been exposed to a pharmaceutical dose that was broadly comparable

with the concentration of these compounds in inland waters (e.g. Álvarez-Muñoz et

al, 2015; Archer et al, 2017; Tousova et al., 2017). Rooibos Tea mass loss

decreased upon EE2 exposure, with no additional/significant effects of trimethoprim

(Figure 2 A, Table 1a). By contrast for green tea, we observed additive and inhibiting

effects of both trimethoprim and EE2 (Figure 2 B, Table 1b). Based on these data we

can demonstrate that EE2 and trimethoprim had significant independent effects upon

both the TBI stabilization factor (Figure 3 A, Table 1c) and initial decomposition rate

(Figure 3 B, Table 1 d).

Trimethoprim decreased mass loss of green tea but not of rooibos, while EE2

decreased both green tea and rooibos mass loss. Together, these pharmaceuticals

had additive negative effects on green tea mass loss but not on rooibos. No

significant interaction was observed for either green tea or rooibos. Further, EE2 and

trimethoprim had significant independent effects upon both the litter stabilization

factor (Figure 3 A, Table 1c) and initial decomposition rate (Figure 3 B, Table 1 d)

Our results demonstrate that exposure to low doses of both an antibiotic

(trimethoprim) and an endocrine disruptor (EE2) inhibit microbial litter decomposition.

Specifically, trimethoprim appears to affect decomposition of lignified material less

compared to more easy to degrade material, whilst EE2 had a broader effect across
litter types. In counter to our hypothesis, EE2 did not reduce the negative impact of
trimethoprim upon aquatic litter decomposition. Both trimethoprim and EE2 had
significant additive positive effects upon the organic matter stabilization factor (S_{TBI})
and its initial decomposition rate (k_{TBI}). Our study, thus, indicates that pollution from
both pharmaceuticals may increase the residence time for organic matter in aquatic
systems. Inland waters (rivers, lakes and streams) typically receive large inputs of
terrestrial organic matter, which is then partially metabolized, temporarily buried
within the sediment or transported to the ocean (Battin et al, 2009). Our study adds
further evidence to that pharmaceutical contaminants directly affect the rates and
pathways for aquatic carbon and nutrient cycling (e.g. Jobling et al. 2003; Hernando
et al. 2006; Gros et al. 2007; Rosi-Marshall and Royer 2012; Rosi-Marshall et al.
2013; Álvarez-Muñoz et al. 2015; Ruhí et al. 2016; Archer et al. 2017; Rosi et al.
2018; Gallagher and Reisinger 2020; McClean and Hunter, 2020).

Given the widespread use of trimethoprim as an antibiotic medication and its high
solubility in water [ranging from 500 mg l^{-1} at pH 8 to 15,500 mg l^{-1} at pH 5.5 (Dahlan
et al. 1987)], its effect upon microbial activity within the teabag experiment was
expected. As a highly soluble contaminant trimethoprim is likely to leach rapidly from
the agar within CES experiments, and thus affect microbial activity during the early
phases of the experiment. Thus the effects of trimethoprim upon labile litter
decomposition are likely to reflect the relatively high solubility of this contaminant. By
contrast, EE2 exhibits relatively low solubility in water, ranging between 2.92 and 4.8
mg l-1 (Lai et al., 2000; Ying et al. 2002; Guo and Hu, 2014). This property of EE2,
combined with its tendency to adsorb onto organic substances (Lai et al., 2000; Ying
et al., 2002; Writer et al., 2012; Zhang et al., 2014), provides a mechanism through
which the EE2 released by the agar would affect the degradation of more recalcitrant
and lignified litter. The additive effects of trimethoprim and EE2 upon litter
decomposition within our experiment were likely to be driven by differences in the
solubility of each compound and their potential to adsorb onto the tea. This highlights
how differences in the physico-chemical properties of pharmaceutical mediate their
potential impacts in aquatic biogeochemical processes.

CES experiments are typically used to determine the effects of a selected
contaminant on biofilm forming bacteria and algae (Tank et al., 2006; Costello et al.,
2016). Integrating the TBI (Keuskemp et al. 2013) into CES experiments provides a
useful low-cost tool to quantify how exposure to pollutants such as pharmaceuticals
affects instream decomposition of organic matter. The inclusion of both a labile
(green tea) and more recalcitrant (rooibos tea) organic matter source within the
experiment allows inferences to made upon the effects of a contaminant upon the
phenology of the litter decomposition process. Refinement of the method is,
however, necessary and should include investigation of agar dissolution and teabag
degradation rates, and how these may vary under differing climatic and hydrological
regimes (following Costello et al., 2016).

Whilst our study was restricted to one stream, the method could easily be replicated
at multiple sites, within more complex experimental designs. We believe this method
will allow the effects of multiple pollutants upon organic matter degradation and
residence times to be made across a range of spatial and temporal scales in both
freshwater and marine systems. As our method is characterized by its low-cost,
simplicity and use of readily available consumer products, it could potentially be used
to achieve high levels replications within experimental designs testing the
interactions between a suite of multiple pollutants and their effects upon aquatic
biogeochemical processes. Furthermore, we propose that the TBI-CES method
could provide a valuable pedagogical tool for research-led environmental science
teaching and in citizen science based initiatives to investigate the impacts of
pollutants in aquatic systems.

Acknowledgements

This work was funded through start-up funds provided to WRH by the University of
Ulster School of Geography and Environmental Science. JMS acknowledges the
Swedish Research Council (Vetenskapsrådet) for funding.

Author Information

Experiments were designed by WRH following discussions with JS, and were
conducted by WRH and AW. All authors contributed to the writing of the manuscript.

Conflict of Interest

The authors declare that we have no conflicts of interest pertaining to this study.

Data Availability

All data are freely available Creative Commons (CC-BY) license through the Zenodo
data repository (DOI: 10.5281/zenodo.3528009).

Animal Research (Ethics)

No animals were used in this research.

Consent to Participate (Ethics)

No human subjects were used in this research.

Consent to Publish (Ethics)

This is not applicable to this study.

Plant Reproducibility

This is not applicable to this study.

Clinical Trials Registration

This is not applicable to this study.

4. References

Álvarez-Muñoz D, Rodríguez-Mozaz S, Maulvault AL, Tediosi A, Fernández-Tejedor M, Van den Heuvel F, *et al.* (2015). Occurrence of pharmaceuticals and endocrine disrupting compounds in macroalgaes, bivalves, and fish from coastal areas in Europe. Environmental Research 143:56-64.

Anderson DI, Levin BR (1999). The biological cost of antibiotic resistance. Current Opinion in Microbiology. 2:489-493.

Archer E, Petrie B, Kasprzyk-Hordern B, Wolfaardt GM (2017). The fate of pharmaceuticals and personal care products (PPCPs), endocrine disrupting contaminants (EDCs), metabolites and illicit drugs in a WWTW and environmental waters. Chemosphere 174:437-446.

Battin TJ, Luyssaert S, Kaplan LA, Aufdenkampe AK, Richter A, Tranvik LJ (2009). The boundless carbon cycle. Nature Geoscience 2:598.

Brodin T, Piovano S, Fick J, Klaminder J, Heynen M, Jonsson M (2014). Ecological effects of pharmaceuticals in aquatic systems: impacts through behavioural alter-ations. Philosophical Transactions of the Royal Society of London B: Biological Sciences 369:20130580.

Costello DM, Rosi-Marshall EJ, Shaw LE, Grace M, Kelly JJ (2016). A novel method to assess effects of chemical stressors on natural biofilm structure and function. Freshwater Biology 61:2129-2140.

Dahlan, R., Mcdonald, C. and Sunderland, V.B. (1987), Solubilities and intrinsic dissolution rates of sulphamethoxazole and trimethoprim. Journal of Pharmacy and Pharmacology, 39: 246-251. https://doi.org/10.1111/j.2042-7158.1987.tb06261.x

Gallagher MT, Reisinger AJ (2020) Effects of ciprofloxacin on metabolic activity and algal biomass of urban stream biofilms. Sci Total Environ 706:135728. https://doi.org/10.1016/j.scitotenv.2019.135728

Gaston L, Lapworth DJ, Stuart M, Arnscheidt J (2019) Prioritization approaches for substances of emerging concern in groundwater: a critical review. Environ Sci Technol 53:6107–6122

Gros M, Petrović M, Barcelo D (2007). Wastewater treatment plants as a pathway for aquatic contamination by pharmaceuticals in the Ebro river basin (northeast Spain). Environmental Toxicology and Chemistry 26:1553-1562.

Hernando MD, Mezcua M, Fernández-Alba AR, Barceló D (2006). Environmental risk assessment of pharmaceutical residues in wastewater effluents, surface waters and sediments. Talanta 69:334-342.

Jansen G, Barbosa C, Schulenburg H (2013). Experimental evolution as an efficient
tool to dissect adaptive paths to antibiotic resistance. Drug Resistance Updates 16:
96-107

Keuskamp JA, Dingemans BJJ, Lehtinen T, Sarneel JM, Hefting MM (2013). Tea
Bag Index: a novel approach to collect uniform decomposition data across
ecosystems. Methods in Ecology and Evolution 4:1070-1075.

Lai KM, Johnson KL, Scrimshaw MD, Lester JN (2000). Binding of waterborne
steroid estrogens to solid phases in river and estuarine systems Environ. Sci.
Technol., 34: 3890-3894

McClean P, Hunter WR (2019). 17α-ethynylestradiol (EE2) limits the impact of
ibuprofen upon respiration by streambed biofilms in a sub-urban stream.
Enviromental Science and Pollution Research. 27: 37149–37154.

Mueller P, Schile-Beers, LM, Mozdzer TJ, Chmura GL, Dinter T, Kuzyakov Y, de
Groot AV, Esselink P, Smit C, D'Alpaos A, Ibáñez C, Lazarus M, Neumeier U,
Johnson BJ, Baldwin AH, Yarwood SA, Montemayor DI, Yang Z, Wu J, Jensen K,
Nolte S (2018). Global-change effects on early-stage decomposition processes in
tidal wetlands: implications from a global survey using standardized litter.
Biogeosciences, 15, 3189–3202, 2018

Raymond PA, Bauer JE (2001). Riverine export of aged terrestrial organic matter to
the North Atlantic Ocean. Nature 409:497-500.

Raymond PA, Saiers JE, Sobczak WV. "Hydrological and biogeochemical controls
on watershed dissolved organic matter transport: pulse-shunt concept." Ecology 97:
15-16.

Robson SV, Rosi EJ, Richmond EK, Grace MR (202 (2016) Environmental
concentrations of pharmaceuticals alter metabolism, denitrification, and diatom
assemblages in artificial streams. Freshwat Sci 39:256–267.
https://doi.org/10.1086/708893

Roose-Amsaleg C, Laverman AM (2016). Do antibiotics have environmental side-
effects? Impact of synthetic antibiotics on biogeochemical processes. Environmental
Science and Pollution Research. 23:4000–4012.

Rosi EJ, Bechtold HA, Snow D, Rojas M, Reisinger AJ, Kelly JJ (2018) Urban stream
microbial communities show resistance to pharmaceutical exposure. Ecosphere
9:e02041

Rosi-Marshall EJ, Kincaid DWL, Bechtold H, Royer TV, Rojas M, Kelly JJ (2013).
Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial
communities in stream biofilms. Ecological Applications 23:583-593.

Rosi-Marshall EJ, Royer TV (2012). Pharmaceutical compounds and ecosystem
function: an emerging research challenge for aquatic ecologists. Ecosystems.
15:867-880.

Seelen LMS, Flaim G, Keuskamp J, Teurlincx S, Arias Font R, Tolunay D*, et al.*
(2019). An affordable and reliable assessment of aquatic decomposition: Tailoring
the Tea Bag Index to surface waters. Water Research 151:31-43,
doi:https://doi.org/10.1016/j.watres.2018.11.081.

Straub JO (2013): An environmental risk assessment for human-use Trimethoprim in
European surface waters. Antibiotics, 2, 115–162

Tank JL, Bernot MJ, Rosi-Marshall EJ (2006). Nitrogen limitation and uptake. In:
Methods in Stream Ecology. Academic Press: San Diego, CA., pp 213-238.

Tell J, Caldwell DJ, Häner A, Hellstern J, Hoeger B, Journel R, Mastrocco F, Ryan J,
Snape J, Straub JO, Vestel J (2019): Science-based targets for antibiotics in
receiving waters from pharmaceutical manufacturing operations. Integr Environ Ass
Manag. 15(3):312–319. https://doi.org/10.1002/ieam.4141

Tousova Z, Oswald P, Slobodnika J, Blaha L, Muzc M, Huc M, Brack W, Krauss M,
Di Paolo C, Tarcai Z, Seilerd T-B, Hollert H, et al. (2017). European demonstration
program on the effect-based and chemical identification and monitoring of organic
pollutants in European surface waters. Science of the Total Environment. 601–602:
1849-1868.

Van Boeckel TP, Gandra S, Ashok A, Caudron Q, Grenfell BT, Levin SA*, et al.*
(2014). Global antibiotic consumption 2000 to 2010: an analysis of national
pharmaceutical sales data. Lancet Infectious Diseases 14:742-750.

Vannote RL, Minshall GW, Cummins KW, Sedell JR, Cushing CE (1980). The River
Continuum Concept. Canadian Journal of Fisheries and Aquatic Sciences 37:130-
137.

Waiser MJ, Swerhone GD, Roy J, Tumber V, Lawrence JR (2016). Effects of
erythromycin, trimethoprim and clindamycin on attached microbial communities from
an effluent dominated prairie stream. Ecotoxicology and Environmental Safety 132:
31-39. doi: 10.1016/j.ecoenv.2016.05.026.

Whigham DF, Walker CM, Maurer J, King, RS, Hauser W, Baird S, Keuskamp JA,
Neale PJ (2017). Watershed influences on the structure and function of riparian
wetlands associated with headwater streams - Kenai Peninsula, Alaska. Science of
the Total Environment. 599–600: 124-134

World Health Organization (2019). World Health Organization model list of essential
medicines: 21st list 2019. Geneva: World Health Organization. hdl:10665/325771.
WHO/MVP/EMP/IAU/2019.06.

Ying GG, Kookana RS, Ru YJ (2002). Occurrence and fate of hormone steroids in
the environment. Environ. Int., 28: 545-551

Yuan K, Xiao S, Jiang X, Yang L, Chen B, Luan T, Lin L, Tam NFY (2017). Effects of
endocrine disrupting chemicals (EDCs) on bacterial communities in mangrove
sediments. Marine Pollution Bulletin 122: 122-128

Tables

Table 1. Two-way analysis of variance testing for significant effects of trimethoprim

and EE2 upon the percentage mass loss of the Green tea and Rooibos teabags, the

organic matter stabilization factor (S_{TBI}) and initial decomposition rate (k_{TBI}).

a) Percentage mass loss of Green tea					
	Df	SSq	MSq	F	p
Trimethoprim	1	105.89	105.889	10.486	0.003 **
EE2	1	108.43	108.428	10.738	0.002 **
Trimethoprim:EE2	1	16.26	16.259	1.610	0.213
Residuals	36	363.52	10.098		
Adjusted R^2	0.3371				
b) Percentage mass loss of Rooibos tea					
	Df	SSq	MSq	F	p
Trimethoprim	1	8.430	8.430	1.342	0.254
EE2	1	70.510	70.510	11.225	0.002 **
Trimethoprim:EE2	1	8.399	8.399	1.337	0.255
Residuals	36	226.119	6.281		
Adjusted R^2	0.219				
c) stabilization factor (S_{TBI})					
	Df	SSq	MSq	F	p
Trimethoprim	1	0.015	0.015	10.486	0.003 **
EE2	1	0.015	0.015	10.738	0.002 **
Trimethoprim:EE2	1	0.002	0.002	1.610	0.213
Residuals	36	0.051	0.001		
Adjusted R^2	0.337				
d) Initial decomposition rate (k_{TBI})					
	Df	SSq	MSq	F	p
Trimethoprim	1	1.073	1.073	5.389	0.026 *
EE2	1	2.594	2.594	13.025	<0.001 ***
Trimethoprim:EE2	1	0.112	0.112	0.564	0.458
Residuals	35	6.972	0.199		
Adjusted R^2	0.29				

409 **Figures**

410 Figure 1. Images of (A) the TBI contaminant exposure experiments, (B) how they are
411 secured prior to the L-shaped metal bars and (C) their deployment at the streambed
412 (right panel)

413
414

Figure 2. Effects of TMP ( ) and EE2 ( ) on the degradation (% mass loss) of A)
Rooibos and B) Green tea, as labile and recalcitrant organic matter sources. Inserts
show pooled data where significant independent effects of either Trimethoprim or
EE2 where detected. Significance levels: *** *p* < 0.001; ** *p* < 0.01; * *p* < 0.05.

421 Figure 3. Effects of TMP (💊) and EE2 (♀) on the A) TBI stabilization factor and B)
422 initial decomposition rate, calculated following Keuskamp et al (2013). Inserts show
423 pooled data where significant independent effects of either Trimethoprim or EE2
424 where detected. Significance levels: *** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$.

426