**Protein Interactome of the Cancerous Inhibitor of Protein Phosphatase 2A (CIP2A) in Th17 Cells**

**Mohd Moin Khan [1,2#], Tommi Välikangas [1,3#], Meraj Hasan Khan[1#], Robert Moulder[1], Ubaid Ullah[1], Santosh Dilip Bhosale [1], Elina Komsi[1], Umar Butt[1], Xi Qiao[1], Jukka Westermarck [1,4,] Laura L Elo [1] and Riitta Lahesmaa[1*]**

[1]Turku Bioscience Centre, University of Turku and Åbo Akademi University, FI-20520 Turku, Finland.

[2]Turku Doctoral Programme of Molecular Medicine (TuDMM), Medical Faculty, University of Turku, Turku, Finland.

[3] Doctoral Programme in Mathematics and Computer Sciences (MATTI), University of Turku, Turku, Finland

[4] Institute of Biomedicine, University of Turku, Turku, Finland

[#] These authors contributed equally to this work

**\*Correspondence**

Professor Riitta Lahesmaa

Turku Bioscience Centre,

Tykistökatu 6A (7th floor)

Turku 20520, FINLAND

Phone: 00358-29-4502415

Email: rilahes@utu.fi

**Running Title: Insights of CIP2A Protein Interactome**

**ABSTRACT**

Cancerous inhibitor of protein phosphatase 2A (CIP2A) is involved in immune

response, cancer progression, and in Alzheimer´s disease. However, an

understanding of the mechanistic basis of its function in this wide spectrum of

physiological and pathological processes is limited due to its poorly characterized

interaction networks. Here we present the first systematic characterization of the

CIP2A interactome by affinity-purification mass spectrometry combined with validation

by selected reaction monitoring targeted mass spectrometry (SRM-MS) analysis in

Th17 cells. In addition to the known regulatory subunit of protein phosphatase PP2A,

the catalytic subunit of protein PP2A was found to be interacting with CIP2A.

Furthermore, the regulatory (PPP1R18, and PPP1R12A) and catalytic (PPP1CA)

subunits of phosphatase PP1 were identified among the top novel CIP2A interactors.

Evaluation of the ontologies associated with the proteins in this interactome revealed

that they were linked with RNA metabolic processing and splicing, protein traffic,

cytoskeleton regulation and ubiquitin-mediated protein degradation processes. Taken

together, this network of protein-protein interactions will be important for

understanding and further exploring the biological processes and mechanisms

regulated by CIP2A both in physiological and pathological conditions.

**Key words:** CIP2A, Interactome, Mass Spectrometry (MS), Selected Reaction

Monitoring (SRM) targeted mass spectrometry, confocal microscopy, Pathway

analysis.

**Highlights:**

▪ The first characterisation of the CIP2A interactome in Th17 cells.

- Key interactions were validated by targeted SRM-MS proteomics, western blot and confocal microscopy.
- Pathway analysis of the interactome revealed interrelationships with proteins across a broad range of processes, in particular associated with mRNA processing.

## INTRODUCTION

PP2A is a recognized tumor suppressor that imparts crucial functions in regulating cell proliferation, differentiation, and apoptosis [1] – [7] . Accordingly, down regulation of PP2A has been observed as a feature in many cancers. Cancerous Inhibitor of PP2A (CIP2A) was first characterized as an endogenous inhibitor of protein phosphatase 2A (PP2A) in cancer cells [8] . Previous studies have shown that many of the known actions of CIP2A are mediated through inhibition of PP2A [9] , [10] . Overexpression of CIP2A has been associated with the poor prognosis in several human malignancies, and CIP2A has thus been seen as a potential therapeutic target for cancer therapy [8] , [11] , [12] . CIP2A supports the activity of many critical onco-proteins (Akt, MYC, E2F1) and promotes malignancy in most cancer types via PP2A inhibition [8] . Structurally, CIP2A forms a homodimer and this dimerization promotes its interaction with the PP2A regulatory subunits B56$\alpha$ and B56$\gamma$ [13] . Interestingly, inhibition of either CIP2A dimerization or B56$\alpha$/$\gamma$ expression destabilizes CIP2A in cancer cells [13] . In Alzheimer's disease (AD), enhanced expression of CIP2A in AD brains and neurons results in PP2A inhibition and tau protein hyperphosphorylation, which leads to abnormal tau localization with memory deficits and other pathological conditions [14] .

Th17 cells are found at mucosal surfaces where they regulate tissue homeostasis [15] – [18] . Dysregulation of Th17 cell differentiation, however, has been associated with a number of autoimmune inflammatory diseases [19] , [20] . Following our earlier report on the involvement of CIP2A in T cell activation [21] , we subsequently observed that CIP2A negatively regulates T helper (Th)17 cell differentiation, whereby CIP2A depletion resulted in enhanced IL17 expression both in human and mouse

Th17 cells (Khan M.M et al., submitted).Thus, CIP2A could have important implications

in the differentiation and regulation of Th17 cells. Although CIP2A has been shown to

regulate a range of pathological and physiological conditions, there has not been any

systematic analysis of the full range of signaling pathways and processes regulated

by CIP2A.

In the current study, we used immunoprecipitation followed by label free quantitative

proteomics to gain insight into the proteins that are associated with CIP2A. We present

the first interactome of CIP2A, with validation of interactors by selected reaction

monitoring (SRM) targeted mass spectrometry as well as by Western blotting and

confocal microscopy. In addition to its known interaction with the regulatory subunit A

and B of phosphatase PP2A, our data demonstrated CIP2A interaction with the PP2A

catalytic subunits namely PPP2CA and PPP2CB. Furthermore, both the regulatory

(PPP1R18, and PPP1R12A) and catalytic (PPP1CA) subunits of phosphatase PP1

were identified among the top interactors of CIP2A. This data further implicates the

involvement of CIP2A in a number of biological processes that it has previously not

been associated to, and provides a basis for comprehensive understanding of CIP2A

for translational studies targeting CIP2A in human diseases.

## RESULTS

### Mass spectrometry analysis identify novel CIP2A binding partners

To identify the proteins interacting with CIP2A and thereby facilitate the understanding of its functions in human Th17 cells, immunoprecipitation (IP) was performed after 72 h of polarization of naïve T cells towards the Th17 lineage **(Fig. 1A; S1A-C).** The cell model for the study was rationalized by our recent findings indicating that CIP2A is involved in T-cell activation [21] and its particular importance for Th17 cell differentiation (Khan MM et al., submitted). Two CIP2A specific antibodies (Ab1 and Ab2) recognizing distinct regions of CIP2A were used in two biological replicates, together with their respective IgGs (IgG1 and IgG2) as control antibodies. Ab1 was a monoclonal antibody targeting the N-terminal structured region of the protein (residues surrounding Val 342) [13] , while Ab2 was of polyclonal origin targeting unstructured C-terminus [22] (Fig. 1B).

The availability of highly specific antibodies against CIP2A facilitated the IP of endogenous CIP2A without the requirement for overexpression. After preparation for proteomic analysis, the digested samples were analyzed by LC-MS/MS to identify the CIP2A associated proteins. Analysis of the data revealed 680 proteins meeting the criteria of identification of two or more unique peptides with a confidence threshold of 99%. In order to distinguish non-specific interactions and common contaminants, statistical analysis of the LFQ intensity data was made using the Significance Analysis of *INTeractome* (*SAINT*) algorithm [23] . Using a threshold for the calculated SAINT probability scores of greater than or equal to 0.95, and exclusion of proteins represented in the contaminant's repository [24] with a frequency of >60%, the list of candidate interacting proteins was reduced to 331 (**Supplementary Table 1A-C)** with

more than 90% overlap in common proteins identified with both antibodies specific to CIP2A **(Fig. 1C; S1D; Supplementary Table 1A-C)**. To exemplify the interactions between CIP2A and its interaction partners, the top fifty (50) common proteins detected with both antibodies based on the detected signal intensity is plotted in the form of heatmap **(Fig. 1D)**.

**Distribution of CIP2A interactome between the nucleus and cytoplasm**

Analysis of the cellular location of CIP2A in epithelial cells has indicated that it is mainly expressed in the cytoplasm [8] , [22] . Further data indicate some representation in the nuclear fraction of epithelial cancer cells [25] . To evaluate the distribution of CIP2A in Th17 cells, confocal microscopy measurements were conducted in this study. For these measurements, 4,6-diamidino-2-phenylindole (DAPI), and Phalloidin were used to stain the nuclear and cytoplasmic fractions, respectively (**Fig 2A**). Statistical analysis of the confocal microscopy images revealed CIP2A is predominantly located in the cytoplasmic compartment of Th17 cells, in addition to some nuclear expression, (**Fig 2B**). Confirmatory measurements were provided with Western Blot (WB) analysis of the cytoplasmic and nuclear fractionations of 72h polarised Th17 cells. For these experiments, vimentin and tubulin were used as control markers for the nuclear and cytoplasmic fractions, respectively (**Fig 2C; S2A**). Whilst the observed expression of CIP2A was greater in the cytoplasmic region, CIP2A was also detected in the nuclear region of 72h differentiated Th17 cells (**Fig 2C; S2A**).

Ingenuity Pathways Analysis (IPA; Qiagen) was used to map cellular locations of the proteins interacting with CIP2A. The analysis revealed that the cytoplasm (42%) and nucleus (43%) constituted the most frequent cellular locations of the associated proteins, whilst the plasma membrane (8%), extracellular space (4%), and undefined

162 (other, 3%) accounted for the remainder (**Fig 2D**). The observed distribution of the

163 proteins detected in the CIP2A interactome is consistent with its association in these

164 compartments in Th17 cells.

**Cellular processes regulated by CIP2A interactome**

To gain an overview of the biological processes associated with CIP2A, we analyzed

the CIP2A interactome data with several Gene Ontology and interaction bioinformatics

tools. To collate the known interactions between the proteins, a network was

constructed using the STRING database [26] . The resulting network was visualized

with Cytoscape [27] revealing "RNA metabolism or splicing" as the process most

frequently linked with the associated proteins (**Fig. 3A; Supplementary Table 2 and**

**3**). The top 20 proteins associated with RNA metabolic process, RNA splicing via

transesterification and RNA splicing via spliceosome are represented in the form of

heatmaps (**Fig. S3A-C**). In agreement with these results, a recent phosphoproteome

screen also identified RNA splicing as one of the most significantly CIP2A-regulated

processes [28] .

To provide additional context and an appropriate milieu, functional enrichment analysis

was made relative to a combined list of the proteins detected together with those found

in a recent cellular proteomics analysis of human Th17 cells [29] rather than relative

to a background of the whole genome. For this comparison the bioinformatics tools

DAVID [30] , [31] and PANTHER [32] , [33] were used. The enriched Th17 pathways

are represented in **Fig. 3B**. Again, mRNA processing and splicing were found to be

the most significant biological process linked with CIP2A interactors. Collectively,

these results suggest the involvement of CIP2A in multiple processes.

IPA analysis was also used to summarize the molecular functions of the CIP2A

interacting proteins (**Fig. 3C**). This revealed enzymes (~24%) as the major functional

class of the proteins. In addition, transcription regulators (~7.4%) and transporters

(6.1%) are the next major class of proteins. A small fraction of proteins was classified

as phosphatases (3.4%), kinases (2.0%), translation regulators and ion channels

(1.0% each), cytokines and peptidases (<1% each). For a major fraction of the proteins

in CIP2A interactome (54%), however, analysis with the IPA database indicated that

no specific function could be assigned (marked as "others" in **Fig 3C**). Overall the IPA

analysis indicated that CIP2A interacts with proteins with a variety of different

functions. It is possible that via its interaction with these proteins CIP2A regulates a

range of processes in different cell types, some of which are as of yet undefined.

Thus, the functional and network analysis of the associated proteins revealed how

CIP2A and its interacting partners participate in a broad set of essential cellular

processes, ranging from RNA metabolic processes and splicing to viral processes.

**Validation of the Th17 cell CIP2A protein interactions**

Validation experiments were performed to verify the interaction of selected proteins

with CIP2A. The choice of targets focused on those with most significant interactions

and nodal positions in CIP2A interactome (**Fig. 1D).** The validation experiments were

carried out by confocal microscopy, WB, or selected reaction monitoring (SRM)

targeted mass spectrometry (MS) analysis. The choice of method depended on the

availability of suitable working antibodies or proteotypic peptide signatures, for

confocal microscopy and WB or SRM-MS analysis, respectively.

SRM-MS analysis provided successful validation of twenty CIP2A interaction partners using immunoprecipitates from three independent biological replicates (**Fig 4A; Supplementary Table 4-5**). The results are summarized in **Supplementary Table 5.** We confirmed the known interaction of CIP2A with PP2A regulatory subunit A (PPP2R1A) **(Fig. 4A).** In addition, we identified and validated among the CIP2A interactors, PP2A catalytic subunits PPP2CA and PPP2CB **(Fig. 4A; S4A and Supplementary Table 5).**

Interestingly, we identified PP1 regulatory (PPP1R18 and PPP1R12A) and catalytic subunits (PPP1CA) as the top CIP2A interacting proteins (**Fig 1D**). By SRM-MS analysis, further confirmed CIP2A interaction with the catalytic subunits (PPP1CA and PPP1CB) of phosphatase PP1 **(Fig. 4A; S4A and Supplementary Table 5)**. Further, in a recent phospho-proteome analysis of CIP2A-regulated proteins, LMNB1 was found to be dephosphorylated in CIP2A silenced cells [28] and here its interaction was identified and validated in CIP2A interactome by SRM MS analysis **(Fig. 4A).** Previous studies have revealed the interaction of with phosphatase PP2A regulatory A subunit [8] . Accordingly, we observed and validated this interaction with PP2A-A in Th17 cells (**Fig 5A).** Additionally, we validated the interaction with the PP2A catalytic subunit PPP2ACA. Notably, we were also able to validate interaction between CIP2A and PP1 catalytic subunit by co-immunoprecipitation followed by WB **(Fig. 5B).** Interaction of CIP2A with both PPP2ACA, and PPP1CA could be because of high structural similarity of the two proteins [34] . Further, PPP1CA interaction was independently validated by Proximity ligation assay (PLA) **(Fig. 5C)**. The negative control, PLA conducted from same samples without primary antibodies, did not show background signals (**Fig. 5C**).

TRIM21 was amongst the most enriched proteins in this interactome, and the

interaction was validated by both SRM-MS and PLA confocal microscopy **(Fig 5D).**

Interaction with the ubiquitin ligase UBR5 was validated by PLA confocal microscopy,

both in Th17 cells and Hela cells (**Fig 5D; Fig S5A**). In addition, WB analysis were

made for STAT1, IRF4 and TRIM21, which were consistent with the targeted SRM-

MS analysis (**Fig 4A; Fig 5E; Fig S5B-C**). Further, NFKB2 was also confirmed to be

a CIP2A interacting protein (**Fig 4A, Fig S4A**) and CIP2A inhibition decreased NF-KB

activation (**Fig 5F**). In CIP2A silenced Th17 cells, reduced phosphorylation of P65 was

detected by WB analysis (**Fig 5F).**

**DISCUSSION**

With these analyses of the CIP2A interactome in the Th17 milieu, and in relation to

phosphatase biology, we verified the well-known interaction with PP2A-A (PPP2R1A)

and PP2A regulatory B subunit **(Supplementary Table 1A-C)**. Previously, Wang et

al. (2017) demonstrated direct interaction between CIP2A and PP2A regulatory B

subunits B56$\alpha$ and B56$\gamma$ by yeast two-hybrid analysis (Y2H), and Junttila et al. (2007)

discovered PP2A-A subunit interaction with CIP2A.  In the present report, we detected

and validated for the first time the interaction of CIP2A with the PP2A catalytic subunit

(PPPCA and PPPCB). All of these interactions were validated with SRM-MS analysis

**(Fig. 4A; and Supplementary Table 4-5).**

The regulatory (PPP1R18 and PPP1R12A) and catalytic (PPP1CA) subunits of

phosphatase PP1 were among the top CIP2A interacting partners. Interaction between

CIP2A and PP1 has not been previously shown and further studies are required to

demonstrate the importance of PP1 association with CIP2A. Notably, the catalytic

subunits of both phosphatases PP2A and PP1 were detected and confirmed in the

CIP2A interactome. This interaction could arise due to the structural similarity between

catalytic subunits of phosphatases PP2A and PP1. It will be an interesting topic for

future studies to determine whether CIP2A engages other catalytic subunits of

serine/threonine phosphatases.

Previously in studies with T regulatory (Treg) cells, PP2A enhanced catalytic activity

was identified as an indispensable element of the suppressive function of Treg cells

[1] . Treg cells maintain tolerance and prevent autoimmune diseases by suppressing

effector cells and modulating the immune system. It remains to be studied whether the

interaction between CIP2A and catalytic subunit of PP2A regulates the suppressive

function of Treg cells.

In relation to CIP2A and  PP1 in autoimmune disease,  in rheumatoid arthritis (RA)

inflamed synovium, increased expression and activity of phosphatase PP1 and

reduced transcriptional activity of FOXP3 through specific dephosphorylation of the

Ser418 site has been observed  [35] . Intriguingly, CIP2A expression in RA has been

associated with the histopathological score of synovitis as well as apoptotic resistance

and  invasive  function  of  fibroblast-like  synoviocytes   [36] ,  [37] .  Thus,  interaction

between CIP2A and PP1 may play a role in regulatory T cell function and inflammation

in RA.

Several kinases related to T cell receptor (TCR) signaling (i.e. CSK and LCK)  [38] ,

[39]  were identified as CIP2A interactors. The initiation of TCR signalling is regulated

by the SRC-related protein tyrosine kinase family, which includes LCK, whereas CSK

is a potent inhibitor of TCR signalling  [38] . Notably, we have observed reduced T cell

activation in CIP2A depleted T cells [21] . It will be interesting to study if the loss of

CIP2A binding to LCK or CSK contributes to impaired T cell activation

The adaptor protein DOCK8 was amongst the most highly enriched proteins detected

in this interactome. Combined immunodeficiency in humans is associated with loss-

of-function mutations in the gene encoding DOCK8 [40] . Notably, the phenotype in

immunodeficiency due to mutations in the gene encoding the transcription factor

STAT3 is very similar to that caused by the loss of DOCK8 [41] . Jabara et al. reported

that in B cells DOCK8 and STAT3 are components of the same signalling cascade

[42] . In Th17 cells, CIP2A interacts with both DOCK8 and Lyn, and regulates STAT3

phosphorylation (Khan MM et al., submitted). It could be revealing to study whether

CIP2A regulation of STAT3 phosphorylation is dependent on its interaction with Lyn

and DOCK8.

Several of the proteins of this interactome have functions related to viral response and

interferon signalling, i.e. IFI16, IRF4, MX2, STAT1 and TRIM21.  Notably amongst

these, the increased activity and DNA binding of IRF4 has previously been found in

association with the increased capacity of PP2Ac transgenic T cells to produce IL-17

in response to TCR stimulation [2] . CIP2A directly interacts and inhibits PP2A in

cancer cells and depletion of CIP2A in T cells results in enhanced IL17 (Khan MM et

al., submitted manuscript). On the basis of these associations, the influence of CIP2A

interaction on the DNA binding of IRF4 in T cells could be an important focus of future

studies.

Amongst the CIP2A interacting transcription factors were RUNX1 and STAT1, which

were validated by WB and SRM. Future studies could address the relationship

between CIP2A and STAT1 in Th cells. RUNX1 is important for the development of

Th17 cells as it directly binds the promoter of RORC and mediates transactivation of

RORC, a transcription factor important for Th17 cell development [43] . In addition,

Runx1 has been reported to bind to the IFNγ locus and induce IFNγ in Th17 cells

resulting in development of pathogenic IL17[+] IFNγ[+] Th17 Cells [44] . In view of the

latter and regarding the detected interaction between CIP2A and RUNX1, it will be of

interest to find out if this interaction plays a role in RUNX1 mediated RORC and IFNγ

induction in Th17 cells [43] , [44] .

TRIM21, an E3 ubiquitin ligase, was prominent in the CIP2A interactome. TRIM21 has

been associated with the pathogenesis of several autoimmune diseases, including

systemic lupus erythematosus (SLE) and inflammatory bowel diseases (IBDs).

TRIM21 plays a protective role in IBD pathogenesis, possibly through inhibition of Th1

and Th17 cell immune responses in the intestinal mucosa [45] – [47] . In view of the

detected interaction, it is possible that pathways involving CIP2A are also active in the

pathogenesis of autoimmune diseases like SLE and IBD.

Beta-spectrin (SPTBN1) was amongst the top most proteins enriched in CIP2A

interactome. Beta spectrin is a molecular scaffold protein that is involved in cell shape

and the organisation of organelles in the cell [48] . Beta spectrin has been identified

as an integral component of beta-amyloid plaques in Alzheimer's disease (AD) and

spectrin breakdown products (SBDPs) have been implicated as potential biomarkers

for neurodegenerative diseases including AD [49] , [50] . In a recent study of AD,

CIP2A was found to be overexpressed and associated with tau hyperphosphorylation,

mis-localization and AD-related cellular pathology [14] . In mice, overexpression of

CIP2A in the brain induced AD-like cognitive deficits [14] . Future studies may

determine whether CIP2A interaction with beta-spectrin modulates protein

aggregation in AD pathology.

In summary, the results of this study present the first characterisation of CIP2A

interactome. In addition to the known interactions with the PP2A protein A and B

regulatory subunits, our results detail several novel CIP2A interacting proteins. These

include the catalytic subunit of PP2A, as well as regulatory and catalytic subunits of

PP1. Gene ontology and pathway analysis of the associated proteins suggest CIP2A

to be involved in several cellular functions that it has not been previously linked to.

Taken together, we anticipate the new information of proteins depicted in this CIP2A

interactome and their associated networks, could be useful in understanding the role

of CIP2A in regulating cell signalling and its role in various human disorders.

**Methods**

**Human CD4+ T cell isolation, activation, and differentiation**

Umbilical cord blood was layered on Ficol (GE health Care # 17-1440-03) for isolation

of white blood cells. CD4+ T cells were then isolated using the Dynal bead CD4+

isolation kit from Invitrogen (Cat #11331D). For activation of T cells, a combination of

plate-bound anti-CD3 (Beckman Coulter REF # IM-1304) and soluble anti-CD28

(Beckman coulter REF # IM1376) antibodies was used. The Th17 cultured cells were

stimulated in presence of IL-6 (20ng/ml, Roche, Cat# 11138600 001); IL-1β (10ng/ml,

R&D Systems Cat# 201 LB); TGF-β1 (10ng/ml, R&D Systems Cat# 240); anti-IL-4 (1

ug/ml) R&D Systems Cat# MAB204); anti-IFN-γ (1 µg/ml R&D Systems Cat#MAB-

285).

**Flow cytometry**

For Th17 and activated T cells, CCR6 surface staining was performed for Th17 cell culture polarization control. Cells were stained with the antibody (eBioscience) in FACS I buffer (1%FBS in PBS) at +4 °C in darkness for 30min. Cells were then washed two times with FACS I, and finally re-suspended in FACS I Buffer or 1% Formalin before acquisition in BD LSR II/ Caliber. Data analysis was performed using FlowJo (BD) or Flowing Software (Version 2.5.1).

**Luminex assay**

As a polarization control, IL17 cytokine protein secretion was detected in 72h polarized human Th17 cells and TCR activated control cells. Luminex assay (Milliplex MAP human cytokine, Luminex 200 by Luminex xMAP technology) was used to detect the IL17 cytokine in the supernatant and measured according to the manufacturer's instructions. To avoid cell proliferation bias, the cytokine concentrations were normalized by the number of cells from each respective culture, as counted by flow cytometry.

**Immunoprecipitation**

CIP2A immunoprecipitation (IP) was carried out with two specific anti-CIP2A antibodies that recognize distinct regions of CIP2A protein and the respective Immunoglobulin G (IgG) were used in control IPs. The Pierce™ MS-Compatible Magnetic IP Kit, (Catalog number: 90409) was used to perform IP. Briefly, cells were first harvested on ice and the culture medium removed. This was followed by washing the cells twice with PBS. Cells were then re-suspended on ice-cold IP-MS Cell Lysis Buffer, as recommended by the manufacturer, and incubated on ice for 10 minutes with periodic mixing. The lysate was centrifuged at ~13,000 × g for 10 minutes to pellet

the cell debris. The supernatant was transferred to a new tube for protein

concentration determination using a NanoDrop™ spectrophotometer (Thermo

Scientific). Cell lysates (500μg) were incubated with the IP antibody (1:50 dilution).

The antibody/lysate solution was diluted with the manufacturer's proprietary IP-MS cell

lysis buffer and incubated with mixing overnight at +4°C to form the immune complex.

On the next day the immune complex was incubated with Protein A/G magnetic beads

for 1h at room temperature. The beads were then washed and the target antigen eluted

with the elution buffer provided and transferred to a new low protein-binding 1.5mL

microcentrifuge tube. A vacuum centrifuge was used to dry the eluate prior to

preparation for MS or SDS-PAGE and Western blot.

**Western Blotting**

T cells were first resuspended in Triton-X-100 lysis buffer (TXLB). The composition of

TXLB used was 50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.5% Triton-X-100, 5%

Glycerol, 1% SDS. For cell lysis, T cells in TXLB were then sonicated in ice for 5

minutes using a Bioruptor® sonicator. After cell lysis, the lysate was centrifuged at

high speed (16,000 g) and the supernatant was transferred to a new tube to remove

cell debris. Protein concentration was estimated using a DC Protein assay (Bio-Rad

Cat#500-0111). Equal amounts of protein were loaded onto the acrylamide gel (Bio-

Rad Mini or Midi PROTEAN® TGX precast gels). For the transfer of proteins to the

PVDF membrane, mini or a midi transfer packs from Bio-Rad were used depending

on the gel size. Primary/ secondary antibody incubations were performed in 5% non-

fat milk or Bovine serum albumin (BSA) for phosphoproteins in TBST buffer

(0.1%Tween 20 in Tris-buffered saline). Quantification of the bands detected was

performed using Image J software after normalization relative to the loading control

intensities. The following antibodies were used for western blotting, CIP2A antibody

1(Ab1) (Cell Signaling Technology; cat#14805) CIP2A antibody 2 (Ab2) provided by

Prof. Jukka Westermarck  [8] , [22] ; PP2A-A/B (sc-15355; Santa Cruz); Trim21

antibody (Santa Cruz); Irf4 antibody (Santa Cruz); Beta-actin antibody (Sigma Cat #.

A5441). PPP1A/PPP1CA antibody (ab137512; abcam), PP1α Antibody (G-4): (sc-

271762; Santa Cruz).

**Immunofluorescence**

Th17 cells were plated onto acid-washed glass coverslips. The cells were then fixed

at room temperature (RT) using 4 % PFA for 10 mins. After fixing, the cells were

permeabilized in PBS containing 30% horse serum and 0.1% Triton X-100 for 10 min

at RT, washed in PBS and then blocked in PBS-30% horse serum for 1h at RT. The

cells were then incubated with the CIP2A primary antibody overnight at 4°C. On the

next day, the cells were washed three times using PBS and incubated for 1h at RT

with AlexaFluor-conjugated secondary antibodies (Life Technologies). Cells were then

mounted with Mowiol® for 30 min at 37°C and Images taken with a Carl Zeiss LSM780

laser scanning confocal microscope equipped with 100×1.40 oil plan-Apochromat

objective (Zeiss). Corrected total cell fluorescence was calculated from confocal

images to quantify the intensities in Th0 and Th17 cells. [51] , [52] .

Immunofluorescence staining of UBR5 in Hela cells was performed similarly, the cells

were cultured on chambered coverslip (80826, Ibidi). 4% paraformaldehyde was used

to fix the cells for 15 minutes under room temperature, and then cells were

permeabilized with 0.5% Triton X-100 in PBS on ice for 5 minutes. Next, the cells were

blocked with 10% normal goat serum (ab7481, Abcam) diluted in PBS for 30 minutes,

and followed by incubating with the primary antibodies anti-UBR5 (ab70311, Abcam)

overnight at 4°C. Subsequently, cells were washed with PBS and incubated with the

secondary antibody, Alex Fluor 488 goat anti-rabbit IgG (A-11008, Invitrogen) for 1h

under room temperature. After secondary antibody incubation, the cells were washed

with PBS and nuclei were stained with DAPI (D1306, Invitrogen) in PBS at RT for 10

min. Images were acquired with confocal microscope (LSM780, Carl Zeiss).

**Proximity ligation assay**

The PLA assay was performed according to the manufacturer's protocol (Duolink®

PLA, Sigma). Briefly, Th17 or HeLa cells were plated on coverslips. The cells were

fixed with 4% paraformaldehyde for 15 minutes at room temperature, and then

permeabilized with 0.5% Triton X-100 in PBS on ice for 5 minutes. Next, cells were

blocked with blocking solution, and incubated in a pre-heated humidity chamber for 30

min at 37°C, followed by incubating with the primary antibodies (in blocking solution)

anti-UBR5 (ab70311, Abcam) and anti-CIP2A (sc-80659, Santa Cruz), anti-PPP1CA

(ab137512, Abcam), anti-UBR5 (ab70311, Abcam), anti-TRIM21 (SC25351, Santa

Cruz), anti-GFP (Mouse, ab1218, Abcam ) and anti-GFP (Rabbit, A11122, Invitrogen)

overnight at 4°C. Subsequently, cells were washed with the kit buffer A, and the PLA

probe was incubated in a pre-heated humidity chamber for 1h at 37°C, followed by

ligase reaction in a pre-heated humidity chamber for 30 minutes at 37°C. Next,

amplification polymerase solution for PLA was added, followed by incubating the cells

in a pre-heated humidity chamber for 100 minutes at 37°C. After amplification, the

coverslips were washed with the kit buffer B, and mounted with DAPI. The PLA signal

was detected by using a confocal microscope (LSM780, Carl Zeiss) ) and 3i CSU-W1

spinning disk microscope equipped with 100x1.4 O Plan-apochromat objective

(Zeiss). PLA signals per cell were calculated by dividing the amount of PLA signal dots

in one field of view, determined using Cell Profiler software  [53]

**Cell transfection with siRNA**

Naïve T cells were isolated form the cord blood and four million cells were transfected

in 100 µl volume of Opti-MEM™ (Gibco by Life Technologies, cat # 31985-047). Lonza

nucleofector was used for siRNA transfections. After nucleofection cells were rested

for 48h in RPMI supplemented with 10% serum.

The siRNAs (synthesized by Sigma) are shown below:

siNT 5´- AAUUCUCCGAACGUGUCACGU -3´ 22;

siCIP2A-1 5´-CUGUGGUUGUGUUUGCACUTT-3´1.

**LC-MS/MS analysis of the proteomic profiles of the immuno-precipitates**

The immuno-precipitates were denatured with 8 M urea and cysteine residues reduced

with dithiothreitol then alkylated with iodoacetamide. After digestion with trypsin (37

$^0$C, 16 hours), the samples were desalted using Empore $C_{18}$ disks (3M, Cat No 2215).

Based on using UV absorption measurements with a NanoDrop-1000 UV

spectrophotometer (Thermo Scientific), equivalent aliquots of the digested peptides

were analyzed in triplicate with an Easy-nLC 1200 coupled to a Q Exactive HF mass

spectrometer (Thermo Fisher Scientific). A 20 x 0.1 mm i.d.  pre-column was used in

conjunction with a 150 mm x 75 $\mu$m i.d. analytical column, both packed with 5 $\mu$m

Reprosil $C_{18}$-bonded silica (Dr Maisch GmbH). A separation gradient from 8 to 45% B

in 78 min was used at a flow rate of 300 nl/min. The mobile phase compositions were,

water with 0.1 % formic acid (A) and 80% acetonitrile 0.1% formic acid (B). The MS/MS

data were acquired in positive ion mode with a data dependent acquisition setting for

higher-energy C-trap dissociation (HCD) of the 10 most intense ions (m/z 300-2000,

charge states > 1+). Further details of the preparation and instrument parameters are

available as supplementary information.

The LC-MS/MS discovery data are available via the PRIDE  [54]   partner repository

from the ProteomeXchange Consortium with the dataset identifier PXD008983.

**LC-MS/MS validation of selected targets**

Selected reaction monitoring (SRM) mass spectrometry was used to validate the

relative abundance of twenty of the proteins that were found to be associated with the

CIP2A immuno- precipitates (**Supplementary Table 4**). Heavy-labeled synthetic

peptides (lysine $^{13}C_6$ $^{15}N_2$ and arginine $^{13}C_6$ $^{15}N_4$) analogues for the targets of interest

were selected on the basis of their consistency and intensity in the discovery data and

obtained from Thermo Scientific.  For these validations, three additional cultures were

prepared from the cord blood of three donors.

Using Skyline software [55]  the top five most intense transitions for each peptide were

evaluated from the MS/MS spectra of heavy labelled peptides and the relative

performance of the native peptides in the spiked validation samples.

The validation samples were prepared using the same digestion and desalting

protocols used for discovery. These were then spiked with synthetic heavy labelled

analogues of the peptide targets and a retention time standard (MSRT1, Sigma) for

scheduled selected reaction monitoring.  The LC-MS/MS analyses were conducted

using Easy-nLC 1000 liquid chromatograph (Thermo Scientific) coupled to a TSQ

Vantage Triple Quadrupole Mass Spectrometer (Thermo Scientific).  The column

configuration was the same as used in the discovery experiments and the peptides

were separated using a gradient of 8% to 43% B in 27 min, then to 100% B in 3 min,

at a flow rate of 300 nl/min (the mobile phase compositions are as indicate above).

The raw SRM data are available through PASSEL  [56]  with the dataset identifier

PASS01186

**Data analysis and statistics**

The peptide/protein identification was performed in Andromeda search engine  [57] ,

incorporated in MaxQuant software  [58] , with a UniProt human protein sequence

database (version June 2016, 20,237 entries). Trypsin digestion with a maximum of

two missed cleavages, carbamidomethylation of cysteine as a fixed modification, and

methionine oxidation and N-terminal acetylation as variable modifications were used

in the search parameters. A false discovery rate (FDR) of 0.01 at the peptide and

protein level was applied. MaxQuant's label-free quantification (LFQ) algorithm  [59]

was used to calculate the relative protein intensity profiles across the samples. The

"match between the runs" option was enabled with a time window of 0.7 min.

The MaxQuant output was further processed using Perseus  [60] . The downstream

processing included the removal of contaminants, reverse hits and proteins with less

than three valid replicate values in the sample groups (IgG or CIP2A pulldown). To

identify the potential interacting proteins, Significance Analysis of *INTeractome*

(*SAINT*) analysis was performed [24] using the Resource for Evaluation of Protein

Interaction Networks (REPRINT) interface (https://reprint-apms.org/). This algorithm

uses the distribution, relative intensity and variation of the prey protein detection to

calculate probability scores for the certainty of their interaction with the bait.

Additionally, the protein lists were compared against a list of common contaminants

identified in a catalogue of immunoprecipitation experiments (CRAPome). The

thresholds applied to define the putatively interacting proteins included identification

on the basis of two or more unique peptides and a SAINT probability score of $\geq 0.95$.

Although there was no direct match for our sample type and IP conditions in the

contaminants repository database, additional filtering was made to exclude proteins

that were more than 60% of the datasets. Skyline was used to design the SRM

experiments and process the data generated. The MS Stats (3.8.4) plugin included in

the Skyline software was used for the group comparison between cases and controls

[61] .

**Network analysis**

Previously reported protein-protein interactions for the detected CIP2A interacting

proteins were downloaded from the STRING functional protein association networks

database version 10.5 [26] . Both predicted and known high confidence (interaction

score $\geq 0.7$) interactions were considered. The resulting network was visualized with

Cytoscape version 3.6.0 [27] . Clusters in the network were identified using the Markov

clustering algorithm implemented in the Cytoscape plug-in clusterMaker v2 algorithm

MCL Cluster [62] with the granularity parameter (inflation value) set to 1.8 as

suggested by [63] . The MS intensity differences between CIP2A immuno-precipitates

and the IgG controls were calculated as median differences across all replicates in the

normalized log2-transformed intensities (color coded as a continuous gradient from

white to grey as node inner color in Figure 3A).

**Enrichment analysis**

The enriched GO biological processes were identified using the Database for

Annotation, Visualization and Integrated Discovery (DAVID) version 6.8 [30] , [31]

and the Protein Annotation Through Evolutionary Relationship (PANTHER)

classification system version 13.0 [32], [33]. The enrichment analyses were performed using the detected CIP2A interactome as the input and the whole detected Th17 proteome from a Th17 profiling study [29] as the background reference. The GO FAT terms from DAVID, which filter out the broadest, non-specific terms, were considered for calculating the most enriched processes within each identified cluster with more than four members. The GO SLIM terms from PANTHER, which give only the broadest terms and filter the more specific ones, were considered for a summary of the enriched biological processes in the CIP2A interactome. A biological process was considered enriched if it had FDR ≤ 0.05. To summarize the cellular locations and protein classes the protein list was annotated using Ingenuity Pathway Analysis (Qiagen).

## ACKNOWLEDGEMENTS

For the cord blood collections, we are thankful to the staff of Turku University Hospital, Maternity Ward, Department of Obstetrics and Gynecology. Marjo Hakkarainen and Sarita Heinonen are acknowledge for excellent technical help. We are thankful to the core facilities at the Turku Bioscience centre namely, the Proteomics facility and Cell Imaging Core (CIC) facility. Biocenter Finland are acknowledged for their support of the latter facilities. The Finnish Centre for Scientific Computing (CSC) is duly acknowledged for their efficient servers and their resources in data analysis. MMK was supported by University of Turku graduate school on Turku Doctoral Programme of Molecular Medicine (TuDMM) as well as a central grant from Finnish Cultural Foundation and T.V. was supported by the Doctoral Programme in Mathematics and Computer Sciences (MATTI) of University of Turku. R.L. was supported by the Academy of Finland, AoF, Centre of Excellence in Molecular Systems Immunology and Physiology

Research (2012-2017) grant 250114; by the AoF grants 292335, 294337, 292482,

31444, by grants from the JDRF, the Sigrid Jusélius Foundation and the Finnish

Cancer Foundation. L.L.E. has received grants from the European Research Council

(ERC; grant 677943), the European Union's Horizon 2020 research and innovation

program (grant 675395), the AoF (grants 296801 and 304995), the Juvenile Diabetes

Research Foundation (JDRF; grant 2-2013-32), the Finnish Funding Agency for

Innovation (TEKES; grant 1877/31/2016), and the Sigrid Jusélius Foundation. J.W.

was funded by the Sigrid Jusélius Foundation.

**ETHICAL APPROVAL**

Usage of the blood of unknown donors is approved by the Ethics Committee, Hospital

District of Southwest Finland.

**AUTHOR CONTRIBUTIONS**

M.M.K designed and performed the experiments, analyzed data and prepared figures

related to WB, TaqMan-PCR and Flowcytometry, and wrote the manuscript. T.V.

analyzed data, prepared figures for network analysis and wrote part of the manuscript.

M.H.K. designed and performed the experiments, prepared figures, and wrote part of

the manuscript. R.M performed SRM-proteomics, analyzed the proteomics data, and

contributed to writing the manuscript. U.U. contributed to the pathway and network

analyses and writing of the manuscript. S.D.B performed the proteomics experiments

data analysis. E.K performed experiments, and analyzed the data. JW provided

scientific expertise, and reagents and edited the manuscript. L.E. designed

experiments, supervised the study and edited the manuscript. R.L. designed and

supervised the study and wrote the manuscript.

## DECLARATION OF INTERESTS

The authors declare no competing interests.

## ACCESSION NUMBERS

The mass spectrometry data for the interactome have been deposited with the ProteomeXchange Consortium via the PRIDE [54] partner repository with the dataset identifier PXD008983. The validation data have similarly been submitted to the Peptide Atlas [64] with the identifier PASS01186.

**REFERENCES**

1. Apostolidis SA, Rodríguez-Rodríguez N, Suárez-Fueyo A, Dioufa N, Ozcan E,

Crispín JC, et al. Phosphatase PP2A is requisite for the function of regulatory

T cells. Nat Immunol. 2016;17(5):556–64.

2. Apostolidis SA, Rauen T, Hedrich CM, Tsokos GC, Crispín JC. Protein

phosphatase 2A enables expression of interleukin 17 (IL-17) through

chromatin remodeling. J Biol Chem. 2013;288(37):26775–84.

3. Eitelhuber AC, Warth S, Schimmack G, Düwel M, Hadian K, Demski K, et al.

Dephosphorylation of Carma1 by PP2A negatively regulates T-cell activation.

EMBO J. 2011;30(3):594–605.

4. Janssens V, Goris J. Protein phosphatase 2A: a highly regulated family of

serine/threonine phosphatases implicated in cell growth and signalling.

Biochem J. 2001 Feb 1;353(Pt 3):417–39.

5. Katsiari CG, Kyttaris VC, Juang Y-T, Tsokos GC. Protein phosphatase 2A is a

negative regulator of IL-2 production in patients with systemic lupus

erythematosus. J Clin Invest. 2005 Nov;115(11):3193–204.

6. Long L, Deng Y, Yao F, Guan D, Feng Y, Jiang H, et al. Recruitment of

phosphatase PP2A by RACK1 adaptor protein deactivates transcription factor

IRF3 and limits Type I interferon signaling. Immunity. 2014;40(4):515–29.

7. Wlodarchak N, Xing Y. PP2A as a master regulator of the cell cycle. Crit Rev

Biochem Mol Biol. 2016;51(3):162–84.

8.    Junttila MR, Puustinen P, Niemelä M, Ahola R, Arnold H, Böttzauw T, et al.

CIP2A Inhibits PP2A in Human Malignancies. Cell. 2007;130(1):51–62.

9.    Laine A, Sihto H, Come C, Rosenfeldt MT, Zwolinska A, Niemelä M, et al.

Senescence Sensitivity of Breast Cancer Cells Is Defined by Positive

Feedback Loop between CIP2A and E2F1. Cancer Discov. 2013 Feb

1;3(2):182–97.

10.   Niemelä M, Kauko O, Sihto H, Mpindi J-P, Nicorici D, Pernilä P, et al. CIP2A

signature reveals the MYC dependency of CIP2A-regulated phenotypes and

its clinical association with breast cancer subtypes. Oncogene. 2012 Sep

16;31(39):4266–78.

11.   Khanna A, Pimanda JE, Westermarck J. Cancerous Inhibitor of Protein

Phosphatase 2A, an Emerging Human Oncoprotein and a Potential Cancer

Therapy Target. Cancer Res. 2013 Nov 15;73(22):6548–53.

12.   Lucas CM, Milani M, Butterworth M, Carmell N, Scott LJ, Clark RE, et al. High

CIP2A levels correlate with an antiapoptotic phenotype that can be overcome

by targeting BCL-XL in chronic myeloid leukemia. Leukemia. 2016

Jun;30(6):1273–81.

13.   Wang J, Okkeri J, Pavic K, Wang Z, Kauko O, Halonen T, et al. Oncoprotein

CIP2A is stabilized via interaction with tumor suppressor PP2A/B56. EMBO

Rep. 2017 Mar 7;18(3):437–50.

14.   Shentu Y-P, Huo Y, Feng X-L, Gilbert J, Zhang Q, Liuyang Z-Y, et al. CIP2A

Causes Tau/APP Phosphorylation, Synaptopathy, and Memory Deficits in

Alzheimer's Disease. Cell Rep. 2018 Jul 17;24(3):713–23.

15.   Guglani L, Khader SA. Th17 cytokines in mucosal immunity and inflammation.

Curr Opin HIV AIDS. 2010 Mar;5(2):120–7.

16.   Kleinewietfeld M, Manzel A, Titze J, Kvakan H, Yosef N, Linker RA, et al.

Sodium chloride drives autoimmune disease by the induction of pathogenic

TH17 cells. Nature. 2013;496(7446):518–22.

17.   Kleinewietfeld M, Hafler DA. The plasticity of human Treg and Th17 cells and

its role in autoimmunity. Semin Immunol. 2013 Nov 15;25(4):305–12.

18.   Romani L. Immunity to fungal infections. Nat Rev Immunol. 2011 Apr

11;11(4):275–88.

19.   Bedoya SK, Lam B, Lau K, Larkin J, III. Th17 cells in immunity and

autoimmunity. Clin Dev Immunol. 2013;2013:986789.

20.   Di Cesare A, Di Meglio P, Nestle FO. The IL-23/Th17 Axis in the

Immunopathogenesis of Psoriasis. J Invest Dermatol. 2009 Jun

1;129(6):1339–50.

21.   Côme C, Cvrljevic A, Khan MM, Treise I, Adler T, Aguilar-Pimentel JA, et al.

CIP2A promotes T-cell activation and immune response to Listeria

monocytogenes infection. PLoS One. 2016;11(4):1–18.

22.   Hoo LS, Zhang JY, Chan EK. Cloning and characterization of a novel 90 kDa

'companion' auto-antigen of p62 overexpressed in cancer. Oncogene. 2002

Jul;21(32):5006–15.

23.     Choi H, Larsen B, Lin Z-Y, Breitkreutz A, Mellacheruvu D, Fermin D, et al.

654             SAINT: probabilistic scoring of affinity purification-mass spectrometry data. Nat

655             Methods. 2011 Jan;8(1):70–3.

656     24.     Mellacheruvu D, Wright Z, Couzens AL, Lambert J-P, St-Denis NA, Li T, et al.

657             The CRAPome: a contaminant repository for affinity purification–mass

658             spectrometry data. Nat Methods. 2013 Aug 7;10(8):730–6.

659     25.     Myant K, Qiao X, Halonen T, Come C, Laine A, Janghorban M, et al. Serine

660             62-Phosphorylated MYC Associates with Nuclear Lamins and Its Regulation

661             by CIP2A Is Essential for Regenerative Proliferation. Cell Rep. 2015 Aug

662             11;12(6):1019–31.

663     26.     Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, et al. The

664             STRING database in 2017: quality-controlled protein-protein association

665             networks, made broadly accessible. Nucleic Acids Res. 2017 Jan

666             4;45(D1):D362–8.

667     27.     Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al.

668             Cytoscape: a software environment for integrated models of biomolecular

669             interaction networks. Genome Res. 2003 Nov;13(11):2498–504.

670     28.     Kauko O, Imanishi SY, Kulesskiy E, Laajala TD, Yetukuri L, Laine A, et al.

671             Rules for PP2A-controlled phosphosignalling and drug responses. bioRxiv.

672             2018 Feb 26;271841.

673     29.     Tripathi SK, Välikangas T, Shetty A, Khan MM, Moulder R, Bhosale SD, et al.

674             Quantitative Proteomics Reveals the Dynamic Protein Landscape during

Initiation of Human Th17 Cell Polarization. iScience. 2019 Jan 26;11:334–55.

30.    Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of

large gene lists using DAVID bioinformatics resources. Nat Protoc. 2009 Jan

1;4(1):44–57.

31.    Huang DW, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths

toward the comprehensive functional analysis of large gene lists. Nucleic Acids

Res. 2009 Jan;37(1):1–13.

32.    Mi H, Huang X, Muruganujan A, Tang H, Mills C, Kang D, et al. PANTHER

version 11: expanded annotation data from Gene Ontology and Reactome

pathways, and data analysis tool enhancements. Nucleic Acids Res. 2017 Jan

4;45(D1):D183–9.

33.    Mi H, Muruganujan A, Casagrande JT, Thomas PD. Large-scale gene function

analysis with the PANTHER classification system. Nat Protoc. 2013 Jul

18;8(8):1551–66.

34.    Shi Y. Serine/Threonine Phosphatases: Mechanism through Structure. Cell.

2009 Oct 30;139(3):468–84.

35.    Nie H, Zheng Y, Li R, Guo TB, He D, Fang L, et al. Phosphorylation of FOXP3

controls regulatory T cell function and is inhibited by TNF-α in rheumatoid

arthritis. Nat Med. 2013 Mar 10;19(3):322–8.

36.    Lee J, Park E-J, Hwang JW, Oh J-M, Kim H, Bae E-K, et al. CIP2A expression

is associated with synovial hyperplasia and invasive function of fibroblast-like

synoviocytes in rheumatoid arthritis. Rheumatol Int. 2012 Jul 9;32(7):2023–30.

37.    Lee J, Jeong H, Park E-J, Hwang JW, Huang B, Bae E-K, et al. CIP2A

facilitates apoptotic resistance of fibroblast-like synoviocytes in rheumatoid

arthritis independent of c-Myc expression. Rheumatol Int. 2013 Sep

2;33(9):2241–8.

38.    Chow LML, Fournel M, Davidson D, Veillette A. Negative regulation of T-cell

receptor signalling by tyrosine protein kinase p50 csk. Nature. 1993 Sep

9;365(6442):156–60.

39.    Veillette A, Latour S, Davidson D. N EGATIVE R EGULATION OF I

MMUNORECEPTOR S IGNALING. Annu Rev Immunol. 2002 Apr;20(1):669–

707.

40.    Zhang Q, Davis JJC, Lamborn IT, Freeman AF, Jing H, Favreau AJ, et al.

Combined Immunodeficiency Associated with *DOCK8* Mutations. N Engl J

Med. 2009 Nov 19;361(21):2046–55.

41.    Su HC. Dedicator of cytokinesis 8 (DOCK8) deficiency. Curr Opin Allergy Clin

Immunol. 2010 Dec;10(6):515–20.

42.    Jabara HH, McDonald DR, Janssen E, Massaad MJ, Ramesh N, Borzutzky A,

et al. DOCK8 functions as an adaptor that links TLR-MyD88 signaling to B cell

activation. Nat Immunol. 2012 May 13;13(6):612–20.

43.    Lazarevic V, Chen X, Shim J-H, Hwang E-S, Jang E, Bolm AN, et al. T-bet

represses TH17 differentiation by preventing Runx1-mediated activation of the

gene encoding RORγt. Nat Publ Gr. 2010;12.

44.    Wang Y, Godec J, Ben-Aissa K, Cui K, Zhao K, Pucsek AB, et al. The

transcription factors T-bet and runx are required for the ontogeny of

pathogenic interferon-γ-producing T helper 17 cells. Immunity.

2014;40(3):355–66.

45.    Espinosa A, Dardalhon V, Brauner S, Ambrosi A, Higgs R, Quintana FJ, et al.

Loss of the lupus autoantigen Ro52/Trim21 induces tissue inflammation and

systemic autoimmunity by disregulating the IL-23–Th17 pathway. J Exp Med.

2009 Aug 3;206(8):1661–71.

46.    Yoshimi R, Ishigatsubo Y, Ozato K. Autoantigen TRIM21/Ro52 as a Possible

Target for Treatment of Systemic Lupus Erythematosus. Int J Rheumatol.

2012;2012:718237.

47.    Zhou G, Wu W, Yu L, Yu T, Yang W, Wang P, et al. Tripartite motif-containing

(TRIM) 21 negatively regulates intestinal mucosal inflammation through

inhibiting TH1/TH17 cell differentiation in patients with inflammatory bowel

diseases. J Allergy Clin Immunol. 2017 Nov 4;

48.    Machnicka B, Grochowalska R, Bogusławska DM, Sikorski AF, Lecomte MC.

Spectrin-based skeleton as an actor in cell signaling. Cell Mol Life Sci. 2012

Jan;69(2):191–201.

49.    Sihag RK, Cataldo AM. Brain beta-spectrin is a component of senile plaques in

Alzheimer's disease. Brain Res. 1996 Dec 16;743(1–2):249–57.

50.    Yan X-X, Jeromin A, Jeromin A. Spectrin Breakdown Products (SBDPs) as

Potential Biomarkers for Neurodegenerative Diseases. Curr Transl Geriatr Exp

Gerontol Rep. 2012 Jun;1(2):85–93.

51.    Burgess A, Vigneron S, Brioudes E, Labbé J-C, Lorca T, Castro A. Loss of

human Greatwall results in G2 arrest and multiple mitotic defects due to

deregulation of the cyclin B-Cdc2/PP2A balance. Proc Natl Acad Sci U S A.

2010 Jul 13;107(28):12564–9.

52.    McCloy RA, Rogers S, Caldon CE, Lorca T, Castro A, Burgess A. Partial

inhibition of Cdk1 in G $_2$ phase overrides the SAC and decouples mitotic

events. Cell Cycle. 2014 May 6;13(9):1400–12.

53.    Carpenter AE, Jones TR, Lamprecht MR, Clarke C, Kang I, Friman O, et al.

CellProfiler: image analysis software for identifying and quantifying cell

phenotypes. Genome Biol. 2006;7(10):R100.

54.    Vizcaíno JA, Csordas A, del-Toro N, Dianes JA, Griss J, Lavidas I, et al. 2016

update of the PRIDE database and its related tools. Nucleic Acids Res. 2016

Jan 4;44(D1):D447–56.

55.    MacLean B, Tomazela DM, Shulman N, Chambers M, Finney GL, Frewen B,

et al. Skyline: an open source document editor for creating and analyzing

targeted proteomics experiments. Bioinformatics. 2010 Apr 1;26(7):966–8.

56.    Farrah T, Deutsch EW, Kreisberg R, Sun Z, Campbell DS, Mendoza L, et al.

PASSEL: the PeptideAtlas SRMexperiment library. Proteomics. 2012

Apr;12(8):1170–5.

57.    Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen J V., Mann M.

Andromeda: A Peptide Search Engine Integrated into the MaxQuant

Environment. J Proteome Res. 2011 Apr 1;10(4):1794–805.

58.   Cox J, Mann M. MaxQuant enables high peptide identification rates,

individualized p.p.b.-range mass accuracies and proteome-wide protein

quantification. Nat Biotechnol. 2008 Dec 30;26(12):1367–72.

59.   Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann M. Accurate Proteome-

wide Label-free Quantification by Delayed Normalization and Maximal Peptide

Ratio Extraction, Termed MaxLFQ. Mol Cell Proteomics. 2014

Sep;13(9):2513–26.

60.   Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, et al. The

Perseus computational platform for comprehensive analysis of (prote)omics

data. Nat Methods. 2016 Sep 27;13(9):731–40.

61.   Choi M, Chang C-Y, Clough T, Broudy D, Killeen T, MacLean B, et al.

MSstats: an R package for statistical analysis of quantitative mass

spectrometry-based proteomic experiments. Bioinformatics. 2014 Sep

1;30(17):2524–6.

62.   Morris JH, Apeltsin L, Newman AM, Baumbach J, Wittkop T, Su G, et al.

clusterMaker: a multi-algorithm clustering plugin for Cytoscape. BMC

Bioinformatics. 2011 Nov 9;12(1):436.

63.   Brohée S, van Helden J. Evaluation of clustering algorithms for protein-protein

interaction networks. BMC Bioinformatics. 2006 Nov 6;7(1):488.

64.   Desiere F, Deutsch EW, King NL, Nesvizhskii AI, Mallick P, Eng J, et al. The

PeptideAtlas project. Nucleic Acids Res. 2006 Jan 1;34(Database

issue):D655-8.

**Figure 1. CIP2A interactome top protein interaction in Th17 cells.**

**(A)** Western Blot (WB) analysis of CIP2A immunoprecipitated (IP) with two antibodies

(Ab1 and Ab2) specific to different regions of CIP2A in 72h polarised Th17 cells. For

each CIP2A targeting antibody, an IgG control antibody (IgG1 and IgG2) was used.

The figure is a representative WB of two experiments. Input, IgG control IP and CIP2A

IP reactions are shown.

**(B)** Schematic diagram of the human CIP2A protein structure. Regions specific to the

antibodies used in the pulldown are shown. Ab1: against residues surrounding valine

343 (monoclonal), Ab2: 618-905 peptide (polyclonal). Protein domain borders are

based on previous publication (Wang et al., 2017) in which the Armadillo repeat, and

the dimerization interface are shown in the CIP2A structured domain studied by Wang

et al., 2017.

**(C)** A Venn diagram showing the number of common and unique CIP2A interacting

proteins identified with two distinct antibodies targeting CIP2A based on the mass

spectrometry (ms) analysis. For details, please see the method section.

**(D)** Heatmap representation of the top fifty CIP2A interactors identified with both

antibodies. R1 and R2 represent data from two biological replicates. Normalized

spectral intensities are plotted as a heatmap to display protein abundance in

respective IP reactions.

**Figure 2. Cellular distribution of CIP2A and the proportions of CIP2A**

**interactome in Th17 cells.**

**(A)** Immunostaining for the study of CIP2A localisation in 72h polarised Th17 cells. A representative image of three replicates are shown. Cells were stained for endogenous CIP2A (green). Phalloidin (red) and DAPI (blue) were used to stain cytoplasmic and nuclear regions, respectively. Scale bar is 7μm.

**(B)** Statistical analyses of CIP2A expression for cellular localization studies from three replicates using GraphPad Prism version 8.0d for Mac OS X (GraphPad Software). Statistical analysis was made by a two-tailed paired Student's T-test, where p-value ** represent <0.01. The error bars represent standard error of the mean.

**(C)** Representative WB analysis of CIP2A localisation in the cytoplasmic and nuclear fractions of 72h polarised Th17 cells. Vimentin and Tubulin were used as controls for the nuclear and cytoplasmic fractions, respectively. WCL: Whole Cell lysate, NF: Nuclear Fraction, and CF: Cytoplasmic Fraction.

**(D)** Enriched cellular locations of the proteins in the CIP2A interactome presented in the form of pie chart. The enrichment analysis was performed using the Ingenuity pathway analysis (IPA; Qiagen).

**Figure 3. The CIP2A protein interaction network.**

**(A)** The interaction network among the CIP2A interacting proteins. Protein-protein interactions for the network were downloaded from the STRING database and visualized using the Cytoscape software. Nodes in the network were clustered using the Markov clustering algorithm. The identified clusters with more than two members are indicated by distinct colours. A representative GO biological process term is shown for each cluster with four or more members. The GO enrichment analysis was

performed using DAVID against a Th17 proteome reference background  (Tripathi et

al., 2019). The median MS intensity differences between the CIP2A immuno-

precipitates and the IgG controls are shown as node inner colour from white to grey

according to increasingly stronger expression in the CIP2A immuno-precipitates

compared to the IgG controls.

**(B)** Enriched Biological processes associated with the proteins in the CIP2A

interactome in human Th17 cells. The enrichment analyses was performed using

PANTHER with the enrichment calculated from a background of proteins detected in

Th17 proteomic reference background (Tripathi et al., 2019). The scale is –log FDR.

**(C)** Functional classification of the CIP2A interactome in Th17 cells. The proportions

of different enriched functional classes in the CIP2A interacting proteins. The

enrichment analysis was performed using IPA (Qiagen).

**Figure 4. CIP2A interactome validation by SRM mass spectrometry.**

Selected reaction monitoring targeted mass spectrometry (SRM-MS) validation of the

CIP2A interactome. CIP2A IP was performed using antibody Ab2 in 72h polarised

Th17 cells and twenty proteins were validated by targeted SRM-MS. Averaged results

from three replicates are presented in the form of a box plot. Statistical analysis was

made by a two-tailed paired Student's T-test, where * p-value <0.05; and *** p-value

<0.001. The error bars represent 95% confidence interval.

**Figure 5. Validation of CIP2A known interaction by Western Blot and Confocal**

**microscopy.**

**(A)** CIP2A IP performed in Th17 cells and WB detection to validates CIP2A known

interaction with phosphatase PP2A-A. Total lysate (Input) and control IP (IgG) and

CIP2A IP using Ab2 are shown on the blot. A conformational specific secondary

antibody was used to probe proteins without interference from the denatured IgG

heavy (50 kDa) and light chains (25 kDa).

**(B)** WB validation of the CIP2A-PPP1CA interaction. CIP2A immunoprecipitated with

two different antibodies (Ab1 and Ab2) in Th17 cells (72h) and detected as in A.

**(C-D)** Proximity ligation assay (PLA) assay validation of the CIP2A-PPP1CA

interaction (C) or selected proteins (TRIM21 and UBR5) interaction with CIP2A in Th17

cells (D). The GFP antibody was used as a negative control in PLA. DAPI was used

to stain the nuclei. Bar is 7$\mu$m. Representation of the interaction between CIP2A-

PPP1CA (C) or TRIM21 and UBR5 with CIP2A (D) by PLA as a univariate dot plot.

Average number of PLA signals (spots) per cell (n=10 images from a representative

of three experiments) was plotted. Statistics by paired two tailed T-test, where *

represents $P< 0.05$. and $**< 0.01$. Scale bar is 7$\mu$m

(**E**) WB validation represent interactions between TRIM21, IRF4, STAT1 and PP2A

with CIP2A in Th17 cells at 72h and CIP2A. IP was performed using Ab2 and

conformational specific secondary antibody was used as in Fig. 2C.

**(F)** CIP2A inhibition leads to decreased TNF$\alpha$-elicited Nf-$\kappa$B activation. Total and

phospho-p65 (pp65) Nf-$\kappa$B subunit expression in CIP2A silenced and control Th17

cells. CIP2A is silenced in three different individual donor cell pools, and CIP2A, pp65,

p65 and Actin are shown on the blot.

 **SUPPLEMENTARY FIGURE LEGENDS**

**Figure S1 (Related to Figure 1). The polarization of the Th17 cultures used in the**

**study and a schematic representation of the interactome study.**

**(A)** The proportion of CCR6+ cells from the flow-cytometry analysis of CCR6 surface

expression at 72h in TCR activated control Th0 cells and polarized Th17 cells. The

IL17A mRNA expression (**B**), and IL17A protein secretion (**C**) from the cultures at 72h

post initiation of Th17 polarization. The data (**A-C**) is from six independent cultures

and each culture consisted of cells from more than four individual donors. Error bars

represent SEM estimates across biological replicates and P-values calculated using

the paired two-tailed student's t-test. (n = 6; P <0.01(**) and P <0.001(***) for Th0 vs

Th17). (**D**) A schematic layout representation of the interactome study.

**Figure S2 (Related to Figure 2).**

WB analysis of two replicates for the cytoplasmic and nuclear fractions for CIP2A sub-

cellular localisation in 72h polarised Th17 cells. WB probes with antibodies as controls

for nuclear and cytoplasmic fractions detecting Vimentin and Tubulin proteins,

respectively. WCL, NF and CF represent Whole Cell lysate, Nuclear Fraction, and

Cytoplasmic Fraction respectively.

**Figure S3 (Related to Figure 3). The normalized expression of the top CIP2A**

**interacting proteins in selected highly enriched GO biological processes in the**

**detected CIP2A interactome.**

The 20 strongest CIP2A interacting proteins in the **A**) RNA metabolic process (**B**) RNA

splicing via transesterification and (**C**) RNA splicing via spliceosome enriched GO

biological processes. The expression values are normalized $\log_2$ transformed protein

intensities from the MS analysis of the CIP2A immuno-precipitates (IP) and the IgG

controls. The expression values for both antibodies (Ab1 and Ab2) specific to different

regions for two individual replicates are shown. The proteins in each heatmap are in

descending order based on the median expression differences between the CIP2A-

IPs and the IgG controls.

**Figure S4 (Related to Figure 4).**

The averaged logarithmic fold changes and logarithmic significance values (log FDR)

for the targeted mass spectrometry (SRM-MS) validations analysis of the twenty

selected (shown in Figure 4A) proteins from the CIP2A interactome.

**Figure S5 (Related to Figure 5).**

**(A)** UBR5 immunostaining in Hela cells and PLA validation for interaction between

CIP2A and UBR5 by confocal microscopy. For PLA, the antibodies were used as

negative controls. DAPI was used to stain nuclei. Bar = 10μm

**(B-C)** WB validations of selected proteins in CIP2A interactome (Two replicates). (**B**)

First replicate showing two separate blots, TRIM21 and IRF4 (first blot) and STAT1

and PP2A (second blot). (**C**) WB validations of the second replicate for the same

proteins. CIP2A pull-down in 72h polarised Th17 cells and CIP2A IP, IgG control IP

and Input reactions are shown on the blot. CIP2A IP was performed using Ab2 and

conformational specific secondary antibody was used to probe proteins without

interference from denatured IgG heavy (50 kDa) and light chains (25 kDa).

 **SUPPLEMENTRY TABLE LEGENDS**

**Supplementary Table S1 (A-C). CIP2A interactome in Th17 cells. (**Related to Fig.

1). Table list proteins associated with the CIP2A interactome (**Supplementary Table**

**S1A)**. The table also list Ingenuity Pathway Analysis (IPA) cellular location and

functional class assigned IPA annotation for the proteins detected in the CIP2A

interactome (**Supplementary Table S1B)**.   The CIP2A interactome associated

proteins information are listed together with the data from the MaxQuant  results

(unique   peptides,   sequence   coverage,   intensity   and   Andromeda   score),

representation in the CRAPome, and SAINT analysis showing Saint Probability (SP)

(**Supplementary Table S1C)**. The data is filtered to only include proteins with a SP

values of > 0.95 and less than 60% representation in the CRAPOME.

**Supplementary Table S2.** (Related to Fig. 3). **The 20 most enriched GO FAT**

**biological processes (BP) in each cluster of the CIP2A interactome network in**

**Figure 3A.** The GO enrichment was performed using DAVID (Huang et al., 2009)

against a Th17 proteome reference background (Tripathi et al., 2019). The GO FAT

BP terms, which filter out the broadest higher level GO terms, were considered for

more specific enrichment information. For each cluster with four or more members, 20

of the most frequently occurring GO FAT terms among the cluster members are listed.

For each term, the number of cluster members (proteins or nodes) with the term, the

proportion of cluster members with the term, cluster number and annotation for the

cluster in Figure 3A, is shown**.**

**Supplementary Table S3. (**Related to Fig. 3). **The enriched GO biological**

**processes in the detected CIP2A interactome.** The enrichment analysis was

performed using PANTHER (Mi et al., 2013, 2017) against a Th17 proteome reference

background  (Tripathi et al., 2019). The broad GO SLIM terms, which include only the

higher level GO terms, were considered for a functional summary of the interactome.

Fold enrichment, significance value (p-value) and false discovery rate is presented for

each term.

**Supplementary Table S4. SRM Targets for the CIP2A interactome validations.**

**(**Associated with Fig. 4 ). The table lists the peptides, associated transitions, collision

energies and retention time windows used for the SRM validations of the CIP2A vs.

control IgG pull downs.

**Supplementary Table S5. SRM MS CIP2A interactome validations. (**Associated

with Fig. 4 ). The table lists the $\log_2$ transformed relative abundance data from the

SRM analysis of the  CIP2A IP vs. control IgG pull downs from three biological

replicates for proteins validated for the CIP2A interactome.

**Figure 1**

**A**

IgG I.P | CIP2A I.P | Input
Ab1 Ab2 | Ab1 Ab2 | Ab1 Ab2

CIP2A

**B**

Structured (1-560) | Un-Structured

1 | 905

Armadillo repeat

Dimerisation interface

CIP2A Ab 1
(residues surrounding Val342;
Monoclonal)

CIP2A Ab 2
(618-905; Polyclonal)

**C**

Ab1 | Ab2

15 | 293 | 23

**D**

CIP2A Ab | IgG Control Ab

R1 | R2 | R1 | R2
Ab1 Ab2 | Ab1 Ab2 | Ab1 Ab2 | Ab1 Ab2

Intensity

KIAA1524
SPTBN1
TRIM21
ACIN1
TMOD3
PNN
PPP1R18
TCOF1
ZC3HAV1
EIF4A3
MYO1B
MYO18A
HNRNPL
PPP1CA
CAPZB
BCLAF1
ATAD3A
PPP1R12A
MYH10
ATAD3B
SRSF3
TAF15
SQOR
EDC4
PLEC
RPA1
IFI16
DOCK8
SNRNP200
RPL18A
RBM8A
LIMA1
PCM1
RPA3
FXR1
DHX15
RBMX
THRAP3
SRRM2
MYL6B
PRPF19
PABPN1
PABPC4
EFTUD2
ACTN4
MYO1C
PCBP2
TRA2B
ADD1
LRCH3

**Figure 2**

**A**

**B**

**C**

**D**

**A**

**Supplementary Figure S1** (Related to Figure 1)

**A**

## Supplementary Figure S3 (Related to Figure 3)

**Supplementary Figure S4** (Related to Figure 4)

**A**

Log2 (Fold Change)

**Supplementary Figure S5** (Related to Figure 5)