Comprehensive molecular characterization of pediatric treatment-induced high-grade glioma:

A distinct entity despite disparate etiologies with defining molecular characteristics and potential

therapeutic targets

John DeSisto[1]*, John T. Lucas Jr.[2]*, Ke Xu[3], Andrew Donson[1], Tong Lin[4], Bridget Sanford[1], Gang Wu[3],

Quynh T. Tran[5], Dale Hedges[2], Chih-Yang Hsu[2], Gregory T. Armstrong[6,8], Michael Arnold[9], Smita

Bhatia[8,10], Patrick Flannery[1], Rakeb Lemma[1], Lakotah Hardie[11], Ulrich Schüller[12], Lindsey M. Hoffman[1,11],

Kathleen Dorris[1,11], Jean Mulcahy Levy[1,11], Todd C. Hankinson[1,11], Michael Handler[1,11], Arthur Liu[11],

Nicholas Foreman[1,11], Rajeev Vibhakar[1,11], Kenneth Jones[1], Sariah Allen[5], Jinghui Zhang[3], Suzanne J.

Baker[7], Thomas E. Merchant[2], Brent A. Orr[5#], Adam L. Green[1,11#]

*These authors contributed equally to this work

#Co-Senior Authors

[1]Morgan Adams Foundation Pediatric Brain Tumor Research Program, University of Colorado School of

Medicine, Aurora, CO

Department of Radiation Oncology[2], Computational Biology[3], Biostatistics[4], Pathology[5], Epidemiology

and Cancer Control[6], and Developmental Neurobiology[7], St. Jude Children's Research Hospital,

Memphis, TN

[8]Childhood Cancer Survivor Study

[9]Department of Pathology, Nationwide Children's Hospital, Columbus, OH

[10]Division of Hematology and Oncology, University of Alabama at Birmingham

[11]Children's Hospital Colorado, Aurora, CO

[12]Institute of Neuropathology and Department of Pediatric Hematology and Oncology, University

Medical Center; Children's Cancer Center, Hamburg, Germany

Author Contributions

ALG and JTL conceived the project and overall experimental design. AD assisted with sample processing and data analysis. JTL and CH designed and analyzed *de novo* pHGG and TIHGG imaging data. ALG and JTL compiled and analyzed TIHGG clinical and imaging data; LH and AL performed analysis of radiation treatment for the Colorado cohort. MH and TCH procured tumor samples. US provided methylation profiling data and clinical information for tumor samples. SA processed samples and performed fluorescence in situ hybridization. TL, QT, and BAO analyzed 450/850K data from TIHGG and pHGG cases. JTL, KX, GW, and BAO conceived of the relevant comparisons to *de novo* pHGG copy-number and sequencing data. JD, BS, KJ, and ALG designed the relevant comparisons and performed analyses of germline and somatic sequencing data, RNA-seq data, drug screening results. SJB provided DNA methylation and sequencing data from PCGP cases with a history of therapeutic ionizing radiation. JD performed all primary cell line experiments, assisted by PF and RL. LMH, KD, JML, NF, and RV assisted with project planning and data analysis. GW, KX, KJ, and DH, performed all computational analyses and interpretation of the germline and somatic sequencing data. JTL, JD, and KX performed statistical analyses. MA, GA, BAO, NF, and SB contributed primary tumor samples and clinical data. KX completed the reconstruction of episomes from WGS data. JTL, AG, BAO, TL, JD, KX, QT and GW contributed to the interpretation of experiments. JTL and JD wrote the manuscript with input from all co-authors; all coauthors were involved with manuscript revision, which was led by JTL, JD, BAO, and ALG.

Acknowledgements

We thank members of the Zhang and Wu laboratories (St. Jude Children's Research Hospital)

and members of the Foreman and Vibhakar laboratories (University of Colorado) for assistance and

discussion. We also thank Matthew Lear (St. Jude Children's Research Hospital) for his help in acquiring

and performing preliminary clinical analysis of primary tumor samples and Scott Olsen, Granger Ridout,

and Emily Walker (Hartwell Center for Bioinformatics and Biotechnology) for their assistance in

sequencing samples. RNA-seq library preparation and high-throughput sequencing were conducted at

the University of Colorado Cancer Center Genomics Shared Resource; microscopy was performed at the

University of Colorado Denver Advanced Light Microscopy Core Facility; and tumor sample preparation

for imaging was performed by the University of Colorado Cancer Center Histology Shared Resource. This

work was supported by the National Cancer Institute (CA55727, G.T. Armstrong, Principal Investigator),

the Fördergemeinschaft Kinderkrebs-Zentrum Hamburg (US), St. Jude Children's Research Hospital

Cancer Center Support (CORE) grant (CA21765, C. Roberts, Principal Investigator), ALSAC, the Morgan

Adams Foundation (ALG), and a St. Baldrick's Foundation Scholarship (ALG).

Data Availability Statement

The datasets generated during and/or analyzed during the current study are currently available

from the corresponding authors upon request and will be made publicly available. Our teams are

current in the process of submitting all sequencing data to EMBL-EBI and will update the editors with

the link(s) as soon as possible.

Abstract

Treatment-induced high-grade gliomas (TIHGGs) are an incurable late complication of cranial

radiation therapy or combined radiation/chemotherapy used to treat pediatric cancer. We assembled a

cohort of 33 TIHGGs from multiple institutions. The primary antecedent malignancies were

medulloblastoma, acute lymphoblastic leukemia, astrocytoma, and ependymoma. We performed

methylation profiling, RNA-seq, and genomic sequencing (whole-genome or whole-exome) on TIHGG

samples. Methylation profiling revealed that TIHGGs cluster primarily with the pediatric receptor

tyrosine kinase I subtype (26/31 samples). Common TIHGG copy-number alterations include

Chromosome (Ch.) 1p loss/1q gain, Ch. 4 loss, Ch. 6q loss, and Ch. 13 and Ch. 14 loss; focal alterations

include *PDGFRA* and *CDK4* gain and loss of *CDKN2A* and *BCOR*. Relative to *de novo* pediatric high-grade

glioma (pHGG), *BCOR* loss (p=0.004) and *CDKN2A* loss (p=0.005) were significantly increased.

Transcriptomic analysis identified two distinct TIHGG subgroups, one with a lesser mutation burden

(0.12 mut/Mb), Ch. 1p loss/1q gain (5/6 samples), and stem cell characteristics, and one with a greater

mutation burden (1.08 mut/Mb, p<0.0002), depletion of DNA repair pathways, and inflammatory

characteristics. We observed increased chromothripsis in TIHGG versus pHGG (67% vs. 31%, p=0.036),

which was associated with extrachromosomal circular DNA-mediated amplification of *PDGFRA* and

CDK4. *In vitro* drug screening in one primary, patient-derived TIHGG cell line from each expression

subgroup identified microtubule inhibitors/stabilizers, DNA-damaging agents, MEK inhibition, and, in the

inflammatory subgroup, proteasome inhibitors as potentially effective therapies. This study provides a

comprehensive molecular profile of TIHGG, including mechanistic insights to TIHGG oncogenesis, and

identifies potentially effective therapeutic modalities for further investigation.

Introduction

Treatment-induced high-grade gliomas (TIHGGs) are pediatric high-grade astrocytomas with an estimated incidence of 3% following cranial radiotherapy[1]. The highest cumulative incidence of TIHGG is in survivors of medulloblastoma, ependymoma, and leukemia[1]. TIHGG represents a rare but significant cause of late mortality in survivors of childhood cancer[2], as few therapeutic options are available at diagnosis[3,4]. Genomic analysis of TIHGG has been restricted to only a few small cohorts, but early evidence suggests that, in comparison with *de novo* pediatric high-grade gliomas (pHGG), TIHGG may be enriched for recurrent copy-number alterations and for amplification of *PDGFRA* and other potential activators of the MAP kinase pathway[5,6]. *TP53* mutations, gain of 1q, deletion of *CDKN2A*, and overexpression of *ERBB3* and *SOX10* may also be more frequent[5,7]. Similarly, alterations in treatment-related second cancers outside the brain show recurrent genomic copy-number variations, some of which contain genes that modify tumor latency *(NF2*, *CHEK2*, and *RET* amplification)[8,9,10]. These divergent oncogenic programs are spread over multiple histologies, limiting inference regarding potential therapeutic options.

In this study, we present an evaluation of the largest TIHGG cohort assembled to date, using comprehensive clinical and molecular analyses. We conducted a clinical review of each case, including the initial tumor location and type, treatment received, and the resulting TIHGG location and histology, followed by DNA methylation analysis using 450/850k arrays, genomic sequencing of TIHGG and matched blood samples, and RNA-seq. We assemble a comprehensive molecular profile of TIHGG, enabling us to identify several previously unknown characteristics of the disease, and propose a method of grouping TIHGG tumors phenotypically based upon a combination of genomic alterations and gene expression characteristics. Our findings differentiate TIHGG from pHGG and provide rationale for alternative therapeutic approaches, which we explore in *in silico* and *in vitro* pre-clinical therapeutic screening studies.

Results

Patient characteristics

The median age at diagnosis of first cancer within the TIHGG cohort was 6.5 years (range, 0.16-

19). The initial diagnoses included medulloblastoma (N=12; 36%), acute lymphoblastic leukemia (ALL)

(N=10; 30.3%), astrocytoma (N=3; 9%), ependymoma (N=2; 6%), germinoma (N=2; 6%), lymphoma (N=1;

3%), craniopharyngioma (N=1; 3%), neuroblastoma (N=1; 3%), and ganglioglioma (N=1; 3%) (Sup. Table

1). Two patients had known mismatch repair deficiencies resulting from a heterozygous germline loss of

MSH2 in one case and *PMS2* in the other (Case 2, Case 32). Both were included in the study because

each had therapeutic radiation exposure to the cranium with sufficient latency (11 and 8 years) to meet

the TIHGG diagnostic criteria. All cases had ionizing radiation exposure, except for one case with a prior

history of high-dose chemotherapy exposure for treatment of high-risk neuroblastoma (further details

on clinical histories are discussed in Methods). The individual clinical histories for the TIHGG cohort with

extended clinical information are presented in Sup. Fig. 1.

Complete clinical-pathologic data regarding treatment history were available in 76% of cases

(Fig. 1A (showing analyses/information available for the entire cohort), Sup. Fig. 1, Sup. Table 1, 2).

Patients received total body irradiation (TBI) (N=1, 3%), cranial irradiation (N=6, 18%), craniospinal

irradiation (N=9, 24%), or focal brain irradiation (N=10, 29%) to treat the initial malignancy (Fig. 1B). One

patient received radiotherapy to the abdomen (Case 44). Radiotherapy field extent was unknown in 24%

(N=7) of cases. The indications for radiotherapy were adjuvant (N=12, 36%), definitive (N=1, 3%), and

salvage (N=9, 26.5%). Radiotherapy technique was 2D (N=13, 38.2%) or 3D conformal radiation therapy

(N=8, 23.5%). Technique was not known in 38% (N=13) of cases. The location of the TIHGG relative to

the initial radiotherapy field was in-field but outside the high-dose region in 24% (N=8) of evaluable

cases and in the high-dose (prescribed dose) region in 52% of cases (N=17).

The median time between initial treatment and TIHGG diagnosis was 8.0 years (95% CI 7.3-13.2

years) (Fig. 1C). The median age at the time of diagnosis of TIHGG was 13.6 years (9-26 years) (Sup.

Table 1). TIHGG tumor grades were WHO grade III (N=3, 9%), grade IV (N=12, 55%), and pHGG not

otherwise specified (N=12, 36%) (Sup. Table 1). Assessment by methylation array identified *MGMT*

promoter methylation in 8 of 31 (25.8%) cases in which methylation status was evaluated during the

clinical course of treatment.

Salvage therapy characteristics for TIHGG were available in 38% of cases (N=13) and included

resection (N=3, 23.1%), repeat radiotherapy (N=12, 92.3%), and/or chemotherapy (N=4, 30.7%) (Sup.

Table 2). The median radiotherapy dose for the second course was 54 Gy (range 30-59.4 Gy). At the time

of this analysis, all TIHGG patients with known outcome were deceased (N=26), and 7 had unknown

clinical status (Fig. 1D). The median survival in the TIHGG cohort was 9 months (95% CI 5-24.7 months)

(Fig. 1D, Sup. Table 2).

The use of radiotherapy, chemotherapy, and/or resection did not substantially improve

treatment outcomes, although a paucity of details available regarding therapy limited power for this

analysis (Sup. Table 2). These outcomes are consistent with prior published clinical series of TIHGG[4,8].

Voxel-wise lesion mapping of TIHGG relative to* de novo *pHGG

An MRI was available at diagnosis in 21 TIHGG cases (Sup. Fig. 2A). In cases for which location

information was available, TIHGGs were located in the infratentorial brain (posterior fossa) (N=11,

32.3%), supratentorial brain (N=12, 35.3%), supra- and infratentorial brain (N=1, 3%), and spine (N=1,

3%) (Fig. 1E, Sup. Fig. 2B). An additional cohort of 84 *de novo* pHGG was used for comparison. Compared

to a cohort of 84 de novo pHGG, TIHGG tumors were more likely to involve the frontal and occipital

lobe, and posterior fossa (cerebellum and inferior brainstem) regions relative to primary pHGG (Fig. 1E,

Sup. Fig. 2B).

TIHGGs cluster into a specific methylation group

We began our molecular analysis using DNA methylation profiling because methylation data (or

a DNA sample sufficient to conduct methylation profiling) were available for the greatest number of

TIHGG samples in our cohort. We first analyzed the methylation data using unsupervised clustering of 34

TIHGG samples from 31 tumors against a combined reference cohort containing pediatric and adult CNS

malignancies[11,12]. Twenty-five of 31 cases clustered among a group of epigenetically similar tumors

consisting of *H3K27M*-negative midline gliomas and gliomas in the pediatric receptor tyrosine kinase I

(pedRTK I) subgroup (Fig. 2A-C and Sup. Fig. 3A-C)[13]. TIHGGs were not readily distinguishable by

methylation from IDH wt, GBM-MID, and pedRTK I cases from the Capper and Korshunov reference

cohorts[11], so the combined group of TIHGG cases is designated as "pedRTK I" going forward, reflecting

their closer overall proximity to the centroids of the pedRTK I reference cases (Sup. Table 3). The

remaining six TIHGGs clustered with the GBM K27M (N=1), control cerebellum (N=2), anaplastic pilocytic

astrocytoma (N=1), HGNET-MN1 (N=1), and pleomorphic xanthoastrocytoma (N=1) methylation groups.

We confirmed the initial classification results using hierarchical and supervised methods (Sup. Table 4

and Sup. Table 5). No trend was noted associating initial cancer diagnosis with subsequent TIHGG

methylation class (Sup. Fig. 4A)[14].The latency and survival of TIHGG tumors classified as pedRTK I did not

differ significantly from that of those classified with other subclasses (Sup. Fig. 4B-C). The odds of being

assigned to the pedRTK I subgroup in the TIHGG cohort were 29.2 times the odds of being assigned to

that subgroup in the *de novo* pHGG cohort (95% CI 9.15-92.9, p<0.0001) (Sup. Fig. 5A, Sup. Table 6).

Survival of the TIHGG-pedRTK I was inferior to pHGG pedRTK I (median survival 9 months, [95% CI, 5-26]

vs. 16 months [95% CI, 10-not reached, p=0.00021] (Sup. Fig. 5B).

Alignment of TIHGGs with pedRTK I suggested the possibility that TIHGGs, despite their

disparate clinical precursors, might share common molecular alterations, which we investigated by

comparing copy-number alteration and transcriptomic profiles discussed in the following sections.

TIHGGs show recurrent copy-number abnormalities

We next reviewed the methylation profiling data for large segment and focal copy-number

alterations. Total copy-number variation was significantly increased in TIHGG when comparing *de novo*

pHGG and TIHGG in the pedRTK I subgroup (Sup. Fig. 5C). Recurrent large segment copy-number

alterations identified from methylation profiling included Ch.1p loss (3/31, 9.7%), Ch.1q gain (3/31,

9.7%), Ch.6q loss (5/31, 16.1%), Ch.13 loss (8/31, 25.8%), and Ch.14 loss (3/31, 9.7%) (Fig. 3A, Sup. Table

5, Sup. Fig. 6). Copy-number alterations in *PDGFRA* gain/amplification (15/31, 48.4%), *CDK4*

amplification (5/31, 12.9%), *CDKN2A* loss (12/31, 38.7%), and *BCOR* loss (7/31, 22.6%) were also noted.

These were increased relative to *de novo* pHGG for *PDGFRA* (p=0.0055) and *BCOR* (p=0.0043) (Sup. Table

7, Sup. Fig. 7). Changes in copy number of *PDGFRA, CDKN2A,* and *BCOR* were associated with an

accompanying change in gene expression in cases with available RNA-Seq data (N=12, 34.3%) (Fig. 3B).

When comparing total genomic copy-number variations by case, TIHGG favored loss over gain (Fig. 3C).

Overall, TIHGGs showed large segment genomic alterations in Ch.1 (1p loss/1q gain) and Ch.13

(loss), and focal alterations in *PDGFRA* (amplification)*, CDKN2A* (loss) and *BCOR* (loss). As further

discussed below, where alterations were identified in both WES/WGS and methylation data, we

obtained consistent results between the two methods. The *PDGFRA* amplification that we observed is

consistent with the alignment of most TIHGGs with the pedRTK I methylation subclass[11].

TIHGG has a high frequency of chromothripsis and episomal amplification of oncogenes

202   To further investigate the potential sources of copy-number amplification, we evaluated the

rate of chromothripsis in the TIHGG samples relative to that in primary non-brainstem and brainstem

pHGG from Wu *et al.*[15]. We observed an increase in the frequency of chromothripsis in the TIHGG cohort

(8/12 cases) relative to that in diffuse intrinsic pontine glioma (DIPG) (66.7% vs. 30.0%, p=0.048), non-

brainstem pHGG (66.7% vs. 33.3%, p=0.091), and all primary pHGG combined (66.7% vs. 31.4%, p=0.036)

(Fig. 4A, Sup. Table 8). Both intra-chromosomal (Case 29-dm2) rearrangements leading to

extrachromosomal circular DNA (eccDNA) structures, as well as inter-chromosomal rearrangements

(Case 29-dm1) leading to eccDNA structures, were observed in case 29 (Fig. 4B, D). *PDGFRA* and *CDK4*

appeared to be amplified separately on the two eccDNA structures (Fig. 4B, D). Amplification was

confirmed in selected cases by fluorescence *in situ* hybridization, which demonstrated a double minute

pattern, a finding typical of episomal amplification (Fig. 4C)[16]. There was a trend toward an increased

frequency of chromothripsis in cases with a coincident pathogenic *TP53* mutation (5/8 cases, 62.5%)

relative to those with wild-type *TP53* (0/4 cases, 0%, p=0.08).

215   In summary, we observed an elevated frequency of chromothripsis in TIHGG relative to *de novo*

216 pHGG, which facilitated extrachromosomal amplification of known oncogenes (*PDGFRA, CDK4).*

218 ***TIHGGs show recurrent abnormalities in genes that regulate cell proliferation, growth, and cell cycle***

219   In addition to copy-number variation, we also identified focal alterations in TIHGG using

WES/WGS data. Eighteen TIHGG cases underwent whole-exome (WES) or whole-genome sequencing

(WGS) (Fig. 5). Tier 1 mutations in the 5 non-hypermutator WES cases without matched germline data

are shown in Sup. Table 9; Tier 1 mutations for the 9 non-hypermutator WGS cases with matched

germline data are shown in Sup. Table 10. The most frequent focal alterations included *PDGFRA*,

CDKN2A, *BCOR*, *NF1*, *TP53*, and *CDK4* (Fig. 5, Sup. Tables 7, 9, 10). Missense mutations were noted in

TP53 and *PDGFRA* in 7 of 8 and 4 of 15 cases with an observed alteration in *TP53* and *PDGFRA,*

respectively. Nonsense and frameshift mutations of *NF1* occurred in 8 of 9 observed cases with *NF1*

alterations. *MET* fusion gene products were identified from RNA-seq samples in three cases (Sup. Table

11). Structural variations and copy-number alterations from 10 WGS cases are shown in Sup. Tables 12

and 13.

Compared to pHGG, TIHGG are enriched for specific somatic and germline abnormalities

We compared select recurrent alterations in the TIHGG cohort to *de novo* pHGG from the HERBY

cohort across each cohort as a whole and among pedRTK I tumors (Sup. Fig. 8, Sup. Table 14)[14].

Statistically significant increases in *BCOR* loss (p=0.0004) and *CDKN2A* loss (p=0.05) were observed in

TIHGG relative to pHGG (Sup. Table 7). No statistically significant differences were observed between

the proportions of molecular alterations in pedRTK I TIHGG relative to pHGG – pedRTK I tumors,

although power was limited by the sparse number of pedRTK I cases in the HERBY cohort (N=8). Of note,

the frequency of *TP53* mutations in pedRTK I TIHGG cases was 43.8% (7/16) and 62% (5/8) in the *de*

novo pHGG pedRTK I cases (p=0.68, Sup. Table 7). We observed a single *H3F3A* mutation (Fig. 5, Case 7)

(1/19, 5.6%) in the TIHGG cohort, compared to 37.8% (28/74) with mutant *H3F3A* (p=0.0053) in the

HERBY cohort (composed of non-brainstem pHGG).

We also compared variant frequency between the TIHGG cohort and pHGG cases in the HERBY

cohort. Germline mutations in the TIHGG cohort were analyzed in 10 cases for which WGS data were

available (Sup. Table 15). The total number of germline and somatic variants per megabase (Mb) were

calculated and stratified by pHGG versus TIHGG and are compared in Sup. Fig. 9A. The number of

variants per Mb was significantly higher in somatic coding regions of TIHGG DNA than in those of pHGG

(median 0.000784 ± 0.000286 (range) variants/Mb vs. median 0.0007 ± 0.000245 (range) variants/Mb,

p=0.031). The relative frequencies of base transitions were calculated per sample in pHGG and TIHGG

(Sup. Fig. 9B). Germline and somatic coding base changes were not significantly different across base

change type. Somatic noncoding region variants showed decreased A to C (median 14.4 ± 4.91 vs. 8.8 ±

4.1, p=0.007) and A to G (34.3 ± 9.67 vs. 20.2 ± 11.3, p=0.03), and increased C to A (6.79 ± 2.2 vs. 16.7 ±

5.47, p=0.004) base changes in TIHGG cases relative to *de novo* pHGG (Sup. Fig. 9B-C).

In summary, TIHGG have comparable burden of somatic and germline alterations to *de novo*

pHGG. Frequent pathogenic alterations were noted in *PDGFRA, BCOR, CDK4, TP53,* and *NF1.*

Transcriptomic analysis demonstrates that TIHGGs cluster into two subgroups that are separate from

pHGG

Transcriptomic data (array and RNA-seq) were available on 14 TIHGG cases and 42 *de novo*

glioblastoma (GBM) cases. TIHGG tumors clustered separately from *de novo* pediatric, infant, and adult

GBM based on transcriptomic data and formed two distinct subgroups (A and B) containing 6 and 8

tumor samples respectively (Fig. 6A). A comparison of the transcriptomic and methylation-based

clustering results showed that the two gene expression-based TIHGG subgroups were, with few

exceptions, recapitulated in the methylation analysis (Fig. 6B). Our data thus suggest two distinct

patterns of gene expression within the pedRTK I methylation-based subgroup into which most of the

TIHGGs cluster.

To further investigate gene expression in TIHGG, we performed Metascape/Cytoscape analysis,

which revealed several differences in gene ontology between TIHGG and the transcriptional reference

group of *de novo* GBM. The differences include fundamental cellular processes, such as RNA processing

and transport, protein translation and catabolism, and cellular signaling, as well as pathways controlling

neurogenesis and gliogenesis (Fig. 6C, Sup. Table 16). To determine the directionality of gene expression

patterns identified in the Metascape/Cytoscape analysis, we performed gene set enrichment analysis

(GSEA) using the gene ontology (GO) gene set collection[17,18]. The results of this analysis showed

similarities in differential gene expression to that identified in the Metascape/Cytoscape analysis, with

TIHGG samples showing relative enrichment of genes involved in DNA metabolism, cell cycle

progression, DNA repair, nervous system development, and protein catabolism compared to *de novo*

GBM, and relative depletion of genes related to immune response, signaling and cellular response to

external stimulus, receptor activity, and neurogenesis (Fig. 6D, Sup. Table 17).

Expression-based characterization of TIHGG subgroups

To characterize the TIHGG gene expression subgroups, we analyzed the differences in their

relative gene expression, mutational status, and copy-number alterations. GSEA revealed that TIHGG

gene expression Group A shows stem cell (proneural) and neuronal characteristics and has enriched

expression of *MYC* pathway genes (Fig. 7A). Expression Group B has mesenchymal and astroglial

characteristics, enriched expression of inflammatory (particularly NF-κB pathway) and immune genes,

and depletion of DNA-repair pathway genes (Fig. 7A, Sup. Table 18). We did not observe relative

expression differences between the TIHGG subgroups in cell cycle or proliferative genes, consistent with

clinical experience showing that TIHGGs, in general, are highly proliferative.

Using the DNA methylation data, we analyzed whether there were differences in copy-number

alterations between the two gene expression subgroups. We found Group A to be enriched in 1p loss/1q

gain (5/6 tumor samples) compared to Group B (0/7 tumor samples, p=0.005). GSEA showed particular

differences at 1p34 (NES = −5.21, FDR=0) and 1p36 (NES = −7.77, FDR=0). The remaining gene- or

chromosome-level amplifications and deletions identified in the TIHGG cohort, including Ch.13 or 14

loss, *PDGFRA* amplification, and *CDKN2A* loss, were relatively evenly distributed between the two TIHGG

subgroups (Sup. Table 5). We did not identify patterns of significant point mutation or similar small-scale

genetic differences between Group A and B tumors, nor did we find any relationship between initial

malignancy and TIHGG subgroup (Fig. 5).

The relative depletion of DNA repair pathway genes that we identified in Group B relative to

Group A led us to investigate whether there were differences in mutation load between the two groups.

Excluding one hypermutator case, WGS data were available on matched blood and tumor samples for 3

cases from Group A and 6 from Group B. We found nearly identical levels of germline mutation load in

Groups A and B (Fig. 7B). We found that Group B tumors, however, had a nine-fold greater somatic

mutation load than Group A tumors (1.08 vs. 0.12 mut/Mb, p<0.002) (Fig. 7B).

As noted previously, relative to Group A, Group B is depleted in DNA repair pathway gene

expression (mean NES = −3.96, p=0.001 vs. expected value of 1), which led us to compare DNA repair

pathway gene expression in group A and B germline samples. We found that germline samples from

Group A patients were likewise enriched in DNA repair pathway gene expression (mean NES = -4.30,

p=0.001) (Fig. 7C). Individual DNA repair pathway genes with lower expression in Group B relative to

Group A include *GTF2H2,* a subunit of RNA polymerase II transcription initiation factor IIH that is

involved in nucleotide excision repair; *PAPD7,* a DNA polymerase involved in DNA repair; *ERCC4,* which is

essential for nucleotide excision repair; and *TP53* (Fig. 7D, Sup. Table 19.). In a pairwise comparison of

the individual DNA repair pathway genes shown in Figure 7D, the mean expression ratio (Group B/Group

A) was 0.76 (p=9.79×10^{-11} vs. expected value of 1) (Sup. Table 19).

In summary, TIHGGs split into two transcriptional subgroups with distinct gene expression

profiles relative to *de novo* GBM. Subgroup A resembles proneural GBM and is enriched in expression of

MYC-pathway genes, while subgroup B resembles mesenchymal GBM and has greater mutational

burden and decreased DNA repair gene expression.

In silico *and* in vitro *drug screening identify potentially effective therapeutic agents for the treatment*

of TIHGG

In order to investigate therapeutic susceptibilities in TIHGG tumors, we used *in silico* modeling

to identify potential TIHGG sensitivity to drug classes compared to a set of 23 *de novo* pediatric and 14

de novo adult GBMs. We also performed an *in vitro* drug screen of two primary tumor-derived cell lines,

one from gene expression Group A (MAF-496) and one from Group B (MAF-145), for sensitivity to a

panel of all FDA-approved anti-cancer agents. The *in silico* modeling predicted several classes of

therapeutic agents to have greater activity in TIHGG when compared with *de novo* GBM. These included

DNA-damaging agents, microtubule polymerization/de-polymerization inhibitors, proteasome inhibitors,

and histone deacetylase (HDAC) inhibitors (Fig. 8A, Sup. Table 20).

Our *in vitro* screen encompassed all FDA-approved anti-cancer agents, including multiple agents

from the classes identified in the *in silico* screen, all dosed at 1 μM concentration (Sup. Figure 10A, Sup.

Table 21). We found, combining results from the two cell lines, that 11/12 DNA intercalator results, 6/12

microtubule agent results, 4/4 proteasome inhibitor results, and 3/6 HDAC inhibitor results showed at

least 50% cell death compared to vehicle (Fig. 8B, Sup. Table 21). The screen also identified

topoisomerase I/II inhibitors (6/8 results) and PI3K-MTOR, BCR-ABL, and RAF-MEK pathway inhibitors

(12/24 results total) as potentially effective using the 50% cell death criterion (Fig. 8B, Sup. Table 21).

Folate analogs (0/6 results with at least 50% cell death), platinum drugs (0/6 results), and alkylators

(3/32 results) performed poorly in the *in vitro* screen (Fig. 8B, Sup. Table 21). We found that, with the

exception of proteasome inhibitors, which are further discussed below, the Group A cell line (MAF-496)

was more susceptible to the effective drugs than the Group B cell line (MAF-145) (Fig. 8C).

To validate the screening results, we performed single-drug assays of selected agents at a range

of concentrations in the two TIHGG cell lines. Drugs were selected for validation based upon the

screening results and included two DNA-damaging agents (aldoxorubicin and etoposide), two

microtubule agents (paclitaxel and vinblastine), three proteasome inhibitors (bortezomib, carfilzomib,

and marizomib), and two kinase inhibitors (sunitinib, a multi-TKI inhibitor, and trametinib, a MEK

inhibitor) (Fig. 8D, Sup. Fig. 10B-G). In the validation testing, aldoxorubicin (an anthracycline drug

capable of blood-brain barrier penetration), etoposide, paclitaxel, and vinblastine showed sub-

micromolar IC_{50} values in both cell lines (Fig. 8D, Sup. Fig. 10B-C), as did the MEK inhibitor trametinib

(Fig. 8D, Sup. Fig. 10D). Sunitinib had an IC_{50} of 5,256 nM in the MAF-145 (Group B) cell line but did not

reach an IC_{50} level in MAF-496 cells (Group A) (Sup. Fig. 10E). Validation testing of bortezomib and

carfilzomib showed *in vitro* effectiveness in Group B cells, consistent with the enrichment of the NF-κB

pathway in Group B tumors (Fig. 8D, Sup. Fig. 10F). In the Group A cell line, however, 25-45% of cells

survived the highest concentration of each drug, suggesting the presence of a substantial drug-resistant

population in Group A (Fig. 8D, Sup. Fig. 10F). We found that the proteasome inhibitor marizomib, which

is capable of penetrating the blood-brain barrier, had IC_{50} values of 366 and 379 nM in the Group A and

Group B lines, respectively, but also had a drug resistant population of approximately 30% of cells in

both cell lines (Sup. Fig. 10G), which was not seen with bortezomib or carfilzomib. Further investigation

showed that proteasome inhibition reduced nuclear levels of NF-κB, a known target of proteasome

inhibition, in the Group B cell line (MAF-145) but had a lesser effect on already low nuclear levels of NF-

κB in the Group A line (MAF-496) (Sup. Fig. 10H-I).

Taken together, our drug screening results in TIHGG tumors showed that drugs that interfere

with S-phase (aldoxorubicin and etoposide) or M-phase (vinblastine and paclitaxel), and the MEK

inhibitor trametinib, were effective *in vitro* in both TIHGG subtypes. In addition, proteasome inhibition

was effective in the Group B cell line.

Discussion

The present TIHGG cohort represents the largest and most comprehensive studied to date and

extends prior knowledge from past studies that largely included targeted analyses of sequencing,

expression, and staining of smaller tumor collections. The use of methylation profiling, WES/WGS of

tumor and matched germline samples, RNA-seq, and drug screening help advance understanding of

long-held questions regarding TIHGG, including similarities and differences from *de novo* pHGG,

distinguishing TIHGG from recurrent tumors, TIHGG oncogenesis, subgroups within TIHGG, and potential

therapeutic susceptibilities.

All but one of our cases had a clinical history of cranial radiation exposure. The exception was a

high-risk neuroblastoma patient who received multiple rounds of multi-agent chemotherapy, including

two cycles of high-dose chemotherapy, but only received radiation to her primary tumor site in the

mediastinum, far removed from her frontal lobe TIHGG. Her latency and tumor findings from our assays

cluster closely with the remainder of the cohort, suggesting chemotherapy was most likely causative.

While it was exceptional that chemotherapy exposure alone should give rise to a TIHGG, the term

treatment-induced is preferred to radiation-induced to reflect this possibility and the known interaction

between combined modality therapy and increased incidence of subsequent malignancies[1]. In addition,

since these tumors were a mix of WHO grade III, IV, and HGG NOS, we have replaced the term

glioblastoma with high-grade glioma.

The results of our methylation and transcriptomic analyses delineate TIHGG as an entity

separate from pHGG and identify two distinct transcriptional subgroups within the TIHGG cohort.

Additional genomic and transcriptomic analyses identify characteristic genetic alterations and gene

expression profiles that define the two subgroups. TIHGG were most frequently assigned to the pedRTK I

methylation subgroup of pHGG, despite their varied clinical origin. Of note, resolving pedRTK I from

IDHwt, GBM-mid subtype pHGG, and TIHGG was not possible in our analysis, thus highlighting the

molecular and epigenetic similarities between these groups. One TIHGG case in our series clustered with

diffuse midline glioma (DMG) but lacked the defining *H3K27M* mutation (Case 13). One recent case

series described three patients who had received prior radiotherapy for medulloblastoma and later

developed tumors clinically consistent with DMG[19]. While one case in our series had the H3K27M

mutation (Case 7), this case lacked methylation data and was not be able to be clustered against

reference cohorts. Because of the close association between the pedRTK I subgroup with TIHGG, specific

consideration may be useful in future brain tumor taxonomy. The relative contribution of *de novo* HGG

and TIHGG remains to be established. Additionally, whether exposure to ionizing radiation outside the

therapeutic setting also predisposes to the tumor subclass is currently unclear.

We compared recurrent genetic alterations in TIHGG and *de novo* pHGG cases in the pedRTK I

group to define potential differentiating characteristics. We observed an increased frequency of

CDKN2A and *BCOR* losses in TIHGG. *BCOR* alterations are infrequently observed in pHGG without co-

segregating *HIST1H3B* K27M mutations and are not described in pedRTK I pHGG[20]. While *BCOR*

alterations noted in the cohort were predominately non-focal, the region showed a high frequency of

overlap across TIHGG cases, suggesting potential biologic relevance. Of note, *BCOR* alterations noted in

the Pediatric Cancer Genome Project were primarily due to frameshift or nonsense mutations and were

often coupled with loss of heterozygosity[21]. Despite the increased frequency of *CDKN2A* deletions, we

did not observe accompanying *BRAF* V600E alterations, a finding common in pediatric low-grade glioma

and some pHGG[22,22]. Because we observed recurrent activation of MAP kinase pathway in our cohort,

concomitant CDKN2A loss may be a necessary mechanism to prevent oncogene induced senescence, as

has been described in other tumor types[23].

The differentiation of TIHGG from recurrent pHGG or transformed low-grade glioma based on

DNA methylation profiling remains challenging. Two cases in our cohort arose following focal

radiotherapy for an optic pathway glioma but were localized to a region within the radiotherapy low

dose region but distant from the original cancer and not coincident with a metastasis. Prior to review,

these were expected to be transformation events, but after reviewing location of the TIHGG in each case

relative to the original tumor and the radiotherapy field, it is clear that the TIHGG arose in previously

uninvolved but irradiated normal brain. It remains unclear whether modern methods such as IMRT,

which expose larger regions of normal tissue to non-therapeutic, albeit potentially mutagenic doses of

radiation, will yield an increased incidence of TIHGG despite the increased conformality of the

therapeutic dose region.

RNA-seq analysis differentiated the overall TIHGG cohort most distinctly from pHGG, which has

been previously documented in array expression analysis[24]. Our results furthermore show distinct

subgroups within the TIHGG cohort: a low mutation load group A that is enriched for Ch.1p loss/1q gain,

has neuronal characteristics, and is enriched in expression of stemness genes and transcriptional targets

of *MYC*, and a higher mutation load group B that is enriched in expression of inflammatory and immune

pathway genes, has astroglial and mesenchymal characteristics, and is impaired in DNA repair processes.

The subgroups do not correlate with original tumor type or histology. Instead, the depleted germline

DNA repair pathway gene expression in the Group B patients suggests their potential susceptibility to

develop TIHGG following mutagenic cancer treatment.

Given the potential for mutational profile to identify malignancy type and the difficulty of

distinguishing recurrent primary from secondary tumors, we evaluated the frequency and type of

somatic variants in TIHGG and pHGG. The relative frequency of pathogenic variants in all TIHGG in our

cohort was not substantially increased relative to pHGG, as other groups have found[7], but Group B

TIHGG had increased mutational burden compared to Group A, suggesting heterogeneity in the pedRTK

I/TIHGG subgroup. We also observed differential rates of select base changes in somatic non-coding

regions. Increased frequency of variations in non-coding regions have also been observed in syndromes

associated with genomic instability[25], as mutations in coding regions are likely to result in negative

selective pressure.

The therapeutic potential of photon radiotherapy in tumor tissue has always come at the cost of

sub-lethal damage in adjacent normal tissue[26]. Irradiated normal tissue damaged directly by secondary

electrons or indirectly by reactive oxygen species is known to produce non-lethal double-stranded DNA

breaks, which lead to impaired function and survival in long-term survivors of childhood cancer[26].

Incomplete error-prone repair of genomic DNA via non-homologous end joining and non-error prone

repair via homologous recombination result in genomic instability, which may be further modified by

host normal tissue radiosensitivity, dictated by germline DNA damage response functionality.

TIHGG exhibits signs of a mutagenic origin, with oncogenic amplifications of Ch.1q, *PDGFRA* and

CDK4, copy number losses in tumor suppressors including *CDKN2A* and *BCOR*, and pathogenic mutations

in *TP53, NF1* and *MET* (fusion events). TIHGGs have more frequent *PDGFRA* gain and *CDKN2A* loss

relative to *de novo* pHGG; TIHGG also displays increased expression of oncogenic pathways indicative of

increased proliferation, including DNA metabolism and cell cycle genes. The characteristics of the two

TIHGG gene expression subgroups suggest differences in tumorigenesis related to differing types of DNA

damage and their respective DNA repair capabilities. Group A TIHGGs all have large-scale, chromosome-

level abnormalities that are associated with poor outcome in pediatric brain tumors, including Ch. 1q

gain and Ch. 13 loss. Gain of Ch. 1q distinguishes pediatric from adult HGG, has been recognized as a

feature of radiation-induced tumors, and is an indicator of progression and poor outcome in

ependymoma, neuroblastoma, and Wilms tumors[5,27-30]. Specific associations between chromosome-level

abnormalities and the other characteristics of group A tumors have not been reported but are an area

for further inquiry. The higher mutation load of group B tumors could arise through the combination of

mutagenic treatment-induced DNA damage and the downregulation of DNA repair pathways in these

cases. Further insight into potential germline susceptibilities in group B patients related to impaired DNA

repair may even lead to identification of patients at risk for TIHGG prior to primary tumor therapy and

allow treatment modification to prevent TIHGG. Based on our germline data, we hypothesize these

germline susceptibilities are different from known mutations causing tumor predisposition syndromes in

that they may only manifest in the setting of highly mutagenic treatment.

Large-scale, localized DNA damage in the form of chromothripsis has also been recognized as a

means of carcinogenesis following radiotherapy, as it can perpetuate a string of subsequent random

molecular alterations. While pHGG exhibit genomic instability secondary to alterations in histone- and

chromatin-modifying machinery, we observed an increased rate of chromothripsis in TIHGG relative to

that of *de novo* pHGG. Chromothripsis was more common in the setting of pathogenic *TP53* mutations.

This has also been observed in pediatric and adult malignancies not associated with therapeutic ionizing

radiation[31]. Although chromothripsis is known to result in complex chromosomal alterations that can

perpetuate a high frequency of copy-number alterations, we also observed characteristic broad and

focal copy-number changes in non-chromothripsis cases (+1q, −6q, −13, −14, *+PDGFRA, +CDK4,*

−CDKN2A, −BCOR) that have known oncogenic potential[27 22,32-34]. We observed several instances of

oncogene amplification, likely facilitated by inclusion in eccDNA, which has been identified as a

mechanism of resistance to targeted therapies. In tumor cells with eccDNA containing the oncogene

EGFR, eccDNA levels increase in response to EGFR inhibition[35]. The eccDNA increase is frequently

independent of and more efficient, in terms of onset and magnitude of response, than chromosomal

copy-number alterations[35-37]. Similarly, serial samples from patients exposed to standard-of-care

therapy show increases in *EGFR* levels during treatment with *EGFR* inhibitors, also likely in part

facilitated by the presence of eccDNA[16]. These findings may explain the lack of observed efficacy of well-

characterized, CNS-penetrant, targeted drugs such as PDGFRA inhibitors in our preclinical drug screens.

In silico and *in vitro* drug screening identified several FDA-approved drugs that merit further

study as potential therapeutic agents to treat TIHGG. Drug classes identified as most effective against

TIHGG (DNA-damaging agents, anti-mitotic drugs that target microtubules, and histone deacetylase

(HDAC) inhibitors) have not been clinically effective in *de novo* pHGG, though clinical studies of HDAC

inhibitors are ongoing.[38,39] Mechanistically, Group B TIHGG may show vulnerability to these DNA-

damaging and anti-mitotic agents because of its DNA repair pathway deficiencies, as cells with

substantial drug-induced DNA damage prevented from completing mitosis would be expected to

undergo cell death associated with mitotic catastrophe.[40] Interestingly, in the *in vitro* drug screen, the

PARP2 inhibitor olaparib enhanced proliferation compared to vehicle in the Group B cell line (Sup. Table

21). Where DNA damage surveillance is intact, PARP2 inhibition is ineffective in inducing apoptosis and

instead induces a choice of repair pathway, favoring error-prone, non-homologous end resection over

homology-directed repair[41,42]. Thus, in Group B TIHGG, in which homology-directed repair pathways are

already deficient, PARP2 inhibition may enhance, rather than retard, proliferation by favoring non-

homologous resection.

Proteasome inhibitors, which have not been studied beyond preclinical phases in pHGG, were

effective *in vitro* in the Group B cell line, possibly through a mechanism involving inhibition of NF-κB-

mediated inflammatory pathways. Clinically, proteasome inhibition as a therapeutic target is established

in multiple myeloma and mantle cell lymphoma; however, its effectiveness is limited by acquired

resistance[43]. This experience suggests that the efficacy of proteasome inhibition for TIHGG may depend

upon identification of effective combination therapies. An interesting possibility for inhibiting the NF-κB

pathway in conjunction with proteasome inhibition is through inhibition of MDA-9/Syntenin, which has

shown promise in preclinical studies in reducing the invasiveness of radiation-induced GBM in adults.[44]

Targeting the MAPK pathway may also be an effective strategy given the frequent *NF1* mutations in

TIHGG; MEK inhibition is a therapeutic strategy of interest in HGG and was effective (trametinib) in both

TIHGG derived cell lines[45,46].

Strengths of the current study include the size and multi-institutional nature of the cohort, as

well as the comprehensive, orthogonal, unbiased assays performed. Recent characterization of large

cohorts of *de novo* HGG provide basis for comparing TIHGG to the HGG landscape in a way that has not

previously been possible. We also characterized two primary human TIHGG cell lines, allowing for

interrogation of therapeutic vulnerabilities. Weaknesses include the heterogeneity of the molecular

assays performed across the cohort samples; this has been partially ameliorated through the

performance of separate assays and analyses at only one institution per modality in order to standardize

results. In addition, *in vivo* models of TIHGG are still in development, and should be the focus of future

work. With cell lines available and common molecular alterations better understood, both patient-

derived xenograft and genetically engineered mouse models are possible. Our study demonstrates

similarities and differences between TIHGG and pHGG, as well as both homogeneity and heterogeneity

within TIHGG. Therefore, more efficacious treatment regimens may include a common backbone

incorporating some aspects of pHGG treatment, in combination with targeted treatments directed at

specific molecular alterations and susceptibilities characterizing TIHGG subgroups and individual patient

tumors.

Methods

TIHGG case review

Fifty-four cases were reviewed across multiple organizations (Children's Hospital Colorado

(CHCO), St. Jude Children's Research Hospital, Childhood Cancer Survivors' Study (CCSS), the University

of Hamburg, and the University of Florida) from 1981 to 2015 to determine cohort eligibility. Initial

query of the CCSS institutional tumor tissue bank was based on the history of prior radiation with

subsequent development of HGG. Patients were enrolled on Institutional Review Board-approved

protocols for the harvest and study of tissue for research, including Colorado Multi-Institutional Review

Board (COMIRB 95-500) and SJCRH IRB Number: Pro00007403; Mnemonic: XPD17-029; Reference

Number: 001628. After clinical review, 7 cases were judged to be recurrent primary pHGG, two cases

were recurrent primary ependymoma, one case was found to be a recurrent vs. malignant

transformation of a juvenile pilocytic astrocytoma, and one case was found to be a recurrent

glioneuronal tumor. Tissue was not available for 6 cases, and tissue was of insufficient quality or

quantity in 2 cases (Sup. Fig. 11).

We used a modified version of Cahan's criteria to determine eligibility for tumors induced by

radiation[47]. All radiation-induced tumors arose within the initial irradiated field. Although Cahan's

criteria specify that TIHGG must have a histologically proven difference between the initial and

subsequent tumors, 7 cases had an initial diagnosis of glial origin (1 anaplastic astrocytoma, 2

astrocytomas, 3 ependymomas, and 1 ganglioglioma). The two cases with astrocytoma both presented

as low-grade pilocytic astrocytomas arising in the optic tract. Both cases were treated with radiotherapy

after progressive visual loss following a failed trial of chemotherapy. The resultant TIHGGs originated in

the posterior fossa and occipital lobe, respectively, both within a region of normal but previously

irradiated brain.

We also included tumors arising in two patients with germline predisposition syndromes in

which clinical evidence supported the diagnosis of induction by radiation. One patient with known

mismatch repair deficiency and germline *PMS2* mutation developed a primary anaplastic astrocytoma

(AA) localized to the right frontal lobe at the age of 14 years and was managed with a gross total

resection followed by erlotinib and adjuvant radiation as a part of SJHG04 (NCT00124657). Following 9.5

years of controlled disease, a new tumor was discovered centered in the left occipital lobe. The area was

judged to be of sufficient latency to be designated a TIHGG. After review of the clinical history,

radiographic imaging, initial radiotherapy plan, and pathology, it was felt that the subsequent GBM was

more likely to be treatment induced, as it arose in the prior radiotherapy field, differed from the initial

AA, arose from a region of normal brain parenchyma (except for previous irradiation), and occurred with

significant delay, making a late recurrence of AA exceedingly unlikely. The second patient had a

mismatch repair deficiency arising from a heterozygous loss of *MSH*. Upon presentation with

Philadelphia chromosome-positive ALL at age 9 years, the patient received a bone marrow transplant

and total body irradiation (12 Gy). At age 20 years, the patient developed a right frontal anaplastic

oligodendroglioma (WHO grade III). The clinical diagnosis of a radiation-induced tumor was based upon

the occurrence of the oligodendroglioma in the radiation field and the likelihood that radiation-induced

DNA damage played a significant role in the tumor formation.

One tumor clinically diagnosed as being induced by prior exposure to chemotherapeutic agents

was also included in the TIHGG cohort (Sup. Table 1)[47]. Details regarding the clinical history of all

reviewed and included cases are in Sup. Fig. 1.

TIHGG material available for analyses

Deidentified tumor tissue specimens from frozen or formalin-fixed paraffin embedded (FFPE)

sections and frozen patient blood samples were processed following appropriate institutional review

board approval (Fig. 1E). Whole-genome methylation analysis was completed in 31 cases. RNA was of

sufficient quality for RNA-seq analysis in 14 cases. We conducted whole-genome sequencing on twelve

frozen tumor and matched blood samples (to obtain germline genomic information) and whole-exome

sequencing on 5 additional FFPE tumor samples[48].

Voxel-wise lesion mapping of TIHGG

T1 images were registered to the Montreal Neurological Institute template[49] by using ANTsR[50]

and analyzed with voxel-based lesion symptom mapping (VLSM)[51] to assess similarity between statistical

maps by calculating the correlation between t-scores, treating lesion voxels as subjects.

Patient-level statistical analysis

Patient- and sample-level statistical analyses were performed using Rstudio Version 1.1.463.

Packages used for the presented analyses include "survminer", "ggplot2", "survival", and "networkD3".

Continuous data are described using non-parametric measures of central tendency and are tested

across strata using the Wilcoxon-Mann-Whitney test. Frequency data across groups were evaluated

using either the Fisher's exact test or Chi-square test. Time to event endpoints are summarized using the

Kaplan Meier estimator. Differences in time to event strata are compared using the log-rank test.

Methylation array processing

Tumor DNA was extracted from FFPE material by using the Maxwell16 FFPE Plus LEV purification

kit and the Maxwell 16 instrument (Promega, Madison, WI) according to the manufacturer's

instructions. Extracted DNA from FFPE tissue underwent quality control assessment using the Illumina

Infinium FFPE QC Assay kit for qPCR. The Delta Cq values for all samples were less than four. DNA

concentration was assessed using PicoGreen. At least 300 ng of DNA was used per sample for the

subsequent bisulfite conversion with the Zymo EZ-96 DNA Methylation kit. After the bisulfite

conversion, the Infinium HD FFPE Restoration kit was used to restore degraded FFPE DNA to a state that

is amplifiable by the Infinium HD FFPE methylation whole-genome amplification kit. Restored DNA was

then plate-purified (with the Zymo ZR-96 DNA Clean & Concentrator-5), amplified, fragmented,

precipitated, re-suspended, and hybridized to an Illumina Infinium Methylation EPIC 850K BeadChip

array for 22 hours and 30 minutes (by using the Illumina Infinium Methylation EPIC assay kit). Following

hybridization, the arrays were manually disassembled and washed. The subsequent X-Staining of the

array features was processed on a Tecan Freedom EVO robotics system. The arrays were then manually

coated and imaged using an Illumina iScan system with autoloader.

DNA methylation data analysis was performed using the open source statistical programming

language R (R Core Team, 2016). Raw data files generated by the iScan array scanner were read and

preprocessed using minfi Bioconductor package[52]. With the minfi package, the same preprocessing steps

as in Illumina's Genomestudio software were performed. In addition, the following filtering criteria were

applied: removal of probes targeting the X and Y chromosomes, removal of probes containing -

nucleotide polymorphism (dbSNP132 Common) within five base pairs of and including the targeted CpG-

site, and removal of probes not mapping uniquely to the human reference genome (hg19), allowing for

one mismatch. In total, 394,848 common probes of Illumina 450K and EPIC arrays were kept for

clustering analysis.

Statistical analysis of DNA methylation

To determine the subgroup affiliation of our TIHGG samples, we used the reference DNA

methylation cohort published by Capper *et al*. and available from the gene expression omnibus

(GSE90496)[12]. TIHGG samples were combined with the 2,801 reference IDATs containing CNS tumors

and control brain tissues for unsupervised hierarchical clustering, as previously described[12]. In brief, the

32,000 most variable methylated CpG probes measured by standard deviation across combined samples

were selected. Pearson correlation was calculated as the distance measured between samples, and the

unsupervised hierarchical clustering was performed by the average linkage agglomeration method. The

probe-level beta values were also analyzed using t-stochastic neighbor embedding (t-SNE)[53]. Hierarchical

clustering and t-SNE analyses were repeated by using the top 12,000 differentially methylated probes

against a reduced reference set of 666 tumors representing 16 different previously described

methylation classes (Sup. Fig. 3, Sup. Table 7)[12]. The reduced set contained normal control methylation

classes, high-grade diffuse astrocytic tumors, high-grade neuroepithelial tumor with MN1 alteration,

pleomorphic xanthoastrocytoma, and anaplastic pilocytic astrocytoma. Supervised analysis was

performed using the random forest DNA methylation class prediction algorithm (V11b2) by uploading

raw IDAT files to www.molecularneuropathology.org[12].

Detection of copy-number alterations with methylation array data

Copy-number variation was analyzed with Illumina methylation arrays using the conumee

Bioconductor package in R using default settings. The combined intensities of all available CpG probes

were normalized against control samples from normal brain tissue using a linear regression approach.

Platform-matched normal brain tissue controls were used to reference against EPIC 850k array or 450k

array tumor samples. Mean segment value of −0.2 and 0.2 were used as thresholds for losses and gains,

respectively. Copy-number plots were manually examined for selected copy-number alterations. When

copy-number information was also available from sequencing data, the two results were compared and

adjudicated. The adjudicated results are shown in Fig. 5 for selected recurrent somatic alterations.

RNA-seq analysis

Library preparation was performed using the TruSeq Library Preparation Kit v2 (Agilent);

directional mRNA sequencing was performed at the University of Colorado Anschutz Medical Campus

Genomics and Microarray Core on a HiSeq 2500 sequencing system (Illumina) using single pass 125 bp

reads (1x125) and approximately 50 million reads per sample. The resulting data were mapped to the

human genome (hg19) by gSNAP, expression (FPKM) was derived by Cufflinks, and differential

expression was analyzed with ANOVA in R. Output files contained read-depth data and FPKM expression

levels for each sample and, where gene expression levels were compared between groups of samples,

the ratio of expression in $\log_2$ format and a p-value for each gene. CICERO was used to detect fusion

genes on the RNA-seq data[54].

Analysis of transcriptomic data

Using micro-array data (Affymetrix Human Genome U133 Plus 2.0 Array) previously acquired

from tumor samples of patients treated at Children's Hospital Colorado, we compared patterns of gene

expression in 14 TIHGG samples (13 radiation-induced and one chemotherapy-induced) to those of a

cohort of non-treatment-induced tumors consisting of 24 primary pHGG, 4 infant HGG, and 14 adult

HGG. We performed clustering analysis using the t-SNE method available in the RTSNE package and

confirmed the TIHGG subgrouping obtained through t-SNE by using non-negative matrix factorization

(NMF)[53,55]. Principal component analysis with 30 initial dimensions preceded the t-SNE analysis, in which

we empirically selected perplexity of 3 and 50,000 iterations as providing optimal results. For the NMF

analysis, we utilized the identical micro-array dataset used in the t-SNE analysis, using k (number of

clusters) of 2–5. We performed Metascape analysis on the micro-array data using as input a list of 2,162

genes differentially expressed (p<0.01) between the TIHGG and HGG samples, followed by Cytoscape to

identify differentially enriched pathways[56,57]. We performed geneset enrichment analysis (GSEA, Broad

Institute) to identify the direction of enrichment between the HGG and TIHGG groups and GSEA and

Ingenuity Pathway Analysis (IPA, Qiagen) to identify gene expression patterns within the TIHGG

cohort[58,59]. GSEA results were evaluated using the normalized enrichment score (NES), in which

increased expression results in a positive score, and reduced expression in a negative score. Scores were

considered potentially informative from a statistical perspective if the false-discovery rate (FDR) was less

than 0.25.

Whole-genome sequencing

WGS library preparation and sequencing were performed by BGI Americas; 100-fold mean

coverage data were acquired. BGI performed initial quality testing of the sample, including

concentration and sample integrity/purity. Concentration was detected by fluorometer or microplate

reader (*e.g.* Qubit Fluorometer, Invitrogen). Sample integrity and purity were detected by agarose gel

electrophoresis (agarose gel concentration: 1%, voltage: 150 V, electrophoresis time: 40 min). Following

quality confirmation, 1 µg genomic DNA was randomly fragmented by Covaris. The fragmented genomic

DNA was selected by Agencourt AMPure XP-Medium kit to an average size of 200-400 bp. Fragments

were end repaired and then 3' adenylated. Adaptors were ligated to the ends of these 3' adenylated

fragments to facilitate amplification by PCR. PCR products were purified using the Agencourt AMPure

XP-Medium kit. The double-stranded PCR products were heat denatured and circularized by the splint

oligonucleotide sequence. The single-strand circular DNA (ssCir DNA) was formed as the final library

qualified via a quality control procedure. The qualified libraries were sequenced by BGISEQ-500. Briefly,

each ssCir DNA molecule was formed into a DNA nanoball (DNB) containing more than 300 copies

through rolling-cycle replication. The DNBs were loaded into the patterned nanoarray using high density

DNA nanochip technology. Finally, pair-end 100-bp reads were obtained by combinatorial Probe-Anchor

Synthesis (cPAS). WGS mapping, quality control, and somatic mutation (SNV and INDEL) calling and

classification have been described previously[60,61].[60,61]. The tumor and germline reads were mapped to

GRCh37-lite and Tier1 mutations (i.e. coding somatic mutations). CNA regions were identified using

CONSERTING on the paired tumor-germline WGS samples with Log Ratio > 0.25 or < -0.25 reported[62].[62].

Structural variants (SVs) of tumor WGS samples were identified using CREST based on soft-clipped reads

evidence[63].[63]. The SVs that have discordant reads support in tumor sample but not in paired germline

sample were reported.

Comparison of mutation frequency in de novo pHGG and TIHGG by expression group

We analyzed mutation load, small scale variants, and structural variants in the TIHGG tumor

samples and matched blood samples. We noticed, based upon our initial review of the genome

sequencing data, that several tumor samples appeared to have anomalously large numbers of mutations

and decided to perform a mutation load analysis based upon that observation. We prepared vcf files

containing QUAL-filtered (threshold of 50) and unfiltered putatively damaging variants. Germline

mutation load was defined as the total number of mutations per sample in the QUAL filtered dataset.

Somatic mutation load was determined from tumor samples without QUAL-filtering. The mutation calls

were filtered to require at least 4 reads per called variant in order to constitute a mutation. This

approach was deemed reasonable because of the likely heterogeneity present in the tumor samples

such that a clone constituting a fraction of the total sample could have an allelic-level mutation that

would be detected in only a small fraction of the overall reads at a particular locus, resulting in a low

QUAL score. Data were sensitivity tested to determine the thresholds for fold requirement and also to

determine whether a QUAL threshold should be imposed. The tumor sample from the patient with

Lynch syndrome was not included in this analysis because of the existence of a known DNA repair defect

unrelated to the therapeutic radiation treatment.

Whole-exome sequencing

Human genomic libraries were generated using the SureSelectXT kit specific for the Illumina

HiSeq instrument (Catalog No. G9611B; Agilent Technologies, Santa Clara, CA), followed by exome

enrichment using the SureSelectXT Human All Exon V6+COSMIC bait set (Catalog No. 5190-9307). The

resulting exome-enriched libraries were then sequenced by the Genome Sequencing Facility on a HiSeq

4000 (Illumina). WES mapping and quality assessment have been described previously[61].[60,61]. The tumor

reads were mapped to GRCh37-lite, and variants were called by Bambino[18] and annotated by Medal

Ceremony as "Gold", "Silver", "Bronze" or "unknown"[64]. "Gold" mutations and variants matching with

COSMIC database were retained[65]. For other coding variants, those that were low-frequency (<0.001) or

absent in ExAC/1000Genome/NHLBI databases were reported if the variant was supported with at least

5 mutant alleles and at least 30% VAF[66,67]. The significance of mutated genes was assessed using the

Significantly Mutated Gene test[68]. Mutation frequency and composition were analyzed by comparing

the number and type of mutations across primary and TIHGG samples. The absolute number of

mutations, and frequencies of base pair substitutions in SNVs were compared across TIHGG and primary

HGG using a *t*-test and Chi-square test, respectively (Fig. 7C, Sup. Table 7, and Sup. Table 8). The

frequencies of commonly altered genes in TIHGG and primary HGG were compared by using Fisher's

exact test and are listed in (Sup. Table 7).

Evaluation of chromothripsis events and structure prediction of eccDNA

730   The presence or absence of chromothripsis was evaluated in twelve samples' WGS data (Sup.

Table 8). Four key criteria were used to infer chromothripsis as described by Korbel *et al*; (oscillating

CNA regions, clustering of breakpoints, randomness of DNA fragment joins, and randomness of DNA

fragment order)[69]. Chromothripsis was called when at least two criteria were satisfied and further

evaluated by manual review. The eccDNA structures were constructed following the procedures

described in Xu *et al.* by identifying the cyclic graphs composed of highly amplified CNA segments and

their associated SVs[16].

FISH analysis

739   Dual-color FISH was performed on 4-μm thick paraffin-embedded tissue sections. Probes were

derived from BAC clones (BACPAC Resources, Oakland, CA) and labeled with either AlexaFluor-488 or

AlexaFluor-555 fluorochromes. BAC clones were used to construct probes for the following genes:

*PDGFRA (*laboratory-developed probe (RP11-231C18 & 601I15); 4p control (CTD-2057N12 & CTD-

2588A19), and *CDK4* (Empire Genomics, Williamsville, New York, Cat# CDK4-CHR12-20-ORGR). Probes

were co-denatured with the target cells on a slide moat at 90°C for 12 minutes. The slides were

incubated overnight at 37°C on a slide moat and then washed in 4M Urea/2xSSC at 25°C for 1

minute. Nuclei were counterstained with DAPI (200ng/mL) (Vector Labs) for viewing on an Olympus

BX51 fluorescence microscope equipped with a 100-watt mercury lamp; FITC, rhodamine, and DAPI

filters; 100X PlanApo (1.40) oil objective; and a Jai CV digital camera. Images were captured and

processed using the Cytovision software from Leica Biosystems (Richmond, IL)[70].

In silico *and* in vitro *drug screening and validations*

In silico *screen.* Micro-array data for 12 TIHGG samples and 37 *de novo* GBM samples (23

pediatric, 14 adult) were compared to target genes of several drug classes. A ratio of mean expression

and p-value for each target gene in the TIHGG and *de novo* GBM samples were computed, and an overall

ratio and p-value of all target genes used for each therapeutic classification for each target gene were

computed and reported (Supp. Table 20).

In vitro *drug screen and validation:* For the drug screen performed in TIHGG cell lines MAF-145

and MAF-496, we utilized the Approved Oncology Drugs Set VI (National Cancer Institute), comprising

129 drugs, supplemented by selinexor (Karyopharm Therapeutics) and AZD2014 (Astra Zeneca). The

complete list of drugs included in the screen is included in Sup. Table 21. Cells were plated at a density

of 5,000 cells per well in in 90 μL medium in a 96-well treated cell culture plate (Corning #3595) and

allowed to gain adhesion overnight. Drugs were applied in 10 μL of medium/1% DMSO at a

concentration of 10 μM, resulting in a final concentration of 1 μM and 0.1% DMSO. Cells were incubated

in drug for 5 days. DMSO (0.1%) was utilized as a control. Cell viability was assayed following 5 days of

treatment using incubation with tritiated thymidine and quantification using a scintillation counter.

Results were collected as counts/min and converted to survival using the formula (sample –

medum)/(DMSO – medium) where "sample" is the scintillation count for each drug-treated sample,

"medium" is the scintillation count for a well containing medium only, and "DMSO" is the scintillation

count for a three-well average of cells treated with 0.1% DMSO only (Sup. Table 21). The drug screen

was conducted twice in MAF-145 cells and once in MAF-496 due to limitations on cell availability.

Validation tests of single drugs were conducted using drug concentrations ranging from 0.316 nM to 10

μM in half-log$_{10}$ increments. Cells were plated as described above and incubated in drug for 120 hours.

Three biological replicates were used for each drug concentration. Results were assessed using CellTiter

96 Aqueous One Solution Cell Proliferation Assay (Promega), according to manufacturer instructions.

Survival was computed as above. IC$_{50}$ values were calculated using a variable slope four parameter non-

linear model with maximum survival constrained at 100% (Prism 7, Graphpad).

Immunofluorescence staining of in vitro samples: Cells were plated at a density of 20,000 cells

per well in BioCoat chamber slides coated with poly-D-lysine or poly-D-lysine and laminin (Corning) and

allowed approximately 24-48 hours in which to develop adhesion before subjecting them to

experimental conditions. Cells to be stained were fixed for 20 minutes in formaldehyde diluted to 3.7%

in PBS (Sigma), permeabilized in 0.1% Triton-X in PBS for 10 minutes, and blocked for 45 minutes in 4%

BSA in PBS supplemented with 0.05% Triton-X. Cells were incubated in primary antibody to the p65

subunit of NF-κB (Cell Signaling, #6956, 1:400) diluted with 4% BSA (in PBS and 0.05% Triton-X) for 1h at

room temperature or overnight at 4°C. Following multiple rinses with PBS, cells were incubated in a

secondary fluorophore (Alexa Fluor 488) for 1 hour, rinsed, and coverslips were then adhered using

ProLong Gold antifade reagent with DAPI (Invitrogen). Confocal imaging was performed at 400x using

405 nM (DAPI) and 488 nM (Alexa Fluor 488) lasers on a 3I Marianas imaging system (Intelligent Imaging

Innovations). Images were captured using an Evolve 16-bit EMCCD camera (Photometrics).

Figures (8 total)

Figure 1. TIHGG cohort characteristics.

A) Venn diagram illustrating the number of cases with each type of historical data/molecular analysis. B)

Radiotherapy field type used for initial cancer. C) Time to diagnosis of TIHGG from initial cancer

diagnosis. D) Time to death following TIHGG diagnosis. *Two patients survived beyond 35 months but

were deceased at study closure. E) Anatomic location of TIHGG. Color scale indicates the number of

cases anatomically overlapping at each point in space.

Figure 2. Methylation-based classification of TIHGG.

A) Circular dendrogram indicating location of TIHGG (black bars) relative to diagnoses. B) Localization of

TIHGG relative to other CNS cancers in t-distributed stochastic neighboring (t-SNE) space. C)

Dendrogram and heatmap of TIHGG cases clustered against other pediatric brain tumor entities. A-IDH=

Astrocytoma, subclass IDH-mutant; A-IDH-HG= High-Grade Astrocytoma, subclass IDH-mutant; ANA PA =

Anaplastic Pilocytic Astrocytoma, CONTR-CEBM= Control Cerebellum; CONTR-HEMI= Control Cerebral

Cortex; DMG-K27 = Diffuse Midline Glioma, H3 K27M-mutant; GBM-G34= Glioblastoma, subclass H3.3

p.G34R-mutant; GBM-MES= Glioblastoma, subclass mesenchymal; GBM-MID = Glioblastoma, IDH-Wild

type, subclass midline; GBM-MYCN= Glioblastoma, subclass MYCN-amplified; GBM-RTK-I = Adult

Glioblastoma, subclass RTK I; GBM-RTK-II= Adult Glioblastoma, subclass RTK II; GBM-RTK-II = Adult

Glioblastoma, subclass RTK III; HGNET-MN1 = High Grade Neuroepithelial Tumor with MN1 alteration;

IHG= Infantile High Grade Glioma; pedRTK I= Pediatric Glioblastoma, subclass RTK I; pedRTK II = Pediatric

Glioblastoma, subclass RTK II; PXA = Pleomorphic Xanthoastrocytoma; TIHGG = Treatment-Induced

High-Grade Glioma.

Figure 3. Frequent copy number alterations in TIHGG.

A) Copy-number frequency plot. B) Expression-based confirmation of high frequency copy number

alterations in *PDGFRα, CDKN2A,* and *BCOR;* mean and standard error of the mean (SEM) shown. C)

Frequency of copy number gain and loss across all TIHGG; boxes show median and 1st/3rd quartiles, with

whiskers representing range limited to 1.5x the interquartile range from the box edge.

Figure 4. Frequency of chromothripsis and resultant eccDNA structures, oncogene incorporation and

FISH analysis in Case_29.

A) Frequency of chromothripsis in primary non-brainstem high-grade glioma (NBS-HGG), primary DIPG,

TIHGG, and combined NBS-HGG and DIPG. B) Circos plot illustrating the intra- and inter-chromosomal

rearrangements of chromosomes 4 and 14, resulting in dm1, and intra-chromosomal rearrangements in

chromosome 12 resulting in dm2. Byproduct of chromothripsis events between chromosomes 4 and 12

(dm1) and intra-chromosomal rearrangements in chromosome 14 (dm2). The innermost circle shows

the highly amplified CNA segments, as evidenced by their extremely high coverage in the tumor sample

(blue circle) compared to that in the paired germline sample (green circle). The outermost circle shows

the involved chromosomes. C) Two-color FISH experiment from Case_29 shows *PDGFRα* (upper panel)

and *CDK4* amplification (lower panel) with diffuse punctate staining relative to the control centromeric

DNA in each cell, suggesting the presence of multiple double-minutes. D) The predicted structures of the

eccDNA dm1 and dm2 showing the directions of each of the joined CNA segments and their break point

positions. "Bridge reads support" means SVs identified through common discordant reads support

between two segment boundaries as described in Xu *et al*[16].

Figure 5. Recurrent molecular alterations in TIHGG.

Oncoprint describing the clinical characteristics, histopathological features, methylation profile, tier1

mutations, genes affected by copy-number gain/loss, and fusion genes. ANA PA = Anaplastic Pilocytic

Astrocytoma; CONTR-CEBM = Control Cerebellum; DMG-K27 = Diffuse Midline Glioma H3.3 K27M; GBM-

MID = Glioblastoma IDH-wild type, subclass midline; HGNET-MN1 = High Grade Neuroepithelial Tumor

with MN1 alteration; PXA = Pleomorphic Xanthoastrocytoma. Sequencing/array platforms for each case

are also shown on the bottom row. The second sample for Case 2 is shown as Case 2B while the second

sample for Case 14 and Case 12 are not shown but are annotated in Sup. Table 5.

Figure 6. Gene expression profiling of TIHGG versus *de novo* GBM.

A) Clustering of TIHGGs with microarray-based transcriptomic data (N=14) versus *de novo* pediatric GBM

(N=24), infant GBM (N=4) and adult GBM (N=14) using t-SNE analysis. B) Methylation clustering showing

locations of TIHGGs included in panel A; all TIHGGs included in panel A cluster in pedRTK I; methylation

and transcriptomic clusters overlap with the exception of two samples. C) Cytoscape analysis based on

gene ontology (GO) genesets show cellular pathways and processes that differ based upon gene

expression between TIHGG and *de novo* GBM (from panel A); color scale represents p-value, and the

comparisons are non-directional between sample sets. D) geneset enrichment analysis (GSEA) using GO

genesets identifies differences in gene expression between TIHGG and *de novo* GBM (from panel A);

number scale represents log of the false discovery rate (with FDR = 0 set to a value of 10 for purposes of

plotting); colors represent the ratio of GSEA normalized enrichment score (NES) of TIHGG/*de novo* GBM.

Figure 7. Gene expression and genetic profiling of Group A versus Group B TIHGG.

A) GSEA results showing GO differences organized by category between Group A (blue) and Group B

(red) TIHGG tumors; horizontal axis is NES for the comparison of TIHGG Group B vs. TIHGG Group A (NES

> 0 means enrichment in Group B and NES < 0 means enrichment in Group A) (FDR values for

comparisons are listed in Sup. Table 18); mean and SEM shown. B) Frequency of 1p loss and 1q gain

based upon methylation profiling in TIHGG gene expression groups A and B; note that one TIHGG tumor

for which there was transcriptomic data lacked a DNA sample and was not included in the methylation

profiling; mean and SEM shown. C) For TIHGG samples with germline somatic genome sequencing data

and transcriptomic data, germline variant load was identical between subgroups, but somatic load was

approximately 9-fold greater in Group B (p<0.002); TIHGG sample with *MSH2* mismatch repair defect

was excluded from this analysis. D) GSEA results for DNA repair pathways for Group B versus Group A

germline (based on blood) and tumor samples; negative numbers represent depletion in Group B;

Individual genes from the GO_DNA_REPAIR geneset with the most significant differences in expression

between Group B and Group A are shown by decreasing p-value (top to bottom); colors represent the

ratio of mean expression (ER) of Group B/Group A.

Figure 8. TIHGG preclinical drug screen.

A) *In silico*-predicted response of TIHGG vs. *de novo* GBM tumors to drug classes based upon gene

expression data. B) *In vitro* drug screening results by drug class for TIHGG cell lines MAF-145 (gene

expression Group B) and MAF-496 (gene expression Group A) showing reduced survival by drug class

relative to mean survival; screen performed using FDA-approved anti-cancer agents at 1 μM

concentration. C) *In vitro* drug screen results in TIHGG cell lines using 1 μM concentration; Group A

(MAF-496) cell line response is plotted on the x-axis and Group B cell line (MAF-145) on the y-axis. Dot

colors correspond to drug class as shown in the legend. D) *In vitro* validation results in TIHGG cell lines

MAF-145 and MAF-496 for candidate drugs identified through *in vitro* drug screen (panels B, D)

Supplementary Figure 8. Recurrent molecular alterations in TIHGG relative to *de novo* pHGG (HERBY

dataset). Oncoprint describing the clinical characteristics, histopathological features, methylation

profile, tier1- mutations, genes affected by copy-number gain/loss, and fusion genes in TIHGG and *de*

novo pHGG.

Supplementary Figure 9. Germline and somatic variants in TIHGG relative to *de novo* pHGG (Wu G

reference cohort) stratified by base transition type. A) Total number of germline, somatic coding, and

somatic non-coding variants per Mb stratified by *de novo* pHGG vs. TIHGG. B) Proportion of base

transitions in somatic non-coding variants per Mb stratified by primary pHGG vs. TIHGG. The relative

frequency of base transitions in germline coding SNVs. C) Somatic coding SNVs and somatic non-

noncoding SNVs. For violin plots, box shows median and interquartile range, whiskers show 95%

confidence interval, total length of violin represents range, and width of violin shows frequency.

Supplementary Figure 10. TIHGG *in vitro* sensitivities by expression profile. A) *In vitro* drug screen

consort diagram; primary cell lines derived from TIHGG tumors were MAF-145 (gene expression TIHGG

Group B) and MAF-496 (gene expression TIHGG Group A). B) MTS assay results for indicated drugs in

MAF-496 (IC$_{50}$ values listed in Fig. 8D). C) MTS results for indicated drugs in MAF-145 (IC$_{50}$ values listed in

Fig. 8D). D) MTS results for trametinib. E) MTS results for sunitinib. F) MTS assay results for proteasome

inhibitors bortezomib and carfilzomib in MAF-145 and MAF-496 cell lines. G) MTS results for marizomib.

H) NF-κB levels in Group A (MAF-496) and B (MAF-145) TIHGG cell lines treated with vehicle and

bortezomib. I) Immunofluorescence of NF-κB nuclear localization in vehicle bortezomib-treated Group A

(MAF-496) and Group B (MAF-145) TIHGG cell lines. Mean and SEM shown.

Supplementary Figure 11. Study consort diagram. CCSS = Childhood Cancer Survivor Study, SJCRH = St.

Jude Children's Research Hospital, U. Colorado = University of Colorado, WGS = whole-genome

sequencing, WES = whole-exome sequencing, RNA-seq = RNA sequencing.

Supplementary Tables (21 total)

Supplementary Table 1. Patient, treatment, initial, and TIHGG disease characteristics.

Supplementary Table 2. TIHGG treatment and outcomes.

Supplementary Table 3. Methylation-based classification probabilities for the reference dataset for

methylation-based classification (Korshunov A and Capper D).

Supplementary Table 4. Methylation-based classification probabilities.

Supplementary Table 5. Methylation-based classification summary.

Supplementary Table 6. Comparison of TIHGG and *de novo* pHGG methylation class.

Supplementary Table 7. Comparison of TIHGG and *de novo* pHGG from the HERBY cohort.

Supplementary Table 8. Chromothripsis evaluation criteria.

Supplementary Table 9. Tier 1 sequence mutations identified from 5 non-hypermutator WES cases

without matched germline data.

Supplementary Table 10. Tier 1 sequence mutations identified from 9 non-hypermutator WGS cases

with matched germline data.

Supplementary Table 11. TIHGG fusion genes in RNA-seq samples.

Supplementary Table 12. TIHGG structural variations in 10 WGS cases.

Supplementary Table 13. TIHGG copy-number alteration segments in 10 WGS cases.

Supplementary Table 14. Methylation-based classification probabilities for the HERBY dataset

Supplementary Table 15. Germline variants.

Supplementary Table 16. Cytoscape/Metascape results

Supplementary Table 17. GSEA results in TIHGG vs. *de novo* GBM for GO, with all genesets

Supplementary Table 18. GSEA results by category as shown in Fig. 7A.

Supplementary Table 19. Expression Ratio (ER) and p-value (Group B TIHGGs/Group A TIHGGs) for

selected genes from GO_DNA_REPAIR geneset.

Supplementary Table 20. *In silico* drug screening results.

Supplementary Table 21. *In vitro* drug screening results.

References

1.   Armstrong, G.T. *et al.* Long-term outcomes among adult survivors of childhood central nervous
system malignancies in the Childhood Cancer Survivor Study. *J Natl Cancer Inst* **101**, 946-58
(2009).
2.   Armstrong, G.T. *et al.* Late mortality among 5-year survivors of childhood cancer: a summary
from the Childhood Cancer Survivor Study. *J Clin Oncol* **27**, 2328-38 (2009).
3.   Maluf, F.C., DeAngelis, L.M., Raizer, J.J. & Abrey, L.E. High-grade gliomas in patients with prior
systemic malignancies. *Cancer* **94**, 3219-24 (2002).
4.   Carret, A.S. *et al.* Outcome of secondary high-grade glioma in children previously treated for a
malignant condition: a study of the Canadian Pediatric Brain Tumour Consortium. *Radiother*
Oncol **81**, 33-8 (2006).
5.   Paugh, B.S. *et al.* Integrated molecular genetic profiling of pediatric high-grade gliomas reveals
key differences with the adult disease. *J Clin Oncol* **28**, 3061-8 (2010).
6.   Behjati, S. *et al.* Mutational signatures of ionizing radiation in second malignancies. *Nat Commun*
7, 12605 (2016).
7.   Lopez, G.Y. *et al.* The genetic landscape of gliomas arising after therapeutic radiation. *Acta*
Neuropathol **137**, 139-150 (2019).
8.   Sholl, L.M., Barletta, J.A. & Hornick, J.L. Radiation-associated neoplasia: clinical, pathological and
genomic correlates. *Histopathology* **70**, 70-80 (2017).
9.   Leeman-Neill, R.J. *et al.* ETV6-NTRK3 is a common chromosomal rearrangement in radiation-
associated thyroid cancer. *Cancer* **120**, 799-807 (2014).
10.   Leeman-Neill, R.J. *et al.* RET/PTC and PAX8/PPARgamma chromosomal rearrangements in post-
Chernobyl thyroid cancer and their association with iodine-131 radiation dose and other
characteristics. *Cancer* **119**, 1792-9 (2013).
11.   Korshunov, A. *et al.* H3-/IDH-wild type pediatric glioblastoma is comprised of molecularly and
prognostically distinct subtypes with associated oncogenic drivers. *Acta Neuropathol* **134**, 507-
516 (2017).
12.   Capper, D. *et al.* DNA methylation-based classification of central nervous system tumours.
Nature **555**, 469-474 (2018).
13.   Sturm, D. *et al.* Paediatric and adult glioblastoma: multiform (epi)genomic culprits emerge. *Nat*
Rev Cancer **14**, 92-107 (2014).
14.   Mackay, A. *et al.* Molecular, Pathological, Radiological, and Immune Profiling of Non-brainstem
Pediatric High-Grade Glioma from the HERBY Phase II Randomized Trial. *Cancer Cell* **33**, 829-842
e5 (2018).
15.   Wu, G. *et al.* The genomic landscape of diffuse intrinsic pontine glioma and pediatric non-
brainstem high-grade glioma. *Nat Genet* **46**, 444-450 (2014).
16.   Xu, K. *et al.* Structure and evolution of double minutes in diagnosis and relapse brain tumors.
Acta Neuropathol **137**, 123-137 (2019).
17.   Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology
Consortium.
18.   Edmonson, M.N. *et al.* Bambino: a variant detector and alignment viewer for next-generation
sequencing data in the SAM/BAM format. *Bioinformatics* **27**, 865-6 (2011).
19.   Gits, H.C. *et al.* Medulloblastoma therapy generates risk of a poorly-prognostic H3 wild-type
subgroup of diffuse intrinsic pontine glioma: a report from the International DIPG Registry. *Acta*
Neuropathol Commun **6**, 67 (2018).

20.   Mackay, A. *et al.* Integrated Molecular Meta-Analysis of 1,000 Pediatric High-Grade and Diffuse
Intrinsic Pontine Glioma. *Cancer Cell* **32**, 520-537 e5 (2017).

21.   Downing, J.R. *et al.* The Pediatric Cancer Genome Project. *Nat Genet* **44**, 619-22 (2012).

22.   Mistry, M. *et al.* BRAF mutation and CDKN2A deletion define a clinically distinct subgroup of
childhood secondary high-grade glioma. *J Clin Oncol* **33**, 1015-22 (2015).

23.   Collado, M. & Serrano, M. Senescence in tumours: evidence from mice and humans. *Nat Rev*
Cancer **10**, 51-7 (2010).

24.   Lassaletta, A. *et al.* Therapeutic and Prognostic Implications of BRAF V600E in Pediatric Low-
Grade Gliomas. *J Clin Oncol* **35**, 2934-2941 (2017).

25.   Hudler, P. *et al.* Mutations in the hMLH1 gene in Slovenian patients with gastric carcinoma. *Clin*
Genet **65**, 405-11 (2004).

26.   De Ruysscher, D. *et al.* Radiotherapy toxicity. *Nat Rev Dis Primers* **5**, 13 (2019).

27.   Kilday, J.P. *et al.* Copy number gain of 1q25 predicts poor progression-free survival for pediatric
intracranial ependymomas and enables patient risk stratification: a prospective European
clinical trial cohort analysis on behalf of the Children's Cancer Leukaemia Group (CCLG), Societe
Francaise d'Oncologie Pediatrique (SFOP), and International Society for Pediatric Oncology
(SIOP). *Clin Cancer Res* **18**, 2001-11 (2012).

28.   Hing, S. *et al.* Gain of 1q is associated with adverse outcome in favorable histology Wilms'
tumors. *The American journal of pathology* **158**, 393-398 (2001).

29.   Gratias, E.J. *et al.* Association of Chromosome 1q Gain With Inferior Survival in Favorable-
Histology Wilms Tumor: A Report From the Children's Oncology Group. *Journal of Clinical*
Oncology **34**, 3189-3194 (2016).

30.   Hirai, M. *et al.* 1q23 gain is associated with progressive neuroblastoma resistant to aggressive
treatment.

31.   Molenaar, J.J. *et al.* Sequencing of neuroblastoma identifies chromothripsis and defects in
neuritogenesis genes. *Nature* **483**, 589-93 (2012).

32.   Roy, A. *et al.* Recurrent internal tandem duplications of BCOR in clear cell sarcoma of the kidney.
Nat Commun **6**, 8891 (2015).

33.   Olanich, M.E. *et al.* CDK4 Amplification Reduces Sensitivity to CDK4/6 Inhibition in Fusion-
Positive Rhabdomyosarcoma. *Clin Cancer Res* **21**, 4947-59 (2015).

34.   Szerlip, N.J. *et al.* Intratumoral heterogeneity of receptor tyrosine kinases EGFR and PDGFRA
amplification in glioblastoma defines subpopulations with distinct growth factor response. *Proc*
Natl Acad Sci U S A **109**, 3041-6 (2012).

35.   Nathanson, D.A. *et al.* Targeted therapy resistance mediated by dynamic regulation of
extrachromosomal mutant EGFR DNA. *Science* **343**, 72-6 (2014).

36.   Turner, K.M. *et al.* Extrachromosomal oncogene amplification drives tumour evolution and
genetic heterogeneity. *Nature* **543**, 122-125 (2017).

37.   L'Abbate, A. *et al.* Genomic organization and evolution of double minutes/homogeneously
staining regions with MYC amplification in human cancer. *Nucleic Acids Res* **42**, 9131-45 (2014).

38.   Zhang, D., Liu, X., Fan, C. & Chen, J. Novel drugs in pediatric gliomas. *Oncol Lett* **13**, 2881-2885
(2017).

39.   Williams, M.J., Singleton, W.G.B., Lowis, S.P., Malik, K. & Kurian, K.M. Therapeutic Targeting of
Histone Modifications in Adult and Pediatric High-Grade Glioma. *Frontiers in oncology* **7**, 45-45
(2017).

40.   Huang, X., Tran, T., Zhang, L., Hatcher, R. & Zhang, P. DNA damage-induced mitotic catastrophe
is mediated by the Chk1-dependent mitotic exit DNA damage checkpoint. *Proc Natl Acad Sci U S*
A **102**, 1065-70 (2005).

41.   Takagi, M. *et al.* Loss of DNA Damage Response in Neuroblastoma and Utility of a PARP
Inhibitor. *JNCI: Journal of the National Cancer Institute* **109**(2017).

42.   Fouquin, A. *et al.* PARP2 controls double-strand break repair pathway choice by limiting 53BP1
accumulation at DNA damage sites and promoting end-resection. *Nucleic Acids Research* **45**,
12325-12339 (2017).

43.   Acosta-Alvear, D. *et al.* Paradoxical resistance of multiple myeloma to proteasome inhibitors by
decreased levels of 19S proteasomal subunits.

44.   Kegelman, T.P. *et al.* Inhibition of radiation-induced glioblastoma invasion by genetic and
pharmacological targeting of MDA-9/Syntenin. *Proc Natl Acad Sci U S A* **114**, 370-375 (2017).

45.   See, W.L., Tan, I.L., Mukherjee, J., Nicolaides, T. & Pieper, R.O. Sensitivity of glioblastomas to
clinically available MEK inhibitors is defined by neurofibromin 1 deficiency. *Cancer Res* **72**, 3350-
9 (2012).

46.   Sunayama, J. *et al.* Crosstalk between the PI3K/mTOR and MEK/ERK pathways involved in the
maintenance of self-renewal and tumorigenicity of glioblastoma stem-like cells. *Stem Cells* **28**,
1930-9 (2010).

47.   Cahan, W.G., Woodard, H.Q. & et al. Sarcoma arising in irradiated bone; report of 11 cases.
Cancer **1**, 3-29 (1948).

48.   Donson, A.M. *et al.* Unique molecular characteristics of radiation-induced glioblastoma. *J*
Neuropathol Exp Neurol **66**, 740-9 (2007).

49.   Fonov, V. *et al.* Unbiased average age-appropriate atlases for pediatric studies. *Neuroimage* **54**,
313-27 (2011).

50.   Avants, B.B. *et al.* A reproducible evaluation of ANTs similarity metric performance in brain
image registration. *Neuroimage* **54**, 2033-44 (2011).

51.   Bates, E. *et al.* Voxel-based lesion-symptom mapping. *Nat Neurosci* **6**, 448-50 (2003).

52.   Aryee, M.J. *et al.* Minfi: a flexible and comprehensive Bioconductor package for the analysis of
Infinium DNA methylation microarrays. *Bioinformatics* **30**, 1363-9 (2014).

53.   van der Maaten, L. & Hinton, G. Visualizing Data using t-SNE. *Journal of Machine Learning*
Research **9**, 2579--2605 (2008).

54.   Parker, M. *et al.* C11orf95-RELA fusions drive oncogenic NF-kappaB signalling in ependymoma.
Nature **506**, 451-5 (2014).

55.   Brunet, J.-P., Tamayo, P., Golub, T.R. & Mesirov, J.P. Metagenes and molecular pattern discovery
using matrix factorization. *Proceedings of the National Academy of Sciences* **101**, 4164-4169
(2004).

56.   Tripathi, S. *et al.* Meta- and Orthogonal Integration of Influenza "OMICs" Data Defines a Role for
UBR4 in Virus Budding. *Cell Host Microbe* **18**, 723-35 (2015).

57.   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular
interaction networks. *Genome Res* **13**, 2498-504 (2003).

58.   Mootha, V.K. *et al.* PGC-1α-responsive genes involved in oxidative phosphorylation are
coordinately downregulated in human diabetes. *Nature Genetics* **34**, 267 (2003).

59.   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for
interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences*
102, 15545-15550 (2005).

60.   Zhang, J. *et al.* The genetic basis of early T-cell precursor acute lymphoblastic leukaemia. *Nature*
481, 157-63 (2012).

61.   Zhang, J. *et al.* A novel retinoblastoma therapy from genomic and epigenetic analyses. *Nature*
481, 329-34 (2012).

62.   Chen, X. *et al.* CONSERTING: integrating copy-number analysis with structural-variation
detection. *Nat Methods* **12**, 527-30 (2015).

63. Wang, J. *et al.* CREST maps somatic structural variation in cancer genomes with base-pair
resolution. *Nat Methods* **8**, 652-4 (2011).
64. Zhang, J. *et al.* Germline Mutations in Predisposition Genes in Pediatric Cancer. *N Engl J Med*
373, 2336-2346 (2015).
65. Tate, J.G. *et al.* COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Res* **47**,
D941-D947 (2019).
66. Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285-91
(2016).
67. Genomes Project, C. *et al.* A global reference for human genetic variation. *Nature* **526**, 68-74
(2015).
68. Dees, N.D. *et al.* MuSiC: identifying mutational significance in cancer genomes. *Genome Res* **22**,
1589-98 (2012).
69. Korbel, J.O. & Campbell, P.J. Criteria for inference of chromothripsis in cancer genomes. *Cell* **152**,
1226-36 (2013).
70. Ellison, D.W. *et al.* Definition of disease-risk stratification groups in childhood medulloblastoma
using combined clinical, pathologic, and molecular variables. *J Clin Oncol* **29**, 1400-7 (2011).

1110

A

RNAseq 1

WGS

Methylation 3

Clinical

8 2

WES

B

RT FIELDS

TBI 3%

CSI 24%

Unknown 24%

Cranial 18%

Focal Brain 31%

C

Latency of Treatment-Related Gliomas

Cumulative event

1.00

0.75

0.50

0.25

0.00

0   5   10   15   20

Time (Years)

Cumulative number of events

0   3   11   16   16

D

Time to Death from 2nd Cancer Diagnosis

Survival probability

1.00

0.75

0.50

0.25

0.00

0   5   10   15   20   25   30

Time (Months)

Number at risk

27   19   10   7   6   5   3

E

1   9

Figure 1

Figure 2

methylation group

- A-IDH
- A-IDH-HG
- ANA-PA
- CONTR-CEBM
- CONTR-HEMI
- DMG-K27
- GBM-G34
- GBM-MES
- GBM-MID
- GBM-MYCN
- GBM-RTK-I
- GBM-RTK-II
- GBM-RTK-III
- HGNET-MN1
- IHG
- pedRTK I
- pedRTK II
- PXA
- TIHGG

Figure 3

Figure 4

Figure 5

Figure 6

Geneset	Germline		Somatic	
	NES	FDR	NES	FDR
GO_DNA_REPAIR	-6.97	0.0	-2.68	5.88e-4
HALLMARK_DNA_REPAIR	-4.37	0.0	-4.84	0.0
KAUFFMANN_DNA_REPAIR_GENES	-4.70	0.0	-5.50	0.0
REACTOME_DNA_REPAIR	-3.78	0.0	-4.46	0.0
REACTOME_NUCLEOTIDE_EXCISION_REPAIR	-2.72	1.25e-4	-3.42	0.0
REACTOME_P53_DEPENDENT_G1_DNA_DAMAGE_RESPONSE	-3.80	0.0	-3.17	0.0

Figure 7

A

Genes	TIHG vs GBM	p-value	Drug Class
DNA Pol & Primase	1.45	1.7e-15	DNA Damage
Mitosis	1.39	1.7e-5	Microtubule
Proteasome	1.33	3.7e-4	Proteasome Inh
Epigenetic Regulation	1.24	2.0 e-5	HDAC Inhibitors

B

Sensitivity by drug class

Folate analog, Anti-angiogenic, Platinum, Misc. supportive drugs, Anti-Steroidal, Misc Anti-neoplastics, DNA Synth/Repair inh., Multi-TKI, PI3K-mToR inh., Bcr-Abl inh., RAF-MEK pathway inh., Topo I/II inh., Protein translation inh., Proteasome inh., DNA Intercalator, Microtubule inh., Microtubule stabilizer, HDAC inh., EGFR/HER2 inh., Alkylator

D

| Drug | IC$_{50}$ | |
	Group A (95% CI)	Group B (95% CI)
Aldoxorubicin	197 (154-251)	182 (112-298)
Bortezomib	5.38 (4.31-6.92)	10.3 (10.0-10.7)
Carfilzomib	0.893 (0.693-1.28)	4.06 (3.79-4.42)
Etoposide	286 (236-345)	797 (367-1780)
Marizomib	366 (182-1136)	379 (260-586)
Paclitaxel	82.9 (51.1-132)	26.9(14.3-50.6)
Trametinib	57.5 (32.4-106)	93.9 (71.2-121)
Vinblastine	3.28 (2.60-4.16)	1.05 (0.73-1.61)

C

In vitro drug screen

MAF-145

MAF-496

Paclitaxel
Vinblastine
Doxorubicin
Etoposide
Dasatinib
Bortezomib
Carfilzomib

- Microtubule inhibitor
- Microtubule stabilizer
- DNA synthesis inhibitor
- Alkylator
- Topo II inhibitor
- DNA intercalator/topo II inhibitor
- Topo I inhibitor
- Protein translation inhibitor
- Multi-TKI
- Proteasome inhibitor
- EGFR/HER2 inhibitor
- Bcr-Abl inhibitor
- RAF-MEK pathway inhibitor
- PI3K-MToR inhibitor
- HDAC inhibitor
- Anti-Hormonal/Anti-Steroidal
- Miscellaneous anti-neoplastics

Figure 8

Supplementary Figure 1

A Imaging Analysis

De Novo Pediatric High Grade Glioma (PMID: 29763623)

Treatment Related High Grade Glioma

CCSS N=2 | SJCRH N=13 | UCol N=16

84 Primary pHGG cases with Sufficient Quality Imaging for Tumor Segmentation

21 Cases with Sufficient Quality Imaging for Tumor Segmentation

Calculation of # cases with voxel involvement across MNI brain atlas

Power Analysis alpha = 0.05

Generation of Raw T map

Review of Voxels meeting a pre-specified threshold (p < 0.05)

B

De Novo — z = -15, x = 0, y = 0 — 1 to 31 — N=84

TlHGG — z = -15, x = 0, y = 0 — 1 to 9 — N=21

All — z = -15, x = 0, y = 0 — 1 to 35 — N=105

Difference — z = -15, x = 0, y = 0 — 1.66 to 7.12

T map with voxelwise threshold of p < 0.05

Supplementary Figure 2

A Supplementary Figure 4

B Latency of Treatment-Induced Gliomas

C Survival of Treatment-Induced Gliomas

Supplementary Figure 5

Supplementary Figure 6

Supplementary
Figure 7

Supplementary Figure 8

Supplementary Figure 9

Supplementary Figure 10

Supplementary Figure 11

Multi-institutional Database Query Identified 53 Cases of Probable Treatment-Induced High-Grade Glioma

5 *de novo* pediatric High-Grade Glioma

2 had insufficient tissue for analysis

8 had no tissue available for analysis

4 had recurrent 1^o brain tumors

1 had no history of High-Grade Glioma

20 Cases were excluded after further review

33 Cases were eligible for inclusion

SJCRH & CCSS N=19

U. Colorado N=14

Clinical Annotation (N=26)

Outcomes Analysis (N=27)

Imaging Review (N=21)

Molecular Characterization (RNAseq, WGS/WES, 850K Methylation Analysis) (N=35)

Comparison Relative to Reference Cohorts (HERBY, Capper, Korshounov)

In Vitro Preclinical Characterization with Patient Derived Cell Lines

HERBY Trial Cohort: Reference Methylation, Copy Number & Outcome Data (N=74)

G. Wu Nature Genetics: Reference Sequencing Data (N=127)

Capper D (N=2801) & Korshunov (N=87): Reference Methylation Data