1   **Centromere scission drives chromosome shuffling and reproductive isolation**

3   Vikas Yadav, Sheng Sun, Marco A. Coelho, and Joseph Heitman*

4   Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham,

5   NC 27710, USA

*Corresponding author

Joseph Heitman
Box 3546, 322 CARL Building, Research Drive
Department of Molecular Genetics and Microbiology
Duke University Medical Center
Durham, NC 27710
Phone: 919-684-2824
Fax: 919-684-2790
Email: heitm001@duke.edu

Classification: Biological Sciences, Genetics

Running title: Chromosome shuffling mirrors species boundaries

Keywords: *Cryptococcus neoformans*; DSB repair; Chromosome translocation; Karyotype

evolution.

Abstract

A fundamental characteristic of eukaryotic organisms is the generation of genetic

variation via sexual reproduction. Conversely, significant large-scale genome structure variations

could hamper sexual reproduction, causing reproductive isolation and promote speciation. The

underlying processes behind large-scale genome rearrangements are not well understood and

include chromosome translocations involving centromeres. Recent genomic studies in the

Cryptococcus species complex revealed that chromosome translocations generated via

centromere recombination have reshaped the genomes of different species. In this study, multiple

DNA double-strand breaks (DSBs) were generated via the CRISPR/Cas9 system at centromere-

specific retrotransposons in the human fungal pathogen *Cryptococcus neoformans*. The resulting

DSBs were repaired in a complex manner, leading to the formation of multiple inter-

chromosomal rearrangements and new telomeres, similar to chromothripsis-like events. The

newly generated strains harboring chromosome translocations exhibited normal vegetative

growth but failed to undergo successful sexual reproduction with the parental wild-type strain.

One of these strains failed to produce any spores, while another produced ~3% viable progeny.

The germinated progeny exhibited aneuploidy for multiple chromosomes and showed improved

fertility with both parents. All chromosome translocation events were accompanied without any

detectable change in gene sequences and thus, suggest that chromosomal translocations alone

may play an underappreciated role in the onset of reproductive isolation and speciation.

Introduction

Chromosomes are prone to undergo several rearrangement events, including fusion, fission, deletion, and segmental duplication. In some cases, one chromosome segment is exchanged with another to generate chromosomal translocations. Such exchanges between homologs are regularly observed during meiosis when homologous chromosomes exchange arms via meiotic recombination (1). Chromosome rearrangements can also occur during mitosis, but in a less well-regulated manner, and sometimes as a result of disease conditions like cancer (2, 3). Additionally, rearrangements can also occur within a single chromosome. As a result, chromosome rearrangements during mitosis can cause mutations, gene disruption, copy number variations, as well as alter the expression of genes near the breakpoints (4). Cancer cells show a high level of chromosome rearrangements as compared to healthy cells, and this contributes to critical pathological conditions observed in these cells, such as activation of oncogenes (5, 6).

Chromosomal translocations are initiated by double-strand breaks (DSBs) in DNA (7). Rearrangements involving a single DSB are mainly repaired by the invasion of the broken DNA molecule into a homologous DNA molecule, in a process termed homologous recombination (HR) (6, 8). This invasion can lead to the exchange of DNA between the two molecules of DNA, leading to reciprocal crossover or gene conversion (2). These types of rearrangements occur during meiosis and are regulated to give rise to an error-free repaired sequence. Other types of chromosomal translocation involve two or more DNA DSB sites, which are then fused in an error-prone mechanism known as non-homologous end joining (NHEJ) (9). The two sites can be present on either the same chromosome or different chromosomes. Repair of two DSBs present on the same chromosome can result in the deletion of the intervening sequence or inversion if the sequence is rejoined in reverse orientation (7). On the other hand, the fusion of DSBs from two

different chromosomes can result in chromosomal translocation. In some instances, other repair

pathways like microhomology-mediated end joining (MMEJ) or alternative End Joining (alt-EJ)

also participate in the repair of DSB ends (9, 10).

The occurrence of multiple DSBs in mitotically growing cells at the same time is rare but

occurs at a relatively higher frequency in cancer cells. Such events could occur in a genome due

to replication defects or exogenous factors such as ionizing radiation or chemotherapeutic agents

(11). Multiple DSBs also occur naturally during processes like V(D)J recombination (12). The

occurrence of multiple DSBs can also be induced in the micronuclei of cancer cells (13).

Micronuclei are small nuclei harboring one or a few chromosomes and are generated as a result

of a mitotic failure. These small nuclei act as hotspots of chromosome fragmentation, where

multiple DSB sites are almost simultaneously generated and subsequently rejoined in a random

order, a process known as chromothripsis. Previous reports have suggested that the occurrence of

multiple DSBs alters HR pathways leading to NHEJ or alt-EJ mediated repair (14, 15). Because

both NHEJ and alt-EJ are error-prone, they lead to a significant increase in mutation at the repair

junctions and also randomly join broken fragments.

Apart from generating mutations and gene disruptions, chromosome translocations can

also result in reproductive isolation during meiosis and facilitate speciation (16). The presence of

multiple rearrangements between the two homologous chromosomes from the parents leads to

failures in chromosome pairing during meiosis or crossovers that result in loss of essential genes

(17). Meiosis that is defective in this way will result in the production of progeny with abnormal

genome content. In fungi, the parental nuclei fuse and undergo meiosis before sporulation that

gives rise to progeny. Thus, defects in meiosis lead to the production of spores with abnormal or

incomplete genetic compositions rendering them inviable. *Cryptococcus neoformans* is a

basidiomycete fungus that largely infects immunocompromised humans causing Cryptococcal

meningoencephalitis (18-20). *C. neoformans* harbors a 19 Mb genome with 14 chromosomes

(21). While most of the genome is devoid of repeat regions, centromeres in *C. neoformans* are

rich in a set of LTR-retrotransposons (251 full-length and truncated copies of 6 elements) that

are shared across multiple centromeres (22, 23). *C. neoformans* also has a meiotic cycle making

it a suitable model organism to study meiosis and sexual development (20, 24). It exhibits two

mating types, α and **a**, defined by the *MAT*α and *MAT***a** alleles at the single mating-type locus

(25, 26).

In this study, we exploited the genomic features of *C. neoformans* to study the impact of

chromosome translocations on reproductive isolation. First, retrotransposons present in

centromeres were targeted with CRISPR, generating multiple DSBs simultaneously. Next, the

presence of chromosome rearrangements was screened by Pulsed-Field Gel Electrophoresis

(PFGE), and isolates with multiple chromosomal translocations were identified. The genomes of

these strains were assembled based on long-read nanopore sequencing to characterize the

chromosome rearrangements. Although the strains with new karyotypes did not exhibit growth

defects as compared to the wild-type, the impact of chromosomal rearrangements had a profound

effect on sexual reproduction. These findings demonstrate that *C. neoformans* can tolerate

multiple chromosomal translocations, but that such large-scale changes can cause reproductive

isolation, and promote incipient speciation.

Results

Simultaneous breaks at multiple centromeres lead to chromosome shuffling

Centromeres in *C. neoformans* were previously identified and shown to possess multiple

retrotransposons named Tcn1-Tcn6 (21, 22). These elements are present only in centromeres,

albeit distributed randomly and non-uniformly across all 14 centromeres. We specifically

targeted the Tcn2 element using a guide RNA (gRNA) that would cleave nine centromeres,

generating a total of 18 DSBs (Fig. 1A). We hypothesized that these DSBs would be repaired to

generate chromosomal translocations (Fig. 1B).

After the transformation of wild-type cells with three DNA fragments, each constitutively

expressing a dominant selection marker (G418), Cas9, and the gRNA, colonies were obtained

and screened via pulsed-field gel electrophoresis (PFGE) (Fig. S1). We found karyotypic

variation for at least one chromosome in three out of twelve colonies screened (Fig. S2A). Of the

three, two (VYD135 and VYD136) harbored more than three changes in chromosome banding

pattern as compared to the wild-type strain *C. neoformans* H99 (Fig. 1C and Fig. S2A). To

characterize these chromosomal alterations, we sequenced the genomes of these two strains

using Oxford Nanopore sequencing and were able to assemble them into 17 contigs each (Fig.

S2B).

A synteny block comparison of these genomes with that of the wild-type revealed the

presence of multiple translocations. However, some of these contigs were broken with the

centromere at one of the ends. Additionally, Illumina sequencing of these strains revealed a near

euploid genome for both of these strains (Fig. S2C) except for the smaller arm of chromosome

13 in VYD136 that was present in two copies. To resolve the status of incompletely assembled

contigs, Southern blots of PFGE-separated chromosomes (chromoblots) followed by

hybridization with chromosome-specific probes were conducted for multiple chromosomes (Fig.

S3). Contigs validated to be arms of the same chromosome by this approach were fused

manually with a 50-bp sequence gap to generate a chromosome-size scaffold. The presence of

telomere repeat sequences at both the ends of these scaffolds suggests that these scaffolds

represent full-length chromosomes. The integrity of each genome assembly was further verified

by mapping the respective nanopore reads to the genome assembly. Thus, using nanopore and

Illumina sequencing, and chromoblot analysis, we were able to assemble the genomes of isolates

VYD135 and VYD136 to the chromosome level (Fig. 1D). This analysis also revealed that the

duplicated arm of chromosome 13 of isolate VYD136 exists as an isochromosome with two

broken centromeres fused with each other. Overall, these results show that multiple breaks at

centromeres can lead to karyotype shuffling in *C. neoformans* that is tolerated by the organism.

Centromere breaks generate new telomeres and increase the number of chromosomes

No species in the *Cryptococcus* species complex has been observed to harbor

chromosomes smaller than 500 kb. However, genome-level assemblies of VYD135 and

VYD136 revealed the presence of 2 to 3 chromosomes that are shorter than the shortest naturally

occurring chromosome of wild-type H99 (chromosome 13 of 757 kb length). Bands of the

expected size for these chromosomes were observed in PFGE, indicating that these

chromosomes are *bona fide* and not a result of assembly error (Fig. 2A). In addition to these

small chromosomes, a few novel features were observed for the genomes of VYD135 and

VYD136. Three of the new chromosomes had generated *de novo* telomere sequences on one of

the ends, next to the broken centromeres (chromosomes 13 and 15 of VYD135; chromosome 13

of VYD136) (Fig. 2B, and Fig. S4A). While in two cases, the telomere repeat sequences were

present next to the Tcn2 elements, in the third multiple copies of the Cas9 sequence were found

to be inserted between the Tcn2 element and telomere repeat sequences (Chr15 of VYD135).

Our analysis of these regions did not reveal any common motif as the most likely target for the

de novo telomere addition. While one (Chr13 of VYD135) of the *de novo* additions was preceded

by a sequence similar to that of a telomere repeat (Table S1), the other two telomeres did not

show such a feature. This suggests that the addition of telomere repeat sequences may randomly

occur on broken chromosome ends.

Centromere locations in these two genome assemblies were defined based on synteny

with centromere flanking regions (Table S2). Specifically, BLASTn analysis with centromere-

flanking ORF sequences as query sequences were used to identify the syntenic regions defining

the boundary for centromeres. Next, Tcn elements (Tcn1-6) were mapped across centromeres of

the new strains, VYD135 and VYD136. Surprisingly, foreign DNA elements, such as Cas9 and

neomycin resistance gene sequence, were found to have integrated into multiple centromeres

(Fig. 2C and 2D). Both of these sequences were introduced as linear DNA molecules during the

CRISPR transformation experiment. In some cases, Cas9 and the neomycin gene sequences were

present in multiple copies and in a random order/orientation. Further analysis revealed that these

elements were present at the junction where two parental centromeres fused with each other.

This result suggests that these foreign sequences may have assisted in joining the broken

centromeres to form hybrid centromeres. Previous studies in *S. cerevisiae* observed similar

events where the Ty1 retrotransposon cDNA sequence was found to be present at DNA break

sites (27, 28).

A comparison of centromere length between H99, VYD135, and VYD136 revealed the

presence of some significantly shorter and longer centromeres (16-83 kb versus 31-64 kb in H99)

in both of these new strains (Fig. S4B and Table S2). However, the variation in centromere

length did not seem to confer a visible growth defect indicating no change in centromere

function due to a decrease or increase of length (Fig. S5A). These strains also did not show any

difference when tested for various stress conditions such as temperature, fluconazole, or DNA

damaging agent like Phleomycin (Fig. S5A). When grown with another strain expressing NAT

resistant gene, to test competitive fitness, both VYD135 and VYD136 did not show any

significant difference as compared to wild-type H99 (Fig. S5B). Additionally, these two strains

do not exhibit any difference in virulence in the *Galleria* model of infection (29) (Fig. S5C).

Both VYD135 and VYD136 lead to lethal infection of *Galleria* with the same efficiency as the

wild-type H99α and KN99**a** isolates. Taken together, these results suggest that the changes in

chromosome structure do not grossly affect mitotic fitness or infectivity of *C. neoformans*.

Chromosome shuffling is driven by the Cas9 induced breaks

Isolates VYD135 and VYD136 show inter-chromosomal rearrangements between seven

and eight chromosomes, respectively. The rearrangements in the two strains are not identical,

although they do involve the same set of chromosomes, suggesting that both of these strains

underwent recombination via different routes (Fig. 3A and 3B). Next, the breakpoints in each of

the centromeres were defined by mapping nanopore reads to the wild-type genome. One caveat

is that long reads from nanopore sequencing might not align precisely with the parental

centromere sequence due to the loss of regions of the original sequence. Many reads were found

to be mapped to a single location next to the gRNA cleavage site in almost all of the centromeres

that underwent recombination (Fig. 3C and Fig. S6). In a few cases, the reads did not map to the

cleavage site flanking sequences suggesting deletions occurred during recombination. This loss

of sequence is prominent in centromeres with three or more cleavage sites (*CEN10* in Fig. 3C).

This mapping pattern suggests these non-essential, small fragments were lost, and their loss does

not compromise centromere function. Interestingly, among the chromosomes that contain Tcn2

elements and could be targeted by Cas9 with the Tcn2-specific gRNA, chromosome 11 was not

found to be involved in recombination in either of the two strains, even though *CEN11* is

predicted to be cleaved once. It is possible that the gRNA cleavage site prediction for *CEN11*

could be a result of an incorrect genome assembly or sequence error. Overall, this analysis

supports that recombination was initiated by the Cas9 DSBs. Also, that centromeres lacking

Tcn2 elements were not involved in recombination reflects the specificity of Cas9 and the repair

machinery.

Multiple types of repair machinery were involved in the DSB repair process

Next, the new centromere sequences were compared with the original centromeres in a

pairwise fashion to understand the repair process (Fig. 4 and Fig. S7). For this purpose, we

utilized our newly generated assembly of the wild-type strain, which showed better coverage for

centromere sequences (See Materials and Methods for details). This analysis provides evidence

that both NHEJ and HR pathways participated in the repair of these broken ends. Insertion of the

CAS9 sequence and the *NEO* gene sequence conferring G418 resistance seems to be the result of

DSB repair via NHEJ in all cases as there was no additional sequence added between the ends of

the breaks (Fig. 4B, 4F, and Fig. S7D, S7E). The presence of these DNA fragments in multiple

copies, due to transformation, might have facilitated their insertion. Additionally, *CEN12* of

VYD136 results from a single fusion event between wild-type *CEN8* and *CEN14* after the DNA

breaks (Fig. S7A). For *CEN14* of VYD135, a sequence of 2.7 kb aligned with both parental

centromeres suggesting that the hybrid centromere formed as a result of homologous

recombination (Fig. 4A). *CEN8* of VYD135 exhibited evidence for repair via multiple

mechanisms, including NHEJ, HR, and invasion into multiple different centromeres (Fig. 4C).

Because all of the sequences involved have a Tcn2 sequence at the ends, it is not possible to infer

the order of these events.

A comparison of wild-type *CEN7* and VYD135 *CEN11* shows evidence of resection

beyond the DSB sites (Fig. 4E). Resection was probably followed by strand invasion into broken

pieces of *CEN7* that were released due to multiple DSBs, and a final fusion event with *CEN8*

leading to the formation of hybrid centromeres. Similarly, *CEN13* of VYD136 seems to have

arisen as a result of resection followed by invasion into multiple centromeres before adding

telomere sequences at the end (Fig. S7A). In addition to these multiple recombination events at

these junctions, we also observed inversion events for sequences that were released due to

multiple DSBs in a single centromere (*CEN15* of VYD135) (Fig. 4F). On the other hand, the

inversion in VYD135 *CEN13* has signatures of invasion into another intact copy of *CEN13,*

resulting in the inversion (Fig. 4A).

Apart from inter-chromosome recombination, we also observed intra-chromosomal

recombination in *CEN12* of VYD135. Wild type *CEN12* has two cleavage sites separated by a 10

kb sequence. In VYD135, *CEN12* is smaller due to the absence of the 10 kb sequence, and two

flanking sequences were joined with an overlap of 640 bp (Fig. 4G). This event also led to a

reduction in centromere length for *CEN12*, shortening it significantly (21 kb versus 31 kb in the

wild type). Combined, these results suggest that both HR and NHEJ processes repair DSBs at the

centromeres. Given the high level of identity shared between Tcn2 elements present among

multiple centromeres, other plausible routes to these rearrangements are possible.

Strains with chromosomal rearrangements fail to undergo normal meiosis

Chromosome shuffled strains, VYD135 and VYD136, exhibit multiple chromosomal

translocations compared to the wild-type karyotype, as described above. We hypothesized that

this would lead to incompatibility during meiosis and defects in producing viable spores. To

study this, the two shuffled strains were crossed with the wild-type strain, KN99**a**, which is

congenic with the parental strain H99α. During *Cryptococcus* sexual reproduction, cells of two

opposite mating types fuse and then produce hyphae, basidia, and long chains of spores over a

two-week course of incubation (26). In the basidium, the two parental haploid nuclei fuse and

undergo meiosis producing four haploid nuclei. These meiotic products subsequently undergo

repeated rounds of mitosis that produce four spore chains from the surface of the basidium by

budding. A pairing defect during meiosis would compromise the segregation of chromosomes,

giving rise to progeny with incomplete or imbalanced genomes.

After two weeks of mating, intact spore chains were observed by light microscopy in the

H99α x KN99**a** wild-type cross. On the other hand, the VYD135α x KN99**a** cross formed basidia

but showed almost no sporulation, whereas the VYD136α x KN99**a** cross was largely

indistinguishable from wild-type and did not show a prominent sporulation defect (Fig. 5A).

Meiotic structures were further examined by scanning electron microscopy (SEM). Abundant

spores were produced in the H99α x KN99**a** cross, whereas no spore chains were observed in the

VYD135 x KN99**a** cross (Fig. 5B). The VYD136 x KN99**a** cross showed defective sporulation,

where some basidia formed fewer spores while others had none (Fig. 5B). Next, the germination

rate (equivalent to spore viability) of spores from the H99α x KN99**a** and VYD136 x KN99**a**

crosses was assessed. The spore germination rate for the VYD136 x KN99**a** cross was only 3%

compared to wild-type with 80 to 90% spore germination (30, 31). The lack of sporulation in the

VYD135 x KN99**a** cross and the reduced spore germination rate for the VYD136 x KN99**a** cross

provide evidence that these strains fail to undergo normal meiosis. The translocations in

VYD135 involve seven chromosomes, and in VYD136 eight chromosomes are involved, yet

these isolates show very different phenotypes during meiosis. We hypothesize that this observed

difference in the sporulation phenotype may be attributable to different chromosome

configurations observed in these two strains. Specifically, different configurations could lead to

differences in meiotic pairing and could cause more severe meiotic defects in one compared to

the other. The failure to undergo meiosis will lead to no or defective sporulation.

VYD136 backcross progeny are aneuploid and exhibit improved sexual reproduction

Three viable F1 progeny were isolated from the cross of VYD136 with KN99**a**. These F1

progeny (P1, P2, P3) were characterized to further understand the impact of chromosomal

translocations on meiosis by assessing mating type and ploidy (Fig. 6A and 6B). Flow cytometry

analysis revealed that two of the progeny (P1 and P2) are aneuploid. To investigate this further,

the relative chromosome copy number for the three progeny was determined from read counts

after Illumina and nanopore sequencing, revealing that they are aneuploid for multiple

chromosomes (Fig. 6C). P1 and P2 are both aneuploid for parts of chromosomes 5, 12, 13, and

14, whereas P2 is also aneuploid for the entire chromosome 6. The third progeny (P3) also

showed aneuploidy but only for the shorter arms of chromosomes 12 and 13.

These three F1 progeny were backcrossed to both parents (VYD136 and KN99**a**) and the

wild-type H99α. All three progeny mated with strains of opposite mating type, as expected.

Surprisingly, spore dissections from these crosses revealed a much higher spore germination rate

for the three progeny (11 to 58%) as compared to their parental cross (3% germination rate) (Fig.

6A). P1 and P2 exhibited a much higher (>50%) germination rate for all crosses as compared to

P3, which showed 11 and 21% germination rates. The segmental aneuploidy of these isolates

may explain their higher germination rate compared to the parent VYD136.

The presence of copy number changes for only one arm of most chromosomes suggested

these progeny may have a mixed karyotype. To address this, genomes for the three progeny were

assembled using nanopore sequencing. Due to aneuploidy, these genomes were not assembled

completely and harbored multiple breaks (Fig. S8A and S8B). However, based on their ploidy

profiles, mapping of nanopore reads, as well as synteny comparison with both parents, their

genome configurations were largely resolved (Fig. 6D). The final karyotype shows that most

chromosomes match either with the wild-type or the VYD136 chromosome profile (Fig. 6D and

Fig. S8A). Based on this, we propose that segmental aneuploidy may contribute to overcome the

reproductive barrier that might otherwise arise due to multiple changes in chromosome

configuration. Genetic transmission via an aneuploid intermediate may yield a transitional

population that is more compatible with the ancestral population and slowly diverges to give rise

to a new species. On the other hand, a new species might also arise because of a single event of

multiple translocations occurring throughout the genome. This process is only one of several

factors contributing to reproductive isolation and could play a greater or lesser role dependent

upon the species distribution and mode of reproduction.

Discussion

Chromosome rearrangements shape the genome in multiple ways by affecting gene

positions, causing mutations, and compromising recombination during meiosis. In this study,

chromosomal rearrangements were generated by severing centromeres in a human fungal

pathogen, *C. neoformans*. The centromeres in this species are rich in retrotransposons that are

common among multiple centromeres (22). CRISPR/Cas9 targeting of retrotransposons cleaved

eight centromeres simultaneously in each case. This approach does not affect gene-rich regions,

thus avoiding the risks involved with the loss or mutation of genes. At the same time, this

allowed us to study a) how multiple breaks in the genome are tolerated, b) how DSBs in

heterochromatic regions such as centromeres are repaired, and c) how the resulting chromosomal

rearrangements impact sexual reproduction. Centromere-mediated chromosomal translocation

events have been observed in the *Cryptococcus* species complex (21, 32) (Fig. 7A-F). Genome

comparisons between non-pathogenic *C. amylolentus* and pathogenic species, such as *C.*

neoformans, show at least six centromere-mediated translocations (32). Previous studies have

suggested that retrotransposons, including centromeric elements, are actively transcribed and can

undergo transposition in *Cryptococcus* species (22, 33, 34). Thus, it is possible that these

elements generate DNA breaks during their insertion in to the centromeres, which can then

catalyze the occurrence of chromosome rearrangements. Our findings suggest these

translocations could have driven speciation in this species complex, in addition to other factors.

Recent studies have suggested that centromeres can undergo recombination, at least

mitotic recombination, as compared to previous models in which centromeres were recalcitrant

to recombination (22, 35-40). Our results further support that centromeres can undergo

recombination when DSBs are generated in these regions. Each of the broken ends was

processed and subjected to HR or NHEJ mediated repair and recombination. We also observed a

case where a centromere, following the generation of DSBs, recombined, causing the loss of

intermediate DNA sequences present between the two ends. In similar events, many fragments of

centromere sequences were either lost or fused within other centromeres, altering the architecture

of centromeres in this species (Fig. 7G). This result suggests that variation in centromere

structure is not critical for centromere function, but might play some other role in genome

organization. This conclusion is also supported by a previous study that showed a significant

reduction in centromere length in *C. deuterogattii*, a closely related species of *C. neoformans*

(22). This variation in centromere architecture also raises questions about the requirement of

longer centromeres in most organisms. According to previously proposed models, longer

centromeres with more centromere repeats might bind more kinetochore proteins (41-43). This

would enable stronger kinetochore-microtubule interactions for longer centromeres and favor

transmission of longer centromeres during meiosis. Our system can be harnessed to understand

the dynamics of centromere transmission and test this hypothesis in future studies.

Interestingly, we also observed the addition of *de novo* telomere repeat sequences

adjacent to the broken centromeres. While the mechanism underlying this process remains to be

elucidated, the frequency of occurrence of *de novo* telomere addition seems to be high at ~10%

(3/32 ends). Additionally, two of the three telomere repeat additions involved invasion or fusion

with other sequences before the addition of repeat sequences. *De novo* telomere formation on

broken DNA ends has been observed in many species and is commonly known as chromosome

healing (44-46). Chromosome healing occurs during mammalian development and has been

observed in mouse embryonic stem cells as well as human germline cells (47-49). It is also

proposed to occur in cancer cells to stabilize the ends of broken chromosome fragments arising

as a result of chromothripsis or telomere crisis (50, 51). This process maintains the genome

content while increasing the number of chromosomes in the case of DSBs that could not be

repaired. *De novo* telomere addition in *S. cerevisiae* was shown to be influenced by the presence

of telomere-like sequences at the broken DNA end. We found a similar scenario only in one of

the three *de novo* telomeres formed, suggesting that the mechanism might differ in *C.*

neoformans. In our study, we did not observe a significant impact of the loss of centromere

sequences on the growth of the chromosomal shuffled strains as compared to the parental strain.

This could suggest that shorter centromere sequences are sufficient to propagate the genome

content. Furthermore, *de novo* telomere formation next to centromeres did not seem to have an

effect on their functions suggesting that centromeres and telomeres do not influence each other's

function in *C. neoformans*, similar to telocentric/acrocentric chromosomes found in other

organisms.

Repair of a DSB site is a complex process and involves multiple repair machineries

including HR and NHEJ. HR mainly takes place during S-phase, whereas NHEJ occurs

throughout the cell cycle (52, 53). HR mainly leads to gene conversion and can also drive

recombination between repetitive sequences leading to translocation. NHEJ is more error-prone

and can join any two sites resulting in translocations. Thus, both of these processes can lead to

translocations but can result in very different types of junctions (8, 54). Our analysis of junctions

exhibited evidence for both HR and NHEJ. Previous reports suggested that the NHEJ pathway is

preferred over HR in the case of multiple DSBs (14, 15). Although the number is small, our

results suggest that both of these pathways might take place at a similar rate in *C. neoformans*.

However, the underlying regulatory mechanisms remain a subject for further investigation.

We also observed the inversion of sequences suggesting either strand invasion or simple

fusion of a broken piece in reverse order by NHEJ. Our analysis also suggests the occurrence of

multi-invasion repair (MIR), a subtype of HR, which can be favored for the repair of repeat

sequences (6, 55). According to the MIR pathway, a single broken end might enter multiple

target DNA molecules based on homology (56, 57). The DNA breaks in our experiments were

made in the Tcn2 element, which is present in multiple copies across centromeres; thus, it is

possible that some of the ends might have been repaired via MIR. Our results also suggest that

breaks in repeat molecules are repaired either by NHEJ or HR, where other repeat sequences

present in the genome may aid in the repair process. Similar observations have also been made

previously in *S. cerevisiae*, where DSBs were generated with gamma radiation, HO-

endonuclease, or CRISPR (58-60). In all of these studies, multiple copies of Ty elements were

found to be involved in the generation of translocations. In the study that employed CRISPR,

similar multiple translocation events were observed (59). However, Ty elements in *S. cerevisiae*

are distributed across the genome, unlike *C. neoformans,* where all Tcn2 elements are clustered

in centromeres.

Multiple concurrent breaks in the genome are rare and occur mainly in pathogenic

conditions such as cancer. A commonly observed outcome is chromothripsis, where one or

several chromosomes are broken into multiple smaller pieces and rejoined randomly to shuffle

the targeted chromosomal region (11, 13). This phenomenon is associated with the generation of

multiple mutations, as well as the activation of oncogenes. Chromothripsis is initiated by the

generation of multiple breaks in a localized manner. The source of these breaks could be either

internal factors such as replication associated breaks or external factors, including ionizing

radiation or chemotherapy. These break sites are then repaired in a random manner. While this

process is well-known, the mechanisms involved in this process are poorly understood (13, 61).

In our experiments, we observed that similar events take place during DSBs repair (Step 3 in Fig.

7G). The connection between these processes needs to be further explored to establish whether

our system can be extended to understand chromothripsis.

Our approach induced multiple simultaneous breaks in the genome, which were then

repaired to generate chromosome shuffling. Centromeres are known to cluster in *C. neoformans*

during mitosis (62, 63), and this may have promoted their interaction during the DSB repair

events that generated these alterations. This centromere clustering in *C. neoformans* also mimics

the clustering of DSB sites observed during the process of chromothripsis. We posit our

approach could provide answers to critical questions regarding chromothripsis. The MIR

pathway has been implicated as one pathway contributing to chromothripsis (6, 57). In our study,

we also found evidence that the MIR pathway contributes to chromosome shuffling, further

suggesting similarities between chromothripsis and the events we observed. Notably,

chromothripsis is mainly observed in chromosome arms, whereas centromeres were targeted in

our studies (13). However, some studies have indicated an association between chromothripsis

break sites and the presence of transposon sequences (11, 64, 65). In our approach, the breaks

were also located within transposon sequences, which are part of the centromeres. Thus,

understanding the factors governing this process in *C. neoformans* could also shed light on facets

of chromothripsis.

Chromosome rearrangements have been implicated in speciation, acting via reducing

fertility or gene flow in the progeny that inherit the translocation (66-69). However, most models

proposing speciation in this manner also consider geographical isolation as other major criteria

(16). Our results, along with other studies, support this line of thought and suggest that

chromosome rearrangements followed by geographical isolation can drive speciation (66, 70-72).

Other models consider chromosome rearrangement as an effect of speciation, which happens as a

result of simultaneous rearrangement after the speciation (67). A number of studies linking

chromosome rearrangements and speciation have been conducted using *S. cerevisiae* and

Schizosaccharomyces pombe. The *S. cerevisiae sensu stricto* group has been the subject of such

studies over a long time period (73). A common approach has been to isolate viable hybrids

between these species and then to study sporulation of these hybrids. These studies have

suggested a role for three factors in the poor spore viability of hybrids: genetic incompatibilities,

sequence divergence affecting mismatch repair system, and chromosome rearrangements (71,

74-76). However, more recent studies have suggested a greater role of chromosome

translocations in this process than previously thought (77, 78). Studies involving a large

collection of *S. pombe* isolates revealed karyotypic diversity among these isolates (79, 80). Some

of these translocation events were found in geographically separated populations, suggesting a

role in speciation. Another study found that *S. pombe* can mate successfully with a close relative,

Schizosaccharomyces kambucha to form viable hybrids but these hybrids give rise to very few

viable spores as compared to the individual parental species (81). The loss in viability was found

to be correlated with two translocations present among these strains as well as three independent

meiotic drive elements. Studies in marsupials suggested a role of chromosome rearrangements

involving centromeres in the process of speciation in this species complex (82-86). These

involved centromere-mediated translocations as well as differences in centromere lengths

between two species. In our study, no genes were found to be affected or mutated because all

rearrangements were generated via centromere recombination. Thus, a meiotic failure of

rearrangement harboring strains suggests that chromosome translocations solely can drive

reproductive isolation in a species. It is important to note that centromeres as such may not have

any role in this phenomenon and rather centromere-localized breaks are just one of the ways to

achieve chromosome rearrangements that in turn facilitate reproductive isolation. A recent study

has provided evidence that even small repositioning of centromeres can generate reproductive

barriers (87).

Here, we find that the presence of multiple centromere-mediated chromosomal

rearrangements dramatically reduces the efficiency of meiosis and generates post-zygotic

reproductive isolation. A complete failure to undergo meiosis with the existing population could

lead to loss of new chromosome configurations, as they will not be passed to the new

generations. We found that the presence of a mix of the original karyotype along with the new

karyotype enhanced meiotic success. Interestingly, all three viable progeny exhibited similar

karyotypes suggesting that this chromosome configuration is probably selected for cell survival

while also keeping most of the genome as haploid. Thus, aneuploidy could also shield the newly

established chromosome rearrangements and allow it to persist and spread in the population,

eventually leading to the fixation of new changes. This hypothesis is supported by our results in

which the aneuploid F1 progeny exhibited a much higher spore germination rate with either

parent as compared to the crosses between the haploid parents. Additionally, the less aneuploid

progeny exhibited less successful meiosis, further suggesting the role of aneuploidy in this

process. Similar observations have also been made in *S. cerevisiae,* and its related species, where

backcrosses and aneuploidy overcame reproductive isolation to increase spore germination (74,

75). Based on this, we propose a model in which the rearranged chromosomes can be present

along with the parent chromosomes in an aneuploid intermediate (Fig. 7G). The aneuploid

intermediate then allows the transmission of rearranged chromosomes to the next generation

until a particular rearrangement becomes fixed in part of the population. Once fixed, the

rearranged population will stably transmit itself and can give rise to a new subpopulation. The

presence of geographical barriers, as well as possible advantageous selective roles for the new

rearrangement, might further promote fixation of this newly acquired rearrangement.

Acknowledgments

We thank Valerie Lapham at North Carolina State University electron microscopy

facility for assistance with scanning electron microscopy. We thank Tom Petes and Sue Jinks-

Robertson for critical reading of this manuscript. This work was supported by NIH/NIAID R37

MERIT award AI39115-21 to JH, R01 grant AI50113-15 to JH, and R01 grant AI112595-04

awarded to JH, David Tobin, and Paul Magwene. We also thank the CIFAR program, Fungal

Kingdom: Threats & Opportunities for their support.

Materials and Methods

Strains and media

C. neoformans wild-type strains H99α and KN99**a** served as the wild-type isogenic

parental lineages for the experiments. Strains were grown in YPD media for all experiments at

30°C unless stated otherwise, and G418 was added at a final concentration of 200 µg/ml for

selection of transformants. MS media was used for all the mating assays, which were performed

as described previously (26). Basidia specific spore dissections were performed after two weeks

of mating, and the spore germination rate was scored after four days of dissection. All strains

used in this study are listed in Table S3.

Genome assembly and synteny comparison

The *C. neoformans* H99 genome was reassembled with Canu v1.7 based on previously

published PacBio and Illumina data to obtain a better resolution of the centromeric regions (see

Table S4 and S5 for details) (21, 22, 88). The resulting draft assembly was improved by

correcting errors using five rounds of Pilon (v1.22) polishing ('--fix all' setting) and the Illumina

reads mapped to the respective assemblies by the use of BWA-MEM (v0.7.17-r1188) (89, 90).

Centromere locations were mapped based on BLAST analysis with centromere flanking genes,

and coordinates for these new locations are provided in Table S1. Because some of the

centromere lengths differed in the new assembly as compared to the previous one, we validated

the new centromere lengths by mapping the Canu-corrected PacBio read using Minimap2,

followed by visual inspection in IGV (91). Because our work involved analysis of centromere

sequences, we utilized this new improved assembly as the reference for all of our analyses. Once

the locations were mapped, we oriented all of the chromosomes such that the longer arm (q arm)

begins the chromosome, and the smaller arm (p arm) is the distal part of the chromosome.

De novo genome assemblies for the VYD135 and VYD136 isolates, and progeny (P1, P2,

and P3) were generated with Canu using Oxford Nanopore reads > 2 kb (see Table S4 and S5 for

details) followed by Pilon polishing as described above. When necessary, and after validating by

chromoblot analysis, broken contigs were joined artificially with a 50 bp sequence gap to

generate a full-length chromosome. Once completed, the chromosomes were numbered based on

their length with the longest chromosome as the first. For progeny P1, P2, and P3, extra

chromosome configuration was inferred to obtain the final karyotype based on their Illumina and

nanopore sequencing reads mapping. Sequence data generated in this study were submitted to

NCBI with the BioProject accession number PRJNA577944.

Synteny comparisons between the genomes were performed with SyMAP v4.2 and

default parameters (92) (http://www.agcol.arizona.edu/software/symap/). The comparison block

maps were exported as .svg files and were then processed using Adobe Illustrator and Adobe

Photoshop for representation purposes.

References

1. Hunter N (2015) Meiotic recombination: The essence of heredity. *Cold Spring Harb Perspect Biol* 7(12):a016618.

2. Andersen SL & Sekelsky J (2010) Meiotic versus mitotic recombination: Two different routes for double-strand break repair. *Bioessays* 32(12):1058-1066.

3. LaFave MC & Sekelsky J (2009) Mitotic recombination: why? when? how? where? *PLoS Genet* 5(3):e1000411.

4. Harewood L & Fraser P (2014) The impact of chromosomal rearrangements on regulation of gene expression. *Hum Mol Genet* 23(R1):R76-82.

5. Hasty P & Montagna C (2014) Chromosomal rearrangements in cancer: Detection and potential causal mechanisms. *Mol Cell Oncol* 1(1):e29904.

6. Piazza A & Heyer WD (2019) Homologous recombination and the formation of complex genomic rearrangements. *Trends Cell Biol* 29(2):135-149.

7. Lysák MA & Schubert I (2013) Mechanisms of chromosome rearrangements. *Plant Genome Diversity Volume 2: Physical Structure, Behaviour and Evolution of Plant Genomes*, eds Greilhuber J, Dolezel J, & Wendel JF (Springer Vienna, Vienna), pp 137-147.

8. Mehta A & Haber JE (2014) Sources of DNA double-strand breaks and models of recombinational DNA repair. *Cold Spring Harb Perspect Biol* 6(9):a016428.

9. Chang HHY, Pannunzio NR, Adachi N, & Lieber MR (2017) Non-homologous DNA end joining and alternative pathways to double-strand break repair. *Nat Rev Mol Cell Biol* 18(8):495-506.

10. Lieber MR (2010) The mechanism of double-strand DNA break repair by the nonhomologous DNA end-joining pathway. *Annu Rev Biochem* 79:181-211.

11. Koltsova AS, *et al.* (2019) On the complexity of mechanisms and consequences of chromothripsis: An update. *Front Genet* 10:393.

12. Arya R & Bassing CH (2017) V(D)J recombination exploits DNA damage responses to promote immunity. *Trends Genet* 33(7):479-489.

13. Ly P & Cleveland DW (2017) Rebuilding chromosomes after catastrophe: Emerging mechanisms of chromothripsis. *Trends Cell Biol* 27(12):917-930.

556   14.   Gudjonsson T, *et al.* (2012) TRIP12 and UBR5 suppress spreading of chromatin
557   ubiquitylation at damaged chromosomes. *Cell* 150(4):697-709.

558   15.   Pellestor F (2019) Chromoanagenesis: cataclysms behind complex chromosomal
559   rearrangements. *Mol Cytogenet* 12:6.

560   16.   Rieseberg LH (2001) Chromosomal rearrangements and speciation. *Trends Ecol Evol*
561   16(7):351-358.

562   17.   Pellestor F, *et al.* (2011) Complex chromosomal rearrangements: origin and meiotic
563   behavior. *Hum Reprod Update* 17(4):476-494.

564   18.   Lin X & Heitman J (2006) The biology of the *Cryptococcus neoformans* species
565   complex. *Annu Rev Microbiol* 60:69-105.

566   19.   Rajasingham R, *et al.* (2017) Global burden of disease of HIV-associated cryptococcal
567   meningitis: an updated analysis. *Lancet Infect Dis* 17(8):873-881.

568   20.   Sun S, Coelho MA, David-Palma M, Priest S, & Heitman J (2019) The evolution of
569   sexual reproduction and the mating-type locus: Links to pathogenesis of *Cryptococcus*
570   human pathogenic fungi. *Annu Rev Genet* 53(1):417-444.

571   21.   Janbon G, *et al.* (2014) Analysis of the genome and transcriptome of *Cryptococcus*
572   *neoformans* var. *grubii* reveals complex RNA expression and microevolution leading to
573   virulence attenuation. *PLoS Genet* 10(4):e1004261.

574   22.   Yadav V, *et al.* (2018) RNAi is a critical determinant of centromere evolution in closely
575   related fungi. *Proc Natl Acad Sci U S A* 115(12):3108-3113.

576   23.   Loftus BJ, *et al.* (2005) The genome of the basidiomycetous yeast and human pathogen
577   *Cryptococcus neoformans*. *Science* 307(5713):1321-1324.

578   24.   Kozubowski L & Heitman J (2012) Profiling a killer, the development of *Cryptococcus*
579   *neoformans*. *FEMS Microbiol Rev* 36(1):78-94.

580   25.   Lengeler KB, *et al.* (2002) Mating-type locus of *Cryptococcus neoformans*: a step in the
581   evolution of sex chromosomes. *Eukaryot Cell* 1(5):704-718.

582   26.   Sun S, Priest SJ, & Heitman J (2019) *Cryptococcus neoformans* mating and genetic
583   crosses. *Curr Protoc Microbiol* 53(1):e75.

584   27.   Garfinkel DJ (1997) Genetic loose change: how retroelements and reverse transcriptase
585   heal broken chromosomes. *Trends Microbiol* 5(5):173-175.

28. Moore JK & Haber JE (1996) Capture of retrotransposon DNA at the sites of
chromosomal double-strand breaks. *Nature* 383(6601):644-646.

29. Mylonakis E*, et al.* (2005) *Galleria mellonella* as a model system to study *Cryptococcus*
neoformans pathogenesis. *Infect Immun* 73(7):3842-3850.

30. Botts MR, Giles SS, Gates MA, Kozel TR, & Hull CM (2009) Isolation and
characterization of *Cryptococcus neoformans* spores reveal a critical role for capsule
biosynthesis genes in spore biogenesis. *Eukaryot Cell* 8(4):595-605.

31. Velagapudi R, Hsueh YP, Geunes-Boyer S, Wright JR, & Heitman J (2009) Spores as
infectious propagules of *Cryptococcus neoformans*. *Infect Immun* 77(10):4345-4355.

32. Sun S*, et al.* (2017) Fungal genome and mating system transitions facilitated by
chromosomal translocations involving intercentromeric recombination. *PLoS Biol*
15(8):e2002527.

33. Janbon G*, et al.* (2010) Characterizing the role of RNA silencing components in
Cryptococcus neoformans. *Fungal Genet Biol* 47(12):1070-1080.

34. Wang X*, et al.* (2010) Sex-induced silencing defends the genome of *Cryptococcus*
neoformans via RNAi. *Genes Dev* 24(22):2566-2582.

35. Jaco I, Canela A, Vera E, & Blasco MA (2008) Centromere mitotic recombination in
mammalian cells. *J Cell Biol* 181(6):885-892.

36. Talbert PB & Henikoff S (2010) Centromeres convert but don't cross. *PLoS Biol*
8(3):e1000326.

37. Thakur J & Sanyal K (2013) Efficient neocentromere formation is suppressed by gene
conversion to maintain centromere function at native physical chromosomal loci in
Candida albicans. *Genome Res* 23(4):638-652.

38. Zafar F*, et al.* (2017) Regulation of mitotic recombination between DNA repeats in
centromeres. *Nucleic Acids Res* 45(19):11222-11235.

39. Liebman SW, Symington LS, & Petes TD (1988) Mitotic recombination within the
centromere of a yeast chromosome. *Science* 241(4869):1074-1077.

40. Symington LS & Petes TD (1988) Meiotic recombination within the centromere of a
yeast chromosome. *Cell* 52(2):237-240.

41. Dawe RK & Henikoff S (2006) Centromeres put epigenetics in the driver's seat. *Trends*
Biochem Sci 31(12):662-669.

42. Kursel LE & Malik HS (2018) The cellular mechanisms and consequences of centromere
drive. *Curr Opin Cell Biol* 52:58-65.

43. Rosin LF & Mellone BG (2017) Centromeres drive a hard bargain. *Trends Genet*
33(2):101-117.

44. Kurzhals RL*, et al.* (2017) Chromosome healing Is promoted by the telomere cap
component Hiphop in *Drosophila*. *Genetics* 207(3):949-959.

45. Melek M & Shippen DE (1996) Chromosome healing: spontaneous and programmed *de*
novo telomere formation by telomerase. *Bioessays* 18(4):301-308.

46. Pennaneach V, Putnam CD, & Kolodner RD (2006) Chromosome healing by *de novo*
telomere addition in *Saccharomyces cerevisiae*. *Mol Microbiol* 59(5):1357-1368.

47. Gao Q*, et al.* (2008) Telomerase-dependent and -independent chromosome healing in
mouse embryonic stem cells. *DNA Repair* 7(8):1233-1249.

48. Kulkarni A, Zschenker O, Reynolds G, Miller D, & Murnane JP (2010) Effect of
telomere proximity on telomere position effect, chromosome healing, and sensitivity to
DNA double-strand breaks in a human tumor cell line. *Mol Cell Biol* 30(3):578-589.

49. Varley H, Di S, Scherer SW, & Royle NJ (2000) Characterization of terminal deletions at
7q32 and 22q13.3 healed by *de novo* telomere addition. *Am J Hum Genet* 67(3):610-622.

50. Maciejowski J & de Lange T (2017) Telomeres in cancer: tumour suppression and
genome instability. *Nat Rev Mol Cell Biol* 18(3):175-186.

51. Murnane JP (2010) Telomere loss as a mechanism for chromosome instability in human
cancer. *Cancer Res* 70(11):4255-4259.

52. Hustedt N & Durocher D (2016) The control of DNA repair by the cell cycle. *Nat Cell*
Biol 19(1):1-9.

53. Murray JM & Carr AM (2018) Integrating DNA damage repair with the cell cycle. *Curr*
Opin Cell Biol 52:120-125.

54. Roukos V & Misteli T (2014) The biogenesis of chromosome translocations. *Nat Cell*
Biol 16(4):293-300.

55. Sakofsky CJ & Malkova A (2017) Break induced replication in eukaryotes: mechanisms,
functions, and consequences. *Crit Rev Biochem Mol Biol* 52(4):395-413.

56. Piazza A & Heyer WD (2018) Multi-invasion-induced rearrangements as a pathway for
physiological and pathological recombination. *Bioessays* 40(5):e1700249.

57.   Piazza A, Wright WD, & Heyer WD (2017) Multi-invasions are recombination

byproducts that induce chromosomal rearrangements. *Cell* 170(4):760-773.e15.

58.   Argueso JL*, et al.* (2008) Double-strand breaks associated with repetitive DNA can

reshape the genome. *Proc Natl Acad Sci U S A* 105(33):11845-11850.

59.   Fleiss A*, et al.* (2019) Reshuffling yeast chromosomes with CRISPR/Cas9. *PLoS Genet*

15(8):e1008332.

60.   Muramoto N*, et al.* (2018) Phenotypic diversification by enhanced genome restructuring

after induction of multiple DNA double-strand breaks. *Nat Commun* 9(1):1995.

61.   Pellestor F & Gatinois V (2019) Chromothripsis, a credible chromosomal mechanism in

evolutionary process. *Chromosoma* 128(1):1-6.

62.   Kozubowski L*, et al.* (2013) Ordered kinetochore assembly in the human-pathogenic

basidiomycetous yeast *Cryptococcus neoformans*. *mBio* 4(5):e00614-00613.

63.   Yadav V & Sanyal K (2018) Sad1 spatiotemporally regulates kinetochore clustering to

ensure high-fidelity chromosome segregation in the human fungal pathogen

Cryptococcus neoformans. *mSphere* 3(4):e00190-00118.

64.   Hancks DC (2018) A role for retrotransposons in chromothripsis. *Methods Mol Biol*

1769:169-181.

65.   Nazaryan-Petersen L*, et al.* (2016) Germline chromothripsis driven by L1-mediated

retrotransposition and Alu/Alu homologous recombination. *Hum Mutat* 37(4):385-395.

66.   Fischer G, James SA, Roberts IN, Oliver SG, & Louis EJ (2000) Chromosomal evolution

in *Saccharomyces*. *Nature* 405(6785):451-454.

67.   Livingstone K & Rieseberg L (2004) Chromosomal evolution and speciation: a

recombination-based approach. *New Phytol* 161(1):107-112.

68.   Noor MA, Grams KL, Bertucci LA, & Reiland J (2001) Chromosomal inversions and the

reproductive isolation of species. *Proc Natl Acad Sci U S A* 98(21):12084-12088.

69.   Brown JD & O'Neill RJ (2010) Chromosomes, conflict, and epigenetics: chromosomal

speciation revisited. *Annu Rev Genomics Hum Genet* 11:291-316.

70.   Bergstrom A*, et al.* (2014) A high-definition view of functional genetic variation from

natural yeast genomes. *Mol Biol Evol* 31(4):872-888.

71.   Liti G, Barton DB, & Louis EJ (2006) Sequence diversity, reproductive isolation and

species concepts in *Saccharomyces*. *Genetics* 174(2):839-850.

72.   Liti G, *et al.* (2009) Population genomics of domestic and wild yeasts. *Nature*
680   458(7236):337-341.

681   73.   Greig D (2009) Reproductive isolation in *Saccharomyces*. *Heredity (Edinb)* 102(1):39-
682   44.

683   74.   Delneri D, *et al.* (2003) Engineering evolution to study speciation in yeasts. *Nature*
684   422(6927):68-72.

685   75.   Greig D, Louis EJ, Borts RH, & Travisano M (2002) Hybrid speciation in experimental
686   populations of yeast. *Science* 298(5599):1773-1775.

687   76.   Louis EJ (2011) Population genomics and speciation in yeasts. *Fungal Biology Reviews*
688   25(3):136-142.

689   77.   Dujon BA & Louis EJ (2017) Genome diversity and evolution in the budding yeasts
690   (Saccharomycotina). *Genetics* 206(2):717-750.

691   78.   Hou J, Friedrich A, de Montigny J, & Schacherer J (2014) Chromosomal rearrangements
692   as a major mechanism in the onset of reproductive isolation in *Saccharomyces cerevisiae*.
693   *Curr Biol* 24(10):1153-1159.

694   79.   Brown WR, *et al.* (2011) A geographically diverse collection of *Schizosaccharomyces*
695   *pombe* isolates shows limited phenotypic variation but extensive karyotypic diversity. *G3*
696   *(Bethesda)* 1(7):615-626.

697   80.   Jeffares DC, *et al.* (2015) The genomic and phenotypic diversity of *Schizosaccharomyces*
698   *pombe*. *Nat Genet* 47(3):235-241.

699   81.   Zanders SE, *et al.* (2014) Genome rearrangements and pervasive meiotic drive cause
700   hybrid infertility in fission yeast. *Elife* 3:e02630.

701   82.   Deakin JE (2018) Chromosome evolution in marsupials. *Genes (Basel)* 9(2).

702   83.   Metcalfe CJ, *et al.* (2007) Genomic instability within centromeres of interspecific
703   marsupial hybrids. *Genetics* 177(4):2507-2517.

704   84.   O'Neill RJ, Eldridge MD, & Metcalfe CJ (2004) Centromere dynamics and chromosome
705   evolution in marsupials. *J Hered* 95(5):375-381.

706   85.   Potter S, *et al.* (2017) Chromosomal speciation in the genomics era: Disentangling
707   phylogenetic evolution of rock-wallabies. *Front Genet* 8:10.

86.   Bulazel KV, Ferreri GC, Eldridge MD, & O'Neill RJ (2007) Species-specific shifts in

centromere sequence composition are coincident with breakpoint reuse in karyotypically

divergent lineages. *Genome Biol* 8(8):R170.

87.   Lu M & He X (2019) Centromere repositioning causes inversion of meiosis and

generates a reproductive barrier. *Proc Natl Acad Sci U S A* 116(43):21580-21591.

88.   Koren S*, et al.* (2017) Canu: scalable and accurate long-read assembly via adaptive k-mer

weighting and repeat separation. *Genome Res* 27(5):722-736.

89.   Li H & Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* 25(14):1754-1760.

90.   Walker BJ*, et al.* (2014) Pilon: an integrated tool for comprehensive microbial variant

detection and genome assembly improvement. *PLoS One* 9(11):e112963.

91.   Li H (2018) Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*

34(18):3094-3100.

92.   Soderlund C, Bomhoff M, & Nelson WM (2011) SyMAP v3.4: a turnkey synteny system

with application to plant genomes. *Nucleic Acids Res* 39(10):e68.

 Figure and Figure legends

Fig. 1. Centromere-specific DSBs mediated by CRISPR lead to chromosome rearrangements. (A) Centromere maps showing the distribution of retrotransposons (Tcn1-Tcn6) in the centromeres of wild-type strain H99 of *C. neoformans*. **(B)** An outline depicting the model for achieving multiple chromosome translocations in *C. neoformans*. **(C)** PFGE revealed many differences in the karyotype of VYD135 and VYD136 as compared with wild-type, H99. **(D)** Chromosome maps for VYD135 and VYD136 compared to the H99 genome revealed multiple chromosome translocations in these strains. Chromosomes are colored with H99 chromosomes as reference. 'q' represents the longer arm while 'p' represents the shorter arm according to the wild-type chromosome configuration.

Fig. 2. Sequence analysis of new centromeres and telomeres. (A) PFGE EtBr staining
showing bands for newly generated short chromosomes and new telocentric chromosomes. M
represents *S. cerevisiae* chromosomes. **(B)** Chromosome map of one of the newly formed
telomeres showing the presence of telomere sequence repeats next to Tcn2 elements present in
the centromere. **(C-D)** Maps showing the distribution of retrotransposons, along with the
integration of foreign sequences from CRISPR/Cas9 and the neomycin resistance gene (NEO[R]),
in the centromeres of VYD135 (C) and VYD136 (D). Numbers in brackets next to *CEN* numbers
represent the wild-type *CEN* numbers that rearranged to form new centromeres.

Fig. 3. Chromosome rearrangements are mediated via the gRNA cleavage site. (A-B)
Simplified outline maps depicting the chromosomes that underwent translocation in VYD135
(A) and VYD136 (B). Black semicircles, telomeres; red semicircles, de novo telomeres, narrow
bands, centromeres; shaded box on chromosome 5, *MAT* locus. **(C)** Nanopore reads mapping to
the wild-type H99 genome revealed the sites of DSB formation or repair junctions at
centromeres. The reads either converge on a single site (*CEN5*, *CEN8*) or exhibit sequence gaps
between sites (*CEN10*) marking the location of junctions. Red bars indicate the centromeres
whereas the black vertical lines mark the site of gRNA cleavage. Cov, Coverage of nanopore
reads; Map, Mapping of nanopore reads.

Fig. 4. Synteny analysis of rearranged centromeres with wild-type H99 centromeres reveals complex rearrangements. (A-G) A pair-wise comparison of newly generated and wild-type centromeres revealed that translocations are mediated by double-strand breaks (DSBs) generated via CRISPR. Centromere specific events are described in the individual panels. Grey shades represent direct synteny, while the cyan shade represents inversion events. In the cases that are shown in detail, the cross represents the evidence for homologous recombination, whereas the connecting grey lines represent joining events marking non-homologous end-joining. VYD135-*CEN7* was generated after artificial fusion of two contigs and hence was not analyzed in detail.

Fig. 5. Chromosome shuffled isolates exhibit defects in sexual reproduction. (A) Light
microscopy images showing hyphae, basidia, and spore chains in crosses between *MAT***a** wild-
type strain KN99**a** and *MAT*α wild-type (H99) and rearranged strains (VYD135 and VYD136).
Scale bar, 100 µm. **(B)** Scanning Electron Microscope (SEM) images depicting a complete or
partial sporulation defect when wild-type KN99**a** was crossed with VYD135 and VYD136,
respectively. Scale bar, 10 µm.

Fig. 6. VYD136 progeny are aneuploid and exhibit mixed karyotypes. (A) Analysis of the mating-type locus and the mating efficiency of three progeny of VYD136 with either parent. The numbers in brackets represent spore germination rates of respective crosses. **(B)** Flow cytometry profiles of wild-type haploid, diploid, and three progeny of VYD136. **(C)** Illumina sequencing data mapped to the wild-type H99 genome revealed aneuploidy for multiple chromosomes in the three VYD136 progeny. **(D)** Karyotypes for three progeny showing synteny as compared to the wild-type H99 genome. The red stars represent breaks that were fused later based on synteny and ploidy. The chromosomes shown with red bar on top were not assembled *de novo* but represent possible chromosomes configuration based on Illumina and nanopore sequencing analysis. Contigs 14 in VYD136.P3 could not be resolved into their chromosome configuration. 'q' represents longer arm while 'p' represents shorter arm according to the wild-type chromosome configuration.

Fig. 7. A model proposing the evolution of reproductive isolation induced by centromere breaks. (A) A schematic representing phylogeny showing the relationships of five *Cryptococcus* species, for which chromosome-level genome assemblies are currently available and published. **(B-F)** Chromosome maps for five species: *C. neoformans*, *C. deneoformans*, *C. amylolentus*, *C. deuterogattii,* and *C. gattii*. The synteny maps for all species were generated with *C. neoformans* as the reference and are colored accordingly. Chromosomal translocations involving centromeres are marked with arrowheads. **(G)** A model showing the events observed in the study. DSBs generated using CRISPR at centromeres (step 2) reshapes the karyotype following complex

repair events. These complex events include the loss of centromere DNA, isochromosome

formation, and *de novo* telomere formation (step 3), similar to what is observed during the

process of chromothripsis. The new karyotype can generate a reproductive barrier with the

parental isolate and lead to speciation. On the other hand, the strain with the rearranged

karyotype could mate with wild-type isolate, albeit at low frequency, leading to aneuploid

progeny (step 4), which can independently establish itself as a new species.