PLASMID CARRIAGE AND THE UNORTHODOX USE OF FITNESS IN MICROBIAL EVOLUTION

CARLOS REDING[1,2*]

[1]*Beckman Center for Molecular and Genetic Medicine, Department of Genetics, Stanford University. USA.*
[2]*School of Cellular and Molecular Medicine, Faculty of Life Sciences, University of Bristol. United Kingdom.*
[*]**Corresponding author:** reding@disroot.org

The link between fitness and reproduction rate is a central tenet in evolutionary biology: Mutants reproducing faster than the dominant wild-type are favoured by selection, but otherwise the mutation is lost. This link is given by Fisher's theorem, under the assumption that fitness can only change through mutations that boost or hinder growth rate, whence the use of logarithms on growth data by experimentalists. Here I show that logarithms are highly sensitive to sampling times, with fitness estimates changing through time, challenging our intuition about the outcome of microbial competitions: Mutants can be less fit, as fit, or fitter than the wild-type genotype by virtue of when the growth data is collected. Contrary to the aforementioned evolutionary principle, this variability reveals that 'unfit' mutants can be co-maintained if they reached their equilibrium. This is what I observed in competition assays between two *Escherichia coli* constructs, one of which harbours a non-transmissible plasmid that protects against tetracycline (pGW155B), without using antibiotic. Despite growing 40% slower than its drug-sensitive counterpart, the construct harbouring the plasmid persisted throughout the competition. And, perhaps more importantly, kept the plasmid. My study suggests that reliance on growth rate masks that selection on plasmid carriage may be stronger than previously thought—explaining the seemingly-paradoxical abundance of plasmids in nature.

I. INTRODUCTION

Population genetics [1–4] and microbiology [5, 6] textbooks define the fitness of microbial populations as their intrinsic growth rate. This notion stems from Fisher's theorem of natural selection where fitness (m) is given by the exponential model $N(t) = N_0 e^{mt}$ as he explicitly wrote in [7, page 26]. Here N_0 is the initial number of individuals of a given genotype, and $N(t)$ the number of individuals at time t. Thus, the ratio $W_{ij} = m_i/m_j$ gives the fitness of a genotype with respect to another (W_{ij}) [8]. Given the simplicity of Fisher's theorem, it is widely used to study microbial ecology and evolution [8–12], where fitness is measured through the use of logarithms on growth data from pairwise competition assays [8, 13–16]. Population genetics says [1] that if fitness, as intrinsic growth rate, is an inherited trait, it is reasonable to expect that mutants with greater fitness will be represented with greater proportion in the next generation—thus replacing the wild-type genotype [1, 7].

But microbes with slow growth abound in nature [17–20]. How is this possible? Take the abundance of plasmids, for example. Plasmids are extra-chromosomal elements that bacteria can acquire through horizontal gene transfer to thrive in unfavourable environments [21]. In favourable environments however, where plasmids do not add an advantage, they impose a fitness cost derived from the synthesis of additional genetic material during the replication and expression of

plasmid-borne genes [22]. Thus, conventional wisdom holds that plasmids should not exist without selection [21]. But they do, and are ubiquitous in nature [23], leading to a so-called *plasmid paradox* [21, 24]. Recent evidence showed that plasmids undergoing horizontal-gene transfer can propagate fast enough to persist despite the costs [25]. The problem is that most of the known plasmids are *not* transmissible [23]. Now the question is whether a myriad of conditions exist to select for plasmid-carriage, or the logic behind fitness estimates is sub-optimal. After all, the availability of data testing the predictions of population genetics theory is extremely scarce [26–28].

A corollary of Fisher's theorem is that the ratio m_i/m_j is constant, following the view that fitness is inheritable and therefore it can only change through mutations that affect m [8]. Below I show it is not and, rather, m_i/m_j changes through time. While this variability is mathematically unsurprising—resulting from the application of the exponential growth model on natural populations—it revealed that slow-growing genotypes could be co-maintained when competing against fitter genotypes that reproduce faster. Even without resorting to trade-offs between fitness and survival [29]. When applying these insights in the laboratory, I co-maintained a construct of *Escherichia coli* harbouring a non-transmissible plasmid, with a tetracycline resistance gene, in competition assays without using antibiotics. The constructs where not only co-maintained, despite the 40% reduction in growth rate imposed by the plasmid with: respect to its plasmid-free counterpart. The bacterium harbouring pGW155B kept the plasmid, even without exposure to tetracycline. In other words, the results suggest that there would be no such thing as a plasmid paradox. Rather, selection for plasmids, commonly studied through the application of Fisher's theorem and use of growth rate to estimate fitness, may be stronger than previously thought.

II.   RESULTS

Let me assume two competing genotypes, mutants A and B, that grow consistently with population genetics theory [1, 2]. The following system of ordinary differential equations describes the change in the number of individuals of each mutant over time:

$$\frac{dA}{dt} = m_a A, \quad \frac{dB}{dt} = m_b B, \tag{1}$$

where m is the population growth rate increase as introduced above, and $A(0) = B(0) = x$ where x is strictly positive. Note that m is derived from random birth (b) and death (d) rates, so that $m = b - d$ [1], and that it is independent of the number of individuals of each genotype. Each individual is equally likely to die or reproduce at any given time, and the age distribution is near to the equilibrium so that b and d are nearly constant. The growth dynamics of both mutants is illustrated in Figure 1A. If b and d are constant, m is a constant and therefore the relative fitness difference between genotypes $W_{ba} = m_b/m_a = k$ is also constant. In other words, if $m_b = \frac{ln[B(t)/B(0)]}{t}$ as derived from the exponential model introduced above [2, 8, 30] (see Methods), and $B(t)$ is the number of individuals of mutant B at an arbitrary time t, the relative fitness W_{ba} is the same regardless of t. The same applies to the fitness of A with respect to B. Indeed, if $m_b > m_a$, the number of mutant B individuals is higher than those of mutant A at all times as Figure 1A shows.

Now suppose that m depends on the number of individuals of each genotype. This is a reasonable assumption given that resources are depleted during microbial growth, and that resources are

90 finite, limiting the abundance of each mutant over time:

$$\frac{dA}{dt} = m_a\left(1 - \frac{A^*}{K_a}\right)A, \quad \frac{dB}{dt} = m_b\left(1 - \frac{B^*}{K_b}\right)B. \tag{2}$$

Here K_a and K_b are the maximal population size attainable (*carrying capacity*) for mutants A and
B, respectively, resulting from the limited availability of resources—note different efficiencies using
the *same* resource can lead to different carrying capacities [29]. Now, the population growth rate
m is corrected by the term $1 - (N_i/K_i)$. The Lotka-Volterra model of competition [31] includes
the term $N_i^* = N_i + \alpha_{ij}N_j$, where $\alpha_{ij}N_j$ is the linear reduction in growth rate—in terms of K—of
species i by competing species j. This formalisation describes the limitation in growth imposed by
the environment, due to finite resources, reducing m and the growth of both genotypes over time
(Figure 1B and C). In this scenario m_b is now corrected by changes in population size, including
that from mutant A, and therefore W_{ba} must depend on time:

$$W_{ba} = \frac{m_b}{m_a} \rightarrow W_{ba}(t) = \overbrace{\frac{m_b}{m_a} \cdot \frac{1}{1+pt}}^{\substack{\text{contribution of} \\ m \text{ to } W}} + \overbrace{\frac{K_b^*}{K_a^*} \cdot \frac{pt}{1+pt}}^{\substack{\text{contribution of} \\ K^* \text{ to } W}} \tag{3}$$

where W_{ba} is now corrected by changes in population size until both mutants reach the equilibrium
K^*—new since their carrying capacities cannot be realised in competition—and by p, a parameter
controlling the steepness of the change. Now, initially $W_{ba}(0) \approx m_b/m_a$ because $\frac{pt}{1+pt} \approx 0$ and $\frac{1}{1+t} \approx$
1. As time progresses W_{ba} declines exponentially, as Figures 1B and C show, with $W_{ba}(\infty) \approx K_b^*/K_a^*$
given that, now, $\frac{pt}{1+pt} \approx 1$ and $\frac{1}{1+pt} \approx 0$ reflecting the shift in the contribution to fitness from m to
K^*. Estimating fitness differences between two genotypes using equation (3), thus, requires prior
knowledge of the carrying capacity.

Curiously, in this scenario, W_{ba} is still calculated using $m_b = \frac{ln[B(t)/B(0)]}{t}$ and $W_{ba} = m_b/m_a$ in
the laboratory [8, 13–16, 32–34]. The result of measuring m_b using logarithms, following Fisher's

theorem, is illustrated in Figures 1B and C: Fitness declines exponentially over time—within the day—and whether a mutant is more or less fit than the reference genotype will be given by the choice of t, an arbitrary period during which genotypes are allowed to grow, and therefore $B(t)$. Thus, depending on the combination of carrying capacities, rates of population increase, and sampling time, mutant A may be fitter, as fit, or less fit than the reference genotype B. Whence the seemingly unpredictability of fitness [35]. While it can be argued that estimations of m aim to *only* capture the exponential growth, the rest of the growth dynamics should not be ignored given the interdependence of the competing mutants.

If, say, $m_b > m_a$ and $W_{ba} > 1$, which mutant, A or B, would be selected in pairwise competition assays? Despite the simplicity of equations (1) and (2), the answer is not straightforward. Figure 2 illustrates the change in relative abundance of two genotypes, A, and B, growing in competition for a common resource. These genotypes are haploid, with continuous growth and overlapping generations where m_a and m_b represent their population growth rates as introduced above. The competition is propagated at regular intervals c where the new initial conditions are given by $N_a(c_{+1}, 0) = N_a(c, t^*)d$, $N_b(c_{+1}, 0) = N_b(c, t^*)d$, with t^* being the time of growth allowed prior to the propagation and $0 \leq d \leq 1$ the dilution factor during the propagation step analog to experimental assays [8, 9, 15]. Following population genetics theory [1, 4, 5, 8], the outcome of this competition is straightforward: if, say, mutant B reproduces faster than mutant A ($m_b > m_a$), then $W_{ab} = m_a/m_b < 1$ and mutant A goes extinct (Figure 2A). Regardless of how long the competition is allowed to progress until the next propagation step, mutant A is always lost in competition with B. Only through the emergence of a new mutant A* with $m_a^* > m_b > m_a$ should this prediction change.

But when fitness changes through time the outcome of the competition becomes unclear, now depending on whether the competing genotypes reach their carrying capacities before the propagation: if the competition is propagated frequently, so that *both* genotypes are in active growth, the genotype reproducing faster will replace that reproducing slower consistently with Fisher's theorem. However, if one or both genotypes are allowed to reach their carrying capacities, both will be be co-maintained regardless of the in rates of population increase (Figures 2B and C). Here the relative frequency will be given by their carrying capacities K, so if reproduction and carrying capacity engage in a trade-off [29] for one genotype, this will be most frequent (Figure 2B). Importantly, these predictions are maintained regardless of the initial frequency of A (Figure S1), consistently with prior data [36].

To test this prediction, I competed two constructs of *Escherichia coli* MC4100, Wyl and GB(c) (see Methods), and measured their variation in fitness through time. Figure 3A shows that both constructs reach their carrying capacicies within 24 hours. The construct GB(c) carries the non-transmissible plasmid pGW155B [37] harbouring *tet(36)*—a ribosome-protection type resistance gene against tetracycline [37]. This plasmid lacks a *rep* gene to control tightly the partition of plasmids after cell division [38–40] (addgene vector database accession #2853). While pGW155B is a synthetic plasmid, many natural plasmids also lack partition systems [40, 41]. Both constructs have identical chromosome with exception of this plasmid, and the fluorescence gene they carry: cyan (*cfp*, GB(c)) or yellow (*yfp*, Wyl), to allow their identification in competition assays. Using a non-transmissible plasmid prevents cross-contamination by horizontal gene transfer between the resistant construct GB(c) and Wyl. Importantly, the use of constructs derived from MC4100 avoids interactions between competitors that may affect the outcome of the competition for reasons

143 **Figure 2. Change in relative frequency during competition depending on the length of the growth cycle.** Changes in
144 frequency over time for mutants A (cyan) and B (black) following propagations at arbitrary times t=6h, t=12h, t=18h, and t=24h.
145 These times correspond to different stages of growth as Figure 1 shows. Each plot illustrate the case for genotypes with density-
146 independent growth (A), or density-dependent growth where a genotype does engage in a trade-off between reproduction and
148 survival (B) and when it does not (C). Parameter used in the model are those from Figure 1.

161 beyond pGW155B, like the production of bacteriocins [42].

pGW155B penalised the population growth rate of construct GB(c) by approximately 40%
compared to Wyl (Figure 3B) as I found out using pure cultures. It might be argued that the
4-parameter logistic model I used does not fully capture the growth data. However, if I measure
the maximum change between the reads (Euler's method, see Methods), the strong difference in
growth rate is maintained. The duration of lag, and carrying capacity were also sensitive to plas-
mid carriage, regardless of whether I measured the growth using fluorescence or light scattering
(Figure S2). Changes in cell size could confound optical density and fluorescence readings, but this

Figure 3. Asymmetric carriage costs of pGW155B. A) Growth curves of constructs Wyl (black) and GB(c) (cyan) in pure cul-
ture, in the absence of tetracycline, over 24h. I estimated the *per capita* growth rate (m), population size in the equilibrium
(K), biomass yield, and lag duration from a 4-parameter logistic model fitted to growth data (see Methods). Data fits for con-
structs Wyl and GB(c) are shown in grey and light cyan, respectively. **B)** Box plots showing the median (centre of the box), 25th
and 75th percentile for each of the aforementioned parameters. The whiskers extend to the most extreme data points that are
not outliers, and these are individually represented. The p value shown refers to a Two-sample t-test with unequal variance (8
replicates) that I used to test differences in the parameters between both constructs. **C)** Alternative metrics for growth rate and
biomass yield: forward Euler approximation (top), data fit predicted yield (middle), and maximal yield across all time points in
growth data (bottom). The p values correspond to Two-sample t-tests with unequal variance (8 replicates).

phenomenon leaves a signature [43] in growth data that is absent in my dataset (Figures S2, S3,
and S4). The change in carrying capacity K can be linked to an increase in biomass yield (y) as
Monod's expression [44] $y = K/S$ suggests, where S the supply of glucose. This metric is consistent
with the data, given that construct GB(c) has to express and translate plasmid-borne genes with
the same supply of glucose (Figures 3B and C). This means that pGW155B triggers a metabolic
trade-off between growth rate and biomass yield redolent of rK-selection theory [29, 45].

Figure 4. Time-dependent fitness data allows co-maintenance of both constructs without tetracycline. Relative fitness
during first 24 hour pairwise-competition of construct Wyl, where both constructs grew in media supplemented with 0.04 μg/mL
of tetracycline (**A**) and without drug (**D**). Fitness for each replicate is presented in grey with respect to the reference GB(c) (black
dashed line), and the prediction from equation 3 in red with an adjusted R^2 of 0.308 (**A**, F-statistic against zero model =640, $p \ll$
10^{-100}) and 0.612 (**D**, F-statistic against zero model = 3360, $p \ll 10^{-100}$). The goodness of fit constrained to the equilibrium
phase can be found in Figure S5. The inset illustrate the growth curves for both constructs, highlighting in grey the period
where the relative fitness of Wyl (W_{sr}) is lower than its tetracycline-resistant competitor's. **B)** and **E)** show the change in relative
frequency of both constructs through time when both constructs grew in media supplemented with tetracycline (B) and without
antibiotic (E). Error bars represent the mean ± 95% confidence intervals, with raw data points shown as dots. **C)** and **F)** show
the change in relative copy numbers of pGW155B borne by construct GB(c) during the pairwise-competition, calculated using
quantitative PCR (see Methods). Error bars represent the mean ± standard deviation, with raw data shown in red.

For the competition assay, I mixed equal proportions (cell/cell) of two pure cultures, from con-
struct GB(c)—harbouring pGW155B—and Wyl respectively, grown overnight, in media containing
no antibiotic or 0.04 μg/mL of tetracycline (see Methods). I incubated the mixed culture at 30^{o}C
until both constructs reached their carrying capacity (K, ~ 24h as per data in Figure 3A), and then
I propagated the competition into a new plate containing fresh media. I repeated the propagation
step seven times totalling between 77 (GB(c)) and 119 (Wyl) generations given the reproduction
rates in Figure 3B, where the generation time per hour is given by $m_i/24$, multiplied times the num-
ber of propagation steps. Considering the mutation rate of enterobacteria [46] and the emergence
of mutants after 3 growth cycles [47], I constrained the fitness measurements in both constructs
over the first 24h of competition and compared the outcome of the competition with predictions
in Figure 2. Indeed, the relative fitness of both constructs changed through time consistently with
Figure 1B and C. In the presence of tetracycline, the relative fitness of drug-sensitive, plasmid-free
construct Wyl was below GB(c), which harbours pGW155B, at all times as Figure 4A illustrates.
This meant that GB(c) increased its relative frequency in following propagation steps and became

the most abundant construct throughout the competition (Figure 4B) consistently with the theo-
retical predictions. Now, the model does not implement any mutation processes but these occur
during this assay. For example, the number of copies of pGW155B harboured by GB(c) increased
5- to 6-fold during the competition (2-sample t-test with unequal variances, p=0.0088, df=2.5826,
t-statistic=-7.2441; with 3 replicates) driven by exposure to tetracycline (Figure 4C) as it would be
reasonable to expect given the exposure to tetracycline. Despite the occurrence of these mutations,
Wyl never goes extinct consistently with Figure 2 and its proportion is determine by its abundance
at the propagation time.

Similarly, when both constructs grew in mixed cultures, now without tetracycline exposure,
the relative fitness of Wyl was higher, equal, or lower than plasmid-harbouring construct GB(c)
through time, as Figure 4D shows. Consistently with the predictions both constructs where co-
maintained, and with similar relative frequencies, despite the difference in growth rate (Figure
4E). Note here that while the initial deviation is small, without strong selection by antibiotic ex-
posure each replicate evolves differently, with some replicates of Wyl being lost, while others are
co-maintained. In other words, GB(c) preserved pGW155B despite the lack of tetracycline and
the difference in growth rates (2-sample t-test with unequal variances, p=0.2305, df=2.2351, t-
statistic=-1.6353; with 3 replicates. Figure 4F). This contrasts with previous studies [15] predicting
a rapid decline, within a similar time frame, in cells harbouring non-transmissible plasmids.

III.   DISCUSSION

It is important to stress that fitness, understood as growth rate following Fisher's rationale, indeed
varies through evolutionary timescales due to mutations [8, 48]. The scope of this manuscript is
the variations in fitness that occur at smaller timescales, where fitness is assumed constant and
often used to make evolutionary predictions [13–16, 49, 50]—which only in the case of chemostats
[51] would be appropriate given the constants state of growth and environment they impose on
cells.

Fitness, as measured in the laboratory, is known to be sensitive to t^* [52] but it is often pre-
sented as an experimental quirk: If t^* is too long one competitor may drive the other to extinction,
whereas if t^* is too short the measurement may not be precise [52]. But this variation has also
been used to suggest the co-existence of multiple populations through frequency-dependent fitness
[36]—which also depends on t^* [36]. To explain this phenomenon, some studies [53, 54] argue
that each growth phase contributes to fitness. But a closer look to the models proposed exposes
that growth rate remains the main contributor. Equation 3 provides a more robust measurement
of relative fitness, given the correction of m by changes in population size, whereas the ratio of
growth rates m_i/m_j would be more appropriate with chemostats where Fisher's assumptions are
closely met.

If the fitness measured experimentally varies within a season or cycle, it means slow-growing
microorganisms are mistakenly assumed to go extinct without positive selection. What I suggest
here is that, if all genotypes reach their equilibrium before the selection event, then they can be
co-maintained even in the absence of mutations—explaining the maintenance of pGW155B. Even
despite differences in carrying capacities, given the persistence of the drug-sensitive construct Wyl
shown in Figure 4B. Indeed, the use of more tetracycline would have cleared Wyl. The point is
that, despite the substantial difference in fitness, the population of Wyl levelled off and was co-

maintained consistently with the predictions in Figure 2. Only emerging mutants that are unable to reach their equilibrium before the selection step, would be lost to selection—which data from Figure 4E suggest might have occurred in one of the replicates. The lack of mutations assumed here is, naturally, unrealistic. But it sets the expectation that a mutant is co-maintained if it can reach an equilibrium, rather than be lost to selection because of its lower fitness as derived from Fisher.

It is noteworthy to mention that rk selection postulates [45] that selection acts on populations in a manner that is contingent on their size—on evolutionary timescales—ultimately leading to fixation or extinction events. But here I argued, and observed, that populations can be co-maintained despite their size at the equilibrium.

The abundance of plasmids in nature is commonly studied through the application of Fisher's theorem [15, 24]. My study suggests that, given the limitations of this theorem and its experimental implementation, selection for plasmids could be higher than previously thought. Importantly, another notion derived from this work is the need to account for the growth stage when using fitness estimates to make evolutionary predictions. While selection events that occur during the growth phase fall within Fisher's theorem, and therefore it would be reasonable to expect the fixation or extinction of the new mutant; selection events that occur after equilibrium are likely to lead to the co-maintenance of mutant and wild-type.

IV. METHODS

Media and Strains. I used the strains of *Escherichia coli* GB(c) and Wyl [55] (a gift from Remy Chait and Roy Kishony), and M9 minimal media supplemented with 0.4% glucose and 0.1% casamino acids (w/v). I made tetracycline stock solutions from powder stock (Duchefa, Ref. #0150.0025) at 5mg/mL in 50% ethanol, filter sterilised, and stored at -20^oC. Subsequent dilutions were made from this stock in supplemented M9 minimal media and kept at 4^oC.

Batch transfer protocol. I inoculated a 96-well microtitre plate containing 150 µg/mL of supplemented M9 media with a mixture of two overnight cultures, one of *E. coli* GB(c) and another of *E. coli* Wyl (1µL containing approx. $2 \cdot 10^6$ cells, Figure S3). The overnight culture for GB(c) was supplemented with 100ng/mL of tetracycline to preserve the plasmid pGW155B carrying *tet(36)* as described elsewhere [9], then centrifuged and removed prior adding to the microtitre plate. I incubated the plate at 30^oC in a commercial spectrophotometer and measured the optical density of each well at 600nm (OD_{600}), yellow florescence for the Wyl strain (YFP excitation at 505nm, emission at 540nm), and cyan fluorescence for the GB(c) strain (CFP at 430nm/480nm) every 20min for 24h. After each day I transferred 1.5µL of each well, using a 96-well pin replicator, into a new microtitre plate containing fresh growth medium and tetracycline.

Growth parameter estimation. Yellow and cyan fluorescence protein genes were constitutively expressed given the constant ratio between fluorescence and optical density (Figure S6). This allowed me to use fluorescence data as a proxy for cell density in mixed culture conditions. I normalised fluorescence readings using a conversion factor, n_f, calculated by growing each construct in a range of glucose concentrations, and regressing the linear model RFU = $n_f \cdot$ OD + c, where RFU is relative fluorescence units data, OD optical density data, n_f the conversion factor between

fluorescence and optical density, and c the crossing point with the y–axis when OD = 0. I imported the resulting time series data set (Figures S4) into `MATLAB R2014b` to subtract background and calculate fitness as described in the main text as follows.

Once corrected, I used the expression $m_i = \frac{ln[N_i(t)/N_i(0)]}{t}$ to estimate the *per capita* growth rate m_i as per in [8, 13–16] although I used different values for t to test the theoretical predictions. This expression derives from the exponential model in equation 1 as it follows. We can isolate the term with m for, say, mutant A, as $A(t) = A_0 e^{m_a t} \rightarrow \frac{A(t)}{A_0} = e^{m_a t}$. To remove the number e we then apply logarithms to this expression leading to $ln\frac{A(t)}{A_0} = m_a t$. Re-arranging the terms we are left with $\frac{ln[A(t)/A_0]}{t} = m_a$, which is the expression used by experimentalists.

To measure the carriage costs of pGW155B, I fitted to data the logistic model

$$g(t) = \frac{b + K}{(1 + C \cdot e^{-m \cdot [t-l]})}.$$

where the term $g(t)$ denotes the change in culture growth as a function of time (in OD, YFP, or CFP units), b the inoculum size used to subtract the background, C is a parameter, m the *per capita* growth rate, K the maximal population size attained and l the duration of the lag phase. Alternatively, I measured the rate of increase using the forward Euler approximation $m_i = max_{0 \le t \le 24h}\left(\frac{N_i(t)/N_i(t-1)}{\Delta t}\right)$, where $\Delta t = 20\text{min} = 1/3$ is the time between reads. To estimate the biomass yield I divided OD data in stationary phase by the glucose supplied [56]. Alternatively, I used the highest density across all time points and the density reported by the data fit, both divided by glucose supply, as alternative metrics for biomass yield.

Finally, I divided the length of the growth cycle (24h) by the *per capita* growth rate, multiplied by the number of propagations (seven) to estimate the total number of duplication events—where one duplication means one generation—for each construct.

DNA material extraction. For each concentration, I sampled three representative 150 µg/mL replicates that I divided in two groups: for chromosome and plasmid DNA extraction. I used 'GeneJet DNA' (ThermoScientific, Ref. #K0729) and 'GeneJet Plasmid' (ThermoScientific, Ref. #K0502) extraction kits to extract chromosomal and plasmid DNA (pDNA) from the samples, respectively, and used Qubit to quantify DNA and pDNA yields. Both extracts were diluted accordingly in extraction buffer, as per in manufacturer instructions, to normalise DNA across samples.

Quantitative PCR and plasmid copy number estimation. I used `primer3` to design two pairs of primers with melting temperature (T_m) of 60^oC and non-overlapping probes with T_m of 70^oC. The amplicon ranges between 100 to 141bp depending on the locus (Table 1). Two reaction mixes were prepared using the kit 'Luminaris Color Probe Low ROX' (ThermoScientific, Ref. #K0342), adding 0.3µM of each primer and 0.2µM of the probe as per manufacturer specifications. Following a calibration curve for each reaction (Figure S7) I added 0.01ng of chromosomal or plasmid DNA material to each of the reaction mixes.

To estimate the relative copies of pGW155B per GB(c) cell, I calculated the corresponding proportion of chromosomal DNA corresponding to the GB(c)-type from data in Figure 4 and used the expression [15]

$$cn = \frac{(1 + E_c)^{Ctc}}{(1 + E_p)^{Ctp}} \times \frac{S_c}{S_p},$$

Table 1. Primers and probes designed using `Primer3`. Amplicon ranging from 100 to 141bp. T_m indicates the estimated melting temperature.

Target gen	Sequence ($5' \rightarrow 3'$)	T_m (^{o}C)	Feature
tatB	CGATGAAGCGTTCCTACGTT	60.27	Forward
	TCATGCGCAGCTTCATTATC	59.94	Reverse
	AAGGCGAGCGATGAAGCGCA	70.70	Probe
tet(36)	ATTGGGCATCTATTGGCTTG	59.22	Forward
	CCGATTCACAGGCTTTCTTG	60.76	Reverse
	AGCCTTTGCCAATTGGGGCG	70.37	Probe

where cn is the number of plasmid copies per chromosome, S_c and S_p are the size of the chromosome and pGW155B amplicon in bp, E_c and E_p the efficiency of the qPCR taken from data in Figure S7, and Ctc and Ctp are the cycles at which I first detected product amplification (C_t).

REFERENCES

1. Crow, J. F., Kimura, M., *et al. An introduction to population genetics theory.* (New York, Evanston and London: Harper & Row, Publishers, 1970).

2. Nagylaki, T. in *Introduction to Theoretical Population Genetics* 5–27 (Springer, 1992).

3. Rice, S. H. *Evolutionary theory: mathematical and conceptual foundations* isbn: 9780878937028 (Oxford University Press, 2004).

4. Templeton, A. *Population Genetics and Microevolutionary Theory* isbn: 9780471409519 (Wiley, 2006).

5. Dawes, I. & Sutherland, I. *Microbial Physiology* isbn: 9780632024636 (Wiley, 1992).

6. Borriello, S., Murray, P. & Funke, G. *Topley and Wilson's Microbiology and Microbial Infections, 2 Volume Set: Bacteriology* isbn: 9780470660263 (Wiley, 2007).

7. Fisher, R. A. *The Genetical Theory of Natural Selection* 308–315 (At The Clarendon Press, 1930).

8. Lenski, R. E., Rose, M. R., Simpson, S. C. & Tadler, S. C. Long-Term Experimental Evolution in Escherichia coli. I. Adaptation and Divergence During 2,000 Generations. *Am. Nat.* **138,** pp. 1315–1341 (1991).

9. Chait, R., Craney, A & Kishony, R. Antibiotic interactions that select against resistance. *Nature* **446,** 668–671 (2007).

10. Wang, H., Avican, K., Fahlgren, A., *et al.* Increased plasmid copy number is essential for Yersinia T3SS function and virulence. *Science* **353,** 492–495 (2016).

11. Adkar, B. V., Manhart, M., Bhattacharyya, S., Tian, J., Musharbash, M. & Shakhnovich, E. I. Optimization of lag phase shapes the evolution of a bacterial enzyme. *Nat. Ecol. Evol.* **1,** 1–6 (2017).

12. Concepción-Acevedo, J., Weiss, H. N., Chaudhry, W. N. & Levin, B. R. Malthusian parameters as estimators of the fitness of microbes: a cautionary tale about the low side of high throughput. *PLoS One* **10,** e0126915 (2015).

13. Andersson, D. I. & Levin, B. R. The biological cost of antibiotic resistance. *Curr. Opin. Microbiol.* **2,** 489–493 (1999).

14. Gullberg, E., Cao, S., Berg, O. G., *et al.* Selection of resistant bacteria at very low antibiotic concentrations. *PLoS Pathog.* **7,** e1002158 (2011).

15. Millán, A. S., Peña-Miller, R., Toll-Riera, M, *et al.* Positive selection and compensatory adaptation interact to stabilize non-transmissible plasmids. *Nat. Commun.* **5** (2014).

16. Ram, Y., Dellus-Gur, E., Bibi, M., *et al.* Predicting microbial growth in a mixed culture from growth curve data. *Proc. Natl. Acad. Sci. U.S.A.* **116,** 14698–14707 (2019).

17. Kussell, E., Kishony, R., Balaban, N. Q. & Leibler, S. Bacterial persistence: a model of survival in changing environments. *Genetics* **169,** 1807–1814 (2005).

18. Pedrós-Alió, C. Marine microbial diversity: can it be determined? *Trends Microbiol.* **14,** 257–263 (2006).

19. Lever, M. A., Rogers, K. L., Lloyd, K. G., *et al.* Life under extreme energy limitation: a synthesis of laboratory-and field-based investigations. *FEMS Microbiol. Revs.* **39,** 688–728 (2015).

20. Flemming, H.-C. & Wuertz, S. Bacteria and archaea on Earth and their abundance in biofilms. *Nat. Rev. Microbiol.* **17,** 247–260 (2019).

21. Harrison, E. & Brockhurst, M. A. Plasmid-mediated horizontal gene transfer is a coevolutionary process. *Trends Microbiol.* **20,** 262–267 (2012).

22. Lili, L. N., Britton, N. F. & Feil, E. J. The persistence of parasitic plasmids. *Genetics* **177,** 399–405 (2007).

23. Smillie, C., Garcillán-Barcia, M. P., Francia, M. V., Rocha, E. P. & de la Cruz, F. Mobility of plasmids. *Microbiol. Mol. Biol. Rev.* **74,** 434–452 (2010).

24. Probing the plasmid paradox. *Nat. Ecol. Evol.* **5,** 1559–1559 (2021).

25. Lopatkin, A. J., Meredith, H. R., Srimani, J. K., Pfeiffer, C., Durrett, R. & You, L. Persistence and reversal of plasmid-mediated antibiotic resistance. *Nat. Commun.* **8,** 1689 (2017).

26. Wakeley, J. The limits of theoretical population genetics. *Genetics* **169,** 1–7 (2005).

27. Kokko, H. & López-Sepulcre, A. The ecogenetic link between demography and evolution: can we bridge the gap between theory and data? *Ecol. Lett.* **10,** 773–782 (2007).

28. Orr, H. A. The population genetics of beneficial mutations. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **365,** 1195–1201 (2010).

29. Reding-Roman, C., Hewlett, M., Duxbury, S., Gori, F., Gudelj, I. & Beardmore, R. The unconstrained evolution of fast and efficient antibiotic-resistant bacterial genomes. *Nat. Ecol. Evol.* **1,** 0050 (2017).

30. Hansen, S. K., Rainey, P. B., Haagensen, J. A. & Molin, S. Evolution of species interactions in a biofilm community. *Nature* **445,** 533–536 (2007).

31. Gilpin, M. E. & Ayala, F. J. Global models of growth and competition. *Proc. Natl. Acad. Sci. U.S.A.* **70,** 3590–3593 (1973).

32. Lenski, R. E. Experimental studies of pleiotropy and epistasis in Escherichia coli. I. Variation in competitive fitness among mutants resistant to virus T4. *Evolution* **42,** 425–432 (1988).

33. Wiser, M. J., Ribeck, N. & Lenski, R. E. Long-term dynamics of adaptation in asexual populations. *Science* **342,** 1364–1367 (2013).

34. Lenski, R. E., Wiser, M. J., Ribeck, N., *et al.* Sustained fitness gains and variability in fitness trajectories in the long-term evolution experiment with Escherichia coli. *Proc. R. Soc. B.* **282,** 20152292 (2015).

35. Hinz, A., Amado, A., Kassen, R., Bank, C. & Wong, A. Unpredictability of the fitness effects of antimicrobial resistance mutations across environments in Escherichia coli. *bioRxiv.* `https://www.biorxiv.org/content/early/2023/10/27/2023.10.25.563906` (2023).

36. Ribeck, N. & Lenski, R. E. Modeling and quantifying frequency-dependent fitness in microbial populations with cross-feeding interactions. *Evolution* **69,** 1313–1320 (2015).

37. Whittle, G, Whitehead, T. R., Hamburger, N, Shoemaker, N. B., Cotta, M. A., *et al.* Identification of a new ribosomal protection type of tetracycline resistance gene, tet(36), from swine manure pits. *Appl. Environ. Microbiol.* **64,** 4151–4158 (2003).

38. Del Solar, G. & Espinosa, M. Plasmid copy number control: an ever-growing story. *Mol. Microbiol* **37,** 492–500 (2000).

39. Garcillán-Barcia, M. P., Alvarado, A. & de la Cruz, F. Identification of bacterial plasmids based on mobility and plasmid population biology. *FEMS Microbiol. Rev.* **35,** 936–956 (2011).

40. Million-Weaver, S. & Camps, M. Mechanisms of plasmid segregation: have multicopy plasmids been overlooked? *Plasmid* **75,** 27–36 (2014).

41. Guynet, C. & de la Cruz, F. Plasmid segregation without partition. *Mob. Genet. Elements* **1,** 236–241 (2011).

42. Dobson, A., Cotter, P. D., Ross, R. P. & Hill, C. Bacteriocin Production: a Probiotic Trait? *Appl. Environ. Microbiol.* **78,** 1–6 (2012).

43. Stevenson, K., McVey, A. F., Clark, I. B., Swain, P. S. & Pilizota, T. General calibration of microbial growth in microplate readers. *Sci. Rep.* **6,** 1–7 (2016).

44. Monod, J. *Recherches sur la croissance des cultures bactériennes* PhD thesis (Paris, 1941).

45. Reznick, D., Bryant, M. J. & Bashey, F. r-and K-selection revisited: the role of population regulation in life-history evolution. *Ecology* **83,** 1509–1520 (2002).

46. Perfeito, L., Fernandes, L., Mota, C. & Gordo, I. Adaptive mutations in bacteria: high rate and small effects. *Science* **317,** 813–815 (2007).

47. Reding, C., Catalán, P., Jansen, G., *et al.* The Antibiotic Dosage of Fastest Resistance Evolution: gene amplifications underpinning the inverted-U. *Mol. Biol. Evol.* **9,** 3847–3863. issn: 0737-4038 (Mar. 2021).

48. Cao, H. & Plague, G. R. The fitness effects of a point mutation ins *Escherichia coli* change with founding population density. *Genetica* **144,** 417–424 (2016).

49. Bergstrom, C. T., Lipsitch, M. & Levin, B. R. Natural selection, infectious transfer and the existence conditions for bacterial plasmids. *Genetics* **155,** 1505–1519 (2000).

50. Blanquart, F., Achaz, G., Bataillon, T. & Tenaillon, O. Properties of selected mutations and genotypic landscapes under Fisher's geometric model. *Evolution* **68,** 3537–3554 (2014).

51. Veldkamp, H & Kuenen, J. The chemostat as a model system for ecological studies. *Bull. Ecol. Res. Comm.,* 347–355 (1973).

52. McDonald, M. J. Microbial experimental evolution–a proving ground for evolutionary theory and a tool for discovery. *EMBO Rep.* **20,** e46992 (2019).

53. Li, Y., Venkataram, S., Agarwala, A., *et al.* Hidden complexity of yeast adaptation under simple evolutionary conditions. *Current Biology* **28,** 515–525 (2018).

54. Lin, J., Manhart, M. & Amir, A. Evolution of microbial growth traits under serial dilution. *Genetics* **215,** 767–777 (2020).

55. Chait, R., Shrestha, S., Shah, A. K., Michel, J.-B. & Kishony, R. A Differential Drug Screen for Compounds That Select Against Antibiotic Resistance. *PLoS One* **5,** e15179 (2010).

56. Monod, J. The Growth of Bacterial Cultures. *Annu. Rev. Microbiol.* **3,** 371–394 (1949).

Competing interests. The author declares no competing interests.

Data availability statement Data used to generate Figures 3 and 4 is available as a supplementary table. An Octave/MATLAB implementation of equations 1 and 2 is available at
https://github.com/rc-reding/papers/tree/master/FisherTheorem.

I. SUPPLEMENTARY FIGURES

Figure S1. Relative fitness of mutant A at different densities. A–B) Relative fitness of mutant A measured after seven prop-
agation steps, analog to those described in the main manuscript, when growth rate and population size engage in a trade-off
(A) and when they do not (B). The frequency on the x-axis is that of mutant A at t=0. The fitness of each competitor (m_i) was
calculated as explained in the main manuscript as $m_i = \frac{ln[N_i(t)/N_i(0)]}{\Delta t}$ where $N_i(t)$ is the density of each mutant at time $t = 7$,
$N_i(0)$ that at time t=0, and Δt = 7. This was used to calculate the relative fitness of A as $W_{ab} = m_a/m_b$ as described in the
text. The code is available in github in the link provided in the text, and the parameters used are those from Figure 1B and C.
C–D) Relative frequency of mutant A at the end of the seventh propagation when growth rate and population size engage in a
trade-off (C) and when they do not (D), shown in the y-axis, as a function of its frequency at t=0.

 Figure S2. Asymmetric carriage costs of pGW155B (optical density data). A) Growth curves of constructs Wyl (black) and
GB(c) (cyan) in pure culture, in the absence of tetracycline, over 24h. I estimated the maximum growth rate (r), population size
in the equilibrium (K), biomass yield, and lag duration from a 4-parameter logistic model fitted to growth data (see Methods).
Data fits for constructs Wyl and GB(c) are shown in grey and light cyan, respectively. **B)** Box plots showing the median (centre
of the box), 25th and 75th percentile for each of the aforementioned parameters. The whiskers extend to the most extreme
data points that are not outliers, and these are individually represented. The p value shown refers to Two-sample t-tests with
unequal variance (8 replicates) that I used to test differences in the parameters between both constructs. **C)** Alternative metrics
for growth rate and biomass yield: forward Euler approximation (top), data fit predicted yield (middle), and maximal yield across
all time points (bottom). The p values correspond to Two-sample t-tests with unequal variance (8 replicates).

Figure S3. Calibration curve to translate optical density data to number of *Escherichia coli* cells. I fitted the linear model
$a = bx + c$ to optical density and colony counting data (dots) to calculate the number of optical density units (OD_{600}) per cell. a
denotes the optical density readings measured at 600nm, c the crossing point with the $y-$axis when $x = 0$, and b the conversion
factor between optical density and number of cells (x). I interpolating optical density readings to calculate the number of cells
within a culture as $x = (a - c)/b$. For the strain Wyl, $b = 1.62 \times 10^{-10} \, OD \cdot mL \cdot CFU^{-1}$ and $c = 1.78 \times 10^{-2} \, OD$, whereas for
GB(c) $b = 1.79 \times 10^{-10} \, OD \cdot mL \cdot CFU^{-1}$ and $c = 1.33 \times 10^{-2} \, OD$.

Figure S4. Raw data of each construct during the competition. Raw optical density data for GB(c) (cyan) and Wyl (black), measured every 20 min and converted from fluorescence readings. Each individual plot represents the change in abundance of each construct over 24 hours. Data corresponding to each condition, with and without tetracycline, are aggregated as columns.

Figure S5. Time-dependent fitness data allows co-maintenance of both constructs without tetracycline. Relative fitness during first 24 hour pairwise-competition of construct Wyl, where both constructs grew in media supplemented with 0.04 µg/mL of tetracycline (**A**) and without drug (**B**). Fitness for each replicate is presented in grey with respect to the reference GB(c) (black dashed line), and the prediction from equation 3 in red with an adjusted R^2 of 0.308 (**A**, F-statistic against zero model =640, $p \ll 10^{-100}$) and 0.612 (**B**, F-statistic against zero model =3360, $p \ll 10^{-100}$). The inset illustrate the growth curves for both constructs, highlighting in grey the period where the relative fitness of Wyl (W_{sr}) is lower than its tetracycline-resistant competitor's.

Figure S6. Changes in relative fluorescence over time in GB(c) and Wyl strains in pure culture conditions. Raw change in florescence, per optical density units, measured every 20min for 24h for the Wyl- (black) and GB(c)-type. Each column represents the data set for each tetracycline concentration used.

Figure S7. Quantitative PCR calibration curves for *tet(36)* and *tatB*. Reaction efficiency for the set of primers and probes, listed in Table 1 the main Methods section, for *tet(36)* (A) and *tatB*. The efficiency was calculated as $E_f = 10^{-1/Slope} - 1$, and the slope term calculated by fitting a linear model to qPCR threshold cycle (C_t) data. The mean $\pm$ standard deviation for the adjusted coefficient of determination R^2 and efficiency are shown in the figures. The amplification curves for each reaction, using 3 replicates, are shown in C) and D), respectively.