PLASMID CARRIAGE AND THE UNORTHODOX USE OF FISHER'S THEOREM IN GENETICS

CARLOS REDING[1,2*]

[1]*Department of Genetics, Stanford University. Stanford, CA 94305, USA.*
[2]*Faculty of Life Sciences, University of Bristol. BS2 8BH Bristol, United Kingdom.*
Corresponding author: reding@disroot.org

The link between fitness and reproduction rate is a central tenet in evolutionary biology: mutants reproducing faster than the dominant wild-type are favoured by selection, otherwise the mutation is lost. This link is given by Fisher's theorem under the assumption that fitness can only change through mutations. Here I show that fitness, as formalised by Fisher, changes through time without invoking new mutations—allowing the co-maintenance of fast- and slow-growing genotypes. The theorem does not account for changes in population growth that naturally occur due to resource depletion, but it is key to this unforeseen co-maintenance of growth strategies. To demonstrate that fitness is not constant, as Fisher's theorem predicates, I co-maintained a construct of *Escherichia coli* harbouring pGW155B, a non-transmissible plasmid that protects against tetracycline, in competition assays without using antibiotics. Despite growing 40% slower than its drug-sensitive counterpart, the construct with pGW155B persisted throughout the competition without tetracycline—maintaining the plasmid. Thus, predicting the selection of microbial genotypes may be more difficult than previously thought.

I. INTRODUCTION

Textbook population genetics [1–3] and textbook microbiology [4, 5] define the fitness of microbial populations as the intrinsic population growth rate. This notion stems from Fisher's theorem of natural selection where fitness (m) is given by the exponential model $N(t) = N_0 e^{mt}$ as he explicitly wrote in [6, page 26]. Here N_0 is the initial number of individuals of a given genotype, and $N(t)$ the number of individuals at time t. Thus, the ratio $W_{ij} = m_i/m_j$ gives the fitness of a genotype with respect to another (W_{ij}). Given the simplicity of Fisher's theorem, it is widely used to study microbial ecology and evolution [7–11], where fitness is measured through pairwise competition assays [7, 12–15]. If fitness is exclusively given by population growth rate, and it is an inherited trait, it is reasonable to expect that mutants with greater fitness will be represented with greater proportion in the next generation thereby replacing the wild-type genotype [1, 6].

However, slow growth strategies abound among microbes [16–19]. The ubiquity of plasmids, for example, cannot be explained with this paradigm given the reduction in fitness they impose [20–22]. Indeed, plasmids undergoing horizontal-gene transfer can

propagate fast enough to exist despite the costs [23], but most of the known plasmids are non-transmissible [20]. This discrepancy can be explained by analysing Fisher's theorem. Fisher assumes the ratio m_i/m_j to be constant, following the view that fitness is inheritable and therefore it can only change through mutations that affect m [7]. Below I show it is not, exposing conditions where slow reproducing genotypes can be maintained through time when competing against genotypes that reproduce faster. Even in the absence of trade-offs between reproduction and survival [24]. Relying on my new analysis of Fisher's theorem, I co-maintained a construct of *Escherichia coli* harbouring a non-transmissible plasmid, with a tetracycline resistance gene, in fitness competition assays without using antibiotics. The plasmid imposed a 40% reduction in growth rate with respect to its plasmid-free counterpart, but the bacterium kept it even without tetracycline. This, in turn, suggests that selection for plasmids, commonly studied through Fisher's theorem, may be stronger than previously thought—explaining their ubiquity in nature.

II.   RESULTS

Let me assume two competing genotypes, mutants A and B, that grow consistently with population genetics theory [1, 2]. The following system of ordinary differential equations describes the change in the number of individuals of each mutant over time:

$$\frac{dA}{dt} = m_a A, \quad \frac{dB}{dt} = m_b B, \tag{1}$$

where m is the population growth rate increase as introduced above, and $A(0) = B(0) = x$ where x is strictly positive. Note that m is derived from random birth (b) and death (d) rates, so that $m = b - d$ [1], and that it is independent of the number of individuals of each genotype. Each individual is equally likely to die or reproduce at any given time, and the age distribution is near to the equilibrium so that b and d are nearly constant. The growth dynamics of both mutants is illustrated in Figure 1A. If b and d are constant, m is a constant and therefore the relative fitness difference between genotypes $W_{ba} = m_b/m_a = k$ is also constant. In other words, if $m_b = \frac{ln[B(t)/B(0)]}{t}$ as derived from the exponential model introduced above [2, 7, 25] (see Methods), and $B(t)$ is the number of individuals of mutant B at an arbitrary time t, the relative fitness W_{ba} is the same regardless of t. The same applies to the fitness of A with respect to B. Indeed, if $m_b > m_a$, the number of mutant B individuals is higher than those of mutant A at all times as Figure 1A shows.

Now suppose that m depends on the number of individuals of each genotype. This is a reasonable assumption given that resources are depleted during microbial growth, and that resources are finite, limiting the abundance of each mutant over time:

$$\frac{dA}{dt} = m_a \left(1 - \frac{A^*}{K_a}\right) A, \quad \frac{dB}{dt} = m_b \left(1 - \frac{B^*}{K_b}\right) B. \tag{2}$$

Here K_a and K_b are the maximal population size attainable (*carrying capacity*) for mutants A and B, respectively, resulting from the limited availability of resources. Now, the

Figure 1. Change in relative fitness when populations have exponential and logistic growth. A) Change in relative fitness of mutant B (W$_{ba}$, black) over time when competing mutants have density-independent, exponential growth−assumed by population genetics theory for populations with continuous growth and overlapping generations [1, 2]. Relative fitness was calculated as [7] $W_{ba} = m_b/m_a$, where m corresponds to the population growth rate for each mutant. The fitness of mutant A, reference, is shown as a black, dotted line. The inset shows the change in cell density, in arbitrary units (A. U.), during the competition. **B-C)** The same information is shown for mutants with density-dependent growth, with (B) and without (C) trade-off between reproduction rate and survival. The sampling time t^* = 24 hours, commonly used in microbiological assays, is noted by a red marker in the main plot, and a vertical, grey line in the insets. Model parameters for A) are m_a = 0.15 h^{-1}, m_b = 0.2 h^{-1}; for B) m_a = 1.0275 h^{-1}, m_b = 1.05 h^{-1}, $K_a = K_b$ = 5 OD, $\alpha_{ab} = \alpha_{ba}$ = 0.15; and for C) m_a = 1.0275 h^{-1}, m_b = 1.05 h^{-1}, K_a = 10 OD, K_b = 5 OD, $\alpha_{ab} = \alpha_{ba}$ = 0.15.

population growth rate m is corrected by the term $1 - (N_i/K_i)$. The Lotka-Volterra model of competition [26] includes the term $N_i^* = N_i - \alpha_{ij}N_j$, where $\alpha_{ij}N_j$ is the linear reduction in growth rate—in terms of K—of species i by competing species j. This formalisation describes the limitation in growth imposed by the environment, due to finite resources, reducing m and the growth of both genotypes over time (Figure 1B and C). In this scenario, m_b depends on the carrying capacity of mutant B *and* mutant A, with the relative fitness changing to

$$W_{ba} = \frac{m_b\left[1 - \left(B^*/K_b\right)\right]}{m_a\left[1 - \left(A^*/K_a\right)\right]} = \frac{m_b^*}{m_a^*}. \tag{3}$$

In other words, W_{ba} will change through time until both mutants reach their carrying capacities, and W_{ba} becomes the ratio of carrying capacities so $W_{ab} = K_b/K_a = k$ is constant. Only after this point fitness can be measured with no variation though time. Curiously, in this scenario m is still calculated as $m_b = \frac{ln[B(t)/B(0)]}{t}$ [7, 12–15]. Estimating fitness differences between two genotypes using equation (3), however, requires prior knowledge of the carrying capacity which can be problematic [7].

The result when m_b is calculated in biological populations, where their growth is limited by resource availability, is illustrated in Figures 1B and C: fitness declines exponentially over time, and whether a mutant is more or less fit than the reference genotype will be given by the choice of t, an arbitrary period during which genotypes are allowed to grow, and therefore $B(t)$. Thus, depending on the combination of carrying capacities, rates of population increase, and sampling time, mutant A may be fitter, as fit, or less fit than the reference genotype B.

If, say, $m_b > m_a$ and $W_{ba} > 1$, which mutant, A or B, would be selected in pairwise competition assays? Despite the simplicity of equations (1) and (2), the answer is not straightforward. Figure 2 illustrates the change in relative abundance of two genotypes, A, and B, growing in competition for a common resource. These genotypes are haploid, with continuous growth and overlapping generations where m_a and m_b represent their population growth rates as introduced above. The competition is propagated at regular intervals c where the new initial conditions are given by $N_a(c_{+1}, 0) = N_a(c, t^*)d$, $N_b(c_{+1}, 0) = N_b(c, t^*)d$, with t^* being the time of growth allowed prior to the propagation and $0 \leq d \leq 1$ the dilution factor during the propagation step analog to experimental assays [7, 8, 14]. Following population genetics theory [1, 3, 4, 7], the outcome of this competition is straightforward: if, say, mutant B reproduces faster than mutant A ($m_b > m_a$), then $W_{ab} = m_a/m_b < 1$ and mutant A goes extinct (Figure 2A). Regardless of how long the competition is allowed to progress until the next propagation step, mutant A is always lost in competition with B. Only through the emergence of a new mutant A^* with $m_a^* > m_b > m_a$ should this prediction change.

But when fitness changes through time the outcome of the competition becomes unclear, now depending on whether the competing genotypes reach their carrying capacities before the propagation: if the competition is propagated frequently, so that *both* genotypes are in active growth, the genotype reproducing faster will replace that reproducing slower consistently with Fisher's theorem. However, if one or both genotypes are allowed to reach their carrying capacities, both will be be co-maintained regardless of the in rates of population increase (Figures 2B and C). Here the relative frequency will be given by their carrying capacities K, so if reproduction and carrying capacity engage in a trade-off [24] for one genotype, this will be most frequent (Figure 2B). Importantly, these predictions are maintained regardless of the initial frequency of A (Figure S1), consistently with prior data [27].

Figure 2. Change in relative frequency during competition depending on the length of the growth cycle. Changes in frequency over time for mutants A (cyan) and B (black) following propagations at arbitrary times t=6h, t=12h, t=18h, and t=24h. These times correspond to different stages of growth as Figure 1 shows. Each plot illustrate the case for genotypes with density-independent growth (A), or density-dependent growth where a genotype does engage in a trade-off between reproduction and survival (B) and when it does not (C). Parameter used in the model are those from Figure 1.

To test this prediction, I competed two constructs of *Escherichia coli* MC4100, Wyl
and GB(c) (see Methods), and measured their variation in fitness through time. Figure
3A shows that both constructs reach their carrying capacicies within 24 hours. The con-
struct GB(c) carries the non-transmissible plasmid pGW155B [28] harbouring *tet(36)*—a
ribosome-protection type resistance gene against tetracycline [28]. This plasmid lacks a
rep gene to control tightly the partition of plasmids after cell division [29–31] (addgene
vector database accession #2853). While pGW155B is a synthetic plasmid, many natural
plasmids also lack partition systems [31, 32]. Both constructs have identical chromo-
some with exception of this plasmid, and the fluorescence gene they carry: cyan (*cfp*,
GB(c)) or yellow (*yfp*, Wyl), to allow their identification in competition assays. Using a
non-transmissible plasmid prevents cross-contamination by horizontal gene transfer be-
tween the resistant construct GB(c) and Wyl. Importantly, the use of constructs derived
from MC4100 avoids interactions between competitors that may affect the outcome of
the competition for reasons beyond pGW155B, like the production of bacteriocins [33].

pGW155B penalised the population growth rate of construct GB(c) by approximately
40% compared to Wyl (Figure 3B) as I found out using pure cultures. It might be argued
that the 4-parameter logistic model I used does not fully capture the growth data. How-
ever, if I measure the maximum change between the reads (Euler's method, see Methods),
the strong difference in growth rate is maintained. The duration of lag, and carrying
capacity were also sensitive to plasmid carriage, regardless of whether I measured the
growth using fluorescence or light scattering (Figure S2). Changes in cell size could con-
found optical density and fluorescence readings, but this phenomenon leaves a signature
[34] in growth data that is absent in my dataset (Figures S2, S3, and S4). The change
in carrying capacity K can be linked to an increase in biomass yield (y) as Monod's ex-
pression [35] $y = K/S$ suggests, where S the supply of glucose. This metric is consistent
the data, given that construct GB(c) has to express and translate plasmid-borne genes
with the same supply of glucose (Figures 3B and C). This means that pGW155B triggers a
metabolic trade-off between growth rate and biomass yield redolent of rK-selection the-
ory [24, 36].

For the competition assay, I mixed equal proportions (cell/cell) of two pure cultures,
from construct GB(c)—harbouring pGW155B—and Wyl respectively, grown overnight, in
media containing no antibiotic or 0.04 μg/mL of tetracycline (see Methods). I incubated
the mixed culture at 30^{o}C until both constructs reached their carrying capacity (K, ~ 24h
as per data in Figure 3A), and then I propagated the competition into a new plate contain-
ing fresh media. I repeated the propagation step seven times totalling between 77 (GB(c))
and 119 (Wyl) generations given the reproduction rates in Figure 3B, where the genera-
tion time per day is given by $m_i/24$, multiplied times the number of propagation steps.
Considering the mutation rate of enterobacteria [37] and the emergence of mutants af-
ter 3 growth cycles [38], I constrained the fitness measurements in both constructs over
the first 24h of competition and compared the outcome of the competition with predic-

Figure 3. Asymmetric carriage costs of pGW155B. A) Growth curves of constructs Wyl (black) and GB(c) (cyan) in pure culture, in the absence of tetracycline, over 24h. I estimated the *per capita* growth rate (*m*), population size in the equilibrium (K), biomass yield, and lag duration from a 4-parameter logistic model fitted to growth data (see Methods). Data fits for constructs Wyl and GB(c) are shown in grey and light cyan, respectively. **B)** Box plots showing the median (centre of the box), 25th and 75th percentile for each of the aforementioned parameters. The whiskers extend to the most extreme data points that are not outliers, and these are individually represented. The *p* value shown refers to a Two-sample *t*-test with unequal variance (8 replicates) that I used to test differences in the parameters between both constructs. **C)** Alternative metrics for growth rate and biomass yield: forward Euler approximation (top), data fit predicted yield (middle), and maximal yield across all time points in growth data (bottom). The *p* values correspond to Two-sample *t*-tests with unequal variance (8 replicates).

tions in Figure 2. Indeed, the relative fitness of both constructs changed through time consistently with Figure 1B and C. In the presence of tetracycline, the relative fitness of drug-sensitive, plasmid-free construct Wyl was below GB(c), which harbours pGW155B, at all times as Figure 4A illustrates. This meant that GB(c) increased its relative frequency in following propagation steps and became the most abundant construct throughout the competition (Figure 4B) consistently with the theoretical predictions. Now, the model does not implement any mutation processes but these occur during this assay. For example, the number of copies of pGW155B harboured by GB(c) increased 5- to 6-fold during the competition (2-sample *t*-test with unequal variances, $p=0.0088$, $df=2.5826$, *t*-statistic=-7.2441; with 3 replicates) driven by exposure to tetracycline (Figure 4C) as it would be reasonable to expect given the exposure to tetracycline. Despite the occurrence of these mutations, Wyl never goes extinct consistently with Figure 2 and its proportion is determine by its abundance at the propagation time.

Similarly, when both constructs grew in mixed cultures, now without tetracycline exposure, the relative fitness of Wyl was higher, equal, or lower than plasmid-harbouring construct GB(c) through time, as Figure 4D shows. Consistently with the predictions both constructs where co-maintained, and with similar relative frequencies, despite the difference in growth rate (Figure 4E). Note here that while the initial deviation is small,

Figure 4. Time-dependent fitness data allows co-maintenance of both constructs without tetracycline. Relative fitness during first 24 hour pairwise-competition of construct Wyl, where both constructs grew in media supplemented with 0.04 μg/mL of tetracycline (**A**) and without drug (**D**). Fitness for each replicate is presented with respect to the reference GB(c) (black dashed line). The inset illustrate the growth curves for both constructs, highlighting in grey the period where the relative fitness of Wyl (W_{sr}) is lower than its tetracycline-resistant competitor's. **B)** and **E)** show the change in relative frequency of both constructs through time when both constructs grew in media supplemented with tetracycline (B) and without antibiotic (E). Error bars represent the mean ± 95% confidence intervals, with raw data points shown as dots. **C)** and **F)** show the change in relative copy numbers of pGW155B borne by construct GB(c) during the pairwise-competition, calculated using quantitative PCR (see Methods). Error bars represent the mean ± standard deviation, with raw data shown in red.

without strong selection by antibiotic exposure each replicate evolves differently, with some replicates of Wyl being lost, while others are co-maintained. In other words, GB(c) preserved pGW155B despite the lack of tetracycline and the difference in growth rates (2-sample t-test with unequal variances, p=0.2305, df=2.2351, t-statistic=-1.6353; with 3 replicates. Figure 4F). This contrasts with previous studies [14] predicting a rapid decline, within a similar time frame, in cells harbouring non-transmissible plasmids.

III. DISCUSSION

Fitness is known to be sensitive to t^* [39] and that it changes through time [27, 40], but it is often presented as an experimental quirk and the root cause is rarely explored. While this variation is used to suggest the co-existence of multiple populations through frequency-dependent fitness [27], here I argue it is something more fundamental: Fitness that is deemed frequency-dependent also changes depending on t^* [27]. Other studies argue that each growth phase contributes to the fitness of a genotype [41], however, even in this case the population rate of increase remains a fundamental parameter.

Despite this, experimentalists extensively use logarithms on experimental data to cal-

culate m_i [7, 12–15] and determine whether mutations are beneficial or deleterious. In other words, experimentalists explicitly rely on Fisher's theorem given that the aforementioned expression $m_b = \frac{ln[B(t)/B(0)]}{t}$ simply results from re-arranging equation 1 and applying logarithms (see Methods). While this may seem reasonable given that populations have a stage where their growth is exponential, such stage is only temporary due to resource depletion. That is, their growth does not fit the assumptions of the exponential growth model. The misconception in Fisher's theorem highlighted in this study is mathematically simple, but the result contests a basic tenet in experimental evolutionary biology: That mutants reproducing faster than their neighbours will be preserved.

Microorganisms must also handle long periods of resource scarcity and therefore multiple strategies may co-exist depending on their natural environment. Indeed, data show that prokaryotes with slow growth are just as abundant as those growing substantially faster [19, 42]. This is what predictions in Figures 2B and C suggest. If the frequency of selection is low, caused for example by long periods of extreme starvation [43, 44], slow- and fast-growing genotypes can be co-maintained. This is particularly relevant to understand the abundance of plasmids in nature, difficult to explain using Fisher's theorem [22]. The co-existence between both Wyl and GB(c) can be surprising given the rationale that plasmids hinder growth rate—fitness—and therefore they cannot be maintained without positive selection. However, selection here was yield. Note that while some replicates are co-maintained at small frequencies without a stronger pressure from antibiotic exposure, most are indeed co-maintained with frequencies ranging between 20–75% as Figure 4E shows. This deviates substantially with the current view where *all* replicates should be lost, or maintained at negligible frequencies [14].

It is noteworthy to mention the co-existence is independent of the advantage provided by pGW155B and indeed any rK trade-offs. Figure 1 shows that such advantage only determines, in the absence of mutations, the relative frequency of each mutant— rather than whether it is co-maintained or lost. While the lack of mutation events is an unrealistic assumption, it suffices to demonstrate that relative fitness changes with time and, therefore, the moment when selection occurs matters to determine whether fast- or slow-growing mutants will be favoured. In this new light, plasmids are not necessarily 'molecular parasites' [45], and the costs of plasmid carriage become relative. Conditions of high frequency, say, caused by repeated exposure to antibiotics during therapies, could indeed favour genotypes with higher growth rates. But the rationale needs not apply when selection is less frequent, where both growth strategies (and plasmid carriage) can be co-maintained—explaining the abundance of plasmids in nature despite the growth penalty they impose on their hosts.

IV. METHODS

Media and Strains. I used the strains of *Escherichia coli* GB(c) and Wyl [46] (a gift from Remy Chait and Roy Kishony), and M9 minimal media supplemented with 0.4% glucose and 0.1% casamino acids (w/v). I made tetracycline stock solutions from powder stock (Duchefa, Ref. #0150.0025) at 5mg/mL in 50% ethanol, filter sterilised, and stored at -20^oC. Subsequent dilutions were made from this stock in supplemented M9 minimal media and kept at 4^oC.

Batch transfer protocol. I inoculated a 96-well microtitre plate containing 150 μg/mL of supplemented M9 media with a mixture of two overnight cultures, one of *E. coli* GB(c) and another of *E. coli* Wyl (1μL containing approx. $2 \cdot 10^8$ cells, Figure S3). The overnight culture for GB(c) was supplemented with 100ng/mL of tetracycline to preserve the plasmid pGW155B carrying *tet(36)* as described elsewhere [8], then centrifuged and removed prior adding to the microtitre plate. I incubated the plate at 30^oC in a commercial spectrophotometer and measured the optical density of each well at 600nm (OD_{600}), yellow florescence for the Wyl strain (YFP excitation at 505nm, emission at 540nm), and cyan fluorescence for the GB(c) strain (CFP at 430nm/480nm) every 20min for 24h. After each day I transferred 1.5μL of each well, using a 96-well pin replicator, into a new microtitre plate containing fresh growth medium and tetracycline.

Growth parameter estimation. Yellow and cyan fluorescence protein genes were constitutively expressed given the constant ratio between fluorescence and optical density (Figure S5). This allowed me to use fluorescence data as a proxy for cell density in mixed culture conditions. I normalised fluorescence readings using a conversion factor, n_f, calculated by growing each construct in a range of glucose concentrations, and regressing the linear model RFU = $n_f \cdot$ OD + c, where RFU is relative fluorescence units data, OD optical density data, n_f the conversion factor between fluorescence and optical density, and c the crossing point with the y–axis when OD = 0. I imported the resulting time series data set (Figures S4) into MATLAB R2014b to subtract background and calculate fitness as described in the main text as follows.

Once corrected, I used the expression $m_i = \frac{ln[N_i(t)/N_i(0)]}{t}$ to estimate the *per capita* growth rate m_i as per in [7, 12–15] although I used different values for t to test the theoretical predictions. This expression derives from the exponential model in equation 1 as it follows. We can isolate the term with m for, say, mutant A, as $A(t) = A_0 e^{m_a t} \rightarrow \frac{A(t)}{A_0} = e^{m_a t}$. To remove the number e we then apply logarithms to this expression leading to $ln\frac{A(t)}{A_0} = m_a t$. Re-arranging the terms we are left with $\frac{ln[A(t)/A_0]}{t} = m_a$, which is the expression used by experimentalists.

307 To measure the carriage costs of pGW155B, I fitted to data the logistic model

$$g(t) = \frac{b + K}{(1 + C \cdot e^{-m \cdot [t-l]})}.$$

308 where the term $g(t)$ denotes the change in culture growth as a function of time (in OD,
309 YFP, or CFP units), b the inoculum size used to subtract the background, C is a parameter,
310 m the *per capita* growth rate, K the maximal population size attained and l the duration
311 of the lag phase. Alternatively, I measured the rate of increase using the forward Euler
312 approximation $m_i = max_{0 \leq t \leq 24h}\left(\frac{N_i(t)/N_i(t-1)}{\Delta t}\right)$, where $\Delta t = 20\text{min}= 1/3$ is the time be-
313 tween reads. To estimate the biomass yield I divided OD data in stationary phase by the
314 glucose supplied [47]. Alternatively, I used the highest density across all time points and
315 the density reported by the data fit, both divided by glucose supply, as alternative metrics
316 for biomass yield.

317 Finally, I divided the length of the growth cycle (24h) by the *per capita* growth rate,
318 multiplied by the number of propagations (seven) to estimate the total number of dupli-
319 cation events—where one duplication means one generation—for each construct.

320 **DNA material extraction.** For each concentration, I sampled three representative 150
321 µg/mL replicates that I divided in two groups: for chromosome and plasmid DNA ex-
322 traction. I used 'GeneJet DNA' (ThermoScientific, Ref. #K0729) and 'GeneJet Plasmid'
323 (ThermoScientific, Ref. #K0502) extraction kits to extract chromosomal and plasmid
324 DNA (pDNA) from the samples, respectively, and used Qubit to quantify DNA and pDNA
325 yields. Both extracts were diluted accordingly in extraction buffer, as per in manufacturer
326 instructions, to normalise DNA across samples.

327 **Quantitative PCR and plasmid copy number estimation.** I used `primer3` to design
328 two pairs of primers with melting temperature (T_m) of 60^oC and non-overlapping probes
329 with T_m of 70^oC. The amplicon ranges between 100 to 141bp depending on the locus (Ta-
330 ble 1). Two reaction mixes were prepared using the kit 'Luminaris Color Probe Low ROX'
331 (ThermoScientific, Ref. #K0342), adding 0.3µM of each primer and 0.2µM of the probe
332 as per manufacturer specifications. Following a calibration curve for each reaction (Fig-
333 ure S6) I added 0.01ng of chromosomal or plasmid DNA material to each of the reaction
334 mixes.

339 To estimate the relative copies of pGW155B per GB(c) cell, I calculated the corre-
340 sponding proportion of chromosomal DNA corresponding to the GB(c)-type from data
341 in Figure 4 and used the expression [14]

342
$$cn = \frac{(1 + E_c)^{Ctc}}{(1 + E_p)^{Ctp}} \times \frac{S_c}{S_p},$$

343 where cn is the number of plasmid copies per chromosome, S_c and S_p are the size of the
344 chromosome and pGW155B amplicon in bp, E_c and E_p the efficiency of the qPCR taken

Table 1. Primers and probes designed using `Primer3`. Amplicon ranging from 100 to 141bp. T$_m$ indicates the estimated melting temperature.

Target gen	Sequence $(5' \rightarrow 3')$	$\mathbf{T}_m$ (^oC)	Feature
tatB	CGATGAAGCGTTCCTACGTT	60.27	Forward
	TCATGCGCAGCTTCATTATC	59.94	Reverse
	AAGGCGAGCGATGAAGCGCA	70.70	Probe
tet(36)	ATTGGGCATCTATTGGCTTG	59.22	Forward
	CCGATTCACAGGCTTTCTTG	60.76	Reverse
	AGCCTTTGCCAATTGGGGCG	70.37	Probe

from data in Figure S6, and *Ctc* and *Ctp* are the cycles at which I first detected product amplification (C_t).

REFERENCES

1. Crow, J. F., Kimura, M., *et al. An introduction to population genetics theory.* (New York, Evanston and London: Harper & Row, Publishers, 1970).

2. Nagylaki, T. in *Introduction to Theoretical Population Genetics* 5–27 (Springer, 1992).

3. Templeton, A. *Population Genetics and Microevolutionary Theory* ISBN: 9780471409519 (Wiley, 2006).

4. Dawes, I. & Sutherland, I. *Microbial Physiology* ISBN: 9780632024636 (Wiley, 1992).

5. Borriello, S., Murray, P. & Funke, G. *Topley and Wilson's Microbiology and Microbial Infections, 2 Volume Set: Bacteriology* ISBN: 9780470660263 (Wiley, 2007).

6. Fisher, R. A. *The Genetical Theory of Natural Selection* 308–315 (At The Clarendon Press, 1930).

7. Lenski, R. E., Rose, M. R., Simpson, S. C. & Tadler, S. C. Long-Term Experimental Evolution in Escherichia coli. I. Adaptation and Divergence During 2,000 Generations. *Am. Nat.* **138,** pp. 1315–1341 (1991).

8. Chait, R., Craney, A & Kishony, R. Antibiotic interactions that select against resistance. *Nature* **446,** 668–671 (2007).

9. Wang, H., Avican, K., Fahlgren, A., *et al.* Increased plasmid copy number is essential for Yersinia T3SS function and virulence. *Science* **353,** 492–495 (2016).

10. Adkar, B. V., Manhart, M., Bhattacharyya, S., Tian, J., Musharbash, M. & Shakhnovich, E. I. Optimization of lag phase shapes the evolution of a bacterial enzyme. *Nat. Ecol. Evol.* **1,** 1–6 (2017).

11. Concepción-Acevedo, J., Weiss, H. N., Chaudhry, W. N. & Levin, B. R. Malthusian parameters as estimators of the fitness of microbes: a cautionary tale about the low side of high throughput. *PLoS One* **10,** e0126915 (2015).

12. Andersson, D. I. & Levin, B. R. The biological cost of antibiotic resistance. *Curr. Opin. Microbiol.* **2,** 489–493 (1999).

13. Gullberg, E., Cao, S., Berg, O. G., *et al.* Selection of resistant bacteria at very low antibiotic concentrations. *PLoS Pathog.* **7,** e1002158 (2011).

14. Millán, A. S., Peña-Miller, R., Toll-Riera, M, *et al.* Positive selection and compensatory adaptation interact to stabilize non-transmissible plasmids. *Nat. Commun.* **5** (2014).

15. Ram, Y., Dellus-Gur, E., Bibi, M., *et al.* Predicting microbial growth in a mixed culture from growth curve data. *Proc. Natl. Acad. Sci. U.S.A.* **116,** 14698–14707 (2019).

16. Kussell, E., Kishony, R., Balaban, N. Q. & Leibler, S. Bacterial persistence: a model of survival in changing environments. *Genetics* **169,** 1807–1814 (2005).

17. Pedrós-Alió, C. Marine microbial diversity: can it be determined? *Trends Microbiol.* **14,** 257–263 (2006).

18. Lever, M. A., Rogers, K. L., Lloyd, K. G., *et al.* Life under extreme energy limitation: a synthesis of laboratory-and field-based investigations. *FEMS Microbiol. Revs.* **39,** 688–728 (2015).

19. Flemming, H.-C. & Wuertz, S. Bacteria and archaea on Earth and their abundance in biofilms. *Nat. Rev. Microbiol.* **17,** 247–260 (2019).

20. Smillie, C., Garcillán-Barcia, M. P., Francia, M. V., Rocha, E. P. & de la Cruz, F. Mobility of plasmids. *Microbiol. Mol. Biol. Rev.* **74,** 434–452 (2010).

21. Harrison, E. & Brockhurst, M. A. Plasmid-mediated horizontal gene transfer is a coevolutionary process. *Trends Microbiol.* **20,** 262–267 (2012).

22. Probing the plasmid paradox. *Nat. Ecol. Evol.* **5,** 1559–1559 (2021).

23. Lopatkin, A. J., Meredith, H. R., Srimani, J. K., Pfeiffer, C., Durrett, R. & You, L. Persistence and reversal of plasmid-mediated antibiotic resistance. *Nat. Commun.* **8,** 1689 (2017).

24. Reding-Roman, C., Hewlett, M., Duxbury, S., Gori, F., Gudelj, I. & Beardmore, R. The unconstrained evolution of fast and efficient antibiotic-resistant bacterial genomes. *Nat. Ecol. Evol.* **1,** 0050 (2017).

25. Hansen, S. K., Rainey, P. B., Haagensen, J. A. & Molin, S. Evolution of species interactions in a biofilm community. *Nature* **445,** 533–536 (2007).

26. Gilpin, M. E. & Ayala, F. J. Global models of growth and competition. *Proc. Natl. Acad. Sci. U.S.A.* **70,** 3590–3593 (1973).

27. Ribeck, N. & Lenski, R. E. Modeling and quantifying frequency-dependent fitness in microbial populations with cross-feeding interactions. *Evolution* **69,** 1313–1320 (2015).

28. Whittle, G, Whitehead, T. R., Hamburger, N, Shoemaker, N. B., Cotta, M. A., *et al.* Identification of a new ribosomal protection type of tetracycline resistance gene, tet(36), from swine manure pits. *Appl. Environ. Microbiol.* **64,** 4151–4158 (2003).

29. Del Solar, G. & Espinosa, M. Plasmid copy number control: an ever-growing story. *Mol. Microbiol* **37,** 492–500 (2000).

30. Garcillán-Barcia, M. P., Alvarado, A. & de la Cruz, F. Identification of bacterial plasmids based on mobility and plasmid population biology. *FEMS Microbiol. Rev.* **35,** 936–956 (2011).

31. Million-Weaver, S. & Camps, M. Mechanisms of plasmid segregation: have multicopy plasmids been overlooked? *Plasmid* **75,** 27–36 (2014).

32. Guynet, C. & de la Cruz, F. Plasmid segregation without partition. *Mob. Genet. Elements* **1,** 236–241 (2011).

33. Dobson, A., Cotter, P. D., Ross, R. P. & Hill, C. Bacteriocin Production: a Probiotic Trait? *Appl. Environ. Microbiol.* **78,** 1–6 (2012).

34. Stevenson, K., McVey, A. F., Clark, I. B., Swain, P. S. & Pilizota, T. General calibration of microbial growth in microplate readers. *Sci. Rep.* **6,** 1–7 (2016).

35. Monod, J. *Recherches sur la croissance des cultures bactériennes* PhD thesis (Paris, 1941).

36. Reznick, D., Bryant, M. J. & Bashey, F. r-and K-selection revisited: the role of population regulation in life-history evolution. *Ecology* **83,** 1509–1520 (2002).

37. Perfeito, L., Fernandes, L., Mota, C. & Gordo, I. Adaptive mutations in bacteria: high rate and small effects. *Science* **317,** 813–815 (2007).

38. Reding, C., Catalán, P., Jansen, G., *et al.* The Antibiotic Dosage of Fastest Resistance Evolution: gene amplifications underpinning the inverted-U. *Mol. Biol. Evol.* **9,** 3847–3863. ISSN: 0737-4038 (Mar. 2021).

39. McDonald, M. J. Microbial experimental evolution–a proving ground for evolutionary theory and a tool for discovery. *EMBO Rep.* **20,** e46992 (2019).

40. Li, Y., Venkataram, S., Agarwala, A., *et al.* Hidden complexity of yeast adaptation under simple evolutionary conditions. *Current Biology* **28,** 515–525 (2018).

41. Lin, J., Manhart, M. & Amir, A. Evolution of microbial growth traits under serial dilution. *Genetics* **215,** 767–777 (2020).

42. Ernebjerg, M. & Kishony, R. Distinct growth strategies of soil bacteria as revealed by large-scale colony tracking. *Appl. Environ. Microbiol.* **78,** 1345–1352 (2012).

43. Gray, D. A., Dugar, G., Gamba, P., Strahl, H., Jonker, M. J. & Hamoen, L. W. Extreme slow growth as alternative strategy to survive deep starvation in bacteria. *Nat. Commun.* **10,** 1–12 (2019).

44. Shoemaker, W. R., Jones, S. E., Muscarella, M. E., Behringer, M. G., Lehmkuhl, B. K. & Lennon, J. T. Microbial population dynamics and evolutionary outcomes under extreme energy limitation. *Proc. Natl. Acad. Sci. U.S.A.* **118,** e2101691118 (2021).

45. Lili, L. N., Britton, N. F. & Feil, E. J. The persistence of parasitic plasmids. *Genetics* **177,** 399–405 (2007).

46. Chait, R., Shrestha, S., Shah, A. K., Michel, J.-B. & Kishony, R. A Differential Drug Screen for Compounds That Select Against Antibiotic Resistance. *PLoS One* **5,** e15179 (2010).

47. Monod, J. The Growth of Bacterial Cultures. *Annu. Rev. Microbiol.* **3,** 371–394 (1949).

Competing interests. The author declares no competing interests.

Acknowledgements. The author thanks Robert Beardmore for laboratory support in the early states of this study.

Data availability statement Data used to generate Figures 3 and 4 is available as a supplementary table. An Octave/MATLAB implementation of equations 1 and 2 is available at `https://github.com/rc-reding/papers/tree/master/FisherTheorem`.

I. SUPPLEMENTARY FIGURES

Figure S1. Relative fitness of mutant A at different densities. A–B) Relative fitness of mutant A measured after seven propagation steps, analog to those described in the main manuscript, when growth rate and population size engage in a trade-off (A) and when they do not (B). The frequency on the x-axis is that of mutant A at t=0. The fitness of each competitor (m_i) was calculated as explained in the main manuscript as $m_i = \frac{ln[N_i(t)/N_i(0)]}{\Delta t}$ where $N_i(t)$ is the density of each mutant at time $t = 7$, $N_i(0)$ that at time t=0, and Δt = 7. This was used to calculate the relative fitness of A as $W_{ab} = m_a/m_b$ as described in the text. The code is available in github in the link provided in the text, and the parameters used are those from Figure 1B and C. **C–D)** Relative frequency of mutant A at the end of the seventh propagation when growth rate and population size engage in a trade-off (C) and when they do not (D), shown in the y-axis, as a function of its frequency at t=0.

Figure S2. Asymmetric carriage costs of pGW155B (optical density data). A) Growth curves of constructs Wyl (black) and GB(c) (cyan) in pure culture, in the absence of tetracycline, over 24h. I estimated the maximum growth rate (r), population size in the equilibrium (K), biomass yield, and lag duration from a 4-parameter logistic model fitted to growth data (see Methods). Data fits for constructs Wyl and GB(c) are shown in grey and light cyan, respectively. **B)** Box plots showing the median (centre of the box), 25th and 75th percentile for each of the aforementioned parameters. The whiskers extend to the most extreme data points that are not outliers, and these are individually represented. The *p* value shown refers to Two-sample *t*-tests with unequal variance (8 replicates) that I used to test differences in the parameters between both constructs. **C)** Alternative metrics for growth rate and biomass yield: forward Euler approximation (top), data fit predicted yield (middle), and maximal yield across all time points (bottom). The *p* values correspond to Two-sample *t*-tests with unequal variance (8 replicates).

Figure S4. Raw data of each construct during the competition. Raw optical density data for GB(c) (cyan) and Wyl (black), measured every 20 min and converted from fluorescence readings. Each individual plot represents the change in abundance of each construct over 24 hours. Data corresponding to each condition, with and without tetracycline, are aggregated as columns.

Figure S5. Changes in relative fluorescence over time in GB(c) and Wyl strains in pure culture conditions. Raw change in florescence, per optical density units, measured every 20min for 24h for the Wyl- (black) and GB(c)-type. Each column represents the data set for each tetracycline concentration used.

A   **B**

C   **D**

Figure S6. Quantitative PCR calibration curves for *tet(36)* and *tatB*. Reaction efficiency for the set of primers and probes, listed in Table 1 the main Methods section, for *tet(36)* (A) and *tatB*. The efficiency was calculated as $E_f = 10^{-1/Slope} - 1$, and the slope term calculated by fitting a linear model to qPCR threshold cycle (C_t) data. The mean ± standard deviation for the adjusted coefficient of determination R^2 and efficiency are shown in the figures. The amplification curves for each reaction, using 3 replicates, are shown in C) and D), respectively.