1
2

Bayesian Evaluation of Temporal Signal in Measurably Evolving Populations

Sebastian Duchene[*,1], Philippe Lemey[2], Tanja Stadler[3], Simon YW Ho[4], David A Duchene[5], Vijaykrishna Dhanasekaran[6], Guy Baele[2]
8
9
[1]Department of Microbiology and Immunology, Peter Doherty Institute for Infection and Immunity, University of Melbourne, Melbourne, VIC, Australia.
[2]Department of Microbiology, Immunology and Transplantation, Rega Institute, KU Leuven, Leuven, Belgium.
[3]Department of Biosystems Science and Engineering, ETH Zürich; and Swiss Institute of Bioinformatics; Basel, Switzerland
[4]School of Life and Environmental Sciences, University of Sydney, Sydney, NSW, Australia
[5]Research School of Biology, Australian National University, Canberra, ACT, Australia.
[6]Department of Microbiology, Biomedicine Discovery Institute, Monash University, Melbourne, VIC, Australia.
20
21
22
*Corresponding author: E-mail: sebastian.duchene@unimelb.edu.au
24

Abstract (250 words max)

Phylogenetic methods can use the sampling times of molecular sequence data to calibrate the molecular clock, enabling the estimation of substitution rates and time scales for rapidly evolving pathogens and data sets containing ancient DNA samples. A key aspect of such calibrations is whether a sufficient amount of molecular evolution has occurred over the sampling time window, that is, whether the data can be treated as being from a measurably evolving population. Here we investigate the performance of a fully Bayesian evaluation of temporal signal (BETS) in molecular sequence data. The method involves comparing the fit of two models: a model in which the data are accompanied by the actual (heterochronous) sampling times, and a model in which the samples are constrained to be contemporaneous (isochronous). We conduct extensive simulations under a range of conditions to demonstrate that BETS accurately classifies data sets according to whether they contain temporal signal or not, even when there is substantial among-lineage rate variation. We explore the behaviour of this classification in analyses of five data sets: modern samples of *A/H1N1 influenza virus*, the bacterium *Bordetella pertussis*, and coronaviruses from mammalian hosts, and ancient DNA data sets of *Hepatitis B virus* and of mitochondrial genomes of dog species. Our results indicate that BETS is an effective alternative to other measures of temporal signal. In particular, this method has the key advantage of allowing a coherent assessment of the entire model, including the molecular clock and tree prior which are essential aspects of Bayesian phylodynamic analyses.
44

Key words: Bayesian phylogenetics, ancient DNA, measurably evolving population, marginal likelihood, molecular clock, temporal signal.
47

Introduction

The molecular clock has become a ubiquitous tool for studying evolutionary processes in rapidly evolving organisms and in data sets that include ancient DNA. In its simplest form, the molecular clock posits that evolutionary change occurs at a predictable rate over time (Zuckerkandl and Pauling 1965). The molecular clock can be calibrated to estimate divergence times by using sampling time information, the timing of known divergence events, or a previous estimate of the substitution rate (Hipsley and Müller 2014). For example, Korber et al. (2000) used sampling times to calibrate the molecular clock and to infer the time of origin of HIV group 1. Their approach consisted of estimating a phylogenetic tree and conducting a regression of the distance from the root to each of the tips as a function of sequence sampling time. In this method, the slope of the regression is an estimate of the substitution rate in substitutions per site per unit of time, the intercept with the time axis is the age of the root node, and the coefficient of determination (R^2) is the degree to which the data exhibit clocklike behaviour (Rambaut et al. 2016). Despite the practicality of root-to-tip regression, its use as a statistical tool for molecular dating has several well-known limitations. In particular, data points are not independent because they have shared ancestry (i.e., internal branches are traversed multiple times) and a strict clocklike behaviour is assumed by necessity.

The past few decades have seen a surge in molecular clock models that explicitly use phylogenetic information. Bayesian methods have gained substantial popularity, largely due to the wide array of complex models that can be implemented and the fact that independent information, including calibrations, can be specified via prior distributions (Nascimento et al. 2017). Of particular importance is the availability of molecular clock models that relax the assumption of strict clock behaviour by explicitly modelling rate variation among lineages (reviewed by Ho and Duchene (2014) and by Bromham et al. (2018)).

Regardless of the methodology used to analyse time-stamped sequence data, a sufficient amount of molecular evolution must have occurred over the sampling time window to allow reliable estimates of substitution rates and timescales. In such cases, the population can be considered to be 'measurably evolving' (Drummond et al. 2003). The degree of 'temporal information' in sequence data is determined by the sequence length, the substitution rate, and the range of available sampling times. Some viruses evolve at a rate of around 5×10^{-3} subs/site/year (Duchene et al. 2014), such that samples collected over a few weeks can be sufficient to calibrate the molecular clock. In more slowly evolving organisms, such as mammals, a sampling window of tens of thousands of years might be necessary; this can be achieved by including ancient DNA sequences (Drummond et al. 2003; Biek et al. 2015).

Testing for temporal signal is an important step for verifying that the molecular clock can be calibrated using the sampling times (Rieux and Balloux 2016). For this purpose a date-randomization test has been proposed that compares actual substitution rate estimates to those obtained by repeatedly permuting the sequence sampling times (Ramsden et al. 2009). A data set is considered to have strong temporal signal if the rate estimated using the correct sampling times does not overlap with those of the permutation replicates (Duchene et al. 2015, 2018; Murray et al. 2015). An implementation of this test is also available that performs the permutation during a single Bayesian inference (Trovão et al. 2015). The interpretation of the date-randomization test is essentially frequentist in nature, which leads to an inconsistent mixture of statistical frameworks when Bayesian phylogenetic methods are used. Moreover, the procedure is not applicable in cases with small numbers of sampling times, owing to the limited number of possible permutations (Duchene et al. 2015).

We propose a full Bayesian model test, which we refer to as BETS (Bayesian Evaluation of Temporal
Signal), to assess temporal signal based on previous analyses by Baele et al. (2012). The approach
involves quantifying statistical support for two competing models: a model in which the data are
accompanied by the actual sampling times (i.e., the data are treated as heterochronous) and a
model in which the sampling times are contemporaneous (i.e., the data are treated as isochronous).
Therefore, the sampling times are treated as part of the model and the test can be understood as a
test of ultrametricity of the phylogenetic tree. If incorporating sampling times improves the
statistical fit, then their use for clock calibration is warranted. The crux of BETS, as with Bayesian
model selection, is that it requires calculating the marginal likelihood of the model in question. The
marginal likelihood measures the evidence for a model given the data, and calculating it requires
integration of its likelihood across all parameter values, weighted by the prior (Kass and Raftery
1995).

Because the marginal likelihood is a measure of model evidence, the ratio of the marginal
likelihoods of two competing models, known as the Bayes factor, is used to assess support for one
model relative to the other. In the case of applying BETS, let M_{het} represent the heterochronous
model, M_{iso} the isochronous model, and Y the sequence data, such that $P(Y|M_{het})$ and $P(Y|M_{iso})$ are
their respective marginal likelihoods. These models differ in the number of parameters; in M_{iso} the
substitution rates and times are nonidentifiable, so the rate is fixed to an arbitrary value, while in
M_{het} it is a free parameter. Differences in the number of parameters do not need to be taken into
account separately, because the marginal likelihood naturally penalizes excessive parameterization.
Kass and Raftery (1995) gave guidelines to interpreting Bayes factors, where a (log) Bayes factor
$\log(P(Y|M_{het})) - \log(P(Y|M_{iso}))$ of at least 5 indicates 'very strong' support for M_{het} over M_{iso}, a value of
3 indicates 'strong' support, and a value of 1 is considered as positive evidence for M_{het} over M_{iso}.

The importance of model selection in Bayesian phylogenetics has prompted the development of
various techniques to calculate marginal likelihoods (reviewed by Baele et al. (2014) and by Oaks et
al. (2019)). These techniques can be broadly classified into prior-based and/or posterior-based
estimators and path-sampling approaches. Prior- and posterior-based estimators, also known as
importance sampling, include the widely used harmonic-mean estimator (Newton and Raftery
1994) and the AICM and BICM (Bayesian analogues to the Akaike information criterion and the
Bayesian information criterion, respectively) (Raftery et al. 2007). These scores are easy to compute
because they only require samples from the posterior distribution as obtained through Markov
chain Monte Carlo (MCMC) integration. However, the harmonic-mean estimator has been shown to
have unacceptably high variance when the prior is diffuse relative to the posterior, and, together
with the AICM, has shown poor performance in practical settings (Baele et al. 2012, 2013). The BICM
requires a sample size to be specified for each parameter, which is far from trivial for phylogenetic
inference and therefore remains unexplored for such applications.

Path-sampling approaches include path sampling (originally introduced in phylogenetics as
'thermodynamic integration') (Lartillot and Philippe 2006), stepping-stone sampling (Xie et al.
2011), and generalized stepping-stone (GSS) sampling (Fan et al. 2011; Baele et al. 2016). These
methods depend on drawing samples using MCMC from a range of power posterior distributions
that represent the path from the posterior to the (working) prior, and therefore require additional
computation. Another numerical technique that was recently introduced to phylogenetics is nested
sampling (NS) (Maturana et al. 2019), which approximates the marginal likelihood by simplifying
the marginal-likelihood function from a multi-dimensional to a one-dimensional integral over the
cumulative distribution function of the marginal likelihood (Skilling 2006). Fourment et al. (2019)
recently compared the accuracy of a range of marginal-likelihood estimation methods and found
GSS to be the most accurate, albeit at increased computational cost. Clearly, the reliability of the
marginal-likelihood estimator is a key consideration for applying BETS.

149
150   We conducted a simulation study to assess the reliability of BETS under a range of conditions that
151   are typical for data sets of rapidly evolving organisms and of those involving ancient DNA. We also
152   analysed five empirical data sets to showcase the performance of the test in practice. Our analyses
153   demonstrate the utility of BETS to provide accurate evaluation of temporal signal across a wide
154   range of conditions.
155
156   **Results**
157
Simulations of Measurably Evolving Populations

In our simulations we considered sequence data from heterochronous and isochronous trees.
Heterochronous trees represent a situation where there is sufficient temporal signal, whereas
isochronous trees lack temporal signal altogether. We simulated heterochronous phylogenetic
trees under a stochastic birth-death process with between 90 and 110 tips. To generate isochronous
trees we used similar settings, but we assumed a single sampling time. We then simulated
substitution rates along the trees according to an uncorrelated relaxed clock with an underlying
lognormal distribution with a mean of 5×10^{-3} subs/site/unit time and a standard deviation, σ, of 0,
0.1, 0.5, or 1, where $\sigma=0$ is equivalent to simulating under a strict clock. We then simulated
sequence evolution using an HKY+Γ substitution model, with parameter values similar to those
estimated for influenza virus (Hedge et al. 2013), to generate alignments of 10,000 nucleotides.
169
Our main simulation conditions produced data sets in which about 50% of the sites were variable.
We refer to this simulation scenario as (i) 'high substitution rate and wide sampling window', and we
considered three other simulation scenarios that involved (ii) a lower substitution rate of 10^{-5}
subs/site/unit time, (iii) a narrower sampling window, and (iv) both of the last two conditions. We
analysed the sequence data using a strict clock and an uncorrelated relaxed clock with an
underlying lognormal distribution (Drummond et al. 2006). We considered three configurations for
sampling times: birth-death sampling times, which are correct for the heterochronous data but not
for the isochronous data; identical sampling times, which is correct for isochronous data but not for
the heterochronous data; and permuted birth-death sampling times, which are incorrect for both
heterochronous and isochronous data.
180
181   We estimated the log marginal likelihoods of these six combinations of sampling times and clock
182   models using NS and GSS as implemented in BEAST 2.5 (Bouckaert et al. 2019) and BEAST 1.10
183   (Suchard et al. 2018), respectively. Our BETS approach ranked the models according to their log
184   marginal likelihoods and computed log Bayes factors of the best heterochronous model (M_{het})
185   compared with the best isochronous model (M_{iso}).
186
(i) Simulations with High Substitution Rate and Wide Sampling Window

Both NS and GSS correctly classified data as being heterochronous or isochronous in 10 out of 10
simulations, including in the presence of a high degree of among-lineage rate variation (i.e., $\sigma=1$;
fig. 1 for heterochronous data and supplementary fig. S1, Supplementary Material online, for
isochronous data). Although both marginal-likelihood estimators detected temporal signal, NS
supported the relaxed clock over the strict clock for three heterochronous data sets simulated
without among-lineage rate variation ($\sigma=0$) and for six data sets simulated with low among-lineage
rate variation ($\sigma=0.1$). In the simulations of isochronous data, NS often favoured the relaxed clock
over the strict clock when there was low among-lineage rate variation ($\sigma=0.0$ and $\sigma=0.1$), albeit
mostly with log Bayes factors below 5 (supplementary fig. S1, Supplementary Material online). In
contrast, GSS always selected the strict clock under these conditions (fig. 1 and supplementary fig.
S1, Supplementary Material online).

For the heterochronous data sets, NS and GSS always displayed very strong support for M_{het} over
M_{iso}, with log Bayes factors of at least 90. For the isochronous data sets, the log Bayes factors for
M_{iso} relative to M_{het} were overall much lower, but still decisive, ranging from 30 to 50. Furthermore,
log Bayes factors tended to decline with an increasing degree of among-lineage rate variation in the
data. Another important observation is that in the heterochronous data, the relaxed clock was
consistently selected over the strict clock when assuming that the data were isochronous, or when
the sampling times had been permuted (fig. 1 and supplementary fig. S1, Supplementary Material
online). Moreover, the strict clock with permuted sampling times yielded the lowest log marginal
likelihoods for heterochronous data. Both of these patterns are likely to be due to an apparently
higher degree of among-lineage rate variation when sampling times are misspecified.

(ii) Simulations with Low Substitution Rate and Wide Sampling Window

Our simulations with a low substitution rate of 10^{-5} subs/site/unit time produced data sets that each
had about 10 variable sites, which provides very little information for the estimation of evolutionary
parameters. Additionally, due to the stochasticity of the simulation process, increased estimator
variance between replicates is to be expected given the small number of variable sites. For the
heterochronous data sets, GSS selected the heterochronous model with correct dates in at least 7
out of 10 simulation replicates (fig. 2). Across the simulations with different clock models (40 in
total), only in five heterochronous data sets did we find models with permuted sampling times to
have the highest log marginal likelihoods. For NS, in 11 out of 40 simulations, either isochronous
models or those with random sampling times were incorrectly selected when heterochronous data
sets were analysed.

Log marginal likelihoods calculated using GSS tended to support models with sampling times
(either permuted or those from the birth-death) for the isochronous data, whereas NS appeared to
support all models with similar frequencies (supplementary fig. S2, Supplementary Material online).
However, a critical feature of the results from the data sets with a low substitution rate is that the
log marginal likelihoods for all models were more similar to one another than those for the data sets
with high substitution rate (note that the log marginal likelihood scale in fig. 2 is smaller than that in
fig. 1). As a case in point, for the isochronous data with σ =0.1 there were log Bayes factors of about
0.1 for the best model with birth-death sampling times relative to those with permuted sampling
times. This result indicates that comparing models with permuted sampling times might be useful
for determining whether the data are informative about a particular set of sampling times.

(iii) Simulations with High Substitution Rate and Narrow Sampling Window

We conducted a set of simulations similar to those described in scenario (i) but where sequence
sampling spanned only the last 10% of the age of the tree (0.5 units of time, compared with 5 units
of time for the simulations with a wide sampling window). These conditions reflect those of
organisms with deep evolutionary histories and for which samples are available for only a small
portion of this time. Since in these trees the samples were collected over a narrower time window,
we used a higher sampling probability to obtain about 100 samples, as in our other simulations (see
examples of trees in supplementary fig. S3, Supplementary Material online). For these analyses we
only considered heterochronous data because the isochronous case is the same as that in scenario
(i).

Both GSS and NS showed excellent performance in detecting temporal signal in this scenario,
almost always selecting models with correct sampling times. The exceptions to this pattern
occurred for one data set with σ =0.5 and for two data sets with σ =1.0 for NS (fig. 3).
Differentiating between the strict clock and relaxed clock appeared somewhat more difficult,

particularly for NS, where the relaxed clock with correct sampling times yielded log marginal
likelihoods very similar to those for the strict clock for data with low among-lineage rate variation
(σ of 0.0 or 0.1). Although NS and GSS performed well in these simulations, the log Bayes factors
for M_{het} relative to M_{iso} were much lower than those for data with a high substitution rate and a wide
sampling window in (i). One obvious example is in the data with σ=0.0, where the mean log Bayes
factors for M_{het} over M_{iso} using GSS was 203.15 with a wide sampling window (fig. 1), but only 35.77
when sampling spanned a narrow time window (fig. 3).
256
257 *(iv) Simulations with Low Substitution Rate and Narrow Sampling Window*

258 We considered data sets with a narrow sampling window, as in scenario (iii), and with a low
259 substitution rate of 10^{-5} subs/site/unit time, as in scenario (ii). We generated only heterochronous
260 trees under these conditions, because the isochronous case would be the same as that in (ii).
261
Estimates of log marginal likelihoods with GSS and NS were very similar among models, with mean
log Bayes factors among data sets of less than 1 for the two models with highest marginal
likelihoods for GSS (fig. 4). In the data sets with σ=0.0, GSS and NS always preferred a
heterochronous model. However, in a few cases (three for GSS and one for NS) the model with
permuted sampling times was selected, indicating that temporal signal was not detected. As with
the data sets with low substitution rate and constant sampling (ii), the relaxed clock was sometimes
preferred over the strict clock, even when the data sets had no rate variation among lineages.
269
Comparison with Root-to-tip Regression

Using a subset of the heterochronous data sets, we conducted root-to-tip regression using
phylogenetic trees inferred using maximum likelihood in PhyML 3.1 (Guindon et al. 2010) with the
same substitution model as in our BEAST analyses, and with the placement of the root chosen to
maximize R^2 in TempEst (Rambaut et al. 2016). We selected data sets generated with a high
substitution rate and with both constant and narrow sampling windows. Because GSS and NS
correctly detected temporal signal under these conditions, these regressions demonstrate the
extent to which this informal regression assessment matches the BETS approach. We did not
attempt to provide a thorough benchmarking of the two methods here.
279
All regressions had R^2 values that matched our expectation from the degree of among-lineage rate
variation, that is, higher values of σ corresponded to lower values of R^2 (fig. 5). The data with a
wide sampling window yielded regression slopes ranging from 7.3×10^{-3} to 5.4×10^{-3} subs/site/unit
time, which is similar to the substitution rate values used to generate the data. Although the root-
to-tip regression is sometimes used to assess temporal signal, it has no cut-off values to confirm
temporal signal. This becomes critical when considering the data with a narrow sampling window,
for which the R^2 was between 0.13 and 0.02. For example, the regression for a data set with σ=1
and narrow sampling window had an R^2 of 0.02, which is sometimes considered sufficiently low as
to preclude molecular clock analyses (Rieux and Balloux 2016). However, BETS supported strong
temporal structure under a relaxed clock in this data set, with log Bayes factors of 5.48 for this
particular data set, which matches the simulation conditions. More importantly, even with such
high rate variation, the substitution rate estimated using a relaxed clock and the correct sampling
times included the true value used to generate the data (5×10^{-3} subs/site/unit time), with a 95%
highest posterior density (HPD) of between 2.15×10^{-3} and 1.90×10^{-2} subs/site/unit time, while the
regression slope was 2.22×10^{-2} subs/site/unit time. A key implication of these comparisons is that
BETS provides a formal assessment of temporal signal, unlike statistics computed from the
regression. Moreover, the root-to-tip regression appears uninformative when the data have been
sampled over a narrow time window and there is some rate variation among lineages.
298

Analyses of Empirical Data Sets

We analysed five empirical data sets with similar configurations of sampling times as in our
simulation study (Table 1). Two data sets consisted of rapidly evolving pathogens: *A/H1N1 influenza*
virus (Hedge et al. 2013) and *Bordetella pertussis* (Bart et al. 2014). We also analysed a data set with
highly divergent sequences of coronaviruses (Wertheim et al. 2013), and two data sets with ancient
DNA: *Hepatitis B virus* (Patterson Ross et al. 2018), and mitochondrial genomes of dog species
(Thalmann et al. 2013). Due to the demonstrated higher accuracy of GSS over NS (Fourment et al.
2019), we applied the BETS approach using the former method only.

The *A/H1N1 influenza virus* data demonstrated clear temporal signal, with the strict clock and
relaxed clock with the correct sampling times having the highest log marginal likelihoods, and a log
Bayes factor of M_{het} with respect to M_{iso} of 150 (fig. 6). The strict clock had higher support than the
relaxed clock for the correct sampling times (log Bayes factor 3.41). Broadly, this result is consistent
with previous evidence of strong temporal signal and clocklike behaviour in this data set (Hedge et
al. 2013). Using the strict clock with correct sampling times we estimated a substitution rate of
3.37×10^{-3} subs/site/year (HPD: 2.98×10^{-3} to 3.78×10^{-3}).

We detected temporal signal in the *Bordetella pertussis* data set (fig. 6). The relaxed clock with the
correct sampling times had the highest log marginal likelihood, with a log Bayes factor relative to
the strict clock of 28.86. The log Bayes factor for M_{het} relative to M_{iso} was 47.40. These results echo
previous assessments of these data using a date-randomization test (Duchene et al. 2016). We
estimated a mean substitution rate using the best model of 1.65×10^{-7} subs/site/year (95% HPD:
1.36×10^{-7} to 2.00×10^{-7}).

Our analyses did not detect temporal signal in the coronavirus data, for which the strict clock and
relaxed clock with no sampling times had the highest log marginal likelihoods. The log Bayes factor
of M_{het} relative to M_{iso} was -16.82, indicating strong support for the isochronous model. The relaxed
clock was supported over the strict clock, with a log Bayes factor of 19.25 (fig. 7). Previous analyses
of this data set suggested an ancient origin for this group of viruses, but here the lack of temporal
signal precludes any interpretation of our estimates of substitution rates and timescales.

The *Hepatitis B virus* data set included several human genotypes with complete genomes, where
135 were modern sequences collected from 1963 to 2013 and two were ancient samples from
human mummies from the 16[th] century. Previous studies have not found any temporal signal in
these data using different approaches, despite the inclusion of ancient sequences. Our estimates of
log marginal likelihoods were consistent with a lack of temporal signal, with a log Bayes factor of -
101.51 for M_{het} relative to M_{iso}.

The dog mitochondrial genome data contained samples from up to 36,000 years before the
present. BETS detected temporal signal in these data, with a log Bayes factor of 38.77 for M_{het}
relative to M_{iso}; this result is consistent with that of a date-randomization test in a previous study
(Tong et al. 2018). The estimated substitution rate for these data using the best model had a mean
of 1.08×10^{-7} subs/site/year (95% HPD: 7.49×10^{-8} to 1.52×10^{-7}).

Discussion

We have proposed BETS, a method that explicitly assesses the statistical support for including
sequence sampling times in a Bayesian framework. It is a test of the strength of the temporal signal
in a data set, which is an important prerequisite for obtaining reliable inferences in phylodynamic
analyses. BETS considers the model ensemble, such that the method can detect temporal signal
using models that account for substitution rate variation among lineages. The results of our

analyses demonstrate that our method is effective in a range of conditions, including when the
substitution rate is low or when the sampling window represents a small portion of the timespan of
the tree.
352
BETS does not require date permutations, which differentiates it from the widely used date-
randomization test for temporal structure. Date-randomization tests address the question of
whether a particular association between sequences and sampling times produces estimates
different from those obtained from data sets with permuted sampling times (Duchene et al. 2015;
Murray et al. 2015). However, such an approach is not a formal test of temporal signal in the data
because the permutations do not necessarily constitute an appropriate null model. In contrast, our
method does not require permutations and so has the benefit of being robust to using a small
number of sampling times.
361
Accurate calculations of marginal likelihoods are essential for BETS. In our simulation study, we
found that GSS and NS correctly assessed the presence and absence of temporal signal in the data
under most conditions. The correct clock model was also identified, although in a few instances NS
preferred an overparameterized model. Conceivably, using different marginal-likelihood estimators
might affect the actual model selected. Murray et al. (2015) also employed a Bayesian model-
testing approach using the AICM to assess temporal signal. In their study, the AICM performed well
in simulations, but failed to detect temporal signal in empirical data. We attribute this finding to the
low accuracy of AICM relative to path-sampling methods (Baele et al. 2012, 2013), and suggest
careful consideration of the marginal-likelihood estimator for tests of temporal signal.
371
A key advantage of BETS is that the complete model is considered, unlike in simpler data-
exploration methods such as root-to-tip regression. Specifically, root-to-tip regression is a visual
tool for uncovering problems with data quality and to inspect clocklike behaviour, but the absence
of appropriate statistics means that there is no clear way of determining whether the data contains
temporal information. Consider the regressions in figure 5 for data with a high substitution rate and
narrow sampling window. Even when among-lineage rate variation is low (σ=0.1), the data points
form a cloud, with a low R^2 of 0.09. However, the apparent 'noise' around the regression line is
probably the result of stochasticity in sequence evolution and of the narrow sampling window
relative to the age of the root of the tree. In fact, for this particular data set the model with the
highest log marginal likelihood is the strict clock with correct sampling times.
382
In all of our analyses, we ensured that the priors for different models and configurations of sampling
times were identical because, as with all Bayesian analyses, model comparison using marginal
likelihoods can depend on the choice of prior (Oaks et al. 2019). For example, the tree prior can
affect inferences of temporal signal, as it is part of the full model specification. Here we used an
exponential-growth coalescent tree prior, which closely matches the demographic dynamics of the
birth-death process under which the data were simulated. The effect of using an inappropriate tree
prior on tests of temporal signal requires further investigation, but previous studies have suggested
that there is only a small impact on estimates of rates and times if the sequence data are
informative (Ritchie et al. 2017; Möller et al. 2018).
392
An interesting finding is that statistical support for isochronous sampling times in truly isochronous
data is lower than that for the correct sampling times in truly heterochronous data. This can
potentially lead to an increased risk of incorrectly concluding the presence of temporal signal, but
we only found this to be a problem in a small number of cases. In particular, in isochronous data
simulated with a low substitution rate, and with very few variable sites, the best models were
sometimes those that included sampling times, albeit with very low log Bayes factors (e.g.,

supplementary fig. S2, Supplementary Material online). This probably occurs because stochastic
error associated with a small amount of evolution leads to low power for model selection.
401
Permuting sampling times led to poor model fit, as expected. This procedure requires substantial
computing requirements, depending on the number of permutations that are performed, and we
find that such date permutations are of limited value for model testing when the data are highly
informative (e.g., figs. 1 and 3). However, in data sets with very low information content, such as
those that were produced by simulation with a low substitution rate here, conducting a small
number of date permutations might offer a conservative approach to determining whether model
fit and parameter estimates are driven by a particular set of sampling times, as one would expect in
the presence of temporal signal.
410
The nature of the BETS approach means that every parameter in the model has a prior probability,
including the substitution rate. Because substitution rates and times are nonidentifiable, it is
conceivable that an informative prior on the rate or on the age of an internal node might have a
stronger effect than the sampling times on the posterior, for example if the samples span a very
short window of time. Such analyses with informative substitution rate priors effectively include
several simultaneous sources of calibrating information (i.e., sampling times, internal nodes, and an
informative rate prior). Using sampling times in addition to other sources of calibration information
might still be warranted if it improves the fit of the model, which can be tested using our proposed
method.
420
Analyses with multiple calibrations can also allow uncertainty in sequence sampling times,
especially in data sets that include ancient DNA, where sampling times can be treated as
parameters in the model (Shapiro et al. 2011). BETS provides a coherent approach to assess
temporal structure in these circumstances, unlike date-randomization tests that typically use point
values for sampling times. In fact, BETS can be used as a means to validate whether a sample is
modern or ancient.
427
428 In general, the uptake of Bayesian model testing in phylogenetics has great potential for improving
429 our confidence in estimates of substitution rates and timescales. The test that we have proposed
430 here, BETS, provides a coherent and intuitive framework to test for temporal information in the
431 data.
432
Materials and Methods

Simulations

We simulated phylogenetic trees under a stochastic birth-death process using MASTER v6.1
(Vaughan and Drummond 2013), by specifying birth rate λ =1.5, death rate μ =0.5, and sampling
rate ϕ =0.5. This corresponds to an exponentially growing infectious outbreak with reproductive
number R_0=1.5 and a wide sampling window. We set the simulation time to 5 units of time, which
corresponds to the time of origin of the process. For isochronous trees, we used similar settings, but
instead of using the sampling rate, we sampled each tip with probability ρ =0.5 when the process
was stopped after 5 units of time (i.e. μ =1 and ϕ =0). Some of our analyses consisted of artificially
specifying sampling times for isochronous trees, which we set to those that we would have
obtained from a birth-death process with μ =0.5 and ϕ =0.5.
444
In a second set of simulations of heterochronous trees, we generated trees with a narrow sampling
window. We specified two intervals for μ and ϕ. The first interval spanned 4.5 units of time with
μ =1.0 and ϕ =0, and the second interval 0.5 units of time with μ =0.1 and ϕ =0.9. As a result, the
process still had a constant become uninfectious rate (μ + ϕ), but samples were only collected in

the second interval. The high sampling rate in the second interval resulted in trees with similar
numbers of tips to those with a wide sampling window, but where their ages only spanned 0.5 units
of time.
452
We only considered the simulated trees that contained between 90 and 110 tips. The trees
generated in MASTER are chronograms (with branch lengths in units of time), so we simulated
substitution rates to generate phylograms (with branch lengths in units of subs/site). To do this we
specified the uncorrelated lognormal relaxed clock with a mean rate of 5×10^{-3} or 10^{-5} subs/site/unit
time and a standard deviation σ of 0 (corresponding to a strict clock), 0.1, 0.5, or 1. We simulated
sequence evolution along these phylograms under the HKY nucleotide substitution model
(Hasegawa et al. 1985). We added among-site rate variation using a discretized gamma distribution
(Yang 1994, 1996) using Phangorn v2.5 (Schliep 2011) to generate sequence alignments of 10,000
nucleotides. We set the transition-to-transversion ratio of the HKY model to 10 and the shape of the
gamma distribution to 1, which is similar to estimates of these parameters in influenza viruses
(Duchene et al. 2014; Hedge and Wilson 2014). For each simulation scenario we generated 10
sequence alignments.
465
Estimation of Marginal Likelihoods Using Nested Sampling
We analysed the data in BEAST 2.5 using the matching substitution model, the exponential-growth
coalescent tree prior, the strict clock or relaxed clock, and different configurations of sampling
times. We chose the exponential-growth coalescent tree prior, instead of the birth-death tree prior,
because it is conditioned on the samples instead of assuming a sampling process; this ensures that
the marginal likelihoods for isochronous and heterochronous trees are comparable.
472
We specified proper priors on all parameters, which is essential for accurate estimation of marginal
likelihoods (Baele et al., 2013). In our heterochronous analyses the prior on the substitution rate had
a uniform distribution bounded between 0 and 1. We made this arbitrary choice to set a somewhat
uninformative prior and because the default prior in BEAST 2.5 is a uniform distribution between 0
and infinity, which is improper. Owing to the nonidentifiability of substitution rates and times,
neither can be inferred in the absence of calibrating information, so in our isochronous analyses we
fixed the value of the substitution rate to 1. The initial NS chain length was chosen so as to draw
20,000 samples, with 20,000 steps, 32 particles, and a subchain length of 5,000 (note that NS is not
equivalent to standard MCMC, nor is the definition of an iteration/step). The chain length and its
accompanying sampling frequency were adjusted to obtain effective sample sizes for key
parameters of at least 200 (computed in the NS output in BEAST 2.5). Examples of MASTER files
and BEAST input files for NS are available online (supplementary data, Supplementary Material
online).
486
Estimation of Marginal Likelihoods Using Generalized Stepping-Stone Sampling
We used BEAST 1.10 with the same model specifications and priors as in BEAST2, except for the
prior on the substitution rate, for which we used the approximate continuous-time Markov chain
reference prior (Ferreira and Suchard 2008). Because our simulation analyses of GSS and NS differ
in this prior, the marginal-likelihood estimates are not directly comparable, so for each simulation
we report log Bayes factors of competing models instead of the log marginal likelihoods. The GSS
implementation in BEAST 1.10 has two different working priors for the tree generative process: a
matching tree prior and a product of exponentials. The latter approach is the most generally
applicable and is the one that we used here (Baele et al. 2016).
496
We used an initial MCMC chain length of 5×10^7 steps sampling every 5000 steps. After discarding
10% of the samples obtained, the remaining samples were used to construct the working
distributions for the GSS analysis. These comprised 100 path steps distributed according to

quantiles from a β distribution with α =0.3, with each of the 101 resulting power posterior
inferences running for 5×10^5 iterations. We assessed sufficient sampling for the initial MCMC
analysis by verifying that the effective sample sizes for key parameters were at least 200 in Coda
v0.19 (Plummer et al. 2006). If this condition was not met, we doubled the length of the MCMC and
reduced sampling frequency accordingly. Examples of MASTER files and BEAST input files for GSS
are available online (supplementary data, Supplementary Material online).
506

Analyses of Empirical Data Sets

We downloaded sequence alignments from their original publications (Table 1): complete genomes
of *A/H1N1 influenza virus* (Hedge et al. 2013), whole genome sequences of *B. pertussis* (Bart et al.
2014; Duchene et al. 2016), RdRP sequences of coronaviruses (Wertheim et al. 2013), complete
genomes of *Hepatitis B virus* (Patterson Ross et al. 2018), and dog mitochondrial genomes
(Thalmann et al. 2013). The data and BEAST input files are available in the Supplementary Material
online.
514

Briefly, we used similar settings as in our simulations to estimate marginal likelihoods using GSS.
For sequence sampling times we considered the correct sampling times, no sampling times (i.e.,
isochronous), and permuted sampling times. We also specified tree priors as follows: an
exponential-growth coalescent for the *A/H1N2 influenza virus*, *Bordetella pertussis*, coronaviruses,
and *Hepatitis B virus* data sets, and a constant-size coalescent for the dog mitochondrial genomes
as used by Tong et al. (2018). We again chose the HKY+Γ substitution model, except in the analysis
of *Hepatitis B virus* data, for which we used the GTR+Γ model (Tavaré 1986), and in the analysis of
the dog data set for which we used the SRD06 substitution model (Shapiro et al. 2006) for coding
regions and the GTR+Γ for noncoding regions.
524

525 **Supplementary Material**

526 Supplementary data are available online.
527

Funding

SD was supported by an Australian Research Council Discovery Early Career Researcher Award
(DE190100805) and an Australian National Health and Medical Research Council grant
(APP1157586). PL acknowledges funding from the European Research Council under the European
Union's Horizon 2020 research and innovation programme (grant agreement no. 725422-
ReservoirDOCS) and the Research Foundation -- Flanders ('Fonds voor Wetenschappelijk
Onderzoek -- Vlaanderen', G066215N, G0D5117N and G0B9317N). SYWH was funded by the
Australian Research Council (FT160100167). GB acknowledges support from the Interne Fondsen
KU Leuven / Internal Funds KU Leuven under grant agreement C14/18/094. VD was supported by
contract HHSN272201400006C from the National Institute of Allergy and Infectious Diseases,
National Institutes of Health, U.S. Department of Health and Human Services, USA.

Acknowledgements

Pending.
542

References

Baele G., Lemey P. 2014. Bayesian model selection in phylogenetics and genealogy-based
population genetics. In: Chen M.-H., Kuo L., Lewis P.O., editors. Bayesian Phylogenetics,
Methods, Algorithms, and Applications. Boca Raton, Florida: CPC Press. p. 59–93.
Baele G., Lemey P., Bedford T., Rambaut A., Suchard M.A., Alekseyenko A. V. 2012. Improving the
accuracy of demographic and molecular clock model comparison while accommodating

phylogenetic uncertainty. Mol. Biol. Evol. 29:2157–2167.

Baele G., Lemey P., Suchard M.A. 2016. Genealogical working distributions for Bayesian model
testing with phylogenetic uncertainty. Syst. Biol. 65:250–264.

Baele G., Li W.L.S., Drummond A.J., Suchard M.A., Lemey P. 2013. Accurate model selection of
relaxed molecular clocks in bayesian phylogenetics. Mol. Biol. Evol. 30:239–243.

Bart M.J., Harris S.R., Advani A., Arakawa Y., Bottero D., Bouchez V., Cassiday P.K., Chiang C.-S.,
Dalby T., Fry N.K. 2014. Global population structure and evolution of Bordetella pertussis and
their relationship with vaccination. MBio. 5:e01074-14.

Biek R., Pybus O.G., Lloyd-Smith J.O., Didelot X. 2015. Measurably evolving pathogens in the
genomic era. Trends Ecol. Evol. 30:306–313.

Bouckaert R., Vaughan T.G., Barido-Sottani J., Duchene S., Fourment M., Gavryushkina A., Heled J.,
Jones G., Kuhnert D., de Maio N. 2019. BEAST 2.5: An Advanced Software Platform for
Bayesian Evolutionary Analysis. PLOS Comput. Biol. 15:e1006650.

Bromham L., Duchêne S., Hua X., Ritchie A.M., Duchêne D.A., Ho S.Y.W. 2018. Bayesian molecular
dating: Opening up the black box. Biol. Rev. 93:1165–1191.

Drummond A.J., Ho S.Y.W., Phillips M.J., Rambaut A. 2006. Relaxed phylogenetics and dating with
confidence. PLOS Biol. 4:e88.

Drummond A.J., Pybus O.G., Rambaut A., Forsberg R., Rodrigo A.G. 2003. Measurably evolving
populations. Trends Ecol. Evol. 18:481–488.

Duchene S., Duchene D.A., Geoghegan J.L., Dyson Z.A., Hawkey J., Holt K.E. 2018. Inferring
demographic parameters in bacterial genomic data using Bayesian and hybrid phylogenetic
methods. BMC Evol. Biol. 18:95.

Duchene S., Duchene D.A., Holmes E.C., Ho S.Y.W. 2015. The performance of the date-
randomization test in phylogenetic analyses of time-structured virus data. Mol. Biol. Evol.
32:1895–1906.

Duchene S., Holmes E.C., Ho S.Y.W. 2014. Analyses of evolutionary dynamics in viruses are
hindered by a time-dependent bias in rate estimates. Proc. R. Soc. London B. 281:20140732.

Duchene S., Holt K.E., Weill F.-X., Le Hello S., Hawkey J., Edwards D.J., Fourment M., Holmes E.C.
2016. Genome-scale rates of evolutionary change in bacteria. Microb. Genomics. 2:e000094.

Fan Y., Wu R., Chen M.-H., Kuo L., Lewis P.O. 2011. Choosing among partition models in Bayesian
phylogenetics. Mol. Biol. Evol. 28:523–532.

Ferreira M.A.R., Suchard M.A. 2008. Bayesian analysis of elapsed times in continuous-time Markov
chains. Can. J. Stat. 36:355–368.

Fourment M., Magee A.F., Whidden C., Bilge A., Matsen I. V, Frederick A., Minin V.N. 2019. 19
dubious ways to compute the marginal likelihood of a phylogenetic tree topology. Syst.
Biol.:syz046.

Guindon S., Dufayard J.-F., Lefort V., Anisimova M., Hordijk W., Gascuel O. 2010. New algorithms
and methods to estimate maximum likelihood phylogenies: assessing the performance of
PhyML 3.0. Syst. Biol. 59:307–321.

Hasegawa M., Kishino H., Yano T. 1985. Dating of the human-ape splitting by a molecular clock of
mitochondrial DNA. J. Mol. Evol. 22:160–174.

Hedge J., Lycett S.J., Rambaut A. 2013. Real-time characterization of the molecular epidemiology
of an influenza pandemic. Biol. Lett. 9:20130331.

Hedge J., Wilson D.J. 2014. Bacterial phylogenetic reconstruction from whole genomes is robust to
recombination but demographic inference is not. MBio. 5:e02158-14.

Hipsley C.A., Müller J. 2014. Beyond fossil calibrations: realities of molecular clock practices in
evolutionary biology. Front. Genet. 5:138.

Ho S.Y.W., Duchene S. 2014. Molecular-clock methods for estimating evolutionary rates and time
scales. Mol. Ecol. 23:5947–5975.

Kass R.E., Raftery A.E. 1995. Bayes factors. J. Am. Stat. Assoc. 90:773–795.

Korber B., Muldoon M., Theiler J., Gao F., Gupta R., Lapedes A., Hahn B.H., Wolinsky S.,

Bhattacharya T. 2000. Timing the ancestor of the HIV-1 pandemic strains. Science. 288:1789–
1796.
Lartillot N., Philippe H. 2006. Computing Bayes factors using thermodynamic integration. Syst.
Biol. 55:195–207.
Maturana P., Brewer B.J., Klaere S., Bouckaert R. 2019. Model selection and parameter inference in
phylogenetics using Nested Sampling. Syst. Biol. 68:219–233.
Möller S., du Plessis L., Stadler T. 2018. Impact of the tree prior on estimating clock rates during
epidemic outbreaks. Proc. Natl. Acad. Sci. USA. 115:4200–4205.
Murray G.G.R., Wang F., Harrison E.M., Paterson G.K., Mather A.E., Harris S.R., Holmes M.A.,
Rambaut A., Welch J.J. 2015. The effect of genetic structure on molecular dating and tests for
temporal signal. Methods Ecol. Evol. 7:80–89.
Nascimento F.F., dos Reis M., Yang Z. 2017. A biologist's guide to Bayesian phylogenetic analysis.
Nat. Ecol. Evol. 1:1446.
Newton M.A., Raftery A.E. 1994. Approximate Bayesian inference with the weighted likelihood
bootstrap. J. R. Stat. Soc. Ser. B. 56:3–26.
Oaks J.R., Cobb K.A., Minin V.N., Leaché A.D. 2019. Marginal likelihoods in phylogenetics: a review
of methods and applications. Syst. Biol. 68:681–697.
Patterson Ross Z., Klunk J., Fornaciari G., Giuffra V., Duchene S., Duggan A.T., Poinar D., Douglas
M.W., Eden J.-S., Holmes E.C., Poinar H.N. 2018. The paradox of HBV evolution as revealed
from a 16thcentury mummy. PLoS Pathog. 14:e1006887.
Plummer M., Best N., Cowles K., Vines K. 2006. CODA: Convergence diagnosis and output analysis
for MCMC. R News. 6:7–11.
Raftery A., Newton M., Satagopan J., Krivitsky P. 2007. Estimating the integrated likelihood via
posterior simulation using the harmonic mean identity. Bayesian Statistics. Oxford University
Press. p. 1–45.
Rambaut A., Lam T.T., Carvalho L.M., Pybus O.G. 2016. Exploring the temporal structure of
heterochronous sequences using TempEst (formerly Path-O-Gen). Virus Evol. 2:vew007.
Ramsden C., Holmes E.C.C., Charleston M.A.A. 2009. Hantavirus evolution in relation to its rodent
and insectivore hosts: no evidence for codivergence. Mol. Biol. Evol. 26:143–153.
Rieux A., Balloux F. 2016. Inferences from tip-calibrated phylogenies: a review and a practical guide.
Mol. Ecol. 25:1911–1924.
Ritchie A.M., Lo N., Ho S.Y.W. 2017. The impact of the tree prior on molecular dating of data sets
containing a mixture of inter-and intraspecies sampling. Syst. Biol. 66:413–425.
Schliep K.P. 2011. phangorn: Phylogenetic analysis in R. Bioinformatics. 27:592–593.
Shapiro B., Ho S.Y.W., Drummond A.J., Suchard M.A., Pybus O.G., Rambaut A. 2011. A Bayesian
phylogenetic method to estimate unknown sequence ages. Mol. Biol. Evol. 28:879–887.
Shapiro B., Rambaut A., Drummond A.J. 2006. Choosing appropriate substitution models for the
phylogenetic analysis of protein-coding sequences. Mol. Biol. Evol. 23:7–9.
Skilling J. 2006. Nested sampling for general Bayesian computation. Bayesian Anal. 1:833–859.
Suchard M.A., Lemey P., Baele G., Ayres D.L., Drummond A.J., Rambaut A. 2018. Bayesian
phylogenetic and phylodynamic data integration using BEAST 1.10. Virus Evol. 4:vey016.
Tavaré S. 1986. Some probabilistic and statistical problems in the analysis of DNA sequences. Lect.
Math. Life Sci. 17:57–86.
Thalmann O., Shapiro B., Cui P., Schuenemann V.J., Sawyer S.K., Greenfield D.L., Germonpré M.B.,
Sablin M. V, López-Giráldez F., Domingo-Roura X. 2013. Complete mitochondrial genomes of
ancient canids suggest a European origin of domestic dogs. Science. 342:871–874.
Tong K.J., Duchêne D.A., Duchêne S., Geoghegan J.L., Ho S.Y.W. 2018. A comparison of methods
for estimating substitution rates from ancient DNA sequence data. BMC Evol. Biol. 18:70.
Trovão N.S., Baele G., Vrancken B., Bielejec F., Suchard M.A., Fargette D., Lemey P. 2015. Host
ecology determines the dispersal patterns of a plant virus. Virus Evol. 1:vev016.
Vaughan T.G., Drummond A.J. 2013. A stochastic simulator of birth–death master equations with

application to phylodynamics. Mol. Biol. Evol. 30:1480–1493.

Wertheim J.O., Chu D.K.W., Peiris J.S.M., Pond S.L.K., Poon L.L.M. 2013. A case for the ancient

origin of coronaviruses. J. Virol. 87:7039–7045.

Xie W., Lewis P.O., Fan Y., Kuo L., Chen M.H. 2011. Improving marginal likelihood estimation for

Bayesian phylogenetic model selection. Syst. Biol. 60:150–160.

Yang Z. 1994. Estimating the pattern of nucleotide substitution. J. Mol. Evol. 39:105–111.

Yang Z. 1996. Among-site rate variation and its impact on phylogenetic analyses. Trends Ecol. Evol.

11:367–372.

Zuckerkandl E., Pauling L. 1965. Evolutionary divergence and convergence in proteins. In: Bryson

V., Vogel H., editors. Evolving Genes and Proteins. New York: Academic press. p. 97–166.

Figure Legends

FIG. 1. Log Bayes factors of heterochronous data simulated with a high substitution rate and wide
sampling window. Each panel shows the results for data sets simulated with a different degree of
among-lineage rate variation, governed by the standard deviation σ of a lognormal distribution. In
each plot the *x*-axis depicts six analysis settings, with two clock models, strict clock (SC) and the
uncorrelated relaxed clock with an underlying lognormal distribution (UCLN), and three settings for
sampling times: generated under the birth-death process (BD), identical sampling times
(Isochronous), and permuted (Permuted). The points have been jittered along the *x*-axis to facilitate
visualization. The *y*-axis shows log Bayes factors relative to the best model. Red points correspond
to estimates using generalized stepping-stone sampling and blue points correspond to estimates
using nested sampling. We conducted 10 simulation replicates, with each replicate data set
analysed under the six analysis settings and two marginal-likelihood estimators, such that
stochastic error might cause differences in the preferred model. The number next to each cloud of
points denotes the number of times (out of 10) that the corresponding model had the highest
marginal likelihood with generalized stepping-stone sampling (red) and nested sampling (blue).

FIG. 2. Log Bayes factors of heterochronous data simulated under a low substitution rate and a wide
sampling window. Symbols and colours are the same as those in figure 1.

FIG. 3. Log Bayes factors of heterochronous data simulated under a high substitution rate and
narrow sampling window. Symbols and colours are the same as those in figure 1.

FIG. 4. Log Bayes factors of heterochronous data simulated under a low substitution rate and
narrow sampling window. Symbols and colours are the same as those in figure 1.

FIG. 5. Root-to-tip regressions for a subset of data sets simulated with varying degrees of among-
lineage rate variation (governed by the standard deviation σ of a lognormal distribution), using a
high substitution rate and either a wide or narrow sampling window. The *y*-axis is the root-to-tip
distance and the *x*-axis is the time from the youngest tip, where 0 is the present. Each point
corresponds to a tip in the tree and the solid line is the best-fit linear regression using least-squares.
The coefficient of determination, R^2, is shown in each case. For comparison, the log Bayes factors of
the best heterochronous model relative the best isochronous model, BF(M_{het}-M_{iso}), are also shown.

FIG. 6. Log marginal likelihoods estimated using generalized stepping-stone sampling for six
analysis settings for sequence data from rapidly evolving pathogens, *A/H1N1 Human influenza virus*
and *Bordetella pertussis*. The *y*-axis is the marginal likelihood and the *x*-axis shows the analysis
settings, with two clock models, strict clock (SC) and the uncorrelated relaxed clock with an
underlying lognormal distribution (UCLN), and three settings for sampling times: generated under
the birth-death process (BD), identical sampling times (Isochronous), and permuted (Permuted).
Solid points and dashed lines correspond to the log marginal-likelihood estimates. The asterisk
denotes the model with the highest marginal likelihood.

FIG. 7. Log marginal likelihoods estimated using generalized stepping-stone sampling for six
analysis settings for data sets with ancient DNA or highly divergent sequences. The *y*-axis is the
marginal likelihood and the *x*-axis shows the analysis settings, with two clock models, strict clock
(SC) and the uncorrelated relaxed clock with an underlying lognormal distribution (UCLN), and
three settings for sampling times: generated under the birth-death process (BD), identical sampling
times (Isochronous), and randomized (Random). Solid points and dashed lines correspond to the log
marginal-likelihood estimates. The asterisk denotes the model with the highest marginal likelihood.

716   **Tables**

719   **Table 1.** Details of empirical data sets used in this study.

Data set	Number of sites (nucleotides)	Number of samples	Sampling time range	Reference
A/H1N1 influenza virus	13,154	329	10 months (March to December 2009)	Hedge et al. (2013)
Bordetella pertussis	4.9×10^6	150	89 years (1920 to 2009)	Bart et al. (2014)
Coronaviruses	1,860	43	70 years (1941 to 2011)	Wertheim et al. (2013)
Hepatitis B virus	3,271	137	445 years (2103 to 1568)	Patterson Ross et al. (2018)
Dog mtDNA	14,596	50	36,000 years (to the present)	Thalmann et al. (2013)

Supplementary Material

FIG. S1. Log Bayes factors of isochronous data simulated with a high substitution rate. Each panel
shows the results for data sets simulated with a different degree of among-lineage rate variation,
governed by the standard deviation σ of a lognormal distribution. The x-axis depicts six analysis
settings, with two molecular clock models, strict clock (SC) and the uncorrelated relaxed clock with
an underlying lognormal distribution (UCLN), and three settings for sampling times: generated
under the birth-death process (BD), identical sampling times (Isochronous), and permuted
(Permuted). The points have been jittered to facilitate visualization. The y-axis shows log Bayes
factors relative to the best model. Red points correspond to estimates using generalized stepping-
stone sampling and blue points correspond to estimates using nested sampling. We conducted 10
simulation replicates, with each replicate data set analysed under the six analysis settings and two
marginal-likelihood estimators, such that stochastic error might cause differences in the preferred
model. The number next to each cloud of points denotes the number of times (out of 10) that the
corresponding model had the highest marginal likelihood with generalized stepping-stone sampling
(red) and nested sampling (blue).

FIG. S2. Log Bayes factors of isochronous data simulated with a low substitution rate. Symbols and
colours are the same as those in figure 1.

Fɪɢ. S3. Example of three phylogenetic trees used in simulations. Red dashed lines indicate the
times of each of the tips and therefore represent the sampling process over time. All trees are
simulated under a birth-death process with time of origin of 5, such that the sum of the tree height
and the length of the stem branch leading to the root is always 5. In all trees, we set the birth rate
λ =1.5, and become uninfectious rate δ =1, where $\delta = \mu + \phi$, where μ is the death rate and ϕ is
the sampling rate upon death. Thus, the population growth rate is constant and the same across all
trees. The top tree assumes a constant sampling process and a wide sampling window (ϕ =0.5
throughout the whole process), whereas in the second tree sampling starts after 4.5. Before this
time the sampling rate, ϕ_0, is zero. After 4.5 time units the sampling rate ϕ_1 is 0.9 (and thus mu_1
= 0.1), resulting in a narrow sampling window. The bottom tree has samples drawn at a single point
in time with a sampling probability at present, ρ, of 0.5 (and thus phi=0).

Coronaviruses

Hepatitis B virus with ancient samples

Dog mtDNA with ancient samples

Log marginal likelihood

| SC | UCLN | SC | UCLN | SC | UCLN |

BD   Isochronous   Permuted