Network-based metrics of community resilience and dynamics in lake ecosystems

David I. Armstrong M^cKay*[1,2,3], James G. Dyke[1,4], John A. Dearing[1],

C. Patrick Doncaster[5], Rong Wang[6]

[1]Geography and Environment, University of Southampton, Southampton, UK, SO17 1BJ

(work started here)

[2]Stockholm Resilience Centre, Stockholm University, SE-10691 Stockholm, Sweden (current

address DIAM)

[3]Bolin Centre for Climate Research, Stockholm University, SE-10691 Stockholm, Sweden

[4]Global Systems Institute, College of Life and Environmental Sciences, University of Exeter,

Exeter, UK (current address JGD)

[5]Biological Sciences, University of Southampton, Southampton, UK, SO17 1BJ

[6]State Key Laboratory of Lake Science and Environment, Nanjing Institute of Geography and

Limnology, Chinese Academy of Sciences, China, 210008

***david.armstrongmckay@su.se**

Paper in review at: Royal Society Biology Letters

Abstract

Some ecosystems can undergo abrupt critical transitions to a new regime after passing

a tipping point, such as a lake shifting from a clear to turbid state as a result of eutrophication.

Metrics-based resilience indicators acting as early warning signals of these shifts have been

developed but have not always been reliable in all systems. An alternative approach is to

focus on changes in the structure and composition of an ecosystem, but this can require long-

term food-web observations that are typically unavailable. Here we present a network-based

algorithm for estimating community resilience, which reconstructs past ecological networks

from palaeoecological data using metagenomic network inference. Resilience is estimated

using local stability analysis and eco-net energy (a neural network-based proxy for

"ecological memory"). The algorithm is tested on model (PCLake+) and empirical (lake

Erhai) data. We find evidence of increasing diatom community instability during

eutrophication in both cases, with changes in eco-net energy revealing complex eco-memory

dynamics. The concept of ecological memory opens a new dimension for understanding

ecosystem resilience and regime shifts, and further work is required to fully explore its drivers

and implications.

Keywords: Lake Eutrophication; Early Warning Signals; Resilience; Palaeoecology; Neural

Networks

1. Introduction

The potential for pressurised ecosystems to abruptly shift to a new regime once tipping points are reached has led to a proliferation of metrics attempting to quantify ecosystem resilience [1–10]. Lake eutrophication is a well-studied example of an ecological tipping point, with evidence of alternative stable states, regime shifts, and strong hysteresis [11–13]. Eutrophication occurs when increasing nutrient loading triggers positive feedbacks driving a rapid shift from clear to turbid conditions [14,15], with recovery requiring nutrient levels to be reduced far below the original threshold [13]. This should be reflected by declining ecosystem resilience, defined here as a relative weakening of negative feedbacks resulting in greater sensitivity to small shocks [16]. However, current metrics for evaluating resilience loss (e.g. time-series metrics of critical slowing down) have mixed reliability when applied as early warning signals (EWS) in freshwater [17] and other ecosystems [18–21]. Methodological issues include the choice of system variable for analysis, false or missed alarms, and subjective parameter choices, while palaeorecord applications presents an additional challenge of variable temporal resolution due to compaction or changing accumulation rates [7,22–25].

An alternative approach to metric-based "classical" EWS is to analyse ecosystem structure in order to detect the changing dynamics of the whole ecosystem without having to understand the full trophic ecology of any one species. To this end Doncaster et al. [26] assessed the compositional disorder of diatoms and chironomids from sediment cores in three Chinese lakes (Erhai, Taibai, and Longgan). The correlation of nestedness and biodiversity in these lakes becomes negative prior to the critical transition before reverting to positive values, which was hypothesised to result from eutrophication shifting the competitive balance between 'keystone' and 'weedy' species. However, the link between this and ecosystem resilience is indirect and the competition dynamics remain theoretical. Another recent network-based method is nodal degree skewness, with increasing human pressure inducing

more negative skewness in Chinese lake ecosystems [27].

An alternative approach is to reconstruct the lake's food web and monitor for signs of

dynamical instability. Kuiper et al. [28] showed that food-web data can be used to reconstruct

the interactions between different species, which correspond to the interaction matrix of a

Lotterra ecosystem model. Given food-web observations, the food web's local stability

can be estimated from the intraspecific interaction strengths or the maximum loop weight of

the ecosystem's longest trophic loop, both of which are closely related to the system's

dominant eigenvalue (λ_d) [29,30]. This approach enabled destabilising food web

reorganisations prior to eutrophication to be tracked in the PCLake model [28]. However, this

requires detailed food-web data to be generated at regular intervals during eutrophication, a

laborious task which is often unavailable for real-world lakes.

Many lakes nevertheless have palaeoecological records of species abundances from

sediment cores. If past ecosystem structure can be reconstructed from abundance data, it

would be possible to estimate changes in ecosystem resilience over time. Here we develop

and apply an algorithm for reconstructing ecological networks (eco-nets) from

palaeoecological data using metagenomic network inference [31–34]. We then estimate

ecosystem resilience using local stability analysis and novel neural network methods.

2.   Methods

2.1.   Eco-Network Inference

All analysis was conducted in R [35]. Abundance data was first transformed by

isometric log ratio to account for compositionality in relative data and imputed to counter data

sparseness [36] to form abundance matrix X. To find the structure of the eco-net, we followed

the methods of metagenomics network inference [33,34]. We first assumed that ecological

interactions follow a generalised Lotka-Volterra model:

$$\frac{dx_i(t)}{dt} = x_i(t)\left[\beta_i - \sum_j \alpha_{ij}x_j(t)\right]$$

<div align="right">**Equation 1**</div>

89   where x is the abundance of species i or j, α_{ij} is the interaction coefficient between species i

90   and j ($<-1 < \alpha_{ij} < 1$), and β_i is the constant unitary net growth rate of species i. Taking the

91   logarithm of a discrete-time Lotka-Volterra model based on Equation 1 and assuming constant

92   time-steps [33] gives:

$$\ln x_i(t+1) - \ln x_i(t) = \zeta_i(t) + \beta_i - \sum_j \alpha_{ij}(x_j(t) - \langle x_j \rangle)$$

<div align="right">**Equation 2**</div>

where ζ is the noise term and $\langle x_j \rangle$ is the median abundance (approximating to equilibrium

abundance and carrying capacity for normalising against; assumes sufficient data to capture

equilibrium population dynamics and ephemeral species). Given abundance time-series, the

interaction coefficients α_{ij} can be found by performing multiple linear regression (lasso

regularised due to data sparseness and limited observations relative to variables, and k-Fold

cross-validated to improve prediction performance) of $[\ln x_i(t+1) - \ln x_i(t)]$ against

$[x_j(t) - \langle x_j \rangle]$. The regression coefficients populate the elements of interaction matrix A,

which represents the eco-net's structure at time t.

We use only non-interpolated data to avoid well-known biases interpolation introduces

to time-series analysis [24], although this results in slightly varying time-steps. Network

inference and linear regression are less sensitive to non-equidistant time points than time-

series analysis [32], but the discrete-time Lotka-Volterra model still assumes constant time-

steps. To minimise this limitation, input data should be scrutinised prior to analysis and

sections with substantial shifts in time resolution excluded.

2.2.   Local Stability

The eco-net's stability was estimated using local stability analysis, by taking the

absolute of the interaction matrix A (which is equivalent to the Jacobian of the Lotka-Volterra

system [28,37,38]; the system is considered stable at its equilibrium point if the real part of

the Jacobian's eigenvalues remain negative, indicating net negative feedbacks), setting α_{ii} to

0 (to satisfy the Perron-Frobenius Theorem [29]), and calculating the dominant eigenvalue λ_d

(which is tightly related to other ecosystem stability metrics [28–30]). This was calculated on

a 50% rolling window to estimate changes in stability over time.

2.3.   Ecological Memory

Recent theory suggests that ecosystems may have a distributed 'memory' of past states

as a result of a process akin to unsupervised Hebbian learning in a neural network [39,40]. In

Hopfield Networks frequent correlations between neurones (nodes) lead to an increase in their

synaptic connection strength (edges) – a process described colloquially as "*neurons that fire*

together, wire together" [41–44]. Over time this allows the emergence of distributive

associative memory of training data that can be recovered when given degraded input. Power

et al. [39,40] proposed that eco-nets experience similar dynamics, with frequently co-

occurring species (nodes) developing stronger interactions (edges), allowing a distributed

associative "ecological memory" (eco-memory) of past environmental forcing (training data)

to emerge (Figure 1). We have no direct metric of Hopfield Network memory strength, but we

can calculate its energy, which is minimised at stable points. By assuming the eco-net's

interaction matrix A is equivalent to a neural network's weight matrix W [39] and treating it

as a continuous Hopfield Network, we posit that the eco-net energy, E_n, can be calculated by:

130
$$E_n = -\frac{1}{2}\mathbf{X(t)}^T \mathbf{A X(t)}$$

Equation 3

where $\mathbf{X(t)}$ is a vector of species abundances at time t [41–44]. We hypothesise that stable

eco-nets 'learning' from long-term environmental conditions leads to lower E_n, and that

destabilisation would conversely result in higher E_n. However, asymmetric matrices with non-

zero diagonals, such as community matrices, can lead to chaotic behaviour and the network

converging to alternative less-stable points, meaning eco-nets may not always completely

minimise their energy.

2.4.   Comparison Metrics

For comparison we also calculated time-series metrics (Z-scores of AR1, standard

deviation (SD), skewness, and kurtosis) on keystone abundance (Cyclostephanos Dubius;

Erhai) or surface chlorophyll concentration (PCLake+) with variance-stabilising square-root

transformation and Gaussian detrending (bandwidth: PCLake+ 1%; Erhai 5%) (R:

generic_ews, "*earlywarnings*"[6]). However, due to methodological limitations [7,22–25]

time-series metrics are shown for comparison only and are not useful results by themselves.

We also calculate compositional disorder metrics [26] (nestedness, biodiversity (inverse

Simpson index), and their correlation; R: *nestedtemp & diversity*, "*vegan*"[45]) where

possible.

3.   Test-Case 1: PCLake+

We first apply the algorithm to output from PCLake+ [46], an extension of the widely

used PCLake model of lake eutrophication, as a test-bed with well-known dynamics and

drivers. PCLake+ is initialised in default clear state with monthly time-steps, run for 200

years to ensure stability, and nutrient input increased (0.0001 to 0.005 $gPm^{-2}d^{-1}$) over 50 years

to induce eutrophication. Dry-weight ecosystem outputs (binned to represent sedimentary

preservation with variable accumulation rates) are used as abundance inputs for the algorithm.

We cannot compute nestedness for PCLake+ as only functional groups rather than individual

species are available.

The impact of nutrient input is clearly visible (Fig. 2a) in both local stability (λ_d) and

eco-net energy (E_n), with λ_d increasing shortly after input begins in conjunction with a peak in

E_n at ~50 cm (along with increases in time-series metrics and biodiversity). However, after a

second λ_d peak (~35 cm) both λ_d and especially E_n decrease unexpectedly (whilst apart from

standard deviation the time-series and biodiversity metrics plateau or decline). This is

contrary to our hypothesis that destabilisation should lead to higher E_n, indicating more

complex resilience and eco-memory dynamics than anticipated. It may result from initial

destabilisation away from memorised conditions being followed by a "relearning" process as

the eco-net adapts to the emergence of a new structure (reflected by higher biodiversity) and

conditions, leading to the latter decrease in E_n. We also simulate the impact of increasing

accumulation rates, compaction, or variable temporal resolution with a non-linear age-depth

model, as these effects are common in palaeorecords. This does not alter the overall signal,

but significantly reduces the resolution of deeper features (Supplementary Figure S1).

4.   Test-Case 2: Lake Erhai

Data from lake Erhai – a lake in south-western China which has undergone

eutrophication after decades of nutrient-enrichment – consists of relative diatom abundances

sampled at regular radioisotope-dated depth intervals down sediment cores [26,47]. We focus

on diatoms as they are well-preserved and all belong to the same trophic level, but this also

means significant parts of the wider ecosystem are not captured. However, we posit that

diatom λ_d indirectly reflects the wider ecosystem, as diatoms play a key role in the trophic

loops involved in eutrophication [28] and are ecologically sensitive to water quality [48,49].

As expected, real-world data exhibits more complexity than model results, but some

phases of activity can be observed (Fig. 2b). Phase 1 corresponds to elevated λ_d and E_n (along

with increasing SD), before both λ_d and E_n decline in phase 2 (along with elevated AR1 and

biodiversity, and negative compositional disorder) prior to the transition, after which E_n rises

sharply (while biodiversity sharply drops and compositional disorder recovers). These

patterns are similar to the PCLake+ results, with initially elevated λ_d and E_n suggesting a shift

away from the memorised state during early destabilisation, followed by a decline in λ_d and

E_n, suggesting the eco-net relearning a new state during pre-transition reorganisation.

5.   Further Development

There are several key methodological limitations that can be improved with future

development. To work best, the algorithm requires long datasets with minimal sparseness that

sufficiently resolve equilibrium population dynamics. Many palaeorecords do not meet these

requirements though, rendering them unsuitable for this analysis. Further development will

attempt to improve the algorithms capacity to analyse limited or sparse data using innovative

techniques from metagenomic network inference [32]. We also assume these ecosystems fit a

generalised Lotka-Volterra model, in which all of the changes in abundance are caused either

by interspecific interactions or by a broad noise term encompassing all other influences. But

interspecific interactions may not fit a simplified model of linear pairwise interaction, and

other important processes are simplified into the noise term [50–53]. Nonlinear interaction

terms are expected in some ecosystems [54] and potentially allow a better representation of

multiple alternative stable states [55], but are harder to parameterise. Future work will assess

the feasibility of allowing nonlinear functional responses.

The assumed model only includes biotic elements with life-environment interactions

implicit, but in reality the environment dynamically interacts with life. Incorporating life-

environment feedbacks explicitly would allow more realistic feedback loops critical to

resilience to emerge, but the eco-net cannot simply be extended to include abiotic elements as

the environment is currently assumed to be its training data. One solution might be using a

neural network that allows training data that is dynamic (e.g. continuous-time recurrent neural

networks) and affected by the eco-net itself (e.g. multi-layer or adversarial networks).

6.   Conclusions

In this paper we develop a novel algorithm to reconstruct eco-networks from palaeoecological abundance data and estimate their changing resilience. With further refinement, this method might allow the past resilience of ecosystems over time to be inferred from palaeoecological data. We also introduce eco-net energy as a proxy of "ecological memory", by treating the eco-network as a neural network that has undergone unsupervised Hebbian learning of past environmental forcing. This concept has only been applied to a simplified ecosystem model before now [39,40], but the development of eco-net energy has allowed us to explore eco-memory in more realistic models and empirical data for the first time. The concept of ecological memory opens a new dimension for understanding ecosystem resilience and regime shifts, with the formation of eco-memory potentially helping to improve ecosystem resilience by allowing past eco-network stable states to be recovered after disruptions. Further work is required to fully understand the drivers and implications of eco-memory dynamics, and to disentangle the effects of eco-memory from other drivers of changing ecosystem resilience.

Acknowledgements

This work was supported by an EPSRC/ReCoVER Pilot Study Project Award (RFFLP021). Erhai data was provided by Rong Wang and the Nanjing Institute of Geography and Limnology. We thank Pete Langdon for comments on preliminary results, and Annette Janssen for advice on using PCLake+.

Figures

Figure 1:   Diagram showing how ecosystem interactions (left) can be represented as an eco-network (middle), the structure of which forms an interaction matrix (right) suitable for resilience analysis. Only symmetric interactions are shown here for simplicity; asymmetric interactions are also likely.

Figure 2:   Results for PCLake+ (left) and lake Erhai (right) plotted against depth (time running left-to-right), showing output for (a) linear instability λ_d and (b) eco-net energy E_n. Also shown are (c) normalised time-series metrics (TSM; shown for comparison only), (d) biodiversity (inverse Simpson index), and (e, Erhai only) compositional disorder (see [26] for details). Vertical dotted lines (green: phase 1, orange: phase 2, red: transition) mark suggested transition phases.

References

1.   Scheffer M *et al.* 2009 Early-warning signals for critical transitions. *Nature* **461**, 53–9.
(doi:10.1038/nature08227)

2.   Dakos V, Carpenter SR, van Nes EH, Scheffer M. 2015 Resilience indicators: prospects
and limitations for early warnings of regime shifts. *Philos. Trans. R. Soc. B Biol. Sci.*
370, 20130263–20130263. (doi:10.1098/rstb.2013.0263)

3.   Lenton TM. 2013 Environmental Tipping Points. *Annu. Rev. Environ. Resour.* **38**, 1–29.
(doi:10.1146/annurev-environ-102511-084654)

4.   Kéfi S, Dakos V, Scheffer M, Van Nes EH, Rietkerk M. 2013 Early warning signals
also precede non-catastrophic transitions. *Oikos* **122**, 641–648. (doi:10.1111/j.1600-
0706.2012.20838.x)

5.   Scheffer M *et al.* 2012 Anticipating critical transitions. *Science* **338**, 344–8.
(doi:10.1126/science.1225244)

6.   Dakos V *et al.* 2012 Methods for Detecting Early Warnings of Critical Transitions in
Time Series Illustrated Using Simulated Ecological Data. *PLoS One* **7**, e41010.
(doi:10.1371/journal.pone.0041010)

7.   Lenton TM. 2011 Early warning of climate tipping points. *Nat. Clim. Chang.* **1**, 201–
209. (doi:10.1038/nclimate1143)

8.   Dakos V, Scheffer M, van Nes EH, Brovkin V, Petoukhov V, Held H. 2008 Slowing
down as an early warning signal for abrupt climate change. *Proc. Natl. Acad. Sci. U. S.*
A. **105**, 14308–12. (doi:10.1073/pnas.0802430105)

9.   Carpenter SR, Brock WA. 2006 Rising variance: a leading indicator of ecological
transition. *Ecol. Lett.* **9**, 311–318. (doi:10.1111/j.1461-0248.2005.00877.x)

10.   Lenton TM, Livina VN, Dakos V, Scheffer M. 2012 Climate bifurcation during the last
deglaciation? *Clim. Past* **8**, 1127–1139. (doi:10.5194/cp-8-1127-2012)

11.   Scheffer M, Van Nes EH. 2007 Shallow lakes theory revisited: Various alternative
regimes driven by climate, nutrients, depth and lake size. *Hydrobiologia* **584**, 455–466.
(doi:10.1007/s10750-007-0616-7)

12.   Scheffer M, Hosper S, Meijer M, Moss B, Jeppesen E. 1993 Alternative equilibria in
shallow lakes. *Trends Ecol. Evol.* **8**, 275–279.

13.   Scheffer M, Carpenter SR. 2003 Catastrophic regime shifts in ecosystems: Linking
theory to observation. *Trends Ecol. Evol.* **18**, 648–656. (doi:10.1016/j.tree.2003.09.002)

14.   Carpenter SR, Ludwig D, Brock WA. 1999 Management of Eutrophication for Lakes
Subject To Potentially Irreversible Change. *Ecol. Appl.* **9**, 751–771. (doi:10.1890/1051-
0761(1999)009[0751:MOEFLS]2.0.CO;2)

15.   Carpenter SR. 2005 Eutrophication of aquatic ecosystems: Bistability and soil
phosphorus. *Proc. Natl. Acad. Sci.* **102**, 10002–10005. (doi:10.1073/pnas.0503959102)

16.   Armstrong McKay DI, Lenton TM. 2018 Reduced carbon cycle resilience across the
268   Palaeocene–Eocene Thermal Maximum. *Clim. Past* **14**, 1515–1527. (doi:10.5194/cp-
269   14-1515-2018)

270   17.   Gsell AS *et al.* 2016 Evaluating early-warning indicators of critical transitions in
271   natural aquatic ecosystems. *Proc. Natl. Acad. Sci.* **113**, E8089–E8095.
272   (doi:10.1073/pnas.1608242113)

273   18.   Bestelmeyer BT *et al.* 2011 Analysis of abrupt transitions in ecological systems.
274   *Ecosphere* **2**, art129. (doi:10.1890/ES11-00216.1)

275   19.   Burthe SJ *et al.* 2016 Do early warning indicators consistently predict nonlinear change
276   in long-term ecological data? *J. Appl. Ecol.* **53**, 666–676. (doi:10.1111/1365-
277   2664.12519)

278   20.   Hastings A, Wysham DB. 2010 Regime shifts in ecological systems can occur with no
279   warning. *Ecol. Lett.* **13**, 464–472. (doi:10.1111/j.1461-0248.2010.01439.x)

280   21.   Litzow MA, Hunsicker ME. 2016 Early warning signals, nonlinearity, and signs of
281   hysteresis in real ecosystems. *Ecosphere* **7**. (doi:10.1002/ecs2.1614)

282   22.   Boettiger C, Ross N, Hastings A. 2013 Early warning signals: The charted and
283   uncharted territories. *Theor. Ecol.* **6**, 255–264. (doi:10.1007/s12080-013-0192-6)

284   23.   Boettiger C, Hastings A. 2012 Early warning signals and the prosecutor's fallacy. *Proc.
285   Biol. Sci.* **279**, 4734–9. (doi:10.1098/rspb.2012.2085)

286   24.   Carstensen J, Telford RJ, Birks HJB. 2013 Diatom flickering prior to regime shift.
287   *Nature* **498**, E11-2. (doi:10.1038/nature12272)

288   25.   Taranu ZE, Carpenter SR, Frossard V, Jenny J-P, Thomas Z, Vermaire JC, Perga M-E.
289   2018 Can we detect ecosystem critical transitions and signals of changing resilience
290   from paleo-ecological records? *Ecosphere* **9**, e02438. (doi:10.1002/ecs2.2438)

291   26.   Doncaster CP, Chávez VA, Viguier C, Wang R, Zhang E, Dong X, Dearing JA,
292   Langdon PG, Dyke JG. 2016 Early warning of critical transitions in biodiversity from
293   compositional disorder. *Ecology* **97**, n/a-n/a. (doi:10.1002/ecy.1558)

294   27.   Wang R *et al.* 2019 Network parameters quantify loss of assemblage structure in
295   human-impacted lake ecosystems. *Glob. Chang. Biol.* , 1–12. (doi:10.1111/gcb.14776)

296   28.   Kuiper JJ, van Altena C, de Ruiter PC, van Gerven LPA, Janse JH, Mooij WM. 2015
297   Food-web stability signals critical transitions in temperate shallow lakes. *Nat.
298   Commun.* **6**, 7727. (doi:10.1038/ncomms8727)

299   29.   Neutel A-M, Heesterbeek JAP, Ruiter PC de. 2002 Stability in Real Food Webs: Weak
300   Links in Long Loops. *Science (80-.).* **296**, 1120–1123. (doi:10.1126/science.1068326)

301   30.   Neutel A-M, Heesterbeek JAP, van de Koppel J, Hoenderboom G, Vos A, Kaldeway C,
302   Berendse F, de Ruiter PC. 2007 Reconciling complexity with stability in naturally
303   assembling food webs. *Nature* **449**, 599–602. (doi:10.1038/nature06154)

304   31.   Faust K, Raes J. 2012 Microbial interactions: From networks to models. *Nat. Rev.
305   Microbiol.* **10**, 538–550. (doi:10.1038/nrmicro2832)

32. Faust K, Lahti L, Gonze D, de Vos WM, Raes J. 2015 Metagenomics meets time series
analysis: Unraveling microbial community dynamics. *Curr. Opin. Microbiol.* **25**, 56–
66. (doi:10.1016/j.mib.2015.04.004)

33. Fisher CK, Mehta P. 2014 Identifying Keystone Species in the Human Gut Microbiome
from Metagenomic Timeseries Using Sparse Linear Regression. *PLoS One* **9**, 1–10.
(doi:10.1371/journal.pone.0102451)

34. Mounier J, Monnet C, Vallaeys T, Arditi R, Sarthou A-S, Helias A, Irlinger F. 2008
Microbial Interactions within a Cheese Microbial Community. *Appl. Environ.*
Microbiol. **74**, 172–181. (doi:10.1128/AEM.01338-07)

35. R Foundation for Statistical Computing. 2016 R: A language and environment for
statistical computing.

36. Aitchison J. 1982 The Statistical Analysis of Compositional Data. *J. R. Stat. Soc. Ser.*
B. Methodol. **44**, 139–177. (doi:10.2307/2345821)

37. de Ruiter PC, Neutel A-M, Moore JC. 1995 Energetics, Patterns of Interaction
Strengths, and Stability in Real Ecosystems. *Science (80-.).* **269**, 1257–1260.
(doi:10.1126/science.269.5228.1257)

38. Lade SJ, Gross T. 2012 Early Warning Signals for Critical Transitions: A Generalized
Modeling Approach. *PLoS Comput. Biol.* **8**, e1002360.
(doi:10.1371/journal.pcbi.1002360)

39. Power DA, Watson RA, Szathmary E, Mills R, Powers ST, Doncaster CP, Czapp B.
2015 What can ecosystems learn? Expanding evolutionary ecology with learning
theory. *Biol. Direct* **10**, 69. (doi:10.1186/s13062-015-0094-1)

40. Watson RA, Szathmáry E. 2016 How Can Evolution Learn? *Trends Ecol. Evol.* **31**,
147–157. (doi:10.1016/j.tree.2015.11.009)

41. Hopfield JJ. 1982 Neural networks and physical systems with emergent collective
computational abilities. *Proc. Natl. Acad. Sci.* **79**, 2554–2558.
(doi:10.1073/pnas.79.8.2554)

42. MacKay DJC. 2005 *Information Theory, Inference, and Learning Algorithms*.
Cambridge University Press. (doi:10.1198/jasa.2005.s54)

43. Rojas R. 1996 *Neural Networks: a Systematic Introduction*. (doi:10.1016/0893-
6080(94)90051-5)

44. Kröse B, Smagt P van der. 1996 *An Introduction To Neural Networks*.

45. Oksanen J *et al.* 2017 vegan: Community Ecology Package.

46. Janssen ABG, Teurlincx S, Beusen AHW, Huijbregts MAJ, Rost J, Schipper AM,
Seelen LMS, Mooij WM, Janse JH. 2019 PCLake+: A process-based ecological model
to assess the trophic state of stratified and non-stratified freshwater lakes worldwide.
Ecol. Modell. **396**, 23–32. (doi:10.1016/j.ecolmodel.2019.01.006)

47. Wang R, Dearing JA, Langdon PG, Zhang E, Yang X, Dakos V, Scheffer M. 2012
Flickering gives early warning signals of a critical transition to a eutrophic lake state.

Nature **492**, 419–22. (doi:10.1038/nature11655)

48.   Hall R, Smol J. 1999 Diatoms as indicators of lake eutrophication. In *The Diatoms:*
347   *Applications for the Environmental and Earth Sciences* (eds EF Stoermer, JP Smol), pp.
348   128–168. Cambridge, UK: Cambridge University Press.

349   49.   Dixit SS, Smol JP, Kingston JC, Charles DF. 1992 Diatoms: Powerful Indicators of
350   Environmental Change. *Environ. Sci. Technol.* **26**, 22–33. (doi:10.1021/es00025a002)

351   50.   Suzuki K, Yoshida K, Nakanishi Y, Fukuda S. 2017 An equation-free method reveals
352   the ecological interaction networks within complex microbial ecosystems. *Methods*
353   *Ecol. Evol.* **8**, 1774–1785. (doi:10.1111/2041-210X.12814)

354   51.   Xiao Y, Angulo MT, Friedman J, Waldor MK, Weiss ST, Liu Y-Y. 2017 Mapping the
355   ecological networks of microbial communities. *Nat. Commun.* **8**, 2042.
356   (doi:10.1038/s41467-017-02090-2)

357   52.   Levine JM, Bascompte J, Adler PB, Allesina S. 2017 Beyond pairwise mechanisms of
358   species coexistence in complex communities. *Nature* **546**, 56–64.
359   (doi:10.1038/nature22898)

360   53.   Cao H-T, Gibson TE, Bashan A, Liu Y-Y. 2016 Inferring human microbial dynamics
361   from temporal metagenomics data: Pitfalls and lessons. *BioEssays* **39**, 1600188.
362   (doi:10.1002/bies.201600188)

363   54.   Neutel A-M, Thorne MAS. 2016 Linking saturation, stability and sustainability in food
364   webs with observed equilibrium structure. *Theor. Ecol.* **9**, 73–81. (doi:10.1007/s12080-
365   015-0270-z)

366   55.   Gunderson LH. 2000 Ecological Resilience—In Theory and Application. *Annu. Rev.*
367   *Ecol. Syst.* **31**, 425–439. (doi:10.1146/annurev.ecolsys.31.1.425)