1 **A virus-encoded protein suppresses methylation of the viral genome in the Cajal body**

2 **through its interaction with AGO4**

3 Liping Wang[1,2], Yi Ding[1,2], Li He[1], Guiping Zhang[1], Jian-Kang Zhu[1], Rosa Lozano-Duran[1]

4 [1]Shanghai Center for Plant Stress Biology, Chinese Academy of Sciences, Shanghai 201602,

5 China; Center for Excellence in Molecular Plant Science, Chinese Academy of Sciences.

6 [2]University of the Chinese Academy of Sciences, Beijing 100049, China

8 ABSTRACT

In plants, establishment of *de novo* DNA methylation is regulated by the RNA-directed DNA

methylation (RdDM) pathway. RdDM machinery is known to concentrate in the Cajal body, but

the biological significance of this localization has remained elusive. Here, we show that the anti-

viral methylation of the *Tomato yellow leaf curl virus* (TYLCV) genome occurs specifically in the

Cajal body of *Nicotiana benthamiana* cells. Methylation of the viral genome is countered by a

virus-encoded protein, V2, which interacts with the central RdDM component AGO4, interfering

with its binding to the viral DNA; Cajal body localization of the V2-AGO4 interaction is required for

the viral protein to exert this function. Taken together, our results draw a long sought-after

functional connection between RdDM, the Cajal body, and anti-viral DNA methylation, paving the

way for a deeper understanding of DNA methylation and anti-viral defences in plants.

INTRODUCTION

DNA methylation in cytosine residues is a conserved epigenetic mark essential for protecting the eukaryotic genome against invading nucleic acids, namely viruses and transposable elements. In plants, establishment of *de novo* DNA methylation is believed to be regulated by the RNA-directed DNA methylation (RdDM) pathway. The canonical RdDM pathway requires the activity of two plant-specific RNA polymerase II-related enzymes, Pol IV and Pol V, and leads to cytosine methylation in a sequence-specific manner. In brief, the current understanding of RdDM is as follows: Pol IV generates RNA transcripts subsequently converted to double-stranded RNA (dsRNA) by RDR2 (Haag et al., 2012, Law et al., 2011), and then diced into 24-nt siRNAs by DCL3 (Xie et al., 2004); the resulting 24-nt siRNAs are loaded into AGO4 (Zilberman et al., 2003), which is guided to scaffold RNA molecules generated by Pol V via sequence complementarity and recruits the *de novo* methyl transferase DRM2 (Böhmdorfer et al., 2014, Chan et al., 2005, Gao et al., 2010, Zhong et al., 2014), which in turn catalyses methylation of adjacent DNA sequences. RdDM generally creates a chromatin environment refractive to gene expression. Part of the RdDM machinery, including AGO4, was found to concentrate in the Cajal body, a subnuclear compartment that is the site of maturation of ribonucleoprotein complexes (Li et al., 2006). This observation led to the suggestion that Cajal bodies might be a centre for the assembly of an AGO4/NRPE1/siRNA complex, which would facilitate RdDM at target loci (Li et al., 2006). However, the biological relevance of the Cajal body localization of RdDM components, and of this compartment on DNA methylation, has so far not been demonstrated.

Geminiviruses are a family of plant viruses with circular single-stranded (ss) DNA genomes infecting multiple crops and causing dramatic yield losses worldwide. The geminiviral genome replicates in the nucleus of the infected cell by using the host DNA replication machinery, made available following the viral re-programming of the cell cycle (reviewed in Hanley-Bowdoin et al., 2013). During viral multiplication, the ssDNA genome generates a double-stranded (ds) DNA intermediate, which then undergoes rolling-circle replication (reviewed in Hanley-Bowdoin et al., 2013). Nevertheless, the cellular and molecular details underlying these essential initial steps of the viral infection cycle, including the subnuclear localization of viral ss and ds DNA accumulation and of the processes leading to their production, are to date mostly unknown.

Notably, the geminiviral genome forms minichromosomes and is subjected to epigenetic modifications, including cytosine DNA methylation and histone modifications (Ceniceros-Ojeda et al., 2016, Deuschle et al., 2016, Jackel et al., 2016, Kushwaha et al., 2017, Raja et al., 2008, Wang et al., 2018). The findings that methylation of viral DNA negatively impacts viral replication

(Brough et al., 1992, Ermak et al., 1993), that different geminivirus-encoded proteins have evolved
to suppress DNA methylation (Buchmann et al., 2009, Ismayil et al., 2018, Raja et al., 2008,
Rodriguez-Negrete et al., 2013, Tu et al., 2017, Wang et al., 2018, Yang et al., 2013, Yang et al.,
2011, Zhang et al., 2011), and that methylation of the viral genome correlates with host resistance
or recovery (Butterbach et al., 2014, Ceniceros-Ojeda et al., 2016, Raja et al., 2008, Torchetti et
al., 2016, Yadav & Chattopadhyay, 2011) strongly support the idea that the plant-mediated
methylation of the viral DNA acts as an anti-viral defence mechanism, underscoring DNA
methylation as an active battlefield in the interaction between plants and geminiviruses.

During the infection by the geminivirus *Tomato yellow leaf curl virus* (TYLCV), the essential virus-
encoded protein V2 has been shown to suppress DNA methylation of the viral genome (Wang et
al., 2014). V2 interacts with the histone deacetylase HDA6 in *Nicotiana benthamiana*, competing
with the recruitment of the maintenance CG methyltransferase MET1 (Woo et al., 2008) and
ultimately reducing viral DNA methylation (Wang et al., 2018). Nevertheless, silencing of *HDA6*
results in limited complementation of a V2 null mutation in the virus and in only a partial reduction
in viral DNA methylation (Wang et al., 2018), suggesting that V2 might counter methylation
through additional interactions with host factors.

In this work, we show that V2 from TYLCV interacts with the central RdDM component AGO4;
that AGO4 plays a role in the defence against TYLCV; and that V2 interferes with the AGO4
binding to the viral DNA. Importantly, our results indicate that the viral DNA gets methylated in
the Cajal body in the absence of V2, since abolishment of Cajal body formation by silencing of its
signature component coilin drastically reduces methylation of the viral genome. This idea is further
supported by the finding that the activity of V2 as a suppressor of viral DNA methylation requires
Cajal body localization of the V2-AGO4 interaction. In summary, our results draw a functional
connection between anti-viral RdDM and the Cajal body in plants, and illustrate how TYLCV has
evolved a protein, V2, to counter this defence mechanism.

RESULTS

V2 from TYLCV interacts with AGO4 from *Nicotiana benthamiana* and tomato

With the aim of gaining insight into the functions of V2 from TYLCV in the plant cell, we used transient expression of GFP-tagged V2 in infected leaf patches of *N. benthamiana* followed by affinity purification and mass spectrometry (AP-MS) to identify plant interactors of this viral protein in the context of the infection (Wang et al., 2017a). Interestingly, we identified the two AGO4 paralogs in *N. benthamiana* (NbAGO4-1 and NbAGO4-2) as associated to V2-GFP (Figure 1A; Wang et al., 2017a); these interactions were confirmed by co-immunoprecipitation (co-IP) and split-luciferase assays (Figure 1B,C).

There are four *AGO4* orthologues in tomato (*SlAGO4a-d*) (Bai et al., 2012), the natural and economically relevant host of TYLCV (Figure 2A,B). All four SlAGO4-encoding genes are expressed in basal conditions in tomato leaves, although *SlAGO4c* and *d* show low expression levels; *SlAGO4b*, *c*, and *d* are slightly up-regulated by TYLCV infection (Supplementary figure 1). *SlAGO4a*, *SlAGO4b*, and *SlAGO4d* were cloned and the encoded proteins confirmed as interactors of V2 in co-IP and split-luciferase assays (Figure 2C,D). Our results therefore show that V2 interacts with AGO4 in two host species, tomato and *N. benthamiana*.

V2 counters the AGO4-dependent methylation of the viral genome to promote virulence

In order to evaluate the contribution of V2 to the viral infection, we generated an infectious TYLCV clone carrying a G-to-A mutation in the fifth nucleotide of the V2 open reading frame (ORF), which converts the second codon (encoding tryptophan) to a stop codon (Supplementary figure 2A), making it unable to produce the V2 protein (TYLCV-V2null). In agreement with previous results (Wartig et al., 1997), V2 is required for full infectivity in both tomato and *N. benthamiana*, since the V2 null mutant accumulates to very low levels and produces no noticeable symptoms (Supplementary figure 2).

Next, we sought out to test whether knock-down of *AGO4* could partially complement the lack of V2 during the TYLCV infection. For this purpose, we employed virus-induced gene silencing (VIGS) to silence both *NbAGO4-1* and *NbAGO4-2*. VIGS efficiently knocked-down both *NbAGO4* orthologues, but did not affect accumulation of the transcript of the close homologue *NbAGO6* (Figure 3A); AGO4-silenced plants did not display any obvious developmental abnormalities (Figure 3B). Expression of *NbAGO4-1* or *NbAGO4-2* was not affected by TYLCV infection, neither

in silenced nor in non-silenced plants (Figure 3C,D). Mutation in V2 does not affect viral replication (Wartig et al., 1997), and therefore viral accumulation in local infections in *N. benthamiana* (leaf patch agroinfiltration assays; see Supplementary figure 3) was not different between the WT virus and the V2 null mutant virus (Figure 3E); in both cases, *AGO4* silencing led to a slight increase in viral accumulation, suggesting an anti-viral role for AGO4 (Figure 3E). Interestingly, in systemic infections (see Supplementary figure 3), *AGO4* silencing mildly increased viral accumulation of the WT TYLCV (1.33-fold), but dramatically improved performance of the V2 null mutant virus (3.23-fold), suggesting that one of the main roles of V2 during the viral infection is the suppression of AGO4 function (Figure 3F).

In light of the role of AGO4 in RdDM and to directly assess the impact of V2 and AGO4 on the methyl-state of the viral DNA, we used bisulfite sequencing (BS-seq) to measure DNA methylation of the intergenic region (IR) of the viral genome, which presents the highest methylation levels during the infection (Piedra-Aguilera et al., 2019). As shown in Figure 4A, cytosine methylation in this region in all contexts (CG, CHG, and CHH) was almost undetectable in the WT viral genome in local infections at 3 or 9 days post-inoculation (dpi), while it reached ~60% and ~80%, respectively, in the V2 null mutant (Figure 4A; Supplementary figure 4A). These results indicate that V2 can prevent or revert methylation of the viral genome during the infection, which occurs rapidly in the absence of this protein.

We then compared the percentage of cytosine methylation in the intergenic region in local infections with the V2 null mutant TYLCV in basal conditions or upon *AGO4* silencing. Strikingly, *AGO4* silencing resulted in a ~30-55% decrease in the percentage of methylated cytosines in all contexts (Figure 4B; Supplementary figure 4B), indicating that knock-down of *AGO4* can partially complement the lack of V2 at the level of the viral methylation state. This complementation suggests that a) methylation of the viral DNA at least partially depends on AGO4 function; and b) V2 can at least partially counter AGO4-dependent methylation of the viral DNA.

As opposed to short-timed local infections, in systemic infections in *N. benthamiana*, which require longer timespans and involve viral cell-to-cell and long-distance movement, methylation of the WT viral genome could be detected at ~8-23% in all contexts (Figure 4C; Supplementary figure 5). Interestingly, the methylation level tends to decrease upon *AGO4* silencing; this reduction (~40-60%) is more prominent in the V2 null mutant genome, again supporting the idea that AGO4-dependent methylation of the viral genome occurs during the infection and is partially countered by V2. Notably, the detected decrease in methylation correlates with the enhanced viral accumulation in the *AGO4*-silenced plants (Figure 3F).

V2 does not hamper production or loading of vsiRNA but interferes with AGO4 binding to the viral genome

The canonical function of AGO4 in the RdDM pathway requires loading of siRNA and association to Pol V, and results in the recruitment of DRM2 to the target loci and the subsequent methylation of the adjacent DNA (Matzke et al., 2015, Matzke & Mosher, 2014). Through physical interaction, V2 could affect AGO4 function on the viral genome in different ways, for example by impairing loading of viral siRNA (vsiRNA) onto this protein or by displacing endogenous interactors, such as Pol V or DRM2. In order to shed light on the molecular mechanism underlying the V2-mediated interference of AGO4-dependent methylation of the viral genome, we tested binding of AGO4 to the viral DNA in the presence or absence of V2 in local infections with TYLCV WT and V2 null mutant, respectively, by Chromatin immunoprecipitation (ChIP). As shown in Figure 5A, 3xFLAG-NbAGO4 could bind both the IR and the V2-encoding region of the viral genome in the absence of V2 (TYLCV-V2null), but the signal decreased to background levels when V2 was present (TYLCV). Therefore, AGO4 has the capacity to bind the viral DNA molecule, but this binding is impaired by the virus-encoded V2 protein. AGO4 binding in the TYLCV V2 null mutant hence correlates with the detected increase in viral DNA methylation (Figure 4A).

Several viral silencing suppressors encoded by different viruses have been shown to inhibit formation of AGO/sRNA complexes (e.g. Burgyán et al., 2011, Rawlings et al., 2011, Schott et al., 2012). To test whether this strategy is also employed by V2, we immunoprecipitated 3xFLAG-NbAGO4 co-expressed with WT or V2 null mutant TYLCV in local infection assays in *N. benthamiana*, and visualized AGO4-bound vsiRNA by sRNA northern blotting. While infected samples contained both 21- and 24-nt vsiRNA, and the occurrence and accumulation of these sRNA species was not affected by the presence of virus-encoded V2, mostly 24-nt vsiRNA co-immunoprecipitated with AGO4 (Figure 5B). Interestingly, a higher amount of vsiRNA associated to AGO4 in the samples infected with the WT virus (Figure 5B). Taken together, these results demonstrate that V2 does not affect the production or accumulation of vsiRNA, nor does it hamper loading of these vsiRNA molecules into AGO4, but interferes with binding of this protein to the viral genome in order to suppress DNA methylation and promote virulence.

Methylation of the viral genome and its suppression by V2 occur in the Cajal body

In Arabidopsis, AGO4 has been shown to co-localize with its interactor NRPE1 (NRPD1b), a subunit of Pol V, in the Cajal body, which was then suggested to be a center for the assembly of AGO4/NRPE1/siRNA complexes, enabling RdDM at target loci (Li et al., 2008, Li et al., 2006). However, the functional significance of this subnuclear localization has so far remained elusive. Interestingly, both V2-GFP and the different RFP-AGO4 orthologues from *N. benthamiana* and tomato co-localize in a distinct subnuclear compartment, identified as the Cajal body by the accumulation of the nucleolus and Cajal body marker fibrillarin (Barneche et al., 2000), upon transient expression in *N. benthamiana* (Figure 6A; Supplementary video 1). Of note, most of nuclear V2-GFP accumulates in the Cajal body, although some fluorescence can be detected in the nucleoplasm. All AGO4 orthologues are distributed throughout the nucleoplasm and absent from the nucleolus; clear Cajal body localization can be detected for NbAGO4-1, NbAGO4-2, SlAGO4a, and SlAGO4b, while Cajal body localization of SlAGO4d is less conspicuous (Figure 6A). Analysis of the V2/AGO4 interaction by bimolecular fluorescence complementation (BiFC), which is based on visualization and hence provides spatial information, unveiled that, strikingly, the association between these two proteins occurs mostly or exclusively in the Cajal body, where V2 homotypic interactions also occur (Figure 6B).

In order to evaluate the relevance of the Cajal body localization of the V2-AGO4 interaction, we took advantage of the fact that a GFP-V2 fusion protein, as opposed to the previously mentioned V2-GFP, does not localize to the Cajal body (Figure 7A; Supplementary video 2), otherwise showing an indistinguishable subcellular distribution pattern. Importantly, GFP-V2 still interacts with AGO4 in co-IP assays (Supplementary figure 6A). Nevertheless, only the Cajal body-localized V2-GFP, but not GFP-V2, can complement a null mutation in V2 in terms of suppression of methylation of the viral genome (Figure 7B; Supplementary figure 6B), indicating that the Cajal body localization of the V2-AGO4 interaction is essential for V2 to exert its function.

Since our results indicate that the suppression of the AGO4-mediated methylation of the viral genome occurs in the Cajal body, we decided to test if the Cajal body is required for this anti-viral response. For this purpose, we eliminated Cajal bodies in the plant by silencing of their signature component coilin through VIGS; the effect of knocking down *coilin* on the formation of Cajal bodies has been previously demonstrated (Shaw et al., 2014). As expected, silencing *coilin* (Figure 7C) resulted in the disappearance of Cajal bodies (Supplementary figure 6C). *Coilin*-silenced plants were then inoculated with TYLCV WT or V2 null mutant, and viral accumulation and methylation of the intergenic region of the viral genome were evaluated three weeks post-inoculation

(Supplementary figure 6D, E; Figure 7D). A tendency towards higher viral accumulation in *coilin*-
silenced plants was observed, but not statistically significant (Supplementary figure 6E). Strikingly,
however, methylation of the IR of the viral genome in the V2 mutant virus was largely decreased
by silencing of *coilin* (Figure 7D; Supplementary figure 6F), suggesting that efficient methylation
of the viral genome requires intact Cajal bodies.

DISCUSSION

The plant DNA viruses geminiviruses and pararetroviruses are both targets and suppressors of
DNA methylation; this possibly extends to the third family of plant DNA viruses, nanoviruses,
although experimental evidence is lacking (reviewed in Poogin, 2013; Pumplin and Voinnet, 2013).
The independent evolution of viral suppressors of DNA methylation argues for an anti-viral effect
of this epigenetic modification. Indeed, seminal experiments by Brough et al. (1992) and Ermak
et al. (1993) demonstrated that methylation of the geminivirus genome interferes with its
replication in transformed protoplasts, likely due to a dual effect on viral gene expression and
function of the replication complex. More specifically, RdDM seems to play a prominent role in
plant defence against geminiviruses, since RdDM mutants or silenced plants display increased
susceptibility to geminivirus infection (Raja et al., 2008, Zhong et al., 2017), and DNA methylation
and repressive histone marks typical of RdDM are deposited on the viral genome (Castillo-
Gonzalez et al., 2015, Ceniceros-Ojeda et al., 2016, Coursey et al., 2018, Dogar et al., 2006,
Jackel et al., 2016, Kushwaha et al., 2017, Wang et al., 2018). However, the subnuclear
distribution of these anti-viral methylation events is currently unknown.

AGO4 is a central component of the canonical RdDM pathway, and as such an obvious target for
viral inhibition. However, AGO4 also affects susceptibility to RNA viruses and viroids, and is
targeted by proteins encoded by RNA viruses, which raises the idea that either RdDM on the host
genome plays a role in modulating plant-virus interactions broadly, or AGO4 has an anti-viral role
beyond RdDM (Brosseau et al., 2016, Ma et al., 2015, Minoia et al., 2014). Supporting the latter,
AGO4-dependent defences against a potexvirus are independent of other RdDM components
and do not require nuclear localization of AGO4 (Brosseau et al., 2016).

The geminivirus TYLCV encodes the essential, multifunctional V2 protein, which acts as a
suppressor of viral DNA methylation (Wang et al., 2014). Here, we show that V2 binds to the plant
AGO4 in the Cajal body, and suppresses the AGO4-dependent methylation of the viral genome,
which requires its Cajal body localization (Figures 1, 2, 4-6), and the AGO4-mediated restriction

of viral accumulation (Figure 3E,F). V2 impairs binding of AGO4 to the viral genome; we
hypothesize that the viral protein might mask a surface required for AGO4 recruitment to the viral
DNA through the association with an endogenous interactor (e.g. NRPE1; Li et al., 2006), or
interfere with the complementarity-based pairing to the nascent Pol V transcript. Our results
indicate that AGO4-dependent methylation of viral DNA occurs quickly in the absence of V2
(Figure 4A,B). Nevertheless, *AGO4* silencing still has a detectable, if minor, positive impact on
the accumulation of the WT virus, which correlates with decreased viral DNA methylation (Figure
3E,F; Figure 4C), suggesting that the V2-mediated suppression of AGO4 function is not complete.
On the other hand, WT levels of viral DNA methylation are not restored in the V2 mutant upon
AGO4 silencing, which raises the idea that AGO4 might not be the only methylation-related target
of V2. In agreement with this, V2 has been shown to bind HDA6 and interfere with its promotion
of MET1-dependent methylation of the viral DNA (Wang et al., 2018).

Recently, V2 encoded by the geminivirus *Cotton leaf curl Multan virus* (CLCuMV) was found to
interact with AGO4 in *N. benthamiana*, leading to enhanced viral accumulation and a reduction in
viral DNA methylation (Wang et al., 2019). It should be noted that the V2 proteins encoded by
CLCuMV and TYLCV are only 65% identical (Supplementary figure 7A); moreover, mutation of a
conserved residue, L76, abolishes the interaction between CLCuMV V2 and NbAGO4 (Wang et
al., 2019), but does not affect the interaction between TYLCV V2 and NbAGO4 or SlAGO4, which
still occurs in the Cajal body (Supplementary figure 7). This mutation, however, negatively affects
V2 self-interaction in the Cajal body (Supplementary figure 7). These results suggest that different
geminivirus species might have convergently evolved to target AGO4, underscoring the relevance
of this host factor for the viral infection.

The finding that the physical association between TYLCV V2 and AGO4 takes places in a specific
nuclear body, the Cajal body, and has an impact on the methyl-state of the viral population in the
cell, suggests that all or most viral DNA molecules must localize in this subnuclear compartment
at some point of the viral cycle. Supporting this notion, the activity of V2 as a suppressor of
methylation of the viral genome requires the Cajal body localization of its interaction with AGO4
(Figure 7A, B). This observation hints at a functional role of the Cajal body during the infection;
whether such a role is linked to gene expression, DNA replication, or some other process remains
to be investigated. Importantly, our results indicate that intact Cajal bodies are required for the
efficient methylation of the viral genome (Figure 7C, D; Supplementary figure 6), strongly
suggesting that anti-viral DNA methylation indeed occurs in this subnuclear compartment. In
agreement with this idea, depletion of Cajal bodies through knock-down of *coilin* was previously

found to promote the infection by the geminivirus *Tomato golden mosaic virus* (Shaw et al., 2014).

The Cajal body has also been connected to systemic infection of plant RNA viruses, and proteins

encoded by RNA viruses can bind coilin, which impacts plant-virus interactions (Kim et al., 2007a,

Kim et al., 2007b, Semashko et al., 2012, Shaw et al., 2014), although these effects are likely

independent on DNA methylation.

Based on our results, we propose a scenario in which anti-viral methylation of invading

geminivirus DNA in plants takes place in the Cajal body. In the context of the arms race between

host and virus, TYLCV has evolved V2, which is required during the infection to interfere with

AGO4 binding to the viral genome in the Cajal body, suppressing methylation of the viral DNA

and promoting virulence (Supplementary figure 8). In summary, taken together, our findings draw

a long sought-after functional connection between RdDM, the Cajal body, and anti-viral DNA

methylation, paving the way to a deeper understanding of DNA methylation and anti-viral defence

strategies in plants.

MATERIALS AND METHODS

Plasmids and cloning

To generate binary vectors to express *AGO4* from *N. benthamiana* and tomato (cv. Money maker),

the full-length coding sequence of *AGO4* genes was amplified using cDNA as template. *NbAGO4-*

1, *NbAGO4-2*, *SlAGO4a*, *SlAGO4b*, *SlAGO4d,* and *SlWRKY75* were cloned into pENTR-D/TOPO

(Invitrogen) following the manufacturer's instructions. The binary plasmids to express RFP and

3xHA N-terminal fusions were generated by Gateway cloning the AGO4 coding sequence into

pGWB555 and pGWB515, respectively (Nakagawa et al., 2007). V2-GFP, GFP-V2 and the

TYLCV infectious clone have been previously described (Rosas-Diaz et al., 2018, Wang et al.,

2017a). The V2 null TYLCV mutant was generated with the QuickChange Lightning Site-Directed

Mutagenesis Kit (Agilent Technologies, Cat #210518) using the WT infectious clone as template.

For the plasmids used for biomolecular fluorescence complementation (BiFC), *NbAGO4, SlAGO4,*

V2/V2$_{L76S}$ were cloned into entry vector pDONR221-P1P4/pDONR221-P3P2 (Invitrogen) and

then Gateway-cloned into the pBiFC-2in1-CN vector (Grefen & Blatt, 2012) as shown in

Supplementary table 2. The binary plasmids for split-luciferase complementation imaging assay

were generated by Gateway cloning the *NbAGO4, SlAGO4, SlWRKY75* or V2 from pENTR-

D/TOPO entry vector into pGWB-N-luc and pGWB-C-luc (Yu et al., 2019). To generate

pCAMBIA1300-3xFLAG-NbAGO4-1, pCAMBIA1300 was digested with *Xba*I and *Xma*I, and then

the 3xFLAG and NbAGO4-1 coding sequences were amplified by PCR and Infusion-cloned into

pCAMBIA1300 with ClonExpress® MultiS One Step Cloning Kit (Vazyme). pDONR207-*Fibrillarin*

(Kim et al., 2007b) was used to Gateway-clone *Fibrillarin* to pGWB545 (Nakagawa et al., 2007).

All primers and plasmids used for cloning are summarized in Supplementary tables 1 and 2,

respectively.

Plant materials and growth conditions

311 *N. benthamiana* and tomato plants (cv. Money maker) were grown in a controlled growth chamber

312 under long day conditions (LD, 16 h light/8 h dark) at 25℃.

Agrobacterium-mediated transient gene expression in *N. benthamiana*

All binary plasmids were transformed into *Agrobacterium tumefaciens* strain GV3101, with the

exception of pBINTRA6, which was transformed into *A. tumefaciens* strain C58c1. *A. tumefaciens*

clones carrying the constructs of interest were liquid-cultured in LB with appropriate antibiotics at

28℃ overnight. Bacterial cultures were then centrifuged at 4,000 g for 10 min and resuspended

in the infiltration buffer (10 mM $MgCl_2$, 10 mM MES pH 5.6, 150 µM acetosyringone) and adjusted

to an OD_{600} = 0.5. Next, the bacterial suspensions were incubated in the buffer at room

temperature and in the dark for 2-4 hours and then infiltrated 3-4-week-old *N. benthamiana* plants.

For co-expression experiments, the different agrobacterium suspensions were mixed at 1:1 ratio

before infiltration.

Protein extraction and immunoprecipitation assays

Protein extraction and co-immunoprecipitation assays were performed as described in Wang et

al., 2017a. Protein extracts were immunoprecipitated with GFP-Trap beads (Chromotek,

Germany), and analyzed by western blot with anti-GFP (Abiocode, M0802-3a) and anti-HA

(12CA5) (Roche, Cat. No. 11 583 816 001) antibodies.

For 3xFLAG-NbAGO4 IP followed by vsiRNA extraction, 6 grams of *N. benthamiana* leaves

transiently expressing 3xFLAG-NbAGO4 were collected, ground in liquid nitrogen and

homogenized in 6x (w:v) extraction buffer (20mM Tris HCl pH7.5, 25mM $MgCl_2$, 300mM NaCl,

5mM DTT, 0.5% NP-40, 1x complete™ Protease Inhibitor Cocktail (Roche)) at 4°C with rotation

for 30 minutes. The extract was subjected to centrifugation (14,000 rpm, 25min) at 4°C. 5 µg anti-

FLAG antibody (Sigma, F3165) per gram of tissue were added to the supernatant in a new tube

and incubated at 4°C overnight. The next day, 20 µl of slurry Protein G beads (Invitrogen) per

gram of tissue were added and subjected to a further incubation for 2 hours with rotation at 4°C.

After incubation, Protein G beads were washed three times in 3x (v:v) homogenate wash buffer (20mM Tris pH7.5, 25mM MgCl$_2$, 300mM NaCl, 0.5% NP-40). The quality of purification was examined by SDS-PAGE followed by immunoblotting.

Split-luciferase complementation imaging assay

Split-luciferase complementation imaging assays were performed as described (Chen et al., 2008). Equal volumes of *A. tumefaciens* harboring V2-N-luc or C-luc-NbAGO4-1/2, C-luc-SlAGO4a/b, or C-luc-SlWRKY75 at OD$_{600}$=0.5 were mixed at 1:1 ratio. Three different combinations of *A. tumefaciens* were infiltrated on the same *N. benthamiana* leaf. 1mM luciferin (in H$_2$O) was infiltrated into the inoculated leaves 2 days after agrobacterium infiltration. A low-light cooled CCD imaging apparatus (NightShade LB985 with IndiGO software) was used to capture and analyse the luciferase signal at 2 dpi.

Confocal imaging

Confocal imaging for co-localization of V2-GFP, RFP-AGO4, and CFP-Fibrillarin upon transient expression in *N. benthamiana* epidermal cells was performed on a Leica TCS SP8 point scanning confocal microscope using the pre-set sequential scan settings for GFP (Ex:488 nm, Em:500–550 nm), RFP (Ex:561 nm, Em:600–650 nm), and CFP (Ex:442 nm, Em:452–482 nm). Confocal imaging for co-localization of Fibrillarin-RFP (Kim et al., 2007b) and V2-GFP or GFP-V2 were performed in the same way.

Bimolecular Fluorescence Complementation

For bimolecular fluorescence complementation (BiFC) analyses, *A. tumefaciens* clones carrying pBiFC-2in1-CN binary constructs (Grefen and Blat, 2012) and CFP-Fibrillarin were mixed at 1:1 ratio and infiltrated on 3-4-week-old *N. benthamiana* plants. Imaging was performed 2 days later under a Leica TCS SP8 confocal microscope by using the pre-set sequential scan settings for YFP (Ex: 514 nm, Em: 525–575 nm) and for CFP (Ex:442 nm, Em:452–500 nm).

Virus-induced gene silencing

The vectors used for virus-induced gene silencing (VIGS) were pBINTRA6 (Ratcliff et al., 2001) and pTRV2-GW (Taylor et al., 2012). A 362-bp fragment of *NbAGO4-1* cDNA (from nt 1920 to 2281) and a 389-bp fragment of *Nbcoilin* cDNA (from nt 2222 to 2610) was amplified using specific primers shown in Supplementary table 1, cloned into pENTR/D-TOPO (Invitrogen), and subcloned into pTRV2-GW through an LR reaction (Invitrogen) to yield TRV-*NbAGO4* and TRV-*Nbcoilin*. VIGS assays were performed as described in Lozano-Duran et al., 2011. For TYLCV

local infection assays, *A. tumefaciens* carrying pBINTRA6 and TRV-*NbAGO4* or TRV-EV were

mixed and inoculated into 18-day-old *N. benthamiana* plants. Two weeks later, fully expanded

young leaves were infiltrated with *A. tumefaciens* carrying the TYLCV infectious clone and

samples were collected at 4 days post-inoculation (dpi) to detect viral accumulation. For TYLCV

systemic infection assays, *A. tumefaciens* carrying pBINTRA6 and TRV-*NbAGO4* or TRV-EV

(empty vector) and the TYLCV infectious clone were mixed and inoculated on 18-day-old *N.*

benthamiana plants. The three most apical leaves of each plant were collected at 3 weeks post-

inoculation (wpi) to detect viral accumulation. TYLCV systemic infection assays in *Nbcoilin-*

silenced plants were performed as previously described for *NbAGO4*-silenced plants.

Quantitative PCR (qPCR) and Reverse Transcription PCR (RT-qPCR)

To determine viral accumulation, total DNA was extracted from *N. benthamiana* leaves (from

infiltrated leaves in local infection assays and from apical leaves in systemic infection assays)

using the CTAB method (Minas et al., 2011). Quantitative PCR (qPCR) was performed with

primers to amplify Rep (Wang et al., 2017b). As internal reference for DNA detection, the 25S

ribosomal DNA interspacer (ITS) was used (Mason et al., 2008). To detect *NbAGO4-1*, *NbAGO4-*

2, *Nbcoilin* and *NbAGO6* transcripts, total RNA was extracted from *N. benthamiana* leaves by

using Plant RNA kit (OMEGA Bio-tek # R6827). RNA was reverse-transcribed into cDNA by using

the iScriptTM cDNA Synthesis Kit (Bio-Rad #1708890) according to the manufacturer's

instructions. *NbTubulin* was used as reference gene (Liu et al., 2012). Relative expression was

calculated by the comparative Ct method (2-ΔΔCt). qPCR and RT-qPCR were performed in a

BioRad CFX96 real-time system as described previously (Wang et al., 2017b). Total RNA was

extracted from the leaves of tomato plants mock-inoculated or infected with TYLCV at 3 weeks

post-inoculation (wpi). *SlActin was used as* reference gene (Exposito-Rodriguez et al., 2008)*.*

Similarly, RT-qPCR was performed on RNA extracted from tomato to detect the expression of

SlAGO4a/d/c/d. All primers used for qPCR and qRT-PCR are listed in Supplementary Table 3.

DNA bisulfite sequencing analysis

DNA from virus-infected plant tissues was extracted by DNeasy Plant Mini Kit (QIAGEN, Cat. No.

69104), and 500 ng of purified DNA was subjected to bisulfite treatment using EpiTect Plus DNA

Bisulfite Kit (QIAGEN, Cat. No. 59124) according to the manufacturer's handbook. The selected

fragment (viral IR) of the bisulfite-treated DNA was amplified by PCR (Fw:

TTTGATGTATTTTTTATTTGTTGGGGTTT, Rv: CCCTTACAACARATATAARATCCCT);

amplified fragments were cloned into the pMD18-T vector by TA ligation and sequenced (>15

clones per experiment). Cytosine methylation analysis was performed with Kismeth

(http://katahdin.mssm.edu/kismeth/revpage.pl) (Gruntman et al., 2008).

Chromatin immunoprecipitation (ChIP) assay

The agrobacterium clone carrying the binary vector to express 3xFLAG-NbAGO4-1 was co-
infiltrated with those carrying the TYLCV or TYLCV-V2null infectious clones in *N. benthamiana*
leaves, and tissues were collected at 2 dpi. Chromatin immunoprecipitation (ChIP) assays were
performed as described (He et al., 2018). In brief, the cross-linking of 2 grams of leaves was
performed with 1% formaldehyde in 1xPBS buffer and stopped with 1/15 volume of 2 M glycine
by vacuum infiltration. Then the tissue was ground to powder and resuspended in HB buffer (2.5%
Ficoll 400, 5% Dextran T40, 0.4 M Sucrose, 25 mM Tris pH 7.4, 10 mM MgCl2, 0.035% β-
mercaptoethanol, 1% Protease Inhibitor Cocktail (Sigma)), homogenized and filtered through
Miracloth (Milli-pore). Triton x-100 was added to the supernatant until final concentration was
0.5%. After spinning at 2,000×g for 20 min at 4°C, the pellet was re-suspended in HB buffer
containing 0.1% Triton x-100 and spun at 2,000×g for 10 min at 4°C. Isolated nuclei were re-
suspended in 500 µl of Nuclei Lysis buffer and sonicated by BioruptorTM UCD-200 sonicator
(diagenode) for 30 min. Following centrifugation at 21,130×g for 5 min at 4°C, the supernatant
was separated and used for input and immunoprecipitation. After adding 9 volume of ChIP dilution
buffer to the supernatant, this was pre-cleared with 10 µl of Dynabeads Protein G (Invirogen) for
1 h at 4°C. After removing the beads from the mixture, the supernatant was incubated with anti-
FLAG antibody (Sigma, F3165), or anti-IgG antibody (Sigma, I5006) overnight at 4°C. The
following day, after adding 20 µl of Dynabeads Protein G, the mixture was incubated for 2 h at
4°C. Beads were sequentially washed with 1 ml of the following buffers: Low Salt Wash buffer
(150 mM NaCl, 0.1% SDS, 1% Triton x-100, 2 mM EDTA, 20 mM Tris pH 8.0), High Salt Wash
buffer (500 mM NaCl, 0.1% SDS, 1% Triton x-100, 2 mM EDTA, 20 mM Tris pH 8.0), LiCl wash
buffer (250 mM LiCl, 1% Igepal, 1% Sodium Deoxycholate, 1 mM EDTA, 10 mM Tris pH 8.0), TE
buffer (10 mM Tris pH 8.0, 1 mM EDTA). Immunocomplexes were eluted with 250 µl of Elution
buffer (1% SDS, 0.1 M NaHCO$_3$) at 65°C for 15 min. After reverse crosslinking, 10 µl of 0.5 M
EDTA, 20 µl of 1 M Tris pH 6.5 and 1 µl of proteinase K (Invitrogen) were added to each sample,
which was incubated at 45 °C for 2 h. DNA was then purified using QIAquick PCR Purification Kit
(QIAGEN, Cat. No. 28106). The products were eluted into 200 µl of ddH$_2$O, and analysed by
qPCR. The primers used in this experiment are listed in Supplementary Table 3; the primers for
Actin are taken from Maimbo et al., 2010.

Small RNA (sRNA) extraction and northern blot analysis

Small RNA (sRNA) extraction and northern blot was performed as described (Yang et al., 2015). Briefly, sRNAs were purified from total extracts or AGO4 immunoprecipitates and subjected to northern blot analysis. For each sample, sRNAs were separated on a 17% polyacrylamide gel, which was electrotransferred to a Hybond N+ membrane (GE Lifesciences). Membranes were cross-linked, incubated for 2 hours at 65°C, and hybridized overnight at 38°C with ^{32}P-labeled probes for the intergenic region (IR) of the viral genome amplified by PCR (Fw: TCCTCTTTAGAGAGAGAACAATTGGGA, Rv: ACAACGAAATCCGTGAACAG) or oligonucleotides in PerfectHyb buffer (Sigma). Washed membranes were exposed to X-ray films at −80 °C for 3 days.

ACKNOWLEDGEMENTS

The authors thank Xinyu Jian, Aurora Luque, and Yujing (Ada) Liu for technical assistance, Gang Yu and Alberto Macho for sharing materials, and all members in Rosa Lozano-Duran's and Alberto Macho's groups for stimulating discussions and helpful suggestions. The authors are grateful to Alberto Macho, Huang Tan, and Tamara Jimenez-Gongora for critical reading of the manuscript. This research was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No. XDB27040206), and by the National Natural Science Foundation of China (NSFC) (grant numbers 31671994 and 31870250). Research in RL-D's lab is funded by the Shanghai Center for Plant Stress Biology of the Chinese Academy of Sciences and the 100 Talent Program of the Chinese Academy of Sciences.

SUPPLEMENTARY MATERIAL

Supplementary figures

- **Supplementary figure 1.** *SlAGO4* expression in TYLCV-infected and control tomato plants.
- **Supplementary figure 2.** V2 is essential for systemic infection in tomato and *N. benthamiana*.
- **Supplementary figure 3.** Experimental design for local and systemic TYLCV infection assays in *NbAGO4*-silenced *N. benthamiana* plants*.*
- **Supplementary figure 4.** Original single-base resolution bisulfite sequencing data of the intergenic region (IR) of TYLCV (WT and V2 null mutant) in local infection assays, related to Figure 4A, B.
- **Supplementary figure 5.** Original single-base resolution bisulfite sequencing data of the intergenic region (IR) of TYLCV and TYLCV-V2null in systemic infection assays, related to Figures 4C.
- **Supplementary figure 6.** Relevance of the Cajal body localization of the V2-AGO4 interaction and TYLCV infection in *Nbcoilin*-silenced plants.
- **Supplementary figure 7.** TYLCV V2$_{L76S}$ interacts with AGO4 in the Cajal body.
- **Supplementary figure 8.** Model for the V2-mediated inhibition of the AGO4-dependent methylation of the viral DNA in the Cajal body.

Supplementary tables

- **Supplementary table 1.** List of primers used for cloning in this study.
- **Supplementary table 2.** List of plasmids used in this study.
- **Supplementary table 3.** List of primers used for qPCR and qRT-PCR in this study.
- **Supplementary table 4.** Values of IR methylation in independent experiments and biological replicates.

Supplementary videos

- **Supplementary video 1.** Z-stack video showing the co-localization of V2-GFP and Fibrillarin-RFP in the Cajal body.
- **Supplementary video 2.** Z-stack video showing the localization of co-expressed GFP-V2 and Fibrillarin-RFP.

REFERENCES

Böhmdorfer G, Rowley MJ, Kuciński J, Zhu Y, Amies I, Wierzbicki AT (2014) RNA-directed DNA methylation requires stepwise binding of silencing factors to long non-coding RNA. *The Plant Journal* 79: 181-191

Bai M, Yang GS, Chen WT, Mao ZC, Kang HX, Chen GH, Yang YH, Xie BY (2012) Genome-wide identification of Dicer-like, Argonaute and RNA-dependent RNA polymerase gene families and their expression analyses in response to viral infection and abiotic stresses in Solanum lycopersicum. *Gene* 501: 52-62

Barneche F, Steinmetz F, Echeverria M (2000) Fibrillarin genes encode both a conserved nucleolar protein and a novel small nucleolar RNA involved in ribosomal RNA methylation in Arabidopsis thaliana. *The Journal of biological chemistry* 275: 27212-20

Brosseau C, El Oirdi M, Adurogbangba A, Ma X, Moffett P (2016) Antiviral Defense Involves AGO4 in an Arabidopsis-Potexvirus Interaction. *Molecular plant-microbe interactions : MPMI* 29: 878-888

Brough CL, Gardiner WE, Inamdar NM, Zhang XY, Ehrlich M, Bisaro DM (1992) DNA methylation inhibits propagation of tomato golden mosaic virus DNA in transfected protoplasts. *Plant Mol Biol* 18: 703-12

Buchmann RC, Asad S, Wolf JN, Mohannath G, Bisaro DM (2009) Geminivirus AL2 and L2 proteins suppress transcriptional gene silencing and cause genome-wide reductions in cytosine methylation. *Journal of virology* 83: 5005-13

Burgyán J, Havelda Z (2011) Viral suppressors of RNA silencing. *Trends Plant Science* 16(5):265-72.

Butterbach P, Verlaan MG, Dullemans A, Lohuis D, Visser RG, Bai Y, Kormelink R (2014) Tomato yellow leaf curl virus resistance by Ty-1 involves increased cytosine methylation of viral genomes and is compromised by cucumber mosaic virus infection. *Proceedings of the National Academy of Sciences of the United States of America* 111: 12942-7

Castillo-Gonzalez C, Liu X, Huang C, Zhao C, Ma Z, Hu T, Sun F, Zhou Y, Zhou X, Wang XJ, Zhang X (2015) Geminivirus-encoded TrAP suppressor inhibits the histone methyltransferase SUVH4/KYP to counter host defense. *eLife* 4

Ceniceros-Ojeda EA, Rodriguez-Negrete EA, Rivera-Bustamante RF (2016) Two Populations of Viral Minichromosomes Are Present in a Geminivirus-Infected Plant Showing Symptom Remission (Recovery). *Journal of virology* 90: 3828-3838

Chan SW, Henderson IR, Jacobsen SE (2005) Gardening the genome: DNA methylation in Arabidopsis thaliana. *Nature reviews Genetics* 6: 351-60

Chen H, Zou Y, Shang Y, Lin H, Wang Y, Cai R, Tang X, Zhou JM (2008) Firefly luciferase complementation imaging assay for protein-protein interactions in plants. *Plant physiology* 146: 368-76

Coursey T, Regedanz E, Bisaro DM (2018) Arabidopsis RNA Polymerase V Mediates Enhanced Compaction and Silencing of Geminivirus and Transposon Chromatin during Host Recovery from Infection. 92

Dereeper A, Audic S, Claverie JM, Blanc G (2010) BLAST-EXPLORER helps you building datasets for phylogenetic analysis. *BMC evolutionary biology* 10: 8

Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, Dufayard JF, Guindon S, Lefort V, Lescot M, Claverie JM, Gascuel O (2008) Phylogeny.fr: robust phylogenetic analysis for the non-specialist. *Nucleic acids research* 36: W465-9

Deuschle K, Kepp G, Jeske H (2016) Differential methylation of the circular DNA in geminiviral minichromosomes. *Virology* 499: 243-258

Dogar AM (2006) RNAi dependent epigenetic marks on a geminivirus promoter. *Virology journal* 3: 5

552 **Ermak G**, Paszkowski U, Wohlmuth M, Mittelsten Scheid O, Paszkowski J (1993) Cytosine
553 methylation inhibits replication of African cassava mosaic virus by two distinct mechanisms.
554 *Nucleic acids research* 21: 3445-50
555

556 **Exposito-Rodriguez M**, Borges AA, Borges-Perez A, Perez JA (2008) Selection of internal
557 control genes for quantitative real-time RT-PCR studies during tomato development process.
558 *BMC plant biology* 8: 131
559

560 **Gao Z**, Liu H-L, Daxinger L, Pontes O, He X, Qian W, Lin H, Xie M, Lorkovic ZJ, Zhang S (2010)
561 An RNA polymerase II-and AGO4-associated protein acts in RNA-directed DNA methylation.
562 *Nature* 465: 106
563

564 **Grefen C**, Blatt MR (2012) A 2in1 cloning system enables ratiometric bimolecular fluorescence
565 complementation (rBiFC). *BioTechniques* 53: 311-14
566

567 **Gruntman E**, Qi Y, Slotkin RK, Roeder T, Martienssen RA, Sachidanandam R (2008) Kismeth:
568 analyzer of plant methylation states through bisulfite sequencing. *BMC bioinformatics* 9: 371
569

570 **Haag JR**, Ream TS, Marasco M, Nicora CD, Norbeck AD, Pasa-Tolic L, Pikaard CS (2012) In
571 vitro transcription activities of Pol IV, Pol V, and RDR2 reveal coupling of Pol IV and RDR2 for
572 dsRNA synthesis in plant RNA silencing. *Molecular cell* 48: 811-818
573

574 **Hanley-Bowdoin L**, Bejarano ER, Robertson D, Mansoor S (2013) Geminiviruses: masters at
575 redirecting and reprogramming plant processes. *Nature reviews Microbiology* 11: 777-88
576

577 **He L**, Wu W, Zinta G, Yang L, Wang D, Liu R, Zhang H, Zheng Z, Huang H, Zhang Q, Zhu J-K
578 (2018) A naturally occurring epiallele associates with leaf senescence and local climate
579 adaptation in Arabidopsis accessions. *Nature communications* 9: 460
580

581 **Ismayil A**, Haxim Y, Wang Y, Li H, Qian L, Han T, Chen T, Jia Q, Yihao Liu A, Zhu S, Deng H,
582 Gorovits R, Hong Y, Hanley-Bowdoin L, Liu Y (2018) Cotton Leaf Curl Multan virus C4 protein
583 suppresses both transcriptional and post-transcriptional gene silencing by interacting with SAM
584 synthetase. *PLoS pathogens* 14: e1007282
585

586 **Jackel JN**, Storer JM, Coursey T, Bisaro DM (2016) Arabidopsis RNA Polymerases IV and V Are
587 Required To Establish H3K9 Methylation, but Not Cytosine Methylation, on Geminivirus
588 Chromatin. *Journal of virology* 90: 7529-40
589
590 **Kim SH**, Macfarlane S, Kalinina NO, Rakitina DV, Ryabov EV, Gillespie T, Haupt S, Brown JW,
591 Taliansky M (2007a) Interaction of a plant virus-encoded protein with the major nucleolar protein
592 fibrillarin is required for systemic virus infection. *Proceedings of the National Academy of Sciences*
593 *of the United States of America* 104: 11115-20
594
595 **Kim SH**, Ryabov EV, Kalinina NO, Rakitina DV, Gillespie T, MacFarlane S, Haupt S, Brown JW,
596 Taliansky M (2007b) Cajal bodies and the nucleolus are required for a plant virus systemic
597 infection. *The EMBO journal* 26: 2169-79
598
599 **Kushwaha NK**, Mansi, Chakraborty S (2017) The replication initiator protein of a geminivirus
600 interacts with host monoubiquitination machinery and stimulates transcription of the viral genome.
601 *PLoS pathogens* 13: e1006587
602
603 **Law JA**, Vashisht AA, Wohlschlegel JA, Jacobsen SE (2011) SHH1, a homeodomain protein
604 required for DNA methylation, as well as RDR2, RDM4, and chromatin remodeling factors,
605 associate with RNA polymerase IV. *PLoS genetics* 7: e1002195
606
607 **Li CF**, Henderson IR, Song L, Fedoroff N, Lagrange T, Jacobsen SE (2008) Dynamic regulation
608 of ARGONAUTE4 within multiple nuclear bodies in Arabidopsis thaliana. *PLoS genetics* 4: e27
609
610 **Li CF**, Pontes O, El-Shami M, Henderson IR, Bernatavichute YV, Chan SW, Lagrange T, Pikaard
611 CS, Jacobsen SE (2006) An ARGONAUTE4-containing nuclear processing center colocalized
612 with Cajal bodies in Arabidopsis thaliana. *Cell* 126: 93-106
613
614 **Liu D**, Shi L, Han C, Yu J, Li D, Zhang Y (2012) Validation of reference genes for gene expression
615 studies in virus-infected Nicotiana benthamiana using quantitative real-time PCR. *PloS one* 7:
616 e46451
617
618 **Lozano-Duran R**, Rosas-Diaz T, Luna AP, Bejarano ER (2011) Identification of host genes
619 involved in geminivirus infection using a reverse genetics approach. *PloS one* 6: e22383

Ma X, Nicole MC, Meteignier LV, Hong N, Wang G, Moffett P (2015) Different roles for RNA silencing and RNA processing components in virus recovery and virus-induced gene silencing in plants. *Journal of experimental botany* 66: 919-32

Maimbo M, Ohnishi K, Hikichi Y, Yoshioka H, Kiba A (2010) S-glycoprotein-like protein regulates defense responses in Nicotiana plants against Ralstonia solanacearum. *Plant physiology* 152: 2023-35

Mason G, Caciagli P, Accotto GP, Noris E (2008) Real-time PCR for the quantitation of Tomato yellow leaf curl Sardinia virus in tomato plants and in Bemisia tabaci. *Journal of virological methods* 147: 282-9

Matzke MA, Kanno T, Matzke AJ (2015) RNA-Directed DNA Methylation: The Evolution of a Complex Epigenetic Pathway in Flowering Plants. *Annual review of plant biology* 66: 243-67

Matzke MA, Mosher RA (2014) RNA-directed DNA methylation: an epigenetic pathway of increasing complexity. *Nature reviews Genetics* 15: 394-408

Minas K, McEwan NR, Newbold CJ, Scott KP (2011) Optimization of a high-throughput CTAB-based protocol for the extraction of qPCR-grade DNA from rumen fluid, plant and bacterial pure cultures. *FEMS microbiology letters* 325: 162-9

Minoia S, Carbonell A, Di Serio F, Gisel A, Carrington JC, Navarro B, Flores R (2014) Specific argonautes selectively bind small RNAs derived from potato spindle tuber viroid and attenuate viroid accumulation in vivo. *Journal of virology* 88: 11933-45

Nakagawa T, Suzuki T, Murata S, Nakamura S, Hino T, Maeo K, Tabata R, Kawai T, Tanaka K, Niwa Y, Watanabe Y, Nakamura K, Kimura T, Ishiguro S (2007) Improved Gateway binary vectors: high-performance vectors for creation of fusion constructs in transgenic analysis of plants. *Bioscience, biotechnology, and biochemistry* 71: 2095-100

Piedra-Aguilera A, Jiao C, Luna AP, Villanueva F, Dabad M, Esteve-Codina A, Diaz-Pendon JA, Fei Z, Bejarano ER, Castillo AG (2019) Integrated single-base resolution maps of transcriptome,

sRNAome and methylome of Tomato yellow leaf curl virus (TYLCV) in tomato. *Scientific reports*
9: 2863

Raja P, Sanville BC, Buchmann RC, Bisaro DM (2008) Viral genome methylation as an epigenetic
defense against geminiviruses. *Journal of virology* 82: 8997-9007

Ratcliff F, Martin-Hernandez AM, Baulcombe DC (2001) Technical Advance. Tobacco rattle virus
as a vector for analysis of gene function by silencing. *The Plant journal : for cell and molecular*
biology 25: 237-45

Rawlings RA, Krishnan V, Walter NG (2011) Viral RNAi suppressor reversibly binds siRNA to
outcompete Dicer and RISC via multiple turnover. *Journal of Molecular Biology* 408(2):262-76.

Rodriguez-Negrete E, Lozano-Duran R, Piedra-Aguilera A, Cruzado L, Bejarano ER, Castillo AG
(2013) Geminivirus Rep protein interferes with the plant DNA methylation machinery and
suppresses transcriptional gene silencing. *The New phytologist* 199: 464-75

Rosas-Diaz T, Zhang D, Fan P, Wang L, Ding X, Jiang Y, Jimenez-Gongora T, Medina-Puche L,
Zhao X, Feng Z, Zhang G, Liu X, Bejarano ER, Tan L, Zhang H, Zhu JK, Xing W, Faulkner C,
Nagawa S, Lozano-Duran R (2018) A virus-targeted plant receptor-like kinase promotes cell-to-
cell spread of RNAi. *Proceedings of the National Academy of Sciences of the United States of*
America 115: 1388-1393

Schott G, Mari-Ordonez A, Himber C, Alioua A, Voinnet O, Dunoyer P (2012) Differential effects
of viral silencing suppressors on siRNA and miRNA loading support the existence of two distinct
cellular pools of ARGONAUTE1. *The EMBO journal* 31: 2553-65

Semashko MA, Rakitina DV, Gonzalez I, Canto T, Kalinina NO, Taliansky ME (2012) Movement
protein of hordeivirus interacts in vitro and in vivo with coilin, a major structural protein of Cajal
bodies. *Doklady Biochemistry and biophysics* 442: 57-60

Shaw J, Love AJ, Makarova SS, Kalinina NO, Harrison BD, Taliansky ME (2014) Coilin, the
signature protein of Cajal bodies, differentially modulates the interactions of plants with viruses in
widely different taxa. *Nucleus* 5: 85-94

Taylor KW, Kim JG, Su XB, Aakre CD, Roden JA, Adams CM, Mudgett MB (2012) Tomato TFT1 is required for PAMP-triggered immunity and mutations that prevent T3S effector XopN from binding to TFT1 attenuate Xanthomonas virulence. *PLoS pathogens* 8: e1002768

Torchetti EM, Pegoraro M, Navarro B, Catoni M, Di Serio F, Noris E (2016) A nuclear-replicating viroid antagonizes infectivity and accumulation of a geminivirus by upregulating methylation-related genes and inducing hypermethylation of viral DNA. *Scientific reports* 6: 35101

Tu YC, Tsai WS, Wei JY, Chang KY, Tien CC, Hsiao HY, Fu SF (2017) The C2 protein of tomato leaf curl Taiwan virus is a pathogenicity determinant that interferes with expression of host genes encoding chromomethylases. *Physiologia plantarum* 161: 515-531

Wang B, Li F, Huang C, Yang X, Qian Y, Xie Y, Zhou X (2014) V2 of tomato yellow leaf curl virus can suppress methylation-mediated transcriptional gene silencing in plants. *The Journal of general virology* 95: 225-30

Wang B, Yang X, Wang Y, Xie Y, Zhou X (2018) Tomato Yellow Leaf Curl Virus V2 Interacts with Host Histone Deacetylase 6 To Suppress Methylation-Mediated Transcriptional Gene Silencing in Plants. *Journal of virology* 92

Wang L, Ding X, Xiao J, Jimenez-Gomicronngora T, Liu R, Lozano-Duran R (2017a) Inference of a Geminivirus-Host Protein-Protein Interaction Network through Affinity Purification and Mass Spectrometry Analysis. *Viruses* 9

Wang L, Tan H, Wu M, Jimenez-Gongora T, Tan L, Lozano-Duran R (2017b) Dynamic Virus-Dependent Subnuclear Localization of the Capsid Protein from a Geminivirus. *Frontiers in plant science* 8: 2165

Wang Y, Wu Y, Gong Q, Ismayil A, Yuan Y, Lian B, Jia Q, Han M, Deng H, Hong Y, Hanley-Bowdoin L, Qi Y, Liu Y (2019) Geminiviral V2 Protein Suppresses Transcriptional Gene Silencing through Interaction with AGO4. *Journal of virology* 93

721 **Wartig L**, Kheyr-Pour A, Noris E, De Kouchkovsky F, Jouanneau F, Gronenborn B, Jupin I (1997)
722 Genetic analysis of the monopartite tomato yellow leaf curl geminivirus: roles of V1, V2, and C2
723 ORFs in viral pathogenesis. *Virology* 228: 132-40
724

725 **Woo HR**, Dittmer TA, Richards EJ (2008) Three SRA-domain methylcytosine-binding proteins
726 cooperate to maintain global CpG methylation and epigenetic silencing in Arabidopsis. *PLoS*
727 *genetics* 4: e1000156
728

729 **Xie Z**, Johansen LK, Gustafson AM, Kasschau KD, Lellis AD, Zilberman D, Jacobsen SE,
730 Carrington JC (2004) Genetic and functional diversification of small RNA pathways in plants.
731 *PLoS biology* 2: e104
732

733 **Yadav RK**, Chattopadhyay D (2011) Enhanced viral intergenic region-specific short interfering
734 RNA accumulation and DNA methylation correlates with resistance against a geminivirus.
735 *Molecular plant-microbe interactions : MPMI* 24: 1189-97
736

737 **Yang D-L**, Zhang G, Tang K, Li J, Yang L, Huang H, Zhang H, Zhu J-K (2015) Dicer-independent
738 RNA-directed DNA methylation in Arabidopsis. *Cell research*
739

740 **Yang LP**, Fang YY, An CP, Dong L, Zhang ZH, Chen H, Xie Q, Guo HS (2013) C2-mediated
741 decrease in DNA methylation, accumulation of siRNAs, and increase in expression for genes
742 involved in defense pathways in plants infected with beet severe curly top virus. *The Plant journal :*
743 *for cell and molecular biology* 73: 910-7
744

745 **Yang X**, Xie Y, Raja P, Li S, Wolf JN, Shen Q, Bisaro DM, Zhou X (2011) Suppression of
746 methylation-mediated transcriptional gene silencing by betaC1-SAHH protein interaction during
747 geminivirus-betasatellite infection. *PLoS pathogens* 7: e1002329
748

749 **Yu G**, Xian L, Xue H, Yu W, Rufian J, Sang Y, Morcillo R, Wang Y, Macho AP (2019) A bacterial
750 effector protein prevents MAPK-mediated phosphorylation of SGT1 to suppress plant immunity.
751 *bioRxiv*: 641241
752

Zhang Z, Chen H, Huang X, Xia R, Zhao Q, Lai J, Teng K, Li Y, Liang L, Du Q, Zhou X, Guo H,

Xie Q (2011) BSCTV C2 attenuates the degradation of SAMDC1 to suppress DNA methylation-

mediated gene silencing in Arabidopsis. *The Plant cell* 23: 273-88

Zhong X, Du J, Hale CJ, Gallego-Bartolome J, Feng S, Vashisht AA, Chory J, Wohlschlegel JA,

Patel DJ, Jacobsen SE (2014) Molecular mechanism of action of plant DRM de novo DNA

methyltransferases. *Cell* 157: 1050-1060

Zhong X, Wang ZQ, Xiao R, Wang Y, Xie Y, Zhou X (2017) iTRAQ analysis of the tobacco leaf

proteome reveals that RNA-directed DNA methylation (RdDM) has important roles in defense

against geminivirus-betasatellite infection. *Journal of proteomics* 152: 88-101

Zilberman D, Cao X, Jacobsen SE (2003) ARGONAUTE4 control of locus-specific siRNA

accumulation and DNA and histone methylation. *Science (New York, NY)* 299: 716-719

A

Identified protein name	Identified protein ID	Bait protein name	Exclusive unique peptide count	Identified protein coverage (%)	Best Mascot Score
NbAGO4-1	NbS00018144g003.1	GFP	- / - / -	- / - / -	- / - / -
		V2-GFP	3 / - / -	6.1 / - / -	44.9 / - / -
NbAGO4-2	NbS00007950g0008.1	GFP	- / - / -	- / - / -	- / - / -
		V2-GFP	16 / 1 / 1	23 / 1.8 / 1.2	116.2 / 35.7 / 59.2

Figure 1. V2 interacts with AGO4 from *N. benthamiana*.

(A) Unique peptide count, protein coverage, and best Mascot Score of NbAGO4-1 and NbAGO4-2 co-immunoprecipitated with V2-GFP, as identified by affinity purification followed by mass spectrometry (AP-MS). Results from three independent biological repeats are shown. "-" indicates no peptide was detected.

(B) 3xHA-NbAGO4-1 and 3xHA-NbAGO4-2 specifically interact with V2-GFP in co-immunoprecipitation (co-IP) assays upon transient expression in *N. benthamiana*. Free GFP was used as negative control. CBB, Coomassie brilliant blue staining. Three independent biological replicates were performed with similar results.

(C) NbAGO4-1 and NbAGO4-2 interact with V2 in split-luciferase assays. V2-N-luc and C-luc-NbAGO4-1/2 were transiently co-expressed in *N. benthamiana*; C-luc-SlWRKY75 is used as negative control. The luciferase bioluminescence from at least three independent leaves per experiment was imaged two days after infiltration. The average bioluminescence, measured in counts per second (cps), as well as an image of a representative leaf are shown. Values represent the mean of three independent biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant difference (according to a Student's *t*-test, **: P<0.01, ***: P<0.001.) compared to the negative control.

Figure 2. V2 interacts with AGO4 from tomato.

(A) Phylogenetic tree of AtAGO4, NbAGO4, and SlAGO4 proteins. The phylogenetic analysis was performed with phylogeny.fr (Dereeper et al., 2010, Dereeper et al., 2008).

(B) Pairwise identity and genetic distance matrix among AtAGO4, NbAGO4 and SlAGO4 proteins. The analysis was performed by Geneious (https://www.geneious.com).

(C) 3xHA-SlAGO4a, 3xHA-SlAGO4b, and 3xHA-SlAGO4d specifically interact with V2-GFP in co-immunoprecipitation (co-IP) assays upon transient expression in *N. benthamiana*. Free GFP was used as negative control. CBB, Coomassie brilliant blue staining. Three independent biological replicates were performed with similar results.

(D) SlAGO4a and SlAGO4b interact with V2 in split-luciferase assays. V2-N-luc and C-luc-SlAGO4a/b were transiently co-expressed in *N. benthamiana*; C-luc-SlWRKY75 is used as negative control. The luciferase bioluminescence from at least three independent leaves per experiment was imaged two days after infiltration. The average bioluminescence, measured in counts per second (cps), as well as an image of a representative leaf are shown. Values represent the mean of three independent biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant difference (according to a Student's *t*-test, ***: P<0.001.) compared to the negative control.

Figure 3. V2 counters the AGO4-dependent anti-viral defence to promote virulence.

(A) Expression of *NbAGO4-1, NbAGO4-2,* and *NbAGO6* in *N. benthamiana* plants infected with
   TRV-EV (empty vector) or TRV-*NbAGO4*, measured by reverse transcription quantitative real-
   time PCR (RT-qPCR). Gene expression was normalized to *NbTubulin*. Values are the mean
   of four independent biological replicates; error bars indicate SEM. Asterisks indicate a
   statistically significant difference according to Student's *t*-test. **: P<0.01, ***: P<0.001, ns:
   not significant.

(B) Representative pictures of *N. benthamiana* plants infected with the indicated combinations of
   viruses. Photographs were taken at 3 weeks post-inoculation (wpi).

(C) *NbAGO4-1* expression in *NbAGO4*-silenced plants and control plants infected with TYLCV, TYLCV-V2null, or mock-inoculated at 3 wpi measured by RT-qPCR. Gene expression was normalized to *NbTubulin*. Values are the mean of six independent biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant difference according to Student's *t*-test. *: P<0.05, **: P<0.01, ****: P<0.0001.

(D) *NbAGO4-2* expression in *NbAGO4*-silenced plants and control plants infected by TYLCV, TYLCV-V2null or mock-inoculated at 3 wpi measured by RT-qPCR. Gene expression was normalized to *Tubulin*. Values are the mean of six independent biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant difference according to Student's *t*-test. *: P<0.05, **: P<0.01, ***: P<0.001.

(E) Viral (TYLCV) accumulation in local infections in *NbAGO4*-silenced or control plants, measured by qPCR. Infiltrated leaf patches from different plants were collected at 4 dpi. The experimental design is shown in Figure S3A. The accumulation of viral DNA is normalized to the *25S ribosomal RNA interspacer (ITS)*. Values are the mean of eight independent biological replicates; error bars indicate SEM.

(F) Viral (TYLCV) accumulation in systemic infections in *NbAGO4*-silenced or control plants, measured by qPCR. Apical leaves from six plants were collected at 3 wpi. The experimental design is shown in Figure S3B. The accumulation of viral DNA is normalized to the *25S ribosomal RNA interspacer (ITS)*. Four independent biological replicates were performed with similar results; one representative result is shown. Values are the mean of six independent biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant difference according to Student's *t*-test. *: P<0.05, ***: P<0.001. The relative fold change of viral accumulation between *NbAGO4*-silenced plants and control plants is shown above each column.

Figure 4. V2 suppresses the AGO4-dependent methylation of viral DNA.

(A) Percentage of methylated cytosines in the intergenic region (IR) of TYLCV in local infection assays with TYLCV WT or V2 null mutant (TYLCV-V2null) in *N. benthamiana* at 3 or 9 days post-inoculation (dpi), as detected by bisulfite sequencing. The original single-base resolution bisulfite sequencing data are shown in Figure S4A. Values are the mean of three independent biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant difference according to Student's *t*-test. ***: P<0.001. The values of cytosine methylation in each biological replicate are shown in Supplemental table 4.

(B) Percentage of methylated cytosines in the intergenic region (IR) of TYLCV in local infection assays with the V2 null mutant TYLCV (TYLCV-V2null) in *AGO4*-silenced (TRV-*NbAGO4*) or control (TRV-EV) *N. benthamiana* plants at 4 dpi, as detected by bisulfite sequencing. Samples come from the same plants used in Figure 3E. The original single-base resolution bisulfite sequencing data are shown in Figure S4B. Values are the mean of four independent biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant difference according to Student's *t*-test. *: P<0.05. The values of cytosine methylation in each biological replicate are shown in Supplemental table 4.

(C) Percentage of methylated cytosines in the intergenic region (IR) of TYLCV in systemic

infection assays with TYLCV WT or V2 null mutant (TYLCV-V2null) in *AGO4*-silenced (TRV-NbAGO4) or control (TRV-EV) *N. benthamiana* plants at 3 weeks post-inoculation (wpi), as detected by bisulfite sequencing. Samples come from the same plants used in Figure 3F. The original single-base resolution bisulfite sequencing data are shown in Figure S5A. Values are the mean of four independent biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant difference according to Student's *t*-test. *: P<0.05. The values of cytosine methylation in each biological replicate are shown in Supplemental table 4.

Figure 5. V2 interferes with AGO4 binding to the viral genome but does not hamper production or loading of vsiRNA.

(A) 3xFLAG-NbAGO4-1 binds the viral (TYLCV) genome. Binding was detected by chromatin immunoprecipitation (ChIP) upon transient expression in *N. benthamiana* followed by qPCR. Two regions of the viral genome, the IR and the V2 ORF, were analyzed; *ACTIN* was used as negative control. Values represent the mean of four independent biological replicates; error bars represent SEM. Asterisks indicate a significant difference according to a Student's *t*-test; the *P*-value for the different comparisons is shown. *: P<0.05.

(B) NbAGO4-1 binds viral small interfering RNA (vsiRNA) independently of V2. Northern blot of vsiRNA in total extracts or 3xFLAG-NbAGO4 immunoprecipitates (NbAGO4-IP) of *N. benthamiana* leaf patches infiltrated with TYLCV WT or V2 null mutant infectious clones (TYLCV, TYLCV-V2null) at two days after infiltration . Detection was performed with a ^{32}P-labeled DNA probe for the intergenic region (IR).

Figure 6. V2 interacts with AGO4 in the Cajal body.

(A) V2-GFP and RFP-AGO4 co-localize in the Cajal body. CFP-Fibrillarin, V2-GFP and RFP-NbAGO4-1/2 or RFP-SlAGO4a/b/d were transiently co-expressed in *N. benthamiana* epidermal cells. CFP-Fibrillarin is used as a nucleolus and Cajal body marker. Confocal images were taken at two days after infiltration. Arrowheads indicate the position of the Cajal body. Bar, 5µm. This experiment was repeated more than three times with same results.

(B) V2 interacts with AGO4 in the Cajal body. The N-terminal half of the YFP fused to V2 (V2-

nYFP) was transiently co-expressed with the C-terminal half of the YFP alone (cYFP, as a negative control), or cYFP-NbAGO4, cYFP-SlAGO4, or cYFP-V2 (as a positive control) in *N. benthamiana* leaves. CFP-Fibrillarin was used as a nucleolus and Cajal body marker. Confocal images were taken at two days after infiltration. Yellow fluorescence indicates a positive interaction. Arrowheads indicate the position of the Cajal body. Bar, 5µm. This experiment was repeated more than three times with similar results.

Figure 7. Methylation of the viral DNA and its suppression by V2 occur in the Cajal body.

(A) V2-GFP co-localizes with Fibrillarin-RFP (Cajal body marker) in the Cajal body, while GFP-V2 does not. RFP-Fibrillarin and V2-GFP or GFP-V2 were transiently co-expressed in *N. benthamiana* epidermal cells. Confocal images were taken at two days after infiltration. Arrowheads indicate the position of the Cajal body. Bar, 5μm. This experiment was repeated three times with similar results.

(B) V2-GFP, but not GFP-V2, can restore the suppression of viral DNA methylation of a V2 null TYLCV mutant in local infection assays. Cytosine methylation in the intergenic region (IR) of the V2 null mutant genome in locally infected leaf patches of *N. benthamiana* expressing V2, V2-GFP, or GFP-V2 was detected by bisulfite sequencing at 3 days post-inoculation (dpi). TYLCV or TYLCV-V2null alone were used as controls. The original single-base resolution bisulfite sequencing data are shown in Figure S6A. This experiment was repeated twice with similar results.

(C) *Nbcoilin* expression in *Nbcoilin*-silenced (TRV-*Nbcoilin*) and control plants (TRV-EV) infected with TYLCV, TYLCV-V2null, or mock-inoculated at 3 wpi measured by RT-qPCR. Gene expression was normalized to *NbTubulin*. Values are the mean of six independent biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant difference

according to Student's *t*-test. **: P<0.01.

(D) Percentage of methylated cytosines in the intergenic region (IR) of TYLCV in systemic infection assays with TYLCV WT or V2 null mutant (TYLCV-V2null) in *Nbcoilin*-silenced (TRV-*Nbcoilin*) or control (TRV-EV) *N. benthamiana* plants at 3 weeks post-inoculation (wpi), as detected by bisulfite sequencing. Values are the mean of three independent biological replicates; error bars indicate SEM. The values of cytosine methylation in each independent biological replicate are shown in Supplemental table 4. The original single-base resolution bisulfite sequencing data are shown in Fig S6F.

Supplementary table 1. List of primers used for cloning in this study.

Plasmid name	Vector name (Source)	Primer sequence
(1) TOPO cloning		
TOPO-*NbAGO4-1* (with stop codon)	pENTR/D-TOPO (Invitrogen)	F: CACCATGGCTGAAGAAGACAATGG R: TTAACAGAAGAACATGGAACTAGAAACAT
TOPO-*NbAGO4-2* (with stop codon)	pENTR/D-TOPO (Invitrogen)	F: CACCATGGCTGAAGAAGATAAGAAT R: TCAACAAAAGAACATGGAACTGGAAACT
TOPO-*SlAGO4a* (with stop codon)	pENTR/D-TOPO (Invitrogen)	F: CACCATGGCTGAAGAAGAAACAAATG R: TTAGCAAAAGAACATAGAACTGGAAAC
TOPO-*SlAGO4b* (with stop codon)	pENTR/D-TOPO (Invitrogen)	F: CACCATGGCTGAAGAAGAGAATGGT R: TCAACAGAAGAACATGGAGCTAGCAAC
TOPO-*SlAGO4d* (with stop codon)	pENTR/D-TOPO (Invitrogen)	F: CACCATGGCCTCTTCAAAAGATGAAG R: TTAGCAGAAGAACATTGAACTGCGAACAT
TOPO-*SlWRKY75* (with stop codon)	pENTR/D-TOPO (Invitrogen)	F: CACCATGGAGAATTATGCAACAATATTTCC R: AAAGGAATTATAGATTTGCATTTGAC
TOPO-VIGS-*NbAGO4*	pENTR/D-TOPO (Invitrogen)	F: CACCATGATTCTTGGAATGGA R: GTGACCACTTCTCATCAAGA
TOPO-VIGS-*Nbcoilin*	pENTR/D-TOPO (Invitrogen)	F：CACCTTTATAATGAAGACGGATCTCTGGA R：GAAGGCCATTGTAGGGCCCAG
(2) BP reaction		
pDONR221-P1P4-V2 / pDONR221-P1P4-V2$_{L76S}$	pDONR221-P1P4 (Invitrogen)	F: GGGGACAAGTTTGTACAAAAAAGCAGGCTTAATGTGGGACCCACTTCTAAATGAA R: GGGGACAACTTTGTATAGAAAAGTTGGGTTCAGGGCTTCGATACATTCTGT
pDONR221-P1P4-*NbAGO4-1*	pDONR221-P1P4 (Invitrogen)	F: GGGGACAAGTTTGTACAAAAAAGCAGGCTTAATGGCTGAAGAAGACAATGGTGG R: GGGGACAACTTTGTATAGAAAAGTTGGGTTTAACAGAAGAACATGGAACTAGAAACATT
pDONR221-P1P4-*NbAGO4-2*	pDONR221-P1P4 (Invitrogen)	F: GGGGACAAGTTTGTACAAAAAAGCAGGCTTAATGGCTGAAGAAGATAAGAAT R: GGGGACAACTTTGTATAGAAAAGTTGGGTTCAACAAAAGAACATGGAACTGGAAACT
pDONR221-P1P4-*SlAGO4a*	pDONR221-P1P4 (Invitrogen)	F: GGGGACAAGTTTGTACAAAAAAGCAGGCTTAATGGCTGAAGAAGAAACAAATG R: GGGGACAACTTTGTATAGAAAAGTTGGGTTTAGCAAAAGAACATAGAACTGGAAAC

pDONR221-P1P4-*SlAGO4b*	pDONR221-P1P4 (Invitrogen)	F: GGGGACAAGTTTGTACAAAAAAGCAGGCTTAATGGCTGAAGAAGAGAATGGTG R: GGGGACAACTTTGTATAGAAAAGTTGGGTTCAACAGAAGAACATGGAGCTAGCAACG
pDONR221-P3P2-V2 / pDONR221-P3P2-V2$_{L76S}$	pDONR221-P3P2 (Invitrogen)	F: GGGGACAACTTTGTATAATAAAGTTGGAATGTGGGACCCACTTCTAAATGAA R: GGGGACCACTTTGTACAAGAAAGCTGGGTGGGGCTTCGATACATTCTGTATATTCT
(3) Infusion cloning		
pCambia1300-3xFLAG-*NbAGO4-1*	pCAMBIA1300 (Cambia)	1300-FLAG-F: CGGGGGACTGTCGACTCTAGAATGGACTACAAAG 1300-NbAGO4-1-FLAG-R: TGTCTTCTTCAGCCATCTTATCGTCATCGTCCTTG NbAGO4-1-F: ATGGCTGAAGAAGACAATGGTGG 1300-NBAGO4-1-R: TCGATCAATCAGGTACCCGGGTTAACAGAAGAACATGGAACT
(4) V2 null mutagenesis		
TOPO-TYLCV-V2null	TOPO-TYLCV (Rosas-Diaz et al., 2018)	F: GTCTTGCAATATGTAGGACCCACTTCTAAATGAATTTCCTG R: CAGGAAATTCATTTAGAAGTGGGTCCTACATATTGCAAGAC
TOPO-TYLCV-V2$_{L76S}$	TOPO-TYLCV (Rosas-Diaz et al., 2018)	F: GCCGAAGGCTGAATCTCGACAGC R: GCTGTCGAGATTCAGCCTTCGGC
TOPO-V2$_{L76S}$	TOPO-V2 (Wang et al., 2017a)	F: GCCGAAGGCTGAATCTCGACAGC R: GCTGTCGAGATTCAGCCTTCGGC

Supplementary table 2. List of plasmids used in this study.

Plasmid name	Entry vector name (Source)	Binary vector (Source)
(1) Confocal microscopy		
RFP-*NbAGO4-1*	TOPO-*NbAGO4-1* (This study)	pGWB555 (Nakagawa et al., 2007)
RFP-*NbAGO4-2*	TOPO-*NbAGO4-2* (This study)	pGWB555 (Nakagawa et al., 2007)
RFP-*SlAGO4a*	TOPO-*SlAGO4a* (This study)	pGWB555 (Nakagawa et al., 2007)
RFP-*SlAGO4b*	TOPO-*SlAGO4b* (This study)	pGWB555 (Nakagawa et al., 2007)
RFP-*SlAGO4d*	TOPO-*SlAGO4d* (This study)	pGWB555 (Nakagawa et al., 2007)
CFP-Fibrillarin	pDONR207-*Fibrillarin* (Kim et al., 2007b)	pGWB545 (Nakagawa et al., 2007)
(2) Luciferase Complementation Imaging assay		
V2-N-luc	TOPO-V2 (Wang et al., 2017a)	pGWB-N-luc (Yu et al., 2019)
C-luc-*NbAGO4-1*	TOPO-*NbAGO4-1* (This study)	pGWB-C-luc (Yu et al., 2019)
C-luc-*NbAGO4-2*	TOPO-*NbAGO4-2* (This study)	pGWB-C-luc (Yu et al., 2019)
C-luc-*SlAGO4a*	TOPO-*SlAGO4a* (This study)	pGWB-C-luc (Yu et al., 2019)
C-luc-*SlAGO4b*	TOPO-*SlAGO4b* (This study)	pGWB-C-luc (Yu et al., 2019)
C-luc-*SlWRKY75*	TOPO-*SlWRKY75* (This study)	pGWB-C-luc (Yu et al., 2019)
(3) Immunoprecipitation		
3xHA-*NbAGO4-1*	TOPO-*NbAGO4-1* (This study)	pGWB515 (Nakagawa et al., 2007)
3xHA-*NbAGO4-2*	TOPO-*NbAGO4-2* (This study)	pGWB515 (Nakagawa et al., 2007)
3xHA-*SlAGO4a*	TOPO-*SlAGO4a* (This study)	pGWB515 (Nakagawa et al., 2007)
3xHA-*SlAGO4b*	TOPO-*SlAGO4b* (This study)	pGWB515 (Nakagawa et al., 2007)
3xHA-*SlAGO4d*	TOPO-*SlAGO4d* (This study)	pGWB515 (Nakagawa et al., 2007)
V2$_{L76S}$-GFP	TOPO-V2$_{L76S}$ (This study)	pGWB505 (Nakagawa et al., 2007)
(4) Virus-induced gene silencing		
TRV-*NbAGO4*	TOPO-VIGS-*NbAGO4* (This study)	pTRV2-Gw (Taylor et al., 2012)
TRV-*Nbcoilin*	TOPO-VIGS-*Nbcoilin* (This study)	pTRV2-Gw (Taylor et al., 2012)
(5) BiFC		
V2-nYFP + cYFP-empty	pDONR221-P3P2-V2 (This study)	pBiFC-2in1-CN (Grefen & Blatt, 2012)
V2-nYFP + cYFP-V2	pDONR221-P3P2-V2 (This study) + pDONR221-P1P4-V2 (This study)	pBiFC-2in1-CN (Grefen & Blatt, 2012)
V2-nYFP + cYFP-*NbAGO4-1*	pDONR221-P3P2-V2 (This study) + pDONR221-P1P4-*NbAGO4-1* (This study)	pBiFC-2in1-CN (Grefen & Blatt, 2012)
V2-nYFP + cYFP-*NbAGO4-2*	pDONR221-P3P2-V2 (This study) + pDONR221-P1P4-*NbAGO4-2* (This study)	pBiFC-2in1-CN (Grefen & Blatt, 2012)

V2-nYFP + cYFP-Sl*AGO4a*	pDONR221-P3P2-V2 (This study) + pDONR221-P1P4-Sl*AGO4a* (This study)	pBiFC-2in1-CN (Grefen & Blatt, 2012)
V2-nYFP + cYFP-Sl*AGO4b*	pDONR221-P3P2-V2 (This study) + pDONR221-P1P4- Sl*AGO4b* (This study)	pBiFC-2in1-CN (Grefen & Blatt, 2012)
V2$_{L76S}$-nYFP + cYFP-empty	pDONR221-P3P2-V2$_{L76S}$ (This study)	pBiFC-2in1-CN (Grefen & Blatt, 2012)
V2$_{L76S}$-nYFP + cYFP-V2$_{L76S}$	pDONR221-P3P2-V2$_{L76S}$ (This study) + pDONR221-P1P4-V2$_{L76S}$ (This study)	pBiFC-2in1-CN (Grefen & Blatt, 2012)
V2$_{L76S}$-nYFP + cYFP-*NbAGO4-1*	pDONR221-P3P2-V2$_{L76S}$ (This study) + pDONR221-P1P4-*NbAGO4-1* (This study)	pBiFC-2in1-CN (Grefen & Blatt, 2012)
V2$_{L76S}$-nYFP + cYFP-*NbAGO4-2*	pDONR221-P3P2-V2$_{L76S}$ (This study) + pDONR221-P1P4-*NbAGO4-2* (This study)	pBiFC-2in1-CN (Grefen & Blatt, 2012)

Supplementary table 3. List of primers used for qPCR or qRT-PCR in this study.

Target gene	Forward Primer (5'-3') Reverse Primer (5'-3')	Primer pair efficiency	Reference
(1) RT-qPCR			
NbAGO4-1	F: TGGTGAAGTGCAAACATCGG	106.7%	This study
	R: TTGCAATGAGACCAACGAGC		
NbAGO4-2	F: GCCTTGCATCAATGTTGGGA	101.1%	This study
	R: TGAGGGCATTGCTTAGGACT		
NbAGO6	F: GCTAGTGCACCGGAAAATGT	88.6%	This study
	R: ATAGAGGTCGCAGTCGTGCT		
NbTubulin	F: CAAGATGCTACTGCAGACGAG	99.7%	Liu et al., 2012
	R: CTGGAAGTTGTGGTTTTGGC		
Nbcoilin	F: TGTGCCCAAGGAAGGTCTTGTG	96.6%	This study
	R: GGCATCAGCAGAACTCTATTTGCTTCA		
SlAGO4a	F: TCAAATGGGACAGTGGATGA	95.9%	This study
	R: TTTCTCCTCGAGCTTTGGAA		
SlAGO4b	F: GGGTCAATGGATGAAATTCG	98.7%	This study
	R: GGGGCAATTGAGGAACAGTA		
SlAGO4c	F: TCAGGGATGGTGTCAGTGAA	103.9%	This study
	R: GGGCGACCACTTTTCATCTA		
SlAGO4d	F: AAGGCAAAGAGGATGCTGAA	95.8%	This study
	R: CAACGTTAATGCAGGGGAGT		
SlActin	F: CCTCAGCACATTCCAGCAG	92.1%	Exposito-Rodriguez et al., 2008
	R: CCACCAAACTTCTCCATCCC		
(2) qPCR for virus accumulation			
Rep	F: TGAGAACGTCGTGTCTTCCG	94.80%	Wang et al., 2017b
	R: TGACGTTGTACCACGCATCA		
ITS	F: ATAACCGCATCAGGTCTCCA	101.1%	Mason et al., 2008
	R: CCGAAGTTACGGATCCATTT		
(3) qPCR following ChIP in *Nicotiana benthamiana*			
IR	F: CGGTGTCCCTCAAAGCTCTA	88.1%	This study
	R: TAGCCATTAGGTGTCCAGGT		
V2	F: AATCATTTCCACGCCCGTCT	89.2%	This study
	R: GTACGTCCATGATCGTCGCT		
Actin	F: CGGAATCCACGAGACTACATAC	108.1%	Maimbo et al., 2010
	R: GGGAAGCCAAGATAGAGC		

Supplementary table 4. Values of IR methylation in independent experiments and replicates.

Percentage of methylated cytosines in the intergenic region (IR) of TYLCV in local infection assays with TYLCV or the V2 null mutant (Figure 4A).

%	3 dpi / TYLCV			3 dpi / TYLCV-V2null			9 dpi / TYLCV			9 dpi / TYLCV-V2null		
	1st	2nd	3rd	1st	2nd	3rd	1st	2nd	3rd	1st	2nd	3rd
CG	0.92	0	1.11	61.8	62.12	47.22	0	0.46	5.32	72	77.58	78.57
CHG	0	0	4	62.5	70	43.33	0	0	6.77	76.15	84.82	80
CHH	0	0.19	1.11	63.02	57.97	49.07	0.28	0.81	7.3	73.55	72.7	73.74
All	0.166	0.143	1.402	62.75	59.93	48.17	0.208	0.666	6.902	73.54	74.79	75.24

Percentage of methylated cytosines in the intergenic region (IR) of TYLCV in local infection assays with the V2 null mutant in *AGO4*-silenced or control plants (Figure 4B).

%	TYLCV-V2null / TRV-EV				TYLCV-V2null / TRV-*NbAGO4*			
	1st	2nd	3rd	4th	1st	2nd	3rd	4th
CG	59.52	55.73	58.78	45.61	46.75	11.56	34.72	34.05
CHG	64.28	50.64	58.79	47.69	48.33	13.33	36.42	34.34
CHH	61.11	52.8	55.55	45.26	48.14	11.21	33.33	32.02
All	61.142	53.113	56.451	45.573	47.916	11.489	33.892	32.622

Percentage of methylated cytosines in the intergenic region (IR) of TYLCV in systemic infection assays with TYLCV or the V2 null mutant in *AGO4*-silenced or control plants (Figure 4C).

%	TYLCV / TRV-EV				TYLCV / TRV-*NbAGO4*				TYLCV-V2null / TRV-EV				TYLCV-V2null / TRV-*NbAGO4*			
	1st	2nd	3rd	4th	1st	2nd	3rd	4th	1st	2nd	3rd	4th	1st	2nd	3rd	4th
CG	49.55	6.34	7.71	27.84	34.6	0	0	0	56.29	71.79	60.58	69.04	39.76	43.82	29.53	56.96
CHG	49.39	6.42	7.77	31.87	35.61	0.86	0	0	52.66	66.92	57.41	73.8	38.94	45.21	26.31	56.36
CHH	44.34	6.55	8.26	30.43	31.85	0.18	0.07	0.08	53.05	67.61	55.75	65.05	39.47	42.68	25.01	54.16
All	45.79	6.504	8.115	30.11	32.72	0.217	0.055	0.058	53.6	68.3	56.79	66.65	39.47	43.14	25.96	54.89

Percentage of methylated cytosines in the intergenic region (IR) of TYLCV in systemic infection assays with TYLCV or the V2 null mutant in *Nbcoilin*-silenced or control plants (Figure 7D).

%	TYLCV / TRV-EV			TYLCV / TRV-*Nbcoilin*			TYLCV-V2null / TRV-EV			TYLCV-V2null / TRV-*Nbcoilin*		
	1st	2nd	3rd	1st	2nd	3rd	1st	2nd	3rd	1st	2nd	3rd
CG	2.77	0	0	3.59	0.3	0.27	18.14	46.6	71.79	5.98	7.07	50.39
CHG	1.66	0	0.52	4.11	0.55	0	22.66	53.88	73.07	6.92	7.27	55
CHH	1.54	0.13	0.21	3.6	0.07	0.06	18.88	45.09	65.52	4.91	7.71	50.74
All	1.778	0.1	0.21	3.651	0.166	0.1	19.133	46.247	67.411	5.311	7.552	51.111