A virus-encoded protein suppresses methylation of the viral genome through its
interaction with AGO4 in the Cajal body

Liping Wang[1,2], Yi Ding[1,2], Li He[1], Guiping Zhang[1], Jian-Kang Zhu[1], Rosa Lozano-Duran[1]

[1]Shanghai Center for Plant Stress Biology, Chinese Academy of Sciences, Shanghai 201602,
China; Center for Excellence in Molecular Plant Science, Chinese Academy of Sciences.
[2]University of the Chinese Academy of Sciences, Beijing 100049, China

ABSTRACT

DNA methylation is a eukaryotic anti-viral defence mechanism. In plants, establishment of *de*
novo DNA methylation is regulated by the RNA-directed DNA methylation pathway, which
requires AGO4 function. The genome of the plant DNA viruses geminiviruses replicates in the
nuclei of infected cells through not yet fully understood mechanisms and is subjected to
methylation, a modification that negatively impacts infectivity. In *Tomato yellow leaf curl virus*, the
virus-encoded V2 protein suppresses methylation of the viral DNA. Here, we identify AGO4 as a
physical interactor of V2. AGO4 mediates methylation of the viral genome, which is countered by
V2. Accordingly, virulence of a V2 mutant virus is partially restored by *AGO4* silencing, hinting at
the inhibition of AGO4 as a crucial virulence function of V2. Virus-produced V2 does not affect
accumulation of viral small interfering RNA nor prevents their loading into AGO4, but impairs
binding of this protein to the viral DNA. Importantly, the association between V2 and AGO4 occurs
in the Cajal body, uncovering this subnuclear compartment as a crucial site in the viral cycle.

Running title: V2 inhibits AGO4 in the Cajal body

Keywords: Geminivirus, AGO4, Cajal body, DNA methylation, virus.

INTRODUCTION

DNA methylation in cytosine residues is a conserved epigenetic mark essential for protecting the eukaryotic genome against invading nucleic acids, namely viruses and transposable elements. In plants, establishment of *de novo* DNA methylation is believed to be regulated by the RNA-directed DNA methylation (RdDM) pathway. The canonical RdDM pathway requires the activity of two plant-specific RNA polymerase II-related enzymes, Pol IV and Pol V, and leads to cytosine methylation in a sequence-specific manner. In brief, the current understanding of RdDM is as follows: Pol IV generates RNA transcripts subsequently converted to double-stranded RNA (dsRNA) by RDR2 (Haag et al., 2012, Law et al., 2011), and then diced into 24-nt siRNAs by DCL3 (Xie et al., 2004); the resulting 24-nt siRNAs are loaded into AGO4 (Zilberman et al., 2003), which is guided to scaffold RNA molecules generated by Pol V via sequence complementarity and recruits the *de novo* methyl transferase DRM2 (Böhmdorfer et al., 2014, Chan et al., 2005, Gao et al., 2010, Zhong et al., 2014), which in turn catalyses methylation of adjacent DNA sequences. RdDM generally creates a chromatin environment refractive to gene expression.

Geminiviruses are a family of plant viruses with circular single-stranded (ss) DNA genomes infecting multiple crops and causing dramatic yield losses worldwide. The geminiviral genome replicates in the nucleus of the infected cell by using the host DNA replication machinery, made available following the viral re-programming of the cell cycle (reviewed in Hanley-Bowdoin et al., 2013). During viral multiplication, the ssDNA genome generates a double-stranded (ds) DNA intermediate, which then undergoes rolling-circle replication (reviewed in Hanley-Bowdoin et al., 2013). However, the cellular and molecular details underlying these essential initial steps of the viral infection cycle, including the subnuclear localization of viral ss and ds DNA accumulation and of the processes leading to their production, are to date mostly unknown.

Notably, the geminiviral genome forms minichromosomes and is subjected to epigenetic modifications, including cytosine DNA methylation and histone modifications (Ceniceros-Ojeda et al., 2016, Deuschle et al., 2016, Jackel et al., 2016, Kushwaha et al., 2017, Raja et al., 2008, Wang et al., 2018). The findings that methylation of viral DNA negatively impacts viral replication (Brough et al., 1992, Ermak et al., 1993), that different geminivirus-encoded proteins have evolved to suppress DNA methylation (Buchmann et al., 2009, Ismayil et al., 2018, Raja et al., 2008, Rodriguez-Negrete et al., 2013, Tu et al., 2017, Wang et al., 2018, Yang et al., 2013, Yang et al., 2011, Zhang et al., 2011), and that methylation of the viral genome correlates with host resistance or recovery (Butterbach et al., 2014, Ceniceros-Ojeda et al., 2016, Raja et al., 2008, Torchetti et al., 2016, Yadav & Chattopadhyay, 2011) strongly support the idea that the plant-mediated

methylation of the viral DNA acts as an anti-viral defence mechanism, underscoring DNA
methylation as an active plant-virus battlefield.

During the infection by the geminivirus *Tomato yellow leaf curl virus* (TYLCV), the essential virus-
encoded protein V2 has been shown to suppress DNA methylation of the viral genome (Wang et
al., 2014). V2 interacts with the histone deacetylase HDA6 in *Nicotiana benthamiana*, competing
with the recruitment of the maintenance CG methyltransferase MET1 (Woo et al., 2008) and
ultimately reducing viral DNA methylation (Wang et al., 2018). However, silencing of *HDA6* results
in limited complementation of a V2 null mutation in the virus and in only a partial reduction in viral
DNA methylation (Wang et al., 2018), suggesting that V2 might counter methylation through
additional interactions with host factors.

In this work, we show that V2 from TYLCV interacts with the central RdDM component AGO4;
that AGO4 plays a role in anti-TYLCV defence; and that V2 and AGO4 have opposite effects on
the methylation of the viral genome. Our results indicate that the viral DNA gets methylated in the
absence of V2, and that increased methylation and reduced accumulation of a V2-defective virus
can be countered by silencing of *AGO4*. The presence of V2 does not affect production of viral
small interfering RNAs (vsiRNA) or prevent their loading into AGO4, but impairs binding of this
protein to the viral genome. Importantly, the physical interaction between V2 and AGO4 occurs
mostly in the Cajal body, unveiling this subnuclear compartment as a crucial site in the viral cycle
and in anti-viral DNA methylation. In summary, this work not only describes AGO4 as a target of
the viral suppressor of DNA methylation V2, but also hints at a functional relevance of subnuclear
compartmentalization of viral processes and components, and suggests a prominent role of the
Cajal body in the viral infection.

RESULTS

V2 from TYLCV interacts with AGO4 from *Nicotiana benthamiana* and tomato in the Cajal
body

With the aim of gaining insight into the functions of V2 from TYLCV in the plant cell, we used
transient expression of GFP-tagged V2 in infected leaf patches of *N. benthamiana* followed by
affinity purification and mass spectrometry (AP-MS) to identify plant interactors of this viral protein
in the context of the infection (Wang et al., 2017a). Interestingly, we identified the two AGO4
paralogs in *N. benthamiana* (NbAGO4-1 and NbAGO4-2) as associated to V2-GFP (Figure 1A;

Wang et al., 2017a); these interactions were confirmed by co-immunoprecipitation and split-
luciferase assays (Figure 1B,C).

There are four *AGO4* orthologues in tomato (*SlAGO4a-d*) (Bai et al., 2012), the natural and
economically relevant host of TYLCV (Figure 2A,B). All four SlAGO4-encoding genes are
expressed in basal conditions in tomato leaves, although *SlAGO4c* and *d* show low expression
levels; *SlAGO4b*, *c*, and *d* are slightly up-regulated by TYLCV infection (Supplementary figure 1).
SlAGO4a, *SlAGO4b*, and *SlAGO4d* were cloned and the encoded proteins confirmed as
interactors of V2 in co-IP and split-luciferase assays (Figure 2C,D).

In Arabidopsis, AGO4 has been shown to co-localize with its interactor NRPE1 (NRPD1b), a
subunit of Pol V, in the Cajal body, which was then suggested to be a center for the assembly of
AGO4/NRPE1/siRNA complexes, enabling RdDM at target loci (Li et al., 2008, Li et al., 2006).
Interestingly, both V2-GFP and the different RFP-AGO4 orthologues from *N. benthamiana* and
tomato co-localize in a distinct subnuclear compartment, identified as the Cajal body by the
accumulation of the nucleolus and Cajal body marker fibrillarin (Barneche et al., 2000), upon
transient expression in *N. benthamiana* (Figure 3A). Of note, most of nuclear V2-GFP
accumulates in the Cajal body, although some fluorescence can be detected in the nucleoplasm.
All AGO4 orthologues are distributed throughout the nucleoplasm and absent from the nucleolus;
clear Cajal body localization can be detected for NbAGO4-1, NbAGO4-2, SlAGO4a, and
SlAGO4b, while Cajal body localization of SlAGO4d is less conspicuous (Figure 3A). Analysis of
the V2/AGO4 interaction by bimolecular fluorescence complementation (BiFC), which is based
on visualization and hence provides spatial information, unveiled that, strikingly, the association
between these two proteins occurs mostly or exclusively in the Cajal body, where V2 homotypic
interactions also occur (Figure 3B).

116   V2 counters the AGO4-dependent methylation of the viral genome to promote virulence

In order to evaluate the contribution of V2 to the viral infection, we generated an infectious TYLCV
clone carrying a G-to-A mutation in the fifth nucleotide of the V2 open reading frame (ORF), which
converts the second codon (encoding tryptophan) to a stop codon (Supplementary figure 2A),
making it unable to produce the V2 protein (TYLCV-V2null). In agreement with previous results
(Wartig et al., 1997), V2 is required for full infectivity in both tomato and *N. benthamiana*, since
the V2 null mutant accumulates to very low levels and produces no noticeable symptoms
(Supplementary figure 2).

Next, we sought out to test whether knock-down of *AGO4* could partially complement the lack of
V2 during the TYLCV infection. For this purpose, we employed virus-induced gene silencing
(VIGS) to silence both *NbAGO4-1* and *NbAGO4-2*. VIGS efficiently knocked-down both *NbAGO4*
orthologues, but did not affect accumulation of the transcript of the close homologue *NbAGO6*
(Figure 4A); AGO4-silenced plants did not display any obvious developmental abnormalities
(Figure 4B). Expression of *NbAGO4-1* or *NbAGO4-2* was not affected by TYLCV infection, neither
in silenced nor in non-silenced plants (Figure 4C,D). Mutation in V2 does not affect viral replication
(Wartig et al., 1997), and therefore viral accumulation in local infections in *N. benthamiana* (leaf
patch agroinfiltration assays; see Supplementary figure 3) was not different between the WT virus
and the V2 null mutant virus (Figure 4E); in both cases, *AGO4* silencing led to a slight increase in
viral accumulation, suggesting an anti-viral role for AGO4 (Figure 4E). Interestingly, in systemic
infections (see Supplementary figure 3), *AGO4* silencing mildly increased viral accumulation of
the WT TYLCV (1.33-fold), but dramatically improved performance of the V2 null mutant virus
(3.23-fold), suggesting that one of the main roles of V2 during the viral infection is the suppression
of AGO4 function (Figure 4F).

In light of the role of AGO4 in RdDM and to directly assess the impact of V2 and AGO4 on the
methyl-state of the viral DNA, we used bisulfite sequencing (BS-seq) to measure DNA methylation
of the intergenic region of the viral genome, which presents the highest methylation levels during
the infection (Piedra-Aguilera et al., 2019). As shown in Figure 5A, cytosine methylation in this
region in all contexts (CG, CHG, and CHH) was almost undetectable in the WT viral genome in
local infections at 3 or 9 days post-inoculation (dpi), while it reached ~60% and ~80%, respectively,
in the V2 null mutant (Figure 5A; Supplementary figure 4). These results indicate that V2 can
prevent or revert methylation of the viral genome during the infection, which occurs rapidly in the
absence of this protein.

We then compared the percentage of cytosine methylation in the intergenic region in local
infections with the V2 null mutant TYLCV in basal conditions or upon *AGO4* silencing. Strikingly,
AGO4 silencing resulted in a ~23% decrease in the percentage of methylated cytosines in all
contexts (Figure 5B; Supplementary figure 5), indicating that knock-down of *AGO4* can partially
complement the lack of V2 at the level of the viral methylation state. This complementation
suggests that a) methylation of the viral DNA at least partially depends on AGO4 function; and b)
V2 can at least partially counter AGO4-dependent methylation of the viral DNA.

As opposed to short-timed local infections, in systemic infections in *N. benthamiana*, which require
longer timespans and involve viral cell-to-cell and long-distance movement, methylation of the

WT viral genome could be detected at ~15% decrease in all contexts (Figure 5C; Supplementary

figure 6). Interestingly, the methylation level tends to decrease upon *AGO4* silencing; this

reduction (~22%) is more prominent in the V2 null mutant genome, again supporting the idea that

AGO4-dependent methylation of the viral genome occurs during the infection and is partially

countered by V2. Notably, the detected decrease in methylation correlates with the enhanced

viral accumulation in the *AGO4*-silenced plants (Figure 4F).

164　V2 does not hamper production or loading of vsiRNA but interferes with AGO4 binding to

165　the viral genome

The canonical function of AGO4 in the RdDM pathway requires loading of siRNA and association

to Pol V, and results in the recruitment of DRM2 to the target loci and the subsequent methylation

of the adjacent DNA (Matzke et al., 2015, Matzke & Mosher, 2014). Through physical interaction,

V2 could affect AGO4 function on the viral genome in different ways, for example by impairing

loading of viral siRNA (vsiRNA) onto this protein or by displacing endogenous interactors, such

as Pol V or DRM2; our previous results demonstrate that V2 does not affect AGO4 accumulation

or localization (Figures 1 and 2). In order to shed light on the molecular mechanism underlying

the V2-mediated interference of AGO4-dependent methylation of the viral genome, we tested

binding of AGO4 to the viral DNA in the presence or absence of V2 in local infections with TYLCV

WT and V2 null mutant, respectively, by Chromatin immunoprecipitation (ChIP). As shown in

Figure 6A, 3xFLAG-NbAGO4 could bind both the intergenic (IR) and the V2-encoding region of

the viral genome in the absence of V2 (TYLCV-V2null), but the signal decreased to background

levels when V2 was present (TYLCV). Therefore, AGO4 has the capacity to bind the viral DNA

molecule, but this binding is impaired by the virus-encoded V2 protein. AGO4 binding in the

TYLCV V2 null mutant hence correlates with the detected increase in viral DNA methylation

(Figure 5A).

Several viral silencing suppressors encoded by different viruses have been shown to inhibit

formation of AGO/sRNA complexes (e.g. Burgyán et al., 2011, Rawlings et al., 2011, Schott et al.,

2012). To test whether this strategy is also employed by V2, we immunoprecipitated 3xFLAG-

NbAGO4 co-expressed with WT or V2 null mutant TYLCV in local infection assays in *N.*

benthamiana, and visualized AGO4-bound vsiRNA by sRNA northern blotting. While infected

samples contained both 21- and 24-nt vsiRNA, and the occurrence and accumulation of these

sRNA species was not affected by the presence of virus-encoded V2, mostly 24-nt vsiRNA co-

immunoprecipitated with AGO4 (Figure 6B). Interestingly, a higher amount of vsiRNA associated

to AGO4 in the samples infected with the WT virus (Figure 6B). Taken together, these results

demonstrate that V2 does not affect the production or accumulation of vsiRNA, nor does it hamper

loading of these vsiRNA molecules into AGO4, but interferes with binding of this protein to the

viral genome in order to suppress DNA methylation and promote virulence.

DISCUSSION

The plant DNA viruses geminiviruses and pararetroviruses are both targets and suppressors of

DNA methylation; this possibly extends to the third family of plant DNA viruses, nanoviruses,

although experimental evidence is lacking (reviewed in Poogin, 2013; Pumplin and Voinnet, 2013).

The independent evolution of viral suppressors of DNA methylation argues for an anti-viral effect

of this epigenetic modification. Indeed, seminal experiments by Brough et al. (1992) and Ermak

et al. (1993) demonstrated that methylation of the geminivirus genome interferes with its

replication in transformed protoplasts, likely due to a dual effect on viral gene expression and

function of the replication complex. More specifically, RdDM seems to play a prominent role in

plant defence against geminiviruses, since RdDM mutants or silenced plants display increased

susceptibility to geminivirus infection (Raja et al., 2008, Zhong et al., 2017), and DNA methylation

and repressive histone marks typical of RdDM are deposited on the viral genome (Castillo-

Gonzalez et al., 2015, Ceniceros-Ojeda et al., 2016, Coursey et al., 2018, Dogar et al., 2006,

Jackel et al., 2016, Kushwaha et al., 2017, Wang et al., 2018).

AGO4 is a central component of the canonical RdDM pathway, and as such an obvious target for

viral inhibition. However, AGO4 also affects susceptibility to RNA viruses and viroids, and is

targeted by proteins encoded by RNA viruses, which raises the idea that either RdDM on the host

genome plays a role in modulating plant-virus interactions broadly, or AGO4 has an anti-viral role

beyond RdDM (Brosseau et al., 2016, Ma et al., 2015, Minoia et al., 2014). Supporting the latter,

AGO4-dependent defences against a potexvirus are independent of other RdDM components

and do not require nuclear localization of AGO4 (Brosseau et al., 2016).

The geminivirus TYLCV encodes the essential, multifunctional V2 protein, which acts as a

suppressor of viral DNA methylation (Wang et al., 2014). Here, we show that V2 binds to the plant

AGO4 in the Cajal body, and suppresses the AGO4-dependent methylation of the viral genome

(Figures 1,2,3,5) and the AGO4-mediated restriction of viral accumulation (Figure 4E,F). Our

results indicate that AGO4-dependent methylation of viral DNA occurs quickly in the absence of

V2 (Figure 5A,B). Nevertheless, *AGO4* silencing still has a detectable, if minor, positive impact

on the accumulation of the WT virus, which correlates with decreased viral DNA methylation

(Figure 4E,F; Figure 5C), suggesting that the V2-mediated suppression of AGO4 function is not

complete. On the other hand, WT levels of viral DNA methylation are not restored in the V2 mutant

upon *AGO4* silencing, which raises the idea that AGO4 might not be the only methylation-related

target of V2. In agreement with this, V2 has been shown to bind HDA6 and interfere with its

promotion of MET1-dependent methylation of the viral DNA (Wang et al., 2018).

Recently, V2 encoded by the geminivirus *Cotton leaf curl Multan virus* (CLCuMV) was found to

interact with AGO4 in *N. benthamiana*, leading to enhanced viral accumulation and a reduction in

viral DNA methylation (Wang et al., 2019). It should be noted that the V2 proteins encoded by

CLCuMV and TYLCV are only 65% identical (Supplementary figure 7A), and they might have

evolved independently to target AGO4. Supporting this notion, mutation of a conserved residue,

L76, abolishes the interaction between CLCuMV V2 and NbAGO4 (Wang et al., 2019), but does

not affect the interaction between TYLCV V2 and NbAGO4 or SlAGO4, which still occurs in the

Cajal body (Supplementary figure 7). This mutation, however, negatively affects V2 self-

interaction in the Cajal body (Supplementary figure 7).

The finding that the physical association between TYLCV V2 and AGO4 takes places in a specific

nuclear body, the Cajal body, and has an impact on the methyl-state of the viral population in the

cell, suggests that all or most viral DNA molecules must localize in this subnuclear compartment

at some point of the viral cycle. This observation hints at a functional role of the Cajal body during

the infection; whether such a role is linked to gene expression, DNA replication, or some other

process remains to be investigated. Interestingly, the Cajal body has been connected to systemic

infection of plant RNA viruses, and proteins encoded by RNA viruses can bind coilin, the signature

protein of this compartment, which impacts plant-virus interactions (Kim et al., 2007a, Kim et al.,

2007b, Semashko et al., 2012, Shaw et al., 2014).

Based on our results, we propose a scenario in which V2 is required to interfere with AGO4

binding to the viral genome, impairing DNA methylation and promoting viral accumulation;

whether this effect is linked to the canonical RdDM pathway will require further investigation. In

the context of the arms race between host and virus, TYLCV has evolved V2 to target AGO4,

impairing its association to the viral DNA and hence suppressing methylation of the viral genome

and promoting virulence (Supplementary figure 8). Although the exact molecular mechanism

underlying this function of V2 is at present unknown, we hypothesize that the virus-encoded

protein might mask a surface required for AGO4 recruitment to the viral DNA through the

association with an endogenous interactor (e.g. NRPE1; Li et al., 2006), or interfere with the complementarity-based pairing to the nascent Pol V transcript. Further work will be necessary to fully elucidate the connections between the Cajal body, DNA methylation, and the geminiviral infection.

MATERIALS AND METHODS

Plasmids and cloning

To generate binary vectors to express *AGO4* from *N. benthamiana* and tomato (cv. Money maker), the full-length coding sequence of *AGO4* genes was amplified using cDNA as template. *NbAGO4-1*, *NbAGO4-2*, *SlAGO4a*, *SlAGO4b*, *SlAGO4d*, and *SlWRKY75* were cloned into pENTR-D/TOPO (Invitrogen) following the manufacturer's instructions. The binary plasmids to express RFP and 3xHA N-terminal fusions were generated by Gateway cloning the AGO4 coding sequence into pGWB555 and pGWB515, respectively (Nakagawa et al., 2007). V2-GFP and the TYLCV infectious clone have been previously described (Rosas-Diaz et al., 2018, Wang et al., 2017a). The V2 null TYLCV mutant was generated with the QuickChange Lightning Site-Directed Mutagenesis Kit (Agilent Technologies, Cat #210518) using the WT infectious clone as template. For the plasmids used for biomolecular fluorescence complementation (BiFC), *NbAGO4*, *SlAGO4*, V2/V2$_{L76S}$ were cloned into entry vector pDONR221-P1P4/pDONR221-P3P2 (Invitrogen) and then Gateway-cloned into the pBiFC-2in1-CN vector (Grefen & Blatt, 2012) as shown in Supplementary table 2. The binary plasmids for split-luciferase complementation imaging assay were generated by Gateway cloning the *NbAGO4, SlAGO4, SlWRKY75* or V2 from pENTR-D/TOPO entry vector into pGWB-N-luc and pGWB-C-luc (Yu et al., 2019). To generate pCAMBIA1300-3xFLAG-NbAGO4-1, pCAMBIA1300 was digested with *Xba*I and *Xma*I, and then the 3xFLAG and NbAGO4-1 coding sequences were amplified by PCR and Infusion-cloned into pCAMBIA1300 with ClonExpress® MultiS One Step Cloning Kit (Vazyme). pDONR207-*Fibrillarin* (Kim et al., 2007b) was used to Gateway-clone *Fibrillarin* to pGWB545 (Nakagawa et al., 2007). All primers and plasmids used for cloning are summarized in Supplementary tables 1 and 2, respectively.

Plant materials and growth conditions

N. benthamiana and tomato plants (cv. Money maker) were grown in a controlled growth chamber under long day conditions (LD, 16 h light/8 h dark) at 25℃.

Agrobacterium-mediated transient gene expression in *N. benthamiana*

All binary plasmids were transformed into *Agrobacterium tumefaciens* strain GV3101, with the exception of pBINTRA6, which was transformed into *A. tumefaciens* strain C58c1. *A. tumefaciens* clones carrying the constructs of interest were liquid-cultured in LB with appropriate antibiotics at 28℃ overnight. Bacterial cultures were then centrifuged at 4,000 g for 10 min and resuspended in the infiltration buffer (10 mM $MgCl_2$, 10 mM MES pH 5.6, 150 µM acetosyringone) and adjusted to an OD_{600} = 0.5. Next, the bacterial suspensions were incubated in the buffer at room temperature and in the dark for 2-4 hours and then infiltrated 3-4-week-old *N. benthamiana* plants. For co-expression experiments, the different agrobacterium suspensions were mixed at 1:1 ratio before infiltration.

Protein extraction and immunoprecipitation assays

Protein extraction and co-immunoprecipitation assays were performed as described in Wang et al., 2017a. Protein extracts were immunoprecipitated with GFP-Trap beads (Chromotek, Germany), and analyzed by western blot with anti-GFP (Abiocode, M0802-3a) and anti-HA (12CA5) (Roche, Cat. No. 11 583 816 001) antibodies.

For 3xFLAG-NbAGO4 IP followed by vsiRNA extraction, 6 grams of *N. benthamiana* leaves transiently expressing 3xFLAG-NbAGO4 were collected, ground in liquid nitrogen and homogenized in 6x (w:v) extraction buffer (20mM Tris HCl pH7.5, 25mM $MgCl_2$, 300mM NaCl, 5mM DTT, 0.5% NP-40, 1x complete™ Protease Inhibitor Cocktail (Roche)) at 4°C with rotation for 30 minutes. The extract was subjected to centrifugation (14,000 rpm, 25min) at 4°C. 5 µg anti-FLAG antibody (Sigma, F3165) per gram of tissue were added to the supernatant in a new tube and incubated at 4°C overnight. The next day, 20 µl of slurry Protein G beads (Invitrogen) per gram of tissue were added and subjected to a further incubation for 2 hours with rotation at 4°C. After incubation, Protein G beads were washed three times in 3x (v:v) homogenate wash buffer (20mM Tris pH7.5, 25mM $MgCl_2$, 300mM NaCl, 0.5% NP-40). The quality of purification was examined by SDS-PAGE followed by immunoblotting.

Split-luciferase complementation imaging assay

Split-luciferase complementation imaging assays were performed as described (Chen et al., 2008). Equal volumes of *A. tumefaciens* harboring V2-N-luc or C-luc-NbAGO4-1/2, C-luc-SlAGO4a/b, or C-luc-SlWRKY75 at OD_{600}=0.5 were mixed at 1:1 ratio. Three different combinations of *A. tumefaciens* were infiltrated on the same *N. benthamiana* leaf. 1mM luciferin

316 (in H$_2$O) was infiltrated into the inoculated leaves 2 days after agrobacterium infiltration. A low-
317 light cooled CCD imaging apparatus (NightShade LB985 with IndiGO software) was used to
318 capture and analyse the luciferase signal at 2 dpi.

Confocal imaging

320 Confocal imaging for co-localization of V2-GFP, RFP-AGO4, and CFP-Fibrillarin upon transient
321 expression in *N. benthamiana* epidermal cells was performed on a Leica TCS SP8 point scanning
322 confocal microscope using the pre-set sequential scan settings for GFP (Ex:488 nm, Em:500–
323 550 nm), RFP (Ex:561 nm, Em:600–650 nm), and CFP (Ex:442 nm, Em:452–482 nm).

Bimolecular Fluorescence Complementation

325 For bimolecular fluorescence complementation (BiFC) analyses, *A. tumefaciens* clones carrying
326 pBiFC-2in1-CN binary constructs (Grefen and Blat, 2012) and CFP-Fibrillarin were mixed at 1:1
327 ratio and infiltrated on 3-4-week-old *N. benthamiana* plants. Imaging was performed 2 days later
328 under a Leica TCS SP8 confocal microscope by using the pre-set sequential scan settings for
329 YFP (Ex: 514 nm, Em: 525–575 nm) and for CFP (Ex:442 nm, Em:452–500 nm).

Virus-induced gene silencing

The vectors used for virus-induced gene silencing (VIGS) were pBINTRA6 (Ratcliff et al., 2001)
and pTRV2-GW (Taylor et al., 2012). A 362-bp fragment of *NbAGO4-1* cDNA (from nt 1920 to
2281) was amplified using primers shown in Supplementary table 1, cloned into pENTR/D-TOPO
(Invitrogen), and subcloned into pTRV2-GW through an LR reaction (Invitrogen) to yield TRV-
NbAGO4. VIGS assays were performed as described in Lozano-Duran et al., 2011. For TYLCV
local infection assays, *A. tumefaciens* carrying pBINTRA6 and TRV-*NbAGO4* or TRV-EV were
mixed and inoculated into 18-day-old *N. benthamiana* plants. Two weeks later, fully expanded
young leaves were infiltrated with *A. tumefaciens* carrying the TYLCV infectious clone and
samples were collected at 3, 4 or 9 days post-inoculation (dpi) to detect viral accumulation. For
TYLCV systemic infection assays, *A. tumefaciens* carrying pBINTRA6 and TRV-*NbAGO4* or TRV-
EV (empty vector) and the TYLCV infectious clone were mixed and inoculated on 18-day-old *N.*
benthamiana plants. The three most apical leaves of each plant were collected at 3 weeks post-
inoculation (wpi) to detect viral accumulation.

Quantitative PCR (qPCR) and Reverse Transcription PCR (RT-qPCR)

345 To determine viral accumulation, total DNA was extracted from *N. benthamiana* leaves (from
346 infiltrated leaves in local infection assays and from apical leaves in systemic infection assays)

using the CTAB method (Minas et al., 2011). Quantitative PCR (qPCR) was performed with primers to amplify Rep (Wang et al., 2017b). As internal reference for DNA detection, the 25S ribosomal DNA interspacer (ITS) was used (Mason et al., 2008). To detect *NbAGO4-1*, *NbAGO4-2*, and *NbAGO6* transcripts, total RNA was extracted from *N. benthamiana* leaves by using Plant RNA kit (OMEGA Bio-tek # R6827). RNA was reverse-transcribed into cDNA by using the iScriptTM cDNA Synthesis Kit (Bio-Rad #1708890) according to the manufacturer's instructions. *NbTubulin* was used as reference gene (Liu et al., 2012). Relative expression was calculated by the comparative Ct method ($2^{-\Delta\Delta Ct}$). qPCR and RT-qPCR were performed in a BioRad CFX96 real-time system as described previously (Wang et al., 2017b). Total RNA was extracted from the leaves of tomato plants mock-inoculated or infected with TYLCV at 3 weeks post-inoculation (wpi). *SlActin was used as* reference gene (Exposito-Rodriguez et al., 2008). Similarly, RT-qPCR was performed on RNA extracted from tomato to detect the expression of *SlAGO4a/d/c/d*. All primers used for qPCR and qRT-PCR are listed in Supplementary Table 3.

DNA bisulfite sequencing analysis

DNA from virus-infected plant tissues was extracted by Dneasy Plant Mini Kit (QIAGEN, Cat. No. 69104), and 500 ng of purified DNA was subjected to bisulfite treatment using EpiTect Plus DNA Bisulfite Kit (QIAGEN, Cat. No. 59124) according to the manufacturer's handbook. The selected fragment (viral IR) of the bisulfite-treated DNA was amplified by PCR (Fw: TTTGATGTATTTTTTATTTGTTGGGGTTT, Rv: CCCTTACAACARATATAARATCCCT); amplified fragments were cloned into the pMD18-T vector by TA ligation and sequenced (>15 clones per experiment). Cytosine methylation analysis was performed with Kismeth (http://katahdin.mssm.edu/kismeth/revpage.pl) (Gruntman et al., 2008).

Chromatin immunoprecipitation (ChIP) assay

The agrobacterium clone carrying the binary vector to express 3xFLAG-NbAGO4-1 was co-infiltrated with those carrying the TYLCV or TYLCV-V2null infectious clones in *N. benthamiana* leaves, and tissues were collected at 2 dpi. Chromatin immunoprecipitation (ChIP) assays were performed as described (He et al., 2018). In brief, the cross-linking of 2 grams of leaves was performed with 1% formaldehyde in 1xPBS buffer and stopped with 1/15 volume of 2 M glycine by vacuum infiltration. Then the tissue was ground to powder and resuspended in HB buffer (2.5% Ficoll 400, 5% Dextran T40, 0.4 M Sucrose, 25 mM Tris pH 7.4, 10 mM MgCl2, 0.035% β-mercaptoethanol, 1% Protease Inhibitor Cocktail (Sigma)), homogenized and filtered through Miracloth (Milli-pore). Triton x-100 was added to the supernatant until final concentration was

0.5%. After spinning at 2,000×g for 20 min at 4°C, the pellet was re-suspended in HB buffer
containing 0.1% Triton x-100 and spun at 2,000×g for 10 min at 4°C. Isolated nuclei were re-
suspended in 500 µl of Nuclei Lysis buffer and sonicated by BioruptorTM UCD-200 sonicator
(diagenode) for 30 min. Following centrifugation at 21,130×g for 5 min at 4°C, the supernatant
was separated and used for input and immunoprecipitation. After adding 9 volume of ChIP dilution
buffer to the supernatant, this was pre-cleared with 10 µl of Dynabeads Protein G (Invirogen) for
1 h at 4°C. After removing the beads from the mixture, the supernatant was incubated with anti-
FLAG antibody (Sigma, F3165), or anti-IgG antibody (Sigma, I5006) overnight at 4°C. The
following day, after adding 20 µl of Dynabeads Protein G, the mixture was incubated for 2 h at
4°C. Beads were sequentially washed with 1 ml of the following buffers: Low Salt Wash buffer
(150 mM NaCl, 0.1% SDS, 1% Triton x-100, 2 mM EDTA, 20 mM Tris pH 8.0), High Salt Wash
buffer (500 mM NaCl, 0.1% SDS, 1% Triton x-100, 2 mM EDTA, 20 mM Tris pH 8.0), LiCl wash
buffer (250 mM LiCl, 1% Igepal, 1% Sodium Deoxycholate, 1 mM EDTA, 10 mM Tris pH 8.0), TE
buffer (10 mM Tris pH 8.0, 1 mM EDTA). Immunocomplexes were eluted with 250 µl of Elution
buffer (1% SDS, 0.1 M NaHCO$_3$) at 65°C for 15 min. After reverse crosslinking, 10 µl of 0.5 M
EDTA, 20 µl of 1 M Tris pH 6.5 and 1 µl of proteinase K (Invitrogen) were added to each sample,
which was incubated at 45 °C for 2 h. DNA was then purified using QIAquick PCR Purification Kit
(QIAGEN, Cat. No. 28106). The products were eluted into 200 µl of ddH$_2$O, and analysed by
qPCR. The primers used in this experiment are listed in Supplementary Table 3; the primers for
Actin are taken from Maimbo et al., 2010.

Small RNA (sRNA) extraction and northern blot analysis

Small RNA (sRNA) extraction and northern blot was performed as described (Yang et al., 2015).
Briefly, sRNAs were purified from total extracts or AGO4 immunoprecipitates and subjected to
northern blot analysis. For each sample, sRNAs were separated on a 17% polyacrylamide gel,
which was electrotransferred to a Hybond N+ membrane (GE Lifesciences). Membranes were
cross-linked, incubated for 2 hours at 65°C, and hybridized overnight at 38°C with [32]P-labeled
probes for the intergenic region (IR) of the viral genome amplified by PCR (Fw:
TCCTCTTTAGAGAGAGAACAATTGGGA,   Rv: ACAACGAAATCCGTGAACAG)   or
oligonucleotides in PerfectHyb buffer (Sigma). Washed membranes were exposed to X-ray films
at −80 °C for 3 days.

ACKNOWLEDGEMENTS

The authors thank Xinyu Jian, Aurora Luque, and Yujing (Ada) Liu for technical assistance, Gang Yu and Alberto Macho for sharing materials, and all members in Rosa Lozano-Duran's and Alberto Macho's groups for stimulating discussions and helpful suggestions. The authors are grateful to Alberto Macho, Huang Tan, and Tamara Jimenez-Gongora for critical reading of the manuscript. This research was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No. XDB27040206), and by the National Natural Science Foundation of China (NSFC) (grant numbers 31671994 and 31870250). Research in RL-D's lab is funded by the Shanghai Center for Plant Stress Biology of the Chinese Academy of Sciences and the 100 Talent Program of the Chinese Academy of Sciences.

AUTHOR CONTRIBUTIONS

RL-D and LW conceived the project; LW, DY, LH, and GZ performed experiments and analyzed results; all authors intellectually contributed to the project; RL-D wrote the manuscript, with input from all authors.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

Böhmdorfer G, Rowley MJ, Kuciński J, Zhu Y, Amies I, Wierzbicki AT (2014) RNA-directed DNA methylation requires stepwise binding of silencing factors to long non-coding RNA. *The Plant Journal* 79: 181-191

Bai M, Yang GS, Chen WT, Mao ZC, Kang HX, Chen GH, Yang YH, Xie BY (2012) Genome-wide identification of Dicer-like, Argonaute and RNA-dependent RNA polymerase gene families and their expression analyses in response to viral infection and abiotic stresses in Solanum lycopersicum. *Gene* 501: 52-62

Barneche F, Steinmetz F, Echeverria M (2000) Fibrillarin genes encode both a conserved nucleolar protein and a novel small nucleolar RNA involved in ribosomal RNA methylation in Arabidopsis thaliana. *The Journal of biological chemistry* 275: 27212-20

Brosseau C, El Oirdi M, Adurogbangba A, Ma X, Moffett P (2016) Antiviral Defense Involves AGO4 in an Arabidopsis-Potexvirus Interaction. *Molecular plant-microbe interactions : MPMI* 29: 878-888

Brough CL, Gardiner WE, Inamdar NM, Zhang XY, Ehrlich M, Bisaro DM (1992) DNA methylation inhibits propagation of tomato golden mosaic virus DNA in transfected protoplasts. *Plant Mol Biol* 18: 703-12

Buchmann RC, Asad S, Wolf JN, Mohannath G, Bisaro DM (2009) Geminivirus AL2 and L2 proteins suppress transcriptional gene silencing and cause genome-wide reductions in cytosine methylation. *Journal of virology* 83: 5005-13

Burgyán J, Havelda Z (2011) Viral suppressors of RNA silencing. *Trends Plant Science* 16(5):265-72.

Butterbach P, Verlaan MG, Dullemans A, Lohuis D, Visser RG, Bai Y, Kormelink R (2014) Tomato yellow leaf curl virus resistance by Ty-1 involves increased cytosine methylation of viral genomes and is compromised by cucumber mosaic virus infection. *Proceedings of the National Academy of Sciences of the United States of America* 111: 12942-7

Castillo-Gonzalez C, Liu X, Huang C, Zhao C, Ma Z, Hu T, Sun F, Zhou Y, Zhou X, Wang XJ,

Zhang X (2015) Geminivirus-encoded TrAP suppressor inhibits the histone methyltransferase

SUVH4/KYP to counter host defense. *eLife* 4

Ceniceros-Ojeda EA, Rodriguez-Negrete EA, Rivera-Bustamante RF (2016) Two Populations of

Viral Minichromosomes Are Present in a Geminivirus-Infected Plant Showing Symptom

Remission (Recovery). *Journal of virology* 90: 3828-3838

Chan SW, Henderson IR, Jacobsen SE (2005) Gardening the genome: DNA methylation in

Arabidopsis thaliana. *Nature reviews Genetics* 6: 351-60

Chen H, Zou Y, Shang Y, Lin H, Wang Y, Cai R, Tang X, Zhou JM (2008) Firefly luciferase

complementation imaging assay for protein-protein interactions in plants. *Plant physiology* 146:

368-76

Coursey T, Regedanz E, Bisaro DM (2018) Arabidopsis RNA Polymerase V Mediates Enhanced

Compaction and Silencing of Geminivirus and Transposon Chromatin during Host Recovery from

Infection. 92

Dereeper A, Audic S, Claverie JM, Blanc G (2010) BLAST-EXPLORER helps you building

datasets for phylogenetic analysis. *BMC evolutionary biology* 10: 8

Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, Dufayard JF, Guindon S, Lefort

V, Lescot M, Claverie JM, Gascuel O (2008) Phylogeny.fr: robust phylogenetic analysis for the

non-specialist. *Nucleic acids research* 36: W465-9

Deuschle K, Kepp G, Jeske H (2016) Differential methylation of the circular DNA in geminiviral

minichromosomes. *Virology* 499: 243-258

Dogar AM (2006) RNAi dependent epigenetic marks on a geminivirus promoter. *Virology journal*

3: 5

Ermak G, Paszkowski U, Wohlmuth M, Mittelsten Scheid O, Paszkowski J (1993) Cytosine
methylation inhibits replication of African cassava mosaic virus by two distinct mechanisms.
Nucleic acids research 21: 3445-50

Exposito-Rodriguez M, Borges AA, Borges-Perez A, Perez JA (2008) Selection of internal control
genes for quantitative real-time RT-PCR studies during tomato development process. *BMC plant*
biology 8: 131

Gao Z, Liu H-L, Daxinger L, Pontes O, He X, Qian W, Lin H, Xie M, Lorkovic ZJ, Zhang S (2010)
An RNA polymerase II-and AGO4-associated protein acts in RNA-directed DNA methylation.
Nature 465: 106

Grefen C, Blatt MR (2012) A 2in1 cloning system enables ratiometric bimolecular fluorescence
complementation (rBiFC). *BioTechniques* 53: 311-14

Gruntman E, Qi Y, Slotkin RK, Roeder T, Martienssen RA, Sachidanandam R (2008) Kismeth:
analyzer of plant methylation states through bisulfite sequencing. *BMC bioinformatics* 9: 371

Haag JR, Ream TS, Marasco M, Nicora CD, Norbeck AD, Pasa-Tolic L, Pikaard CS (2012) In
vitro transcription activities of Pol IV, Pol V, and RDR2 reveal coupling of Pol IV and RDR2 for
dsRNA synthesis in plant RNA silencing. *Molecular cell* 48: 811-818

Hanley-Bowdoin L, Bejarano ER, Robertson D, Mansoor S (2013) Geminiviruses: masters at
redirecting and reprogramming plant processes. *Nature reviews Microbiology* 11: 777-88

He L, Wu W, Zinta G, Yang L, Wang D, Liu R, Zhang H, Zheng Z, Huang H, Zhang Q, Zhu J-K
(2018) A naturally occurring epiallele associates with leaf senescence and local climate
adaptation in Arabidopsis accessions. *Nature communications* 9: 460

Ismayil A, Haxim Y, Wang Y, Li H, Qian L, Han T, Chen T, Jia Q, Yihao Liu A, Zhu S, Deng H,
Gorovits R, Hong Y, Hanley-Bowdoin L, Liu Y (2018) Cotton Leaf Curl Multan virus C4 protein
suppresses both transcriptional and post-transcriptional gene silencing by interacting with SAM
synthetase. *PLoS pathogens* 14: e1007282

Jackel JN, Storer JM, Coursey T, Bisaro DM (2016) Arabidopsis RNA Polymerases IV and V Are
Required To Establish H3K9 Methylation, but Not Cytosine Methylation, on Geminivirus
Chromatin. *Journal of virology* 90: 7529-40
536
537 Kim SH, Macfarlane S, Kalinina NO, Rakitina DV, Ryabov EV, Gillespie T, Haupt S, Brown JW,
538 Taliansky M (2007a) Interaction of a plant virus-encoded protein with the major nucleolar protein
539 fibrillarin is required for systemic virus infection. *Proceedings of the National Academy of Sciences*
540 *of the United States of America* 104: 11115-20
541
542 Kim SH, Ryabov EV, Kalinina NO, Rakitina DV, Gillespie T, MacFarlane S, Haupt S, Brown JW,
543 Taliansky M (2007b) Cajal bodies and the nucleolus are required for a plant virus systemic
544 infection. *The EMBO journal* 26: 2169-79
545
546 Kushwaha NK, Mansi, Chakraborty S (2017) The replication initiator protein of a geminivirus
547 interacts with host monoubiquitination machinery and stimulates transcription of the viral genome.
548 *PLoS pathogens* 13: e1006587
549
550 Law JA, Vashisht AA, Wohlschlegel JA, Jacobsen SE (2011) SHH1, a homeodomain protein
551 required for DNA methylation, as well as RDR2, RDM4, and chromatin remodeling factors,
552 associate with RNA polymerase IV. *PLoS genetics* 7: e1002195
553
554 Li CF, Henderson IR, Song L, Fedoroff N, Lagrange T, Jacobsen SE (2008) Dynamic regulation
555 of ARGONAUTE4 within multiple nuclear bodies in Arabidopsis thaliana. *PLoS genetics* 4: e27
556
557 Li CF, Pontes O, El-Shami M, Henderson IR, Bernatavichute YV, Chan SW, Lagrange T, Pikaard
558 CS, Jacobsen SE (2006) An ARGONAUTE4-containing nuclear processing center colocalized
559 with Cajal bodies in Arabidopsis thaliana. *Cell* 126: 93-106
560
561 Liu D, Shi L, Han C, Yu J, Li D, Zhang Y (2012) Validation of reference genes for gene expression
562 studies in virus-infected Nicotiana benthamiana using quantitative real-time PCR. *PloS one* 7:
563 e46451
564
Lozano-Duran R, Rosas-Diaz T, Luna AP, Bejarano ER (2011) Identification of host genes
involved in geminivirus infection using a reverse genetics approach. *PloS one* 6: e22383

Ma X, Nicole MC, Meteignier LV, Hong N, Wang G, Moffett P (2015) Different roles for RNA
silencing and RNA processing components in virus recovery and virus-induced gene silencing in
plants. *Journal of experimental botany* 66: 919-32

Maimbo M, Ohnishi K, Hikichi Y, Yoshioka H, Kiba A (2010) S-glycoprotein-like protein regulates
defense responses in Nicotiana plants against Ralstonia solanacearum. *Plant physiology* 152:
2023-35

Mason G, Caciagli P, Accotto GP, Noris E (2008) Real-time PCR for the quantitation of Tomato
yellow leaf curl Sardinia virus in tomato plants and in Bemisia tabaci. *Journal of virological*
methods 147: 282-9

Matzke MA, Kanno T, Matzke AJ (2015) RNA-Directed DNA Methylation: The Evolution of a
Complex Epigenetic Pathway in Flowering Plants. *Annual review of plant biology* 66: 243-67

Matzke MA, Mosher RA (2014) RNA-directed DNA methylation: an epigenetic pathway of
increasing complexity. *Nature reviews Genetics* 15: 394-408

Minas K, McEwan NR, Newbold CJ, Scott KP (2011) Optimization of a high-throughput CTAB-
based protocol for the extraction of qPCR-grade DNA from rumen fluid, plant and bacterial pure
cultures. *FEMS microbiology letters* 325: 162-9

Minoia S, Carbonell A, Di Serio F, Gisel A, Carrington JC, Navarro B, Flores R (2014) Specific
argonautes selectively bind small RNAs derived from potato spindle tuber viroid and attenuate
viroid accumulation in vivo. *Journal of virology* 88: 11933-45

Nakagawa T, Suzuki T, Murata S, Nakamura S, Hino T, Maeo K, Tabata R, Kawai T, Tanaka K,
Niwa Y, Watanabe Y, Nakamura K, Kimura T, Ishiguro S (2007) Improved Gateway binary vectors:
high-performance vectors for creation of fusion constructs in transgenic analysis of plants.
Bioscience, biotechnology, and biochemistry 71: 2095-100

Piedra-Aguilera A, Jiao C, Luna AP, Villanueva F, Dabad M, Esteve-Codina A, Diaz-Pendon JA,
Fei Z, Bejarano ER, Castillo AG (2019) Integrated single-base resolution maps of transcriptome,

sRNAome and methylome of Tomato yellow leaf curl virus (TYLCV) in tomato. *Scientific reports* 9: 2863

Raja P, Sanville BC, Buchmann RC, Bisaro DM (2008) Viral genome methylation as an epigenetic defense against geminiviruses. *Journal of virology* 82: 8997-9007

Ratcliff F, Martin-Hernandez AM, Baulcombe DC (2001) Technical Advance. Tobacco rattle virus as a vector for analysis of gene function by silencing. *The Plant journal : for cell and molecular biology* 25: 237-45

Rawlings RA, Krishnan V, Walter NG (2011) Viral RNAi suppressor reversibly binds siRNA to outcompete Dicer and RISC via multiple turnover. *Journal of Molecular Biology* 408(2):262-76.

Rodriguez-Negrete E, Lozano-Duran R, Piedra-Aguilera A, Cruzado L, Bejarano ER, Castillo AG (2013) Geminivirus Rep protein interferes with the plant DNA methylation machinery and suppresses transcriptional gene silencing. *The New phytologist* 199: 464-75

Rosas-Diaz T, Zhang D, Fan P, Wang L, Ding X, Jiang Y, Jimenez-Gongora T, Medina-Puche L, Zhao X, Feng Z, Zhang G, Liu X, Bejarano ER, Tan L, Zhang H, Zhu JK, Xing W, Faulkner C, Nagawa S, Lozano-Duran R (2018) A virus-targeted plant receptor-like kinase promotes cell-to-cell spread of RNAi. *Proceedings of the National Academy of Sciences of the United States of America* 115: 1388-1393

Schott G, Mari-Ordonez A, Himber C, Alioua A, Voinnet O, Dunoyer P (2012) Differential effects of viral silencing suppressors on siRNA and miRNA loading support the existence of two distinct cellular pools of ARGONAUTE1. *The EMBO journal* 31: 2553-65

Semashko MA, Rakitina DV, Gonzalez I, Canto T, Kalinina NO, Taliansky ME (2012) Movement protein of hordeivirus interacts in vitro and in vivo with coilin, a major structural protein of Cajal bodies. *Doklady Biochemistry and biophysics* 442: 57-60

Shaw J, Love AJ, Makarova SS, Kalinina NO, Harrison BD, Taliansky ME (2014) Coilin, the signature protein of Cajal bodies, differentially modulates the interactions of plants with viruses in widely different taxa. *Nucleus* 5: 85-94

Taylor KW, Kim JG, Su XB, Aakre CD, Roden JA, Adams CM, Mudgett MB (2012) Tomato TFT1

is required for PAMP-triggered immunity and mutations that prevent T3S effector XopN from

binding to TFT1 attenuate Xanthomonas virulence. *PLoS pathogens* 8: e1002768

Torchetti EM, Pegoraro M, Navarro B, Catoni M, Di Serio F, Noris E (2016) A nuclear-replicating

viroid antagonizes infectivity and accumulation of a geminivirus by upregulating methylation-

related genes and inducing hypermethylation of viral DNA. *Scientific reports* 6: 35101

Tu YC, Tsai WS, Wei JY, Chang KY, Tien CC, Hsiao HY, Fu SF (2017) The C2 protein of tomato

leaf curl Taiwan virus is a pathogenicity determinant that interferes with expression of host genes

encoding chromomethylases. *Physiologia plantarum* 161: 515-531

Wang B, Li F, Huang C, Yang X, Qian Y, Xie Y, Zhou X (2014) V2 of tomato yellow leaf curl virus

can suppress methylation-mediated transcriptional gene silencing in plants. *The Journal of*

general virology 95: 225-30

Wang B, Yang X, Wang Y, Xie Y, Zhou X (2018) Tomato Yellow Leaf Curl Virus V2 Interacts with

Host Histone Deacetylase 6 To Suppress Methylation-Mediated Transcriptional Gene Silencing

in Plants. *Journal of virology* 92

Wang L, Ding X, Xiao J, Jimenez-Gomicronngora T, Liu R, Lozano-Duran R (2017a) Inference of

a Geminivirus-Host Protein-Protein Interaction Network through Affinity Purification and Mass

Spectrometry Analysis. *Viruses* 9

Wang L, Tan H, Wu M, Jimenez-Gongora T, Tan L, Lozano-Duran R (2017b) Dynamic Virus-

Dependent Subnuclear Localization of the Capsid Protein from a Geminivirus. *Frontiers in plant*

science 8: 2165

Wang Y, Wu Y, Gong Q, Ismayil A, Yuan Y, Lian B, Jia Q, Han M, Deng H, Hong Y, Hanley-

Bowdoin L, Qi Y, Liu Y (2019) Geminiviral V2 Protein Suppresses Transcriptional Gene Silencing

through Interaction with AGO4. *Journal of virology* 93

Wartig L, Kheyr-Pour A, Noris E, De Kouchkovsky F, Jouanneau F, Gronenborn B, Jupin I (1997)
Genetic analysis of the monopartite tomato yellow leaf curl geminivirus: roles of V1, V2, and C2
ORFs in viral pathogenesis. *Virology* 228: 132-40

Woo HR, Dittmer TA, Richards EJ (2008) Three SRA-domain methylcytosine-binding proteins
cooperate to maintain global CpG methylation and epigenetic silencing in Arabidopsis. *PLoS*
genetics 4: e1000156

Xie Z, Johansen LK, Gustafson AM, Kasschau KD, Lellis AD, Zilberman D, Jacobsen SE,
Carrington JC (2004) Genetic and functional diversification of small RNA pathways in plants.
PLoS biology 2: e104

Yadav RK, Chattopadhyay D (2011) Enhanced viral intergenic region-specific short interfering
RNA accumulation and DNA methylation correlates with resistance against a geminivirus.
Molecular plant-microbe interactions : MPMI 24: 1189-97

Yang D-L, Zhang G, Tang K, Li J, Yang L, Huang H, Zhang H, Zhu J-K (2015) Dicer-independent
RNA-directed DNA methylation in Arabidopsis. *Cell research*

Yang LP, Fang YY, An CP, Dong L, Zhang ZH, Chen H, Xie Q, Guo HS (2013) C2-mediated
decrease in DNA methylation, accumulation of siRNAs, and increase in expression for genes
involved in defense pathways in plants infected with beet severe curly top virus. *The Plant journal :*
for cell and molecular biology 73: 910-7

Yang X, Xie Y, Raja P, Li S, Wolf JN, Shen Q, Bisaro DM, Zhou X (2011) Suppression of
methylation-mediated transcriptional gene silencing by betaC1-SAHH protein interaction during
geminivirus-betasatellite infection. *PLoS pathogens* 7: e1002329

Yu G, Xian L, Xue H, Yu W, Rufian J, Sang Y, Morcillo R, Wang Y, Macho AP (2019) A bacterial
effector protein prevents MAPK-mediated phosphorylation of SGT1 to suppress plant immunity.
bioRxiv: 641241
Zhang Z, Chen H, Huang X, Xia R, Zhao Q, Lai J, Teng K, Li Y, Liang L, Du Q, Zhou X, Guo H,
Xie Q (2011) BSCTV C2 attenuates the degradation of SAMDC1 to suppress DNA methylation-
mediated gene silencing in Arabidopsis. *The Plant cell* 23: 273-88

Zhong X, Du J, Hale CJ, Gallego-Bartolome J, Feng S, Vashisht AA, Chory J, Wohlschlegel JA,

Patel DJ, Jacobsen SE (2014) Molecular mechanism of action of plant DRM de novo DNA

methyltransferases. *Cell* 157: 1050-1060

Zhong X, Wang ZQ, Xiao R, Wang Y, Xie Y, Zhou X (2017) iTRAQ analysis of the tobacco leaf

proteome reveals that RNA-directed DNA methylation (RdDM) has important roles in defense

against geminivirus-betasatellite infection. *Journal of proteomics* 152: 88-101

Zilberman D, Cao X, Jacobsen SE (2003) ARGONAUTE4 control of locus-specific siRNA

accumulation and DNA and histone methylation. *Science (New York, NY)* 299: 716-719

FIGURE LEGENDS

Figure 1. V2 interacts with AGO4 from *N. benthamiana.*

A   Unique peptide count, protein coverage, and best Mascot Score of NbAGO4-1 and NbAGO4-2 co-immunoprecipitated with V2-GFP, as identified by affinity purification followed by mass spectrometry (AP-MS). Results from three independent biological repeats are shown. "-" indicates no peptide was detected.

B   3xHA-NbAGO4-1 and 3xHA-NbAGO4-2 specifically interact with V2-GFP in co-immunoprecipitation (co-IP) assays upon transient expression in *N. benthamiana*. Free GFP was used as negative control. CBB, Coomassie brilliant blue staining. Three independent biological replicates were performed with similar results.

C   NbAGO4-1 and NbAGO4-2 interact with V2 in split-luciferase assays. V2-N-luc and C-luc-NbAGO4-1/2 were transiently co-expressed in *N. benthamiana*; C-luc-SlWRKY75 is used as negative control. The luciferase bioluminescence from at least three independent leaves per experiment was imaged two days after infiltration. The average bioluminescence, measured in counts per second (cps), as well as an image of a representative leaf are shown. Values represent the mean of three independent biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant difference (according to a Student's *t*-test, **: $P<0.01$, ***: $P<0.001$.) compared to the negative control.

Figure 2. V2 interacts with AGO4 from tomato*.*

A   Phylogenetic tree of AtAGO4, NbAGO4, and SlAGO4 proteins. The phylogenetic analysis was performed with phylogeny.fr (Dereeper et al., 2010, Dereeper et al., 2008).

B   Pairwise identity and genetic distance matrix among AtAGO4, NbAGO4 and SlAGO4 proteins. The analysis was performed by Geneious (https://www.geneious.com).

C   3xHA-SlAGO4a, 3xHA-SlAGO4b, and 3xHA-SlAGO4d specifically interact with V2-GFP in co-immunoprecipitation (co-IP) assays upon transient expression in *N. benthamiana*. Free GFP was used as negative control. CBB, Coomassie brilliant blue staining. Three independent biological replicates were performed with similar results.

D   SlAGO4a and SlAGO4b interact with V2 in split-luciferase assays. V2-N-luc and C-luc-SlAGO4a/b were transiently co-expressed in *N. benthamiana*; C-luc-SlWRKY75 is used as

negative control. The luciferase bioluminescence from at least three independent leaves per
experiment was imaged two days after infiltration. The average bioluminescence, measured
in counts per second (cps), as well as an image of a representative leaf are shown. Values
represent the mean of three independent biological replicates; error bars indicate SEM.
Asterisks indicate a statistically significant difference (according to a Student's *t*-test, ***:
P<0.001.) compared to the negative control.

Figure 3. V2 interacts with AGO4 in the Cajal body.

A   V2-GFP and RFP-AGO4 co-localize in the Cajal body. CFP-Fibrillarin, V2-GFP and RFP-
NbAGO4-1/2 or RFP-SlAGO4a/b/d were transiently co-expressed in *N. benthamiana*
epidermal cells. CFP-Fibrillarin is used as a nucleolus and Cajal body marker. Confocal
images were taken at two days after infiltration . Arrowheads indicate the position of the Cajal
body. Bar, 5μm. This experiment was repeated more than three times with similar results.

B   V2 interacts with AGO4 in the Cajal body. The N-terminal half of the YFP fused to V2 (V2-
nYFP) was transiently co-expressed with the C-terminal half of the YFP alone (cYFP, as a
negative control), or cYFP-NbAGO4, cYFP-SlAGO4, or cYFP-V2 (as a positive control) in *N.*
benthamiana leaves. CFP-Fibrillarin was used as a nucleolus and Cajal body marker.
Confocal images were taken at two days after infiltration. Yellow fluorescence indicates a
positive interaction. Arrowheads indicate the position of the Cajal body. Bar, 5μm. This
experiment was repeated more than three times with similar results.

Figure 4. V2 counters the AGO4-dependent defence to promote virulence

A   Expression of *NbAGO4-1, NbAGO4-2,* and *NbAGO6* in *N. benthamiana* plants infected with
TRV-EV (empty vector) or TRV-*NbAGO4*, measured by reverse transcription quantitative real-
time PCR (RT-qPCR). Gene expression was normalized to *NbTubulin*. Values are the mean
of four independent biological replicates; error bars indicate SEM. Asterisks indicate a
statistically significant difference according to Student's *t*-test. **: P<0.01, ***: P<0.001, ns:
not significant.

B   Representative pictures of *N. benthamiana* plants infected with the indicated combinations of
viruses. Photographs were taken at 3 weeks post-inoculation (wpi).

C   *NbAGO4-1* expression in *NbAGO4*-silenced plants and control plants infected with TYLCV,
TYLCV-V2null, or mock-inoculated at 3 wpi measured by RT-qPCR. Gene expression was
normalized to *NbTubulin*. Values are the mean of six independent biological replicates; error
bars indicate SEM. Asterisks indicate a statistically significant difference according to
Student's *t*-test. *: P<0.05, **: P<0.01, ****: P<0.0001.

D   *NbAGO4-2* expression in *NbAGO4*-silenced plants and control plants infected by TYLCV,
TYLCV-V2null or mock-inoculated at 3 wpi measured by RT-qPCR. Gene expression was
normalized to *Tubulin*. Values are the mean of six independent biological replicates; error bars
indicate SEM. Asterisks indicate a statistically significant difference according to Student's *t*-
test. *: P<0.05, **: P<0.01, ***: P<0.001.

E   Viral (TYLCV) accumulation in local infections in *NbAGO4*-silenced or control plants,
measured by qPCR. Infiltrated leaf patches from different plants were collected at 4 dpi. The
experimental design is shown in Figure S3A. The accumulation of viral DNA is normalized to
the *25S ribosomal RNA interspacer (ITS)*. Values are the mean of eight independent biological
replicates; error bars indicate SEM.

F   Viral (TYLCV) accumulation in systemic infections in *NbAGO4*-silenced or control plants,
measured by qPCR. Apical leaves from six plants were collected at 3 wpi. The experimental
design is shown in Figure S3B. The accumulation of viral DNA is normalized to the *25S*
ribosomal RNA interspacer (ITS). Four independent biological replicates were performed with
similar results; one representative result is shown. Values are the mean of six independent
biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant
difference according to Student's *t*-test. *: P<0.05, ***: P<0.001. The relative fold change of
viral accumulation between *NbAGO4*-silenced plants and control plants is shown above each
column.

Figure 5. V2 suppresses the AGO4-dependent methylation of viral DNA.

A   Percentage of methylated cytosines in the intergenic region (IR) of TYLCV in local infection
assays with TYLCV WT or V2 null mutant (TYLCV-V2null) in *N. benthamiana* at 3 or 9 days
post-inoculation (dpi), as detected by bisulfite sequencing. The original single-base resolution
bisulfite sequencing data are shown in Figure S4. Values are the mean of three independent
biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant

difference according to Student's *t*-test. ***: P<0.001.

B   Percentage of methylated cytosines in the intergenic region (IR) of TYLCV in local infection
assays with the V2 null mutant TYLCV (TYLCV-V2null) in *AGO4*-silenced (TRV-*NbAGO4*) or
control (TRV-EV) *N. benthamiana* plants at 4 dpi, as detected by bisulfite sequencing.
Samples come from the same plants used in Figure 4E. The original single-base resolution
bisulfite sequencing data are shown in Figure S5. Values are the mean of four independent
biological replicates; error bars indicate SEM. Asterisks indicate a statistically significant
difference according to Student's *t*-test. *: P<0.05.

C   Percentage of methylated cytosines in the intergenic region (IR) of TYLCV in systemic
infection assays with TYLCV WT or V2 null mutant (TYLCV-V2null) in *AGO4*-silenced (TRV-
NbAGO4) or control (TRV-EV) *N. benthamiana* plants at 3 weeks post-inoculation (wpi), as
detected by bisulfite sequencing. Samples come from the same plants used in Figure 4F.
The original single-base resolution bisulfite sequencing data are shown in Figure S6. Values
are the mean of four independent biological replicates; error bars indicate SEM. Asterisks
indicate a statistically significant difference according to Student's *t*-test. *: P<0.05.

Figure 6. V2 interferes with AGO4 binding to the viral genome but does not hamper
production or loading of vsiRNA.

A   3xFLAG-NbAGO4-1 binds the viral (TYLCV) genome. Binding was detected by chromatin
immunoprecipitation (ChIP) upon transient expression in *N. benthamiana* followed by qPCR.
Two regions of the viral genome, the IR and the V2 ORF, were analyzed; *ACTIN* was used
as negative control. Values represent the mean of four independent biological replicates;
error bars represent SEM. Asterisks indicate a significant difference according to a Student's
t-test; the *P*-value for the different comparisons is shown. *: P<0.05.

B   NbAGO4-1 binds viral small interfering RNA (vsiRNA) independently of V2. Northern blot of
vsiRNA in total extracts or 3xFLAG-NbAGO4 immunoprecipitates (NbAGO4-IP) of *N.*
benthamiana leaf patches infiltrated with TYLCV WT or V2 null mutant infectious clones
(TYLCV, TYLCV-V2null) at two days after infiltration . Detection was performed with a ^{32}P-
labeled DNA probe for the intergenic region (IR).

 SUPPLEMENTARY MATERIALS

Supplemental Table S1. List of primers used for cloning in this study.

Supplemental Table S2. List of plasmids used in this study.

Supplemental Table S3. List of primers used for qPCR and qRT-PCR in this study.

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. *SlAGO4* expression in TYLCV-infected and control tomato plants.

SlAGO4a/b/c/d expression in TYLCV-infected or control (mock-inoculated) tomato plants at 3
weeks post-inoculation (wpi), as measured by qRT-PCR. Gene expression was normalized to
SlActin. Values are the mean of three independent biological replicates; error bars indicate SEM.

Figure S2. V2 is essential for systemic infection in tomato and *N. benthamiana*.

A   Design of the V2 null TYLCV mutant used in this work. The second codon of the V2 ORF,
originally encoding a Trp (TGG), is mutated to STOP codon (TAG).

B   Representative pictures of tomato plants infected with TYLCV WT or V2 null mutant (TYLCV-
V2null) or mock-inoculated. Photographs were taken at 3 weeks post-inoculation (wpi). Bar,
5cm.

C   Height of tomato plants infected with TYLCV WT or V2 null mutant (TYLCV-V2null) or mock-
inoculated at 3 wpi. Values are the mean of five independent biological replicates; error bars
indicate SEM.

D   Viral (TYLCV) accumulation in tomato plants infected with TYLCV WT or V2 null mutant
(TYLCV-V2null) or mock-inoculated at 3 wpi, measured by qPCR. Each sample corresponds
to the apical leaves from six plants. The accumulation of viral DNA is normalized to the *25S*
ribosomal RNA interspacer (*ITS*). Values are the mean of six independent biological
replicates; error bars indicate SEM.

E   Representative pictures of *N. benthamiana* plants infected with TYLCV WT or V2 null mutant
(TYLCV-V2null) or mock-inoculated. Photographs were taken at 3 weeks post-inoculation
(wpi). Bar, 5cm.

F Viral (TYLCV) accumulation in *N. benthamiana* plants infected with TYLCV WT or V2 null
mutant (TYLCV-V2null) or mock-inoculated at 3 wpi, measured by qPCR. Each sample
corresponds to the apical leaves from six plants. The accumulation of viral DNA is normalized
to the *25S ribosomal RNA interspacer (ITS)*. Values are the mean of six independent
biological replicates; error bars indicate SEM.

Figure S3. Experimental design for local and systemic TYLCV infection assays in *NbAGO4-*
silenced *N. benthamiana* plants*.*

A Experimental design for local TYLCV infection assays. *A. tumefaciens* carrying the TRV-EV
or TRV-*NbAGO4* infectious clones were inoculated on 18-day-old *N. benthamiana* cotyledons*.*
2 weeks later, young leaves were infiltrated with *A. tumefaciens* carrying the TYLCV infectious
clone (WT or V2null) and leaf patches were collected at 3, 4, or 9 days post-inoculation (dpi).

B Experimental design for systemic TYLCV infection assays. *A. tumefaciens* carrying the TRV-
EV or TRV-*NbAGO4* infectious clones were inoculated on 18-day-old *N. benthamiana*
cotyledons. At the same time, *A. tumefaciens* carrying the TYLCV infectious clone (WT or
V2null) were injected into plant stems. The top three leaves were collected at 3 weeks post-
inoculation (wpi).

Figure S4. Original single-base resolution bisulfite sequencing data of the intergenic
region (IR) of TYLCV in local infection assays at 3 dpi or 9 dpi (Figure 5A).

At least 5 individual clones were sequenced per replicate and sample at 3 dpi, and >18 individual
were sequenced per replicate and sample at 9 dpi. Each single circle, corresponding to a cytosine,
is colored in blue, red, or green, representing the CHG, CG or CHH contexts, respectively.
Methylated cytosines are represented by filled circles, while unmethylated cytosines are
represented by empty circles.

Figure S5. Original single-base resolution bisulfite sequencing data of the intergenic
region (IR) of TYLCV-V2null in local infection assays at 4 dpi in *NbAGO4*-silenced plants
(Figure 5B). At least 7 individual clones were sequenced per sample in the first replicate, and >
15 individual clones were sequenced per sample in replicates second to fourth. Each single circle,

corresponding to a cytosine, is colored in blue, red, or green, representing the CHG, CG or CHH
contexts, respectively. Methylated cytosines are represented by filled circles, while unmethylated
cytosines are represented by empty circles.

Figure S6. Original single-base resolution bisulfite sequencing data of the intergenic
region (IR) of TYLCV and TYLCV-V2null in systemic infection assays in *NbAGO4*-silenced
plants (Figure 5C).

> 14 individual clones were sequenced per sample and replicate. Each single circle,
corresponding to a cytosine, is colored in blue, red, or green, representing the CHG, CG or CHH
contexts, respectively. Methylated cytosines are represented by filled circles, while unmethylated
cytosines are represented by empty circles.

Figure S7. TYLCV V2$_{L76S}$ interacts with AGO4 in the Cajal body.

A   Alignment of the amino acid sequences of V2 from *Cotton Leaf Curl Multan virus* (CLCuMuV)
and V2 from TYLCV. The alignment was performed by Geneious (https://www.geneious.com).
Black background indicates conservation. The identity of these two proteins is 65%.

B   V2$_{L76S}$-GFP and RFP-AGO4 co-localize in the Cajal body. CFP-Fibrillarin, V2$_{L76S}$-GFP and
RFP-NbAGO4-1/2 or RFP-SlAGO4a/b/d were transiently co-expressed in *N. benthamiana*
epidermal cells. CFP-Fibrillarin is used as a nucleolus and Cajal body marker. Confocal
images were taken at two days after infiltration. Arrowheads indicate the position of the Cajal
body. Bar, 5µm. This experiment was repeated three times with similar results.

C   3xHA-NbAGO4-1 interacts with V2-GFP and V2$_{L76S}$-GFP in co-immunoprecipitation (co-IP)
assays upon transient expression in *N. benthamiana*. Free GFP was used as negative control.
CBB, Coomassie brilliant blue staining. The V2-GFP sample was diluted 1/20 for western blot
to reach a protein amount comparable to that of V2$_{L76S}$-GFP. Three independent biological
replicates were performed with similar results.

D   V2$_{L76S}$ interacts with AGO4 in the Cajal body. The N-terminal half of the YFP fused to V2$_{L76S}$
(V2$_{L76S}$-nYFP) was transiently co-expressed with the C-terminal half of the YFP alone (cYFP,
as a negative control), or cYFP-NbAGO4, cYFP-SlAGO4, or cYFP- V2$_{L76S}$ in *N. benthamiana*
leaves. CFP-Fibrillarin was used as a nucleolus and Cajal body marker. V2 was used as a

control. Yellow fluorescence indicates a positive interaction. Arrowheads indicate the position

of the Cajal body. Bar, 5µm. This experiment was repeated three times with similar results.

Figure S8. Model for the V2-mediated inhibition of the AGO4-dependent methylation of the

viral DNA.

During the viral infection, the ssDNA TYLCV genome forms dsDNA replicative intermediates,

which could be targeted by the host AGO4-dependent RNA-directed DNA methylation (RdDM)

pathway as an antiviral defence mechanism. Viral small interfering RNA (vsiRNA) are generated

and loaded into AGO4. In the absence of the virus-encoded V2 protein, the AGO4-vsiRNA

complex could be effectively guided towards the viral genome by complementary base pairing to

the scaffold RNA and association with Pol V, and recruit the methyl transferase DRM2 to catalyze

methylation of the viral genome. When V2 is present, however, V2 interacts with AGO4 and

interferes with the binding of this protein to the viral DNA, enabling viral evasion from the AGO4-

dependent DNA methylation.

Figure 1

A

Identified protein name	Identified protein ID	Bait protein name	Exclusive unique peptide count	Identified protein coverage (%)	Best Mascot Score
NbAGO4-1	NbS00018144g003.1	GFP	- / - / -	- / - / -	- / - / -
		V2-GFP	3 / - / -	6.1 / - / -	44.9 / - / -
NbAGO4-2	NbS00007950g0008.1	GFP	- / - / -	- / - / -	- / - / -
		V2-GFP	16 / 1 / 1	23 / 1.8 / 1.2	116.2 / 35.7 / 59.2

B

C

Figure 2

A

B

Identity (%) / Distance matrix							
Distance \ Identity	AtAGO 4	NbAGO 4-1	NbAGO 4-2	SlAGO 4a	SlAGO 4b	SlAGO 4c	SlAGO 4d
AtAGO4		0.302	0.303	0.305	0.308	0.390	0.384
NbAGO4-1	73.5		0.133	0.137	0.056	0.177	0.339
NbAGO4-2	73.4	86.8		0.071	0.148	0.242	0.329
SlAGO4a	73.5	86.8	92.5		0.140	0.248	0.336
SlAGO4b	73.0	94.4	85.4	86.4		0.146	0.332
SlAGO4c	67.3	83.1	77.7	77.6	85.7		0.432
SlAGO4d	67.8	70.8	71.6	71.1	71.2	64.9	

C

D

Figure 3

Figure 4

Figure 5

Figure 6

A

B