Identifying Sialylation Linkages at the Glycopeptide Level by Glycosyltransferase Labeling Assisted Mass Spectrometry (GLAMS)

He Zhu[†], Shuaishuai Wang[†], Ding Liu[†], Lang Ding[†], Congcong Chen[†], Yunpeng Liu[†], Zhigang Wu[†], Roni Bollag[‡], Kebin Liu[§], Jun Yin[†], Cheng Ma[†], Lei Li[*†], Peng George Wang[*†,¶]

[†] Department of Chemistry, Georgia State University, Atlanta, GA 30303, United States

[‡] Georgia Cancer Center, Augusta University, Augusta, GA 30912, United States

[§] Department of Biochemistry and Molecular Biology, Augusta University, Augusta, GA 30912, United States

[¶] Current address: Department of Chemistry, Southern University of Science and Technology, Shenzhen 518055, China.

ABSTRACT

Precise assignment of sialylation linkages at the glycopeptide level is of importance in bottom-up glycoproteomics, and is also an indispensable step to understand the function of glycoproteins in pathogen-host interactions and cancer progression. Even though some efforts have been dedicated to the discrimination of $\alpha2,3/\alpha2,6$-sialylated isomers, unambiguous identification of sialoglycopeptide isomers is still needed. Herein, an innovative strategy of glycosyltransferase labeling assisted mass spectrometry (GLAMS) were developed. After specific enzymatic labeling, oxonium ions from higher-energy C-trap dissociation (HCD) fragmentation of $\alpha2,3$-sailoglycopeptides generate unique reporters to distinctly differentiate those of $\alpha2,6$-sailoglycopeptide isomers. Using this strategy, a total of 1,138 linkage-specific sialoglycopeptides were successfully identified from 126 glycoproteins in human serum.

INTRODUCTION

Glycosylation is one of the most important post-translational modifications (PTMs) of proteins[1]. It plays an indispensable role in a wide variety of biological and physiological processes including protein folding[2], cellular communication and signaling[3], immune response[4], and pathogen-host interactions[5]. It was estimated that more than half of proteins in the human proteome are glycosylated[6]. Glycosylation is generally classified into N-glycosylation and mucin-type O-glycosylation. In N-glycosylation, the glycans are attached to asparagine residues of proteins in the consensus motif Asn-X-Ser/Thr/Cys (X can be any amino acid except for proline)[7], whereas the glycan moieties are linked to serine or threonine residues of proteins in O-glycosylation[1]. N-acetylneuraminic acid (Neu5Ac) is most common form of diverse sialic acids found on glycan chains in humans[8]. Neu5Ac is the frequently the terminal monosaccharide residue that linked to galactose (Gal) via an α2,3 or an α2,6 linkage[9]. The two Neu5Ac linkage isomers have important roles in pathogen binding. For example, avian influenza A viruses preferentially bind to respiratory epithelial cells that contain α2,3-linked Neu5Ac moieties, while human influenza viruses selectively bind to α2,6-linked Neu5Ac moieties[10]. In addition, altered sialylation linkages is closely correlated with health status. In patients diagnosed with prostate cancer, the structure of sialylated glycans released from the prostate-specific antigen (PSA) is rich in α2,3 linkages, whereas the structure of sialylated glycans from healthy people's PSA is predominantly via α2,6 linkages[11]. Without detailed information on sialylation linkages, it is challenging to decipher the biological functions of sialoglycoproteins.

A variety of methods, including sialidase treatment[12], chemical derivatization[13-21], and mass spectrometry (MS)[22-24], have been used to distinguish α2,3 and α2,6-sialylated glycans, but the location of corresponding glycosylation sites is not evident. The emerging technology of MS-

based glycoproteomics[25-29] provides a high-throughput approach to analyze site-specific protein

glycosylation. Consequently, strategies for identifying sialylation linkages at the glycopeptide

level are required. Established chemical approaches[30, 31] modify not only sialic acids of the

glycan moiety but also acidic amino acid residues and the C-terminus of the peptide backbone,

which increases the complexity of spectra analysis, especially in relation to large scale

glycoproteomics data[32]. Pseudo MS^3 fragmentation[22] and ion mobility-mass spectrometry (IM-

MS)[33] have also been applied to distinguish sialoglycopeptide isomers but this technology has

not been **demonstrated (**?effective) for large-scale glycopeptides analysis. Even though the

fragmentation of high energy C-trap dissociation (HCD) can achieve the high-throughput

assignment of α2,3/α2,6-sialoglycopeptide isomers, there are still many ambiguous

glycopeptides without sialylation linkage information[34].

Enzymatic labeling has become a promising tool for probing proteins with specific

glycosylation because of its high specificity and mild reaction condition[35-41]. In our previous

research, a β1,4-N-acetylgalactosaminyltransferase from *Campylobacter jejuni* (CgtA) was

demonstrated to specifically recognize the Neu5Ac(α2,3)Gal epitope and transfer N-

acetylgalactosamine (GalNAc) or N-azidoacetylgalactosamine (GalNAz) to Gal residue via β1,4

linkage[42]. Taking advantage of the specificity of CgtA, we herein report a new strategy for

identification of α2,3 and α2,6 sialoglycopeptide isomers by glycosyltransferase labeling assisted

mass spectrometry (GLAMS). As shown in **Scheme 1**, sialoglycopeptides (monosialylation for

example) without linkages information were treated with GalNAz labeling by CgtA. After HCD

fragmentation, if only the oxonium ions of HexNAc (N-acetylhexosamine, m/z 204) and Neu5Ac

(m/z 292) are observed in the spectrum, this identifies an α2,6 linkage; if one more oxonium ion

m/z 245 from GalNAz is included, then it represents an α2,3 linkage. We demonstrated the

capacity of GLAMS to effectively differentiate α2,3/α2,6 isomers using synthetic

sialoglycopeptides, and applied it to profile sialylated N-glycopeptides digested from serum

samples, in which we identified 1,138 linkage specific sialoglycopeptides with the modified

pGlyco 2.0 software[27].

EXPERIMENTAL SECTION

Chemicals and materials

Amicon Ultra-0.5 Centrifugal Filter Unit with NMWL of 30 kD was obtained from

Millipore Sigma (St. Louis, MO). iSPE® HILIC SPE Cartridge was purchased from The Nest

Group (Southborough, MA). XAmide medium was provided by Dr. Xinmiao Liang's lab (Dalian

Institute of Chemical Physics, Chinese Academy of Sciences). Sequencing grade porcine trypsin

was purchased from Promega (Madison, WI); Deionized water was produced by a Milli-Q A10

system from Millipore (Bedford, MA). Optimal LC-MS grade quality acetonitrile (ACN), formic

acid (FA) and water were purchased from Thermo Fisher Scientific, Inc. (Waltham, MA).

Dithiothreitol (DTT), 2-Iodoacetamide (IAA) and trifluoroacetic acid (TFA) were bought from

Alfa Aesar (Ward Hill, MA). Ammonium bicarbonate (NH_4HCO_3) was purchased from

Millipore Sigma (St. Louis, MO). All other chemicals of analytical grade were commercially

available.

Sialylated oligosaccharide standards including 3'-sialyllactose (3SL) and 6'-sialyllactose

(6SL) were enzymatically synthesized using *Pasteurella multocida* α2,3-sialyltransferase 1 and

Photobacterium damselae α2,6-sialyltransferase as described before[43].

Serum samples from four healthy people, ten patients with colon cancer and four patients

with liver cancer were provided by Georgia Cancer Center at Augusta University and stored at -

80 °C until use. The protocol for serum sample preparation was approved by the Institutional

Review Board of Augusta University and was performed in accordance with the Helsinki

Declaration. All participants gave written informed consents.

Protein digestion

The protein digestion was performed using a filter aided sample preparation (FASP)[44]

method with minor modification. Approximately 1 mg of total proteins from human serum was

subjected to FASP procedures. Samples were denatured in 20 μL of lysis buffer (4% SDS, 100

mM DTT, 20 mM Tris-HCl, pH 7.6) at 95 °C for 10 min. When the solution was cooled down to

room temperature, 1 mL of UA solution (8 M urea, 20 mM Tris-HCl, pH 8.5) was added. The

solution was gradually transferred into a 30 kD filter and centrifugated at 15,000 rpm at 20 °C

for 15 min. Then 100 μL of IAA solution (50 mM IAA in UA solution) was added to the filter

and incubated for 30 min in the darkness followed by centrifugation at 15,000 rpm for 10 min.

After that, 100 μL of UA solution and 200 μL of digestion buffer (100 μL of 50 mM NH_4HCO_3

each time) were added and centrifuged at 15,000 rpm for 10 min, respectively. Finally, 20 μg of

trypsin was added into the filter and incubated at 37 °C for 12 h. The tryptic peptides were eluted

using 100 μL of 50 mM NH_4HCO_3 for six times. The concentration of the collected peptides was

measured based on absorbance at 280 nm with Nanodrop. Five hundred μg of the peptides were

taken and subsequently dried in Vacufuge at room temperature, then were lyophilized twice to

remove NH_4HCO_3. The dried samples were stored at −20 °C for further use.

Enrichment of glycopeptides

The enrichment of glycopeptides was performed using an iSPE HILIC cartridge by the

procedure of ZIC-HILIC[45]. Briefly, the HILIC cartridge was pre-washed with 300 μL of 0.1%

TFA and conditioning with 600 μL of 80% ACN containing 0.1% TFA. Peptide samples

dissolved in 400 µL of 80% ACN containing 0.1% TFA were loaded onto the cartridge. The

flow-through was reloaded onto the column for another two times. Then the column was washed

with 1.2 mL of 80% ACN containing 1% TFA. The glycopeptides were eluted using 750 µL of

0.1% TFA, 60 µL of H_2O, 60 µL of 25 mM NH_4HCO_3 and 60 µL of 50% ACN. The fractions

were combined followed by lyophilization and stored at -20 °C.

Enzymatic labeling of glycopeptides

The β1,4-N-acetylgalactosaminyltransferase from *C. jejuni* (CgtA) and sugar nucleotide

UDP-N-azidoacetylgalactosamine (UDP-GalNAz) were prepared as described previously[42]. For

enzymatic labeling, the 40 µL reaction mixture containing the enriched glycopeptides, 50 mM

Tris-HCl buffer (pH 7.5), 5 mM $MgCl_2$, 200 µM UDP-GalNAz and 10 µg CgtA was incubated

in a 37 °C water bath for 1 h. The reaction was quenched by adding 60 µL of ACN, 6 µL of 10%

TFA and 215 µL of ACN containing 1% TFA. Then the glycopeptides were purified with an in-

house packed Amide HILIC microcolumn (6 mg of XAmide HILIC medium in a 200 µL tip)

which was preconditioned with 100 µL of 0.1% TFA and 200 µL of 85% ACN containing 0.1%

TFA. After loading the reaction mixture onto the tip column, the column was washed with 100

µL of 85% ACN, 0.1% TFA to remove salts and proteins. The glycopeptides were collected

using 200 µL of 0.1% TFA and lyophilized.

Nano LC-MS/MS analysis

Nano RP HPLC-MS experiments were performed on an LTQ-Orbitrap Elite mass

spectrometer (Thermo Fisher) equipped with EASY-spray source and nano-LC UltiMate 3000

high-performance liquid chromatography system (Thermo Fisher). The separation was

performed on an EASY-Spray PepMap C18 Column (75 µm × 50 cm, 2 µm, ThermoFisher, US)

with a linear gradient from 3% to 40% mobile phase B for 120 min at a flow rate of 300 nL/min

(mobile phase A: 1.95% ACN, 97.95% H_2O, 0.1% FA; mobile phase B: 79.95% ACN, 19.95%

H_2O, 0.1% FA). The mass spectrometer was operated in the data-dependent mode. A full-scan

survey MS experiment (m/z range from 375 to 2000; automatic gain control target, 1,000,000

ions; resolution at 400 m/z, 60,000; maximum ion accumulation time, 50 ms) was acquired by

the Orbitrap mass spectrometer, and ten most intense ions were fragmented by higher-energy C-

trap dissociation (HCD) with an optimized stepped collision energy (SCE) of $30 \pm 15\%$ NCE.

Data analysis

A modified version of pGlyco 2.0 was used to identify N-glycopeptides[27]. In the glycan

database, we defined an unnatural monosaccharide GalNAz (Gaz, Z). We created all the possible

"Gaz" containing human N-glycan sequences following the motif H(A)(Z), where "H" represents

Galactose, "A" represents Neu5Ac and "Z" represents GalNAz. For the sialoglycopeptides

identification, the raw data was first converted to an mgf file using pParse[46]. The potential

sialoglycopeptides spectra containing oxonium ions of both GlcNAc (m/z 204.09, m/z 186.08)

and Neu5Ac (m/z 292.10, m/z 274.09) were extracted and merged into a new mgf file using an

in-house program. The new mgf file was searched against an Uniprot-Swiss human protein

database (20,417 reviewed entries on March 20, 2019) and a modified N-glycan database. Other

searching parameters were used including carbamidomethyl (Cys) as a fixed modification,

deamination (Asn) and oxidation (Met) as variable modifications, and trypsin as the digestion

enzyme with maximum two missed cleavage sites allowed. The mass tolerance for the precursor

ions and the fragment ions was set to 10 ppm and 20 ppm, respectively. A total false discovery

rate (FDR) of 1% was applied to all data sets at the glycopeptide level. The identified

glycoproteins were annotated using Uniprot ID mapping[47].

RESULTS AND DISCUSSION

The specificity of CgtA for Sialoglycopeptides

Our previous study showed that CgtA had a high specificity for (Neu5Ac-α2,3-Gal)-containing glycans[42]. To test its selectivity for sialoglycopeptides, we synthesized 23SP and 26SP (synthesis and data not published) as shown in **Figure 1A**. The activity of CgtA was validated with sialolactose prior to the enzymatic reaction of sialoglycopeptides. After labeling with GalNAz by CgtA, only α2,3-sialylated lactose showed a mass shift of 244 Da in the MS spectrum and no substrate was obviously observed (**Figure S1**), which indicated that CgtA had good catalytic activity and high specificity as well.

ESI-MS was used to monitor the GalNAz labeling reaction for sialoglycopeptides. For the α2,3-sialoglycopeptide 23SP, the monoisotopic peak at m/z 1032.0808 is from the triply charged glycopeptide (**Figure 1B**). After incubating with CgtA and UDP-GalNAz for 30 min, a new dominant peak at m/z 1113.4422 with three positive charges was observed, which correspond to the GalNAz labeled 23SP (**Figure 1D**). However, we could still detect minor substrate 23SP. The α2,6-sialoglycopeptide 26SP has the same exact mass (**Figure 1C**) as 23SP. After GalNAz labeling by CgtA, the strongest peak is located at the same exact position, and a trace amount of product at m/z 1113.4422 was detected (**Figure 1E**). This may result from the very weak activity of CgtA for 2,6-sialylated glycans[42].

The reaction time was further optimized to increase the conversion rate of GalNAz labeling of 23SP. The reaction mixture at different time points was analyzed by LC-MS and the reaction efficiency was calculated based on the peak area. For 23SP, in the first half an hour, the reaction efficiency was dramatically increased. After that, the increase slowed down and the reaction efficiency approached 100% at 60 min. Meanwhile, the conversion rate for the 26SP

185   remained unchanged (less than 1%) over time (**Figure 2**). Finally, we chose 60 min as the

186   reaction time for the subsequent GalNAz labeling reactions.

Identifying Sialoglycopeptide Linkage Isomers by GalNAz Labeling and Tandem MS

Although discrimination of α2,3/α2,6-sialoglycopeptide isomers can be achieved by the

mass shift in primary MS as shown above, we cannot accurately determine the glycopeptide

sequence using only molecular weight information. Tandem MS is an effective way for

sequencing glycopeptides, and HCD-MS/MS is commonly used to analyze glycopeptides

because of the high accuracy and detectable diagnostic oxonium ions of monosaccharides[26-28, 48].

It was also demonstrated that HCD with stepped collision energy (SCE) can simultaneously

determine the sequences of both glycans and peptides in a single run[27, 48]. To achieve a better

sequence coverage of glycopeptides on the Orbitrap we used, SCE was optimized with

glycopeptides enriched from human serum. The identification of N-glycopeptides was performed

using pGlyco 2.0 software[27]. A series of two-step SCEs and three-step SCEs were employed to

compare the number of identified glycopeptides. As shown in **Figure S2**, with a three-step SCE

of "15-30-45", more glycopeptides can be identified from the same sample.

The optimized SCE-HCD-MS/MS was applied for sequencing 23SP and 26SP. After

GalNAz labeling, 23SP (**Figure 3A**) showed diagnostic oxonium ions of GlcNAc and Neu5Ac at

m/z 204.09 and 292.10, which are common features of sialoglycopeptides. Apart from these

oxonium ions, 23SP also had a strong unknown peak at m/z 245.09. Compared with the spectrum

of 23SP before GalNAz labeling (**Figure S3**), we can obviously see that the peak with m/z

245.09 corresponds to the oxonium ion of GalNAz. Following the dissociation pathway of

GlcNAc[39, 49, 50] in **Scheme S1A**, a series of fragmentation ions of GalNAz in HCD were

illustrated in **Scheme S1B**. The peaks at m/z 245.09, m/z 227.08, m/z 209.07 and m/z 179.06,

which match the theoretical values very well, are diagnostic ions of GalNAz (**Figure S4**). For

26SP (**Figure 3B**), its spectrum showed that it shared similar fragmentation patterns with

GalNAz labeled 23SP. In the low m/z region, only diagnostic ions of GlcNAc and Neu5Ac were

observed, which clearly demonstrated that the GLAMS is an effective approach to distinguish

intact linkage specific sialoglycopeptides.

Linkage Specific Sialoglycopeptides in Human Serum

We then applied this method to analyze the N-linked α2,3/α2,6-sialoglycopeptides of

complex biological samples. Serum samples from healthy individuals (n = 4) and patients

diagnosed with liver cancer (n = 4) and colon cancer (n = 10) were used in this study. In the

beginning, we tried to perform the enzymatic labeling of the mixture of tryptic peptides and

glycopeptides. After drying from the FASP eluents, the mixture was not completely dissolved in

the Tris buffer (pH 7.5) for the enzymatic reaction, which may result from the existence of some

basic peptides. To solve the solubility issue, we adopted step-by-step glycopeptides enrichment

and enzymatic labeling as shown in the workflow **Figure 4**. It turned out that the enriched

glycopeptides were dissolved in the basic buffer very well. This may be attributed to that

conjugated glycans increase the hydrophilicity of peptides and some sialoglycans decrease the

isoelectric points of peptides according to the previous findings[51, 52].

All the serum samples were analyzed in three replicated LC-MS/MS runs for a large-

scale study of N-linked sialoglycopeptides. Using the oxonium ions of GalNAz as a tag to mark

α2,3-sialylation, a total of 18,822 sialoglycopeptide spectra were identified with a 1% total FDR,

corresponding to 1,138 unique site-specific sialoglycans on 220 N-glycosylation sites from 126

glycoproteins (**Table S1**), among of which 63% was also identified previously using a

deglycosylation method[53] as shown in **Figure S5**. For the distribution analysis of α2,3 or α2,6-

sialylation (**Figure 5A**), only 71 distinct site-specific sialoglycopeptides contained pure α2,3-

sialylation linkage, 186 sialoglycopeptides contained hybrid sialylation, and all the rest were

uniquely 2,6-sialylated, which accounts for more than 77% of the identified sialoglycopeptides.

This is consistent with the previous result that α2,6-sialylation is dominant in sialylated N-

glycans released from human serum samples[54]. Taking a closer look at the distribution of

sialylation levels, monosialylation is the major component in strictly α2,3 or α2,6-

sialoglycopeptides, and 83% of the hybrid sialoglycopeptides are disialylated.

The percentage of the overall overlapped glycosites (**Figure 5C**) and glycoproteins

(**Figure 5D**) in a specific type of serum is 76-84% and 75-80%, respectively. The percentage of

overlapping sialoglycopeptides (**Figure 5B**) in each kind of serum sample was significantly

lower than those of the glycosites and glycoproteins, reflecting the diversity of site-specific

glycosylation, especially for samples from colon cancer and liver cancer patients. A highly

glycosylated protein P01876 (IGHA1) was selected to demonstrate the diversity of site-specific

glycosylation in the three different sera. The previous study showed that IgA was α2,3 and α2,6-

sialylated on N-glycans[55]. Here, we identified 59 unique sialoglycopeptides including α2,3 and

α2,6 linkages on three known N-glycosylation sites of IGHA1 (**Table S1**). In addition, we found

some unique sialoglycopeptides of IGHA1 expressed in a serum-specific manner. For example,

both unique α2,6-sialoglycopeptides and hybrid sialoglycopeptides of IGHA1 were identified in

serum from colon cancer patients (representative spectra in **Figure S6**) and liver cancer patients

(representative spectra in **Figure S7**), whereas only one unique α2,6-sialoglycopeptide was

detected from control sera (**Figure S8**).

CONCLUSIONS

Elucidating linkage-specific sialylation of glycoproteins is significant to understand

pathogen-host interactions and disease development. By integrating enzymatic labeling and MS-

based glycoproteomics, we developed a high-throughput glycosyltransferase labeling assisted

mass spectrometry (GLAMS) to identify sialoglycopeptide linkage isomers. Facilitated by the

acceptor specificity of the glycosyltransferase CgtA toward the Neu5Ac(α2,3)Gal epitope, we

have demonstrated the distinct diagnostic oxonium ions as tags to effectively discriminate α2,3-

and α2,6-sialoglycopeptide isomers using LC separation coupled with SCE-HCD-MS/MS

fragmentation. We also successfully applied this approach to a large-scale profiling of linkage-

specific sialoglycopeptides in human serum samples. The strategy of GLAMS should be of great

benefit in investigating differential changes in response to disease states from a glycoproteomics

perspective, and has potential to contribute to biomarker discovery in many diseases.

AUTHOR INFORMATION

Corresponding Authors

lli22@gsu.edu (L. L.) or pwang11@gsu.edu (P. G. W.)

Author Contributions

H.Z., L. L., and P. G. W. conceived the project; H.Z. designed and performed the experiments, analyzed the data, and wrote the manuscript; S. W. synthesized the sialoglycopeptides; D. L. and L. D. prepared the serum glycopeptides for collision energy optimization; C. C., Y. L. and Z. W. synthesized the UDP-GalNAz; R. B. and K. L. collected and prepared the human serum samples; C. M., J. Y., L. L., and P. G. W. edited the manuscript. All the authors revised and approved the manuscript.

ACKNOWLEDGEMENTS

We sincerely thank Dr. Wenfeng Zeng from the Key Lab of Intelligent Information Processing of Chinese Academy of Sciences for the help of parameter setting of pGlyco 2.0, Georgia Cancer Center Biorepository for providing human serum samples, and Georgia Research Alliance for financial support to purchase mass spectrometers. This work is supported by National Institute of Health (U01GM116263 and U54HL142019).

CONFLICT OF INTERESTS

The authors declare no conflict of interest.

REFERENCES

1.   Ohtsubo, K.; Marth, J. D., Glycosylation in cellular mechanisms of health and disease. *Cell* **2006,** *126* (5), 855-67.

2.   Banerjee, S.; Vishwanath, P.; Cui, J.; Kelleher, D. J.; Gilmore, R.; Robbins, P. W.; Samuelson, J., The evolution of N-glycan-dependent endoplasmic reticulum quality control factors for glycoprotein folding and degradation. *P Natl Acad Sci USA* **2007,** *104* (28), 11676-11681.

3.   Furukawa, K.; Ohkawa, Y.; Yamauchi, Y.; Hamamura, K.; Ohmi, Y.; Furukawa, K., Fine tuning of cell signals by glycosylation. *J Biochem* **2012,** *151* (6), 573-578.

4.   Rudd, P. M.; Elliott, T.; Cresswell, P.; Wilson, I. A.; Dwek, R. A., Glycosylation and the immune system. *Science* **2001,** *291* (5512), 2370-2376.

5.   Karlsson, K. A., Pathogen-host protein-carbohydrate interactions as the basis of important infections. *Adv Exp Med Biol* **2001,** *491*, 431-443.

6.   Apweiler, R.; Hermjakob, H.; Sharon, N., On the frequency of protein glycosylation, as deduced from analysis of the SWISS-PROT database. *Bba-Gen Subjects* **1999,** *1473* (1), 4-8.

7.   Zielinska, D. F.; Gnad, F.; Wisniewski, J. R.; Mann, M., Precision Mapping of an In Vivo N-Glycoproteome Reveals Rigid Topological and Sequence Constraints. *Cell* **2010,** *141* (5), 897-907.

8.   Varki, N. M.; Varki, A., Diversity in cell surface sialic acid presentations: implications for biology and disease. *Lab Invest* **2007,** *87* (9), 851-7.

9.   Varki, A.; Schauer, R., Sialic Acids. In *Essentials of Glycobiology*, nd; Varki, A.; Cummings, R. D.; Esko, J. D.; Freeze, H. H.; Stanley, P.; Bertozzi, C. R.; Hart, G. W.; Etzler, M. E., Eds. Cold Spring Harbor (NY), 2009.

10.   Stevens, J.; Blixt, O.; Paulson, J. C.; Wilson, I. A., Glycan microarray technologies: tools to survey host specificity of influenza viruses. *Nat Rev Microbiol* **2006,** *4* (11), 857-64.

11.   Hatakeyama, S.; Yoneyama, T.; Tobisawa, Y.; Ohyama, C., Recent progress and perspectives on prostate cancer biomarkers. *Int J Clin Oncol* **2017,** *22* (2), 214-221.

12.   Mancera-Arteu, M.; Gimenez, E.; Barbosa, J.; Sanz-Nebot, V., Identification and characterization of isomeric N-glycans of human alfa-acid-glycoprotein by stable isotope labelling and ZIC-HILIC-MS in combination with exoglycosidase digestion. *Anal Chim Acta* **2016,** *940*, 92-103.

13.   Mechref, Y.; Kang, P.; Novotny, M. V., Differentiating structural isomers of sialylated glycans by matrix-assisted laser desorption/ionization time-of-flight/time-of-flight tandem mass spectrometry. *Rapid Commun Mass Sp* **2006,** *20* (8), 1381-1389.

14.   Alley, W. R., Jr.; Novotny, M. V., Glycomic analysis of sialic acid linkages in glycans derived from blood serum glycoproteins. *J Proteome Res* **2010,** *9* (6), 3062-72.

15.   Jiang, K.; Zhu, H.; Li, L.; Guo, Y. X.; Gashash, E.; Ma, C.; Sun, X. L.; Li, J.; Zhang, L. W.; Wang, P. G., Sialic acid linkage-specific permethylation for improved profiling of protein glycosylation by MALDI-TOF MS. *Anal Chim Acta* **2017,** *981*, 53-61.

16.   Wheeler, S. F.; Domann, P.; Harvey, D. J., Derivatization of sialic acids for stabilization in matrix-assisted laser desorption/ionization mass spectrometry and concomitant differentiation of $\alpha(2\rightarrow3)$ and $\alpha(2\rightarrow6)$-isomers. *Rapid Commun Mass Spectrom* **2009,** *23* (2), 303-12.

17.   Reiding, K. R.; Blank, D.; Kuijper, D. M.; Deelder, A. M.; Wuhrer, M., High-Throughput Profiling of Protein N-Glycosylation by MALDI-TOF-MS Employing Linkage-Specific Sialic Acid Esterification. *Anal Chem* **2014,** *86* (12), 5784-5793.

18.   Holst, S.; Heijs, B.; de Haan, N.; van Zeijl, R. J. M.; Briaire-de Bruijn, I. H.; van Pelt, G. W.; Mehta, A. S.; Angel, P. M.; Mesker, W. E.; Tollenaar, R. A.; Drake, R. R.; Bovee, J. V. M. G.; McDonnell, L. A.; Wuhrer, M., Linkage-Specific in Situ Sialic Acid Derivatization for N-Glycan Mass Spectrometry Imaging of Formalin-Fixed Paraffin-Embedded Tissues. *Anal Chem* **2016,** *88* (11), 5904-5913.

19.   Yang, S.; Jankowska, E.; Kosikova, M.; Xie, H.; Cipollo, J., Solid-Phase Chemical Modification for Sialic Acid Linkage Analysis: Application to Glycoproteins of Host Cells Used in Influenza Virus Propagation. *Anal Chem* **2017,** *89* (17), 9508-9517.

20.   Li, H. H.; Gao, W. J.; Feng, X. J.; Liu, B. F.; Liu, X., MALDI-MS analysis of sialylated N-glycan linkage isomers using solid-phase two step derivatization method. *Anal Chim Acta* **2016,** *924*, 77-85.

21.   Nishikaze, T.; Tsurnoto, H.; Sekiya, S.; Iwamoto, S.; Miura, Y.; Tanaka, K., Differentiation of Sialyl Linkage Isomers by One-Pot Sialic Acid Derivatization for Mass Spectrometry-Based Glycan Profiling. *Anal Chem* **2017,** *89* (4), 2353-2360.

22.   Chen, C. H.; Lin, Y. P.; Lin, J. L.; Li, S. T.; Ren, C. T.; Wu, C. Y.; Chen, C. H., Rapid Identification of Terminal Sialic Acid Linkage Isomers by Pseudo-MS3 Mass Spectrometry. *Isr J Chem* **2015,** *55* (3-4), 412-422.

23.   Guttman, M.; Lee, K. K., Site-Specific Mapping of Sialic Acid Linkage Isomers by Ion Mobility Spectrometry. *Anal Chem* **2016,** *88* (10), 5212-5217.

24.   Depland, A. D.; Renois-Predelus, G.; Schindler, B.; Compagnon, I., Identification of sialic acid linkage isomers in glycans using coupled InfraRed Multiple Photon Dissociation (IRMPD) spectroscopy and mass spectrometry. *Int J Mass Spectrom* **2018,** *434*, 65-69.

25.   Woo, C. M.; Iavarone, A. T.; Spiciarich, D. R.; Palaniappan, K. K.; Bertozzi, C. R., Isotope-targeted glycoproteomics (IsoTaG): a mass-independent platform for intact N- and O-glycopeptide discovery and analysis. *Nat Methods* **2015,** *12* (6), 561-7.

26.   Sun, S. S.; Shah, P.; Eshghi, S. T.; Yang, W. M.; Trikannad, N.; Yang, S.; Chen, L. J.; Aiyetan, P.; Hoti, N.; Zhang, Z.; Chan, D. W.; Zhang, H., Comprehensive analysis of protein glycosylation by solid-phase extraction of N-linked glycans and glycosite-containing peptides. *Nat Biotechnol* **2016,** *34* (1), 84-88.

27.   Liu, M. Q.; Zeng, W. F.; Fang, P.; Cao, W. Q.; Liu, C.; Yan, G. Q.; Zhang, Y.; Peng, C.; Wu, J. Q.; Zhang, X. J.; Tu, H. J.; Chi, H.; Sun, R. X.; Cao, Y.; Dong, M. Q.; Jiang, B. Y.; Huang, J. M.; Shen, H. L.; Wong, C. C. L.; He, S. M.; Yang, P. Y., pGlyco 2.0 enables precision N-glycoproteomics with comprehensive quality control and one-step mass spectrometry for intact glycopeptide identification. *Nat Commun* **2017,** *8*.

28.   Stadlmann, J.; Taubenschmid, J.; Wenzel, D.; Gattinger, A.; Durnberger, G.; Dusberger, F.; Elling, U.; Mach, L.; Mechtler, K.; Penninger, J. M., Comparative glycoproteomics of stem cells identifies new players in ricin toxicity. *Nature* **2017,** *549* (7673), 538-+.

29.   Riley, N. M.; Hebert, A. S.; Westphall, M. S.; Coon, J. J., Capturing site-specific heterogeneity with large-scale N-glycoproteome analysis. *Nat Commun* **2019,** *10*.

30.   Nishikaze, T.; Nakamura, T.; Jinmei, H.; Amano, J., Negative-ion MALDI-MS2 for discrimination of alpha2,3- and alpha2,6-sialylation on glycopeptides labeled with a pyrene derivative. *J Chromatogr B Analyt Technol Biomed Life Sci* **2011,** *879* (17-18), 1419-28.

31.   de Haan, N.; Reiding, K. R.; Haberger, M.; Reusch, D.; Falck, D.; Wuhrer, M., Linkage-Specific Sialic Acid Derivatization for MALDI-TOF-MS Profiling of IgG Glycopeptides. *Anal Chem* **2015,** *87* (16), 8284-8291.

32.   Yang, S.; Wu, W. W.; Shen, R. F.; Bern, M.; Cipollo, J., Identification of Sialic Acid
Linkages on Intact Glycopeptides via Differential Chemical Modification Using IntactGIG-
HILIC. *J Am Soc Mass Spectr* **2018,** *29* (6), 1273-1283.
33.   Hinneburg, H.; Hofmann, J.; Struwe, W. B.; Thader, A.; Altmann, F.; Silva, D. V.;
Seeberger, P. H.; Pagel, K.; Kolarich, D., Distinguishing N-acetylneuraminic acid linkage
isomers on glycopeptides by ion mobility-mass spectrometry. *Chem Commun* **2016,** *52* (23),
4381-4384.
34.   Pett, C.; Nasir, W.; Sihlbom, C.; Olsson, B. M.; Caixeta, V.; Schorlemer, M.; Zahedi,
R. P.; Larson, G.; Nilsson, J.; Westerlind, U., Effective Assignment of alpha 2,3/alpha 2,6-
Sialic Acid Isomers by LC-MS/MS-Based Glycoproteomics. *Angew Chem Int Edit* **2018,** *57* (30),
9320-9324.
35.   Khidekel, N.; Arndt, S.; Lamarre-Vincent, N.; Lippert, A.; Poulin-Kerstien, K. G.;
Ramakrishnan, B.; Qasba, P. K.; Hsieh-Wilson, L. C., A chemoenzymatic approach toward the
rapid and sensitive detection of O-GlcNAc posttranslational modifications. *J Am Chem Soc* **2003,**
125 (52), 16162-3.
36.   Chaubard, J. L.; Krishnamurthy, C.; Yi, W.; Smith, D. F.; Hsieh-Wilson, L. C.,
Chemoenzymatic probes for detecting and imaging fucose-alpha(1-2)-galactose glycan
biomarkers. *J Am Chem Soc* **2012,** *134* (10), 4489-92.
37.   Mbua, N. E.; Li, X.; Flanagan-Steet, H. R.; Meng, L.; Aoki, K.; Moremen, K. W.;
Wolfert, M. A.; Steet, R.; Boons, G. J., Selective exo-enzymatic labeling of N-glycans on the
surface of living cells by recombinant ST6Gal I. *Angew Chem Int Ed Engl* **2013,** *52* (49), 13012-
5.
38.   Li, Q.; Li, Z.; Duan, X.; Yi, W., A tandem enzymatic approach for detecting and
imaging tumor-associated Thomsen-Friedenreich antigen disaccharide. *J Am Chem Soc* **2014,**
136 (36), 12536-9.
39.   Zheng, J.; Xiao, H.; Wu, R., Specific Identification of Glycoproteins Bearing the Tn
Antigen in Human Cells. *Angew Chem Int Ed Engl* **2017,** *56* (25), 7107-7111.
40.   Wen, L.; Liu, D.; Zheng, Y.; Huang, K.; Cao, X.; Song, J.; Wang, P. G., A One-Step
Chemoenzymatic Labeling Strategy for Probing Sialylated Thomsen-Friedenreich Antigen. *ACS*
Cent Sci **2018,** *4* (4), 451-457.
41.   Wu, Z. L. L.; Person, A. D.; Anderson, M.; Burroughs, B.; Tatge, T.; Khatri, K.; Zou,
Y. L.; Wang, L. C.; Geders, T.; Zaia, J.; Sackstein, R., Imaging specific cellular glycan
structures using glycosyltransferases via click chemistry. *Glycobiology* **2018,** *28* (2), 69-79.
42.   Wen, L. Q.; Zheng, Y.; Jiang, K.; Zhang, M. Z.; Kondengaden, S. M.; Li, S. S.;
Huang, K.; Li, J.; Song, J.; Wang, P. G., Two-Step Chemoenzymatic Detection of N-
Acetylneuraminic Acid-alpha(2-3)-Galactose Glycans. *J Am Chem Soc* **2016,** *138* (36), 11473-
11476.
43.   Wu, Z.; Liu, Y.; Ma, C.; Li, L.; Bai, J.; Byrd-Leotis, L.; Lasanajak, Y.; Guo, Y.;
Wen, L.; Zhu, H.; Song, J.; Li, Y.; Steinhauer, D. A.; Smith, D. F.; Zhao, B.; Chen, X.;
Guan, W.; Wang, P. G., Identification of the binding roles of terminal and internal glycan
epitopes using enzymatically synthesized N-glycans containing tandem epitopes. *Org Biomol*
Chem **2016,** *14* (47), 11106-11116.
44.   Wisniewski, J. R.; Zougman, A.; Nagaraj, N.; Mann, M., Universal sample preparation
method for proteome analysis. *Nat Methods* **2009,** *6* (5), 359-62.

45.   Parker, B. L.; Thaysen-Andersen, M.; Solis, N.; Scott, N. E.; Larsen, M. R.; Graham,
M. E.; Packer, N. H.; Cordwell, S. J., Site-specific glycan-peptide analysis for determination of
N-glycoproteome heterogeneity. *J Proteome Res* **2013,** *12* (12), 5791-800.
46.   Yuan, Z. F.; Liu, C.; Wang, H. P.; Sun, R. X.; Fu, Y.; Zhang, J. F.; Wang, L. H.; Chi,
H.; Li, Y.; Xiu, L. Y.; Wang, W. P.; He, S. M., pParse: a method for accurate determination of
monoisotopic peaks in high-resolution mass spectra. *Proteomics* **2012,** *12* (2), 226-35.
47.   UniProt, C., The Universal Protein Resource (UniProt). *Nucleic Acids Res* **2007,** *35*
(Database issue), D193-7.
48.   Pioch, M.; Hoffmann, M.; Pralow, A.; Reichl, U.; Rapp, E., glyXtool(MS): An Open-
Source Pipeline for Semiautomated Analysis of Glycopeptide Mass Spectrometry Data. *Anal*
Chem **2018,** *90* (20), 11908-11916.
49.   Zhao, P.; Viner, R.; Teo, C. F.; Boons, G. J.; Horn, D.; Wells, L., Combining high-
energy C-trap dissociation and electron transfer dissociation for protein O-GlcNAc modification
site assignment. *J Proteome Res* **2011,** *10* (9), 4088-104.
50.   Halim, A.; Westerlind, U.; Pett, C.; Schorlemer, M.; Ruetschi, U.; Brinkmalm, G.;
Sihlbom, C.; Lengqvist, J.; Larson, G.; Nilsson, J., Assignment of saccharide identities through
analysis of oxonium ion fragmentation profiles in LC-MS/MS of glycopeptides. *J Proteome Res*
2014, *13* (12), 6024-32.
51.   Willey, K. P., An elusive role for glycosylation in the structure and function of
reproductive hormones. *Hum Reprod Update* **1999,** *5* (4), 330-55.
52.   Hayakawa, S.; Nakai, S., Relationships of Hydrophobicity and Net Charge to the
Solubility of Milk and Soy Proteins. *Journal of Food Science* **1985,** *50* (2), 486-491.
53.   Zhu, H.; Li, X.; Qu, J.; Xiao, C.; Jiang, K.; Gashash, E.; Liu, D.; Song, J.; Cheng, J.;
Ma, C.; Wang, P. G., Diethylaminoethyl Sepharose (DEAE-Sepharose) microcolumn for
enrichment of glycopeptides. *Anal Bioanal Chem* **2017,** *409* (2), 511-518.
54.   Jiang, K.; Zhu, H.; Li, L.; Guo, Y.; Gashash, E.; Ma, C.; Sun, X.; Li, J.; Zhang, L.;
Wang, P. G., Sialic acid linkage-specific permethylation for improved profiling of protein
glycosylation by MALDI-TOF MS. *Anal Chim Acta* **2017,** *981*, 53-61.
55.   Maurer, M. A.; Meyer, L.; Bianchi, M.; Turner, H. L.; Le, N. P. L.; Steck, M.;
Wyrzucki, A.; Orlowski, V.; Ward, A. B.; Crispin, M.; Hangartner, L., Glycosylation of
Human IgA Directly Inhibits Influenza A and Other Sialic-Acid-Binding Viruses. *Cell Rep* **2018,**
23 (1), 90-99.
467

Captions for Figures, Tables and Schemes

Scheme 1. Strategy for distinguishing α2,3/α2,6-sialoglycopeptides by glycosyltransferase labeling assisted mass spectrometry.

Figure 1. (A) Structure of the synthesized α2,3-sialoglycopeptides (23SP) and α2,6-sialoglycopeptides (26SP); ESI-MS spectra of 23SP (B) and 26SP (C) before enzymatic labeling, and 23SP (D) and 26SP (E) after enzymatic labeling.

Figure 2. Time optimization for CgtA labeling of 23SP and 26SP.

Figure 3. HCD-MS/MS spectra of 23SP (A) and 26SP (B) after enzymatic labeling.

Figure 4. Workflow for identification of linkage specific sialoglycopeptides from human serum.

Figure 5. (A) Distribution of the sialoglycopeptides from human serum; Venn diagrams of identified glycopeptides (B), N-glycosylation sites (C), and glycoproteins (D) in serum samples from colon cancer patients, liver cancer patients and normal people.

Figure S1. MALDI-MS spectra of 6SL (A) and 3SL (C) before enzymatic labeling and 6SL (B) and 3SL (D) after enzymatic labeling.

Figure S2. Identification number of N-glycopeptides in normal human serum under different collision energies.

Figure S3. Tandem MS spectrum of 23SP with HCD fragmentation before enzymatic labeling.

Scheme S1. Dissociation pathways for the GlcNAc (A) and GalNAz (B) in HCD mode.

Figure S4. Zoomed-in spectra of 23SP before enzymatic labeling (A) and 23SP after enzymatic labeling (B).

Figure S5. Venn diagram of identified glycoproteins from current study and previous study in
which deglycosylation method was used.

Figure S6. Representative annotated spectra of α2,6-sialylated glycopeptide (top) and α2,3-
sialylated glycopeptide (bottom) in the protein P01876 from colon cancer patients' serum. Green
circles indicate Hexose, blue squares indicate GlcNAc, olive squares represent GalNAz, purple
diamonds indicate Neu5Ac, and red triangles indicate Fuc.

Figure S7. Representative annotated spectra of α2,6-sialylated glycopeptide (top) and α2,3-
sialylated glycopeptide (bottom) in the protein P01876 from liver cancer patients' serum.

Figure S8. Representative annotated spectrum of α2,6-sialylated glycopeptide in the protein
P01876 from normal human sera.

Table S1. Identified N-glycosite-specific sialoglycans in serum samples from cancer patients and
normal people. SC: serum for colon cancer patients; SL: serum from liver cancer patients; SN:
serum from normal people. "X" indicates detected glycopeptides in a specific sample; "H"
indicates Hexose; "N" indicates GlcNAc; "A" indicates Neu5Ac; "Z" indicates GalNAz; "F"
indicates Fuc.

Abstract

Scheme 1

Figure 1

Figure 2

Figure 3

Figure 4

Trypsin Digestion

HILIC Enrichment

GalNAz Labeling by Glycosyltransferase CgtA

SCE-HCD-MS/MS

α2,3-sialylation

204.09
245.09
274.09
292.10
m/z

α2,6-sialylation

204.09
274.09
292.10
m/z

= Peptides ■ = GlcNAc □ = GalNAc ◯ = Gal ● = Man N₃ = GalNAz ◆ = Neu5Ac

SCE = Stepped Collision Energy

Figure 5

Figure S1

Figure S2

Figure S3

Scheme S1

A

204.08665 →(-H2O)→ 186.07608
- →(-H2O)→ 168.06552 →(-CH2O)→ 138.05495
- →(-C2H2O)→ 144.06552 →(-H2O)→ 126.05495

B

245.08805 →(-H2O)→ 227.07748
- →(-H2O)→ 209.06692 →(-CH2O)→ 179.05635
- →(-C2HN3O)→ 144.06552 →(-H2O)→ 126.05495

Figure S4

Figure S5

Figure S6

Figure S7

Figure S8