Evolutionary relationships among *Massospora* spp. (Entomophthorales), obligate pathogens of cicadas

Angie M. Macias[1], David M. Geiser[2], Jason E. Stajich[3], Piotr Łukasik[4,5], Claudio Veloso[6], DeAnna C. Bublitz[5], Matthew C. Berger[1], Greg R. Boyce[1], Kathie Hodge[7], Matt T. Kasson[1]

[1] Division of Plant and Soil Sciences, West Virginia University, Morgantown, West Virginia, 26506, USA.

[2] Department of Plant Pathology & Environmental Microbiology, The Pennsylvania State University, University Park, PA 16802, USA.

[3] Department of Microbiology and Plant Pathology and Institute for Integrative Genome Biology, University of California, Riverside, California 92521, USA.

[4] Institute of Environmental Sciences, Jagiellonian University, 30-387 Kraków, Poland.

[5] Division of Biological Sciences, University of Montana, Missoula, Montana, 59812, USA

[6] Department of Ecological Sciences, Science Faculty, University of Chile, Santiago, Chile.

[7] Plant Pathology & Plant-Microbe Biology, School of Integrative Plant Science, Cornell University, Ithaca, New York 14853, USA.

Keywords: Zoopagomycota, Entomophthoraceae, Entomopathogen, Invertebrate pathology, *Magicicada*, *Okanagana*, *Platypedia*, *Tettigades*, *Diceroprocta*

Abstract

The fungal genus *Massospora* (Zoopagomycota: Entomophthorales) includes more than a dozen obligate, sexually transmissible pathogenic species that infect cicadas (Hemiptera) worldwide. At least two species are known to produce psychoactive compounds during infection, which has garnered considerable interest for this enigmatic genus. As with many Entomophthorales, the evolutionary relationships and host associations of *Massospora spp.* are not well understood. The acquisition of *M. diceroproctae* from AZ, *M. tettigatis* from Chile, and independent populations of *M. platypediae* from CA and CO provided an opportunity to conduct molecular phylogenetic analyses and morphological studies to investigate if these cicada-infecting fungi represent a monophyletic group and delimit species boundaries in *Massospora*. In a three-locus (EFL+LSU+TUB) phylogenetic analysis, *Massospora* was resolved in a strongly supported monophyletic group containing four well-supported genealogically exclusive lineages, based on two of three methods of phylogenetic inference. There was incongruence among the single-gene trees: two methods of phylogenetic inference recovered trees with either the same topology as the 3-gene concatenated tree (EFL), or a basal polytomy (LSU, TUB). *Massospora levispora* and *M. platypediae* isolates formed a single lineage in all analyses and are synonymized here as a single species, *M. levispora*. *Massospora diceroproctae* was sister to *M. cicadina* in all 3 single gene trees and on an extremely long branch relative to the other *Massospora* species, and even the outgroups, which may reflect either an accelerated rate of molecular evolution, incomplete taxa sampling or some combination of the two. The results of the morphological study presented here indicate that spore measurements are not phylogenetically or diagnostically informative. Despite recent advances in understanding the ecology of *Massospora*, much about its host-range

and diversity is yet to be discovered. The emerging phylogenetic framework can provide a foundation for exploring co-evolutionary relationships with cicada hosts as well as the evolution of behavior-altering compounds.

Introduction

The Entomophthorales (Zoopagomycota) are among the most important arthropod-destroying fungi (Spatafora *et al.* 2016). Many North American Entomophthorales were first described by Thaxter (1888) more than a century ago. Well-known examples include *Entomophthora muscae*, causal agent of "summit disease" of numerous fly genera (Fresenius 1856, Elya *et al.* 2018) and *Entomophaga maimaiga*, a virulent pathogen and biological control agent of gypsy moth (Soper *et al.* 1988, Hajek *et al.* 1990).

Due to the ephemeral nature, obligate lifestyle, and large genome size of the Entomophthorales, these fungi are grossly underrepresented in phylogenetic studies (Spatafora *et al.* 2016, Gryganskyi *et al.* 2017). Only recently have a select few been formally investigated using molecular phylogenetics (Gryganskyi *et al.* 2012, 2013), including the recently described *Arthrophaga myriapodina*, a lethal summit disease pathogen of polydesmid millipedes (Hodge *et al.* 2017) and *Massospora*, an active host transmission pathogen of numerous cicada species (Boyce *et al.* 2019). In total, the Entomophthorales includes some 12 accepted genera with 237 species, including *Massospora* with 13 established species (Index Fungorum and MycoBank).

Massospora was first described anecdotally by Dr. Joseph Leidy (1851), who noted an undescribed fungal disease of periodical cicadas in the eastern U.S.: "[*Magi*]*cicada septendecim* was subject to a fungous disease" and observed that "the posterior part of the abdomen…in several instances [was] filled with a mass of oval spore-like bodies" (Leidy 1851). *Massospora*

was formally established by Peck (1879) with the description of *M. cicadina* from a periodical cicada (*Magicicada septendecim*) collected in New York, USA in 1877. Following Peck's description, Thaxter (1888) recognized *Massospora* as a member of the Entomophthorales. Research on *Massospora* gained momentum in the twentieth century with spore development studies (Speare 1921, Goldstein 1929) and the description of ten new species in the western Hemisphere (Ciferri *et al.* 1957, Soper 1963, 1974), two species from Australia and Afghanistan (Soper 1981), plus undescribed *Massospora* species from *Platypleura* sp. (Kobayashi 1951) and *Meimua* sp. (Ohbayashi *et al.* 1999) in Japan. Today, *Massospora* includes more than a dozen obligate, sexually transmissible, pathogenic species that attack at least 24 cicada species worldwide (Soper 1963, 1974, 1981, Cooley *et al.* 2018) (Table 1). Nearly all extant *Massospora* species are associated with a single cicada genus with two exceptions. *Massospora cicadettae* is reported from *Plerapsalta incipiens, Chelapsalta puer,* and *Cicadetta. spp.* (Table 1) and *M. platypediae / M. levispora,* based on existing phylogenetic data, represent a single species infecting two genera of annual cicadas, *Platypedia* sp. and *Okanagana* sp. (Boyce *et al.* 2019). Generally, specimens of *Massospora* have been identified based on the cicada host they are found on, but this method of identification has proven unreliable given the recent finding that *M. levispora* and *M. platypediae* comprise a single species that occupies a broader geographic and host range than previously reported (Boyce *et al.* 2019). Until the host associations and fungus names are confirmed with detailed molecular studies, IDs based solely on host associations should be viewed with skepticism.

The life cycles of individual *Massospora* species are closely tied to the life cycle of the cicada host. Mature cicada nymphs are believed to be infected by resting spores encountered underground during construction of their vertical emergence burrows (Soper *et al.* 1976a). These

nymphs then emerge, eclose into adults, and over a period of days develop infections in their abdomen. These infections become more conspicuous as the fungus destroys the cicada's abdominal intersegmental membranes, inciting a progressive sloughing off of sclerites that reveals a large fungal mass (Fig. 1). Conidia are passively disseminated during mating attempts or flights, or possibly in crowded settings where high densities of cicadas promote close contact (Soper 1963, Cooley *et al.* 2018). Cicadas infected by conidia develop secondary infections (Soper *et al.* 1976b, Cooley *et al.* 2018), resulting in the production of resting spores inside cicada hosts. These resting spores are incorporated back into the soil to infect new cohorts of cicadas as they emerge in later years (Soper *et al.* 1976a).

Complex infection and transmission strategies that involve manipulation of host behavior are notable in the Entomophthorales, including several cases of summiting behaviors and active host transmission (Roy *et al.* 2006, Hughes *et al.* 2016, Gryganskyi *et al.* 2017, Hodge *et al.* 2017, Boyce *et al.* 2019). The rarer of these two transmission behaviors, active host transmission (AHT), involves infected living hosts that directly transmit spores to new hosts (Roy *et al.* 2006). AHT behavior in *Massospora* is thought to be chemically induced (Boyce *et al.* 2019), and includes hypersexual behavior where infected male cicadas mimic female-specific behaviors to attract copulation attempts from other males (Cooley *et al.* 2018). *Massospora* and *Strongwellsea,* a fly pathogen, are the only two genera where all species are known to induce AHT behavior in their hosts, although AHT has also been reported in select species of *Entomophthora* (*E. erupta* and *E. thripidum*) and *Entomophaga* (*E. kansana*) (Roy *et al.* 2006). However, the identity and phylogenetic placement of these latter species have not been molecularly resolved (Gryganskyi *et al.* 2012, 2013). Given this taxonomic uncertainty coupled with the occurrence of both AHT and summit disease in *Entomophthora* and *Entomophaga,* the

evolutionary history of AHT among members of the Entomophthoraceae should be further investigated (Boyce *et al.* 2019). More specifically, is AHT the ancestral state for the Entomophthoraceae or has it evolved multiple times among *Massospora, Strongwellsea, Entomophthora,* and *Entomophaga*?

Multi-locus phylogenetics using few loci can serve as a rapid, cost-effective screening tool to inform further research using genomic, transcriptomic and metabolomic approaches. Ultimately, genomics-based approaches offer superior phylogenetic resolution, but Entomophthorales genomes are difficult to obtain for several reasons. Compared to other fungi, some Entomophthorales genomes are massive in size, including the publically available *Entomophthora muscae* genome (600 Mb for NCBI: PRJNA479887) and *Zoophthora radicans* genome (655 Mb for JGI: ATCC 208865) (Nordberg *et al.* 2014, Elya *et al.* 2018). Additionally, many Entomophthorales are unculturable and therefore impure and potentially degraded environmental samples must be used. Phylogenetic studies can also help populate NCBI sequence data repositories, which are significantly underpopulated for members of the Entomophthoraceae. In total, GenBank's Nucleotide sequence repository has 616 DNA sequences, excluding genomes, representing only about 20% of described species. More than 45% of these sequences are from just three taxa: *Pandora neoaphidis*, *Entomophthora muscae* and *Zoophthora radicans*. Additionally, 30% of the 616 sequences deposited are ITS or SSU ribosomal RNA gene sequences, which are not suitable for accurate phylogenetic analyses (Tang *et al.* 2007, Schoch *et al.* 2012, Demirel 2016).

In this study, we used molecular phylogenetics and morphology to further investigate three findings of Boyce *et al.* (2019): 1) *Massospora* is monophyletic; 2) *M. levispora* and *M. platypediae* are not genealogically exclusive; and 3) *M. levispora* and *M. platypediae* are not

distinguishable based on spore measurements. To test the monophyly of *Massospora*, we included newly obtained specimens from Chile (*M. tettigatis*) and Arizona (*M. diceroproctae*) as well as additional outgroup taxa. To determine if synonymization of *M. levispora* / *M. platypediae* is warranted, we sequenced additional isolates from previously sampled sites and isolates of *M. platypediae* from new sites. Last, we measured spores for new isolates of *M. platypediae*, *M. tettigatis,* and *M. diceroproctae,* and compared these measurements to historical spore measurements (Soper 1963, 1974, 1981).

Materials and Methods

Sample collection & DNA extraction

The following designations are used throughout the remainder of the methods: *M. cicadina = Mc*, *M. diceroproctae = Md*, *M. levispora = Ml*, *M. platypediae = Mp*, and *M. tettigatis = Mt*.

Infected cicadas were obtained from various locations and collectors (Table 2). Samples from each collector were stored differently, with some samples stored dry at room temperature, some frozen in RNAlater (Invitrogen, Grand Island, NY, USA) or 95% ethanol, and some frozen dry immediately upon collection. The fungal plug on each infected cicada was sampled using a sterile scalpel, or by centrifuging a solution of loose spores into a pellet. DNA was extracted using a modified Wizard kit (Short *et al*. 2015). Samples were macerated in 1.5-mL microcentrifuge tubes (Eppendorf, Hamburg, Germany) with 600 µl of Nuclei Lysis Solution (Promega, Madison, WI, USA) and incubated at 65°C for 30 minutes, vortexing at 15 minutes. After cooling briefly, 200 µl of Protein Precipitation Solution (Promega, Madison, WI, USA) was added, and samples were vortexed vigorously for 10 seconds. Then, samples were centrifuged for 3 minutes at 14,000 RPM, and the supernatant was collected and moved to fresh

1.5-mL tubes with 600 µl of 99.9% isopropanol. Tubes containing the protein pellet were discarded. Sample tubes containing isopropanol were gently inverted several times and centrifuged again for 1 minute at 14,000 RPM. The supernatant was discarded, leaving a DNA pellet behind. Tubes were then loaded with 600 µl of 70% ethanol and centrifuged for 1 minute at 14,000 RPM. Supernatant was again discarded, and the DNA pellets were left to dry at room temperature for 20-30 minutes. Finally, the DNA was resuspended in 100 µl of warmed (65°C) Elution Buffer (Alfa Aesar, Ward Hill, MA, USA) and stored at -20°C.

PCR and sequencing

We targeted sequencing of 28S rDNA (LSU), 18S rDNA (SSU), elongation factor 1-α-like gene (EFL), and β-tubulin (TUB) for each sample listed in Table 2. We used existing data from GenBank for 6 reference strains. Additionally, six gene sequences were extracted from 2 assembled metagenomes from Boyce *et al.* (2019). The PCR reaction volumes are as follows: 12.5 µl MyTaq™ Master Mix (Bioline, London, UK), 10 µl molecular-grade water (G-Biosciences, St. Louis, MO, USA), 1 µl (10 µM in IDTE, pH 8.0) each of forward and reverse primers (IDT, Coralville, IA, USA), and 1 µl of DNA template for a final reaction volume of 25.5 µl. Primer names, sequences, and full PCR protocols are listed in Supplemental Table 1. PCR products were visualized via gel electrophoresis on a 1.5% w/v agarose (Amresco, Solon, OH, USA) gel with 0.5% Tris-Borate-EDTA buffer (Amresco, Solon, OH, USA). SYBR Gold (Invitrogen, Grand Island, NY, USA) was used as the nucleic acid stain, and bands were visualized on a UV transilluminator (Bio-Rad, Hercules, CA, USA). Prior to sequencing, PCR products were purified using ExoSAP-IT (Affymetrix, Santa Clara, CA, USA): 2.2 µl of ExoSAP and 6 µl of PCR product in a 2-step reaction of 15 minutes at 37°C, followed by 15

minutes at 80°C. Purified products were Sanger-sequenced (Eurofins, Huntsville, AL, USA) with

the same primers used for PCR. Sequences generated during this study are deposited in GenBank

(Table 2).

Alignments, model selection, and phylogenetic analyses

Chromatograms were quality-checked using default parameters, clipped, and manually

corrected in CodonCode Aligner v. 5.1.5. Each gene was aligned separately using MAFFT

(Katoh and Standley 2013) on the Guidance2 server (http://guidance.tau.ac.il/ver2/, Landan and

Graur 2008, Sela *et al.* 2015), and individual residues with Guidance scores <0.5 were masked.

An intron in LSU (positions 299-478) was deleted. Alignments are available here:

https://github.com/ammacias18/MassoPhylogenetics.

Nucleotide substitution models were chosen using AICc scores in Model Test in MEGA

v. 7.0.16 (Kumar *et al*. 2016). Each individual gene, and a concatenated alignment of the three

genes, were used in a maximum likelihood (ML) analysis (RAxML v. 8.2.12, Stamatakis 2014),

a maximum parsimony (MP) analysis (PAUP* v. 4.0a build 166, Swofford 2002), and a

Bayesian inference (BI) analysis (MrBayes v. 3.2.5, Ronquist *et al.* 2012), for a total of 12

analyses. The default parameters of each software package were used, unless otherwise noted

(see code and notes in Supplemental File 1). In brief, for ML analyses, an appropriate model was

chosen, partitions were applied (for each gene in the concatenated analysis only), 1,000 bootstrap

replicates were used, and the best-scoring tree was identified and bootstrapped in a single run.

For MP analyses, a heuristic search with TBR swapping was used to identify the set of most-

parsimonious trees, and 1,000 bootstrap replicates were used. For BI analyses, MrBayes was

allowed to select a substitution model for each dataset, and rates were set based on results from

Model Test. Runs were stopped when the standard deviation of split frequencies fell below 0.01, then checked for convergence in Tracer v. 1.7.1 (Rambaut *et al.* 2018).

One additional tree was generated: a single-gene SSU tree using the same isolates as the 3-gene dataset, which was generated using all three methods of phylogenetic inference (see detailed methods above).

All resulting trees are available as .tre files (https://github.com/ammacias18/MassoPhylogenetics). Trees were viewed and prepared for publication using FigTree v. 1.4.4 (Rambaut 2017) and Inkscape v. 0.92.2 (https://www.inkscape.org/).

Morphological study

To examine spore morphology, a portion of select fungal plugs (n = 63) was harvested with a sterile scalpel and mounted on a slide in lactophenol or lactophenol+cotton blue for examination with light field microscopy. Cover slips were fastened with nail polish to allow slides to be archived and re-examined when necessary. Slides were examined and photographed using a Nikon Eclipse E600 compound microscope (Nikon Instruments, Melville, NY, USA) equipped with a Nikon Digital Sight DS-Ri1 high-resolution microscope camera. A sample of 25 spores from each slide mount were measured using Nikon NIS-Elements BR3.2 imaging software. For conidial samples, the length and width of 25 conidia were recorded, and for resting spore samples, two diameter measurements were taken and averaged for 25 resting spores. Conidial measurements were taken from 45 isolates: $Mc = 12$, $Md = 4$, $Ml = 8$, $Mp = 20$, and $Mt = 1$. Resting spore measurements were taken from 18 isolates: $Mc = 9$, $Md = 2$, $Ml = 1$, $Mp = 2$, and $Mt = 4$. Raw spore measurements are available in Supplemental Table 2.

Spore measurement data was analyzed using packages dplyr (Wickham *et al.* 2019), ggplot2 (Wickham 2016), car (Fox and Weisberg 2012), userfriendlyscience (Peters *et al.* 2018), and gplots (Warnes *et al.* 2019) in R v. 3.6.1 (R Core Team, 2019). Normality was assessed using density plots and the Shapiro-Wilkes test, and equality of variance was assessed using Levene's Test and the Fligner-Killeen test. ANOVAs and Welch's ANOVAs (where appropriate) were performed to check for differences in spore measurements across species, and the Tukey or Games-Howell multiple-comparisons post-hoc tests (respectively) were used to identify the significant pairwise differences. A p-value < 0.05 was considered significant for all analyses. Reported P-values are Bonferroni-corrected. R code and outputs are available in Supplemental File 1.

Results

Phylogenetics

The following designations are used throughout the remainder of the results: *M. cicadina* = *Mc*, *M. diceroproctae* = *Md*, *M. levispora* = *Ml*, *M. platypediae* = *Mp*, and *M. tettigatis* = *Mt*.

To infer evolutionary relationships among sampled taxa, several phylogenetic analyses were performed. The three individual gene trees (EFL, LSU, TUB) as well as the concatenated 3-gene tree resolved *Massospora* as a monophyletic ingroup. In a separate analysis, SSU placed *Md* among the outgroup taxa, and the remainder of *Massospora* was left monophyletic. In all trees, *Md* resolved as a very long branch, and we attribute its occasional displacement to be a long-branch artefact, disproportionately based on signal from the SSU locus. A visual scan of all alignments indicated that differences between *Md* and other *Massospora* were distributed across all four loci, in a somewhat patchy distribution, with no indication of insertions, deletions or

alignment errors being the basis of its apparent divergence. This observation together with other indications that SSU performs poorly as a phylogenetic marker for fungi (Tang *et al.* 2007, Schoch *et al.* 2012, Demirel 2016) led us to remove SSU from the analyses shown (For SSU results see: https://github.com/ammacias18/MassoPhylogenetics).

Two of three methods of phylogenetic inference for the three-gene (EFL+LSU+TUB) concatenated dataset resolved all five *Massospora* species into a strongly supported monophyletic group (Fig. 2). The third method of phylogenetic inference, MP, showed very weak bootstrap support (27%) for the genus (Fig. 2, https://github.com/ammacias18/MassoPhylogenetics). An additional MP analysis constraining *Massospora* to be monophyletic resulted in a tree 1259 steps in length (data not shown), only four steps longer than the unconstrained analysis. Three of the *Massospora* species, *Mc*, *Md*, and *Mt*, were all genealogically exclusive and had strong bootstrap support. *Massospora levispora* and *M. platypediae* did not resolve as genealogically exclusive, and instead together formed a single well-supported lineage. Within *Massospora, Mc* and *Md* formed a clade sister to the *Ml* / *Mp* lineage. *Massospora tettigatis* was recovered as the earliest diverging species in our sample of species.

Additional LSU sequences from specimens of *Mp* from *P. putnami* collected in 2013 in CO were compared using NCBI BLAST to *Mp* LSU sequences used in the 3-gene concatenated dataset. Isolates from CO were identical to isolates from CA and NM (Supplemental Table 3). Additional LSU sequences from specimens of *Mt* from 3 additional *Tettigades* spp. from Chile were compared using NCBI BLAST to *Mt* LSU sequences used in the 3-gene concatenated dataset. These comparisons revealed *Mt* is a single species capable of infecting diverse *Tettigates*

species (Supplemental Table 3). These *Mp* and *Mt* isolates were excluded from the phylogenetic analyses due to insufficient sequence data for the other loci used.

Morphological study

Morphological studies were conducted to permit comparisons between isolates used in this study and previously reported measurements (Soper 1963, 1974, 1981) and among species. Conidia and resting spore measurements were acquired from *Mp*-infected wing-banger cicadas (*Platypedia putnami*) from CA and CO, and from *Mt*-infected *Tettigades* cicadas and *Md*-infected *Diceroprocta* cicadas. Spore measurements for *Mc*, *Mp*, and *Ml* previously reported by Boyce *et al.* (2019) were also included in this study.

Conidial measurements are summarized in Fig. 3 with raw spore measurements available in Supplemental Table 2. Mean, standard deviation, minimum, and maximum values for each species are reported in Supplemental Table 4. Conidial length measurements are presented as follows: Mean conidial length ± standard deviation for each species. *Mc* = 16.6 µm ± 1.92 µm, *Md* = 14.7 µm ± 2.15 µm, *Ml* = 14.3 µm ± 2.00 µm, *Mp* = 12.7 µm ± 1.78 µm, and *Mt* = 15.8 µm ± 1.87 µm. Conidial widths are reported in the same format as above, and are as follows: *Mc* = 14.9 µm ± 1.57 µm, *Md* = 7.1 µm ± 0.95 µm, *Ml* = 8.8 µm ± 0.80 µm, *Mp* = 8.0 µm ± 0.84 µm, and *Mt* = 11.3 µm ± 1.08 µm. Comparisons of mean conidial length and width among species and their statistical significance are shown in Fig. 3. Overall, mean conidial width was significantly affected by species ($p < 0.001$, Welch's ANOVA), and each individual pairwise comparison was also significant (all $p < 0.01$, Games-Howell post-hoc test). Mean conidial length was also significantly affected by species ($p < 0.001$, ANOVA), but mean lengths

overlapped among several species (*Mt-Mc* p = 0.55, *Ml-Md* p = 1.00, *Mt-Md* p = 0.09, all others p < 0.01; Tukey's post-hoc test) (Fig. 3).

Unfortunately, our spore measurements cannot be statistically compared with those reported by Soper (1963, 1974, 1981) due to the fact that Soper only reported measurement means, minimums, and maximums, but not standard deviation or sample size (raw data is also unavailable). Regardless, our study found that all Soper's mean conidial measurements fell within our reported range for each *Massospora* species (Supplemental Table 4), but not always within one standard deviation of our mean: not for *Md* conidia length, *Mp* conidia length and width, or *Mt* conidial length and width.

Resting spore measurements are summarized in Figure 3 with raw measurements available in Supplemental Table 2. Mean, standard deviation, minimum and maximum values for each species are reported in Supplemental Table 4. Resting spore diameter is reported in the same format as above, and are as follows: $Mc = 42.7\,\mu m \pm 2.71\,\mu m$, $Md = 37.8\,\mu m \pm 3.50\,\mu m$, $Ml = 39.8\,\mu m \pm 1.99\,\mu m$, $Mp = 38.3\,\mu m \pm 3.26\,\mu m$, and $Mt = 44.0\,\mu m \pm 3.05\,\mu m$. Overall, mean resting spore diameter was significantly affected by species (p < 0.001, Welch's ANOVA), but mean resting spore diameters overlapped among several species (*Ml-Md* p = 0.28, *Mp-Md* p = 1.00, *Mp-Ml* p = 0.44, all others p ≤ 0.02; Games-Howell post-hoc test) (Fig. 3). Relative to Soper's measurements, mean resting spore diameters fell within our reported range for *Mc* but not *Ml* and *Mt*, and not always within one standard deviation of our mean: not for *Ml* resting spore diameter or *Mt* resting spore diameter. Soper did not observe a resting spore stage for *Md* and *Mp* (Supplemental Table 4).

In addition to spore measurements, conidial plug color varied among species: *Md* plugs from specimens were violet to purple in color, compared to creamy white to brown plugs from all other species included in this study (Fig. 1).

Discussion

In a recent study (Boyce *et al.* 2019), three species of *Massospora* were found to form a monophyletic group containing two genealogically exclusive lineages. In this work, we confirmed the monophyly of *Massospora*, even with the addition of two previously unavailable described *Massospora* species. At least 4 *Massospora* species are now well-supported according to the criteria of genealogical concordance and non-discordance (Taylor *et al.* 2000; Dettman *et al.* 2003). One of these lineages is composed of two previously described *Massospora* species, *Ml* and *Mp*, and is synonymized here as *M. levispora* based on the findings of this study and previous work (Boyce *et al.* 2019).

The incongruence between spore morphology and molecular phylogenetics regarding the *Ml* / *Mp* lineage is intriguing. *Ml* conidia from *O. rimosa* are significantly longer ($p < 0.01$) and wider ($p < 0.01$), compared to *Mp* counterparts from *P. putnami*. Additionally, Soper's mean conidial length and width measurements for *Mp* are not within one standard deviation of our measurements, nor are his resting spore measurements for *Ml* (Supplemental Table 4). The mountant used for spore measurements may affect spore shape and size but it is not known what mountant was used by Soper (1963, 1974, 1981). Other studies of Entomophthorales used either lactophenol or aceto-orcein (Humber 1976, Soper *et al.* 1988, Gryganskyi *et al.* 2013, Hodge *et al.* 2017). Differences among species (Fig. 3) and in comparison to Soper's measurements (Supplemental Table 4) may also be due to differing sample ages and storage: in our study, *M.*

levispora samples were stored in ethanol for 20 years while *M. platypediae* were stored dry and only for a few years. It is not known how Soper's samples were stored or for how long (Soper 1963, 1974, 1981). One study examining sample age and mountant used in *Strongwellsea* found that these factors have an interacting effect on spore dimensions (Humber 1976). Sample size may also be important (n = 8 for *Ml*, n = 20 for *Mp*). Previous work by Boyce *et al*. (2019) used fewer populations of *Mp* (14 isolates from one population), and found considerable overlap in both conidia and resting spore measurements for *Mp* and *Ml*, although these measurements were not statistically compared. Taken together, these studies suggest that there may be population-level variation in *Mp* spore sizes, such that sampling too few populations will result in misleading conclusions. However, this does not explain the incongruence of our phylogenetic study and morphologic study, with respect to *Mp* and *Ml*. Further sampling is needed.

Massospora diceroproctae was on an extremely long branch relative to the other *Massospora* species in both the 3-gene concatenated tree and the single-gene trees (Fig. 2), sometimes longer than even the total branch length separating the genus *Massospora* from the most distantly related outgroups. In several parsimony-derived trees, *Md* fell among the outgroups. Some of the incongruence between MP and the other methods of phylogenetic inference observed in this study can be explained by long branch attraction or LBA (Felsenstein 1978) acting on the *Md* clade and the outgroup clade. This result is not entirely surprising, given that MP is often more susceptible to LBA than other phylogenetic methods (O'Connor *et al.* 2010). In the 3-gene ML and BI concatenated trees (https://github.com/ammacias18/MassoPhylogenetics), LBA cannot explain *Mc* and *Md* forming a clade, because *Mc* is not on a long branch in this study, and did not appear on a long branch in Boyce *et al*. (2019) either, in a tree with only *Mc*, *Ml*, and *Mp*.

One of the possible explanations for the long branch lengths and inconsistent resolution of *Massospora* in this study is that, *Md* may have experienced an accelerated rate of molecular evolution compared to all other *Massospora* species. A second, perhaps more likely explanation for long branches associated with *Md* is that the closest relatives of *Md* were not sampled here, due either to unavailability of samples, their undiscovered status, or extinction. Only five of the twelve described *Massospora* were available for this study, and there may be undiscovered extant taxa that would disrupt the long branches associated with *Md*. *Massospora* is not the only member of the Entomophthorales where long branches have been observed: *Batkoa* was recovered on a longer branch compared to other taxa in two separate analyses (Gryganskyi *et al*. 2012, Hodge *et al*. 2017). Similar long-branch taxa have been observed in other early diverging fungi outside the Entomophthorales, which can be partially explained by the limited taxon sampling compared to members of Basidiomycota and Ascomycota (James *et al.* 2006b, Jones *et al.* 2011).

Two *Massospora* species treated in this study, *Mt* and *Ml* / *Mp* have cicada hosts both belonging to the subfamily Tibicinae, while the hosts of *Md* and *Mc* belong to two other subfamilies, Cicadinae and Cicadettinae, respectively (Sanborn 2013, Łukasik *et al.* 2018, Marshall *et al*. 2018). Our results indicate that all three cicada subfamilies are susceptible to *Massospora*, but *Massospora* has only been molecularly confirmed from cicadas in the New World. All three subfamilies contain dozens of genera and species that have never been formally surveyed for *Massospora*. Before cophylogenetic analyses of *Massopora* and their cicada hosts can be performed to test for evidence of parallel cladogenesis, the relationships among *Massospora* species need to be better resolved through the addition of more taxa and other loci.

Given the previous findings by Boyce *et al.* (2019) that two species of *Massospora*, *Mc*

and the single lineage containing *Ml* / *Mp*, produce psychoactive compounds during host infection, and the findings of this study that within *Massospora*, *Md*, *Mc*, and *Ml* / *Mp* form a clade, *Md* is a likely candidate worth investigating for similar biologically active compounds. Observations of *Md*-infected *Diceroprocta semicincta* in AZ revealed altered calling patterns in these cicadas despite continued mating attempts (Dr. DeAnna Bublitz, unpublished observations). A separate observation of *M. diceroproctae*-infected *Diceroprocta* sp. from Big Bend National Park showed behavioral changes in infected individuals: elevated mating effort in terms of time spent signaling in males and increased likelihood to be in contact with a conspecific for males and females (Dr. Jon Hastings, unpublished observations). Few observations exist on the behavior of *Massospora*-infected *Tettigades* cicadas, although *Mt*-infected cicadas continue mating attempts (Dr. Piotr Łukasik, unpublished observations).

The results of the morphological study presented here indicate that spore measurements may not be useful for species level identifications. Unfortunately, the numbers of isolates sampled for many of these species were insufficient to confidently conclude whether differences truly exist. In general, trends observed across spore measurements were incongruent with the evolutionary relationships proposed by molecular phylogenetics. For example, comparisons between *Ml* isolates and *Mp* isolates uncovered significant differences in conidia length ($p < 0.01$) and width ($p < 0.01$) (Fig. 3) despite forming a single lineage based on multi-locus sequence data (Fig. 2). However, resting spore diameter was not significantly different between *Ml* and *Mp* ($p = 0.44$).

In less than a decade, the research on Entomophthorales has grown significantly, leading to breakthrough discoveries on the biology and ecology of several members of this long-neglected group (Grell *et al.* 2011; Małagocka *et al.* 2015, De Fine Licht *et al.* 2017, Arnesen *et*

al. 2018, Elya *et al.* 2018, Wronska *et al.* 2018, Boyce *et al.* 2019). Still, the vast majority of the Entomophthorales remain understudied. Despite recent advances in understanding the ecology of *Massospora*, much about the host-range and diversity of this genus is yet to be discovered. The emerging phylogenetic framework for *Massospora* provides a starting point for co-evolutionary studies with their cicada hosts as well as lays a foundation for deciphering the evolution of behavior-altering compounds among *Massosopora* and close allies.

Acknowledgements

AM was supported by The Ruby Distinguished Doctoral Fellows Program, Morgantown, WV, P.L by National Geographic Society grant 9760-15, J.E.S. by NSF DEB 1441715, and M.T.K. by funds from West Virginia Agricultural and Forestry Experiment Station.

Literature Cited

Arnesen JA, Małagocka J, Gryganskyi A, Grigoriev IV, Voigt K., Stajich JE, De Fine Licht, HH. 2018. Early diverging insect pathogenic fungi of the order Entomophthorales possess diverse and unique subtilisin-like serine proteases. G3:Genes|Genomics|Genetics 8:3311–3319.

Boyce GR, Gluck-Thaler E, Slot JC, Stajich JE, Davis WJ, James TY, Cooley JR, Panaccione DG, Eilenberg J, De Fine Licht HH, Macias AM, Berger MC, Wickert KL, Stauder CM, Spahr EJ, Maust MD, Metheny AM, Simon C, Kritsky G, Hodge KT, Humber RA, Gullion T, Short DPG, Kijimoto T, Mozgai D, Arguedas N, Kasson MT. 2019. Psychoactive plant- and

mushroom-associated alkaloids from two behavior modifying cicada pathogens. Fungal Ecology 41:147–164.

Ciferri R, Machado AA, Vital AF. 1957. A new species of the genus *Massospora* with an *Allomyces* species. Atti dell'Instituto Botanico della Università e Laboratorio Crittogamico di Pavia 5:15–22.

Cooley JR, Marshall DC, Hill KB. 2018. A specialized fungal parasite (*Massospora cicadina*) hijacks the sexual signals of periodical cicadas (Hemiptera: Cicadidae: *Magicicada*). Scientific Reports 8:1432.

De Fine Licht HH, Jensen AB, Eilenberg J. 2017. Comparative transcriptomics reveal host-specific nucleotide variation in entomophthoralean fungi. Molecular Ecology 26:2092–2110.

Demirel R. 2016. Comparison of rDNA regions (ITS, LSU, and SSU) of some *Aspergillus, Penicillium,* and *Talaromyces* spp. Turkish Journal of Botany 40:576–583.

Dettman JR, Jacobson DJ, Taylor JW. 2003. A multilocus genealogical approach to phylogenetic species recognition in the model eukaryote *Neurospora*. Evolution 57:2703–2720.

Elya C, Lok TC, Spencer QE, McCausland H, Martinez CC, Eisen MB. 2018. Robust manipulation of the behavior of *Drosophila melanogaster* by a fungal pathogen in the laboratory. eLife 7:34414.

Felsenstein J. 1978. Cases in which parsimony or compatibility methods will be positively misleading. Systematic Zoology 27:401–410.

Fox J, Weisberg S, Adler D, Bates D, Baud-Bovy G, Ellison S, Firth D, Friendly M, Gorjanc G, Graves S, Heiberger R. 2012. Package 'car'. Vienna: R Foundation for Statistical Computing.

Fresenius G. 1856. Notiz, Insecten-Pilze betreffrend. Bot. Zeitg. 14: 882.

Gargas A, Taylor JW. 1992. Polymerase chain reaction (PCR) primers for amplifying and sequencing nuclear 18S rDNA from lichenized fungi. Mycologia 84:589–592.

Goldstein B. 1929. A cytological study of the fungus *Massospora cicadina*, parasitic on the 17□year cicada, *Magicicada septendecim*. American Journal of Botany 16:394–401.

Grell MN, Jensen AB, Olsen PB, Eilenberg J, Lange L. 2011. Secretome of fungus-infected aphids documents high pathogen activity and weak host response. Fungal Genetics and Biology 48: 343–352.

Gryganskyi AP, Humber RA, Smith ME, Miadlikovska J, Wu S, Voigt K, Walther G, Anishchenko IM, Vilgalys R. 2012. Molecular phylogeny of the Entomophthoromycota. Molecular Phylogenetics and Evolution 65:682–694.

Gryganskyi AP, Humber RA, Smith ME, Hodge K, Huang B, Voigt K, Vilgalys R. 2013. Phylogenetic lineages in Entomophthoromycota. Persoonia 30:94–105.

Gryganskyi AP, Mullens BA, Gajdeczka MT, Rehner SA, Vilgalys R, Hajek AE. 2017. Hijacked: co-option of host behavior by entomophthoralean fungi. PLoS Pathogens 13:e1006274.

Hajek AE, Humber RA, Elkinton JS, May B, Walsh SR, Silver JC. 1990. Allozyme and restriction fragment length polymorphism analyses confirm *Entomophaga maimaiga* responsible for 1989 epizootics in North American gypsy moth populations. Proceedings of the National Academy of Sciences 87:6979–6982.

Hodge KT, Hajek AE, Gryganskyi A. 2017. The first entomophthoralean killing millipedes, *Arthrophaga myriapodina* n. gen. n. sp., causes climbing before host death. Journal of Invertebrate Pathology 149:135–140.

Hughes DP, Araújo JP, Loreto RG, Quevillon L, De Bekker C, Evans HC. 2016. From so simple a beginning: the evolution of behavioral manipulation by fungi. In: Lovett B, St. Leger RJ, eds. Advances in Genetics. New York: Academic Press. p. 437–469.

Humber RA. 1976. The systematics of the genus *Strongwellsea* (Zygomycetes: Entomophthorales). Mycologia 68:1042–1060.

Index Fungorum. Web application at http://www.indexfungorum.org/names/names.asp. (Accessed 17 May 2019).

James TY, Kauff F, Schoch CL, Matheny PB, Hofstetter V, Cox CJ, Celio G, Gueidan C, Fraker E, Miadlikowska J, Lumbsch HT. 2006a. Reconstructing the early evolution of Fungi using a six-gene phylogeny. Nature 443:818–822.

James TY, Letcher PM, Longcore JE, Mozley-Standridge SE, Porter D, Powell MJ, Griffith GW, Vilgalys R. 2006b. A molecular phylogeny of the flagellated fungi (Chytridiomycota) and description of a new phylum (Blastocladiomycota). Mycologia 98: 860–871.

Jones MDM, Forn I, Gadelha C, Egan MJ, Bass D, Massana R, Richards TA. 2011. Discovery of novel intermediate forms redefines the fungal tree of life. Nature 474: 200–203.

Kauff F, Lutzoni F. 2002. Phylogeny of the Gyalectales and Ostropales (Ascomycota, Fungi): among and within order relationships based on nuclear ribosomal RNA small and large subunits. Molecular Phylogenetics and Evolution 25:138–156.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Molecular Biology and Evolution 30:772–780.

Keeling PJ, Luker MA, Palmer JD. 2000. Evidence from beta-tubulin phylogeny that microsporidia evolved from within the fungi. Molecular Biology and Evolution 17:23–31.

Kobayashi Y. 1951. Notes on fungi. (1) On the newly found genus *Massospora* from Japan. The Journal of Japanese Botany 26:117–119.

Kumar S, Stecher G, Tamura K. 2016. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. Molecular Biology and Evolution 33:1870–1874.

Landan G, Graur D. 2008. Local reliability measures from sets of co-optimal multiple sequence alignments. Pacific Symposium on Biocomputing 13:15–24.

Leidy J. 1851. (Note on fungus disease of *Cicada septendecim*.) Proceedings of the Academy of Natural Sciences Philadelphia 5:235.

Łukasik P, Chong RA, Nazario K, Matsuura Y, Bublitz DA, Campbell MA, Meyer MC, Van Leuven JT, Pessacq P, Veloso C, Simon C. 2018. One hundred mitochondrial genomes of cicadas. Journal of Heredity 110:247–256.

Małagocka, J, Grell MN, Lange L, Eilenberg J, Jensen AB. 2015. Transcriptome of an entomophthoralean fungus (*Pandora formicae*) shows molecular machinery adjusted for successful host exploitation and transmission. Journal of Invertebrate Pathology 128:47–56.

Marshall DC, Moulds M, Hill KB, Price BW, Wade EJ, Owen CL, Goemans G, Marathe K, Sarkar V, Cooley JR, Sanborn AF. 2018. A molecular phylogeny of the cicadas (Hemiptera: Cicadidae) with a review of tribe and subfamily classification. Zootaxa.4424:1–64.

MycoBank. Web application at http://www.mycobank.org/quicksearch.aspx. (Accessed 17 May 2019).

Nordberg H, Cantor M, Dusheyko S, Hua S, Poliakov A, Shabalov I, Smirnova T, Grigoriev IV, Dubchak, I. 2014. The genome portal of the Department of Energy Joint Genome Institute: 2014 updates. Nucleic Acids Research 42:26–31.

Ohbayashi T, Sato H, Igawa S. 1999. An epizootic on *Meimuna boninensis* (Distant) (Homoptera: Cicadidae), caused by *Massospora* sp. (Zygomycetes: Entomophthorales) and *Nomuraea cylindrospora* (Deuteromycotina: Hyphomycetes) in the Ogasawara (Bonin) Islands, Japan. Applied Entomology and Zoology 34:339–343.

O'Connor T, Sundberg K, Carroll H, Clement M, Snell Q. 2010. Analysis of long branch extraction and long branch shortening. BMC Genomics 11:S14.

Peck C. 1879. *Massospora cicadina* n.g. et sp. Thirty-first Report of State Botanist of New York, 44.

Peters GJ, Verboon P, Green J. 2018. Package 'userfriendlyscience'. Vienna: R Foundation for Statistical Computing.

R Core Team. 2019. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Rambaut A. 2017. FigTree-version 1.4.3, a graphical viewer of phylogenetic trees.

Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. 2018. Posterior summarization in Bayesian phylogenetics using Tracer 1.7. Systematic Biology 67:901–904.

Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61(3):539–542.

Roy HE, Steinkraus DC, Eilenberg J, Hajek AE, Pell JK. 2006. Bizarre interactions and endgames: entomopathogenic fungi and their arthropod hosts. Annual Review of Entomology 51:331–357.

Sanborn A. 2013. Catalogue of the Cicadoidea (Hemiptera: Auchenorrhyncha). New York: Academic Press. p. 1–1002.

Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, Chen W. 2012. Fungal Barcoding Consortium. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. Proceedings of the National Academy of Sciences 109:6241–6246.

Sela I, Ashkenazy H, Katoh K, Pupko T. 2015. GUIDANCE2: accurate detection of unreliable alignment regions accounting for the uncertainty of multiple parameters. Nucleic Acids Research 43: W7–W14.

Short DP, Double M, Nuss DL, Stauder CM, MacDonald W, Kasson MT. 2015. Multilocus PCR assays elucidate vegetative incompatibility gene profiles of *Cryphonectria parasitica* in the United States. Applied Environmental Microbiology 81:5736–5742.

Soper RS. 1963. *Massospora levispora*, a new species of fungus pathogenic to the cicada, *Okanagana rimosa*. Canadian Journal of Botany 41:875–878.

Soper RS. 1974. The genus *Massospora*, entomopathogenic for cicadas, Part I, Taxonomy of the genus. Mycotaxon 1:13–40.

Soper RS, Delyzer AJ, Smith LF. 1976a. The genus *Massospora* entomopathogenic for cicadas, Part II Biology of *Massospora levispora* and its host *Okanagana rimosa*, with notes on *Massospora cicadina* on the periodical cicadas. Annals of the Entomological Society of America 69:89–95.

Soper RS, Smith LF, Delyzer AJ. 1976b. Epizootiology of *Massospora levispora* in an isolated population of *Okanagana rimosa*. Annals of the Entomological Society of America 69:275–83.

Soper RS. 1981. New cicada pathogens: *Massospora cicadettae* from Australia and *Massospora pahariae* from Afghanistan. Mycotaxon 13: 50–58.

Soper RS, Shimazu M, Humber RA, Ramos ME, Hajek AE. 1988. Isolation and characterization of *Entomophaga maimaiga* sp. nov., a fungal pathogen of gypsy moth, *Lymantria dispar,* from Japan. Journal of Invertebrate Pathology 51:2292–2241.

Spatafora JW, Chang Y, Benny GL, Lazarus K, Smith ME, Berbee ML, Bonito G, Corradi N, Grigoriev I, Gryganskyi A, James TY. 2016. A phylum-level phylogenetic classification of zygomycete fungi based on genome-scale data. Mycologia 108: 1028–1046.

Speare AT. 1921. *Massospora cicadina* Peck: a fungous parasite of the periodical cicada. Mycologia 13:72–82.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30:1312–1313.

Swofford DL. 2002. PAUP* Phylogenetic Analysis Using Parsimony (* and Other Methods). Sunderland, MA. Sinauer Associates, v. 4.0a build 166.

Tang AM, Jeewon R, Hyde KD. 2007. Phylogenetic utility of protein (RPB2, β-tubulin) and ribosomal (LSU, SSU) gene sequences in the systematics of Sordariomycetes (Ascomycota, Fungi). Antonie van Leeuwenhoek 91:327–349.

Taylor JW, Jacobson DJ, Kroken S, Kasuga T, Geiser DM, Hibbett DS, Fisher MC. 2000. Phylogenetic species recognition and species concepts in fungi. Fungal Genetics and Biology 31:21–32.

Thaxter R. 1888. The Entomophthoreae of the United States. Memoirs of the Boston Society of Natural History 4:190.

Tretter ED, Johnson EM, Benny GL, Lichtwardt RW, Wang Y, Kandel P, Novak SJ, Smith JF, White MM. 2014. An eight-gene molecular phylogeny of the Kickxellomycotina, including the first phylogenetic placement of Asellariales. Mycologia 106:912–935.

Vilgalys R, Hester M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. Journal of Bacteriology 172:4238–4246.

Warnes GR, Bolker B, Bonebakker L, Gentleman R, Huber W, Liaw A, Lumley T, Maechler M, Magnusson A, Moeller S, Schwartz M, Venables B. 2019. gplots: Various R Programming Tools for Plotting Data. R package version 3.0.1.1.

Wickham H. 2016. ggplot2: elegant graphics for data analysis. New York, NY: Springer-Verlag. 260 p.

Wickham H, François R, Henry L, Müller K. 2019. dplyr: a grammar of data manipulation. R package version 0.8.3.

Wronska AK, Bogus MI, Kaczmarek A, Kazek M. 2018. Harman and norharman, metabolites of entomopathogenic fungus *Conidiobolus coronatus* (Entomopthorales), disorganize development of *Galleria mellonella* (Lepidoptera) and affect serotonin-regulating enzymes. PLoS One 13:e0204828.

Figure 1: Photographs of cicada hosts (top), conidia (middle), and resting spores (bottom) of the *Massospora* species used in this study. Cicada hosts, from left to right: *Magicicada septendecim*, *Okanagana rimosa*, *Platypedia putnami*, *Diceroprocta semicincta*, *Tettigades* sp. Images are not to scale.

EFL+LSU+TUB

- 100/100/1.0 — *M. tettigatis* PL504.2
- *M. tettigatis* PL373
- 100/100/1.0
- 100/27/1.0
- *M. platypediae* NM4
- *M. platypediae* NM2
- *M. levispora* ML4
- *M. levispora* ML2
- *M. platypediae* CA3
- 44/---/---
- 100/---/1.0
- 100/100/1.0 — *M. diceroproctae* DB128
- *M. diceroproctae* DB176
- *M. diceroproctae* DB138
- 83/75/0.99
- *M. cicadina* GK2
- *M. cicadina* MC17
- 96/100/0.99 — *M. cicadina* MC7
- 81/78/0.80 — *M. cicadina* MICH231384
- *Arthrophaga myriapodina* ARSP1
- *Entomophthora muscae* ARSEF3074
- 100/27/1.0
- 69/---/0.71
- *Zoophthora radicans* Voucher45
- 100/100/1.0
- *Pandora neoaphidis* ARSEF7939
- 52/98/0.83 — *Pandora formicae* V1
- 22/56/0.53 — *Furia ithacensis* FiMD2

0.05 substitutions / site

EFL

- 97/100/1.0 — *M. tettigatis* PL504.2
- *M. tettigatis* PL373
- 100/46/1.0
- *M. platypediae* NM4
- *M. platypediae* NM2
- *M. platypediae* CA3
- 100/100/1.0
- *M. levispora* ML2
- *M. levispora* ML4
- 68/100/0.90
- 48/---/0.84
- *M. cicadina* MC7
- *M. cicadina* GK2
- 88/---/0.98
- *M. cicadina* MC17
- 37/---/---
- *M. diceroproctae* DB176
- *Entomophthora muscae* ARSEF3074
- 87/81/0.98
- 89/86/0.99 — *Pandora formicae* V1
- *Furia ithacensis* FiMD2
- 100/100/1.0
- *Pandora neoaphidis* ARSEF7939
- 100/46/1.0
- *Arthrophaga myriapodina* ARSP1

0.06 substitutions / site

Figure 2: Phylogenetic trees from the three-gene analysis (Concatenated, EFL, LSU, TUB). Topology and branch lengths shown are from the ML analysis. Bootstrap support and posterior probabilities are indicated for each node supported in the ML analysis (ML/MP/BI). Dashes indicate that the node did not appear in the indicated analysis. Arrow indicates the most recent common ancestor of the genus *Massospora.*

Figure 3: Boxplots of spore measurements used in this study. Letters indicate statistically significant differences among species. Top: Conidial length; middle: conidial width; bottom: resting spore diameter.

Supplemental Figure 1: SSU single gene tree of *Massospora* spp. and outgroups. Topology and

branch lengths shown are from the ML analysis. Bootstrap support and posterior probabilities are

indicated for each node supported in the ML analysis (ML/MP/BI). Dashes indicate that the

node did not appear in the indicated analysis

Table 1: Information about the currently accepted *Massospora* species, including host information and historic collection localities. Bold font indicates the species used in this study. *: Soper is not sure if these collections represent *M. diceroproctae* or a novel *Massospora* species, as they were found far from the type locality (Soper 1974).

Species	Authority	Reference	Known cicada hosts	Known localities
Massospora carinetae	R.S. Soper 1974	Soper 1974	*Carineta* sp.	Sao Paulo, Brazil; Misiones Province, Argentina
Massospora cicadettae	R.S. Soper 1981	Soper 1981	*Plerapsalta incipiens, Chelapsalta puer, Cicadetta* spp.	New South Wales, Queensland, Tasmania, Australia
Massospora cicadina	Peck 1878	Peck 1879	*Magicicada* spp. (all)	Eastern USA (entire range of the cicada genus)
Massospora diceroproctae	R.S. Soper 1974	Soper 1974	*Diceroprocta delicata, D. cinctifera, D. vitripennis*, D. biconica**	Texas, Louisiana*, Florida*, USA
Massospora diminuta	R.S. Soper 1974	Soper 1974	*Cicada* sp.	Amapa, Brazil
Massospora dorisianae	R.S. Soper 1974	Soper 1974	*Dorisiana semilata*	Paraiba, Brazil
Massospora fidicinae	R.S. Soper 1974	Soper 1974	*Fidicina* sp.	Guaimas District, Honduras; Chiapas, Mexico
Massospora levispora	R.S. Soper 1963	Soper 1963	*Okanagana rimosa, O. sperata*	California, USA; Ontario, Canada
Massospora ocypetes	R.S. Soper 1974	Soper 1974	*Guyalna bonaerensis*	Gualeguaychu, Argentina
Massospora pahariae	R.S. Soper 1981	Soper 1981	*Paharia casyapae*	Paghman District, Afghanistan
Massospora platypediae	R.S. Soper 1974	Soper 1974	*Platypedia putnami*	California, Utah, New Mexico, USA
Massospora spinosa	Cif., A.A. Machado & Vittal 1956	Ciferri *et al.* 1957	*Quesada gigas*	Paraiba, Brazil; Nuevo Leon, Mexico; Caracas, Venezuela
Massospora tettigatis	R.S. Soper 1974	Soper 1974	*Tettigades* spp.	Santiago, Aconcagua, Cautin Provinces, Chile

Table 2: Isolates used in phylogenetic analyses and associated metadata. Novel sequences are underlined.

Genus	Species	Strain	Source	Host	Host sex	Brood	Year	Location	Collector	Concat?	LSU	SSU	EFL	TUB
Massospora	*cicadina*	GK2	Reference & Novel	*Magicicada tredecassini*	F	23	2015	IN	G. Kritsky	Y	MH483015	MH483019	TBD	TBD
Massospora	*cicadina*	MC17	Novel	*Magicicada septendecim*	F	5	2016	WV	Kasson Lab	Y	TBD	TBD	TBD	TBD
Massospora	*cicadina*	MC7	Novel	*Magicicada cassini*	M	5	2016	WV	Kasson Lab	Y	TBD	TBD	TBD	TBD
Massospora	*cicadina*	MICH 231384	Genome	*Magicicada* sp.	-----	5	2016	OH	W. Davis	Y	TBD	TBD	-----	TBD
Massospora	*diceroproctae*	DB128	Novel	*Diceroprocta semicincta*	M	N/A	2016	AZ	D. Bublitz, P. Lukasik	Y	TBD	TBD	-----	TBD
Massospora	*diceroproctae*	DB138	Novel	*Diceroprocta semicincta*	M	N/A	2016	AZ	D. Bublitz, P. Lukasik	Y	TBD	-----	-----	TBD
Massospora	*diceroproctae*	DB176	Novel	*Diceroprocta semicincta*	F	N/A	2017	AZ	D. Bublitz, P. Lukasik	Y	TBD	TBD	TBD	TBD
Massospora	*levispora*	ML2	Novel	*Okanagana rimosa*	F	N/A	1998	MI	J. Cooley	Y	TBD	TBD	TBD	TBD
Massospora	*levispora*	ML4	Reference	*Okanagana rimosa*	F	N/A	1998	MI	J. Cooley	Y	MH483017	MH483021	MH483010	-----
Massospora	*platypediae*	CA3	Novel	*Platypedia putnami*	-----	N/A	2018	CA	M. Berger	Y	TBD	TBD	TBD	-----
Massospora	*platypediae*	NM2	Reference	*Platypedia putnami*	M	N/A	2017	NM	M. Berger	Y	MH483016	MH483020	MH483009	-----
Massospora	*platypediae*	NM4	Novel	*Platypedia putnami*	M	N/A	2017	NM	M. Berger	Y	TBD	TBD	TBD	TBD
Massospora	*tettigatis*	PL373	Novel	*Tettigades* sp.	M	N/A	2014	Chile	P. Lukasik, C. Veloso	Y	TBD	TBD	TBD	TBD
Massospora	*tettigatis*	PL504.2	Novel	*Tettigades* sp.	M	N/A	2014	Chile	P. Lukasik, C. Veloso	Y	TBD	TBD	TBD	TBD
Arthrophaga	*myriapodina*	Arsp1	Novel	*Xystocheir dissecta*	-----	N/A	2019	CA	S. Wolny	Y	TBD	TBD	TBD	-----
Entomophthora	*muscae*	ARSEF3074 and A	Reference	*Delia antiqua*	-----	N/A	1990	NY	R. Carruthers, S. Krasno	Y	NG_027647	AY635820	DQ275343	KF848865
Furia	*ithacensis*	FiMD2	Novel	*Rhagio mystaceus*	-----	N/A	2018	MD	Kasson Lab	Y	TBD	TBD	TBD	-----
Pandora	*formicae*	v1	Genome	*Formica polyctena*	-----	N/A	-----	Denmark	-----	Y	TBD	TBD	TBD	-----
Pandora	*neoaphidis*	ARSEF7939	Reference	*Microlophium carnosum*	-----	N/A	-----	-----	-----	Y	GU132790	EU267193	GU132771	-----
Zoophthora	*radicans*	Voucher 45	Reference	*Plutella xylostella*	-----	N/A	-----	Argentina	-----	Y	MG256490	MG252997	-----	-----