Among-individual diet variation within a lake trout ecotype: lack of stability of niche use.

Chavarie, L.[1,2,3], K.L. Howland[4,5], L.N., Harris[4], C.P. Gallagher[4], M.J. Hansen[6], W.M. Tonn[5],

A.M., Muir[7], C.C. Krueger[2].

[1]University of British Columbia, Biodiversity Center, 2212 Mall, Vancouver, Canada

[2]Department of Fisheries and Wildlife, Center for Systems Integration and Sustainability,

Michigan State University, 115 Manly Miles Building, East Lansing, Michigan, USA

[3] Scottish Centre for Ecology and the Natural Environment, IBAHCM, Rowardennan, Loch

Lomond, Glasgow G63 0AW, UK

[4] Fisheries and Oceans Canada, 501 University Crescent, Winnipeg, Canada

[5]Department of Biological Sciences, University of Alberta, CW 405 Biological building,

Edmonton, Canada

[6]U.S. Geological Survey, Hammond Bay Biological Station, 11188 Ray Road, Millersburg,

Michigan, USA (retired)

[7]Great Lakes Fishery Commission, 2100 Commonwealth Blvd., Suite 100, Ann Arbor,

Michigan, USA

[*]<u>Corresponding author:</u>

Louise Chavarie

Email:<u>chavarie@ualberta.ca</u>

Abstract:

In a polymorphic species, stable differences in resource use are expected among ecotypes, and homogeneity in resource use is predicted within an ecotype. Yet, using a broad resource spectrum has been identified as a strategy for fishes living in unproductive northern environments, where food is patchily distributed and ephemeral. We investigated whether individual specialization of trophic resources occurred within the generalist piscivore ecotype of lake trout from Great Bear Lake, Canada, reflective of a form of diversity. Four distinct dietary patterns of resource use within the lake trout ecotype were detected from fatty acid composition, with some variation linked to spatial patterns within Great Bear Lake. Feeding habits of different groups within the ecotype were not associated with detectable morphological or genetic differentiation, suggesting that behavioral plasticity caused the trophic differences. A low level of genetic differentiation was detected between exceptionally large-sized individuals and other individuals. Investigating a geologically young system that displays high levels of intraspecific diversity and focusing on individual variation in diet suggested that individual trophic specialization can occur within an ecotype. The characterization of niche use among individuals, as done in this study, is necessary to understand the role that individual variation can play at the beginning of differentiation processes.

Keywords: Resource use, Arctic, *Salvelinus namaycush*, generalist, specialist, fatty acids, genetic, morphology.

Introduction:

Phenotypic diversity within fish species that have colonized post-glacial lakes often represent early stages of species diversification (Snorrason et al., 2004). Many fishes that have colonized post-glacial freshwater systems are assumed to have been plastic generalists (i.e., flexible in use of habitat and food resources) at the time of colonization (Skúlason et al., 2019; Snorrason et al., 2004). Given the novel environment and new ecological opportunities, a newly established population may begin to display among-individual differences in behavior and other phenotypic characteristics (Svanbäck et al., 2007). Phenotypic plasticity, the capacity for one genotype to produce different phenotypes in response to environmental cues, could be a character subject to selection, facilitating the process of diversification (De Jong, 2005). Despite uncertainties of how phenotypic plasticity promotes divergence, plasticity appears to serve as an important element in early phases of diversification (Handelsman et al., 2013; Nonaka et al., 2015; Snorrason et al., 2004). Theory predicts that stable and predictable recently colonized systems would favor foraging and habitat specialization and increase the probability of eco-morphological diversification (Skúlason et al., 1999; Snorrason et al., 2004; Van Kleunen et al., 2005).

Phenotypic plasticity in temporally and spatially variable environments has been demonstrated repeatedly within and among populations (Skúlason et al., 2019). Whether niche expansion of a population is achieved by a general increase in niche widths for all individuals overall or by an increase of among-individual variation (i.e., expression of multiple individual specializations within a population) is a question in evolutionary ecology that remains unanswered (Bolnick et al., 2003; Roughgarden, 1972; Svanbäck et al., 2012). Several apparent generalist populations have been reported to be composed of combinations of specialized

individuals using several narrow niches that together yield an overall wide population niche

(Araújo et al., 2011; Araújo et al., 2008; Bolnick et al., 2003). Post-glacial lakes and co-

inhabiting species offer a wide range of characteristics that may favor or constrain individual

specialization. Post-glacial lakes are depauperate ecosystems with low interspecific competition,

which provides ecological opportunities that likely favor niche expansion (Bolnick et al., 2010;

Costa et al., 2008; Parent et al., 2014). Additionally, the large flexibility within post-glacial

colonizing species, with individuals having the potential to exploit a wide range of resources, can

facilitate the evolution of individual resource specialization and population divergence. Yet,

northern ecosystem food-webs are subjected to strong seasonal and episodic influences of

climate and the environment (McMeans et al., 2015). Accordingly, using a broad resource

spectrum has been identified as a useful strategy for fishes living in Arctic environments, where

food can be patchily distributed and ephemerally available. From all the facets of niche use that

are possible in northern lakes, understanding the magnitude and effect of individual

specialization in species and trophic positions is necessary to understand the role that variation

among individuals can play at the beginning of differentiation processes (Cloyed et al., 2016; De

León et al., 2012; Svanbäck et al., 2015).

Great Bear Lake (Northwest Territories, Canada), straddling the Arctic Circle, provides

an excellent opportunity to investigate the role of among-individual diet variation in

diversification processes in post-glacial lakes. Lake trout, *Salvelinus namaycush,* in this lake

show a high degree of intraspecific diversity within a geologically young system (8,000–10,000

yr BP; Johnson, 1975; Pielou, 2008). Specifically, extensive sympatric divergence has occurred

for this species with four ecotypes inhabiting the shallow-water (≤ 30 m) zone of Great Bear

Lake (Fig. A1; Chavarie et al., 2016a; Chavarie et al., 2015; Chavarie et al., 2013; Harris et al.,

2015). Three of these four shallow-water lake trout ecotypes are described as trophic generalists

with differing degrees of omnivory along a weak benthic-pelagic gradient (Chavarie et al.,

2016a; Chavarie et al., 2016b). Despite habitat and dietary overlap, significant differences in

morphological, genetic, and life-history variation have been reported (Chavarie et al., 2016 ;

Chavarie et al., 2013; Harris et al., 2015). The suggested resource use among the three ecotypes

could be caused by the combination of individual specialists along a resource continuum

(Chavarie et al., 2016b). In other words, although ecotype resource use may appear similar,

individuals within an ecotype may differ in their resource use. One of these three generalist

ecotypes (Ecotype 2; generalist with a tendency to consume more fish than other ecotypes,

referred to here as the piscivorous ecotype; Fig. 1) showed at least two different feeding

strategies, benthic cannibalism and interspecific piscivory in the pelagic zone (Chavarie et al.,

2016c).

To characterize niche use and individual variation within an ecotype in relation to

observed differentiation of feeding strategies, we focused this study solely on the piscivorous

lake trout ecotype. Fatty acid analysis assumes that dietary lipids are broken down into their

constituent fatty acids and incorporated relatively unchanged into consumer tissues (Howell et

al., 2003; Iverson, 2009; Iverson et al., 2004), allowing spatial and temporal diet comparison

among individuals (Duerksen et al., 2014; Eloranta et al., 2011; Hoffmann, 2017; Iverson, 2009;

Scharnweber et al., 2016). Fatty acids have been assessed to be a robust tool to characterize

lake trout diets (Happel et al., 2017; Happel et al., 2016; Iverson, 2009). Thus, fatty acids were

used as trophic bio-indicators to better understand dietary patterns of piscivorous lake trout and

investigate whether variation occurred among individuals in this ecotype and if individual

specialization may be contributing to the trophic breadth of the ecotype. Specifically, our aims

were to 1) compare resource use among lake trout individuals within Ecotype 2 (piscivores) by

characterizing their fatty acids profiles, 2) determine whether resource-use differences were

influenced by life-history traits (e.g., size and age), 3) characterize and compare morphological

variation among groups that expressed different feeding strategies, and 4) determine if genetic

differences existed among groups. In addition, we examined a sub-set of large lake trout of this

ecotype from our collections ($\geq$ 900 mm in fork length) referred to locally as "Giants" (Fig. 1), to

determine if they showed any ecological and genetic differences from others of this ecotype.

These exceptionally large individuals comprise $< 1\%$ of the lake trout population sampled in

Great Bear Lake, and are among the largest lake trout in the world (Chavarie et al., 2016).

Except for their large body-size, these individuals show no major morphological or spatial and

temporal distribution differences relative to other co-occurring piscivorous lake trout.

Methods

Study area and field sampling

Great Bear Lake is an oligotrophic Arctic freshwater system, 250 km south of the Arctic

Ocean, in Northwest Territories, Canada (N66° 06' W120° 35') (Johnson, 1975). As the world's

ninth largest and 19[th] deepest lake, the lake has a complex, multi-armed surface area of 31,790

km^2 and a maximum depth of 446 m (mean depth = 90 m). Great Bear Lake was formed by

scouring from the Laurentide ice-sheet during the Pleistocene and was originally part of glacial

Lake McConnell 8,000–10,000 yr BP (Johnson, 1975; Pielou, 2008). The lake has

characteristics typical of an arctic lake: ultra-oligotrophic, short ice-free season, and a simple

food web supporting only 15 fish species (Alfonso, 2004; Johnson, 1975; MacDonald et al.,

2004). Great Bear Lake lacks a commercial fishery but plays an important role in the local

economy, supporting a fly-in sport fishery for tourists and a subsistence fishery for the small

Sahtu community of Déline. Great Bear Lake has considerable intraspecific diversity within lake

trout, lake whitefish (*Coregonus clupeaformis*), and cisco (*C. artedi*) (Chavarie et al., 2013;

Howland et al., 2013).

Piscivorous lake trout were caught at depths ≤ 30 m using paired bottom sets (ca. 24 h) of

140-mm and multi-mesh (38–140 mm) stretched-mesh gill nets from late-July through August

over multiple years (2002–2011) among all five arms of the lake (Chavarie et al., 2016b;

Chavarie et al., 2015; Chavarie et al., 2013). During 2012-2014, multi-mesh gill nets (38 to 140

mm), with a typical soak time of 24 hours, were distributed across random depth-stratified sites

(0–150 m) among Keith, McVicar, and McTavish arms (Table A1). Compared to the other

ecotypes, piscivores have a streamlined body, large gape, and high growth rates throughout life,

similar to other piscivores (Chavarie et al., 2016 ; Chavarie et al., 2013). The piscivorous ecotype

also displayed a modest level of genetic differentiation from the three other ecotypes (Harris et

al., 2015).

We focused on adult trout due to the difficulty of classifying juveniles into ecotypes

(Chavarie et al., 2013; Zimmerman et al., 2006; Zimmerman et al., 2007) and to avoid the

confounding effects of ontogenetic shifts in morphology and diet. Of 79 fish analyzed herein, 35

piscivourous lake trout (Ecotype 2) were previously analyzed for fatty acids by Chavarie et al.

(2016b) and 44 fish were new additions to the diet analyses presented here. Fish were selected

from collections analyzed morphologically by Chavarie et al. (2015) to include a range of sizes

and ages within the piscivorous ecotype. For analyses invloving giant individuals, we selected

lake trout with fork lengths ≥ 900 mm.

A left lateral full-body digital image was taken for each lake trout caught according to the

procedures in Muir et al. (2012). Measurements, tissues, and structures were sampled to

determine biological characteristics related to life-history, including otoliths, fork length, round

weight, sex, and stage of maturity (i.e., immature, current year spawner, or resting) (Chavarie et

al., 2016 ; Chavarie et al., 2013). A dorsal muscle sample was collected and frozen at −20ºC for

fatty acid analysis (Budge et al., 2006; Kavanagh et al., 2010; Loseto et al., 2009) and tissue

from pectoral fins was collected and preserved in 95% ethanol for genetic analyses.

Fatty Acids

Analysis of 41 dietary fatty acids was carried out using procedures described by Chavarie

et al. (2016b) (Table 1). Muscle samples were freeze-dried and subsequently homogenized with

a mortar and pestle. Lipids were extracted overnight from 1 g of the homogenate in a 2:1

chloroform-methanol solution containing 0.01% BHT (v/v/w) at −20ºC (Folch et al., 1957).

After extraction, samples were filtered through Whatman Grade 1 Qualitative filter paper and the

filter paper/sample was rinsed twice with 2 ml of the 2:1 chloroform:methanol. Sample extract

was collected in a test tube and 7 ml of 0.88 N NaCl solution was added to encourage fatty acids

to move into the organic (chloroform) layer. The aqueous layer was discarded after which the

chloroform was dried with sodium sulfate prior to total lipid measurement. The extracted lipid

was used to prepare fatty acid methyl esters (FAME) by transesterification with Hilditch reagent

(0.5 N H_2SO_4 in methanol) (Morrison et al., 1964). Samples were heated for 1 h at 100 °C. Gas

chromatographic (GC) analysis was performed on an Agilent Technologies 7890N GC equipped

with a 30 m J&W DB-23 column (0.25 mm I.D; 0.15 μm film thickness). The GC was coupled

to a Flame Ionization Detector operating at 350 °C. Hydrogen was used as carrier gas flowing at

1.25 ml/min for 14 minutes, and increased to 2.5 ml/min for 5 min. The split/splitless injector

was heated to 260 °C and run in splitless mode. The oven program was as follows: 60 °C for 0.66

min, increasing by 22.82 °C/min to 165 °C with a 1.97 min hold; increasing by 4.56 °C/min to

174 ^{o}C and by 7.61 ^{o}C/min to 200 ^{o}C with a six min hold. Peak areas were quantified using

Agilent Technologies ChemStation software. Fatty acids standards were obtained from Supelco

(37 component FAME mix) and Nuchek (54 component mix GLC-463).

All fatty acid values were converted to a mass percentage of the total array, and were

named according the IUPAC nomenclature as X:Y n-z, where X is the number of carbon atoms

in the fatty acids, Y is the number of methylene-interrupted double bonds in the chain, and n-z

denotes the position of the last double bond relative to the methyl terminus (Ronconi et al.,

2010). Fatty acids suggested by Iverson et al. (2004) as important dietary fatty acids, which

transfer from prey to predator, were used in our analyses. Fatty acids profiles (% of fatty acids)

were transformed using arcsin square-root function. Fatty acids groups were identified using a

multivariate analysis R Package (Team, 2017), FactoMineR, using a hierarchical clustering

analysis based on principal components (Husson et al., 2012). To reduce the number of variables

used, A SIMPER (similarity percentage routine) was used to assess which fatty acids were

primarily responsible for observed differences among groups (King et al., 1999). A principal

components analysis (PCA) was performed on the fatty acid profiles with PC-ORD version 6

(McCune et al., 2011) among piscivorous groups to provide inferences about patterns of resource

use as defined by Chavarie et al. (2016b). Two-way Permutational Multivariate Analysis of

Variance (PERMANOVA), a non-parametric analog of Multivariate analysis of variance

(MANOVA), was used to test for differences in fatty acid composition among the groups

identified by FactoMineR and among arms (i.e., to investigate any spatial variations within the

piscivorous ecotype). Two-way PERMANOVA were performed in PAST 3 (Hammer et al.,

2001) using 9999 permutations. Pairwise *post-hoc* comparison (Bonferroni corrected) followed

to test differences among groups defined by FactoMineR and among arms. Pairwise *post-hoc*

comparison (Bonferroni corrected) also followed to test differences among arms (i.e., spatial

variation). Finally, the fatty acid groups determined by FactoMineR were tested for differences

in depth of capture using one-way analysis of similarities (ANOSIM) with 9999 permutations

using PAST 3.

Life-history

To determine if fatty acid groups differed in size-at-age, length vs. age was modeled

using the Von Bertalanffy length-age model fit to length at age-of-capture of individual fish

(Quinn et al., 1999):

$$L_t = L_\infty \left(1 - e^{-K(t-t_0)}\right)^\varepsilon$$

The length-age model describes length L_t at age-of-capture t as a function of theoretical

maximum length (L_∞ = mm), instantaneous rate at which L_t approaches L_∞ (K = 1/year),

theoretical age-at-zero length (t_0 = years), and multiplicative error (ε). Model parameters, L_∞, K,

and t_0, and associated standard errors were estimated using nonlinear regression. Residual sums-

of-squares were compared between a full model (separate models for each group) to a reduced

model (a single model for all groups) in a likelihood-ratio test (Hosmer Jr et al., 2000). If the

likelihood-ratio test was significant ($P \leq 0.05$), we concluded that growth differed among groups

identified by fatty acids. If the likelihood-ratio test was not significant ($P > 0.05$), we concluded

that growth did not differ among groups. The same test was repeated for each pair of groups,

with and without giant individuals (fork length ≥900 mm) included in each group, to isolate the

influence of this subset in our size-at-age comparison due to the prevalence of giants in one of

the groups (see Results).

 Genetic analyses

To determine if genetic differences existed among individuals expressing different

feeding strategies, the 79 lake trout classified by fatty acid composition into four groups were

genotyped to determine genetic variation and structure within and among groups. To allow a

sample size sufficient for making a genetic comparison of giants to the other dietary groups, 22

additional individuals determined non-randomly by their size ($\geq$ 900 mm ; giant sub-set) from

the 2002-2015 collections were added to giants processed for fatty acids, for a total of 39 giants.

Lake trout DNA was extracted from pectoral fin tissue preserved in ethanol using DNEasy

extraction kits (Qiagen Inc., Valencia, CA) following manufacturer protocols. Piscivorous

groups were assayed using a suite of 23 putatively neutral microsatellite markers amplified in

four multiplexes previously described in Harris et al. (2015). Amplified microsatellite fragments

were analyzed using an automated sequencer (ABI 3130xl Genetic Analyzer; Applied

Biosystems, Foster City, CA). The LIZ 600 size standard was incorporated for allele base-size

determination. All genotypes were scored using GeneMapper software ver. 4.0 (Applied

Biosystems) and then manually inspected to ensure accuracy.

The program MICROCHECKER ver. 2.2.0.3 (Van Oosterhout et al., 2004) was used to

identify genotyping errors, specifically null alleles and large allele dropout. Observed and

expected heterozygosity (H_E and H_O) were calculated using GENEPOP ver. 4.2 (Rousset, 2008).

The program HP-RARE ver. 1.1 (Kalinowski, 2005) was used to determine the number of

alleles, allelic richness, and private allelic richness for each group, sampling 22 genes in each

sample. Tests of departure from Hardy-Weinberg equilibrium and genotypic linkage

disequilibrium within each sample (i.e., for each fatty acid grouping and the Giant subset) were

conducted in GENEPOP using default values for both. Results from all tests were compared with

an adjusted alpha ($\alpha = 0.05$) following the False Discovery Rate procedure (Narum, 2006).

We used the POWSIM V. 4.1 analysis to assess the statistical power of our microsatellite

data set given the observed allelic frequencies within our samples in detecting significant genetic

differentiation between sampling groups (Ryman et al., 2006). For POWSIM analyses, we

assumed that lake trout within our study diverged from a common baseline population with the

same allelic frequencies as observed in our contemporary samples. Simulations were performed

with an effective population size of 5000 to yield values of F_{ST} of 0.01, 0.005 and 0.001. The

significance of tests in POWSIM were evaluated using Fisher's exact test and the $\chi2$ test and the

statistical power was determined as the proportion of simulations for which these tests showed a

significant deviation from zero. All simulations were performed with 1000 iterations.

Genetic structuring was tested among lake trout groups using several different methods.

First, genotypic differentiation among lake trout groups was calculated using log-likelihood (G)

based exact tests (Goudet et al., 1996) implemented in GENEPOP. Global F_{ST} (θ) (Weir et al.,

1984) was calculated in FSTAT ver. 2.9.3 (Goudet, 1995) and pairwise comparisons of F_{ST}

between groups were calculated in ARLEQUIN ver. 3.5 (Excoffier et al., 2005) using 10,000

permutations. We then employed the Bayesian clustering program STRUCTURE V. 2.3.2

(Pritchard et al., 2000) to resolve the putative number of populations (i.e., genetic clusters (K))

within our samples. Owing to the remarkably low levels of genetic differentiation among lake

trout in the Great Bear Lake (Harris et al., 2015; Harris et al., 2013), we employed the

LOCPRIOR algorithm (Hubisz et al., 2009). The LOCPRIOR algorithm considered the

location/sampling information as a prior in the model, which may perform better than the

traditional STRUCTURE model when the genetic structure is weak (Hubisz et al., 2009). We

also incorporated an admixture model with correlated allelic frequencies and the model was run

with a burn-in period of 500,000 iterations and 500,000 Markov chain Monte Carlo iterations.

We varied the potential number of populations (K) from 1 to 10 and we ran 20 iterations for each

value of K. The STUCTURE output was first processed in the program STRUCTURE

HARVESTER (Earl, 2012), followed by the combination of results of independent runs of the

program and compilation of results based on lnP(D) and the post hoc ΔK statistic of Evanno et

al. (2005), to infer the most likely number of clusters. The best alignment of replicate runs was

assessed with CLUMPP V. 1.1 (Jakobsson et al., 2007) and DISTRUCT V. 1.1 (Rosenberg,

2004) was then used to visualize the results. For STRUCTURE analyses, we reported both

lnP(D) and the post hoc ΔK statistic.

Finally, Discriminant Analysis of Principal Components (DAPC) (Jombart et al., 2010)

was implemented in the Adegenet package (Jombart, 2008) in R (Team, 2015). The number of

clusters was identified using the *find.clusters* function (a sequential K-means clustering

algorithm) and subsequent Bayesian Information Criterion (BIC), as suggested by Jombart et al.

(2010). Stratified cross-validation (carried out with the function *xvalDapc*) was used to

determine the optimal number of principal components to retain in the analysis.

Morphology

Morphological variation was quantified for the 79 lake trout and used to compare fatty

acid groupings (different feeding strategies) identified within the piscivorous ecotype. Twenty-

three landmarks and 20 semi-landmarks, based on Chavarie et al. (2015), and fourteen linear

measurements based on Muir et al. (2014), were used to characterize body and head shape from

digital images. The combination of traditional and geometric ecotype metrics was used because

relationships of phenotype morphology with foraging (e.g., jaw size) and swimming (e.g., fin

lengths and caudal peduncle depth) (Kahilainen et al., 2004; Kristjánsson et al., 2002; Webb,

1984). Landmarks and semi-landmarks were digitized in x and y coordinates using TPSDig2

software (http://life.bio.sunysb.edu/ecotype). Subsequently, digitized landmarks and semi-

landmarks were processed in a series of Integrated Morphometrics Programs (IMP) version 8

(http://www2.canisius.edu/;sheets/ecotypesoft), using partial warp scores, which are thin-plate

spline coefficients. Morphological methods and programs are described in Zelditch et al. (2012)

and specific procedures were described in further detail by Chavarie et al. (2013). All

morphological measurements were size-free, using centroid sizes or residuals from regressions

on standard length (Zelditch et al., 2012).

Canonical Variate Analyses (CVA) were conducted on all morphological data, including

body shape, head shape, and linear measurements, to determine relationships among fatty acid

groups. Body and head shape were analysed using CVAGen8 from the IMP software (Zelditch et

al., 2012) and for linear measurements, CVA was analyzed with SYSTAT (Systat Software Inc.,

Chicago, IL, USA). Single Factor Permutation MANOVA with 10 000 permutations tested for

differences among groups and determined the percentage of variation explained for a grouping if

a CVA was significant. For linear measurements, a Bonferroni-corrected post-hoc test followed

MANOVA to identify measurements that differed among group. Principal component analyses

(PCA) were performed on body- and head-shape data using PCAGen8 (IMP software) among

groups to visualize morphological variation within the dataset. PC-ORD version 6 software

(McCune et al., 2011) was used to perform a PCA on the linear measurements.

Results

Fatty acids

On the basis of fatty acid composition, piscivorous lake trout were divided along a

resource use axis into four groups (1-4; Fig. A2), containing 14, 16, 21, and 28 individuals,

respectively (Figs. 2 and A2; Table 1). Average dissimilarity was 14.61 (SIMPER analysis);

whereas, the most discriminating 26 fatty acids, explaining together ~89% of the separation

among groups, were: 22:6n-3 (12.5 %), 18:1n-9 (10.8 %), 16:1n-7 (6.8 %), 20:5n-3 (5.0 %),

20:4n-6 (3.9 %), 18:2n-6 (3.8 %), 22:4n-3 (3.7 %), 16:0 (3.5%), 20:4n-3 (3.3%), 18:1n7 (3.3%),

20:2n-6 (3.1%), 14:0 (2.8%), 20:1n-9 (2.7%), 22:5n-6 (2.7%), 20:3n-3 (2.3%), 22:2n-6 (2.1%),

18:0 (2.0%), 18:3n-3 (1.9%), 18:4n-3 (1.8%), 22:4n-6 (1.7%), 20:1n-7 (1.5%), 22:5n-3 (1.4%),

21:5n-3 (1.3%), 22:1n-11 (1.2%), 20:0 (1.2%), 16:4n-3 (1.2%), and 16:2n-4 (1.1%) (Table 1).

The first two axes of the fatty acids PCA explained 65.2 % of the variation in diet and the four

groups were supported by PERMANOVA ($F_{3,76}$ = 23.9, $P < 0.01$) and pairwise comparisons

between all pairs (all $P < 0.01$; Bonferroni corrected). Spatial differences in fatty acids

composition were found among arms ($F_{4,76}$ = 3.2, $P < 0.01$). Pairwise comparisons identified

differences between Smith and McVicar arms ($P = 0.02$; Bonferroni corrected; Fig. A3).

Interaction between fatty acids groups and arms was not significant ($p > 0.05$). Finally, depth of

capture did not differ among fatty acid groups ($p > 0.05$). For all groups, most lake trout were

caught between 0 and 20 m (Fig. A4).

Life-history

Overall, life history parameters did not differ among lake trout groups grouped by fatty

acid composition, including length-age models (Fig. 3; $F_{9, 63}$ = 1.58; $P = 0.141$). With the giant

sub-set included, growth differed between only between Group 3 and Group 4 ($F_{3, 41}$ = 3.958; P

= 0.014), but not between any other pairs ($P > 0.1$). Without Giants included (prevalence of

Giants was higher in Group 3 than Group 1, Group 2, and Group 4), none of the pairs differed for

length-at-age (P > 0.1), suggesting growth rate similarities among groups.

Genetic differentiation

Piscivorous lake trout groups displayed little genetic differentiation, except for the Giant

sub-set, which differed slightly from other groups that were defined by fatty acids.

MICROCHECKER identified two loci (OtsG253b and Sco102) that contained null alleles. These

loci, along with non-variable loci Sco218 and SSOSL456, were removed, leaving 19 informative

loci for subsequent analyses. Descriptive statistics of genetic variation were similar among

groups. The number of alleles per locus ranged from four (Smm21) to 41 (SnaMSU10) and

averaged 28.75 across all loci. Averaged observed heterozygosity ranged from 0.78 (Giant) to

0.83 (Group 1) while expected heterozygosity was 0.84 for all groups except Group 1 (0.85;

Table 2). Allelic richness ranged from 9.57 (Group 2 and 4) to 9.87 (Group 1), while expected

private allelic richness ranged from 0.87 (Group 3) to 1.08 (Group 2; Table 2). Only five of 95

tests (all of which involved different loci) showed significant departures from Hardy-Weinberg

equilibrium after adjustment for False Discovery Rate (adjusted $\alpha = 0.01$). Of those five, all were

heterozygote deficits and three involved the Giant sub-set. Only nine of 885 tests revealed

significant linkage disequilibrium after adjusting for False Discovery Rate (adjusted $\alpha = 0.0068$).

No locus-pair linkage disequilibrium combinations were consistently significant, but seven of

nine departures were in the Giant sub-set.

Using our microsatellite data set, the POWSIM analysis indicated a 100% power of

detecting F_{ST} values as low of 0.01 and 0.005. However, power was reduced to 77% when

assessing genetic differentiation at a F_{ST} of 0.001. Overall, our microsatellite data set (including

the number of loci, alleles per locus, and sample sizes) had sufficient power to detect relatively

low levels of genetic differentiation.

Global genetic differentiation was extremely low ($\theta = 0.001$, 95% c.i. $= -0.002$–0.005)

among the groups of piscivorous lake trout. Pairwise F_{ST} ranged from -0.004 to 0.016 (Table 3);

comparisons that included Giants always differed the most from the other fatty acid groups, and

they were involved in the only significant pairwise comparisons ($P < 0.05$, Table 3). The F_{ST}

values for the Giant vs. Groups 1 and 4 were generally similar to genetic differentiation among

the four original lake trout ecotypes in Great Bear Lake, except for Ecotype 1 vs Ecotype 2

(Table 3). Bayesian clustering implemented in STRUCTURE provided evidence for two genetic

clusters when evaluating both lnP(D) or ΔK (Table A2). The admixture plot based on K=2

showed no clear genetic structure between groups defined by fatty acid analysis; however, some

differentiation of the Giant sub-set from the fatty acid groups was observed (Fig. 4).

Finally, the Bayesian information criterion in the DAPC analysis (BIC = 185.42, Table

A3, Fig. A5 A) suggested that two clusters best explained genetic structure in our study (30 PCs

retained as suggested by the cross-validation procedure; Fig. A5 B). A compoplot (barplot

showing the probabilities of assignment of individuals to the different clusters) for K=2 revealed

no clear genetic structure between two groups identified by the DAPC analysis except for the

Giant group, which appeared to have more individuals assigned to cluster two (Fig. 4). Density

plots of the discriminant function, however, suggested that the two clusters identified through the

DAPC analysis are mostly non-overlapping (Fig. A5 C).

Morphology

Morphological variation was low among the four dietary groups within the piscivorous

ecotype. The first canonical axis for body shape CVA was significant ($P > 0.05$), but head shape

CVA revealed no significant canonical axes (P > 0.05) in groupings (Fig. 5 a, b, c). MANOVAs

for body and head shape were not significant (P > 0.05). Linear measurements CVA revealed

one significant canonical axis (P > 0.05). MANOVA permutation tests confirmed differences in

linear measurements among groups (P = 0.047). Most distinctions were related to linear

measurements of heads, with upper and lower jaws, head depth, and snout-eye lengths differing

between Group 3 and Group 4 (P ≤ 0.05), and head length differing between Group 1 and 4 (P =

0.03; Fig. 6). Caudal peduncle length and anal fin length differed marginally between Groups 2

vs 3 (P = 0.068) and Groups 1 vs 3 (P = 0.075), respectively. The first two PCA axes explained

44.3% and 12.3 % of variation for body shape, 35.1% and 30.7 % of variation for head shape,

and 39.6 % and 20.9 % for linear measurements (Fig. 5 d, e, f).

Discussion

A common assumption in polyphenism is that partitioning and variability of resource use will

occur predominantly among ecotypes rather than within ecotypes. In contrast, homogeneity of

resource use is anticipated to occur within ecotypes, be spatially and temporally stable, and

provide the selection opportunity for specialization (Amundsen et al., 2008; Knudsen et al.,

2010; Svanbäck et al., 2004). However, this study provided evidence that variation occurred

within an ecotype due to diet specialization among individuals, possibly a precursor to further

population diversification via fine scale ecological selection (Richardson et al., 2014;

Vonlanthen et al., 2009). The co-existence of multiple generalist ecotypes in Great Bear Lake

(Chavarie et al., 2016a), combined with the individual specialization shown here in the

piscivorous generalist ecotype, expands our understanding of niche use and expansion, plasticity,

individual specialization, and intraspecific diversity in evolutionarily young populations.

Using fatty acids as dietary biomarkers, four distinct patterns of resource use were identified

within the piscivorous lake trout of Great Bear Lake (Fig. 2). Groups 3 and 4 had the most

overlap and these groups were characterized by C20 and C22 monounsaturates, biomarkers of a

food web based on pelagic or deep-water copepods (Ahlgren et al., 2009; Happel et al., 2017;

Hoffmann, 2017; Loseto et al., 2009; Stowasser et al., 2006). Specifically, 20:1n-9 is associated

with calanoid copepods known to be particularly important in northern pelagic food webs

(Ahlgren et al., 2009; Budge et al., 2006; Kattner et al., 1998; Loseto et al., 2009). High levels of

14:0, 18:3n-3 and 18:4n-3 fatty acids within groups 3 and 4 are also associated with pelagic

environments (Scharnweber et al., 2016; Tucker et al., 2008), although high levels of 18:2n-6

and 18:3n-3 have also been associated with terrestrial markers (Budge et al., 2001; Budge et al.,

1998; Hoffmann, 2017).

Groups 1 and 2 were characterized by high concentrations of 16:4n-3, 20:4n-6 and 22:6n-3

found in diatom and dinoflagellate-based food webs, respectively. The fatty acid 20:4n-6 reflects

a benthic feeding strategy (from benthic invertebrates to fish) (Stowasser et al., 2006; Tucker et

al., 2008), whereas 22:6n-3 in pennate diatoms (Iverson, 2009) and filter feeders links planktonic

dinoflagellates to benthic filter-feeding bivalves in a food web (Alfaro et al., 2006; Virtue et al.,

2000). Relatively high concentrations of 16:0, 18:0 and 22:6n-3 and low concentrations of 16:1n-

7 supported the interpretation of carnivorous (or cannibalistic) dietary patterns (Dalsgaard et al.,

2003; Iverson, 2009; Iverson et al., 2004; Piché et al., 2010). Individuals positioned between

ends of principal components suggests a clinal pattern of resource use or habitat coupling

(Vonlanthen et al., 2009), where borders among groups are neither abrupt nor obvious as they are

part of a continuum (Hendry et al., 2009b). Overall, observed trophic patterns could reflect prey

associated with different microhabitat patches; however, the key assumption of disparity of prey

associated with habitat heterogeneity (Skulason et al., 1995; Svanbäck et al., 2005) may not be

applicable to Great Bear Lake (Chavarie et al., 2016a; Chavarie et al., 2020).

Sympatric divergence, in which barriers to gene flow are driven by selection between

ecological niches, has been implicated in the evolution of ecological and morphological variation

in fishes (Chavarie et al., 2016d; Hendry et al., 2007; Præbel et al., 2013). Despite the limited

ability of neutral microsatellite markers to detect patterns of functional divergence (Berg et al.,

2016; Lamichhaney et al., 2016; Roesti et al., 2015), the significant genetic differentiation based

on comparisons with Giant sub-set suggests some deviation from panmixis within the

piscivorous ecotype. Such a genetic pattern displayed by the Giant sub-set, despite a lack of

ecological discreteness, perhaps resulted from size-assortative mating and/or differences in

timing and location of spawning (Nagel et al., 1998; Rueger et al., 2016; Servedio et al., 2011).

Great Bear Lake is not the only lake in North America with an apparent divergence in lake trout

body size; in Lake Mistassini, "Giant" individuals also differed genetically from other lake trout

groups (Marin et al., 2016). The similarity based on lake trout body size between both lakes

suggests analogous variables favoring partial reproductive isolation. Although alternative causes

of genetic differentiation may be possible, due to the short time since the onset of divergence,

post-zygotic isolation seems unlikely in this system (e.g., prezygotic isolation generally evolves

more rapidly Coyne et al., 2004) and we therefore favor assortative mating based on size and

location as an explanation for the low-level genetic divergence observed. Nonetheless, putative

partial reproductive isolation within an ecotype adds to the complexity of diversification and

speciation processes potentially occurring within lake trout in Great Bear Lake (Hendry, 2009;

Nosil et al., 2009).

A central question arising from our analysis is what are the mechanisms behind these

patterns of variation? As individual specialization can result in dietary sub-groups and perhaps

differences in habitat use among sections of a population, such inter-individual variation within

ecological sub-groups could substantially influence processes of diversification (Araújo et al.,

2008; Cloyed et al., 2016). Among-individual resource specialization within an ecotype in a

species-poor ecosystem like Great Bear Lake could reflect the diversifying force of intraspecific

competition, lack of constraining effects of interspecific competition, the abundance and

distribution of resources (e.g., temporal and spatial variation of resources), or some combination

of these variables (Bolnick et al., 2007; Cloyed et al., 2016). Multiple patterns of resource

specialization within a single ecotype, as we see for lake trout in Great Bear Lake, contrasts with

the expected pattern of trophic divergence among ecotypes and homogenization in habitat use or

diet within an ecotype, a key assumption guiding the development of functional ecological

theory (Svanbäck et al., 2004; Violle et al., 2012). Expression of intraspecific divergence through

habitat and foraging specialization is thought to drive selection on traits that enable more

efficient use of resources (Schluter, 2000; Skulason et al., 1995; Snorrason et al., 2004).

In Great Bear Lake, multiple trophic generalists (which include the piscivores studied

herein) coexist with one specialist lake trout ecotype. This contrasts with the more commonly

reported observation of multiple specialist ecotypes (Chavarie et al., 2016a; Elmer, 2016;

Kassen, 2002). A generalist population, however, can be composed of several subsets of

specialized individuals (Bolnick et al., 2009; Bolnick et al., 2007; Bolnick et al., 2002). This

broad distribution of trophic variation within a population appears to be the case within the Great

Bear Lake piscivores. The among-individual specialization may result, to some degree, from

variable use of spatially separated resources and possibly temporally variable resources, both of

which could be expected in a large northern lake (Fig. A4; Costa et al., 2008; Cusa et al., 2019;

Quevedo et al., 2009). Ecologically, among-individual resource specialization within an ecotype

is another form of diversity (Araújo et al., 2008; Bolnick et al., 2003; Pires et al., 2011). Such

diversity may increase stability and persistence of an ecotype within a system where energy

resources are scarce and ephemeral (Cloyed et al., 2016; Pfennig et al., 2012; Smith et al., 2011).

Whether the level of among-individual specialization within this ecotype is stable or not is a

question that cannot be answered with our data.

Realized niche expansions are often linked to individuals of different morphologies and

body sizes, with evidence of efficiency trade-offs among different resources (Cloyed et al., 2016;

Parent et al., 2014; Roughgarden, 1972; Svanbäck et al., 2004). When a resource gradient exists,

niche expansion can be achieved via genetic differentiation, phenotypic plasticity, or a

combination of these processes (Bolnick et al., 2020; Parent et al., 2014). The apparent

segregation of resource use, based on our fatty acid analyses, despite a lack of major

morphological, body size, and genetic differentiation among the four dietary groups within the

piscivorous ecotype, suggests that behavioral plasticity is causing the observed patterns of

dietary differentiation. Plasticity may promote diversification by expanding the range of

phenotypes on which selection can act (Nonaka et al., 2015; Pfennig et al., 2010; West-Eberhard,

2003). Theoretical models suggest that exploiting a wide range of resources is either costly or

limited by constraints, but plasticity is favored when 1) spatial and temporal variation of

resources are important, 2) dispersal is high, 3) environmental cues are reliable, 4) genetic

variation for plasticity is high and 5) cost/limits of plasticity are low (Ackermann et al., 2004;

Hendry, 2016).

The expression of plasticity in response to ecological conditions (e.g., habitat structure,

prey diversity) can increase fitness. While most studies of diet variation focus on morphological

differences among ecotypes in a population, diet variation can also arise from behavioral,

biochemical, cognitive, and social-rank differences that cause functional ecology to be expressed

at a finer scale than at the ecotype level (McGill et al., 2006; Svanbäck et al., 2005; Violle et al.,

2012; Zhao et al., 2014). Indeed, behavioral plasticity likely has a temporal evolutionary

advantage due to relatively reduced reliance on ecologically beneficial morphological adaptation

(Smith et al., 2011; Svanbäck et al., 2009). The only detectable morphological differences among

piscivorous groups we identified in Great Bear Lake were associated with jaw lengths, snout-eye

distance, and head length and depth, which are strongly related to foraging opportunities (Adams

et al., 2002; Sušnik et al., 2006; Wainwright et al., 2016). Some morphological characters likely

express different degrees of plastic responses (adaptive or not), and thus may be expressed

differently depending on the magnitude and time of exposure to heterogeneous environments

(Hendry, 2016; Sharpe et al., 2008). For example, environmental components (e.g., habitat

structure) appear to have stronger and faster effects on linear characters (e.g., jaw length) than on

body shape (Chavarie et al., 2015; Sharpe et al., 2008). Trophic level might also limit the scope

for morphological variation in lake trout because piscivory can limit diversification of feeding

morphology in fishes (Collar et al., 2009; Svanbäck et al., 2015).

Conclusion

Understanding ecological mechanisms of diversification is challenging (Ackermann et

al., 2004). Divergence occurs along a continuum and in early stages, such as in post-glacial

lakes, morphological and dietary variation may not always be features that are related (Bolnick et

al., 2020; Bolnick et al., 2007). The debate around diversification sequence, (which diverges

first, behaviour, morphology, or ecology?) highlights the mosaic nature of intraspecific variation

(Hendry et al., 2009a). In this study, we asked whether among-individual diet variation could be

occurring within an ecotype by examining the fine-scale trophic variation of an early stage of

sympatric divergence of lake trout in Great Bear Lake (i.e., postglacial, representing ~350

generations; Harris et al., 2015). Due to presumed homogeneity, few studies have investigated

dietary patterns and groupings within an ecotype. Thus, this study provides evidence that among-

individual resource specialization can occur within an ecotype.

Rapid divergence within relatively few generations and among-individual diet variation

have both been demonstrated to be a strong driver of population dynamics (Ashley et al., 2003;

Bolnick et al., 2020; Fussmann et al., 2007; Turcotte et al., 2011). In this study, the fine-grained

trophic patterns shown within this ecotype suggested that ecological drivers (i.e., spatial

variation, habitat use, prey diversity, and abundance) could have important effects on plasticity

expression in early stages of divergence. Theory and experiments have demonstrated that

among-individual diet variation can increase stability within a system (Agashe, 2009). Using a

broad resource spectrum has been identified as an adaptive strategy for fishes living in Arctic

environments, where food availability is patchily distributed and ephemeral (Dill, 1983; Kassen,

2002; Smith et al., 2011). Thus, it is no surprise that the trophic individual specialization within

an ecotype was discovered within a northern lake.

List of abbreviations:

BP = before present

m = meter

mm = millimeter

h = hour

ca. = around

i.e., = stands for

e.g., = for example

NaCl = Sodium Chloride

FAME = fatty acid methyl esters

H_2SO_4 = Sulfuric acid

GC = Gas chromatographic

^{o}C = degree Celsius

^{o}C/min = degree Celsius/minutes

UPGMA = Unweighted Pair Group Method with Arithmetic Mean

PCA = principal component analysis

PERMANOVA = Permutational Multivariate Analysis of Variance

MANOVA = Multivariate analysis of variance

SIMPER= similarity percentage routine

ANOSIM = analysis of similarities

N =Number of individuals genotyped

N_A = number of alleles

H_E=expected heterozygosity

H_O = observed heterozygosity

A_R= allelic richness

PA_R = private allelic richness

A = alpha

FCA = Factorial correspondence analysis

k=number of alleles

DAPC = Discriminant Analysis of Principal Components

IMP = Integrated Ecotypeometrics Programs

CVA = Canonical Variate Analyses

Acknowledgements

We thank one anonymous reviewer and Alexandra Tyers for their insightful readings and
constructive suggestions. Déline Renewable Resources Council, Déline Lands and Finance
Corporation, the community of Déline, DFO in Hay River, and the Department of Environment
and Natural Resources in Déline provided valuable help with field planning and logistics. We
especially thank J. Chavarie, S. Buckley, L. Harris, G. Lafferty, M. Lindsey, M. Low, Z. Martin,
S. Wiley, and C. Yukon, who helped lead sampling teams and coordinate logistics.   The
following individuals who helped conduct field sampling in various years are gratefully
acknowledeged: J. Baptiste, D. Betsidea, D. Baton, L. Dueck, R. Eshenroder, G. Menacho, N.
Modeste, I. Neyelle, L. Neyelle, M. Smirle, A. Swietzer, C. Takazo, A. Vital, F. Vital, B. Yukon,
M. Yukon, T. Yukon and Charity, Cameron, and Cyre Yukon. D. Phillibert, K. Laubriat, S.
Thacker, and K. Theoret conducted isotopic lab work.

Declarations:

Authors' contributions

LC, KH, WT, CK, and AM conceived and funded the study. LC and CG carried out the field

work. LC, CG, LH, and MH participated in the data analyses. LC wrote the manuscript. All

authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article. Raw data

are available from the corresponding author upon reasonable request.

Consent to publish

Not applicable.

Ethics approval and consent to participate

We declare that our experiments were performed in the respect of ethical rules. This protocol

was approved by Department of Fisheries and Ocean Canada, Freshwater Institute Animal Care

Committee Science Laboratories.

Funding

Funds are from Fisheries and Oceans Canada (DFO), Northern Development Canada Northwest

Territories Cumulative Impacts Monitoring Program grants, Polar Continental Shelf Program,

Sahtu Renewable Resource Board, and the Great Lakes Fishery Commission. Funding bodies

had no role in the study design, in the collection, analysis and interpretation of data, in writing

the manuscript or the decision to submit the paper for publication.

Literature:

Ackermann, M., Doebeli, M., & Gomulkiewicz, R. (2004). Evolution of niche width and adaptive diversification. *Evolution, 58*(12), 2599-2612.

Adams, C. E., & Huntingford, F. A. (2002). The functional significance of inherited differences in feeding morphology in a sympatric polymorphic population of Arctic charr. *Evolutionary Ecology, 16*(1), 15-25.

Agashe, D. A. (2009). Consequences of intraspecific genetic variation for population dynamics and niche expansion.

Ahlgren, G., Vrede, T., & Goedkoop, W. (2009). Fatty Acid Ratios in Freshwater Fish, Zooplankton and Zoobenthos–Are There Specific Optima? In *Lipids in aquatic ecosystems* (pp. 147-178): Springer.

Alfaro, A. C., Thomas, F., Sergent, L., & Duxbury, M. (2006). Identification of trophic interactions within an estuarine food web (northern New Zealand) using fatty acid biomarkers and stable isotopes. *Estuarine, Coastal and Shelf Science, 70*(1), 271-286.

Alfonso, N. R. (2004). Evidence for two morphotypes of lake charr, *Salvelinus namaycush*, from Great Bear Lake, Northwest Territories, Canada *Environmental Biology of Fishes, 71*, 21-32.

Amundsen, P.-A., Knudsen, R., & Klemetsen, A. (2008). Seasonal and ontogenetic variations in resource use by two sympatric Arctic charr morphs. *Environmental Biology of Fishes, 83*(1), 45-55. doi:10.1007/s10641-007-9262-1

Araújo, M. S., Bolnick, D. I., & Layman, C. A. (2011). The ecological causes of individual specialisation. *Ecology Letters, 14*(9), 948-958. doi:10.1111/j.1461-0248.2011.01662.x

Araújo, M. S., Guimarães, P. R., Svanbäck, R., Pinheiro, A., Guimarães, P., Reis, S. F. d., & Bolnick, D. I. (2008). Network analysis reveals contrasting effects of intraspecific competition on individual vs. population diets. *Ecology, 89*(7), 1981-1993.

Ashley, M. V., Willson, M. F., Pergams, O. R., O'Dowd, D. J., Gende, S. M., & Brown, J. S. (2003). Evolutionarily enlightened management. *Biological Conservation, 111*(2), 115-123.

Berg, P. R., Star, B., Pampoulie, C., Sodeland, M., Barth, J. M. I., Knutsen, H., . . . Jentoft, S. (2016). Three chromosomal rearrangements promote genomic divergence between migratory and stationary ecotypes of Atlantic cod. *Scientific Reports, 6*, 23246. doi:10.1038/srep23246

Bolnick, D. I., & Ballare, K. M. (2020). Resource diversity promotes among-individual diet variation, but not genomic diversity, in lake stickleback. *Ecology Letters, 23*(3), 495-505. doi:10.1111/ele.13448

Bolnick, D. I., Ingram, T., Stutz, W. E., Snowberg, L. K., Lau, O. L., & Paull, J. S. (2010). Ecological release from interspecific competition leads to decoupled changes in population and individual niche width. *Proceedings of the Royal Society B: Biological Sciences, 277*(1689), 1789-1797. doi:10.1098/rspb.2010.0018

Bolnick, D. I., & Paull, J. S. (2009). Morphological and dietary differences between individuals are weakly but positively correlated within a population of threespine stickleback. *Evolutionary Ecology Research, 11*(8), 1217-1233.

Bolnick, D. I., Svanbäck, R., Araújo, M. S., & Persson, L. (2007). Comparative support for the niche variation hypothesis that more generalized populations also are more heterogeneous. *Proceedings of the National Academy of Sciences, 104*(24), 10075-10079.

Bolnick, D. I., Svanbäck, R., Fordyce, J. A., Yang, L. H., Davis, J. M., Hulsey, C. D., & Forister,
M. L. (2003). The ecology of individuals: incidence and implications of individual
specialization. *American Naturalist, 161*(1), 1-28.
Bolnick, D. I., Yang, L. H., Fordyce, J. A., Davis, J. M., & Svanbäck, R. (2002). Measuring
individual-level resource specialization. *Ecology, 83*(10), 2936-2941. doi:10.1890/0012-
9658(2002)083[2936:MILRS]2.0.CO;2
Budge, S., Parrish, C., & Mckenzie, C. (2001). Fatty acid composition of phytoplankton, settling
particulate matter and sediments at a sheltered bivalve aquaculture site. *Marine
Chemistry, 76*(4), 285-303.
Budge, S. M., Iverson, S. J., & Koopman, H. N. (2006). Studying trophic ecology in marine
ecosystems using fatty acids: a primer on analysis and interpretation. *Marine Mammal
Science, 22*(4), 759-801.
Budge, S. M., & Parrish, C. C. (1998). Lipid biogeochemistry of plankton, settling matter and
sediments in Trinity Bay, Newfoundland. II. Fatty acids. *Organic Geochemistry, 29*(5),
1547-1559.
Chavarie, L., Harford, W. J., Howland, K. L., Fitzsimons, J., Muir, A. M., Krueger, C. C., &
Tonn, W. M. (2016a). Multiple generalist morphs of Lake Trout: Avoiding constraints on
the evolution of intraspecific divergence? *Ecology and Evolution, 6*, 7727–7741.
Chavarie, L., Hoffmann, J., Muir, A. M., Krueger, C. C., Bronte, C. R., Howland, K. L., . . .
Swanson, H. K. (2020). Dietary versus nondietary fatty acid profiles of lake trout
ecotypes from Lake Superior and Great Bear Lake: Are fish really what they eat?
Canadian Journal of Fisheries and Aquatic Sciences, 1-12. doi:10.1139/cjfas-2019-0343
Chavarie, L., Howland, K., Gallagher, C., & Tonn, W. (2016b). Fatty acid signatures and
stomach contents of four sympatric Lake Trout: assessment of trophic patterns among
morphotypes in Great Bear Lake. *Ecology of Freshwater Fish, 25*, 109-124.
doi:10.1111/eff.12195
Chavarie, L., Howland, K., Gallagher, C., & Tonn, W. (2016c). Fatty acid signatures and
stomach contents of four sympatric Lake Trout: assessment of trophic patterns among
morphotypes in Great Bear Lake. *Ecology of Freshwater Fish, 25*(1), 109-124.
doi:10.1111/eff.12195
Chavarie, L., Howland, K., Harris, L., & Tonn, W. (2015). Polymorphism in lake trout in Great
Bear Lake: intra□lake morphological diversification at two spatial scales. *Biological
Journal of the Linnean Society, 114*(1), 109-125.
Chavarie, L., Howland, K., Venturelli, P., Kissinger, B. C., Tallman, R., & Tonn, W. (2016).
Life-history variation among four shallow-water morphotypes of lake trout from Great
Bear Lake, Canada. *Journal of Great Lakes Research*,
DOI:10.1016/j.jglr.2015.1007.1006. doi:doi:10.1016/j.jglr.2015.07.006
Chavarie, L., Howland, K. L., & Tonn, W. M. (2013). Sympatric polymorphism in lake trout: the
coexistence of multiple shallow-water morphotypes in Great Bear Lake. *Transactions of
the American Fisheries Society, 142*(3), 814-823. doi:10.1080/00028487.2013.763855
Chavarie, L., Muir, A. M., Zimmerman, M. S., Baillie, S. M., Hansen, M. J., Nate, N. A., . . .
Krueger, C. C. (2016d). Challenge to the model of lake charr evolution: shallow- and
deep-water morphs exist within a small postglacial lake. *Biological Journal of the
Linnean Society*, n/a-n/a. doi:10.1111/bij.12913

Cloyed, C. S., & Eason, P. K. (2016). Different ecological conditions support individual
specialization in closely related, ecologically similar species. *Evolutionary Ecology,*
30(3), 379-400.
Collar, D. C., O'Meara, B. C., Wainwright, P. C., & Near, T. J. (2009). Piscivory Limits
Diversification of Feeding Morphology in Centrarchid Fishes. *Evolution, 63*(6), 1557-
1573.
Costa, G., xa, C, Mesquita, D., xa, O, . . . Editor: Monica, G. (2008). Niche Expansion and the
Niche Variation Hypothesis: Does the Degree of Individual Variation Increase in
Depauperate Assemblages? *The American Naturalist, 172*(6), 868-877.
doi:10.1086/592998
Coyne, J. A., & Orr, H. A. (2004). *Speciation* (Vol. 37): Sinauer Associates Sunderland, MA.
Cusa, M., Berge, J., & Varpe, Ø. (2019). Seasonal shifts in feeding patterns: Individual and
population realized specialization in a high Arctic fish. *Ecology and Evolution, 0*(0).
doi:10.1002/ece3.5615
Dalsgaard, J., St. John, M., Kattner, G., Müller-Navarra, D., & Hagen, W. (2003). Fatty acid
trophic markers in the pelagic marine environment. In *Advances in Marine Biology* (Vol.
46, pp. 225-340): Academic Press.
De Jong, G. (2005). Evolution of phenotypic plasticity: patterns of plasticity and the emergence
of ecotypes. *New Phytologist, 166*(1), 101-118.
De León, L. F., Rolshausen, G., Bermingham, E., Podos, J., & Hendry, A. P. (2012). Individual
specialization and the seeds of adaptive radiation in Darwin's finches. *Evolutionary*
Ecology Research, 14(4), 365-380.
Dill, L. M. (1983). Adaptive flexibility in the foraging behavior of fishes. *Canadian Journal of*
Fisheries and Aquatic Sciences, 40(4), 398-408.
Duerksen, S. W., Thiemann, G. W., Budge, S. M., Poulin, M., Niemi, A., & Michel, C. (2014).
Large, omega-3 rich, pelagic diatoms under Arctic sea ice: sources and implications for
food webs. *PloS one, 9*(12), e114070.
Earl, D. A. (2012). STRUCTURE HARVESTER: a website and program for visualizing
STRUCTURE output and implementing the Evanno method. *Conservation genetics*
resources, 4(2), 359-361.
Elmer, K. R. (2016). Genomic tools for new insights to variation, adaptation, and evolution in
the salmonid fishes: a perspective for charr. *Hydrobiologia*, DOI:10.1007/s10750-10015-
12614-10755. doi:10.1007/s10750-015-2614-5
Eloranta, A. P., Siwertsson, A., Knudsen, R., & Amundsen, P. A. (2011). Dietary plasticity of
Arctic charr (*Salvelinus alpinus*) facilitates coexistence with competitively superior
European whitefish (*Coregonus lavaretus*). *Ecology of Freshwater Fish, 20*(4), 558-568.
Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals
using the software STRUCTURE: a simulation study. *Molecular Ecology, 14*(8), 2611-
2620.
Excoffier, L., Laval, G., & Schneider, S. (2005). Arlequin (version 3.0): an integrated software
package for population genetics data analysis. *Evolutionary bioinformatics, 1*, 47-50.
Folch, J., Lees, M., & Sloane-Stanley, G. (1957). A simple method for the isolation and
purification of total lipids from animal tissues. *Journal of Biological Chemistry, 226*(1),
497-509.

Fussmann, G. F., Loreau, M., & Abrams, P. A. (2007). Eco-evolutionary dynamics of
communities and ecosystems. *Functional Ecology, 21*(3), 465-477. doi:10.1111/j.1365-
2435.2007.01275.x
Goudet, J. (1995). FSTAT (version 1.2): a computer program to calculate F-statistics. *Journal of*
Heredity, 86(6), 485-486.
Goudet, J., Raymond, M., de Meeüs, T., & Rousset, F. (1996). Testing Differentiation in Diploid
Populations. *Genetics, 144*(4), 1933.
Hammer, Ø., Harper, D., & Ryan, P. (2001). Past: paleontological statistics software package for
education and data analysis. *Palaeontologia Electronica, 4*, 1-9.
Handelsman, C. A., Broder, E. D., Dalton, C. M., Ruell, E. W., Myrick, C. A., Reznick, D. N., &
Ghalambor, C. K. (2013). Predator-induced phenotypic plasticity in metabolism and rate
of growth: rapid adaptation to a novel environment. *Integrative and Comparative*
Biology, 53(6), 975-988.
Happel, A., Jonas, J. L., McKenna, P., Rinchard, J., He, J. X., & Czesny, S. J. (2017). Spatial
Variability of Lake Trout Diets in Lakes Huron and Michigan Revealed by Stomach
Content and Fatty Acid Profiles. *Canadian Journal of Fisheries and Aquatic Sciences*.
doi:10.1139/cjfas-2016-0202
Happel, A., Stratton, L., Pattridge, R., Rinchard, J., & Czesny, S. (2016). Fatty-acid profiles of
juvenile lake trout reflect experimental diets consisting of natural prey. *Freshwater*
Biology, 61(9), 1466-1476. doi:10.1111/fwb.12786
Harris, L. N., Chavarie, L., Bajno, R., Howland, K. L., Wiley, S. H., Tonn, W. M., & Taylor, E.
B. (2015). Evolution and origin of sympatric shallow-water morphotypes of Lake Trout,
Salvelinus namaycush, in Canada's Great Bear Lake. *Heredity, 114*(1), 94-106.
doi:10.1038/hdy.2014.74
Harris, L. N., Howland, K. L., Kowalchuk, M. W., Bajno, R., Lindsay, M. M., & Taylor, E. B.
(2013). Microsatellite and mtDNA analysis of lake trout, Salvelinus namaycush, from
Great Bear Lake, Northwest Territories: impacts of historical and contemporary
evolutionary forces on Arctic ecosystems. *Ecology and Evolution, 3*(1), 145-161.
doi:10.1002/ece3.439
Hendry, A., Bolnick, D., Berner, D., & Peichel, C. (2009a). Along the speciation continuum in
sticklebacks. *Journal of Fish Biology, 75*(8), 2000-2036.
Hendry, A. P. (2009). Ecological speciation! Or the lack thereof?This Perspective is based on the
author's J.C. Stevenson Memorial Lecture delivered at the Canadian Conference for
Fisheries Research in Halifax, Nova Scotia, January 2008. *Canadian Journal of Fisheries*
and Aquatic Sciences, 66(8), 1383-1398. doi:10.1139/F09-074
Hendry, A. P. (2016). Key questions on the role of phenotypic plasticity in eco-evolutionary
dynamics. *Journal of Heredity, 107*, 25-41.
Hendry, A. P., Bolnick, D. I., Berner, D., & Peichel, C. L. (2009b). Along the speciation
continuum in stickleback. *Journal of Fish Biology, 75*. doi:10.1111/j.1095-
8649.2009.02419.x
Hendry, A. P., Nosil, P., & Rieseberg, L. H. (2007). The speed of ecological speciation.
Functional Ecology, 21(3), 455-464.
Hoffmann, J. M. (2017). *Investigating trophic ecology and dietary niche overlap among morphs*
of Lake Trout in Lake Superior. (MS thesis), University of Waterloo,
Hosmer Jr, D. W., Lemeshow, S., & Sturdivant, R. X. (2000). Model building strategies and
methods for logistic regression. *Applied Logistic Regression, Third Edition*, 89-151.

Howell, K. L., Pond, D. W., Billett, D. S., & Tyler, P. A. (2003). Feeding ecology of deep-sea
798   seastars (Echinodermata: Asteroidea): a fatty-acid biomarker approach. *Marine Ecology*
799   *Progress Series, 255*, 193-206.
800   Howland, K. L., Gallagher, C., Boguski, D., Chavarie, L., Reist, J., Rosenburg, B., & Wiley, S.
801   (2013). *Variation in morphology, life history and ecology of cisco in Great Bear Lake,*
802   *Northwest Territories, Canada*. (2013/106). Sci. Advis. Sec. Res. Doc. .
803   Hubisz, M. J., Falush, D., Stephens, M., & Pritchard, J. K. (2009). Inferring weak population
804   structure with the assistance of sample group information. *Molecular Ecology Resources,*
805   *9*(5), 1322-1332.
806   Husson, F., Josse, J., Le, S., & Mazet, J. (2012). FactoMineR: Multivariate Exploratory Data
807   Analysis and Data Mining with R. . *R package version 1.18*.
808   Iverson, S. J. (2009). Tracing aquatic food webs using fatty acids: from qualitative indicators to
809   quantitative determination. In *Lipids in Aquatic Ecosystems* (pp. 281-308): Springer.
810   Iverson, S. J., Field, C., Bowen, W. D., & Blanchard, W. (2004). Quantitative fatty acid
811   signature analysis: a new method of estimating predator diets. *EcologicalMonographs,*
812   *74*, 211–235.
813   Jakobsson, M., & Rosenberg, N. A. (2007). CLUMPP: a cluster matching and permutation
814   program for dealing with label switching and multimodality in analysis of population
815   structure. *Bioinformatics, 23*(14), 1801-1806.
816   Johnson, L. (1975). Distribution of fish species in Great Bear Lake, Northwest Territories, with
817   reference to zooplankton, benthic invertebrates, and environmental conditions. *Journal of*
818   *the Fisheries Research Board of Canada, 32*, 1989-2004.
819   Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers.
820   *Bioinformatics, 24*(11), 1403-1405.
821   Jombart, T., Devillard, S., & Balloux, F. (2010). Discriminant analysis of principal components:
822   a new method for the analysis of genetically structured populations. *BMC Genetics,*
823   *11*(1), 94.
824   Kahilainen, K., Malinen, T., Tuomaala, A., & Lehtonen, H. (2004). Diel and seasonal habitat and
825   food segregation of three sympatric *Coregonus lavaretus* forms in a subarctic lake.
826   *Journal of Fish Biology, 64*(2), 418-434.
827   Kalinowski, S. T. (2005). hp☐rare 1.0: a computer program for performing rarefaction on
828   measures of allelic richness. *Molecular Ecology Notes, 5*(1), 187-189.
829   Kassen, R. (2002). The experimental evolution of specialists, generalists, and the maintenance of
830   diversity. *Journal of Evolutionary Biology, 15*(2), 173-190. doi:10.1046/j.1420-
831   9101.2002.00377.x
832   Kattner, G., Hagen, W., Graeve, M., & Albers, C. (1998). Exceptional lipids and fatty acids in
833   the pteropod *Clione limacina* (Gastropoda) from both polar oceans. *Marine Chemistry,*
834   *61*(3), 219-228.
835   Kavanagh, A. S., Cronin, M. A., Walton, M., & Rogan, E. (2010). Diet of the harbour seal
836   (*Phoca vitulina vitulina*) in the west and south-west of Ireland. *Journal of the Marine*
837   *Biological Association of the United Kingdom, 90*(08), 1517-1527.
838   King, J. R., & Jackson, D. A. (1999). Variable selection in large environmental data sets using
839   principal components analysis. *Environmetrics, 10*(1), 67-77. doi:10.1002/(SICI)1099-
840   095X(199901/02)10:1<67::AID-ENV336>3.0.CO;2-0

Knudsen, R., Primicerio, R., Amundsen, P.-A., & Klemetsen, A. (2010). Temporal stability of
individual feeding specialization may promote speciation. *Journal of Animal Ecology,*
79(1), 161-168. doi:10.1111/j.1365-2656.2009.01625.x
Kristjánsson, B. K., Skúlason, S., & Noakes, D. L. G. (2002). Morphological segregation of
Icelandic threespine stickleback (*Gasterosteus aculeatus* L). *Biological Journal of the*
Linnean Society, 76. doi:10.1046/j.1095-8312.2002.00063.x
Lamichhaney, S., Fan, G., Widemo, F., Gunnarsson, U., Thalmann, D. S., Hoeppner, M. P., . . .
Andersson, L. (2016). Structural genomic changes underlie alternative reproductive
strategies in the ruff (*Philomachus pugnax*). *Nature Genetics, 48*(1), 84-88.
doi:10.1038/ng.3430
http://www.nature.com/ng/journal/v48/n1/abs/ng.3430.html#supplementary-information
Loseto, L., Stern, G., Connelly, T., Deibel, D., Gemmill, B., Prokopowicz, A., . . . Ferguson, S.
(2009). Summer diet of beluga whales inferred by fatty acid analysis of the eastern
Beaufort Sea food web. *Journal of Experimental Marine Biology and Ecology, 374*(1),
12-18.
MacDonald, D., Levy, D., Czarnecki, A., Low, G., & Richea, N. (2004). State of the aquatic
knowledge of Great Bear Lake watershed. Report to Indian and Northern Affairs Canada.
Water Resources Division, MacDonald Environmental Sciences, Nanaimo, British
Columbia.
Marin, K., Coon, A., Carson, R., Debes, P. V., & Fraser, D. J. (2016). Striking Phenotypic
Variation yet Low Genetic Differentiation in Sympatric Lake Trout (*Salvelinus*
namaycush). *PloS one, 11*(9), e0162325. doi:10.1371/journal.pone.0162325
McCune, B., & Mefford, M. (2011). PC-ORD v. 6.255 beta. *MjM Software. Gleneden Beach,*
Lincoln.
McGill, B. J., Enquist, B. J., Weiher, E., & Westoby, M. (2006). Rebuilding community ecology
from functional traits. *Trends in Ecology & Evolution, 21*(4), 178-185.
doi:http://dx.doi.org/10.1016/j.tree.2006.02.002
McMeans, B. C., McCann, K. S., Humphries, M., Rooney, N., & Fisk, A. T. (2015). Food Web
Structure in Temporally-Forced Ecosystems. *Trends in Ecology & Evolution, 30*(11),
662-672. doi:https://doi.org/10.1016/j.tree.2015.09.001
Morrison, W. R., & Smith, L. M. (1964). Preparation of fatty acid methyl esters and
dimethylacetals from lipids with boron fluoride–methanol. *Journal of Lipid Research,*
5(4), 600-608.
Muir, A. M., Bronte, C. R., Zimmerman, M. S., Quinlan, H. R., Glase, J. D., & Krueger, C. C.
(2014). Ecomorphological diversity of Lake Charr *Salvelinus namaycush* at Isle Royale,
Lake Superior. *Transactions of the American Fisheries Society, 143*(4), 972-987.
Muir, A. M., Vecsei, P., & Krueger, C. C. (2012). A perspective on perspectives: a method
toward reducing variation in digital shape analysis. *Transactions of the American*
Fisheries Society, 141(4), 1161-1170. doi:10.1080/00028487.2012.685823
Nagel, L., & Schluter, D. (1998). Body size, natural selection, and speciation in sticklebacks.
Evolution, 209-218.
Narum, S. R. (2006). Beyond Bonferroni: less conservative analyses for conservation genetics.
Conservation Genetics, 7(5), 783-787.
Nonaka, E., Svanbäck, R., Thibert-Plante, X., Englund, G., & Brännström, Å. (2015).
Mechanisms by Which Phenotypic Plasticity Affects Adaptive Divergence and
Ecological Speciation. *Am. Nat., 186*(5), 126–143.

Nosil, P., Harmon, L. J., & Seehausen, O. (2009). Ecological explanations for (incomplete)
speciation. *Trends in Ecology & Evolution, 24*(3), 145-156.
Parent, C. E., Agashe, D., & Bolnick, D. I. (2014). Intraspecific competition reduces niche width
in experimental populations. *Ecology and Evolution, 4*(20), 3978-3990.
Pfennig, D. W., & Pfennig, K. S. (2012). *Evolution's wedge: competition and the origins of*
diversity: Univ of California Press.
Pfennig, D. W., Wund, M. A., Snell-Rood, E. C., Cruickshank, T., Schlichting, C. D., &
Moczek, A. P. (2010). Phenotypic plasticity's impacts on diversification and speciation.
Trends in Ecology & Evolution, 25(8), 459-467.
Piché, J., Iverson, S. J., Parrish, F. A., & Dollar, R. (2010). Characterization of forage fish and
invertebrates in the northwestern Hawaiian Islands using fatty acid signatures: species
and ecological groups. *Marine Ecology Progress Series, 418*, 1-15.
doi:10.3354/meps08814
Pielou, E. C. (2008). *After the ice age: the return of life to glaciated North America*: University
of Chicago Press.
Pires, M. M., Guimarães, P. R., Araújo, M. S., Giaretta, A. A., Costa, J. C. L., & dos Reis, S. F.
(2011). The nested assembly of individual-resource networks. *Journal of Animal*
Ecology, 80(4), 896-903. doi:10.1111/j.1365-2656.2011.01818.x
Præbel, K., Knudsen, R., Siwertsson, A., Karhunen, M., Kahilainen, K. K., Ovaskainen, O., . . .
Amundsen, P. A. (2013). Ecological speciation in postglacial European whitefish: rapid
adaptive radiations into the littoral, pelagic, and profundal lake habitats. *Ecology and*
Evolution, 3(15), 4970-4986.
Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using
multilocus genotype data. *Genetics, 155*(2), 945-959.
Quevedo, M., Svanbäck, R., & Eklöv, P. (2009). Intrapopulation niche partitioning in a
generalist predator limits food web connectivity. *Ecology, 90*(8), 2263-2274.
doi:10.1890/07-1580.1
Quinn, T. J., II, & Deriso, R. B. (1999). *Quantitative fish dynamics*. New York: Oxford
University Press.
Richardson, J. L., Urban, M. C., Bolnick, D. I., & Skelly, D. K. (2014). Microgeographic
adaptation and the spatial scale of evolution. *Trends in Ecology & Evolution, 29*(3), 165-
176.
Roesti, M., Kueng, B., Moser, D., & Berner, D. (2015). The genomics of ecological vicariance in
threespine stickleback fish. *6*, 8767. doi:10.1038/ncomms9767
http://dharmasastra.live.cf.private.springer.com/articles/ncomms9767#supplementary-
information
Ronconi, R., Koopman, H. N., McKinstry, C., Wong, S., & Westgate, A. J. (2010). Inter-annual
variability in diet of non-breeding pelagic seabirds *Puffinus* spp. at migratory staging
areas: evidence from stable isotopes and fatty acids. *Marine Ecology Progress Series,*
419, 267-282.
Rosenberg, N. A. (2004). DISTRUCT: a program for the graphical display of population
structure. *Molecular Ecology Notes, 4*(1), 137-138.
Roughgarden, J. (1972). Evolution of niche width. *American Naturalist*, 683-718.
Rousset, F. (2008). genepop'007: a complete re□implementation of the genepop software for
Windows and Linux. *Molecular Ecology Resources, 8*(1), 103-106.

Rueger, T., Gardiner, N. M., & Jones, G. P. (2016). Size matters: male and female mate choice leads to size-assortative pairing in a coral reef cardinalfish. *Behavioral Ecology, 27*(6), 1585-1591.

Ryman, N., & Palm, S. (2006). POWSIM: a computer program for assessing statistical power when testing for genetic differentiation. *Molecular Ecology Notes, 6*(3), 600-602.

Scharnweber, K., Strandberg, U., Karlsson, K., & Eklöv, P. (2016). Decrease of Population Divergence in Eurasian Perch (Perca fluviatilis) in Browning Waters: Role of Fatty Acids and Foraging Efficiency. *PloS one, 11*(9), e0162470.

Schluter, D. (2000). *The ecology of adaptive radiation*. Oxford: Oxford University Press.

Servedio, M. R., Doorn, G. S. V., Kopp, M., Frame, A. M., & Nosil, P. (2011). Magic traits in speciation: 'magic' but not rare? *Trends in Ecology & Evolution, 26*(8), 389-397. doi:http://dx.doi.org/10.1016/j.tree.2011.04.005

Sharpe, D. M., Räsänen, K., Berner, D., & Hendry, A. P. (2008). Genetic and environmental contributions to the morphology of lake and stream stickleback: implications for gene flow and reproductive isolation. *Evolutionary Ecology Research, 10*(6), 849-866.

Skúlason, S., Parsons, K. J., Svanbäck, R., Räsänen, K., Ferguson, M. M., Adams, C. E., . . . Snorrason, S. S. (2019). A way forward with eco evo devo: an extended theory of resource polymorphism with postglacial fishes as model systems. *Biological Reviews, 0*(0). doi:10.1111/brv.12534

Skulason, S., & Smith, T. B. (1995). Resource polymorphisms in vertebrates. *Trends in Ecology & Evolution, 10*, 366-370.

Skúlason, S., Snorrason, S. S., & Jonsson, B. (1999). Sympatric morphs, populations and speciation in freshwater fish with emphasis on arctic charr. In A. E. Magurran & R. M. May (Eds.), *Evolution of biological diversity* (pp. 329). London: Oxford University Press.

Smith, J., Baumgartner, L., Suthers, I., & Taylor, M. (2011). Generalist niche, specialist strategy: the diet of an Australian percichthyid. *Journal of Fish Biology, 78*(4), 1183-1199.

Snorrason, S. S., & Skúlason, S. (2004). Adaptive speciation in northern freshwater fishes. *Adaptive speciation. Cambridge University Press, Cambridge*, 210-228.

Stowasser, G., Pierce, G. J., Moffat, C. F., Collins, M. A., & Forsythe, J. W. (2006). Experimental study on the effect of diet on fatty acid and stable isotope profiles of the squid Lolliguncula brevis. *Journal of Experimental Marine Biology and Ecology, 333*(1), 97-114.

Sušnik, S., Knizhin, I., Snoj, A., & Weiss, S. (2006). Genetic and morphological characterization of a Lake Ohrid endemic, Salmo (Acantholingua) ohridanus with a comparison to sympatric Salmo trutta. *Journal of Fish Biology, 68*(A), 2-23. doi:10.1111/j.0022-1112.2006.00902.x

Svanbäck, R., & Bolnick, D. I. (2005). Intraspecific competition affects the strength of individual specialization: an optimal diet theory method. *Evolutionary Ecology Research, 7*(7), 993-1012.

Svanbäck, R., & Bolnick, D. I. (2007). Intraspecific competition drives increased resource use diversity within a natural population. *Proceedings of the Royal Society of London B: Biological Sciences, 274*(1611), 839-844.

Svanbäck, R., Mario Pineda☐Krch, a. M. D., Krch, M., Doebeli, M., Associate Editor: Claire de, M., & Editor: Donald, L. D. (2009). Fluctuating Population Dynamics Promotes the Evolution of Phenotypic Plasticity. *American Naturalist, 174*(2), 176-189. doi:10.1086/600112

Svanbäck, R., & Persson, L. (2004). Individual diet specialization, niche width and population
dynamics: implications for trophic polymorphisms. *Journal of Animal Ecology, 73*(5),
973-982.
Svanbäck, R., Quevedo, M., Olsson, J., & Eklöv, P. (2015). Individuals in food webs: the
relationships between trophic position, omnivory and among-individual diet variation.
Oecologia, 178(1), 103-114.
Svanbäck, R., & Schluter, D. (2012). Niche specialization influences adaptive phenotypic
plasticity in the threespine stickleback. *The American Naturalist, 180*(1), 50-59.
Team, R. C. (2015). R: a language and environment for statistical computing. Vienna, Austria;
2015. In.
Team, R. C. (2017). *R: A language and environment for statistical computing. R Foundation for
Statistical Computing*. Vienna, Austria.
Tucker, S., Bowen, W. D., & Iverson, S. J. (2008). Convergence of diet estimates derived from
fatty acids and stable isotopes within individual grey seals. *Marine Ecology Progress
Series, 354*, 267.
Turcotte, M. M., Reznick, D. N., & Hare, J. D. (2011). The impact of rapid evolution on
population dynamics in the wild: experimental test of eco□evolutionary dynamics.
Ecology Letters, 14(11), 1084-1092.
Van Kleunen, M., & Fischer, M. (2005). Constraints on the evolution of adaptive phenotypic
plasticity in plants. *New Phytologist, 166*(1), 49-60.
Van Oosterhout, C., Hutchinson, W. F., Wills, D. P., & Shipley, P. (2004).
MICRO□CHECKER: software for identifying and correcting genotyping errors in
microsatellite data. *Molecular Ecology Notes, 4*(3), 535-538.
Violle, C., Enquist, B. J., McGill, B. J., Jiang, L., Albert, C. H., Hulshof, C., . . . Messier, J.
(2012). The return of the variance: intraspecific variability in community ecology. *Trends
in Ecology & Evolution, 27*(4), 244-252.
Virtue, P., Mayzaud, P., Albessard, E., & Nichols, P. (2000). Use of fatty acids as dietary
indicators in northern krill, *Meganyctiphanes norvegica*, from northeastern Atlantic,
Kattegat, and Mediterranean waters. *Canadian Journal of Fisheries and Aquatic
Sciences, 57*(S3), 104-114.
Vonlanthen, P., Roy, D., Hudson, A. G., LargiadÈR, C. R., Bittner, D., & Seehausen, O. (2009).
Divergence along a steep ecological gradient in lake whitefish (*Coregonus* sp.). *Journal
of Evolutionary Biology, 22*(3), 498-514. doi:10.1111/j.1420-9101.2008.01670.x
Wainwright, P. C., & Price, S. A. (2016). The Impact of Organismal Innovation on Functional
and Ecological Diversification. *Integrative and Comparative Biology, 56*(3), 479-488.
doi:10.1093/icb/icw081
Webb, P. W. (1984). Body form, locomotion and foraging in aquatic vertebrates. *American
Zoologist, 24*, 107-120.
Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the analysis of population
structure. *Evolution*, 1358-1370.
West-Eberhard, M. J. (2003). *Developmental plasticity and evolution*: Oxford University Press.
Zelditch, M. L., Swiderski, D. L., & Sheets, H. D. (2012). *Geometric morphometrics for
biologists: a primer*: Academic Press.
Zhao, T., Villéger, S., Lek, S., & Cucherousset, J. (2014). High intraspecific variability in the
functional niche of a predator is associated with ontogenetic shift and individual
specialization. *Ecology and Evolution, 4*(24), 4649.

Zimmerman, M. S., Krueger, C. C., & Eshenroder, R. L. (2006). Phenotypic diversity of lake
trout in Great Slave Lake: differences in morphology, buoyancy, and habitat depth.
Transactions of the American Fisheries Society, 135, 1056-1067.
Zimmerman, M. S., Krueger, C. C., & Eshenroder, R. L. (2007). Morphological and ecological
differences between shallow- and deep-water lake trout in Lake Mistassini, Quebec.
Journal of Great Lakes Research, 33, 156-169.

1030
1031

1032   Table 1. Mean composition (% ± SD) of 41 fatty acids for the four groups of piscivorous Lake

1033   Trout morph identified from Great Bear Lake.

Fatty acids	Group 1	Group 2	Group 3	Group 4
14:0	6.8 ± 1.0	7.0 ± 1.0	9.2 ± 1.0	9.9 ± 1.1
16:0	28.1 ± 1.0	28.13 ± 1.7	24.2 ± 1.7	26.1 ± 1.2
16:1n-7	15.9 ± 3.7	10.1 ± 2.0	19.5 ± 3.4	15.6 ± 2.1
16:2n-6	2.0 ± 0.5	2.4 ± 0.6	2.6 ± 0.2	3.1 ± 0.2
16:2n-4	2.6 ± 0.7	1.5 ± 0.4	2.7 ± 0.9	2.3 ± 0.4
17:0	2.7 ± 0.5	2.8 ± 0.3	2.4 ± 0.4	2.8 ± 0.2
16:3n-4	1.5 ± 0.7	1.4 ± 0.5	1.9 ± 0.6	1.8 ± 0.9
16:4n-3	2.6 ± 1.2	0.8 ± 0.3	1.3 ± 0.6	1.2 ± 0.4
16:4n-1	1.6 ± 0.7	1.5 ± 0.8	0.9 ± 0.6	1.0 ± 0.6
18:0	14.2 ± 1.6	13.1 ± 0.8	11.6 ± 0.7	11.7 ± 0.5
18:1n-9	20.6 ± 4.1	18.5 ± 3.4	32.3 ± 3.9	27.9 ± 3.5
18:1n-7	11.9 ± 2.4	9.5 ± 1.0	13.9 ± 1.2	12.5 ± 0.8
18:2n-6	8.6 ± 1.5	9.2 ± 1.6	12.4 ± 1.2	12.9 ± 1.0
18:2n-4	2.0 ± 0.4	1.5 ± 0.2	2.1 ± 0.2	2.1 ± 0.2
18:3n-6	2.2 ± 0.8	1.5 ± 0.4	2.5 ± 0.4	2.3 ± 0.2
18:3n-4	2.2 ± 0.7	1.5 ± 0.3	2.4 ± 0.4	2.0 ± 0.3
18:3n-3	6.6 ± 1.4	6.9 ± 0.9	7.9 ± 0.6	8.7 ± 0.7
18:3n-1	1.2 ± 0.7	1.2 ± 0.3	1.1 ± 0.3	1.5 ± 0.3
18:4n-3	3.5 ± 0.7	4.0 ± 1.2	4.9 ± 0.7	5.6 ± 0.7
18:4n-1	1.3 ± 0.6	0.4 ± 0.5	0.9 ± 0.5	1.2 ± 0.6
20:0	2.1 ± 0.7	2.8 ± 0.7	3.1 ± 0.6	2.8 ± 0.8
20:1n-11	1.7 ± 1.0	0.8 ± 0.5	1.9 ± 0.8	1.4 ± 0.4
20:1n-9	6.0 ± 1.4	4.2 ± 0.8	7.9 ± 0.9	7.1 ± 0.9
20:1n-7	2.5 ± 0.4	2.5 ± 0.3	3.8 ± 0.4	4.1 ± 0.6
20:2n-9	0.8 ± 0.6	1.4 ± 0.8	1.3 ± 0.4	1.2 ± 0.4
20:2n-6	3.8 ± 0.9	4.7 ± 0.9	6.8 ± 1.3	7.5 ± 1.0
20:3n-6	3.4 ± 0.5	3.6 ± 0.4	4.4 ± 0.5	4.0 ± 0.4
20:4n-6	13.8 ± 1.7	14.2 ± 1.3	10.1 ± 1.1	10.0 ± 1.2
20:3n-3	3.5 ± 0.7	4.5 ± 0.9	5.1 ± 0.6	6.6 ± 0.7
20:4n-3	6.1 ± 1.2	8.2 ± 1.3	8.8 ± 1.1	10.8 ± 0.9
20:5n-3	18.0 ± 2.9	15.7± 1.2	11.8 ± 2.1	12.2 ± 1.8
22:1n-11	1.8 ± 1.7	0.9 ± 0.5	1.0 ± 1.3	0.9 ± 0.4
22:1n-9	2.2 ± 0.5	2.4 ± 0.4	3.3 ± 0.4	3.1 ± 0.4
22:1n-7	1.2 ± 0.6	1.0 ± 0.5	1.1 ± 0.3	1.6 ± 0.4
22:2n-6	1.4 ± 0.5	1.7 ± 0.6	3.0 ± 0.5	4.0 ± 0.8
21:5n-3	0.9 ± 0.6	1.8 ± 0.6	2.2 ± 0.6	1.6 ± 0.9
22:4n-6	0.2 ± 0.5	1.0 ± 1.6	0.3 ± 0.6	1.6 ± 1.7
22:5n-6	7.6 ±1.1	10.7 ± 1.4	7.7 ± 0.7	9.6 ± 1.4
22:4n-3	2.3 ± 0.9	4.2 ± 1.3	5.1 ± 0.9	7.2 ± 1.7
22:5n-3	10.4 ± 0.9	10.8 ± 0.6	10.4 ± 2.4	11.1 ± 0.7
22:6n-3	33.9 ± 5.6	38.9 ± 4.3	23.1 ± 3.7	26.3 ± 4.7

1034

1035   Table 2. Number of individuals genotyped (N), number of alleles (N_A), expected heterozygosity

1036   (H_E), observed heterozygosity (H_O), allelic richness (A_R) and private allelic richness (PA_R) within

1037   fatty acid groups identified within a piscivorous morphotype of Lake Trout from Canada's Great

1038   Bear Lake.

	N	N_A	H_E	H_O	A_R	PA_R
Group 1	12	10.16	0.85	0.83	9.87	1.08
Group 2	16	11.26	0.84	0.82	9.57	0.99
Group 3	20	12.32	0.84	0.81	9.70	0.87
Group 4	28	14.11	0.84	0.81	9.57	0.98
Giant	39	15.95	0.84	0.78	9.69	1.05

1039

1040

 Table 3. Pairwise F_{ST} based on variation at microsatellite loci among Lake Trout morphs from

1042 Harris et al. (2015) and piscivorous fatty acids dietary groups from Great Bear Lake. Significant

1043 results are represented as follow: * values are significant at an initial α of 0.05 and ** values are

1044 significant at an α of 0.02 subsequent False Discovery Rate adjustments for multiple

1045 comparisons.

	Morph 1	Morph 2	Morph 3		Group 1	Group 2	Group 3	Group 4
Morph 1				**Group 1**				
Morph 2	0.063**			**Group 2**	0.003			
Morph 3	0.004**	0.007**		**Group 3**	0.001	-0.01		
Morph 4	0.012**	0.017**	0.009**	**Group 4**	0.005	-0.004	-0.002	
				Giant	0.016**	0.001	-0.002	0.006**

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

analysis (based on lnP[D] and the ΔK statistic of Evanno et al. (2005)) and DAPC analysis

(based on the lowest BIC score and with 30 PCs retained).

Fig. 5. Canonical Variate Analyses (95% ellipses) and Principal Components Analysis of body

shape (a, d), head shape (b, e) and linear measurements (c, f), respectively, of piscivorous Lake

Trout represented as: square = Group 1, circle = Group 2, triangle = Group 3, and diamond =

Group 4. The first two PCA axes explained 44.3% and 12.3 % of variation for body shape,

35.1% and 30.7 % of variation for head shape, and 39.6 % and 20.9 % for linear measurements

(Fig. 6 d, e, f). To visualize their variation within and among groups, individuals longer than 900

mm FL, which are considered the Giant sub-set in this study, are depicted by larger symbols.

Fig. 6. Residuals of mean (± 95%CI) size-standardized upper and lower jaw lengths, head depth

and length, and snout-eye length among piscivorous Lake Trout groups. Grouping symbols are

as follows: square = Group 1, circle = Group 2, triangle = Group 3, and diamond = Group 4.

1095

1096

1097

1098

1099

1100

1101

1102

1103
1104

1105

1106

1107
1108
1109   Fig. 1.

PC2: 15,8%

1131

1132
1133   Fig. 2

$$L_t = 1{,}129 \times (1 - e^{-0.035 \times (t+8.84)})$$

$$L_t = 849 \times (1 - e^{-0.086 \times (t+2.11)})$$

1134

1135   Fig. 3.

1136

1137

1148

1149   Fig. 4.

1175

1176

1177   Fig. 5

1178

1179
1180   Fig. 6

1204 **Appendix**

1205

1206 Table A1. Spatial and temporal information for the 79 Lake Trout classified as piscivorous

1207 morph from Great Bear Lake and analyzed for fatty acids. Sample sizes are in brackets.

Sample information		
Group 1 (14)	Dease 2005 (1)	Smith 2011 (1)
	Dease 2010 (1)	
	McTavish 2009 (6)	
	McVicar 2008 (1)	
	Smith 2006 (4)	
Group 2 (16)	Dease 2010 (4)	Smith 2006 (1)
	Keith 2012 (2)	Smith 2011 (5)
	McTavish 2009 (1)	
	McTavish 2014 (2)	
	McVicar 2008 (1)	
Group 3 (21)	Dease 2005 (2)	McVicar 2003 (2)
	Dease 2010 (3)	McVicar 2008 (2)
	Keith 2012 (3)	McVicar 2013 (3)
	McTavish 2004 (1)	Smith 2006 (1)
	McTavish 2014 (2)	Smith 2011 (2)
Group 4 (28)	Dease 2005 (5)	McTavish 2014 (1)
	Dease 2010 (2)	McVicar 2003 (2)
	Keith 2002 (4)	McVicar 2008 (5)
	Keith 2003 (4)	Smith 2006 (1)
	MCTavish 2004 (2)	Smith 2011

Table A2. Bayesian clustering (i.e., STRUCTURE, Pritchard et al. 2000) results for piscivorous
morphotypes of lake trout from Great Bear Lake assessed using variation at 19 microsatellite
markers. Shown are the mean log$\square$likelihood values (LnP[D]) for different hypothesized
numbers of genetic populations (K) and the mean value of ΔK statistic of Evanno et al. (2005).
Bold values represent the most likely number of genetic groups indicated by ΔK. Dashes = not
applicable given that ΔK cannot be calculated for these values of K. For all STRUCTURE
analyses, we employed an admixture model with the LOCPRIOR algorithm, correlated allelic
frequencies, 100,000 burn-in and MCMC iterations and 10 iterations per K value were
completed.

K	Reps	Mean LnP(D)	Delta K
1	10	-10271.83	—
2	10	**-10266.25**	**9.26**
3	10	-10572.68	0.03
4	10	-10868.33	1.39
5	10	-10739.53	0.45
6	10	-10806.97	0.84
7	10	-10678.37	0.66
8	10	-10739.90	0.13
9	10	-10862.98	1.09
10	10	-10553.04	—

1222

1223

1224
1225
1226
1227

Table A3. Results of the discriminant analysis of principal components (DAPC, Jombart et al.
2010) implemented in the Adegenet package (Jombart et al. 2008) to determine the most likely
number of genetic clusters (K) within the piscivorous Lake Trout form Great Bear Lake. The
number of groups was identified using the find.clusters function (a sequential K-means
clustering algorithm) and subsequent Bayesian Information Criterion (BIC), as suggested by
Jombart et al. (2010). Stratified cross-validation carried out with the function *xvalDapc* was
employed to determine the optimal number of PCs to retain in the analysis.
1246

K	BIC
1	185.98
2	185.42
3	185.89
4	186.51
5	187.40
6	189.10
7	190.64
8	191.99
9	193.61
10	195.67

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

1257

1258

1259   Table A4. Microsatellite loci used in this study and F_{is} values for each group per locus.

Locus	Group 1	Group 2	Group 3	Group 4	Giant
OtsG83b	-0.021	0.017	0.046	0.090	-0.013
Sco215	0.061	0.042	-0.011	-0.029	0.071
Smm17	-0.433	-0.069	-0.038	-0.304	-0.012
Smm21	-0.143	-0.286	-0.266	0.023	0.028
SnaMSU1	0.012	-0.069	-0.024	-0.028	-0.074
SnaMSU8	-0.031	0.002	0.048	0.023	0.081
OMM1105	0.094	-0.075	-0.041	-0.098	-0.065
Smm22	-0.014	0.055	0.082	-0.088	0.137
SnaMSU13	-0.105	0.053	0.136	-0.073	-0.049
SnaMSU5	0.088	0.065	-0.032	0.039	0.159
Sco19	-0.082	0.190	0.067	-0.009	0.051
Sco202	0.107	-0.166	0.047	0.115	-0.080
SnaMSU10	-0.108	-0.030	0.086	0.096	0.203
SnaMSU12	0.122	0.069	0.123	0.201	0.072
SnaMSU6	0.008	-0.090	0.002	0.007	0.207
Sal38	-0.056	0.121	-0.016	-0.012	0.041

Sco200	-0.015	0.098	-0.096	0.041	0.244
SnaMSU11	-0.060	-0.108	-0.012	0.083	-0.011
SnaMSU3	0.059	-0.019	0.065	0.012	0.085
Overall	-0.027	-0.011	0.009	0.005	0.057

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269 Fig. A1. The four shallow-water morphotypes of Lake Trout from Great Bear Lake identified in
1270 Chavarie et al. (2013, 2015, 2016a, 2016b): the generalist, the piscivore, the benthic-oriented,
1271 and the pelagic specialist, Morphs1-4, respectively.

1272

1273

1274

1275

1276

Hierarchical clustering on the factor map

1277
1278 Fig. A2. Hierarchical clusters of Great Bear Lake Lake Trout fatty acids profiles overlaid on the
1279 first two principal component axes (PCA) using FactoMineR.

1297
1298 Fig. A3. Principal Component Analysis (PCA) of fatty acids of 79 Lake Trout classified as
1299 piscivorous morph from Great Bear Lake, based on the proportions of 41 fatty acids in dorsal
1300 muscle tissue. Spatial variations (5 arms; 1=Keith, 2=McVicar, 3=McTavish, 4=Dease, and
1301 5=Smith) are represented, based on the fatty acids profile of each lake trout analyzed in this
1302 study.

1325
1326 Fig. A4. Depth of capture for four groups of piscivorous Lake Trout from Great Bear Lake
1327 (Groups identified by fatty acids profiles of individuals). Outliers are represented by a circle.
1328
1329
1330
1331

1332

1333
Fig. A5. Summary of the DAPC analysis. (A) Results of the cross-validation analysis used to
determine the number of PCs to retain in the DAPC analysis. Cross-validation analysis
determined the most appropriate number of PCs retained was 30. (B) Inference of the number of
clusters in the DAPC performed on piscivorous Lake Trout from Great Bear Lake. The function
find.clusters was run with a maximum number of clusters of 10 to identify the optimal number of
clusters based on the BIC values. A K value of 2 (the lowest BIC value) represents the best
summary of the data (most probable number of (K)). (C) The results of the discriminant function
that shows that the two clusters are mostly non-overlapping.