A transcriptional regulatory atlas of human pancreatic islets reveals non-coding functional signatures at GWAS loci

Arushi Varshney[1,2], Yasuhiro Kyono[1,3], Venkateswaran Ramamoorthi Elangovan[1], Collin Wang[1,4], Michael R. Erdos[5], Narisu Narisu[5], Ricardo D'Oliveira Albanus[1], Peter Orchard[1], Michael L. Stitzel[6], Francis S. Collins[5], Jacob O. Kitzman[1,2], Stephen C. J. Parker[1,2,*]

Department of Computational Medicine & Bioinformatics, University of Michigan, Ann Arbor, MI, USA

Department of Human Genetics, University of Michigan, Ann Arbor, MI, USA

Current address: Tempus Labs, Inc., Chicago, IL, USA

Current address: Columbia University, New York, NY, USA

National Human Genome Research Institute, NIH, Bethesda, MD, USA

The Jackson Laboratory for Genomic Medicine, Farmington, CT, USA

* Corresponding author scjp@umich.edu

Abstract

Identifying the tissue-specific molecular signatures of active regulatory elements is critical to understand gene regulatory mechanisms. Active enhancers can be transcribed into enhancer RNA. Here, we identify transcription start sites (TSS) using cap analysis of gene expression (CAGE) across 71 human pancreatic islet samples. We identify 9,954 CAGE tag clusters (TCs) in islets, ~20% of which are islet-specific and occur mostly distal to known gene TSSs. We integrated islet CAGE data with histone modification and chromatin accessibility profiles to identify epigenomic signatures of transcription initiation. Using a massively parallel reporter assay, we observe significant enhancer activity (5% FDR) for 2,279 of 3,378 (~68%) tested CAGE elements. TCs within accessible enhancers show higher enrichment to overlap type 2 diabetes genome-wide associated loci than existing accessible enhancer annotations, which emphasizes the utility of mapping CAGE profiles in disease-relevant tissue. This work provides a high-resolution transcriptional regulatory map of human pancreatic islets with utility for dissecting functional enhancers at GWAS loci.

Introduction
Type 2 diabetes (T2D) is a complex disease that results from an interplay of factors such as pancreatic islet dysfunction and insulin resistance in peripheral tissues such as adipose and muscle. Genome wide association studies (GWAS) to date have identified >240 loci that modulate risk for T2D (Mahajan *et al.* 2018). However, these SNPs mostly occur in non protein-coding regions and are highly enriched to overlap islet-specific enhancer regions (Parker *et al.* 2013; Pasquali *et al.* 2014; Quang *et al.* 2015; Thurner *et al.* 2018; Cebola 2019). These findings suggest the variants likely affect gene expression rather than directly altering protein structure or function. Moreover, due to the correlated structure of common genetic variations across the genome, GWAS signals are usually marked by numerous SNPs in high linkage disequilibrium (LD). Therefore, identifying causal SNP(s) is extremely difficult using genetic information alone. These factors have impeded our understanding of the molecular mechanisms by which genetic variants modulate gene expression in orchestrating disease.

In order to understand gene regulatory mechanisms, it is essential to identify regulatory elements at high genomic
resolution. Active regulatory elements can be delineated by profiling covalent modifications of the histone H3
subunit such as H3 lysine 27 acetylation (H3K27ac) which is associated with enhancer activity (Creyghton *et al.*
2010; Zhou *et al.* 2011), and H3 lysine 4 trimethylation (H3K4me3) which is associated with promoter activity
among others (Mikkelsen *et al.* 2007; Adli *et al.* 2010; Zhou *et al.* 2011). However, such histone modification
based profiling methods identify regions of the genome that typically span hundreds of base pairs. Since TF
binding can affect gene expression, TF accessible regions of the chromatin within these broader regulatory
elements enable higher-resolution identification of the functional DNA bases. Numerous studies have utilized
diverse chromatin information in pancreatic islets to nominate causal gene regulatory mechanisms (Fadista *et*
al. 2014; Bunt *et al.* 2015; Varshney *et al.* 2017; Roman *et al.* 2017; Thurner *et al.* 2018; Mahajan *et al.* 2018).
50

Studies have shown that a subset of enhancers are transcribed into enhancer RNA (eRNA), and that transcription
is a robust predictor of enhancer activity (Andersson *et al.* 2014; Mikhaylichenko *et al.* 2018). eRNAs are nuclear,
short, mostly-unspliced, 5' capped, usually non-polyadenylated and usually bidirectionally transcribed (Kim *et al.*
2010; Melgar *et al.* 2011; Andersson *et al.* 2014). Previous studies have indicated that eRNA transcripts could
be the stochastic output of Pol2 and TF machinery at active regions, or in some cases, the transcripts could
serve important functions such as sequestering TFs or assisting in chromatin looping (Li *et al.* 2013; Kaikkonen
et al. 2013; Hsieh *et al.* 2014; Yang *et al.* 2016). Therefore, identifying the location of transcription initiation can
pinpoint active regulatory elements.
59

Genome-wide sequencing of 5′ capped RNAs using CAGE can detect TSSs and thereby profile transcribed
promoter and enhancer regions (Kim *et al.* 2010; Andersson *et al.* 2014). CAGE-identified enhancers are two to
three times more likely to validate in functional reporter assays than non-transcribed enhancers detected by
chromatin-based methods (Andersson *et al.* 2014). An advantage of CAGE is that it can be applied on RNA
samples from hard to acquire biological tissue such as islets and does not require live cells that are imperative
for other TSS profiling techniques such as GRO-cap seq (Core *et al.* 2008, 2014; Lopes *et al.* 2017). The
functional annotation of the mammalian genome (FANTOM) project (The FANTOM Consortium *et al.* 2014) has
generated an exhaustive CAGE expression atlas across 573 primary cell types and tissues, including the
pancreas. However, pancreatic islets that secrete insulin and are relevant for T2D and related traits, constitute
only ~1% of the pancreas tissue. Therefore, the bulk pancreas transcriptome does not accurately represent the
islet enhancer transcription landscape. Motivated by these reasons, we profiled the islet transcriptome using
CAGE. Here, we present the CAGE TSS atlas of pancreatic islets, validate this atlas using a massively parallel
reporter assay, and perform integrative analyses across diverse epigenomic datasets to reveal molecular
signatures of non-coding islet elements and their role in T2D and related traits.
74

Results
The CAGE landscape in human pancreatic islets
We analyzed transcriptomes in 71 human pancreatic islet total RNA samples obtained from unrelated organ
donors by performing CAGE. To enrich for the non poly-adenylated and smaller (<1kb) eRNA transcripts, we
performed polyA depletion and fragment size selection (<1kb, see Methods). Strand-specific CAGE libraries
were prepared according to the no-amplification non-tagging CAGE libraries for Illumina next-generation
sequencers (nAnT-iCAGE) protocol (Murata *et al.* 2014), and an 8 bp unique molecular identifier (UMI) was

added during reverse transcription to identify PCR duplicates. We sequenced CAGE libraries, performed pre-
alignment quality control (QC), mapped to the hg19 reference genome, performed post-alignment QC, and
identified CAGE tags. We selected 57 samples that passed our QC measures (see Methods) for all further
analyses. We identified regions with a high density of transcription initiation events or TCs using the paraclu
(Frith *et al.* 2008) method in each islet sample in a strand-specific manner. We then identified a consensus set
of aggregated islets TCs by merging overlapping TCs on the same strand across samples and retaining
segments that were supported by at least 10 individual samples (see Methods, Supplementary Figure 1). We
identified 9,954 tag clusters with median length of 176 bp (Supplementary Figure 2), spanning a total genomic
territory of ~2.4 Mb. As a resource to identify transcribed gene isoforms in Islets, Supplementary table 1 includes
the closest identified TCs to known gene TSSs (Gencode V19, (Harrow *et al.* 2012)). To analyze characteristics
of islet TCs and explore the chromatin landscape underlying these regions, we utilized publicly available ChIP-
seq data for various histone modifications along with ATAC-seq data in islets (Varshney *et al.* 2017). We
integrated ChIP-seq datasets for promoter associated H3K4me3, enhancer associated H3K4me1, active
promoter and enhancer associated H3K27ac, transcribed gene-associated H3K36me3 and repressed chromatin
associated H3K27me3 in islets along with corresponding publicly available ChIP-seq datasets for Skeletal
Muscle, Adipose and Liver (included for other ongoing projects) using ChromHMM (Ernst and Kellis 2010, 2012;
Ernst *et al.* 2011). This analysis produced 11 distinct and recurrent chromatin states (Supplementary Figure 3),
including promoter, enhancer, transcribed, and repressed states. Figure 1A shows an example locus in the
intronic region of the *ST18* gene where a TC identified in islets overlaps the active TSS chromatin state and an
ATAC-seq peak. The regulatory activity of this element was validated by the VISTA project in an *in vivo* reporter
assay in mouse embryos (Visel *et al.* 2007).
104
We next compared the islet TCs with TCs in across 118 human tissues where CAGE data was available through
the FANTOM project (The FANTOM Consortium *et al.* 2014). We identified TCs in 118 FANTOM human tissues
using the paraclu method with the same parameters as Islets (see Methods). For each islet TC segment, we
then calculated the number of FANTOM tissues in which TCs overlapped the segment. We observed that ~20%
of islet TCs were unique to islets (1,974 TCs had no overlap with TCs in any FANTOM tissues), whereas about
~60% of islet TCs were shared across 60 or more FANTOM tissues (Figure 1B). We then asked if the more
islet-specific TCs occurred more distal to known protein coding gene TSSs. We categorized islet TC segments
based on the number of FANTOM tissues in which they overlap TCs, namely no overlap, overlap in 1-40, 41-
100 or 100-118 FANTOM tissues (colored bars in Figure 1B). We calculated the distance to the nearest known
protein-coding gene TSS (Gencode V19) and observed a clear pattern where islet-specific TCs (no overlap with
FANTOM) occurred farthest from known TSSs, and TCs overlapping more FANTOM tissues occurred closer to
gene TSSs (Figure 1C). We highlight an example locus where an islet TC in the *AP1G2* gene occurs in active
TSS chromatin states across multiple tissues, and overlaps shared ATAC-seq peaks in islet and the
lymphoblastoid cell line GM12878 (Buenrostro *et al.* 2013) (Figure 1D, blue box). TCs across FANTOM tissues
are identified in this region (Figure 1D, FANTOM TCs track). The islet TC segment (Figure 1D, blue dashed box)
overlaps TCs in 88 FANTOM tissues. Another islet TC ~34kb away, however, occurs in a region lacking gene
annotations, and overlaps a more islet-specific active enhancer chromatin state and ATAC-seq peak (Figure 1D,
orange dashed box). This region was not identified as a TC in any of the 118 analyzed FANTOM tissues. At
other islet-relevant loci such as the insulin (*INS)* gene locus, we observe TCs in islets and also in Pancreas and
Duodenum tissues from the FANTOM data (Supplementary Figure 4). Collectively, these results highlight that
CAGE profiling in islets identifies islet-specific sites of active transcription initiation.
126

We next asked if islet TCs preferentially overlap certain genomic annotations. We computed the enrichment of
islet TCs to overlap islet annotations such as active TSS, enhancer and other chromatin states and islet ATAC-
seq peaks. We also included 'common' annotations such as known gene promoters, coding, untranslated (UTR)
regions, or annotations such as super enhancers, or histone-modification ChIP-seq peaks that were aggregated
across multiple cell types. We observed that islet TCs were highly enriched to overlap islet active TSS chromatin
states (fold enrichment = 69.72, P value = 0.0001, Figure 1E, Supplementary Table 2). This result is expected
since CAGE profiles transcription start sites where the underlying chromatin is more likely to resemble the 'active
TSS' chromatin state. TCs were also enriched to overlap islet ATAC-seq peaks (fold enrichment = 53.8, P value
= 0.0001, Figure 1E), signifying that the identified transcription initiation sites constitute accessible chromatin
where TFs can bind.
137
To gauge if these transcribed elements could be relevant for diverse disease/traits, we computed enrichment for
islet TCs to overlap GWAS loci for 116 traits from the NHGRI catalog (Buniello *et al.* 2019). We observed that
traits such as Fasting Glucose (FGlu) (fold enrichment = 7.05, P value = 3.30×10^{-4}), metabolic traits (fold
enrichment = 6.46, P value = 2.03×10^{-4}) were among the most highly enriched, highlighting the relevance of
these transcribed elements for islet biology (Figure. 1F, Supplementary Table 3). GWAS loci for T2D were also
enriched in islet TCs (fold enrichment = 2.48, P value = 0.02). Because T2D is orchestrated through a complex
interplay between islet beta cell dysfunction and insulin resistance in peripheral tissues, we reasoned that some
underlying pathways in T2D might be more relevant in islets than others. To explore this rationale, we utilized
results from a previous study that analyzed established GWAS data for T2D along with 47 other diabetes related
traits and identified clusters of T2D GWAS signals (Udler *et al.* 2018). Interestingly, we observe that GWAS loci
in the islet beta cell and proinsulin cluster were highly enriched to overlap islet TCs (fold enrichment = 5.62, P
value = 0.004), whereas loci in the insulin resistance cluster were not enriched (fold enrichment = 0.97, P value
= 0.672; Figure 1F, Supplementary Table 3). These results suggest that our islet TC atlas is comprised of active
regulatory elements relevant for traits specifically related to islet function.
152
Integrating CAGE TCs with epigenomic information
We further explored CAGE profiles relative to the underlying chromatin landscape to identify signatures of
transcription initiation. We first overlayed CAGE profiles over ATAC-seq data. Aggregated CAGE signal over
ATAC-seq narrow peak summits highlighted a bidirectional pattern of transcription initiation flanking the ATAC-
seq peak summit on both strands (Figure 2A). Conversely, on anchoring the ATAC-seq signal over islet TC
centers we observed that the summit of the ATAC-seq signal lies upstream of the TC center (Figure 2B). In these
analyses, upstream or downstream positions were determined from the strand the CAGE signal mapped to. We
next asked if TF binding sites were more enriched to occur upstream or downstream of the TC. We utilized TF
footprint motifs previously identified using islet ATAC-seq data and TF DNA binding position weight matrices
(PWMs) (Varshney *et al.* 2017). These footprint motifs represent putative TF binding sites that are also supported
by accessible chromatin profiles, as opposed to TF motif matches that are only informed by DNA sequence. We
observe that most TF footprint motifs were more enriched to overlap the 500 bp TC upstream region compared
to the 500 bp downstream region relative to TCs (Figure 2C). These observations show that, as expected, the
region just upstream of the TC is highly accessible where more TF binding events occur, and indicate the high
quality of our islet TC atlas.
168
We next explored the characteristics of TCs that occur in the two main regulatory classes - promoter and
enhancers, relative to each other. We focussed on transcribed, accessible regions in promoter or enhancer

chromatin states - namely TCs overlapping ATAC-seq peaks in promoter (active, weak or flanking TSS)
chromatin states or enhancer (active, weak or genic enhancer) chromatin states. We considered the proximity
of these elements to known gene TSSs and further classified the segments as TSS proximal or distal using a
5kb distance threshold from the nearest protein coding genes (Gencode V19). We then explored the chromatin
landscape at these regions across 98 Roadmap Epigenomics cell types for which chromatin state annotations
are publicly available (18 state 'extended model', see Methods) (The Roadmap Epigenomics Consortium *et al.*
2015). We observed that TSS proximal islet TCs in accessible islet promoter chromatin states (N = 7,064
segments) were nearly ubiquitously identified as promoter chromatin states across roadmap cell types (Figure
2D, left). A subset of TSS distal islet TCs in accessible islet promoter chromatin states (out of total N = 443
segments) however were more specific for pancreatic islets (Figure 2D, right). In contrast, we observed that islet
TCs in accessible islet enhancer chromatin states, both proximal (N = 254 segments) and distal (N = 289
segments) to known gene TSSs were more specifically identified as enhancer states in pancreatic islets (Figure
2E). This pattern was more clear for Roadmap pancreatic islet segmentations compared to the whole pancreas
segmentations (Figure 2D and E, labelled) which further emphasizes the differences between epigenomic
profiles for pancreatic islets compared to whole pancreas tissue.
186
Having observed differences in cell type-specificities in islet TCs in promoter vs enhancer states, we next asked
if TFs displayed preferences to bind in these regions. We observed that footprint motifs for the regulatory factor
X (RFX) TF family were highly enriched (>3 fold, P value = 0.0001) in TCs in accessible enhancers (Figure 2F).
On the other hand, TCs in accessible promoter regions were highly enriched to overlap footprint motifs of the
E26 transformation-specific (ETS) TF family (Figure 2F). We observe divergent aggregate CAGE profiles over
TF footprint motifs enriched in enhancers; for example RFX5_known8 footprint motifs in 5kb TSS distal regions
and ELK4_1 motif (Figure 2G and H). Taken together, these results show the differences in the characteristics
of transcription initiation sites based on the underlying chromatin context.
195
Experimental validation of transcribed regions
We next sought to experimentally validate the transcriptional activity of islet CAGE-profiled regions. We utilized
a massively parallel reporter assay (MPRA) platform wherein thousands of elements can be simultaneously
tested by including unique barcode sequences for each element and determining the transcriptional regulatory
activity using sequencing-based barcode quantification (Melnikov *et al.* 2012; Arnold *et al.* 2013; Neumayr *et al.*
2019). We generated a library of 7,188 CAGE elements (198 bp each, see Methods) and cloned these along
with a library of 16 bp barcode sequences into the MPRA vector 3' to the polyA signal of the GFP gene. Our
experimental setup therefore assays the 'enhancer' activity of the CAGE element where the readout is the
transcript levels of the multiple barcode sequences corresponding to the CAGE element. We transfected the
MPRA libraries into the rat beta cell insulinoma (INS1 832/13) cell line in triplicate, extracted DNA and RNA and
sequenced the barcodes. We added 6 bp unique molecular identifier (UMI) sequences before the PCR
amplification of the RNA libraries to enable accounting for PCR duplicates while quantifying true biological RNA
copies. We selected barcodes that were observed with at least 10 DNA counts and non-zero RNA counts in at
least one replicate, and selected CAGE elements that were observed with at least two such qualifying barcodes.
This filtering procedure resulted in 3,378 CAGE elements. We observed high correlations between the
normalized sum of RNA counts of the CAGE element barcodes across the three biological replicates (Pearson r
= 0.97 Supplementary Figure 5). We quantified the enhancer reporter activity of the 3,378 CAGE elements by
modeling the RNA and DNA barcode counts using generalized linear models (GLMs) implemented in the
MPRAnalyze package (Ashuach *et al.* 2019) (Supplementary table 4). We then tested for significant

transcriptional activity against a null model (see Methods). We observed that ~68% (N = 2,279) of the testable
CAGE elements showed significant regulatory activity (5% FDR) (Figure 3A, top). On classifying CAGE elements
based on the underlying chromatin landscape such as - promoter (active, weak or flanking TSS) chromatin state,
enhancer (active, weak or genic enhancer) chromatin state or other chromatin state overlap in islets, we
observed that a larger fraction of CAGE elements overlapping the promoter states had significant transcriptional
activity compared to elements overlapping enhancer states, which was in turn higher than CAGE elements in
other chromatin states (Figure 3A, bottom). We also observed that the CAGE elements in promoter chromatin
states had higher MPRA activity Z scores compared to the elements in enhancer chromatin states (Wilcoxon
rank sum test P = 1.02×10^{-6}) (Figure 3B). Z scores for the CAGE elements that overlapped ATAC-seq peaks
were significantly higher than the elements that did not occur in peaks (Wilcoxon rank sum test P = 5.50×10^{-16})
(Figure 3C). Z scores for CAGE elements 5kb proximal to protein-coding gene TSSs (Gencode V19) were higher
than CAGE elements that were distal to gene TSS locations (Wilcoxon rank sum test P = 5.38×10^{-9})
(Supplementary Figure 6). These results highlight that accessible elements in promoter chromatin states show
higher 'enhancer' reporter activities.

We then asked if sequence-based features of the CAGE elements such as the occurrence of TF motifs could
predict the activities of these elements in the MPRA assay. We used linear regression to model transcriptional
activity as a function of TF motif overlap. Since thousands of TF motifs are available and many are correlated
with each other, it is useful to identify a subset of more relevant motifs. For this purpose, we utilized the least
absolute shrinkage and selection operator (LASSO) procedure which uses shrinkage, constraining the model
parameters such that some regression coefficients are shrunk to zero. This results in simpler model with a small
subset of selected features. We trained the lasso regression model with the CAGE element MPRA Z scores as
the response variable and the TF motif scan scores for 540 representative TF motifs in each CAGE element as
predictors (see Methods). We report the TF motifs with the top 30 lasso regression coefficients in (Figure 3D,
full results in Supplementary Table 5). We observed that TF motifs from the ETS family showed positive lasso
coefficients, indicating that these sequence elements are associated with high transcriptional activity. Indeed,
we earlier observed that these motifs were also highly enriched to occur in TCs in accessible promoter chromatin
state regions (Figure 2E). Other TF motifs with positive lasso coefficients included CEBP, YY1, and NRF-1
(Figure 3D). NRF-1, for instance, has been identified to be relevant for islet biology, as its target genes are
downregulated in diabetic individuals (Patti *et al.* 2003); knockdown of this gene in the mouse insulinoma cell
line (MIN6) and beta cell specific Nrf1-knockout mice resulted in impaired glucose responsiveness, elevated
basal insulin release and decreased glucose-stimulated insulin secretion (GSIS) (Zheng *et al.* 2015).

In Figure 3E, we highlight an islet TC for which we tested three tiled elements (Figure 3E, MPRA elements track),
which occurred in active TSS and enhancer states in islets and overlapped an islet ATAC-seq peak. All three
elements showed significant transcriptional activity in our assay (Z score > 2.94, P values < 0.001). Interestingly,
while there are no known gene TSS annotations in this region, clear islet polyA+ mRNA-seq profiles overlapping
the CAGE signal can be observed here. The active elements overlapped more TF motifs with positive lasso
regression coefficients (Figure 3E, positive lasso motifs track) compared to TF motifs with negative lasso
regression coefficients (Figure 3E, negative lasso motifs track).

Through these analyses we experimentally validated ~68% of testable CAGE elements for significant
transcriptional regulatory activity in a rodent islet beta cell model system, and identify TF motifs associated with
high transcriptional activities.

259
CAGE profiles augment functional genomic annotations to better fine-map GWAS loci
Observing the molecular signature of islet TCs in different epigenomic contexts and validating the activities of
these elements, we next asked if islet TCs taken as an additional layer of functional genomic information could
supplement our understanding of GWAS or islet expression quantitative loci (eQTL) associations. We classified
genomic annotations based on layers of epigenomic data such as a) histone modification based chromatin
states, b) accessible regions within the chromatin states and c) transcribed and accessible regions within the
chromatin states. We then computed enrichment for T2D GWAS loci (Mahajan *et al.* 2018) to overlap these
annotations using a Bayesian hierarchical model implemented in the fGWAS tool (Pickrell 2014). This method
utilizes not only the genome wide significant loci but leverages full genome wide association summary statistics
such that marginal associations can also be accounted for. We observed that TCs in accessible enhancer regions
were the most highly enriched for T2D GWAS signals among the annotations tested (Figure 4A, left). We also
computed enrichment for annotations to overlap islet eQTL (Varshney *et al.* 2017) and observed that TCs in
accessible regions in both enhancers and promoters were the most highly enriched (Figure 4A, right). These
data suggest that including TC information with other functional genomics data help illuminate relevant regions
for the genetic control of gene expression and trait association signals.

We next asked to what extent TCs or ATAC-seq annotations add information above and beyond chromatin state
annotations in the GWAS and eQTL enrichment models. We performed conditional analyses in fGWAS where,
first, the enrichment parameters for active TSS or active enhancer chromatin states were modeled and fixed to
their maximum likelihood values. Second, an additional parameter for either TCs or ATAC-seq peaks was
estimated. We then asked if the added enrichment parameter was significant (confidence intervals above zero).
We observed that TCs had a higher conditional enrichment over enhancer states for T2D (Figure 4B) compared
to ATAC-seq peaks. TCs also showed a higher conditional enrichment over enhancer and promoter states for
islet eQTL loci compared to ATAC-seq peaks (Figure 4B).

We then sought to utilize this new set of islet TC functional annotations to fine-map GWAS loci and reweight the
SNP posterior probabilities of association (PPAs). We performed functional reweighting of T2D GWAS (BMI-
unadjusted) using annotations such as chromatin states, ATAC-seq peaks and TCs with fGWAS. We observed
that including functional annotations resulted in higher maximal SNP PPA at a locus when compared to maximal
SNP PPAs from genetic fine mapping alone in many cases (Supplementary Figure 7), consistent with other
studies (Pickrell 2014; Mahajan *et al.* 2018). We then compared functional reweighting when using chromatin
states and ATAC-seq peaks without or with TCs and observed that some loci achieve a higher maximal
reweighted SNP PPA when TC annotations are included in the model, suggesting that TCs add valuable
information in fine-mapping (Figure 4C). We highlight one such GWAS locus named LCORL (lead SNP
rs12640250, p value = 3.7×10^{-8}). The 99% genetic credible set at this locus includes 74 variants (Mahajan *et al.*
2018), where the lead SNP rs12640250 obtains the maximum PPA = 0.15 (Supplementary Figure 8A). This
GWAS signal co-localizes with an eQTL signal in islets for the gene *DCAF16* (lead eQTL SNP rs2074974, eQTL
beta distribution-adjusted p values = 7.7×10^{-4}, COLOC (Giambartolomei *et al.* 2014) q value = 5.44×10^{-4}) (Viñuela
et al. 2019). Functional reweighting using Islet chromatin states, ATAC-seq peaks and TCs resulted in 44 SNPs
in the 99% credible set, where rs7667864 (genetic PPA = 0.12, LD r_2 0.97 with the lead GWAS SNP) obtained
the maximum reweighted PPA = 0.62 (Supplementary Figure 8B). This SNP overlaps an ATAC-seq peak and
a TC in islets (Figure 4D, E). The eQTL lead SNP rs2074974 (genetic PPA = 0.026, LD r2 = 0.96 with lead
GWAS SNP) occurs upstream of the TC and overlaps the ATAC-seq peak, and obtains a reweighted PPA =
0.096 (Figure 4E). An element overlapping this TC showed significant activity in our MPRA assay (Z score =

304 18.48, p value = 1.56x10^{-76}) and several TF motifs that showed positive MPRA lasso regression coefficients also
305 occur in this region (Figure 4D). These analyses demonstrate that transcription initiation sites demarcate active
306 regulatory elements in islets and this information can be useful in fine mapping and prioritizing variants.
307
Discussion
We profiled TSSs in human pancreatic islets using CAGE. Islet TCs were enriched to occur in promoter
chromatin states and ATAC-seq peaks, which expectedly reflects the chromatin landscape at regions where
transcription initiation occurs. Comparison of islet CAGE TCs with those identified across 118 diverse tissues
revealed that 20% of islet TCs were islet-specific. The islet-specific TCs occurred more distal to known protein-
coding gene TSSs. Also, the chromatin landscape underlying gene TSS-distal TCs was comprised of more islet-
specific promoter and enhancer chromatin states. Our analyses highlighted the differences in the chromatin
architecture underlying islet TC in islets compared to whole pancreas tissues. Islet TCs were enriched to overlap
GWAS loci of fasting glucose and specifically the islet beta cell related components of T2D GWAS signals, while
being depleted for the insulin resistance related components of T2D GWAS signals. These analyses
demonstrate that islet TCs mark active, specific and relevant islet regulatory elements.
319
Surveying the TF footprint motifs occurring in TCs revealed that several ETS family footprint motifs were highly
enriched in transcribed and accessible promoter regions, and that these motifs were also strong predictors of
the elements' activity in the MPRA assay. The regulatory potential of ETS family motifs has been described in
the literature. One study demonstrated that orienting for islet eQTL SNPs occurring in ETS footprint motifs, the
base preferred in the motifs was significantly more often associated with increased expression of the target gene
(Viñuela *et al.* 2019). Another study utilizing an MPRA assay with tiled sequences in HepG2 and K562 cell lines
also observed high regulatory activities of ETS motifs (Ernst *et al.* 2016). Our concordant findings with these
studies using orthogonal datasets highlights the robustness and utility of our islet TC atlas. Our work also
revealed that transcribed and accessible enhancer regions were most enriched to overlap TF footprint motifs for
the RFX family of TFs. We previously showed that RFX footprint motifs are confluently disrupted by T2D GWAS
risk alleles (Varshney *et al.* 2017), which are enriched to occur in islet-specific enhancer regions. These
observations together highlight the role of islet specific enhancer regions, and the potential of ATAC-seq and
CAGE profiling to pinpoint the active regulatory nucleotides within enhancer regions.
333
Utilizing an enhancer MPRA approach, we observed 2,279 CAGE elements showed significant enhancer activity
which represents ~68% of the testable elements and again highlights how CAGE profiling identifies active
regulatory elements. A larger proportion of CAGE elements that occurred in promoter chromatin states showed
significant enhancer activity and higher activity Z scores than CAGE elements occurring in active enhancer or
other chromatin states. It is interesting to note that our MPRA platform where the tested element is cloned
downstream of the GFP gene is an 'enhancer' reporter assay. While MPRA vectors are episomal and do not
recapitulate the underlying chromatin context, our results nevertheless show that sequences associated with
native promoter chromatin state landscapes can show strong enhancer activity when cloned downstream of a
reporter gene. We note that only a small fraction of CAGE TCs identified in our work (0.4%) overlapped with the
active enhancer chromatin state. Studies have shown that gene distal transcripts are more unstable, which would
therefore be difficult to profile from a total RNA sample. Of course, given the relative instability of enhancer
RNAs, enhancer chromatin sites may be actively transcribed but fall below the limits of detection of CAGE.
Therefore, it is plausible that islet CAGE profiling from total RNA, as we have performed, would comprise more
stable promoter-associated RNA transcripts and have a lesser representation of weaker transcripts originating

from enhancer regions. Recent new technologies such as native elongating transcript CAGE (NET-CAGE) also show promise in more efficiently identifying more unstable transcripts from fixed tissues (Hirabayashi *et al.* 2019). In our previous work (Varshney *et al.* 2019), we showed that genetic variants in more cell type-specific enhancer regions have lower effects on gene expression than the variants occurring in more ubiquitous promoter regions. This finding is consistent with our observation that enhancer chromatin state regions comprise a smaller proportion of active transcription initiation sites and lower transcriptional activities relative to promoter chromatin state regions. The transcription initiation landscape could also change from the basal cell state under stimulatory conditions where relevant enhancers help orchestrate a cellular response.

Integrating CAGE TC information with GWAS and eQTL data revealed the potential of the CAGE dataset to better understand the mechanisms underlying these associations. We reasoned that if CAGE TCs represent active sites within regulatory elements and are functional for gene regulation and disease, T2D GWAS and islet eQTL variants would be enriched at these sites and SNPs occurring at these sites would more likely be causal. Indeed, regions supported by TCs, ATAC-seq peaks and enhancer chromatin states (transcribed, accessible enhancer regions) were most enriched to overlap T2D GWAS loci. This enrichment was higher than in regions only informed by ATAC-seq peaks and enhancer chromatin states, indicating that the small set of TCs in enhancer regions delineate highly relevant elements. Our work demonstrates that transcription initiation information profiled using CAGE in islets can be used in addition to other relevant epigenomic information such as histone mark informed chromatin states and chromatin accessibility in nominating causal variants and biological mechanisms.

Data Availability

Raw and processed Islet CAGE data has been submitted to dBGaP (accession no. phs001188.v1.p1), raw and processed MPRA data has been submitted to GEO (GSE137693). A UCSC genome browser track hub with Islet CAGE tag clusters and tracks, chromatin states, mRNA-seq, ATAC-seq is available at https://theparkerlab.med.umich.edu/gb/custom-tracks/2019_CAGE/hub.txt A UCSC browser session is also available at http://genome.ucsc.edu/s/arushiv/cage_2019. Code to run analyses presented in the paper is shared on GitHub (https://github.com/ParkerLab/islet_cage) and the accompanying data files are shared on Zenodo (https://zenodo.org/record/3524578).

Competing interests

The authors declare no competing interests.

Materials and Methods

Islet Procurement and Processing

Islet samples from organ donors were received from the Integrated Islet Distribution Program, the National Disease Research Interchange (NDRI), and Prodo-Labs. Islets were shipped overnight from the distribution centers. Upon receipt, we prewarmed islets to 37 °C in shipping media for 1–2 h before harvest; ~2,500–5,000 islet equivalents (IEQs) from each organ donor were harvested for RNA isolation. We transferred 500–1,000 IEQs to tissue culture-treated flasks and cultured them as in the work in (Gershengorn *et al.* 2004).

RNA isolation, CAGE-seq library preparation and sequencing

Total RNA from 2000-3000 islet equivalents (IEQ) was extracted and purified using Trizol (Life Technologies). RNA quality was confirmed with Bioanalyzer 2100 (Agilent); samples with RNA integrity number (RIN) > 6.5 were prepared for CAGE sequencing. 1ug Total RNA samples were sent to DNAFORM, Japan where CAGE libraries were generated. The library preparation included polyA negative selection and size selection (<1000bp). Stranded CAGE libraries were generated for each islet sample using the no-amplification non-tagging CAGE libraries for Illumina next-generation sequencers (nAnT-iCAGE) protocol (Murata *et al.* 2014). Each islet CAGE library was barcoded, pooled into 24-sample batches, and sequenced over multiple lanes of HiSeq 2000 to obtain paired-end 126 bp sequences. All procedures followed ethical guidelines at the National Institutes of Health (NIH).

CAGE data mapping and processing

We processed Islet CAGE data uniformly with CAGE data for other tissues included in separate ongoing projects. Because read lengths differed across libraries, we trimmed all reads to 51 bp using fastx_trimmer (FASTX Toolkit v. 0.0.14). Adapters and technical sequences were trimmed using trimmomatic (v. 0.38; paired-end mode, with options ILLUMINACLIP:adapters.fa:1:30:7:1:true). To remove potential *E. coli* contamination, we mapped to the *E. coli* chromosome (genome assembly GCA_000005845.2) with bwa mem (v. 0.7.15; options: -M). We then removed read pairs that mapped in a proper pair (with mapq >= 10) to *E. coli.* We mapped the remaining reads to hg19 using STAR (v. 2.5.4b; default parameters) (Dobin *et al.* 2013). We pruned the mapped reads to high quality autosomal read pairs (using samtools view v. 1.3.1; options -f 3 -F 4 -F 8 -F 256 -F 2048 -q 255)(Li *et al.* 2009). We then performed UMI-based deduplication using umitools dedup (v. 0.5.5; --method directional).

We selected 57 Islet samples with strandedness measures >0.85 calculated from QoRTS (Hartley and Mullikin 2015) for all downstream analyses.

Tag cluster identification

We used the paralu method to identify clusters of CAGE start sites (CAGE tag clusters) (Frith *et al.* 2008). The algorithm uses a density parameter d and identifies segments that maximize the value of (Number of events - d * size of the segment (bp)). Here, large values of d would favor small, dense clusters and small values of d would favor larger more rarefied clusters. The method identifies segments over all values of d beginning at the largest scale, where d = 0, where all of the events are merged into one big cluster. It then calculates the density (events per nucleotide) of every prefix and suffix of the big cluster. The lowest value among all of these densities is the maximum value of d for the big cluster because at higher values of d the big cluster will no longer be a maximal-scoring segment (because zero-scoring prefixes or suffixes are not allowed).

We called TCs in each individual sample using raw tag counts, requiring at least 2 tags at each included start site and allowing single base-pair tag clusters ('singletons') if supported by >2 tags. We then merged the tag clusters on each strand across samples. For each resulting segment, we calculated the number of islet samples in which TCs overlapped the segment. We included the segment in the consensus TCs set if it was supported by independent TCs in at least 10 individual islet samples. This threshold was selected based on comparing the number of tag clusters with the number of samples across which support was required to consider the segment (Figure 1 - figure supplment 1). We then filtered out regions blacklisted by the ENCODE consortium due to poor

433 mappability (wgEncodeDacMapabilityConsensusExcludable.bed and
434 wgEncodeDukeMapabilityRegionsExcludable.bed) using bedtools subtract to obtain the final set of Islet tag
435 cluster regions used in all downstream analyses.
436
FANTOM CAGE datasets
We obtained the set of 'robust CAGE peaks' identified by the FANTOM 5 consortium (The FANTOM Consortium
et al. 2014) using CAGE libraries (CAGE sequencing on HeliScope Single Molecule Sequencer (hCAGE))
of 988 human cell lines or tissues
([http://fantom.gsc.riken.jp/5/datafiles/latest/extra/CAGE_peaks/hg19.cage_peak_phase1and2combined_coord.](http://fantom.gsc.riken.jp/5/datafiles/latest/extra/CAGE_peaks/hg19.cage_peak_phase1and2combined_coord.bed.gz)
[bed.gz](http://fantom.gsc.riken.jp/5/datafiles/latest/extra/CAGE_peaks/hg19.cage_peak_phase1and2combined_coord.bed.gz)). These peaks were identified using the decomposition-based peak identification (DPI) method (The
FANTOM Consortium *et al.* 2014), followed by filtering 'robust' peaks that included a CAGE tag (TSS) with more
than 10 read counts in at least 1 sample and 1 tags per million (TPM). For a more direct comparison of islet TCs
with TCs from other tissues, we downloaded the CAGE transcription start site (CTSS) data for 118 tissue types
(from http://fantom.gsc.riken.jp/5/datafiles/latest/basic/human.tissue.hCAGE/) and called tag clusters for each
tissue sample using the paraclu method (Frith *et al.* 2008) as described above, with the same parameters.
448
Chromatin state analysis
We collected publicly available cell/tissue ChIP-seq data for H3K27ac, H3K27me3, H3K36me3, H3K4me1, and
H3K4me3 and input for Islets, Adipose and Skeletal Muscle (Supplementary Table 6). Data for Adipose, Skeletal
Muscle and Liver tissues were included in the joint model for other ongoing projects. We performed read mapping
and integrative chromatin-state analyses in a manner similar to that of our previous reports and followed quality
control procedures reported by the Roadmap Epigenomics Study (The Roadmap Epigenomics Consortium *et al.*
2015). Briefly, we trimmed reads across datasets to 36bp and overrepresented adapter sequences as shown by
FASTQC (version v0.11.5) using cutadapt (version 1.12) (Martin 2011). We then mapped reads using BWA
(version 0.5.8c), removed duplicates using samtools (Li *et al.* 2009), and filtered for mapping quality score of at
least 30. To assess the quality of each dataset, we performed strand cross-correlation analysis using
phantompeakqualtools (v2.0; code.google.com/p/phantompeakqualtools) (Landt *et al.* 2012). We converted bam
files for each dataset to bed using the bamToBed tool. To more uniformly represent datasets with different
sequencing depths across histone marks and tissues, we randomly subsampled each dataset bed file to the
mean depth for that mark across the four included tissues. This allowed comparable chromatin state territories
across tissues and ensured that chromatin state territories were not heavily driven by high sequencing depth.
Chromatin states were learned jointly for the three cell types using the ChromHMM (version 1.10) hidden Markov
model algorithm at 200-bp resolution to five chromatin marks and input (Ernst and Kellis 2010, 2012; Ernst *et al.*
2011). We ran ChromHMM with a range of possible states and selected a 11-state model, because it most
accurately captured information from higher-state models and provided sufficient resolution to identify biologically
meaningful patterns in a reproducible way. We have used this state selection procedure in previous analyses
(Scott *et al.* 2016; Varshney *et al.* 2017). To assign biological function names to our states that are consistent
with previously published states, we performed enrichment analyses in ChromHMM comparing our states with
the states reported previously (Varshney *et al.* 2017) for the four matched tissues. We assigned the name of the
state that was most strongly enriched in each of our states (Supplementary Figure 3).
473
ATAC-seq data analysis
We used previously published chromatin accessibility data profiled using ATAC-seq in islets from two human
organ donor samples (Varshney *et al.* 2017). For each sample, we trimmed reads to 36 bp (to uniformly process

ATAC-seq from other tissues for ongoing projects) and removed adapter sequences using Cutadapt (version
1.12) (Martin 2011), mapped to hg19 used bwa-mem (version 0.7.15-r1140) (Li 2013), removed duplicates using
Picard (http://broadinstitute.github.io/*picard*) and filtered out regions blacklisted by the ENCODE consortium due
to poor mappability (wgEncodeDacMapabilityConsensusExcludable.bed and
wgEncodeDukeMapabilityRegionsExcludable.bed). For each tissue we subsampled both samples to the same
depth (18M reads for Islet samples) so that each tissue had overall similar genomic region called as peaks. We
used MACS2 (https://github.com/taoliu/MACS), version 2.1.0, with flags "-g hs–nomodel–shift -100–extsize 200
-B–broad–keep-dup all," to call peaks and retained all broad-peaks that satisfied a 1% FDR.
485
Overlap enrichment between TCs and annotations
We calculated the enrichment for Islet TCs to overlap annotations such as different Islet chromatin states, Islet
ATAC-seq peaks and various 'common' annotations. Common annotations imply annotations that don't vary
across cell types such as coding gene regions, intronic regions or annotations created by merging epigenomic
data such as histone modification peaks across cell types. We utilized 29 total static annotation bed files supplied
by (Finucane *et al.* 2015) (https://data.broadinstitute.org/alkesgroup/LDSCORE/baseline_bedfiles.tgz). These
included coding, un-translated regions (UTRs), promoter and intronic regions obtained from UCSC (Kent *et al.*
2002); the histone marks monomethylation (H3K4me1) and trimethylation (H3K4me3) of histone H3 at lysine 4
and acetylation of histone H3 at lysine 9 (H3K9ac) (The ENCODE project Consortium 2012; Trynka *et al.* 2013;
The Roadmap Epigenomics Consortium *et al.* 2015) and acetylation of histone H3 at lysine 27 (H3K27ac) (Hnisz
et al. 2013; Schizophrenia Working Group of the Psychiatric Genomics Consortium 2014); open chromatin, as
reflected by DNase I hypersensitivity sites (DHSs) (Trynka *et al.* 2013; Gusev *et al.* 2014); combined chromHMM
and Segway predictions(Hoffman *et al.* 2013), which partition the genome based on distinct and recurring
patterns of histone marks into seven underlying chromatin states; regions that are conserved in
mammals(Lindblad-Toh *et al.* 2011; Ward and Kellis 2012); super-enhancers, which are large clusters of highly
active enhancers(Hnisz *et al.* 2013); and enhancers with balanced bidirectional capped transcripts identified
using CAGE in the FANTOM5 panel of samples, (called Enhancer (Andersson)) (Andersson *et al.* 2014). Histone
marks included in the static annotation set included merged histone mark data from different cell types into a
single annotation.
505
Enrichment for overlap between each Islet tag clusters and regulatory annotations was calculated using the
Genomic Association Tester (GAT) tool (Heger *et al.* 2013). To ask if two sets of regulatory annotations overlap
more than that expected by chance, GAT randomly samples segments of one regulatory annotation set from the
genomic workspace (hg19 chromosomes) and computes the expected overlaps with the second regulatory
annotation set. We used 10,000 GAT samplings for each enrichment run. GAT outputs the observed overlap
between segments and annotation along with the expected overlap and an empirical p-value.
512
Enrichment of GWAS loci in TCs
We downloaded the GWAS data for various traits from the NHGRI website on June 12, 2018 (file
gwas_catalog_v1.0.2-associations_e92_r2018-05-29.tsv from https://www.ebi.ac.uk/gwas/docs/file-
downloads). We selected genome-wide significant GWAS SNPs (P <5x10-8) for traits for which the study included
European samples. To retain independent signals, we LD pruned the list of SNPs to retain SNPs with the most
significant P values that had LD r2 <0.2 between each pair. This procedure was performed using the PLINK
(v1.9) tool (Purcell *et al.* 2007; Chang *et al.* 2015) –clump option and 1000 genomes phase 3 vcf files
(downloaded from ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/release/20130502), subsetted to the European

samples as reference. We selected traits that had >30 independent signals for following analyses. We also
downloaded data from (Udler *et al.* 2018) which included clusters of T2D GWAS loci such as beta cell function,
insulin resistance etc. This study clustered T2D GWAS loci by analyzing GWAS for T2D along with 47 other
diabetes related traits.
525
Enrichment for GWAS variants for different traits in Islet TCs was calculated using the Genomic Regulatory
Elements and Gwas Overlap algoRithm (GREGOR) tool (version 1.2.1) (Schmidt *et al.* 2015). Since the causal
SNP(s) for the traits are not known, GREGOR allows considering the input lead SNP along with SNPs in high
LD (based on the provided R2THRESHOLD parameter) while computing overlaps with genomic features (such
as islet TCs). Therefore, as input to GREGOR, we supplied SNPs that were not in high LD with each other. For
each input SNP, GREGOR selects ~500 control SNPs that match the input SNP for minor allele frequency (MAF),
distance to the nearest gene, and number of SNPs in LD. Fold enrichment is calculated as the number of loci at
which an input SNP (either lead SNP or SNP in high LD) overlaps the feature over the mean number of loci at
which the matched control SNPs (or SNPs in high LD) overlap the same features. This process accounts for the
length of the features, as longer features will have more overlap by chance with control SNP sets. Specific
parameters for the GWAS enrichment were: GREGOR: r2 threshold = 0.8. LD window size = 1Mb; minimum
neighbor number = 500, population = European.
538
539 **Aggregate signal**
540 We generated the ATAC-seq density plot over islet TC midpoints using the Agplus tool (version 1.0) (Maehara
541 and Ohkawa 2015). We used the ATAC-seq signal track for reads per 10 Million to aggregate over stranded
542 TCs.
543
544 To obtain CAGE tracks, we merged CAGE bam files for islet samples that passed QC (see CAGE data
545 processing section) and obtained the read 1 start sites or TSSs. To better visualise the CAGE signal, we then
546 flanked each TSS 10bp upstream and downstream and normalized the TSS counts to 10M mapped reads. We
547 generated CAGE density plots over ATAC-seq narrow peak summits by using the agplus tool.
548
549 To obtain aggregate CAGE signal over TF footprint motifs, we oriented the CAGE signal with respect to the
550 footprint taken on the plus strand. We used HTSeq GenomicPosition method (Anders *et al.* 2015) to obtain the
551 sum of CAGE signal at each base pair relative to the footprint motif mid point.
552
Enrichment for Islet TF footprint motifs to overlap TC-related annotations
We compared the enrichment of Islet TF footprint motifs in several TC-related annotations such as upstream
and downstream 500bp regions of TCs and Islet TCs that occurred in accessible enhancer states vs those that
occurred in accessible promoter states using the GAT tool similarly as described above (Heger *et al.* 2013). TF
footprint motifs are occurrences of TFs motifs (obtained from databases of DNA binding motifs for several TFs)
in accessible chromatin regions (identified from assays such as ATAC-seq). We utilized previously published
islet TF footprint motifs (Varshney *et al.* 2017), which were generated using ATAC-seq data in two islet samples
and DNA binding motif information for 1,995 publicly available TF motifs (Jolma *et al.* 2013; Kheradpour and
Kellis 2014; Mathelier *et al.* 2016).
562
We obtained the 500bp upstream or downstream regions of each TC (upstream/downstream regions were
determined based on TC strand; TC region itself was not included). We generated the list of regions where islet

TCs overlapped ATAC-seq peaks and any enhancer states (Active enhancer, Weak enhancer, or Genic
enhancer) using BEDTools intersect - we referred to these as 'TCs in accessible enhancers'. Similarly, we also
generated the list of regions where islet TCs overlapped ATAC-seq peaks and any TSS/promoter states (Active
TSS, Weak TSS, Flanking TSS) - we referred to these as 'TCs in accessible promoters'. as segments.
569
In the GAT analyses, the TC-related annotations were considered as 'annotations' (argument -a) and footprint
motif occurrences for each known motif were considered as 'segments' (argument -s). Since the TF footprint
motifs only occur in ATAC-seq peaks, we considered these peaks as the 'workspace' (argument -w) to sample
segments. We used 10,000 GAT samplings for each enrichment run. We accounted for the 1,995 footprint motifs
being tested against each TC-related annotation by performing an FDR correction with the Benjamini-Yekutieli
method (Benjamini and Yekutieli 2001) using the stats.multitest.multipletests function from the statsmodels
library in Python (Seabold and Perktold 2010). Significant enrichment was considered at 5% FDR threshold.
577

Comparison of features with Roadmap chromatin states
We downloaded the chromatin state annotations identified in 127 human cell types and tissues by the Roadmap
epigenomics project (The Roadmap Epigenomics Consortium *et al.* 2015) after integrating ChIP-seq data for five
histone 3 lysine modifications (H3K4me3, H3K4me1, H3K36me3, H3K9me3 and H3K27me3) that are associated
with promoter, enhancer, transcribed and repressed activities, across each cell type. For each TC feature, for
example, TCs in ATAC-seq peaks within islet enhancer chromatin states, we identified segments occurring
proximal to (within 5kb) and distal from (further than 5kb) known protein coding gene TSS (gencode V19). For
each such segment, we identified the maximally overlapping chromatin state across 98 cell types publicly
available from the Roadmap Epigenomics project in their 18 state 'extended' model using BEDtools intersect.
We then ordered the segments using clustering (hclust function in R) based on the gower distance metric (daisy
function in R) for the roadmap state assignments across 127 cell types.
589

Experimental validation using MPRA
1.   **Selection of CAGE elements**
We generated a library of islet CAGE elements to test in the MPRA assay by using two approaches. First, we
identified clusters of CAGE tags in each islet sample by simply concatenating tags that occurred within 20bp.
We retained clusters with at least two tags in each islet sample. We then merged these cluster coordinates
across samples and retained clusters supported by at least 15 samples, representing a highly reproducible set.
Second, we complemented this approach by also including the set of FANTOM 5 'robust' CAGE peaks that were
also supported by CAGE tags in at least 15 samples. 94% of the selected CAGE robust peak regions were
already included in the selected CAGE 20 bp clusters; we reasoned that the remaining 6% of CAGE peaks
represented relevant and reproducible CAGE elements missed by the 20 bp concatenation approach. We
therefore took the union of these two sets of CAGE elements and created 198 bp oligo sequences centered on
each element for cloning into the MPRA vector. When a CAGE element was longer than 198 bp, we tiled 198 bp
oligos over the element, offset by 100 bp. Through this approach, we included a total of 7,188 CAGE elements
(each 198 bp long). We note that these CAGE elements represent slightly different coordinates from the TCs
coordinates presented elsewhere in the paper that were identified using the paraclu method. While the paraclu
approach of calling TCs was adopted after the MPRA experiments were already performed, 6,810 (94.7%) of
the CAGE elements included in MPRA experiment overlapped the final set of TCs presented in the manuscript.
607

We synthesized 230-bp oligos (198bp CAGE element flanked by 16bp anchor sequences) (Agilent
Technologies). We PCR-amplified oligos to add homology arms for Gibson assembly cloning, and gel-purified
the PCR products (i.e. inserts) using the Zymoclean Gel DNA Recovery Kit (Zymo Research). We used the
NEBuilder HiFi DNA Assembly Kit (NEB) to assemble the purified inserts and the backbone of MPRA plasmid
(previously digested with EcoRV). After column purification of the reaction using the DNA Clean and
Concentrator-5 kit (Zymo), we transformed it into 10beta electrocompetent cells (NEB), and obtained 1.39 million
unique transformants.
615
616 We post-barcoded the library by first digesting the library with PmeI, and then by setting up Gibson assembly to
617 insert 16-bp random nucleotides ('barcodes') at the PmeI restriction site. After column purification, we
618 transformed the reaction into electrocompetent cells (NEB), and obtained 1.44 million unique transformants. We
619 prepared the library plasmid for transfection using the ZymoPURE Plasmid Maxiprep Kit (Zymo).
620

2. Electroporation, RNA isolation and cDNA synthesis
We electroporated 50 ug of barcoded MPRA library into 25 million the 832/13 rat insulinoma cell line for each
biological replicate (3 replicates), and harvested the cells twenty-four hours later. We isolated total RNA using
TRIZOL reagent (Life Technologies) following the manufacturer's protocol up to phase separation. After phase
separation, we transferred the aqueous phase of the solution to a new 1.5 mL Eppendorf tube, added 1:1 volume
of 100% ethanol, and then column-purified using the Direct-zol RNA Miniprep Kit (Zymo Research Corporation,
Irvine, CA) following the manufacturer's protocol. We further purified mRNA using Dynabeads oligo(dT) beads
(Thermo Fisher, Carlsbad, CA). We treated 2 ug of mRNA with RNase-free DNaseI (Invitrogen, Carlsbad CA) to
eliminate possible plasmid DNA contamination, and then reverse transcribed 1ug into cDNA using the
SuperScript III First-Strand Synthesis kit (Invitrogen) with a custom primer that specifically recognizes 'MPRA
transcripts' (i.e. mRNA that had been transcribed from the MPRA plasmids). The other 1ug of mRNA was used
in a enzyme-negative reaction to determine DNA/plasmid contamination in cDNA, which we did not. To eliminate
any residual plasmid contamination in cDNA samples, we treated cDNA with DpnI (NEB), and purified the
reaction using the DNA Clean and Concentrator-5 kit (Zymo).
635

3. Construction of Illumina sequencing library
We constructed Illumina sequencing library via two serial rounds of PCR. In the first round, we used a primer set
to specifically amplify STARR transcripts using cDNA as starting material. In the second round, we used a primer
set to append the P5/P7 Illumina sequences using the PCR product from the first round as starting material. In
both rounds, we PCR-amplified the fragments until the amplification curve reached a mid-log phase, and then
purified the products for subsequent steps using the DNA Clean and Concentrator-5 kit (Zymo).
642

4. **CAGE element-barcode pairing**
To identify the barcodes corresponding to each CAGE element in the MPRA plasmid, 1ng of each library
constructed was used in a polymerase chain reaction with primers flanking the allele and the barcode to generate
fragments which were subsequently gel verified and extracted using Zymo gel extraction kit (Zymo). 25ng of the
purified product was subjected to self-ligation at 16 °C overnight in a total volume of 50uls and column purified
using Qiagen (Qiagen) PCR purification kit as per manufacturers recommendations. The purified fragments were
subsequently treated with 10U of Plasmid-Safe ATP-Dependent DNase for 1 hour in the presence of 25mM ATP
to remove unligated linear DNA fragments. 1ul of the recircularized fragments were subjected to another round
of PCR resulting in a smaller fragment. Briefly an aliquot of this product was diluted 1:10 in DNAse/RNase free

water and amplified in a PCR reaction, with Illumina P5 and P7 adapters, until saturation to generate libraries. The libraries were subsequently column purified, quantified and sequenced.

5. Data analysis
The MPRA barcode sequencing data included the input DNA barcode library along with three cDNA barcode libraries representing three biological replicated. We processed this data through a custom pipeline which quantified barcode counts while accounting for sequencing errors. We extracted the DNA barcodes from the input DNA library (first 16 bp of the read-1 fastq file) and clustered these at an edit distance of 0 followed by computing the DNA counts for each read group of DNA sequencing files. We then aggregated the read groups and collapsed counts for barcodes using the sequence clustering algorithm Starcode (https://github.com/gui11aume/starcode) (Zorita *et al.* 2015). We repeated this process for the cDNA barcode counts for each replicate, with the added step of removing PCR-duplicated barcodes using the UMI information (UMI sequence was the reverse complement of the first 6bp of the read-2 fastq file). The pipeline is shared at https://github.com/ParkerLab/STARR-seq-Analysis-Pipeline.

We matched the barcodes with CAGE inserts using results from CAGE insert-barcode pairing experiment (data file "cage_insert_barcode_pairing.tsv" in GEO GSE137693). We first retained barcodes with at least 10 DNA counts, and further retained CAGE elements that had at least two such qualifying barcodes. This was the set of N=3,446 CAGE elements quantifiable in our assay. To quantify MPRA activities from these count-based data, we used the tool MPRAnalyze (version 1.3.1) (https://github.com/YosefLab/MPRAnalyze) (Ashuach *et al.* 2019) that models DNA and RNA counts in a negative binomial generalized linear model. This approach is more robust than using metrics such as the aggregated ratio, which is the ratio of the sum of RNA counts across barcodes divided by the sum of DNA counts across barcodes and loses the statistical power provided by multiple barcodes per tested element; and the mean ratio, which is the mean of the observed RNA/DNA ratios across barcodes which can be quite sensitive to low counts and noise. We corrected for library depth for the three replicates using upper quartile normalization via the 'estimateDepthFactors' function in MPRAnalyze. MPRA activity is quantified by estimating the transcription rate for each element in the dataset, followed by identifying active elements that induce a higher transcription by testing against a null. MPRAnalyze fits two nested generalized linear models - the DNA model estimates plasmid copy numbers, and the RNA model is estimates transcription rate. We included barcode information in the DNA model which allows different estimated counts for each barcode, and increases the statistical power of the model. Replicate information was included in the RNA model. MPRAnalyze then tests the transcriptional activity of each element against a null distribution and computing Z and Median-Absolute-Deviation (MAD) scores. The null is based on the assumption that the mode of the distribution of transcription rate estimates is the center of the null distribution, and that values lower than the mode all belong to the null. Thus, values lower than the mode are used to estimate the variance of the null.

Lasso regression
We used lasso regression to model TC element MPRA activity z scores as a function of TF motif occurrences within the TC elements. Lasso regression is useful when a large number of features such as hundreds of TF motifs in this case are included because it imposes a constraint on the model parameters causing regression coefficients for some variables to shrink toward zero. Features with non-zero regression coefficients are most strongly associated with the response variable.

We utilized a set of 1,995 TF motifs including their position weight matrices (PWMs), available from ENCODE,
JASPAR and Jolma datasets (Jolma *et al.* 2013; Kheradpour and Kellis 2014; Mathelier *et al.* 2016), which we
have also used previously (Scott *et al.* 2016; Varshney *et al.* 2017). In order to reduce motif redundancy, we
performed PWM clustering in our motif database using the matrix-clustering tool from RSAT (Castro-Mondragon
et al. 2017), with parameters -lth cor 0.7 -lth Ncor 0.7. For each of the 540 clusters obtained, we retained the
motif with the highest total PWM information content. Because MPRA is an episomal assay and doesn't
recapitulate the native chromatin context, we quantified overlaps of each TC element with sequence motif scans
rather than ATAC-seq informed footprint motifs. We scanned each of these motifs on the hg19 reference using
FIMO (Grant *et al.* 2011). We used the nucleotide frequencies from the hg19 reference and the default p value
cutoff of 10_{-4}.
705
To quantify motif occurrences within each TC element, we considered the -log10(P value) of each motif
occurrence from FIMO. Since the FIMO motif scan p-values depend on the motif length and information content
etc., these log transformed P values are not directly comparable across motifs. We therefore inverse normalized
the -log10(P values) for occurrences of each motif using the RNOmni package (version 0.7.1) to obtain motif
'scores' on a comparable normal scale. P value = 1 was included for each motif to obtain the score corresponding
to no motif occurrence on the transformed scale. For each TF motif, we aligned the hg19 scan occurrences with
each islet TC MPRA element using BedTools intersect and recorded the corresponding motif scores. We added
the scores for TC elements that overlapped multiple occurrences for the said motif. We again inverse normalized
the motif overlap score vector across the input CAGE elements for each TF motif so that the regression
coefficients could be comparable across motifs. The lasso regression was run using the glmnet package (version
2.0-16) with default parameters (specifically, alpha=1, which corresponds to the lasso regression). Lambda was
determined automatically by glmnet as the lambda that belonged to the model with the lowest mean cross
validated error.
719
fGWAS analyses and fine-mapping
We used the fGWAS (version 0.3.6) (Pickrell 2014) tool to compute enrichment of GWAS and islet eQTL data in
TC-related annotations along with computing conditional enrichment and fine mapping analyses. fGWAS
employs a Bayesian hierarchical model to determine shared properties of loci affecting a trait. The model uses
association summary level data, divides the genome into windows generally larger than the expected LD patterns
in the population. The method assumes that there is either a single causal SNP in a window or none. The model
defines the prior probabilities that an association lies in a genomic window and that a SNP within it is causal.
These probabilities are allowed to depend on genomic annotations, and are estimated based on enrichment
patterns of annotations across the genome using a Bayes approach.
729
We obtained publicly available summary data for T2D GWAS (Mahajan *et al.* 2018) and islet eQTL (Varshney *et*
al. 2017) and organized it in the format required by fGWAS. We used fGWAS with default parameters for
enrichment analyses for individual annotations in Figure 4 A. For each individual annotation, the model provided
the maximum likelihood enrichment parameter. Annotations were considered as significantly enriched if the
log2(parameter estimate) and respective 95% confidence intervals were above zero or significantly depleted if
the log2(parameter estimate) and respective 95% confidence intervals were below zero. We performed
conditional analyses using the '-cond' option.
737

To reweight GWAS summary data based on functional annotation overlap, we used the '-print' option in addition
in fGWAS while including multiple annotations in the model that were individually significantly enriched or
depleted. We included active TSS, active enhancer, quiescent and polycomb repressed annotations and ATAC-
seq peaks with or without or TCs. To compare with T2D genetic credible sets (Mahajan *et al.* 2018), we created
1Mb windows centered on the lead variant at each primary GWAS signal. We then partitioned the rest of the
genome into 1Mb windows as well. Specifying these regions using the --bed option in fGWAS enabled
constraining each primary signal in a single window. The model derived enrichment priors to evaluate both the
significance and functional impact of associated variants in GWAS regions, such that variants overlapping more
enriched annotations carried extra weight.
747

References
Adli M., J. Zhu, and B. E. Bernstein, 2010 Genome-wide chromatin maps derived from limited numbers of

hematopoietic progenitors. Nat. Methods 7: 615–618. https://doi.org/10.1038/nmeth.1478

Anders S., P. T. Pyl, and W. Huber, 2015 HTSeq--a Python framework to work with high-throughput

sequencing data. Bioinforma. Oxf. Engl. 31: 166–169. https://doi.org/10.1093/bioinformatics/btu638

Andersson R., C. Gebhard, I. Miguel-Escalada, I. Hoof, J. Bornholdt, *et al.*, 2014 An atlas of active enhancers

across human cell types and tissues. Nature 507: 455. https://doi.org/10.1038/nature12787

Arnold C. D., D. Gerlach, C. Stelzer, Ł. M. Boryń, M. Rath, *et al.*, 2013 Genome-Wide Quantitative Enhancer

Activity Maps Identified by STARR-seq. Science 339: 1074–1077.

https://doi.org/10.1126/science.1232542

Ashuach T., D. S. Fischer, A. Kreimer, N. Ahituv, F. J. Theis, *et al.*, 2019 MPRAnalyze: statistical framework

for massively parallel reporter assays. Genome Biol. 20: 183. https://doi.org/10.1186/s13059-019-1787-

z

Benjamini Y., and D. Yekutieli, 2001 The control of the false discovery rate in multiple testing under

dependency. Ann. Stat. 29: 1165–1188. https://doi.org/10.1214/aos/1013699998

Buenrostro J. D., P. G. Giresi, L. C. Zaba, H. Y. Chang, and W. J. Greenleaf, 2013 Transposition of native

chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and

nucleosome position. Nat. Methods 10: 1213–1218. https://doi.org/10.1038/nmeth.2688

Buniello A., J. A. L. MacArthur, M. Cerezo, L. W. Harris, J. Hayhurst, *et al.*, 2019 The NHGRI-EBI GWAS

Catalog of published genome-wide association studies, targeted arrays and summary statistics 2019.

Nucleic Acids Res. 47: D1005–D1012. https://doi.org/10.1093/nar/gky1120

Bunt M. van de, J. E. M. Fox, X. Dai, A. Barrett, C. Grey, *et al.*, 2015 Transcript Expression Data from Human

Islets Links Regulatory Signals from Genome-Wide Association Studies for Type 2 Diabetes and

Glycemic Traits to Their Downstream Effectors. PLOS Genet. 11: e1005694.

https://doi.org/10.1371/journal.pgen.1005694

Castro-Mondragon J. A., S. Jaeger, D. Thieffry, M. Thomas-Chollier, and J. van Helden, 2017 RSAT matrix-

clustering: dynamic exploration and redundancy reduction of transcription factor binding motif

collections. Nucleic Acids Res. 45: e119–e119. https://doi.org/10.1093/nar/gkx314

Cebola I., 2019 Pancreatic Islet Transcriptional Enhancers and Diabetes. Curr. Diab. Rep. 19: 145.

https://doi.org/10.1007/s11892-019-1230-6

Chang C. C., C. C. Chow, L. C. Tellier, S. Vattikuti, S. M. Purcell, *et al.*, 2015 Second-generation PLINK: rising

to the challenge of larger and richer datasets. GigaScience 4: 1–16. https://doi.org/10.1186/s13742-

015-0047-8

Core L. J., J. J. Waterfall, and J. T. Lis, 2008 Nascent RNA Sequencing Reveals Widespread Pausing and

Divergent Initiation at Human Promoters. Science 322: 1845–1848.

https://doi.org/10.1126/science.1162228

Core L. J., A. L. Martins, C. G. Danko, C. Waters, A. Siepel, *et al.*, 2014 Analysis of nascent RNA identifies a

unified architecture of initiation regions at mammalian promoters and enhancers. Nat. Genet. 46: 1311–

1320. https://doi.org/10.1038/ng.3142

Creyghton M. P., A. W. Cheng, G. G. Welstead, T. Kooistra, B. W. Carey, *et al.*, 2010 Histone H3K27ac

separates active from poised enhancers and predicts developmental state. Proc. Natl. Acad. Sci. U. S.

A. 107: 21931–21936. https://doi.org/10.1073/pnas.1016071107

Dobin A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, *et al.*, 2013 STAR: ultrafast universal RNA-seq

aligner. Bioinforma. Oxf. Engl. 29: 15–21. https://doi.org/10.1093/bioinformatics/bts635

Ernst J., and M. Kellis, 2010 Discovery and characterization of chromatin states for systematic annotation of

the human genome. Nat. Biotechnol. 28: 817–825. https://doi.org/10.1038/nbt.1662

Ernst J., P. Kheradpour, T. S. Mikkelsen, N. Shoresh, L. D. Ward, *et al.*, 2011 Mapping and analysis of

chromatin state dynamics in nine human cell types. Nature 473: 43–49.

https://doi.org/10.1038/nature09906

Ernst J., and M. Kellis, 2012 ChromHMM: automating chromatin state discovery and characterization. Nat.

Methods 9: 215–216. https://doi.org/10.1038/nmeth.1906

Ernst J., A. Melnikov, X. Zhang, L. Wang, P. Rogov, *et al.*, 2016 Genome-scale high-resolution mapping of

activating and repressive nucleotides in regulatory regions. Nat. Biotechnol. 34: 1180–1190.

https://doi.org/10.1038/nbt.3678

Fadista J., P. Vikman, E. O. Laakso, I. G. Mollet, J. L. Esguerra, *et al.*, 2014 Global genomic and

transcriptomic analysis of human pancreatic islets reveals novel genes influencing glucose metabolism.

Proc. Natl. Acad. Sci. U. S. A. 111: 13924–13929.

Finucane H. K., B. Bulik-Sullivan, A. Gusev, G. Trynka, Y. Reshef, *et al.*, 2015 Partitioning heritability by

functional annotation using genome-wide association summary statistics. Nat. Genet. 47: 1228–1235.

https://doi.org/10.1038/ng.3404

Frith M. C., E. Valen, A. Krogh, Y. Hayashizaki, P. Carninci, *et al.*, 2008 A code for transcription initiation in

mammalian genomes. Genome Res. 18: 1–12. https://doi.org/10.1101/gr.6831208

Gershengorn M. C., A. A. Hardikar, C. Wei, E. Geras-Raaka, B. Marcus-Samuels, *et al.*, 2004 Epithelial-to-

Mesenchymal Transition Generates Proliferative Human Islet Precursor Cells. Science 306: 2261–

2264.

Giambartolomei C., D. Vukcevic, E. E. Schadt, L. Franke, A. D. Hingorani, *et al.*, 2014 Bayesian Test for

Colocalisation between Pairs of Genetic Association Studies Using Summary Statistics. PLOS Genet.

10: e1004383. https://doi.org/10.1371/journal.pgen.1004383

Grant C. E., T. L. Bailey, and W. S. Noble, 2011 FIMO: scanning for occurrences of a given motif.

Bioinformatics 27: 1017–1018. https://doi.org/10.1093/bioinformatics/btr064

Gusev A., S. H. Lee, G. Trynka, H. Finucane, B. J. Vilhjálmsson, *et al.*, 2014 Partitioning heritability of

regulatory and cell-type-specific variants across 11 common diseases. Am. J. Hum. Genet. 95: 535–

552. https://doi.org/10.1016/j.ajhg.2014.10.004

Hartley S. W., and J. C. Mullikin, 2015 QoRTs: a comprehensive toolset for quality control and data processing

of RNA-Seq experiments. BMC Bioinformatics 16: 224. https://doi.org/10.1186/s12859-015-0670-5

Heger A., C. Webber, M. Goodson, C. P. Ponting, and G. Lunter, 2013 GAT: a simulation framework for testing

the association of genomic intervals. Bioinformatics 29: 2046–2048.

https://doi.org/10.1093/bioinformatics/btt343

Hirabayashi S., S. Bhagat, Y. Matsuki, Y. Takegami, T. Uehata, *et al.*, 2019 NET-CAGE characterizes the

dynamics and topology of human transcribed cis -regulatory elements. Nat. Genet. 51: 1369–1379.

https://doi.org/10.1038/s41588-019-0485-9

Hnisz D., B. J. Abraham, T. I. Lee, A. Lau, V. Saint-André, *et al.*, 2013 Super-Enhancers in the Control of Cell

Identity and Disease. Cell 155: 934–947. https://doi.org/10.1016/j.cell.2013.09.053

Hoffman M. M., J. Ernst, S. P. Wilder, A. Kundaje, R. S. Harris, *et al.*, 2013 Integrative annotation of chromatin

elements from ENCODE data. Nucleic Acids Res. 41: 827–841. https://doi.org/10.1093/nar/gks1284

Hsieh C.-L., T. Fei, Y. Chen, T. Li, Y. Gao, *et al.*, 2014 Enhancer RNAs participate in androgen receptor-driven

looping that selectively enhances gene activation. Proc. Natl. Acad. Sci. 111: 7319–7324.

https://doi.org/10.1073/pnas.1324151111

Jolma A., J. Yan, T. Whitington, J. Toivonen, K. R. Nitta, *et al.*, 2013 DNA-Binding Specificities of Human

Transcription Factors. Cell 152: 327–339. https://doi.org/10.1016/j.cell.2012.12.009

Kaikkonen M. U., N. J. Spann, S. Heinz, C. E. Romanoski, K. A. Allison, *et al.*, 2013 Remodeling of the

Enhancer Landscape during Macrophage Activation Is Coupled to Enhancer Transcription. Mol. Cell

51: 310–325. https://doi.org/10.1016/j.molcel.2013.07.010

Kent W. J., C. W. Sugnet, T. S. Furey, K. M. Roskin, T. H. Pringle, *et al.*, 2002 The human genome browser at

UCSC. Genome Res. 12: 996–1006. https://doi.org/10.1101/gr.229102

Kheradpour P., and M. Kellis, 2014 Systematic discovery and characterization of regulatory motifs in ENCODE

TF binding experiments. Nucleic Acids Res. 42: 2976–2987. https://doi.org/10.1093/nar/gkt1249

Kim T.-K., M. Hemberg, J. M. Gray, A. M. Costa, D. M. Bear, *et al.*, 2010 Widespread transcription at neuronal

activity-regulated enhancers. Nature 465: 182–187. https://doi.org/10.1038/nature09033

Landt S. G., G. K. Marinov, A. Kundaje, P. Kheradpour, F. Pauli, *et al.*, 2012 ChIP-seq guidelines and practices

of the ENCODE and modENCODE consortia. Genome Res. 22: 1813–1831.

https://doi.org/10.1101/gr.136184.111

Li H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, *et al.*, 2009 The Sequence Alignment/Map format and

SAMtools. Bioinformatics 25: 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Li H., 2013 Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. ArXiv13033997

Q-Bio.

Li W., D. Notani, Q. Ma, B. Tanasa, E. Nunez, *et al.*, 2013 Functional roles of enhancer RNAs for oestrogen-

dependent transcriptional activation. Nature 498: 516–520. https://doi.org/10.1038/nature12210

Lindblad-Toh K., M. Garber, O. Zuk, M. F. Lin, B. J. Parker, *et al.*, 2011 A high-resolution map of human

evolutionary constraint using 29 mammals. Nature 478: 476–482. https://doi.org/10.1038/nature10530

Lopes R., R. Agami, and G. Korkmaz, 2017 GRO-seq, A Tool for Identification of Transcripts Regulating Gene

Expression. Methods Mol. Biol. Clifton NJ 1543: 45–55. https://doi.org/10.1007/978-1-4939-6716-2_3

Maehara K., and Y. Ohkawa, 2015 agplus: a rapid and flexible tool for aggregation plots. Bioinformatics 31:

3046–3047. https://doi.org/10.1093/bioinformatics/btv322

Mahajan A., D. Taliun, M. Thurner, N. R. Robertson, J. M. Torres, *et al.*, 2018 Fine-mapping type 2 diabetes

loci to single-variant resolution using high-density imputation and islet-specific epigenome maps. Nat.

Genet. 50: 1505. https://doi.org/10.1038/s41588-018-0241-6

Martin M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal 17: 10–12. https://doi.org/10.14806/ej.17.1.200

Mathelier A., O. Fornes, D. J. Arenillas, C. Chen, G. Denay, *et al.*, 2016 JASPAR 2016: a major expansion and

update of the open-access database of transcription factor binding profiles. Nucleic Acids Res. 44:

D110–D115. https://doi.org/10.1093/nar/gkv1176

Matys V., O. V. Kel-Margoulis, E. Fricke, I. Liebich, S. Land, *et al.*, 2006 TRANSFAC and its module

TRANSCompel: transcriptional gene regulation in eukaryotes. Nucleic Acids Res. 34: D108-110.

https://doi.org/10.1093/nar/gkj143

Melgar M. F., F. S. Collins, and P. Sethupathy, 2011 Discovery of active enhancers through bidirectional

expression of short transcripts. Genome Biol. 12: R113. https://doi.org/10.1186/gb-2011-12-11-r113

Melnikov A., A. Murugan, X. Zhang, T. Tesileanu, L. Wang, *et al.*, 2012 Systematic dissection and optimization

of inducible enhancers in human cells using a massively parallel reporter assay. Nat. Biotechnol. 30:

271–277. https://doi.org/10.1038/nbt.2137

Mikhaylichenko O., V. Bondarenko, D. Harnett, I. E. Schor, M. Males, *et al.*, 2018 The degree of enhancer or

promoter activity is reflected by the levels and directionality of eRNA transcription. Genes Dev. 32: 42–

57. https://doi.org/10.1101/gad.308619.117

Mikkelsen T. S., M. Ku, D. B. Jaffe, B. Issac, E. Lieberman, *et al.*, 2007 Genome-wide maps of chromatin state

in pluripotent and lineage-committed cells. Nature 448: 553–560. https://doi.org/10.1038/nature06008

Murata M., H. Nishiyori-Sueki, M. Kojima-Ishiyama, P. Carninci, Y. Hayashizaki, *et al.*, 2014 Detecting

expressed genes using CAGE. Methods Mol. Biol. Clifton NJ 1164: 67–85. https://doi.org/10.1007/978-

1-4939-0805-9_7

Neumayr C., M. Pagani, A. Stark, and C. D. Arnold, 2019 STARR-seq and UMI-STARR-seq: Assessing

Enhancer Activities for Genome-Wide-, High-, and Low-Complexity Candidate Libraries. Curr. Protoc.

Mol. Biol. 128: e105. https://doi.org/10.1002/cpmb.105

Parker S. C. J., M. L. Stitzel, D. L. Taylor, J. M. Orozco, M. R. Erdos, *et al.*, 2013 Chromatin stretch enhancer

states drive cell-specific gene regulation and harbor human disease risk variants. Proc. Natl. Acad. Sci.

110: 17921–17926. https://doi.org/10.1073/pnas.1317023110

Pasquali L., K. J. Gaulton, S. A. Rodríguez-Seguí, L. Mularoni, I. Miguel-Escalada, *et al.*, 2014 Pancreatic islet

enhancer clusters enriched in type 2 diabetes risk-associated variants. Nat. Genet. 46: 136–143.

https://doi.org/10.1038/ng.2870

Patti M. E., A. J. Butte, S. Crunkhorn, K. Cusi, R. Berria, *et al.*, 2003 Coordinated reduction of genes of

oxidative metabolism in humans with insulin resistance and diabetes: Potential role of PGC1 and

NRF1. Proc. Natl. Acad. Sci. U. S. A. 100: 8466–8471. https://doi.org/10.1073/pnas.1032913100

Pickrell J. K., 2014 Joint Analysis of Functional Genomic Data and Genome-wide Association Studies of 18 Human Traits. Am. J. Hum. Genet. 94: 559–573. https://doi.org/10.1016/j.ajhg.2014.03.004

Purcell S., B. Neale, K. Todd-Brown, L. Thomas, M. A. R. Ferreira, *et al.*, 2007 PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. Am. J. Hum. Genet. 81: 559–575. https://doi.org/10.1086/519795

Quang D. X., M. R. Erdos, S. C. J. Parker, and F. S. Collins, 2015 Motif signatures in stretch enhancers are enriched for disease-associated genetic variants. Epigenetics Chromatin 8: 23. https://doi.org/10.1186/s13072-015-0015-7

Roman T. S., M. E. Cannon, S. Vadlamudi, M. L. Buchkovich, B. N. Wolford, *et al.*, 2017 A Type 2 Diabetes–Associated Functional Regulatory Variant in a Pancreatic Islet Enhancer at the ADCY5 Locus. Diabetes 66: 2521–2530. https://doi.org/10.2337/db17-0464

Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014 Biological insights from 108 schizophrenia-associated genetic loci. Nature 511: 421–427. https://doi.org/10.1038/nature13595

Schmidt E. M., J. Zhang, W. Zhou, J. Chen, K. L. Mohlke, *et al.*, 2015 GREGOR: evaluating global enrichment of trait-associated variants in epigenomic features using a systematic, data-driven approach. Bioinformatics 31: 2601–2606. https://doi.org/10.1093/bioinformatics/btv201

Scott L. J., M. R. Erdos, J. R. Huyghe, R. P. Welch, A. T. Beck, *et al.*, 2016 The genetic regulatory signature of type 2 diabetes in human skeletal muscle. Nat. Commun. 7: ncomms11764. https://doi.org/10.1038/ncomms11764

Seabold S., and J. Perktold, 2010 Statsmodels: Econometric and Statistical Modeling with Python. Proc. 9th Python Sci. Conf. 5.

The ENCODE project Consortium, 2012 An Integrated Encyclopedia of DNA Elements in the Human Genome. Nature 489: 57–74. https://doi.org/10.1038/nature11247

The FANTOM Consortium, A. R. R. Forrest, H. Kawaji, M. Rehli, J. K. Baillie, *et al.*, 2014 A promoter-level mammalian expression atlas. Nature 507: 462. https://doi.org/10.1038/nature13182

The Roadmap Epigenomics Consortium, A. Kundaje, W. Meuleman, J. Ernst, M. Bilenky, *et al.*, 2015

924   Integrative analysis of 111 reference human epigenomes. Nature 518: 317–330.

925   https://doi.org/10.1038/nature14248

926 Thurner M., M. van de Bunt, J. M. Torres, A. Mahajan, V. Nylander, *et al.*, 2018 Integration of human

927   pancreatic islet genomic data refines regulatory mechanisms at Type 2 Diabetes susceptibility loci.

928   eLife 7: e31977. https://doi.org/10.7554/eLife.31977

929 Trynka G., C. Sandor, B. Han, H. Xu, B. E. Stranger, *et al.*, 2013 Chromatin marks identify critical cell types for

930   fine mapping complex trait variants. Nat. Genet. 45: 124–130. https://doi.org/10.1038/ng.2504

931 Udler M. S., J. Kim, M. von Grotthuss, S. Bonàs-Guarch, J. B. Cole, *et al.*, 2018 Type 2 diabetes genetic loci

932   informed by multi-trait associations point to disease mechanisms and subtypes: A soft clustering

933   analysis. PLOS Med. 15: e1002654. https://doi.org/10.1371/journal.pmed.1002654

934 Varshney A., L. J. Scott, R. P. Welch, M. R. Erdos, P. S. Chines, *et al.*, 2017 Genetic regulatory signatures

935   underlying islet gene expression and type 2 diabetes. Proc. Natl. Acad. Sci. 114: 2301–2306.

936   https://doi.org/10.1073/pnas.1621192114

937 Varshney A., H. VanRenterghem, P. Orchard, A. P. Boyle, M. L. Stitzel, *et al.*, 2019 Cell Specificity of Human

938   Regulatory Annotations and Their Genetic Effects on Gene Expression. Genetics 211: 549–562.

939   https://doi.org/10.1534/genetics.118.301525

940 Viñuela A., A. Varshney, M. van de Bunt, R. B. Prasad, O. Asplund, *et al.*, 2019 Influence of genetic variants

941   on gene expression in human pancreatic islets – implications for type 2 diabetes. bioRxiv 655670.

942   https://doi.org/10.1101/655670

943 Visel A., S. Minovitsky, I. Dubchak, and L. A. Pennacchio, 2007 VISTA Enhancer Browser—a database of

944   tissue-specific human enhancers. Nucleic Acids Res. 35: D88–D92. https://doi.org/10.1093/nar/gkl822

945 Ward L. D., and M. Kellis, 2012 Evidence of abundant purifying selection in humans for recently acquired

946   regulatory functions. Science 337: 1675–1678. https://doi.org/10.1126/science.1225057

947 Yang Y., Z. Su, X. Song, B. Liang, F. Zeng, *et al.*, 2016 Enhancer RNA-driven looping enhances the

948   transcription of the long noncoding RNA DHRS4-AS1, a controller of the DHRS4 gene cluster. Sci.

949   Rep. 6: 20961. https://doi.org/10.1038/srep20961

Zheng H., J. Fu, P. Xue, R. Zhao, J. Dong, *et al.*, 2015 CNC-bZIP Protein Nrf1-Dependent Regulation of

Glucose-Stimulated Insulin Secretion. Antioxid. Redox Signal. 22: 819–831.

https://doi.org/10.1089/ars.2014.6017

Zhou V. W., A. Goren, and B. E. Bernstein, 2011 Charting histone modifications and the functional organization

of mammalian genomes. Nat. Rev. Genet. 12: 7–18. https://doi.org/10.1038/nrg2905

Zorita E., P. Cuscó, and G. J. Filion, 2015 Starcode: sequence clustering based on all-pairs search.

Bioinformatics 31: 1913–1919. https://doi.org/10.1093/bioinformatics/btv053

Figure 1: Islet CAGE tag cluster identification. A: Genome browser view of the intronic region of the *ST18* gene as an example locus where an islet TC overlaps an islet ATAC-seq peak and active TSS chromatin state. This TC also overlaps an enhancer element which was validated by the VISTA project (Visel *et al.* 2007). Also shown is the human-mouse-rat conserved TF binding site (TFBS) track from the Transfac Matrix Database (Matys *et al.* 2006). B: Cumulative fraction of islet TC segments overlapping with TCs identified in x FANTOM tissues. C: Distribution of the log10(distance to the nearest known protein coding gene TSS + 1bp) while classifying Islet TC segments by the number of FANTOM tissues where TCs overlap. Number of TC segments in each category are shown in parentheses. D: Genome browser view of an example locus near the *AP1G2* gene that highlights an islet TC (blue box) that is also identified in FANTOM tissues (FANTOM TCs track is a dense depiction of TCs called across 118 human tissues), occurs in a ATAC-seq peak region in both islets and lymphoblastoid cell line GM12878 (ATAC-seq track) and overlaps active TSS chromatin states across numerous tissues. Another islet TC (orange box) ~34 kb distal to the *AP1G2* gene is not identified as a TC in other FANTOM tissues, occurs in an islet ATAC-seq peak and a more islet-specific active enhancer chromatin state region. E: Enrichment of islet TCs to overlap islet chromatin state and other common annotations. Error bars represent the 95% confidence intervals. Colors represent significant enrichment after Bonferroni correction accounting for 40 total annotations (see Methods for references of common annotations, Supplementary Table 1), nominal enrichment (P value < 0.05) or non-significant enrichment. F: Enrichment of islet TCs to overlap GWAS loci of various disease/traits. Number of independent loci for each trait are noted in

parentheses. Colors represent significant enrichment after Bonferroni correction accounting for total 116 traits (see Methods for GWAS data sources, Supplementary Table 2), nominal enrichment (P value < 0.05) or non-significant enrichment. Acronyms: T2D = type 2 diabetes, ASD = Autism Spectrum Disorder, LDL = Low density lipoprotein, HDL = High density lipoprotein, BMI = Body mass index.

Figure 2. Integrating Islet CAGE TCs with other epigenomic information reveals characteristics of transcription initiation. A: Aggregate CAGE profiles over ATAC-seq peak summits. B: Aggregate ATAC-seq profile over TC midpoints. C: Enrichment of TF footprint motifs to overlap 500 bp upstream region (y axis) vs 500 bp

downstream region (x axis) of islet TCs. Colors denote if a TF footprint motif was significantly enriched (5% FDR correction, Benjajmini-Yekutieli method) to overlap only upstream regions, only downstream regions, both or none. D: Chromatin state annotations across 98 Roadmap Epigenomics cell types (using the 18 state 'extended model' (The Roadmap Epigenomics Consortium *et al.* 2015) for TC segments that occur in islet promoter chromatin states and overlap ATAC-seq peaks. These segments were segregated into those occurring 5kb proximal (left, N=7,064 TC segments) and distal (right, N=443 TC segments) to known protein coding gene TSS (Gencode V19). E: Chromatin state annotations across 98 Roadmap Epigenomics cell types for TC segments that occur in islet enhancer chromatin states and overlap ATAC-seq peaks, segregated into those occurring 5kb proximal (left, N=254 TC segments) and distal (right, N=289 TC segments) to known protein coding gene TSS. F: Enrichment of TF footprint motifs to overlap TCs occurring in accessible enhancer chromatin states (y axis) vs TCs occurring in accessible promoter chromatin states (x axis). Colors denote if a TF footprint motif was significantly enriched (5% FDR correction, Benjajmini-Yekutieli method) to overlap only TCs in accessible enhancer regions, only TCs in accessible promoter regions, both or none. G: Aggregate CAGE profiles centered and oriented relative to RFX5_known8 footprint motifs occurring in 5kb TSS distal regions. H: Aggregate CAGE profiles centered and oriented relative to ELK4_1 footprint motifs.

Figure 3. Experimental validation of CAGE elements using MPRA assay: A: (Top) Number and fraction of CAGE elements that show significant (5% FDR), nominal (P < 0.05) or non-significant transcriptional activity in the MPRA assay performed in rat beta cell insulinoma (INS1 832/13) cell line model. (Bottom) Proportion of CAGE elements overlapping promoter (active, weak or flanking TSS), enhancer (active, weak or genic enhancer) or other chromatin states that showed significant transcriptional activity in the MPRA assay. B. MPRA activity Z scores for CAGE elements overlapping in promoter, enhancer or other chromatin states. C:

MPRA activity Z scores for CAGE elements that overlap ATAC-seq peak vs CAGE elements that do not overlap peaks. D: (Top) An overview of the lasso regression model to predict the MPRA activity Z scores of CAGE elements as a function of the TF motif scan scores within the element. (Bottom) Top 30 TF motifs with non-zero coefficients from the model. E: An example locus on chr17, where the nearest gene *RPH3AL* lies ~6kb away, an islet TC overlaps active TSS and enhancer chromatin states and an ATAC-seq peak. Elements overlapping this TC showed significant transcriptional activity in the MPRA assay. The CAGE profile coincides with islet mRNA profile that is detected despite no known gene annotation in the region and the nearest protein coding gene is ~6kb away. Also shown are occurrences of TF motifs with positive or negative lasso regression coefficients from the analysis in D.

Figure 4. Islet TCs supplement functional understanding of GWAS and eQTL associations and help nominate causal variants. A: Enrichment of T2D GWAS (left) or islet eQTL (right) loci in annotations that comprise different levels of epigenomic information, including including chromatin state, ATAC-seq and TCs. Annotations defined using combinations of these datasets are depicted with different colors on the y axis. Enrichment was calculated using fGWAS (Pickrell 2014) using summary statistics from GWAS (left) (Mahajan *et al.* 2018) or islet eQTL (right) (Varshney *et al.* 2017). Error bars denote the 95% confidence intervals. B:

fGWAS conditional enrichment analysis testing the contribution of islet TC or ATAC-seq peak annotations after conditioning on histone-only based annotations such as active TSS and active enhancer chromatin states in islets. C: Maximum SNP PPA per T2D (BMI-unadjusted) GWAS locus after functional re-weighting using a model with islet chromatin states and ATAC-seq peak annotations (x axis) or chromatin states, ATAC-seq peaks and TC (y axis) annotations. D: The LCORL T2D GWAS locus showing SNPs in the 99% credible set from genetic fine-mapping. This locus comprises genes *DCAF16, NCAPG and LCORL.* The lead GWAS SNP is labelled in red, along with LD r^2 > 0.8 proxy SNPs in the top track. Also shown are CAGE, TC, ATAC-seq and chromatin state tracks. E: Browser shot of the *DCAF16* and *NCAPG* promoter regions where rs7667864 and eQTL lead SNP rs2074974 overlap an ATAC-seq peak. An overlapping CAGE element showed significant activity in the MPRA assay. Also shown are TF motifs with positive or negative coefficients from the MPRA lasso regression analysis.

Supplementary Figure 1: Islet TC identification using CAGE data across multiple samples. TC segments called using the paraclu method in each of the 57 selected islet samples were merged in a strand specific manner. Shown here is the number of merged TC segments that overlap TCs in x or more islet samples. We required TC overlap in a minimum of 10 islet samples to include a segment in the aggregate list of islet TCs.

Supplementary Figure 2: Distribution of islet TC lengths.

Supplementary Figure 3: 11 chromatin state model. Shown are the emission probabilities of each of the five histone marks, chromatin state annotation and the percent genome coverage of each state

Supplementary Figure 4: INS gene locus. Shown are Islet TCs along with TCs called in FANTOM tissues at the *INS* locus.

Supplementary Figure 5. Correlation between replicates for normalized total RNA counts for CAGE inserts.

Supplementary Figure 6: MPRA activity Z scores for CAGE elements based on position relative to known protein coding gene TSSs (5kb TSS proximal or distal, Gencode V19).

Supplementary Figure 7: The maximal SNP PPA from genetic fine-mapping for a locus vs the maximal SNP PPA after functional re-weighting using islet chromatin states, ATAC-seq peaks and TC annotations.

A

T2D GWAS lead SNP rs12640250

T2D GWAS lead SNP + r2>0.8proxies

T2D 99% genetic credible set SNPs

100 kb   hg19

Islet CAGE   0

-20

Islet TCs

Islet ATAC-seq   0

Islet ATAC-seq peaks

Chromatin states

Islets
Skeletal Muscle
Adipose
Liver

Islet mRNA-seq   0

-5

DCAF16   LCORL
DCAF16   LCORL
DCAF16   LCORL
NCAPG   LCORL
LCORL

Chromatin States

Active TSS	Active Enhancer	Strong Transcription	Repressed Polycomb
Weak/Flanking TSS	Weak Enhancer	Weak Transcription	Weak Repressed Polycomb
Bivalent/Poised TSS	Genic Enhancer		Quiescent/Low Signal

B

Re-weighthed SNP PPA using islet chromatin states + ATAC-seq peaks + TCs

rs7667864

eQTL lead rs2074974

GWAS lead rs12640250

SNP genetic PPA

Supplementary figure 8: A: LCORL GWAS locus showing all SNPs in the 99% credible set after genetic fine-mapping. B: Genetic fine-mapping PPA and functionally re-weighted PPA for SNPs at the LCORL locus.