Title: Single Cell Transcriptional Profiling of Phox2b-Expressing Geniculate Ganglion Neurons

Authors: Catherine B. Anderson and Eric D. Larson

Affiliations: Department of Otolaryngology, University of Colorado Anschutz Medical Campus, Aurora CO

80045

Article Type: Original Report

Author Roles: CBA performed experiments and analyzed data. EDL conceived and executed the study,

performed analyses, and wrote the manuscript. All authors reviewed and approved the final manuscript.

Corresponding Author:

Eric D. Larson

University of Colorado, Department of Otolaryngology

12700 E. 19th Ave, MS 8606

Aurora, CO 80045

Key Words: Phox2b, geniculate ganglion, single cell sequencing, taste

Abstract

The sense of taste is fundamental for survival as harmful substances can be discriminated and prevented from entering the body. Taste buds act as chemosensory sentinels and detect bitter, salty, sweet, sour, and umami substances. Taste receptor cells within taste buds detect these sapid molecules and transmit signals to afferent nerve fibers. Whether a single gustatory nerve fiber selectively is responsive to a single taste modality (through taste receptor cell activation) is a point of contention in the field. Two studies have attempted to molecularly classify individual gustatory neurons using single cell RNA sequencing and two others have attempted to classify neurons based on physiological properties, with some conflicting results. In the present study, we present a method for single cell RNA sequencing of gustatory geniculate ganglion neurons and compare the results obtained to prior published works. We find multiple classes of ganglion neurons, but in conjunction with published works, the only clusters that can be clearly attributed to specific taste modalities are those involved in sweet or sour detection.

Introduction:

Taste buds detect sapid molecules in the mouth and are the initiators of gustation. Each taste bud contains 50-100 taste receptor cells, which are classified as three separate types. Type I cells are the least understood, but act as supporting cells and may transduce come portion of salty taste. Type II cells are well-understood and transduce signals in response to bitter, sweet, or umami stimuli. Type II cells communicate with afferent nerve fibers through release of ATP via non-traditional synaptic mechanisms(Finger et al., 2005; Huang et al., 2007; Romanov et al., 2007, 2018; Taruno et al., 2013). Type III cells form classical synapses with afferent nerve fibers and are responsible for sour and portions of salt taste detection. Due to the different nature of how Type II and Type III cells communicate with afferent fibers, it is likely that the innervating fibers have different molecular and physiological properties. Indeed, we have shown that afferent fibers that express the serotonin receptor 5-HT3A preferentially contact serotonergic Type III cells (Stratford et al., 2017). This subset of afferent fibers responds to exogenous serotonin as well as ATP while other gustatory fibers respond only to ATP (Larson et al., 2015). These data suggest at least one subpopulation of gustatory afferent nerve fibers.

Lingual taste fields are innervated by branches of the facial (cranial nerve VII) and glossopharyngeal nerve (cranial nerve IX). The posterior tongue receives input from the lingual branch of the glossopharyngeal nerve while the anterior tongue receives input from the chorda tympani branch of the facial nerve. The cell bodies of the chorda tympani reside in the geniculate ganglion. In addition to the chorda tympani, this ganglion houses the cell bodies of the greater superficial petrosal and the nervus intermedius which receive gustatory input from the soft palate and somatosensory input from the external auditory meatus (Mtui et al., 2011).

We previously showed that the majority of geniculate ganglion neurons show excitatory responses to exogenously applied ATP and about 25% to serotonin via P2X2/P2X3 and 5HT3A receptors, respectively (Larson et al., 2015; Vandenbeuch et al., 2015). Older studies using patch clamp electrophysiology of rat gustatory geniculate ganglion neurons also showed excitatory responses to ACh, serotonin, substance P, and GABA in a small percentage of cells (King and Bradley, 2000; Koga and Bradley, 2000). These data suggest that multiple classes of neurons exist at the physiological and that likely, these classes could be reflected at the molecular level.

Gustatory geniculate ganglion neurons are delineated by expression of the transcription factor, *Phox2b* (Dvoryanchikov et al., 2017; Ohman-Gault et al., 2017; Rios-Pilier and Krimm, 2019) which can molecular define a neuron as gustatory or non-gustatory. The first study to characterize the transcriptome of single geniculate ganglion neurons utilized the Fluidigm C1 to profile 96 neurons from wildtype mice (Dvoryanchikov et al., 2017). They found about 2/3 of the cells express *Phox2b*, consistent with histological studies. Within the gustatory cells, three main clusters were identified, all of which expressed *Phox2b* and *P2rx3*. Other cluster markers included T1: *Olfm3*, *Itm2a*, *Hspb3*; T2: *Cd302*, *Kcnip1*, *Slc39a11*; T3: *Lypd1*, *Penk*, and *Trhr*. The authors then characterized the physiological response profiles of isolated ganglion neurons to exogenous ATP and 5-HT to match functional profiling with transcriptional profiling and demonstrated multiple physiological profiles that nicely matched expression of P2X and 5-HT receptor mRNA. This study was the first to characterize geniculate ganglion neurons at the molecular level using scRNA-seq.

More recently, Zhang et al. performed scRNA-sequence analysis on a larger cohort of geniculate ganglion neurons (Zhang et al., 2019). They profiled over 400 neurons using the Cel-seq (Hashimshony et al., 2012) method and identified 7 clusters. Based on cluster markers, they made multiple transgenic and knockout mice to test the role of certain transcripts in taste detection. Knockout of *Cdh4* and *Cdh13* eliminated detection of umami and bitter molecules, respectively, as assessed by two-bottle taste preference testing. Inhibition of *Egr2*-neurons using a *Egr2*-cre/Tetanus toxin model eliminated detection of NaCl. However, evidence for exactly how these transcripts are responsible for detecting taste modalities was not presented. However, more convincing evidence is presented for a class of sweet-best and sour-best neurons, marked by expression of *Spon1* and *Penk* transcripts, respectively. Using a floxed Gcamp model, the authors recorded in vivo calcium signals in geniculate ganglion neurons and showed that *Spon1* and *Penk* neurons selectively responded to sweet and sour lingual stimuli, respectively.

In the present study, we performed transcriptional profiling of geniculate ganglion neurons using an alternative method. We show that the neurons we collected expressed many transcripts overlapping with currently published datasets. Additionally, secondary analyses were performed on published datasets to further explore geniculate ganglion neuron clusters in attempt to understand more about the role of ganglion neurons in taste perception. Lastly, we show that combining all three datasets preserves cell clustering using the new analysis methods in Seurat V3.

Materials and Methods:

Animals

All animal procedures were conducted under a protocol approved by the University of Colorado Animal Care and Use Committee. Phox2bcre mice were purchased from Jax (B6(Cg)-Tg(Phox2b-cre)3Jke/J; stock # 016223) and bred to Rosa26/tdTomato mice (B6.Cg-Gt(ROSA)26Sortm14(CAG-tdTomato)Hze/J; stock # 007914). Crossed mice of both gender were used between the ages of 8 and 24 weeks.

Immunofluorescence

Animals were anesthetized using urethane and transcardially perfused with 4% paraformaldehyde in phosphate buffer (0.1M; PB) or euthanized with carbon dioxide. Geniculate ganglia and tongues were extracted and fixed for 1 or 4 hours at room temperature before incubating in 20% sucrose in PB overnight at 4C. Tissue was embedded and frozen in Optimal Cutting Temperature (OCT) and sectioned at a thickness of 12-16 um using a cryostat. After > 24 hours at -20C, slides were thawed, rehydrated with PBS, and blocked using 2% normal donkey serum in blocking buffer (PBS plus 0.3% TritonX100, 1% BSA). Primary antibodies were applied overnight at 4C. After thorough washing fluorescence secondary antibodies were applied at room temperature for 3 hours before counterstaining with DAPI and mounting with FluoroMountG (Southern Biotech). Slides were imaged using a Leica SP8 using 20x (NA 0.75) and 63x (NA 1.4) oil-immersion objectives.

Single cell preparation

Geniculate ganglia of three carbon dioxide-euthanized mice were rapidly extracted an placed in minimum

essential medium with Earle's balanced salts (MEM/EBSS; Hyclone) supplemented with 1.25 mg/ml trypsin

(Sigma-Aldrich) and 2.5 mg/ml collagenase A (Roche Diagnostics) for 30 minutes at 35C. Digested ganglia

were washed with MEM/EBSS, and gently triturated with a fire-polished glass pipette. The suspension was

passed through a 50 um filter and loaded as the top layer of a Percoll gradient (12.5 and 28%, Sigma-Aldrich;

Malin et al., 2007). The gradient was spun at 1300g for 10 minutes using a swinging bucket rotor. The cell

pellet at the bottom of the tube was washed with MEM/EBSS before resuspending in 2 mL MEM/EBSS.

Single cell capture and RNA extraction

A QiaScout Microraft Array (Qiagen) was coated with poly-D-lysine (0.02 mg/ml) for 2 hours and washed with

MEM/EBSS. The geniculate ganglion cell suspension was placed in the microraft array and the cells allowed to

settle for 20 minutes. The microraft array was secured using a custom set up to the stage of an Olympus IX71

inverted microscope equipped with a 10X objective and the QiaScout instrument (Qiagen). Cells were visually

inspected and microrafts containing a single tdTomato-positive (or negative in some cases) geniculate

ganglion neuron were extracted and transferred to a PCR tube containing lysis buffer as part of the QIAseq FX

Single Cell RNA Library Kit (Qiagen). Once in lysis buffer, cells were lysed by heating the tube to 95C for 3

minutes followed by an infinite hold at 4C until 24 individual samples were collected. The collection protocol

was performed across two independent experiments, resulting in 48 individual sequencing libraries (42

tdTomato+, 6 tdTomato-).

Library preparation and sequencing

Sequencing libraries were prepared using the QIAseq FX Single Cell RNA Library Kit (Qiagen) according to

manufacturer instructions. Cells were lysed, followed by gDNA removal, reverse transcription, ligation, and

whole transcriptome amplification. After amplification, libraries were enzymatically fragmented (~300 bp

fragments) followed by adapter ligation. Library size was evaluated using a DNA tapestation and concentration

was determined using qPCR (QIAseq Library Quant Array). Libraries were pooled at equimolar concentration

and sequenced using an Illumina MiSEQ to generate 2x151 bp reads.

Sequencing analysis

Quality and adapter trimming was performed using BBDuk (Bushnell). Salmon (v0.14.0) was used to quantify

transcript expression using trimmed reads and the Ensembl GRCm38.p6 (version 92) transcriptome (Patro et

al., 2017; Zerbino et al., 2018). Transcript expression was summarized at the gene level and input to R using

TxImport (Soneson et al., 2016; R Core Team, 2019). A custom R script was used to visualize gene

expression in the dataset. Samples were kept if expression of *Snap25* or *Uclh1* was > 5 TPM. Discarded

samples were further interrogated for expression of neuron-specific genes, but they were rarely present.

Sequencing analysis of published datasets

Single cell sequencing FASTQ files were obtained from NCBI Gene Expression Omnibus (GSE102443

(Dvoryanchikov et al., 2017) and GSE135801 (Zhang et al., 2019)). Raw reads from Dvoryanchikov et al. were

filtered, trimmed, mapped, and summarized at the gene-level using the same tools as described above. A

processed gene count matrix was obtained from Zhang et al. Gene count matrices were used as the input for

Seurat (V3; ref) where samples were filtered, scaled, and clustered.

Meta analysis of datasets

Individual Seurat objects were read in to R and processed using Seurat following the "Multiple Dataset

Integration and Label Transfer" vignette available on the program website

(https://satijalab.org/seurat/v3.1/integration.html).

Data visualization

Data were visualized using the pheatmap package in R or with Seurat plotting features DimPlot and

DoHeatmap and FeaturePlot (Butler et al., 2018; Kolde, 2019; Stuart et al., 2019).

Results:

Characterization of Phox2b-cre mouse

To confirm that the *Phox2b*-cre driver faithfully drives CRE recombinase in PHOX2B-expressing cells,

geniculate ganglia and gustatory taste fields of *Phox2b*-cre/Rosa-tomato mice were examined using confocal

immunofluorescence microscopy. A subset of geniculate ganglion neurons expresses robust tdTomato

fluorescence consistent with previous reports(Ohman-Gault et al., 2017; Rios-Pilier and Krimm, 2019). The

majority of tdTomato-expressing neurons exhibited P2X3 immunoreactivity, and conversely, nearly all P2X3-

immunoreactive neurons exhibited tdTomato fluorescence (Figure 1). To confirm that tdTomato is being driven

in PHOX2B-expressing ganglion cells, sections were labeled with an antibody against PHOX2B. Indeed, every

tdTomato-expressing neuron showed positive nuclear labeling of the transcription factor (Figure 1).

Within taste papillae, tdTomato positive nerve fibers enter taste buds consistent with the *Phox2b* as a

marker of gustatory nerve fibers. Most tdTomato-expressing nerve fibers exhibit strong P2X3-immunoreactivity.

Additionally, these fibers are rarely outside of taste buds but when present are not immunoreactive for P2X3

(Figure 2)

scRNA-seq of Phox2b geniculate ganglion neurons

To understand how gustatory nerve fibers communicate with taste receptor cells, single cell RNA sequencing

was performed on manually isolated geniculate ganglion neurons (48 samples from 6 total mice across 2

collection days; 40 tdTomato+, 6 tdTomato-, 2 blanks). Ganglion neurons were enzymatically digested and

plated on a Qiagen QiaScout Microraft array. Using the QiaScout device cells were visually inspected for

presence or absence of tdTomato, dislodged from the microraft array, and transferred to cell lysis buffer where

lysis was initiated followed by reverse transcription, whole transcriptome amplification, and sequencing library

preparation. Post processing of the sequencing data excluded samples that had < 50% mapping rate,

expression of *Snap25* or *Uchl1* < 5 TPM, and >2000 unique genes resulted in 31 cells (17 tdTomato+, 4

tdTomato-). *Phox2b* expression correlated with tdTomato fluorescence (Figure 3) and taste-related transcripts

(*P2rx2, P2rx3*) were present in all *Phox2b+* neurons. A subset of *Phox2b+* neurons also expressed *Htr3a*

(Figure 3). Additionally, genes such as *Penk and Spon1*, markers of sour-best and sweet-best neurons (Zhang

2019) were identified in some neurons. Unfortunately, the low sample size prevented use of unbiased

clustering algorithms such as Seurat (Stuart et al., 2019).

Analysis of other datasets

Zhang et al 2019 manually collected over 800 *P2rx3*-tdTomato expressing geniculate ganglion cells and

performed scRNA-sequencing using the Cel-seq technology. Of the 454 neurons available from GEO:

GSE135801, 258 had expression of >4000 unique genes. Zhang et al reported identifying clusters enriched for

Spon1, Penk, Cdh4, Cdh13, and Egr2 and showed that neurons from those classes were selective for sweet,

sour, umami, bitter, and salty taste modalities, respectively. Analysis of raw RPM from these cells shows

similar patters to the data in this paper including expression of *P2rx3, P2rx3, Phox2b, Gabra1, Penk, Spon1,*

and Htr3a (Figure 4A)*.* Seurat cluster analysis reveals 7 molecularly distinct neuronal clusters including those

marked by *Spon1* (cluster 2)*, Penk* (cluster 3)*, Cdh13* (clusters 1 and 5)*,* and *Piezo2* (clusters 0 and 6; Figure

4B,C)*.* However, *Cdh4* and *Egr2* were not detected as significant markers. The top 5 cluster makers (FDR-

pvalue <0.001, weighted by log fold change) for each cluster are plotted in Figure 4D. All cluster markers can

be found in Table S1.

An earlier report used wildtype mice and the Fluidigm C1 platform to perform expression profiling and

demonstrated 4 cell clusters (3 gustatory, 1 non-gustatory). Analysis of 96 cells using Seurat reveals 6 clusters

of cells (Figure 5B). The discrepancy among cluster numbers likely arises due to 1) different clustering

software/algorithms and 2) less conservative clustering variables. Indeed, the cluster identified by expression

of *Olfm3* in the original report was subdivided to three clusters in this current analysis (clusters 1,3,5). About

60% of profiled cells were gustatory based on expression of *Phox2b*. Consistent with Zhang et al 2019, *Penk*

(cluster 2), *Spon1* (cluster 5), *Piezo2* (clusters 0 and 4), and *Cdh13* (clusters 1 and 3) were significant clusters

identifiers (Figure 5C, D). Similar to above, neither *Egr2* nor *Cdh4* were identified as cluster markers. All cluster

markers can be found in Table S2.

Meta analysis of all datasets

Generally, in RNA-sequence analysis, it is against best-practices to combine datasets from multiple sources

given the range of variability in sample preparation, sequencing, and data processing. However, with recent

the release of Seurat V3 a method for combining datasets from single cell sequencing experiments across

multiple technologies has been developed (Stuart et al., 2019). This method finds integration anchors common

amongst the datasets, merges the data, and scales the data based on the anchors. This meta analysis allows for integration of data on similar cell types from multiple sources.

The three datasets in this paper can be combined, scaled, and clustered together. Interestingly, UMAP plotting of the integrated data shows a strong similarly among three datasets, with cells from each dataset being represented throughout the plot (Figure 6A,B). This suggests that despite three different methods and technologies used to collect scRNA-seq data, the genetic information was similar between studies. Clustering of the integrated data reveals 7 cell clusters. *Penk* (cluster 3), *Spon1* (cluster 4), *Piezo2* (clusters 0 and 6), and *Cdh13* (clusters 5), *Egr2* (cluster 2), and *Cdh4* (cluster 2) were significant cluster identifiers (Figures C,D). All cluster markers can be found in Table S3.

Discussion:

Do neuron clusters represent specific taste modalities?

Single cell RNA-seq allows for the determination of heterogeneity within a cell population. By clustering cells based on similarities, markers that are unique to those clusters can be identified, which can give strong insight to their function. While Dvoryanchikov et al. were the first reveal molecular heterogeneity by identifying clusters of geniculate ganglion neurons, Zhang et al. were the first to attribute the role of specific cell clusters to taste detection. Convincing evidence demonstrates that neurons in the *Penk* and *Spon1*- expressing cell clusters represent sour- and sweet-best neurons. Thus, it is plausible that these neurons receive specific input from Type III and Type II taste receptor cells, respectively.

Sour and sweet: All three datasets analyzed in the present study showed expression of *Penk* and *Spon1* in gustatory fibers and that they marked specific cell clusters. In Zhang et al 2019, *Penk* and *Spon1* were cluster markers. Dvoryanchikov et al 2017 reported *Penk* as a cluster marker, but not *Spon1*. Interestingly, secondary analysis reveals that *Spon1* neurons were a subset of the cluster labeled 'T1' in the original report, which was subdivided in to three clusters in this current study (Figure 5C). These findings together with the in vivo imaging of Zhang et al 2019 strongly suggest that sour-best and sweet-best fibers are molecularly distinct from other ganglion neurons.

Bitter, salty, and umami: Zhang et al reports that *Cdh4*, *Egr2*, and *Cdh13* are cluster markers for bitter, salty, and umami-best neurons. However, in the present analyses, these markers did not clearly define clusters as well as *Penk* and *Spon1*. Additionally, the behavioral assays performed by Zhang et al relied on constitutive knockout of *Cdh4* and *Cdh13* and tetanus toxin inactivation of *Egr2* neurons. While the experiments showed that umami, bitter, and salty taste preference rely on these transcripts, they do not show that this requirement is at the level of the geniculate ganglion. Indeed, according to the Allen Brain Map, these three transcripts are expressed in a variety of cell types throughout the central nervous system. Additionally, according to secondary analysis of Qin et al, *Cdh13* is expressed in both Type II and Type III taste receptor cells and *Egr2* is expressed in some Type III cells (not shown; Qin et al., 2018). Thus, any ganglion cell clusters responsible for the detection of bitter, salty, and umami signals from taste receptor cells remains enigmatic.

Does Htr3a mark a specific cluster?

We previously showed that nerve fibers expressing the serotonin receptor 5-HT3 preferentially contact

Type III taste receptor cells in all taste fields (Stratford et al 2017) suggesting that these fibers may be

molecularly distinct within the geniculate ganglion. However, Dvoryanchikov et al. argued that *Htr3a* did not

delineate a specific cluster in their analyses. In the present analyses, all three datasets show clustering of

Htr3a-expressing neurons. While *Htr3a* was in multiple clusters, it had strongest expression in the cluster

marked by *Penk* – the sour-best fibers which support our findings that these cells wire to sour-sensing Type III

taste receptor cells. Conversely, *Htr3a* is absent in the *Spon1* cluster (sweet-best fibers). Interestingly, in all

datasets, *Htr3a* is also present in a gustatory (*Phox2b*+) cluster expressing *Piezo2*, a transcript for a

mechanosensitive channel. This cluster expressed low levels of *P2rx3*, and no *P2rx2*. Moayedi et al showed

that *Piezo2*-GFP expressing nerve fibers are present in fungiform papillae, just adjacent to taste buds, and that

these fibers enter the papillae in a bundle with other gustatory fibers (Moayedi et al., 2018). Interestingly,

P2X2/3 double knockout mice (which lack nerve response to all tastants) still show nerve responses to

mechanical stimulation of the tongue (Finger et al., 2005). Perhaps this *Piezo2*/*Htr3a*/*Phox2b* neuronal cluster

is responsible for this ATP-independent response.

Conclusions

Overall, the data analyzed in this paper suggest molecular heterogeneity among the neurons of the geniculate

ganglion. The datasets clearly separate gustatory vs non-gustatory neurons by expression of *Phox2b*. Within

the *Phox2b* expressing fibers, clear sub clusters are formed. Consistent among the datasets are clusters

defined by *Penk*, *Spon1*, and *Piezo2*. Zhang et al showed that *Penk* clusters represent sour-best fibers while

Spon1 clusters represent sweet-best fibers. The *Piezo2* cluster could be responsible for mechanosensation of

the fungiform papillae. The remaining clusters were more variable between the datasets and markers identified

by Zhang et al (*Cdh4*, *Cdh13*, *Egr2*) were not clearly observed in this study. While it is likely that neurons

responsible for detecting each taste modality are molecularly distinct, more work needs to be done to elucidate

the molecular phenotypes of bitter, salty, and umami sensitive fibers.

Figure Legends:

Figure 1. tdTomato expression in PHOX2B-immunoreactive neurons. Sections of Phox2bcre/tdTomato

geniculate ganglion labeled with antibodies against PHOX2B and P2X3. Nearly all tdTomato-expressing cells

are immunoreactive for PHOX2B and P2X3. Non-nuclear "green" signal is due to binding of anti-mouse

secondary antibodies to endogenous mouse IgG; this signal persists in no primary controls ,and no nuclear

signal is observed. Sections are maximal z-projections of ~12 μm image stacks.

Figure 2. tdTomato-expressing nerve fibers enter taste buds. Sections of lingual taste papillae showing

immunoreactivity to GNAT3 (α-gustducin; Type II cells) and P2X3 with endogenous tdTomato fluorescence.

Within taste buds, all tdTomato+ fibers exhibit P2X3 immunoreactivity. In some cases, tdTomato+ fibers were

found in the intergemmal area and were P2X3 negative. Images are maximal z-projections of ~12-16 μm

image stacks.

Figure 3. Expression of select transcripts in tdTomato-expressing neurons. Geniculate ganglion neurons were

collected using the QiaScout and processed for single cell RNA sequencing using the Qiagen FX single cell

RNA kit. Heatmap shows log(transcripts per million) of select transcripts. *Phox2b* expression correlates with

presence or absence of tdTomato.

Figure 4. Secondary analysis of Zhang et al. 2019 scRNA-seq data. Analysis of cells publicly available from

GSE135801. A. Heatmap showing log(reads per million) of select transcripts across all cells. B. Seurat UMAP

clustering of cells expressing greater than 2000 genes. C. Heatmap showing transcript expression of same

genes as in A, but after scaling and clustering in Seurat. D. Heatmap showing transcript expression of top 5

markers of each cluster. Clusters in C and D correlate with clusters in B.

Figure 5. Secondary analysis of Dvoryanchikov et al. 2017 scRNA-seq data. Analysis of cells publicly available

from GSE102443. Raw FASTQ files were analyzed to quantify transcript expression. A. Heatmap showing

log(reads per million) of select transcripts across all cells. B. Seurat UMAP clustering of cells expressing

greater than 2000 genes. C. Heatmap showing transcript expression of same genes as in A, but after scaling

and clustering in Seurat. D. Heatmap showing transcript expression of top 5 markers of each cluster. Clusters

in C and D correlate with clusters in B.

Figure 6. Combination of data from this paper, Dvoryanchikov et al 2017, and Zhang et al 2019. A. Seurat

UMAP clustering showing each dataset. Each cluster has cells from each dataset, though some cluster lack

cells from 'this study' likely due to the small sample size. B. Overall clusters of the combined datasets. C.

Heatmap showing transcript expression of select genes after scaling and clustering in Seurat. D. Heatmap

showing transcript expression of top 5 markers of each cluster. Clusters in C and D correlate with clusters in B.

E. Gene expression overlay of UMAP cluster plot shows expression of each gene. Each point represents a

single cell.

Table S1. Seurat-identified cluster markers for Zhang et al 2018.

Table S2. Seurat-identified cluster markers for Dvoryanchikov et al 2017.

Table S3. Seurat-identified cluster markers for meta analysis.

Conflict of Interest:

The authors have no financial conflicts of interest to declare.

Funding:

This work was supported by NIH/NIDCD R21DC015115-01 to Sue C. Kinnamon (University of Colorado

Anschutz Medical Campus).

Acknowledgments:

We thank Qiagen for allowing us to demo the QiaScout device and supplying samples of the QiaScout

Microraft Arrays.

References:

Bushnell, B. BBMap.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. 2018. Integrating single-cell transcriptomic
data across different conditions, technologies, and species. Nature Biotechnology. 36:411–420.

Dvoryanchikov, G., Hernandez, D., Roebber, J.K., Hill, D.L., Roper, S.D., and Chaudhari, N. 2017.
Transcriptomes and neurotransmitter profiles of classes of gustatory and somatosensory neurons in the
geniculate ganglion. Nature Communications. 8:760.

Finger, T.E., Danilova, V., Barrows, J., Bartel, D.L., Vigers, A.J., Stone, L., Hellekant, G., and Kinnamon, S.C.
2005. ATP Signaling Is Crucial for Communication from Taste Buds to Gustatory Nerves. Science. 310:1495–
1499.

Hashimshony, T., Wagner, F., Sher, N., and Yanai, I. 2012. CEL-Seq: Single-Cell RNA-Seq by Multiplexed
Linear Amplification. Cell Reports. 2:666–673.

Huang, Y.-J., Maruyama, Y., Dvoryanchikov, G., Pereira, E., Chaudhari, N., and Roper, S.D. 2007. The role of
pannexin 1 hemichannels in ATP release and cell–cell communication in mouse taste buds. PNAS. 104:6436–
6441.

King, M.S., and Bradley, R.M. 2000. Biophysical properties and responses to glutamate receptor agonists of
identified subpopulations of rat geniculate ganglion neurons. Brain Research. 866:237–246.

Koga, T., and Bradley, R.M. 2000. Biophysical Properties and Responses to Neurotransmitters of Petrosal and
Geniculate Ganglion Neurons Innervating the Tongue. J Neurophysiol. 84:1404–1413.

Kolde, R. 2019. pheatmap: Pretty Heatmaps.

Larson, E.D., Vandenbeuch, A., Voigt, A., Meyerhof, W., Kinnamon, S.C., and Finger, T.E. 2015. The Role of
5-HT3 Receptors in Signaling from Taste Buds to Nerves. J Neurosci. 35:15984–15995.

Malin, S.A., Davis, B.M., and Molliver, D.C. 2007. Production of dissociated sensory neuron cultures and
considerations for their use in studying neuronal function and plasticity. Nat Protocols. 2:152–160.

Moayedi, Y., Duenas-Bianchi, L.F., and Lumpkin, E.A. 2018. Somatosensory innervation of the oral mucosa of
adult and aging mice. Sci Rep. 8:1–14.

Mtui, E., Gruener, G., and FitzGerald, M. 2011. Clinical Neuroanatomy and Neuroscience - 6th Edition.

Ohman-Gault, L., Huang, T., and Krimm, R. 2017. The transcription factor Phox2b distinguishes between oral
and non-oral sensory neurons in the geniculate ganglion. J Comp Neurol. 525:3935–3950.

Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. 2017. Salmon: fast and bias-aware
quantification of transcript expression using dual-phase inference. Nat Methods. 14:417–419.

Qin, Y., Sukumaran, S.K., Jyotaki, M., Redding, K., Jiang, P., and Margolskee, R.F. 2018. Gli3 is a negative
regulator of Tas1r3-expressing taste cells. PLOS Genetics. 14:e1007058.

R Core Team. 2019. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation
for Statistical Computing.

Rios-Pilier, J., and Krimm, R.F. 2019. TrkB expression and dependence divides gustatory neurons into three
subpopulations. Neural Dev. 14:3.

Romanov, R.A., Lasher, R.S., High, B., Savidge, L.E., Lawson, A., Rogachevskaja, O.A., Zhao, H.,
Rogachevsky, V.V., Bystrova, M.F., Churbanov, G.D., et al. 2018. Chemical synapses without synaptic

vesicles: Purinergic neurotransmission through a CALHM1 channel-mitochondrial signaling complex. Sci
Signal. 11:eaao1815.

Romanov, R.A., Rogachevskaja, O.A., Bystrova, M.F., Jiang, P., Margolskee, R.F., and Kolesnikov, S.S. 2007.
Afferent neurotransmission mediated by hemichannels in mammalian taste cells. The EMBO Journal. 26:657–
667.

Soneson, C., Love, M.I., and Robinson, M.D. 2016. Differential analyses for RNA-seq: transcript-level
estimates improve gene-level inferences. F1000Res. 4.

Stratford, J., Larson, E., Yang, R., Salcedo, E., and Finger, T. 2017. 5-HT3a-driven GFP delineates Gustatory
Fibers innervating Sour-responsive Taste Cells: A Labeled Line for Sour Taste? J Comp Neurol. 525:2358–
2375.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao, Y., Stoeckius, M.,
Smibert, P., and Satija, R. 2019. Comprehensive Integration of Single-Cell Data. Cell. 177:1888-1902.e21.

Taruno, A., Vingtdeux, V., Ohmoto, M., Ma, Z., Dvoryanchikov, G., Li, A., Adrien, L., Zhao, H., Leung, S.,
Abernethy, M., et al. 2013. CALHM1 ion channel mediates purinergic neurotransmission of sweet, bitter and
umami tastes. Nature. 495:223–226.

Vandenbeuch, A., Larson, E.D., Anderson, C.B., Smith, S.A., Ford, A.P., Finger, T.E., and Kinnamon, S.C.
2015. Postsynaptic P2X3-containing receptors in gustatory nerve fibres mediate responses to all taste qualities
in mice. J Physiol. 593:1113–1125.

Zerbino, D.R., Achuthan, P., Akanni, W., Amode, M.R., Barrell, D., Bhai, J., Billis, K., Cummins, C., Gall, A.,
Girón, C.G., et al. 2018. Ensembl 2018. Nucleic Acids Res. 46:D754–D761.

Zhang, J., Jin, H., Zhang, W., Ding, C., O'Keeffe, S., Ye, M., and Zuker, C.S. 2019. Sour Sensing from the
Tongue to the Brain. Cell.

Figure 1

GNAT3/P2X3   **GNAT3/tdTomato**   **Merge**

Circumvallate

Fungiform

Soft Palate

25 μm

25 μm

25 μm

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6